



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search


Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



PRIVATE LIBRARY
—o—
DUDLEY SMITH
No. _____
Cont. _____ Date _____




Gift of
Dr. Dudley Smith



~~1/2~~

Handwritten notes, possibly a signature or initials, including the letters "H", "N", and "F".



THE STUDENT'S
MEDICAL DICTIONARY
—
GOULD

RECOMMENDATIONS AND PRESS NOTICES

OF PREVIOUS EDITIONS OF

GOULD'S STUDENT'S MEDICAL DICTIONARY

"One pleasing feature of the book is that the reader can almost invariably find the definition under the word he looks for, without being referred from one place to another, as is too commonly the case in medical dictionaries. The tables of the bacilli, micrococci, leucomains and ptomains are excellent, and contain a large amount of information in a limited space. The anatomical tables are also concise and clear. . . . We should unhesitatingly recommend this dictionary to our readers, feeling sure that it will prove of much value to them."—*American Journal of Medical Science*.

"Again and again we have submitted the book to tests, and we have found it reliable and full, every page giving proof of careful editing and research. It is sufficiently large to meet the requirements of any ordinary practitioner. Of course, the bespectacled, polysyllabic, home-from-Berlin young man could not take pleasure or feel satisfied in anything short of a multi-voluminous work; but, like the Heathen, he is a law unto himself, and may safely be left out of calculation in practical therapeutics. In addition to the dictionary proper we have elaborate tables of bacilli, micrococci, leucomains and ptomains, and all such infinitesimal creatures as we to-day build laboratories to hatch; tables with analyses of American mineral waters, some of which waters will soon, we believe, be found in our midst; and tables of vital statistics. All this great mass of information is excellently arranged, so that the reader has no difficulty in at once finding what he wants, and the type is beautifully clear, there being no blurring, so that reading is a pleasure. As we turn over the pages we are grateful for the free-trade system that admits, free of duty, such a useful and desirable book."—*The Dublin Journal of Medical Science*.

"We know of no work in which so many important and yet isolated facts may be obtained without great expenditure of time. The tables of muscles, nerves, arteries, etc., seem to be very complete, and in looking through the dictionary, we were struck by the good presswork, clear type, and handsome paper, while no one of the many words which it occurred to us to look for have been found absent."—*Therapeutic Gazette*.

"As a handy, concise and accurate, and complete medical dictionary it decidedly claims a very high place among works of this description. In fact, taking handiness and cheapness into account, we certainly think this is the general practitioner's model dictionary, and we cordially recommend it to our readers. The definitions are for the most part terse and accurate, and the derivations up to modern lights."—*British Medical Journal*.

"I find it an excellent work, doing credit to the learning and discrimination of the author."—*Dr. J. M. Da Costa, Prof. of Practice of Medicine, Jefferson Medical College*.

"In gynecology, ophthalmology, otology, and laryngology; in biology, embryology, physiology, and pathology; in electro-therapeutics, and in the newly-developed fields of bacteriology, ptomains and leucomains, the aim has been evidently to issue an authoritative text-book, one that should be ample in its vocabulary, concise in its definitions, compact in its arrangement, and convenient of size for the everyday use of busy practitioners and as a handbook for medical students. The author, in this respect, is to be congratulated upon his success, and so far as a careful examination enables us to judge, it faithfully represents the medical literature of to-day."—*Journal of American Medical Association*.

"The work of Dr. Gould claims to be essentially a new work, all definitions being framed 'by the direct aid of new, standard, and authoritative text-books.' It certainly bears very little resemblance to previous works of the kind, in nearly all of which a great deal of space is devoted to obsolete terms. . . . More complete and more up to date than any other medical dictionary of similar dimensions in our own, or, indeed, as far as we know, in any other language."—*London Lancet*.

THE STUDENT'S MEDICAL DICTIONARY

INCLUDING ALL THE WORDS AND PHRASES GENERALLY
USED IN MEDICINE, WITH THEIR PROPER PRO-
NUNCIATION AND DEFINITIONS.

BASED ON RECENT MEDICAL LITERATURE

BY

GEORGE M. GOULD, A.M., M.D.

AUTHOR OF "AN ILLUSTRATED DICTIONARY OF MEDICINE, BIOLOGY, AND ALLIED SCIENCES," "12,000
MEDICAL WORDS PRONOUNCED AND DEFINED," "THE MEANING AND THE METHOD OF LIFE,"
"BORDERLAND STUDIES;" FORMERLY EDITOR OF "THE MEDICAL NEWS;"
PRESIDENT, 1893-1894, AMERICAN ACADEMY OF MEDICINE

WITH ELABORATE TABLES OF THE BACILLI, MICROCOCCI, LEUCOMAINS, PTOMAINS,
ETC. OF THE ARTERIES, GANGLIA, MUSCLES, AND NERVES; OF WEIGHTS
AND MEASURES, ANALYSES OF THE WATERS OF THE MINERAL
SPRINGS OF THE UNITED STATES, ETC., ETC.

TENTH EDITION, REWRITTEN AND ENLARGED

NEW YORK

PHILADELPHIA
P. BLAKISTON, SON & CO.
1012 WALNUT STREET

1896

COPYRIGHTED, 1896, BY P. BLAKISTON, SON & Co.

YASSEL 3851

WM. F. FELL & CO.,
ELECTROTYPERS AND PRINTERS,
1220-24 SANBOM STREET,
PHILADELPHIA.

PREFACE TO TENTH EDITION.

THE present volume is an entirely new one and is designed to take the place of the NEW MEDICAL DICTIONARY and the STUDENT'S MEDICAL DICTIONARY, the plates of which have been destroyed. Despite their popularity, the older editions were unsatisfactory to me, and I determined to replace them by one that should more perfectly represent the recent rapid progress in medical science, and that would more adequately meet the ordinary lexicographic demands of the student and practicing physician. The ILLUSTRATED MEDICAL DICTIONARY issued in 1894 (the third edition in 1896) is more encyclopedic in character than the foregoing volume, has many new and large tables or groupings of systematized medical facts, such as those of Stains, Parasites, Pigments, Tests, etc., etc., that could not be included in the present book. The prominent feature of illustrations is also here omitted. Thus to the larger work one is referred for the more complete information desired by the physician, the educated layman, lawyer, and the student of the more highly specialized branches of medicine.

This, the STUDENT'S MEDICAL DICTIONARY, is especially adapted to the wants of medical students. It cannot be used as a substitute for, but rather as an introduction to, "The Illustrated," or for a brief suggestion of a more extended or half-forgotten definition. In it one will find correct and succinct definitions of all the more common words liable to be found in the usual text-books, compends, lectures, etc., of those engaged in the study of medicine or in the early years of practice.

Much care has been given to the difficult subject of pronunciation, and what has seemed as the most commendable method of pronouncing each word has been sought, and a simple, easily-understood manner of expressing the

same has been adopted, that does away with all diacritic marks and bothersome as well as doubtful means, so that "he who runs may read."

A vigorous endeavor has been made to keep the volume within the limit of a "handy" size, and the cordial cooperation of both publisher and printer has in this respect resulted in success, although there is here perhaps twice the amount of matter contained in the old editions.

I am unable to express my gratitude to Dr. David Riesman, who has unreservedly devoted to the work a scholarship and a painstaking conscientiousness of which every line bears evidence. I also wish to acknowledge my indebtedness to Dr. Clarence W. Lincoln, Dr. W. A. N. Dorland, and Dr. Emma Billstein for valuable assistance.

GEORGE M. GOULD.

May, 1896.

ABBREVIATIONS.

ABBREVIATIONS USED IN GENERAL MEDICINE.

aa	ana	Of each.	Cent.	Centigrade.
A. c.	Ante cibum	Before meals.	Cg.	Centigram.
Abdom.	Abdomen	The belly.	C. m.	Cras mane
Aba. feb.	Absente febre	When fever is absent.	cm.	Centimeter.
Abstr.	Abstractum	Abstract.	C. m. s.	Cras mane sumendus,
Ad.	Adde	Add.	C. n.	Cras nocte
Ad lib.	Ad libitum	At pleasure, as desired.	Co.	Cobalt.
Admov.	Admoveatur	Let it be applied.	Cochl.	Cochleare
Ad pond.	Ad pondus omnium,	To the weight of the whole.	Cochl. ampl.	Cochleare amplum,
Adv.	Adversum	Against.	Cochl. infant.	Cochleare infantis,
Aggred.	Aggrediente febre,	While fever is coming on.	Cochl. mag.	Cochleare magnum,
Al.	Aluminum	Aluminum.	Cochl. med.	Cochleare medium,
Al. dieb.	Alterius diebus	Every other day.	Cochl. parv.	Cochleare parvum,
Alt. hor.	Alterius horis	Every other hour.	Col.	Cola
Alv. adstrict.	Alvo adstricto	The bowels being confined.	Colat.	Colatus
Alv. deject.	Alvi dejectiones,	The intestinal evacuations.	Colet.	Coletur
Anat.	Anatomy	Anatomy.	Color.	Coloretur
Aq.	Aqua	Water.	Comp.	Compositus
Aq. astr.	Aqua astricta	Ice.	Cong.	Congius
Aq. bull.	Aqua bulliens	Boiling water.	Cons.	Conserva
Aq. com.	Aqua communis	Common water.	Coq.	Coque
Aq. dest.	Aqua destillata	Distilled water.	Cort.	Cortex
Aq. ferv.	Aqua fervens	Hot water.	Crast.	Crastinus
Aq. font.	Aqua fontana	Spring water.	Crys.	Crystal.
Aq. mar.	Aqua marina	Ocean water.	Cs.	Cesium.
Aq. pur.	Aqua pura	Pure water.	Cu.	Cuprum
As.	Arsenic	Arsenic.	Cuj.	Cujus
At. wt.	Atomic weight	Atomic weight.	C. v.	Cras vespere
Av.	Avoirdupois	Avoirdupois.	Cyath.	Cyathus
B.	Boron	Boron.	D.	Dosis; Da
Ba.	Barium	Barium.	Decub.	Decubitus
B. a.	Balneum arenae,	Sand bath.	De d. in d.	De die in diem
Bals.	Balsamum	Balsam.	Deg.	Degree.
Be.	Beryllium	Beryllium.	De R.	Reaction of degeneration.
Bis.	Bismuth	Bismuth.	Destil.	Destilla
Bib.	Drink	Drink.	Det.	Detur
B. i. d.	Bis in die	Twice daily.	Dieb. alt.	Diebus alterius
Biol.	Biology	Biology.	" tert.	" tertius
B. m.	Balneum maris	Sea-water bath.	Dil.	Dilue
Bol.	Bolus	A large pill.	Dilut.	Dilutus
B. p.	Boiling-point	Boiling-point.	Dim.	Dimidius
Br.	Bromin	Bromin.	D. in p.	Divide in partes
Bull.	Bulliat	Let it boil.	sq.	aequales,
B. v.	Balneum vaporis	Vapor-bath.	Div.	Divide
C.	Carbon; Cubic; a Gallon (Congius); Centigrade.	Carbon; Cubic; a Gallon (Congius); Centigrade.	Ejusd.	Ejusdem
Ca.	Calcium	Calcium.	Elec.	Electuarium
Cap.	Capiat	Let him take.	Enem.	Enema
c. c.	Cubic centimeter.	Cubic centimeter.	Exhib.	Exhibeatur
Cd.	Cadmium	Cadmium.	Ext.	Extractum
Ce.	Cerium	Cerium.		
Cel.	Celsius	Celsius.		

ABBREVIATIONS.

F.	{ Fahrenheit; Fluo-	P.	{ Phosphorus, Pulse,
Far.	rinn.	Part. æq.,	Partes æquales	Pharmacopeia.
Fe.	Faradic.	Pb.	Equal parts.
Feb. dur.	Febre durante	Iron.	P. c.	Lead.
F. Ft.	Fac, Fiat	{ The fever continu-	P. C.	After meals.
Filt.	ing.	P. C.	A voir du pois
Fld.	Make.	Pd.	weight.
Flor.	Filter.	Phar.	Palladium.
F. m.	Fluid.	Pil.	Pharmacopeia.
Fol.	Flowers.	Pil.	Pill.
F. m.	Make a mixture.	Pocul.	A cup.
F. p.	Leaves.	Pond.	By weight.
F. p.	Make a potion.	Pot.	Potion; Potassa.
F. pil.	Make pills.	Ppt.	Precipitate.
Galv.	Galvanic.	P. rat.	{ In proportion to
Gl.	Glucinum.	ætat.,	Pro rata ætatis	age.
Gm.	Gram.	P. r. n.	Pro re nata	When required.
Gr.	Grain.	Pt.	Pint.
Gtt.	Drops.	Pulv.	Powder.
Guttat.	By drops.	Q. d.	Four times a day.
H.	Hydrogen.	Q. l.	According as re-
Hg.	Mercury.	Q. p.	quired.
Hl.	Hectoliter.	Q. a.	At will.
Hm.	Hectometer.	Q. s.	A sufficient quanti-
Hor.	At bed-time.	Qt.	ty.
decub.,	Hora decubitus		Q. v.	Quart.
H. s.	Hora somni	At bed-time.			As much as you
I.	Iodin.			wish.
In.	Indium.	R.	Reaumur's ther-
In d.	In die	Daily.			момeter; Respir-
Inf.	Infunde	Pour in.			ation.
Ir.	Iridium.	R.	Take.
K.	Potassium.	Rad.	Root.
K. J., K. J.	Knee-jerk.	Rb.	Rubidium.
Kl.	Kiloliter.	R. D.	Reaction of degen-
Km.	Kilometer.			eration.
L.	Liter.	Rect.	Rectified.
Li.	Lithium.	Rep.	Let it be repeated.
Liq.	Liquor.	S.	Half; Sulphur.
M.	Noon; Mix.	S.	Label.
Mac.	Macerate.	S. a.	According to art.
Mass. pil.	Massa pilularum	Pill-mass.	Sb.	Antimony.
M. ft.	Mistura fiat	Let a mixture be	Scr.	Scruple.
		made.	Sig.	Let it be labeled.
Mg.	Magnesium; Milli-	Sing.	Of each.
		gram.	Solv.	Dissolve.
Ml.	Milliliter.	Sp. gr.	Specific gravity.
Mm.	Millimeter.	Sp., or	
Mol. wt.	Molecular weight.	Spir.,	Spirit.
Muc.	Mucilage.	Sa.	Semi, semissis
Myg.	Myriagram.	Su.	Let him take.
My.	Myrialiter.	S. v.	Alcoholic spirit.
Mym.	Myriameter.	S. v. r.	Rectified spirit of
N.	Nitrogen.			wine.
Na.	Sodium.	S. v. t.	Dilute alcohol,
Ni.	Nickel.			proof-spirit.
No.	Number.	Syr.	Syrup.
Noct.	By night.	T.	Temperature.
O.	A pint.	T. d.	Three times a day.
Ol.	Oil.	Tr., Tinct.	Tincture.
O. m.	Every morning.	Ung.	Ointment.
Omn. bih.	Every two hours.	Ur.	Urine.
Omn. hor.	Every hour.	Vesic.	A blister.
Omn.	Every night.	Wt.	Weight.
noct.,	Omni nocte		℥.	Minim.
Os.	Osmium.	ʒ.	Dram.
Ov.	An egg.	ʒ.	Scruple.
Oz.	Ounce.	ʒ.	Ounce.

ABBREVIATIONS USED IN OPHTHALMOLOGY.

Acc.	Accommodation.	B. U.	Base (of prism) up.
Ah.	Hyperopic Astigmatism.	Cyl.	Cylinder; Cylindric Lens.
Am.	Myopic Astigmatism.	D.	Diopter.
As.	Astigmatism.	E.	Emmetropia; Emmetropic.
Ax.	Axis.	F.	Formula.
B. D.	Base (of prism) down.	H.	Hyperopia; Hyperopic; Horizontal.
B. I.	" " in.	L. E.	Left Eye.
B. O.	" " out.	M.	Myopia; Myopic.

ABBREVIATIONS.

ix

mm.	Millimeter.
O. D.	Oculus dexter—Right Eye.
O. S.	Oculus sinister—Left Eye.
P. p.	Punctum proximum—Near Point.
P. r.	Punctum remotum—Far Point.
R. E.	Right Eye.
Sph.	Spheric; Spheric Lens.

Sym.	Symmetric.
V.	Vision; Visual Acuity; Vertical.
+, —, =	Plus; Minus; Equal to.
∞	Infinity, 20 ft. distance.
o	Combined with.
°	Degree.

ABBREVIATIONS USED IN OTOTOLOGY.

a.	{ Applied to or in contact with Au- ricle.
A. C.	Air-conduction.
A. D.	Auris dextra—Right Ear.
A. S.	Auris sinistra—Left Ear.
B. C.	Bone-conduction.
c.	Contact.
d.	Dentes—applied to Teeth.
F.	Tuning-fork.
gl.	Glabella—applied to Forehead.
h.	Hearing Power.
L.	Left Ear.
m.	Applied to Mastoid.
Meat. Aud.	{ External Auditory Meatus.
Ext.; M. E.	
Meat. Aud.	{ Internal Auditory Meatus.
Int.; M. I.	
M. flac.	{ Membrana flaccida; Shrapnell's Membrane.
M. T.; Mt.	Membrana tympani.
O.	{ Complete Lack of Perception of Sound.
ot. ext. ac.	Otitis externa acuta.

ot. ext. chron.	Otitis externa chronica.
ot. ext. diff.	Otitis externa diffusa.
ot. med. sup.	{ Otitis media suppurativa acuta.
ac.	
ot. med. sup.	{ Otitis media suppurativa chronica.
chron.	
Pol.	Pollitzer's.
Pol. Ac.	Pollitzer's Acoumeter.
R.	Right Ear.
S.	Susurrus—a Whisper.
t.	Applied to Temple.
Tymp.	Tympanum.
ub.	Ubique—when applied at all points.
v.	Voice.
V.	Applied to Vertex.
vib.	Vibration.
z.	Applied to Zygoma.
'	Foot.
"	Inches.
∞	Heard, but Not Understood.
+R.	Rinné's Test Positive.
-R.	Rinné's Test Negative.

ABBREVIATIONS USED IN ELECTROTHERAPEUTICS.

A., or An.	Anode.
Amp.	Ampère.
A. C.	Anodal Closing.
A. C. C.	Anodal Closure Contraction.
A. C. O.	Anodal Closing Odor.
A. C. P.	Anodal Closing Picture.
A. C. S.	Anodal Closing Sound.
A. D.	Anodal Duration.
A. D. C.	Anodal Duration Contraction.
A. M.	Ampère-meter.
A. O.	Anodal Opening.
A. O. C.	Anodal Opening Contraction.
A. O. O.	Anodal Opening Odor.
A. O. P.	Anodal Opening Picture.
A. O. S.	Anodal Opening Sound.
B.	Magnetic Induction.
B. A. U.	British Association Unit.
C.	Centigrade; Current; Cathode.
C. C.	Cathodal Closure.
C. C. C.	Cathodal Closure Contraction.
C. C. C."	Various Degrees of Contraction.
C. C. T.	Cathodal Closing Tetanus.
C. G. S. Units	Centimeter-gram-second Units.
C. O.	Cathodal Opening.
C. O. C.	Cathodal Opening Contraction.
C. S.	Current-strength.
D.	Duration; Density.
De. R.	Reaction of Degeneration.
E.	Earth; Electromotive Force.
E. M. D. P.	{ Electromotive Difference of Poten- tial.
E. M. F.	Electromotive Force.
F. M.	Field Magnet.
H.	{ Horizontal Intensity of the Earth's Magnetism; One Unit of Self- induction.
H	Intensity of Magnetic Force.
I.	Intensity of Magnetism.
J.	Joule.
K.	Electrostatic Capacity.
K.	Kathode.

Kl.	Klang (sound).
K. C.	Kathodal Closing.
K. C. C.	Kathodal Closing Contraction.
K. C. T.	Kathodal Closing Tetanus.
K. D.	{ Kathodal Duration (or Period of Closure of Circuit).
K. D. C.	Kathodal Duration Contraction.
K. D. T.	Kathodal Duration Tetanus.
K. W.	Kilo-watt.
L.	{ Inductance (Coefficient of); Length.
M.	Strength of Pole.
Ma.	Milliampère.
Mfd.	Microfarad.
O.	Opening of Circuit.
P. D.	Potential Difference.
Q.	Electric Quantity.
R.	Ohmic Resistance.
T.	Time.
Te.	Tetanic Contraction.
U.	Unit.
V.	Volume; Velocity.
v.	Volt.
V. A.	Voltaic Alternative.
V. M.	Volt-meter.
W.	Work; Weight; Watt.
Z.	Contraction (Zuckung).
Z. Z. Z."	{ Increasing Strengths of Contrac- tion.
κ	Magnetic Susceptibility.
μ	Magnetic Permeability.
Ω	Ohm.
ρ	Specific Resistance.
o	Megohm (one-millionth part of an ohm).
- -	Battery.
+	Anode or Positive Pole.
-	Kathode or Negative Pole.
>	Greater than. $a > b < 4$.
<	Less than.

PREFIXES AND SUFFIXES USED IN CHEMISTRY.

Am- indicates the group NH_2 .
Azo-, **diazo-**, and **hydrazo-** indicate compounds in which nitrogen atoms are linked in various ways.
Di- is applied as a prefix to signify *two*.
Im- indicates the group NH .
Ket- indicates the molecule CO in certain structural relations.
Mon- is employed as a prefix to signify *one*.
Nitro- indicates the group NO_2 .
Pent- is applied as a prefix to signify *five*.
per- denotes in a rather vague sense an indefinitely large amount of the body to which it is prefixed, or to which it is referred.
Sequi- indicates the proportion of *two to three*.
Sub- is employed in a rather vague sense to indicate deficiency of the body to which it is prefixed.
Tetr- is applied as a prefix to signify *four*.
Thio- indicates sulphur, especially replacing oxygen.
Tri- (sometimes "*ter-*") is applied as a prefix to signify *three*.
-al indicates aldehydic structure.
-an is applied to a class of bodies related to the starch and sugar group.
-ane indicates a saturated hydrocarbon.
-ase indicates an enzyme, or nonorganized ferment, *e. g.*, diastase. This termination is at present restricted generally to enzymes of vegetable origin, but it should also be used with animal enzymes—which, however, usually end in "*in*." It would thus be better to say *pepsase* and *trypcase*, rather than *pepsin* and *trypsin*.
-ate. A suffix to nouns in chemistry signifying any salt formed by an acid acting on a base; *e. g.*, *sulphate*, *phosphate*.
-ic denotes the higher of two valencies assumed by an element, and incidentally in many cases a larger amount of oxygen.

-in is of no precise significance, and is mostly applied to bodies the structure of which is not yet known.
-yl, **-ene**, **-enyl**, and **-ine** indicate hydrocarbons. According to the American system of orthography, the only case in which "*ine*" is used is as a termination for a series of hydrocarbons, beginning with Ethine, C_2H_2 . English writers and some American chemists use it to signify basic properties, regarding **-in** as the proper term for nonbasic bodies. They thus distinguish between salicin, which forms no salts with acids, and morphin (which under such system is spelled "morphine"), which does. It must be noted that such methods are not in accordance with the tendency of modern chemie nomenclature, which seeks to express structure, not properties. The organic bases or alkaloids are not all of the same type, and when their structure is elucidated systematic names will be found for them. Until then there is no particular gain in indicating them as a group.
-id is similar to "*in*."
-ol indicates alcoholic structure, *i. e.*, presence of the group HO (hydroxyl), *e. g.*, alcohol, glycerol, phenol.
-one is applied to bodies related to the starches and sugars. It is, however, not used with this significance in "peptone," which word is not formed according to any established system.
-ose indicates a carbohydrate, *e. g.*, glucose, although it is also occasionally applied to the results of digestion of proteids, *e. g.*, albumose.
-ous denotes the lower of two degrees of valency assumed by an element and incidentally indicates, in many cases, a small amount of oxygen.
The word *sulphonic* indicates the group HSO_3 .

SUNDRY ABBREVIATIONS.

Am. American.
Arab. Arabic.
A. S. Anglo-Saxon.
Beng. Bengalese.
B. P., Br. P. British Pharmacopeia.
B. Ph., Br. Ph. "
Bret. Breton.
Celt. Celtic.
Comp. Comparative.
D., Du. Dutch.
Dim. Diminutive.
Elec. Electricity.
Eng. English.
F., Fem. Feminine.
Fr. French.
Gael. Gaelic.
Ger. German.
G. Ph. German Pharmacopeia.
Gr. Greek.
Heb. Hebrew.
Icel. Icelandic.
It. Italian.
Jav. Javanese.
L., or Lat. Latin.

Masc. Masculine.
M.E. Middle English.
Med. Medicine; Medical.
Neg. Negative.
N. F. National Formulary.
Obs. Obsolete.
O. D., O. Du. Old Dutch.
O. Fr. Old French.
O. L. G. Old Low German.
Ophth. Ophthalmology; Ophthalmologic.
Pers. Persian.
Peruv. Peruvian.
Pg. Portuguese.
pl. Plural.
priv. Privative.
Sax. Saxon.
Scand. Scandinavian.
Skt. Sanskrit.
Sp. Spanish.
Superl. Superlative.
Surg. Surgery; Surgical.
Teut. Teutonic.
Unof. Unofficial.
U. S. Ph., U. S. P. United States Pharmacopeia.

MATHEMATIC MEMORANDA.

To reduce grams to grains, multiply by 15.432. To reduce grains to grams, multiply by 0.0648. To reduce kilograms to pounds, multiply by 2.2046. To reduce ounces to grams, multiply by 28.349. To reduce inches to meters, multiply by 0.0254. To reduce inches to centimeters, multiply by 2.540. To reduce centimeters to inches, multiply by 0.3937. To reduce pints to cubic centimeters, multiply by 567.936. To reduce liters to gallons, multiply by 0.22. To reduce gallons to liters, multiply by 4.548. 1 grain = 0.064799 gram. 1 gram = 15.43235 grains. 1 millimeter = 0.03937 inch. 1 liter = 1.76077 pints. 1 minim = 0.91 grain of water. 1 line = $\frac{1}{16}$ inch. 1 μ = one-thousandth of a millimeter, 1 micron, or 1 micromillimeter, or 0.001 mm. 1 oz. (*Avoirdupois*) = 28.34954 grams. 1 oz. (*Troy*) = 31.10349 grams. 1 cc. of water at 4° C. = 1 gram. 30 in. (barometer) = 761.986 millimeters.

VALUES OF LETTERS USED IN PRONUNCIATION.

In giving the pronunciation, the word has been divided into syllables according to the following plan: When a vowel is long and is followed by a consonant, the latter is considered as belonging to the succeeding syllable; *e. g.*, pet-ro'-le-um. When the vowel is short the consonant is considered as of the syllable containing the vowel; *e. g.*, man-ip-u-la'-shun.

The vowels have then the following values: a, e, i, o, u, when immediately followed by a hyphen or at the end of a word, are long—a = *ay* in may; e = *ee* in feel; i = *i* in mine; o = *ow* in flow; u = *ew* in new. When followed by a consonant and not marked with a - above, a = *a* in cat; e = *e* in met; i = *i* in sit; o = *o* in not; u = *u* in but; oo = *oo* in boot. Consonants: ch = *ch* in cherry; g = *g* in get; j = *j* in joy; n(g) = *ng* in gong.

INDEX OF TABLES.

	PAGE
Acids,	24
Anesthetics,	55
Arteries,	79
Bacteria,	103
Baths,	137
Batteries, Electric,	139
Breath-sounds,	154
Canals,	164
Convolutions of Brain,	205
Craniometric Points,	210
Diseases, Eponymic,	230
Elements, Chemic,	252
Exanthemata,	266
Fissures,	276
Foramina,	278
Ganglia,	286
Laws,	355
Leukomains,	358
Ligaments,	360
Lines or Lineæ,	362
Membranes,	383
Metric System,	391
Monstrosities,	399
Murmurs,	405, 406
Muscles,	407
Nerves,	432
Nuclei,	449
Operations, Eponymic,	456
Pelves,	482
Points,	506
Poisons,	507
Positions,	515
Pregnancy, Duration of,	519
Processes,	521
Ptomains,	531
Rales,	543
Reflexes,	547
Respiration,	552
Signs and Symptoms of Diseases,	576
Sinuses,	583
Sutures,	608
Triangles,	631
Tumors,	637
Units,	642
Veins,	650
Weights and Measures,	659

Appendix :—

Mineral Springs of the United States,	669
---	-----

THE NEW STUDENTS' MEDICAL DICTIONARY.

A

A [*ā, av, or au*, without]. 1. The Greek letter *alpha*, called alpha privative, equivalent to the prefix *un* or *in*. It denotes absence or want of the thing or quality expressed by the root of the word. 2. Symbol of *Argon*.

Aa [*āvā*, of each]. An abbreviation, written *āā*, used in prescriptions to denote repetition of the same quantity for each item.

Ab [*ab*, from]. A Latin preposition signifying *from*.

Abaca (*ab'-ak-ah*, Sp. pron. *ah-vah-kah'*). Manilla hemp. See *Hemp*. Also *Musa textilis*, the plant which produces it.

Abadie's Sign. See *Signs and Symptoms, Table of*.

Abaptiston (*ah-bap-tis'-ton*) [*ā* priv.; *βάντιστος*, immersed]. A trephine so shaped that penetration of the brain is impossible.

Abarthrosis (*ab-ar-thro'-sis*) [*ab*, from; *arthrosis*, a joint]. Same as *Diarthrosis*, or *Abarticulation*.

Abarticular (*ab-ar-tik'-u-lar*) [*ab*, from; *articulus*, joint]. Not connected with or not situated near a joint.

Abarticulation (*ab-ar-tik-u-la'-shun*) [*ab*, from; *articulatio*, joint]. 1. Same as *Diarthrosis*; sometimes also a synonym of *Synarthrosis*. 2. A luxation.

Abasia (*ah-ba'-ze-ah*) [*ā* priv.; *βάσις*, a step]. Motor incoordination in walking. See *Astasia*.

Abasic (*ah-ba'-sik*) [*ā* priv.; *βάσις*, a step]. Pertaining to, or affected with, abasia.

Abattoir (*ah-bat-war'*) [Fr.]. A slaughterhouse or establishment for the killing and dressing of animals.

Abaxial (*ab-ak'-se-al*) [*ab*, from; *axis*, an axle]. Not situated in the line of the axis.

Abbé's Catgut Rings, rings composed of eight or ten turns of heavy catgut in the shape

of an oval, with inside diameter of two inches, for use in intestinal anastomosis. **A. Condenser**. See *Illuminator*, *Abbé*. **A. Illuminator**. See *Illuminator*.

Abdomen (*ab-do'-men*) [*abdere*, to hide]. The large inferior cavity of the trunk, extending from the pelvic cavity to the diaphragm, and bounded in front and at the sides by the lower ribs and abdominal muscles, and behind by the vertebral column, the psoas and the quadratus lumborum muscles. It is artificially divided into nine regions by two circular lines, the upper parallel with the cartilages of the ninth ribs, the lower with the iliac crests, and by two lines from the cartilages of the eighth rib to the center of Poupart's ligament. The regions thus formed are, above, the right hypochondriac, the epigastric, and the left hypochondriac; in the middle, the right lumbar, umbilical, and left lumbar; and below, the right inguinal, the hypogastric, and the left inguinal. **A., Pendulous**, a relaxed condition of the abdominal walls in which the latter hang down over the pubis.

Abdominal (*ab-dom'-in-al*) [*abdere*, to hide]. Pertaining to or connected with the abdomen.

A. Aneurysm. See *Aneurysm*. **A. Aorta**, the part of the aorta below the diaphragm.

A. Aponeurosis. See *Aponeurosis*.

A. Bandage. See *A. Binder*.

A. Binder, a broad bandage of muslin or flannel applied to the abdomen for making pressure after delivery or after an operation. Sometimes a many-tailed bandage is used.

A. Brain, the solar plexus.

A. Breathing. See *A. Respiration*.

A. Compress, a form of local pack, made by forming folds of a coarse linen towel of sufficient breadth to reach from the ensiform cartilage to the pubis; one of the folds is then wrung out of cold water, applied,

and the remainder is rolled around the body so as to retain it in position. **A. Dropsy**, ascites. **A. Gestation**. See *Pregnancy*, *Extra-uterine*. **A. Hysteria**, an hysterical condition simulating peritonitis, in which the abdomen becomes extremely painful to the touch, swollen, and distended with gas. **A. Line**, the *linea alba*. **A. Muscles**, the internal and external obliques, the transversalis, rectus, pyramidalis, and quadratus lumborum. **A. Press**. See *Prelum Abdominale*. **A. Reflex**. See *Reflexes*, *Table of*. **A. Regions**. See *Abdomen*. **A. Respiration**, R. carried on chiefly by the diaphragm and abdominal muscles. **A. Ring, External**, a triangular opening in the fibers of the aponeurosis of the external oblique muscle transmitting the spermatic cord of the male and the round ligament of the female. **A. Ring, Internal**, an oval aperture in the fascia transversalis that transmits the spermatic cord of the male and the round ligament of the female. **A. Section**. See *Celiotomy*. **A. Surgery**, the branch of surgery that deals with the lesions of the abdominal viscera and the operations performed upon them through incisions in the abdominal walls. **A. Typhus**, Enteric Fever.

Abdomino-anterior (*ab-dom' in-o-an-te'-re-or*). Having the belly forward (used of the fetus in utero).

Abdomino-genital (*ab-dom' in-o-jen'-it-al*). Relating to the abdomen and the genitalia. **A. Nerve, Inferior**, the ilio-inguinal nerve. **A. Nerve, Superior**, the ilio-hypogastric nerve.

Abdomino-posterior (*ab-dom' in-o-pos-te'-re-or*). Having the belly toward the mother's back (used of the fetus in utero).

Abdomino-scrotal (*ab-dom' in-o-skro'-tal*). Relating to the abdomen and the scrotum. **A. Muscle**, the cremaster muscle.

Abdomino-thoracic (*ab-dom' in-o-tho-ras'-ik*). Relating to the abdomen and thorax.

Abdomino-vesical (*ab-dom' in-o-ves'-ik-al*). Relating to the abdomen and the urinary bladder. **A. Pouch**, a fold of the peritoneum in which are comprised the urachal fossae.

Abducens (*ab-du'-sens*) [L., "leading away"]. A term applied to certain muscles, or their nerves, that draw the related part from the median line of the body. Also, the sixth pair of nerves supplying the external recti of the eyes. **A. oculi**, the external rectus muscle of the eye.

Abducent (*ab-du'-sent*). Abducting.

Abduct (*ab-duk't*) [*abducere*]. To draw away from the median line, as to abduct a limb.

Abduction (*ab-duk'-shun*) [*ab*, from; *ducere*, to lead]. 1. The withdrawal of a part from

the axis of the body. 2. The recession or separation from each other of the parts of a fractured bone.

Abductor (*ab-duk'-tor*). Same as *Abducens*.

A. Auris. See *Muscles*, *Table of*.

Aberrant (*ab-er'-ant*) [*ab*, from; *errare*, to wander]. Deviating from the normal or regular type, in appearance, structure, course, etc., as the aberrant duct of the testis or liver, aberrant arteries, etc.

Aberration (*ab-er'-a'-shun*) [*ab*, from; *errare*, to wander]. Deviation from the normal; mental derangement; fetal malformation; vicarious menstruation; escape of the fluids of the body by an unnatural channel. In optics, any imperfection of focalization or refraction of a lens. **A., Chromatic**, the dispersion arising from unequal refraction of light of different parts of the spectrum. The violet rays, being more refrangible than the red rays, are brought to a focus nearer the lens, and the image is surrounded by a halo of colors. **A., Mental**, a degree of paranoia that may or may not amount to insanity. **A., Spheric**, the excess of refraction of the peripheral part of a convex lens over the central part, producing an imperfect focus and a blurred image.

Abies (*a'-be-ēs*) [L.]. A genus of coniferous plants, including the fir, hemlock, and spruce. **A. balsamea**, Silver Fir, Balsam Fir, or Balm of Gilead, a tree of the nat. ord. *Conifera*, from which is derived the Terebinthina Canadensis. **A. canadensis**, Hemlock Spruce; bark of the Canadian Fir-tree. It is used as an astringent in various local and internal conditions. It yields Pix Canadensis. **A. excelsa**, Norway Spruce. It yields Pix Burgundica. **A. pectinata**, the European Silver Fir. Its buds are resinous, balsamic, and sudorific. Unof. **A. Preparations**, Extractum Abietis Canadensis Liquidum (B. P.). Dose m xv-lx (1-4). Glycerinum Abietis (B. P.). Dose ʒ j-ij (4.0-8.0). Geddes' Fluid-Extract of Hemlock Bark. Dose m v-x (0.3-0.6).

Abietene (*a-bi'-et-en*), $C_{11}H_{18}$. A hydrocarbon, obtained from the *Pinus sabiniana*, a California nut pine. It is an aromatic, volatile liquid, agreeing in composition with normal heptane. It is extensively used as a popular remedy; called, also, *Erasene*. Unof.

Abietic, **Abietinic** (*ab-i-et'-ik*, *ab-i-et-in'-ik*) [*Abies*]. Pertaining to the genus *Abies*, as *Abietic Acid*, $C_{14}H_{10}O_2$ or $C_{20}H_{30}O_2$, occurring in the resin of *Abies excelsa* and *Larix europea*.

Abietin (*a-bi'-et-in*) [*Abies*]. A resinous principle obtained from the turpentine of various species of pine and fir. **A. Anhydrid**, $C_{44}H_{62}O_2$, the main constituent of *Resin*.

Abietite (*ab'-i-et-it*), $C_4H_8O_3$. A sugar re-

- sembling mannite, found in the needles of the European Silver Fir, *Abies pectinata*.
- Abiogenesis** (*ab-i-o-jen'-es-is*) [*â* priv.; *βίος*, life, and *genesis*]. The (theoretic) production of living by non-living matter. The older term was *Spontaneous Generation*.
- Abirritant** (*ab-ir'-it-ant*) [*ab*, from; *irritare*, to irritate]. 1. Tending to diminish irritation; soothing. 2. Relating to diminished sensitiveness. 3. A remedy or agent that allays irritation.
- Abirritation** (*ab-ir-it-a'-shun*) [*ab*, from; *irritare*, to irritate]. Diminished tissue-irritability; atony or asthenia.
- Ablactation** (*ab-lak-ta'-shun*) [*ab*, from; *lactare*, to give suck]. The weaning of a child. The end of the suckling period.
- Ablate** (*ab-late'*) [*ab*, from; *latum*, from *ferre*, to bear]. To remove, to cut off.
- Ablation** (*ab-la'-shun*) [*ab*, from; *latum*, from *ferre*, to bear]. Removal of a part, as a tumor, by amputation, excision, etc.
- Ablatio retinæ** (*ab-la'-she-o ret-in'-e*). Detachment of the retina.
- Ablepharia** (*ah-blef-a'-re-ah*) [*â* priv.; *βλέφαρον*, the eyelid]. The condition of the absence of the eyelids.
- Ablepharon** (*ah-bleff-ar-on*) [*â* priv.; *βλέφαρον*, the eyelid]. Absence of the eyelids.
- Ablepharous** (*ah-bleff-ar-us*) [*â* priv.; *βλέφαρον*, the eyelid]. Without eyelids.
- Ablepsia** (*ah-blep'-se-ah*) [*â* *βλεψία*, without sight]. 1. Blindness. 2. Dulness of perception.
- Abluent** (*ab'-lu-ent*) [*abluerē*, to wash away]. Detergent. That which cleanses or washes away.
- Ablution** (*ab-lu'-shun*) [*abluerē*, to wash away]. Washing or cleansing the body. Separation of chemic impurities by washing.
- Abnormal** (*ab-nor'-mal*) [*ab*, away from; *norma*, a law]. Not normal; not conformable with nature or with the general rule.
- Abnormality** (*ab-nor-mal'-it-e*) [*ab*, away from; *norma*, a law]. The quality of being abnormal; a deformity or malformation.
- Abnormity** (*ab-nor'-mit-e*). Same as *Abnormality*.
- Abomasum** (*ab-o-ma'-sum*) [*ab*, away; *omasum*, paunch]. The reed or proper digestive stomach of ruminating mammals. Also called "fourth," or "true," stomach.
- Aboral** (*ab-o'-ral*) [*ab*, away from; *os*, the mouth]. Opposite to, or remote from, the mouth.
- Abort** (*ab-ort'*) [*ab*, from; *ortus*, from *oriri*, to grow]. 1. To miscarry; to expel the fetus before it is viable. 2. To prevent the full development, as of a disease. 3. To come short of full development.
- Abortient** (*ab-ort'-shent*) [*ab*, from; *oriri*, to grow]. Abortive; abortifacient.
- Abortifacient** (*ab-ort-te-fa'-shent*) [*abortus*; *facere*, to make]. 1. Causing abortion. 2. A drug or agent inducing the expulsion of the fetus.
- Abortion** (*ab-ort'-shun*) [*abortus*, a miscarriage]. The expulsion of the ovum before the child is viable, that is, occurring any time before the end of the sixth month. By some authors expulsion of the ovum during the first three months is termed *abortion*; from this time to viability it is termed *immature delivery*, or *miscarriage*, and from the period of viability to that of maturity, *premature delivery*. **A.**, **Accidental**. See **A.**, **Spontaneous**. **A.**, **Artificial**, that produced intentionally. **A.**, **Criminal**, when not demanded for therapeutic reasons. **A.**, **Habitual**, repeated **A.** in successive pregnancies, usually due to syphilis. **A.**, **Incomplete**, when the membranes or the placenta is retained. **A.**, **Induced**. See **A.**, **Artificial**. **A.**, **Inevitable**, when the embryo or fetus is dead, or when there is an extensive detachment or rupture of the ovum. **A.**, **Missed**, the death of the fetus and not followed within two weeks by its expulsion. **A.**, **Spontaneous**, that not induced by artificial means.
- Abortionist** (*ab-ort'-shun-ist*) [*abortus*, a miscarriage]. One who criminally produces abortions; especially one who follows the business of producing abortions.
- Abortive** (*ab-ort'-tiv*) [*abortus*, a miscarriage]. Prematurely born; coming to an untimely end; incompletely developed; cutting short the course of a disease; abortifacient.
- Abortus** (*ab-ort'-tus*) [*L.*]. An aborted fetus; abortion.
- Aboulia** (*ah-boo'-le-ah*). See *Abulia*.
- Aboulomania** (*ah-boo lo-ma'-ne-ah*). See *Abulomania*.
- Abrachia** (*ah-bra'-ke-ah*) [*â* priv.; *βραχίον*, arm]. The condition of an armless monster.
- Abrachiocephalia** (*ah-bra-ke-o-sef-a'-le-ah*) [*â* priv.; *βραχίον*, arm; *κεφαλή*, head]. Absence of the head and arms.
- Abrachius** (*ah-bra'-ke-us*). A monster without arms. See *Abrachia*.
- Abrasio** (*ab-ra'-ze-o*) [*L.*]. An abrasion. **A.** *corneæ*, a scraping off of the superficial epithelium of the cornea.
- Abrasion** (*ab-ra'-zhun*) [*ab*, from; *radere*, to rub]. The rubbing off of the cutaneous or mucous surface by an injury.
- Abrin** (*a'-brin*). The chemic ferment or poisonous principle of jequirity. It has been employed in the study of immunity.
- Abrotanum** (*ab-rot'-an-um*) [*ἀβρότονον*, an aromatic plant]. The plant called Southern Wood, *Artemisia abrotanum*.
- Abrus** (*a'-brus*) [*ἀβρός*, pretty]. Jequirity; Indian Licorice. The seeds of *A. precatorius*, or Wild Licorice. Its properties are thought

to be due to the presence of certain ferments. See *Abrin*. Infusions applied to the conjunctiva or to any mucous surface induce violent purulent inflammation with growth of false membrane. It is used in producing artificial conjunctivitis.

Abscess (*ab'-ses*) [*abscessus*, a departure or separation]. A localized collection of pus surrounded by a wall of lymph. According to location, abscesses are named *Dorsal*, *Iliac*, *Mammary*, *Ischio-rectal*, *Perityphlitic*, *Retro-pharyngeal*, *Urethral*, etc. **A.**, *Alveolar*, abscess in the gum or alveolus. **A.**, *Atheromatous*, an area of softening in the wall of a vessel the result of sclerotic endarteritis. **A.**, *Bursal*, abscess in the bursæ. **A.**, *Canalicular*, mammary abscess that communicates with a milk duct. **A.**, *Chronic*, or *Cold Abscess*, one of slow and apparently non-inflammatory development, generally about a bone, joint, or gland. It is usually tuberculous and contains cheesy material. **A.**, *Cold*. See *A. Chronic*. **A.**, *Congestive*, the pus appears at a point distant from where it is formed. **A.**, *Embolic*, formed at the seat of a septic embolus. **A.**, *Metastatic*. See *A. Embolic*. **A.**, *Miliary*, a small embolic abscess. **A.**, *Primary*, one formed at the seat of pyogenic infection. **A.**, *Psoas*, one arising from disease of the lumbar or lower dorsal vertebrae, the pus descending in the sheath of the psoas muscle, and usually pointing beneath Poupart's ligament. **A.**, *Pyemic*. See *Pyemia*. **A.**, *Residual*, one formed in or about the residues of former inflammation. **A.**, *Tuberculous*. Same as *A.*, *Chronic*.

Abscissæ (*ab-sis'-se*) [*ab*, away; *scindere*, to cut]. The transverse lines cutting vertical ones at right angles, to show by a diagram the relations of two series of facts, as, *e. g.*, the number of pulse-beats, or the temperature record in given periods of time.

Abscission (*ab-sish'-un*) [*ab*, from; *scindere*, to cut]. Removal of a part by cutting.

Absinthe (*ab'sinth*). See *Absinthium*.

Absinthin (*ab-sin'-thin*) [*absinthium*]. A bitter crystalline principle obtainable from wormwood. See *Absinthium*.

Absinthism (*ab-sinth'-izm*). A disease similar to alcoholism, the result of the excessive use of absinthe. It is characterized by general muscular debility and mental disturbances, that may proceed to convulsions, acute mania, or general paralysis.

Absinthium (*ab-sinth'-e-um*) [L.]. Wormwood. The leaves and tops of *Artemisia absinthium*. **A.** contains a volatile oil and an intensely bitter principle, *Absinthin*, $C_{10}H_{16}O_4$, which is a narcotic poison. **A.** increases cardiac action, produces tremor and epileptiform convulsions. Dose gr. 2x-xi (1.3-2.6),

in infusion. It is used as a stomachic tonic. *Absinthe*, a French liquor, is an alcoholic solution of the oil exhibited with oils of anise, marjoram, and other aromatic oils.

Absinthol (*ab-sinth'-ol*), $C_{10}H_{16}O$. The principal constituent of oil of wormwood; it is isomeric with ordinary camphor.

Absorb (*ab-sorb'*) [*ab*, from; *sorbere*, to suck up]. To suck up or imbibe; to take within one's self.

Absorbent (*ab-sor'-bent*) [*absorbere*, to suck in]. 1. Absorbing, capable of absorbing. 2. An organ or part that absorbs. 3. A term applied to the *Lacteals* and *Lymphatics*.

4. In materia medica, a drug or medicine that produces absorption of diseased tissue.

A. Glands. See *Lymphatics*. **A. System**, the lacteals and lymphatics, with their associated glands.

Absorptiometer (*ab-sorp-te-om'-et-er*) [*absorption*; *μετρον*, measure]. A device for measuring the thickness of the layer of liquid that is taken up between two glass plates by capillary attraction. Used in conjunction with a spectro-photometer, it serves as a hematoscope.

Absorption (*ab-sorp'-shun*) [*absorbere*, to suck in]. The permeation or imbibition of one body by another. **A. Lines or Bands**, dark lines of the spectrum, called Fraunhofer's lines, caused by the arrest or absorption of the ethereal waves of certain lengths and rapidities, mainly by vapors of the sun's atmosphere. **A.**, *Interstitial*, the removal by the absorbent system of effete matters.

Absorptive (*ab-sorp'-tiv*) [*absorbere*, to suck in]. Having the power or function of absorbing.

Abstergent (*ab-ster'-jent*) [*abs*, from; *tergere*, to cleanse]. Cleansing, detergent. See *Detergent*.

Abstersive (*ab-ster'-siv*) [*abstersivus*]. Abstergent.

Abstract (*ab'-strakt*) [*abstrahere*, to draw away]. In pharmacy, a solid preparation in which two parts of the drug are represented by one part of the abstract (which is compounded with milk-sugar). Abstracts are double the strength of the fluid extracts.

Abterminal (*ab-ter'-min-al*) [*ab*, from; *terminus*, end]. Passing from tendinous into muscular tissue (used of electric currents).

Abulia (*ah-bu'-le-ah*) [*â priv.*; *βούλη*, will]. Loss or defect of will-power.

Abulic (*ah-bu'-lik*) [*â priv.*; *βούλη*, will]. Characterized by or affected with abulia.

Abulomania (*ah-bu-lo-ma'-ne-ah*) [*â priv.*; *βούλη*, will; *μανία*, madness]. A disease of the mind characterized by imperfect or lost will-power.

Acacia (*ah-ka'-she-ah*) [L.]. 1. A large genus of leguminous trees, shrubs, and

herbs, many of them Australian or African. A number of the species are medicinal, and some are poisonous. The bark is usually very astringent. Gum arabic is produced by various species. 2. *A. anthelmintica*. See *Mussanin*. *A. catechu*. See *Catechu*. *A. lebbek*, *A. nilotica*, *A. vera*, and *A. vereck*, are among the species that furnish gum arabic. 3. Gum Arabic. A nearly white, transparent gum, exuding from *Acacia Senegal*. Soluble in water. It is used in the manufacture of mucilage, and contains *Arabin*, $C_{12}H_{22}O_{11}$, identical in composition with cane sugar. **A.**, Mucilago, acacia 34, water, to make 100 parts; incompatible with alcoholic tinctures. **A.**, Syrup., mucilage 25, syrup. simp. 75. It is used in various mixtures as a demulcent and to suspend insoluble powders.

Acalypha (*ak-kal'-if-ah*) [*ἀκαλύφης*, unveiled]. A genus of euphorbiaceous plants. **A.**, Ext. Liq. Dose π x-3j (0.6-4.0). *A. fruticosa*, of India, is useful in dyspepsia and diarrhea, and is tonic and alterant. *A. hispida* has similar uses. *A. indica*, a euphorbiaceous plant, common in India. The leaves are expectorant, emetic, laxative. **A.**, Succus. Dose for an infant 3j (4.0). Unof. *A. virginica*, of North America, is diuretic and expectorant.

Acanthia lectularia (*ak-an'-the-ah lek-chu-la'-re-ah*) [L.]. The common bedbug.

Acanthial (*ak-an'-the-al*) [*ἀκάνθιον*, a little thorn]. Pertaining to the acanthion.

Acanthion (*ak-an'-the-on*) [*ἀκάνθιον*, a little thorn]. A point at the base of the nasal spine.

Acanthoma (*ak-an-tho'-mah*) [*ἀκανθώμα*, a spine]. A localized excessive growth in any part of the prickle-cell layer of the skin.

Acanthosis (*ak-an-tho'-sis*) [*ἀκανθώσις*, thorn]. Any skin disease marked by abnormalities in the prickle-cell layer. **A. nigricans**, a general pigmentation of the skin, with papillary, mole-like growths; a rare condition.

Acardia (*ak-kar'-de-ah*) [*ἄ καρδία*, heart]. Congenital absence of the heart.

Acariasis (*ak-ar-i'-as-is*) [*ἄ καρδία*, to cut]. See *Mange*.

Acarid, **Acaridan** (*ak'-ar-id*, *ak-ar'-id-an*) [*ἀκαρίς*, small, tiny]. Pertaining to *Acarus*.

Acarinosis (*ak-ar-in-o'-sis*) [*acarus*, a mite]. Any disease, as the itch, produced by a mite, or acarid.

Acarodermatitis (*ak-ar-o-der-mat-i'-tis*) [*acarus*, a mite; *dermatitis*]. Dermatitis caused by acari, or mites.

Acaroid (*ak'-ar-oid*) [*Acarus*, a mite]. Mite-like. **A. Gum**, Botany Bay Gum; *Resina lutea*. An aromatic resin used in Australia as a remedy for gastric troubles, intestinal catarrhs, diarrheas, etc. Dose grs. viij-xvj, (0.5-1.0), in alcoholic solution. Unof. **A.**

Resin, Black Boy Gum. The resin of various species of *Xanthorrhoea*. Benzoic acid is prepared from it, and it is said to have the properties of storax, and balsam of Peru.

Acarus (*ak'-ar-us*) [*ἄ* priv.; *καίρειν*, to cut (because so small)]. The mite, or tick, a parasite of man and animals.

Acataphasia (*ak-kat-af-a'-ze-ah*) [*ἄ* priv.; *κατά*, after; *φάσις*, utterance]. A disorder in the syntactical arrangement of uttered speech, due to some central lesion.

Acathectic (*ak-ah-thek'-tik*) [*ἀκαθεκτός*, unrestrained]. A term applied by Liebermeister, to that form of icterus caused by pathologic changes in the liver cells through which they become unable to retain their secretion.

Acaudal, **Acaudate** (*ak-kaw'-dal*, *ak-kaw'-dat*) [*ἄ* priv.; *cauda*, a tail]. Tailless.

Acceleration (*ak-sel-er-a'-shun*) [*accelerare*, to hasten]. Quickening, as of the rate of the pulse, or of the respiration.

Accelerator (*ak-sel'-er-a-tor*) [L.]. That which accelerates. **A. Nerves**, nerves passing from the medulla to the heart and conducting stimuli that cause acceleration of the heart's action. **A. Urinæ**, a muscle of the penis the function of which is to expel the last drops in urination, to expel the semen, and to assist erection. The sphincter vaginæ is its analogue in the female.

Accentuation (*ak-sen-tu-a'-shun*) [*accentuare*]. Increased loudness or distinctness.

Accessory (*ak'-ses-o-re*, or *ak-ses'-o-re*) [*accessorius*]. A term applied to certain glands, muscles, ducts, nerves, arteries, etc., that are auxiliary in function, course, etc., to the principal. Certain small muscles, as the lumbricales, are regarded as accessory to more important muscles.

Accidental (*ak-se-dent'-al*) [*accidentalis*]. 1. Due to, or caused by, an accident. 2. Intercurrent; having no essential connection with other conditions or symptoms.

Accipiter (*ak-sip'-it-er*) [L., a hawk]. A facial bandage with tails radiating like the claws of a hawk.

Acclimatation, **Acclimation**, **Acclimatization** (*ak-kli-mat-a'-shun*, *ak-lim-a'-shun*, *ak-kli-mat-is-a'-shun*) [*ad*, to; *clima*, climate]. The process of becoming accustomed to the climate, soil, water, etc., of a country to which a plant, animal, person, or a people has removed.

Accommodation (*ak-om-o-da'-shun*) [*accommodare*, to adjust]. Adaptation or adjustment, particularly the adjustment of the eye for different distances. **A.**, **Absolute**, the accommodation of either eye separately. **A.**, **Histologic**, the occurrence of changes in the morphology and function of cells following changed conditions. **A.**, **Negative**, the condition of the eye at rest. **A. of the Eye**, that

function of the ciliary muscle and lens whereby objects at different distances are clearly seen. It depends upon the inherent elasticity of the lens, which when the ciliary muscle of an emmetropic eye is at rest, is adapted to the proper focalization of theoretically parallel rays of light. Objects nearer, to be clearly seen, require a greater refracting power on the part of the eye because the rays from such objects are more divergent. This additional refracting power is gained by an increased antero-posterior diameter of the lens, brought about by the contraction of the ciliary muscle, which occasions a loosening of the suspensory ligament and a thickening of the lens by its own elasticity. **A. Phosphenes**, the peripheral light-streak seen in the dark after the act of accommodation. **A., Range of**, the distance between the nearest point of distinct vision, and the most distant point.

Accouchement (*ak-koozh-mon(g')*) [Fr.]. The French term for childbirth. **A. Forcé**, rapid and forcible delivery with the hand.

Accoucheur (*ak-koo-shur'*) [Fr.]. A male midwife.

Accoucheuse (*ak-koo-shu(r)z'*) [Fr.]. A wife.

Accretion (*ak-re'-shun*) [*ad*, to; *erescere*, to increase]. A term denoting the manner by which crystalline and certain organic forms increase their material substance. Also, the adherence of parts normally separate.

Accumulator (*ak-u'-mu-la-tor*) [*accumulare*, to heap up]. An apparatus to store electricity.

A. C. E. Mixture. An anesthetic mixture composed of alcohol 1 part, chloroform 2 parts, ether 3 parts. See *Anesthetic*.

Acentric (*ah-sen'-trik*) [*á* priv.; *κέντρον*, center]. Not eccentric; not originating in, or pertaining to, a nerve center; peripheric.

Acephalia (*ah-sef'-al-ah*) [*á* priv.; *κεφαλή*, head]. The absence of the head.

Acephalism (*ah-sef'-al-izm*) [*a* priv.; *κεφαλή*, head]. See *Acephalia*.

Acephalobrachia (*ah-sef'-al-o-brá'-ke-ah*) [*á* priv.; *κεφαλή*, head; *βραχίον*, arm]. Absence of the head and arms.

Acephalobrachius (*ah-sef'-al-o-brá'-ke-us*) [*á*; *κεφαλή*; *βραχίον*]. A monster with neither head nor arms.

Acephalocardia (*ah-sef'-al-o-kar'-de-ah*) [*á* priv.; *κεφαλή*, head; *καρδία*, heart]. Absence of the head and heart.

Acephalocardius (*ah-sef'-al-o-kar'-de-us*) [*á*; *κεφαλή*; *καρδία*]. A monster with neither head nor heart.

Acephalochiria (*ah-sef'-al-o-ki'-re-ah*) [*á* priv.; *κεφαλή*, head; *χείρ*, hand]. Absence of the head and hands.

Acephalochirus (*ah-sef'-al-o-ki'-rus*) [see

Acephalochiria]. A monster with neither head nor hands.

Acephalocyst (*ah-sef'-al-o-sist*) [*á* priv.; *κεφαλή*, head; *κυστίς*, a bladder]. The bladder-worm. A headless, sterile hydatid, found in the liver and other organs. **A. racemosa**, the hydatid mole of the uterus.

Acephalogaster (*ah-sef'-al-o-gas'-ter*) [*á* κεφαλή, headless; *γαστήρ*, belly]. A monster with neither head nor stomach.

Acephalogasteria (*ah-sef'-al-o-gas-te'-re-ah*). Absence of the head and stomach. See *Acephalogaster*.

Acephalopodia (*ah-sef'-al-o-po'-de-ah*) [*á* priv.; *κεφαλή*, head; *πούς*, foot]. Absence of the head and feet.

Acephalopodius (*ah-sef'-al-o-po'-de-us*). A monster with neither head nor feet. See *Acephalopodia*.

Acephalorrhachia (*ah-sef'-al-or-a'-ke-ah*) [*á* priv.; *κεφαλή*, head; *ράχις*, spine]. Absence of the head and vertebral column.

Acephalostomia (*ah-sef'-al-o-sto'-me-ah*) [*á* priv.; *κεφαλή*, head; *στόμα*, mouth]. Absence of the head, with a mouth-like opening on the superior aspect.

Acephalostomus (*ah-sef'-al-o-sto'-mus*) [*á* priv.; *κεφαλή*; *στόμα*]. A monster without a head, but with a mouth-like aperture.

Acephalous (*ah-sef'-al-us*) [*á* κεφαλή, headless]. Headless.

Acephalus (*ah-sef'-al-us*) [*á* priv.; *κεφαλή*, head]. 1. A species of omphalositic monsters characterized by complete absence of the head and usually of the upper extremities. It is the commonest condition among the omphalositic. 2. A variety of the foregoing species marked by the highest form of development. The head is absent, but there is at least one superior extremity, and the thorax is generally fairly well-developed.

Acervulus, or **Acervulus cerebri** (*as-er'-vu-lus ser'-e-bri*). Concretionary matter near the base of the pineal gland, consisting of alkaline phosphates and carbonates, with amyloid matter; brain-sand.

Acescence (*as-es'-ens*) [*acescere*, to grow sour]. 1. The process of becoming sour; the quality of being somewhat sour. 2. A disease of wines, whereby they become sour owing to the agency of *Mycoderma aceti*.

Acetabular (*as-et-ab'-u-lar*) [*acetabulum*, a cup]. Pertaining to the acetabulum.

Acetabulum (*as-et-ab'-u-lum*) [*acetabulum*, a small cup]. A cup-shaped depression on the outer aspect of the innominate bone for the reception of the head of the femur.

Acetal (*as'-et-al*) [*acetum*, vinegar]. $C_2H_4O_2$. Ethidene diethylate, a colorless, liquid, with an ethereal odor, produced by the imperfect oxidation of alcohol under the influence of platinum black. It is sparingly soluble in

water; boils at $104^{\circ}\text{C}.$; sp. gr. at 20° is 0.8304. Its action is that of a soporific. Dose \mathfrak{zj} (4 o).

Acetaldehyd (*as-et-al'-de-hid*). The normal aldehyd; ethaldehyd. See *Aldehyd*.

Acetamid (*as-et'-am-id*), $\text{C}_2\text{H}_5\text{NO}$. A white crystalline solid produced by distilling ammonium acetate, or by heating ethyl acetate with strong aqueous ammonia. It combines with both acids and metals to form unstable compounds.

Acetanilid (*as-et-an'-il-id*), $\text{C}_6\text{H}_5\text{NO}$. Phenylacetamid. A white, crystalline solid, produced by boiling anilin and glacial acetic acid together for several hours, the crystalline mass being then distilled. It melts at 114° and boils at 295° . It is soluble in hot water, alcohol, and ether. Under the name *antifebrin* it is prescribed as an antipyretic. Dose gr. ij-x (0.13-0.65), not exceeding gr. xxx (2 o) in the 24 hours.

Acetate (*as'-et-at*) [*acetum*, vinegar]. Any salt of acetic acid.

Acetic (*as-e'-tik*) [*acetum*, vinegar]. Pertaining to *acetum* or vinegar; sour. See *Acid*, *Acetic*. A. Fermentation, the development of acetic acid by the activity of the *Mycoderma aceti*.

Acetin (*as'-et-in*) [*acetum*, vinegar], $\text{C}_2\text{H}_5(\text{C}_2\text{H}_5\text{O})_2$. A chemic compound formed by the union of glycerol and acetic acid.

Aceto-acetic Acid. See *Acid*, *Diacetic*.

Acetometer (*as-et-om'-et-er*) [*acetum*, vinegar; μέτρον, measure]. An instrument used in the quantitative determination of acetic acid.

Acetone (*as'-et-on*) [*acetum*, vinegar], $\text{C}_3\text{H}_6\text{O}$. Dimethyl ketone. A colorless, mobile liquid, of peculiar odor and burning taste, present in crude wood-spirit; it occurs in small quantities in the blood and in normal urine, and in considerable quantities at times in the urine of diabetic patients. It is miscible with ether, alcohol, and water.

Acetonemia (*as-et-o-ne'-me-ah*) [*acetone*; αίμα, blood]. The presence of acetone in the blood.

Acetones (*as'-et-ons*). A class of compounds that may be regarded as consisting of two alcoholic radicals united by the group CO, or as aldehyds in which hydrogen of the group COH has been replaced by an alcoholic radicle.

Acetonitril (*as-et-o-ni'-tril*), CH_3CN . Methyl cyanid. It is a colorless liquid.

Acetono-resorcin. A combination of two molecules of resorcin and one molecule of acetone, obtained by heating together 15 gm. resorcin, 100 gm. acetone, and 50 gm. concentrated hydrochloric acid. It appears as small, anhydrous, prismatic crystals, insoluble in water, alcohol, ether, or chloroform, but readily soluble in alkalis.

Acetonuria (*as-et-on-u'-re-ah*) [*acetone*; ού-

ρον, urine]. The presence of acetone in the urine.

Acetophenone (*as-et-o-fe'-nōn*), $\text{C}_6\text{H}_5(\text{CO})(\text{CH}_3)$. "Hypnone;" an hypnotic and antiseptic. It results from the action of zinc methyl upon benzoyl chlorid and crystallizes in large plates, melts at 20.5° , and boils at 202° . It is without satisfactory action. Dose $\mathfrak{m}\text{iv}$ -xv (0.26-1 o).

Acetous (*as-e'-tus*) [*acetum*, vinegar]. Resembling vinegar; pertaining to, or charged with vinegar or acetic acid.

Acetphenetidid (*as-et-fe-net'-id-in*). See *Phenacetin*.

Acet-toluide (*as-et-tol'-u-ed*), $\text{C}_7\text{H}_7\text{NH}\cdot\text{C}_2\text{H}_5\text{O}$. Aceto-orthotoluide. An antipyretic resembling acetanilid. The dose is not accurately determined.

Acetum (*as-e'-tum*) [*gen.*, *Aceti*; *pl.*, *Aceta*]. [L.]. Vinegar. An impure, dilute acetic acid produced by acetous fermentation of wine, cider, or other fruit-juice. In pharmacy, a solution of the active principles of certain drugs in dilute acetic acid.

A. aromaticum (N. F.) ["aromatic vinegar"], a mixture of alcohol, water, and acetic acid, aromatized with the oils of rosemary, lavender, juniper, peppermint, cassia, lemon, and cloves. **A. britannicum**, an aromatic vinegar consisting of glacial acetic acid 600 o, camphor 60 o, oil of cloves 2 o, oil of cinnamon 1 o, oil of lavender 0.5.

Acetyl (*as'-et-il*) [*acetum*, vinegar], $\text{C}_2\text{H}_3\text{O}$. A univalent radicle supposed to exist in acetic acid and its derivatives. Aldehyd may be regarded as the hydrid and acetic acid as the hydrate, of acetyl. **A. Chlorid**, $\text{C}_2\text{H}_3\text{OCl}$, a colorless liquid, used as a reagent. **A. Peroxid**, $(\text{C}_2\text{H}_3\text{O})_2\text{O}_2$, a thick liquid, insoluble in water, but readily dissolved by ether and alcohol. It is a powerful oxidizing agent. It is decomposed in sunlight and explodes violently when heated.

Acetylene (*as-et'-il-ēn*) [*acetum*, vinegar], C_2H_2 . A colorless gas, with a characteristic, unpleasant odor, burning with a luminous, smoky flame. It is formed by the imperfect combustion of illuminating gas and other hydrocarbons. The *acetylene series* of hydrocarbons has the general formula, C_nH_{n-2} .

Acetylphenylhydrazin (*as-et-il-fe-nil-hi'-dra-zin*), $\text{C}_6\text{H}_5\text{N}_2\text{H}_2\text{C}_2\text{H}_3\text{O}$. See *Pyrodrin*.

Ache (*āk*) [A.S., *acan*, to ache]. Any continuous or throbbing pain.

Acheilia (*ah-ki'-le-ah*) [*ā* priv.; χείλος, a lip]. The congenital absence of lips.

Acheilous (*ah-ki'-lus*) [*ā* priv.; χείλος, a lip]. Without lips.

Acheiria (*ah-ki'-re-ah*) [*ā* priv.; χείρ, a hand]. The congenital absence of hands.

Acheirous (*ah-ki'-rus*) [*ā* priv.; χείρ, a hand]. Affected with acheiria.

Achilia (*ah-ki'-le-ah*). See *Acheilia*.

Achillea (*ah-il'-e-ah*) [*Achilles*, its reputed discoverer]. Milfoil, Yarrow. The herb *A. millefolium*. Its properties are due to a bitter, aromatic, astringent, tonic extractive, *achillein*, and a volatile oil. It has long been used as a vulnerary, and has been highly recommended for intermittent and low exanthematous fevers. Dose $\frac{3}{4}$ -Oj, infusion *ad lib.*; of the extractive, $\frac{3}{4}$ -ij (3.0-12.0); of the volatile oil, gtt. v-xv (0.3-1.0). Unof. To the genus *Achillea* belong various other unofficial medicinal plants, as *A. moschata*, of the Alps, used in preparing cordials and a diaphoretic medicine, and *A. pharmica*, or sneezewort, a strong sialagogue.

Achillein (*ah-il'-e-in*). An extractive from *Achillea millefolium*.

Achilles Tendon (*ah-il'-ez ten'-don*). The tendon of the gastrocnemius and soleus muscles, inserted into the back of the heel.

Achillodynia (*ah-il'-o-din'-e-ah*) [*Achilles*; *ὀδύνη*, pain]. Pain referred to the insertion of the tendo Achillis.

Achlorhydria (*a-klor-hi'-dri-a*) [*ἄ* priv.; *χλωρός*, green; *ὕδωρ*, water]. Absence of free hydrochloric acid from the gastric juice.

Achloropsia (*ah-klo-rop'-se-ah*) [*ἄ* priv.; *χλωρός*, green; *ὄψις*, vision]. Green-blindness.

Acholia (*ah-ko'-le-ah*) [*ἄ* priv.; *χολή*, bile]. Absence of biliary secretion.

Achorion (*a-ko'-re-on*) [dim. of *ἄχωρ*, chaff]. A genus of fungous organisms including several species (possibly modified forms of *Penicillium glaucum*), found in the skin, especially the hair follicles. **A. keratophagus**, the form causing *Onychomycosis*. **A. lebertii**, the parasite of *Tinea tonsurans*. **A. Schönleini**, the species occurring in ringworm, or *Tinea favosa*.

Achroma (*ah-kro'-mah*) [*ἄ* priv.; *χρώμα*, color]. Absence of color. **A.**, Congenital. See *Albinism*. **A. Cutis**. See *Leukoderma*.

Achromatic (*ah-kro-mat'-ik*) [*ἄ* priv.; *χρώμα*, color]. Without color. **A. Lens**, one the dispersing power of which is exactly neutralized by another lens with the same curvature but having a different refractive index.

Achromatin (*ah-kro'-mat-in*) [*ἄ* priv.; *χρώμα*, color]. The groundwork of the nucleus of a cell; it is so called because it is not readily stained by coloring agents.

Achromatism (*ah-kro'-mat-izm*) [*ἄ* priv.; *χρώμα*, color]. Absence of chromatic aberration.

Achromatopsia (*ah-kro-mat-op'-se-ah*) [*ἄ* priv.; *χρώμα*, color; *ὄψις*, eyesight]. Color-blindness; Daltonism.

Achromatosis (*ah-kro-mat'-o'-sis*) [*ἄ* priv.; *χρώμα*, color]. Any disease characterized by deficiency of pigmentation in the integumentary tissues.

Achromia (*ah-kro'-me-ah*) [*ἄ* priv.; *χρώμα*, color]. Albinism; achroma.

Achromoderma (*ah-kro-mo-der'-me-ah*) [*ἄ* priv.; *χρώμα*, color; *δέρμα*, skin]. An albinotic, or colorless state of the skin.

Achromotrichia (*ah-kro-mo-trik'-e-ah*) [*ἄ* priv.; *χρώμα*, color; *τριχίς*, hair]. Absence of pigment from the hair.

Achroodextrin (*ah-kro-o-deks'-trin*) [*ἄ* priv.; *χρῶμα*, colorless; *dexter*, right]. A reducing dextrin formed by the action of the diastatic ferment of saliva upon starch. It is a modification of dextrin and may be precipitated by alcohol; it is not converted into sugar by ptyalin, nor colored by iodine.

Achyilia (*ah-ki'-le-ah*) [*ἄ* priv.; *χυλός*, juice]. Absence of chyle.

Achylosis (*ah-ki-lo'-sis*). Syn. of *Achyilia*.

Achylous (*ah-ki'-lus*) [*ἄ* priv.; *χυλός*, juice]. Deficient in chyle.

Achymosis (*ah-ki-mo'-sis*) [*ἄ* priv.; *χυμός*, chyme]. Deficient formation of chyme.

Acicular (*as-ik'-u-lar*) [*acus*, a needle]. Needle-like.

Acid (*as'-id*) [*acere*, to be sour]. 1. A name applied to any substance having a sour taste. 2. A compound of an electro-negative element with one or more atoms of hydrogen which can be replaced by electro-positive or basic atoms. The majority of acids contain oxygen, and are known as *oxyacids*; those not containing oxygen are termed *hydrogen acids*. Acids vary in their terminations according to the quantity of oxygen or other electro-negative constituent. Those having the maximum of oxygen end in *-ic*; those of a lower degree in *-ous*. When there are more than two combinations the preposition *hyper-* is prefixed to the highest, and *hypo-* to the lowest. Acids that end in *-ic*, as sulphuric acid, form salts terminating in *-ate*; those ending in *-ous* form salts terminating in *-ite*. **A.**, Abietic. See *Abietic*. **A.**, Abrie, $C_{12}H_{24}N_2O$, a crystallizable acid, said to exist in jequirity. **A.**, Acetic, an acid solution composed of 36 parts of absolute acetic acid, $C_2H_4O_2$, and 64 parts of water. It has strongly acid properties. **A.**, Acetic, Dilute, contains six per cent. of absolute acid. Dose $\frac{3}{4}$ -ij (4.0-8.0). An impure form obtained by the destructive distillation of wood is known as wood vinegar, or pyroligneous acid. **A.**, Acetic, Glacial, the absolute acid occurring in crystals melting at 22.5° C. It is an escharotic. **A.**, Aconitic, $C_6H_8O_6$, occurs in several plants, as *Aconitum napellus*, sugar cane, and beetroots. It crystallizes in small plates, that

dissolve readily in alcohol, ether, and water, and melt at $186^{\circ}-7^{\circ}$. **A.**, **Adipic**, $C_6H_{10}O_4$, obtained by oxidizing fats with nitric acid. It crystallizes in shining leaflets, or prisms; is soluble in thirteen parts of cold water; melts at 148° . It is dibasic. **A.**, **Agaric** or **Agaricic**, $C_{16}H_{30}O_5 + H_2O$, a resin acid obtained from the fungus *Polyporus officinalis*, growing on larch trees. The acid has been recommended for the checking of night-sweats. It also checks the other excretions and diminishes thirst. It is mildly cathartic. **Unof.** **A.**, **Aldepalmitic**, $C_{16}H_{30}O_5$, the chief component of the butter of the cow. **A.**, **Alloxanic**, $C_4H_2N_2O_4$, a crystalline acid, obtained by treating alloxan with alkalis. **A.**, **Amido-acetic**. See *Glycin*. **A.**, **Amido-benzoic**, $C_7H_5NO_3$, occasionally found in the urine. **A.**, **Amido-succinamic**, same as *Asparagin*. **A.**, **Angelic**, $C_5H_8O_3$, a crystalline monobasic acid. It exists free along with valeric and acetic acids in the roots of *Angelica archangelica*, and as butyl and amyl esters in Roman oil of cumin. It crystallizes in shining prisms, melts at 45° , and boils at 185° . It has a peculiar smell and taste. **A.**, **Anisic**, $C_6H_8O_3$, obtained by oxidizing anisol and anethol with HNO_3 , and from aniseed by the action of oxidizing substances. **A.**, **Anticylic**, a white, fragrant powder with pleasant, acid taste, readily soluble in water, alcohol, and glycerol; it is used as an antipyretic. Dose gr. $\frac{1}{10}$ (.0006). **A.**, **Arabic**. See *Arabin*. **A.**, **Aromatic**, a name applied to certain organic acids occurring in the balsams, resins, and other odoriferous principles. Also, in pharmacy, a dilute mineral acid reinforced by aromatic substances in order to modify its flavor. **A.**, **Arsenic**, and **Arsenous**. See *Arsenic*. **A.**, **Aspartic**, $C_4H_7NO_4$, occurs in the vinasse obtained from the beet root, and is procured from albuminous bodies in various reactions. It is prepared by boiling asparagin with alkalis and acids, crystallizes in rhombic dibasic prisms, or leaflets, and dissolves with difficulty in water. **A.**, **Auric**, $Au(OH)_3$, gold trihydroxid. **A.**, **Benzoic**, $C_7H_6O_2$, occurs free in some resins, chiefly in gum benzoin, and in coal tar. It crystallizes in white, shining needles, or leaflets, melts at 120° , and distils at 250° . It volatilizes readily, its vapor possessing a peculiar odor. **A.**, **Boracic**, or **Boric**. See *Boron*. **A.**, **Butyric**, $C_4H_8O_2$, an acid having a viscid appearance and rancid smell. It is obtained commercially by the fermentation of a mixture of sugar and butter or cheese in the presence of an alkaline carbonate, but occurs in various plants, in cod-liver oil, in the juice of meats, and in the perspiration. Combined with glycerol as glyceryl butyrate,

it is essentially butter. **A.**, **Caffeic**, $C_9H_8O_4$, obtained when the tannin of coffee is boiled with potassium hydroxid. **A.**, **Camphoric**, $C_{10}H_{16}O_4$, a dibasic acid, obtained by boiling camphor with HNO_3 ; it crystallizes from hot water in colorless leaflets; melts at 178° , and decomposes into water and its anhydrid, $C_8H_{14}(CO)_2O$. It is used in night-sweats of phthisis. Dose gr. x-xxx (0.65-2.0). **A.**, **Capric**, $C_9H_{18}CO.OH$, occurs in small quantity, as a glycerid in cow's butter. It crystallizes in fine needles, melting at $30^{\circ}C.$, and is very insoluble in boiling water. **A.**, **Caproic**, $C_8H_{16}O_2$, the sixth in the series of fatty acids; a clear, mobile oil, colorless, inflammable, and with a very acid and penetrating taste. **A.**, **Caprylic**, $C_7H_{14}CO.OH$, an acid combined with glycerol, forming a glycerid existing in various animal fats; it is liquid at ordinary temperatures. **A.**, **Carbamic**, $H_2N.CO.OH$, carbonic acid in which NH_2 replaces OHI ; it is not known in the free state; its ammonium salt is contained in commercial ammonium carbonate. The esters of carbamic acid are called urethanes. **A.**, **Carbazotic**. See *A*, *Picric*. **A.**, **Carbolic**, C_6H_5OH , phenol,—the correct designation of this substance—is procured from coal tar by fractional distillation. It has a very peculiar and characteristic odor, a burning taste, is poisonous, and has antiseptic properties. The sp. gr. at 0° is 1.084; it crystallizes in colorless rhombic needles that melt at 42.2° , boiling at 180° , and it is not decomposed upon distillation. At ordinary temperatures it dissolves in water with difficulty (1 : 15), but is soluble in alcohol, ether, glacial acetic acid, and glycerol, in all proportions. It unites with bases to form salts, known as *Carbolates*. Upon exposure to light and air it deliquesces and acquires a pinkish color. It is used in the manufacture of many of the artificial coloring matters, e. g., picric acid. It is a powerful antiseptic and germicide. Internally it is useful in vomiting, fermentation in the stomach, and as an intestinal antiseptic; locally, as a caustic. Dose, internally, gr. $\frac{1}{2}$ -ij (0.03-0.13). **A.** **Carb.**, **Aqua**, contains 10 drams of the glycerite to 1 pint of water. Dose $\frac{3}{4}$ ss (4.0-16.0). **A.** **Carb.**, **Gargarisma**, grs. ij- $\frac{3}{4}$ j (0.13-32.0) for fetid sore throat. **A.** **Carb.**, **Glyceritum**, contains acid 1, glycerol 4 parts. **A.** **Carb.**, **Liquefactum** (B. P.). Dose $\frac{1}{2}$ j-ij (0.06-0.13). **A.**, **Carb.**, **Solutions**, vary from 1 to 5 per cent. in water. **A.** **Carb.**, **Suppos. cum Sapone** (B. P.), each contains gr. j (0.06) of carbolic acid. **A.** **Carb.**, **Unguent.**, contains acid 10, ointment 90 parts. **A.** **Carbol.**, **Injectio Hypoderm.**, 2-5 per cent., for anthrax and erysipelas. **A.**, **Carbonic**, CO_2 , carbon

dioxid; an ultimate product of the combustion of carbon compounds; a colorless, odorless gas, heavier than air, incapable of sustaining respiration. **A., Carminic**, $C_{17}H_{18}O_{10}$, a coloring matter found in the buds of certain plants and especially in cochineal, an insect inhabiting different varieties of cactus. It is an amorphous purple-red mass, readily soluble in water and alcohol, and yields red salts with the alkalis. **A., Cathartic**, or **Cathartinic**, an active principle from several species of *Cassia*. **A., Cerotic**, $C_{27}H_{54}O_2$, a fatty acid, existing in beeswax and in Chinese wax. **A., Chloracetic** [*chlorin* and *acetic*], an acid, called also *monochloracetic* acid, produced by the substitution of chlorin for the hydrogen of the radicle in acetic acid. It is sometimes used as a caustic. **A., Chloric**, $HClO_3$, an acid known only in its compounds (*chlorates*) and its aqueous solution. **A., Cholic**. See *A., Cholic*. **A., Cholic**, $C_{26}H_{48}O_5$, *Cholic Acid*, from glycocholic and taurocholic acids; it crystallizes from hot water in small anhydrous prisms, sparingly soluble in water, and melting at 195° . **A., Chromic**, strictly the compound H_2CrO_4 ; it forms salts called chromates. In most books the chromium trioxid, CrO_3 , is called by this name. It is crystalline solid; escharotic. **A., Chrysophanic**, $C_{15}H_{10}O_5$, *Rheinic Acid*, exists in the lichen, *Parmelia parietina*, in senna leaves, and in the rhubarb root. It crystallizes in golden yellow needles or prisms, melting at 162° . See *Chrysarbin*. **A., Cinnamic**, $C_9H_8O_2$, occurs in Peru and Tolu balsams, in storax, and in some benzoin resins. It has been used in tuberculosis, both internally and externally. Dose $m\bar{j}$ -x (0.06-0.65) hypodermatically. **A., Citric**, $C_6H_8O_7$, occurs free in lemons, black currants, bilberry, beets, and in various other acid fruits. It crystallizes with one molecule of water in large rhombic prisms, that melt at 100° , are colorless, inodorous, and extremely sharp in taste. It is refrigerant, antiseptic, and diuretic. **A., Cresolsulphuric**, $C_6H_7O_2SO_3OH$, exists in the urine in small traces. **A., Cresylic**. See *Cresol*. **A., Cyanic**, $CONH$, obtained by heating polymeric cyanuric acid. **A., Cyanuric**. See *A., Tricyanic*. **A., Diacetic**, $C_4H_4O_3$, an acid present in the urine in certain stages of diabetes and other diseased conditions. **A., Fatty**, a monobasic acid formed by the oxidation of a primary alcohol. The fatty acids have a general formula of $C_nH_{2n}O_2$. **A., Fluoric**, HF , gaseous and soluble in water; a strong escharotic. **A., Formic**, CH_2O_2 , an acid obtained from a fluid emitted by ants when irritated; it is also found in stinging nettles, in shoots of the pine, and in various

animal secretions. It is prepared by heating oxalic acid and glycerol. It is a colorless, mobile fluid, with a pungent odor, and vesicates the skin. **A., Gallic**, $C_7H_6O_5$, occurs free in nutgalls, in tea, and in the fruit of various other plants. It is obtained from ordinary tannic acid by boiling it with dilute acids. It crystallizes in fine, silky needles, containing one molecule of water. It dissolves slowly in water and readily in alcohol and ether; has a faintly acid, astringent taste; melts near 220° . It is astringent and disinfectant; useful in night sweats, diabetes, and chronic diarrhea. **A., Gallici**, *Unguent.*, benzoated lard 90, gallic acid 10. **A., Glycocholic**, $C_{26}H_{48}NO_6$, a monobasic acid found in bile; sparingly soluble in water, and crystallizing in minute needles. **A., Glycuronic**, $C_6H_{10}O_7$. This acid has been found in urine; it probably does not exist there normally, but appears after taking certain drugs, as benzol, indol, nitro-benzol, and the quin derivatives. **A., Hippuric**, $C_9H_8NO_3$, *Benzoyl Glycoll*, occurs in considerable amount in the urine of herbivorous animals, sometimes in that of man. It crystallizes in rhombic prisms, and dissolves readily in hot water and alcohol. **A., Hydriodic**, *III*, *Acidum Hydriodicum*; a gaseous acid. Its solution and a syrup prepared from it, *Syrupus acidi hydriodici* (U. S. Ph.), are used as alteratives, with the general effects of iodine. Dose of the syrup, $\bar{f}\bar{3}\bar{j}$ -iv (4.0-16.0). **A., Hydrobromic**, HBr ; the dilute acid, which is the chief form used, consists of 10 parts acid and 90 parts water. It is a solvent for quinin, is useful in hysteria, congestive headaches, and neuralgia, and is recommended as a substitute for potassium and sodium bromids. Dose $m\bar{x}$ - $\bar{f}\bar{3}\bar{j}$ (1.3-8.0). **A., Hydrochloric**, *Muriatic Acid*, HCl , a liquid consisting of 31.9 per cent. by weight of HCl gas in 68.1 per cent. of water. It is colorless, pungent, and intensely acid. **A., Hydrochloric**, *Dilute*, a ten per cent. solution of absolute acid in water. Valuable as an aid to digestion. Dose $m\bar{i}\bar{i}\bar{j}$ -x (0.10-0.65). **A., Hydrocyanic**, *Dilute*, HCN , *Prussic Acid*, a liquid consisting of two per cent. of the acid with 98 per cent. of water and alcohol. It possesses an odor like that of bitter almonds. Prussic acid is found in the bitter almond, the leaves of the peach, and in the cherry laurel, from the leaves of which it is distilled. It is one of the most active poisons known, death from complete asphyxia being almost instantaneous. It is valuable for its sedative effects in vomiting, whooping-cough, and spasmodic affections. Dose $m\bar{j}$ - $\bar{i}\bar{i}\bar{j}$ (0.06-0.2). **A., Hydrocyanic**, *Vapor*, 1 part of dilute acid in 4-6 parts

of water, warmed, and the vapor inhaled to relieve irritable coughs. **A., Hydrofluoric**, HF, an aqueous solution of hydrofluoric gas, powerfully corrosive, used for etching on glass. **A., Hydrosulphuric**, H_2S , a gas formed during the putrefaction of albuminous substances; it occurs in sulphur mineral waters, and is produced by the action of mineral acids on metallic sulphids. It has the odor of rotten eggs. It is also called *Hydrogen Sulphid*, *Sulphuretted Hydrogen*, and *Sulphydric Acid*. **A., Hypochlorous**, $HClO$, an unstable compound, important as a disinfecting and bleaching agent. **A., Hypophosphorous**, H_3PO_2 , its salts (hypophosphites), also the dilute acid, and a syrup prepared from it, are used as remedial agents. (See *Calcii Hypophosphis*, *Ferri Hypophosphis*, *Sodii Hypophosphis*, *Potassii Hypophosphis*.) **A., Indoxylsulphuric**, an acid that, combined with potassium, occurs in the urine as indican. **A., Iodic**, HIO_3 , a monobasic acid. Its solution (two per cent.) has been recommended as an alternative by subcutaneous injection. **A., Lactic**, $HC_2H_3O_2$, a liquid containing 75 per cent. of absolute acid in 25 per cent. of water, produced in the fermentation of milk. It is useful in aiding digestion, in diabetes, in tuberculosis of the larynx, and as a solvent of false membrane in diphtheria. Dose ʒss – ʒss (2.0–16.0) in the 24 hours. **A., Lactic**, Dil. (B. P.). Dose ʒss – ij (2.0–8.0). **A., Linoleic**, $C_{18}H_{34}O_2$, occurs as glycerid in drying oils, such as linseed oil, hemp oil, poppy oil, and nut oil. **A., Malic**, $C_4H_5O_5$, a bibasic acid, occurring free or in the form of salts in many plant juices, in unripe apples, in grapes, and in mountain-ash berries. It forms deliquescent crystals, that dissolve readily in alcohol, slightly in ether, and melt at 100° ; it has a pleasant acid taste. **A., Malonic**, $C_3H_4O_4$, occurs in the deposit found in the vacuum pans employed in the beet-sugar manufacture; it may be obtained by the oxidation of malic acid with chromic acid. **A., Meconic**, $C_7H_4O_7$, a tribasic acid, occurring in opium in union with morphin. It crystallizes with $3H_2O$ in white laminae. **A., Metaphosphoric**, HPO_3 , a glassy solid, freely soluble in cold water, and converted by boiling into orthophosphoric acid. It is used as a test for albumin in the urine. **A., Muriatic**. See *A., Hydrochloric*. **A., Myronic**, $C_{10}H_{18}NS_2O_{10}$, an acid that occurs as a potassium salt in the seeds of black mustard. **A., Nitric**, HNO_3 , a liquid consisting of 68 per cent. absolute acid in 32 per cent. of water. The pure acid is colorless, fuming, and highly caustic. It is used in cauterization of chancres and phagedenic ulcers and as a reagent. **A.,**

Nitric, Dilute, contains 10 per cent. absolute acid. It is used internally to aid digestion; to stimulate the hepatic function, etc. Dose ʒijj – xv (0.2–1.0), well diluted. **A., Nitrohydrochloric**, **A., Nitromuriatic**, *Aqua Regia*, a golden-yellow, fuming mixture of 4 parts of nitric and 15 of hydrochloric acid. It is a solvent of gold, and is valuable in affections of the liver. Dose ʒj – vij (0.06–0.45), very dilute. **A., Nitrohydrochloric**, Dilute, consists of 4 parts nitric and 18 hydrochloric acid, and 78 parts water. Dose ʒv – xx (0.3–1.3), well diluted. **A., Oleic**, $C_{18}H_{34}O_2$, an acid present in many fats and oils. It is a colorless oil, crystallizing on cooling, soluble in alcohol, benzol, and the essential oils; insoluble in water. It saponifies when heated with alkaline bases. It is used in making the oleates. **A., Organic**, an acid characterized by the presence of the carboxyl group, $CO.OH$. **A., Orthophosphoric**, H_3PO_4 , ordinary phosphoric acid, as distinguished from metaphosphoric and pyrophosphoric acids. **A., Osmic**, OsO_4 , the oxid of osmium, one of the rarer elements; it occurs as yellow, acrid, burning crystals, yielding an intensely irritating vapor; it has been recommended for hypodermatic use in sciatica, strumous glands, and cancer. It is used in histology as a fixing agent and as a stain for fat. **A., Oxalic**, $C_2H_2O_4$, a colorless crystalline solid obtained by treating sawdust with caustic soda and potash. It occurs in many plants chiefly as potassium oxalate; with two parts of water it crystallizes in fine transparent monoclinic prisms. It is soluble in nine parts of water at moderate temperature and quite easily in alcohol. It has been recommended in amenorrhea. Dose gr. $\frac{1}{2}$ – $\frac{3}{4}$ (0.032–0.048). In large doses it is a violent poison. **A., Palmitic**, $C_{16}H_{32}O_2$, an acid existing as a glycerin ether in palm-oil and in most of the solid fats. **A., Phosphoric**, H_3PO_4 , *Orthophosphoric Acid*, contains 50 per cent. each of acid and water; it is obtained from bones or by oxidation of phosphorus. **A., Phosphoric**, Dilute, contains 10 per cent. of absolute acid. It is employed in digestive disturbances, in strumous diseases, and to dissolve phosphatic deposits. Dose ʒv – xxx (0.32–2.0). **A., Phosphorous**, H_3PO_3 , a tribasic oxyacid of phosphorus, containing one atom of oxygen less than phosphoric acid. **A., Picric**, $C_6H_2(NO_2)_3OH$, *Carbazotic Acid*, *Trinitrophenol*, obtained by the nitration of phenol. It crystallizes from hot water and alcohol in yellow leaflets or prisms which possess a very bitter taste. It is readily soluble in hot water, its solution imparting a beautiful yellow color to silk and wool. It

is recommended as an antiperiodic and anthelmintic. It is used as a test for albumin and sugar. Dose gr. v-xv (0.32-1.0) *per diem*. **A., Prussic.** See *A., Hydrocyanic*. **A., Pyrogallie,** $C_6H_6O_3$, pyrogallol, formed by heating gallic acid with water to 210° . It forms white leaflets or needles, is readily soluble in water, less so in alcohol and ether. It is useful in the treatment of certain skin diseases, but is poisonous, and must be used with caution. **A., Pyroligneous,** the crude acid obtained in the destructive distillation of wood. It is a clear liquid of reddish-brown color and strong acid taste, with a peculiar penetrating odor described as empyreumatic, due largely to the furfural it contains. It contains from four to seven per cent. of real acetic acid. **A., Pyrophosphoric,** the dihydric phosphate, $2H_2O.P_2O_5$, one of the forms of phosphoric acid. It is poisonous. Its iron salt is used in medicine. The pure acid is a soft, glassy mass. **A., Salicylic,** $C_7H_6O_3$, *Ortho-oxobenzoic Acid*, occurs in the buds of *Spirea Ulmaria*, in the oil of wintergreen, and other varieties of gaultheria. It consists of four-sided prisms and crystallizes readily from hot water in long needles. It is soluble in water and in chloroform, and is antiseptic; it is used in the treatment of acute articular rheumatism and myalgia. Dose grs. v-xx (0.3-1.3), not exceeding \mathfrak{zj} (4.0), daily. **A., Sarcolactic,** $C_6H_8O_6$, occurs in blood and in muscles, to which it gives their acid reaction, especially after the muscles have been in a state of activity. It is also found in urine in phosphorus poisoning. **A., Sclerotinic,** an acid found in ergot, of which it is one of the active principles. **A., Stearic,** $C_{18}H_{36}O_2$, associated with palmitic and oleic acids as a mixed ether, in solid animal fats, the tallow. **A., Succinic,** $C_4H_4O_4$, an acid obtained in the distillation of amber, and also prepared artificially. **A., Sulphanilic,** $C_6H_4(NH_2).SO_3H$, obtained by heating anilin (1 part) with fuming H_2SO_4 (2 parts) to 180° until SO_3 appears. It crystallizes in rhombic plates which effloresce in the air. It is used as a reagent. **A., Sulphocarbolic,** $C_6H_5HSO_4$, phenyl bisulphate, formed by the union of carbolic and sulphuric acids. Its salts, the sulphocarbonates, are used in medicine as intestinal antiseptics, etc. **A., Sulphuric,** H_2SO_4 , *Oil of Vitriol*, a heavy, oily, corrosive acid, consisting of not less than 92.5 per cent. sulphuric anhydrid and 7.5 per cent. of water. It is used as a reagent and as a caustic. **A., Sulph., Aromatic,** contains 20 per cent. acid, diluted with alcohol and flavored with cinnamon and ginger. It is used as an astringent in diarrhea and in

night-sweats; also in hemoptysis. Dose $\mathfrak{m}\mathfrak{v}$ -xv (0.32-1.0). **A., Sulph., Dilute,** contains 10 per cent. strong acid to 90 of water. It is used as an astringent. Dose $\mathfrak{m}\mathfrak{x}$ -xv (0.65-1.0), well diluted. **A., Sulphurous,** H_2SO_3 , a colorless acid containing about 6.4 per cent. of sulphurous anhydrid in 93.6 per cent. of water. The gas, SO_2 , is a valuable disinfectant. The acid is used as a spray or lotion in diphtheria, stomatitis, and as a wash for indolent and syphilitic ulcers. The various hyposulphites are mainly valuable in that they decompose and give off sulphur dioxide. Dose $\mathfrak{m}\mathfrak{v}$ - \mathfrak{zj} (0.32-4.0). **A., Sulphydic.** See *A., Hydrosulphuric*. **A., Tannic,** $C_{12}H_{10}O_8$, *Tannin*, an astringent acid obtained from nutgalls, and occurring in yellowish, scaly crystals. It is soluble in water and alcohol. It is an antidote in poisoning by alkaloids and tartar emetic, and is used as an astringent in catarrh of mucous membranes, and externally in many skin diseases. Dose gr. j-xx (0.065-1.3). **A., Tann., Glyceritum,** one part tannin in four of glycerol. **A., Tann., Suppositories,** one part of tannin to five of butter of cacao. **A., Tann., Troches,** each contains $\frac{1}{2}$ gr. (0.032) of tannic acid. **A., Tann., Unguent,** a ten per cent. ointment of the acid incorporated with benzoated lard. **A., Tartaric,** $C_4H_4O_6$, an astringent acid widely distributed in the vegetable world, occurring principally in the juice of the grape, from which it deposits after fermentation in the form of acid potassium tartrate (argol). It is chiefly employed in refrigerant drinks and in baking powders; 20 grains neutralize 27 of potassium bicarbonate, 22 of sodium bicarbonate, and $15\frac{1}{2}$ of ammonium carbonate. Dose gr. x-xxx (0.65-1.3). **A., Taurocholic,** $C_{24}H_{45}NO_5$, occurs in bile; it is very soluble in water and alcohol and crystallizes in fine needles. **A., Trichloroacetic,** $HC_2Cl_3O_2$, an acid formed from acetic acid, three atoms of the hydrogen of which are (in the new acid) replaced by chlorine. It is used as a reagent for the detection of albumin in the urine, and as a caustic. **A., Uric,** $C_5H_4N_4O_6$, an acid found in the urine of all animals, especially man and the carnivora—rarely in the herbivora—abundantly in the excrement of birds, reptiles, and molluscs. It exists usually in combination with the metals of the alkaline group. It is separated from urine by adding hydrochloric acid and allowing the crystals to settle. **A., Valeric,** $C_5H_{10}O_2$, is formed by oxidizing normal amyl alcohol. It is a mobile liquid with caustic acid taste and the pungent smell of old cheese. **Acid-Albumin** (*as'id-al-bu'-min*). A proteid acted upon or dissolved in the stronger acids, and yielding an acid reaction.

Acidifiable (*as-id'-if-i-a-bl'*) [*acidum*, acid; *fieri*, to become]. Capable of becoming an acid, or of becoming sour.

Acidification (*as-id'-if-ik-a'-shun*) [*acidum*, acid; *facere*, to make]. Conversion into an acid; the process of becoming sour.

Acidimeter (*as-id-im'-et-er*) [*acidum*, acid; *μετρον*, a measure]. An instrument for performing acidimetry.

Acidimetry (*as-id-im'-et-re*) [*acidum*, acid; *μετρον*, a measure]. Determination of the free acid in a solution by an acidimeter, or by chemic reactions.

Acidity (*as-id'-it-e*) [*acidum*, acid]. The quality of being acid; sourness; excess of acid.

Acidulated (*as-id'-u-la-ted*) [*acidulare*, to make sour]. Somewhat sour or acid.

Acidulous (*as-id'-u-lus*) [*acidulare*, to make sour]. Moderately sour.

Acidum (*as'-id-um*) [L.]. See *Acid*.

Acinesia (*as-in-e'-se-ah*) [*ἀκίνησις*, motion]. Motor paralysis.

Acinetic (*as-in-et'-ik*) [*ἀκίνητος*, motionless]. Relating to, or affected with, acinesia.

Aciniform (*as-in'-if-orm*) [*acinus*, a grape]. Grape-like.

Acinus (*as'-in-us*) [*acinus*, a grape; *pl.*, *Acini*]. Any one of the smallest lobules of a compound gland, as an acinus of the liver.

Acme (*ak'-me*) [*ἀκμή*, a point]. The highest point of anything. The critical stage of a disease; the crisis.

Acmon (*ak'-mon*) [*ἀκμων*, an anvil]. The incus.

Acne (*ak'-ne*) [*ἀκνή*, a point]. *Acne vulgaris*; *Varus*; a common, usually chronic, inflammatory disease of the sebaceous glands, occurring mostly about the face, chest, and back. The lesions may be papular, pustular, or tubercular. It occurs usually between the ages of puberty and twenty-four years, is generally worse in winter, and is associated with menstrual and gastrointestinal troubles. The individual lesions consist of minute pink, acuminate papules or pimples, in the center of which is a black-topped comedo (*A. punctata*, *A. papulosa*).

A., **Adenoid**. See *Lupus*, *Disseminated Follicular*. **A. adolescentium**. Synonym of *A. vulgaris*. **A. albida**. Synonym of *Milium*. **A. artificialis**, that form that disappears when the cause is removed. **A. atrophica**. Synonym of *A. varioliformis*. **A. cachecticum**, a form occurring in starved or debilitated persons after long wasting diseases, as phthisis. **A. ciliaris**, acne at the edges of the eyelids. **A. disseminata**. Synonym of *A. vulgaris*. **A. erythematosa**. Synonym of *A. rosacea*. **A. frontalis**. Synonym of *A. varioliformis*. **A. generalis**, acne that has become general

over the surface of the body. **A. hypertrophica**, a stage of *A. rosacea* in which there is a permanent, intensely red, non-inflammatory, nodulated thickening of the tips and sides of the nose, expanding it, both laterally and longitudinally. **A. indurata**, a variety of *A. vulgaris*, characterized by chronic, livid indurations, the result of extensive perifollicular infiltration. It is especially seen in strumous subjects. **A. keratosa**, a rare form of acne in which a horny plug takes the place of the comedo, and by its presence excites inflammation. **A. mentagra**. See *Sycosis*. **A. necrotica**. Synonym of *A. varioliformis*. **A. papulosa**. See *Acne*. **A. picealis**, *Tar Acne*; a form of dermatitis common in fiber-dressers who work with paraffin and in persons otherwise brought in contact with tar or its vapor. It involves chiefly the extensor surfaces of the limbs. **A. punctata**, a variety of *A. vulgaris*. **A. pustulosa**, a variety of *A. vulgaris*, characterized by abscesses. **A. rodens**. Synonym of *A. varioliformis*. **A. rosacea**. *Rosacea*; *Telangiectasis faciei*; *Nervus araneus*; *Brandy Nose*; *Whisky Nose*; *Spider Nevus*; *Spider Cancer*. A chronic, hyperemic, or inflammatory affection of the skin, situated usually upon the face, especially the nose, cheeks, forehead, and chin. **A. scrofulosa**, a variety of *A. cachecticum*, occurring in strumous children. **A. sebacea**. Synonym of *Seborrhea*. **A. simplex**, a variety of *A. vulgaris*. **A. sycosiformis**. Same as *Sycosis non-parasitica*. **A. tarsi**, an inflammatory affection of the large sebaceous glands of the eye-lashes (Meibomian glands). **A. varioliformis**, a rather rare disease, situated chiefly about the forehead, at the junction with the hairy scalp, and extending into the hair. The pustules appear in groups. Its etiology is unknown. **A. vulgaris**. See *Acne*.

Aconite (*ak'-on-it*). See *Aconitum*.

Aconitic Acid (*ak-on-it'-ik*). See *Acid*.

Aconitum (*ak-on-ŭ'-tum*) [L.]. The root of *Aconitum napellus*. It possesses a bitter, pungent taste, and produces numbness and persistent tingling in the tongue and lips. It is very poisonous. It depresses the heart, respiration, circulation, and paralyzes the sensory nerves. It is antipyretic, diaphoretic, and diuretic. The active principle is *Aconitin*. As a diaphoretic and depressant to the circulation, it is highly beneficial in fevers, acute throat affections, and inflammation of the respiratory organs. Dose gr. ss-ij (0.03-0.13). **A.**, **Abstractum**, has double the strength of the powdered drug, or its fluid extract. Dose gr. 1/6-j (0.016-0.065). **A. Extractum**. Dose gr. 1/6-1/3 (0.011-0.022). **A.**, **Ext. Fld.**, has a strength of one drop

to the grain of powdered drug. Dose $\text{m}\frac{1}{2}$ -ij (0.032-0.13). **A., Liniment.** (B. P.), aconite root, camphor, and rectified spirit. **A., Tinct.,** contains, aconite 35, alcohol and water, each *q. s.* to make 100 parts. Dose $\text{m}\frac{1}{2}$ -iv (0.032-0.26). The following are unofficial: **Aconitina**, $\text{C}_{35}\text{H}_{45}\text{NO}_{12}$ (aconitum), an amorphous solid. Dose $\text{gr.}\frac{1}{15}$ - $\frac{1}{30}$ (0.0003-0.0013). **A., Injectio Hypodermica**, one grain in $\frac{1}{2}$ ounce. Dose $\text{m}\frac{1}{2}$ -iv (0.065-0.26). **A. Oleatum**, a two per cent. solution of aconite in oleic acid. **A., Ung.** (B. P.), eight grains to the ounce.

Acorea (*ah-ko'-re-ah*) [*â* priv.; *κόρη*, pupil]. Absence of the pupil.

Acoria (*ah-ko-re'-ah*) [*â* priv.; *κόρος*, satisfaction]. A greedy or insatiable appetite.

Acormus (*ah-kor'-mus*) [*â* priv.; *κόρμος*, the trunk]. A monster without a trunk or body.

Acorus (*ah'-o-rus*) [*â* priv.; *κόρη*, the pupil]. See *Calamus*.

Acoumeter, Acouometer (*ah-koo'-met-er*, *ah-koo-om'-et-er*) [*ἀκούειν*, to hear; *μέτρον*, a measure]. An instrument for measuring the acuteness of hearing.

Acoustic (*ah-koo'-tik* or *ah-kooz'-tik*) [*ἀκουστικός*]. Relating to the ear or sense of hearing.

A. Duct, the external meatus of the ear. **A. Nerve**, the eighth cranial nerve.

A. Tetanus, the rapidity of the induction shocks in a frog's nerve-muscle preparation, as measured by the pitch of a vibrating rod. **A. Tubercle**, a rounded elevation on either side of the floor of the fourth ventricle.

Acoustics (*ah-koo'-tik-s* or *a-kooz'-tik-s*) [*ἀκουστικός*]. The science of sound.

Acquired (*ah-wei'-erd*) [*acquerere*, to acquire]. Obtained; especially obtained after birth, not inherited.

Acraconitin (*ah-kra-con'-it-in*). See *Pseudoconitin*.

Acrania (*ah-kra'-ne-ah*) [*â* priv.; *κράνιον*, skull]. The condition of a monster with partial or complete absence of the cranium.

Acranial (*ah-kra'-ne-al*) [*â* priv.; *κράνιον*, skull]. Without cranium.

Acraturesis (*ah-kra-tu-re'-sis*) [*ἀκράτεια*, lack of strength; *αἵρρασις*, micturition]. Inability to micturate, from atony of the bladder.

Acrid (*ah'-rid*) [*acer*, *acris*, sharp]. Pungent, irritating.

Acridin (*ah'-rid-in*) [*acer*, *acrid*], $\text{C}_{13}\text{H}_9\text{N}$. A substance produced by heating anilin and salicylic aldehyd to 260° with ZnCl_2 . It dissolves in dilute acids with a beautiful green fluorescence, and has a very pungent odor.

Acritical (*ah-kri'-ik-al*) [*â* priv.; *κρίσις*, a crisis]. Without a crisis; not relating to a crisis.

Acroesthesia (*ah-ro-es-the'-ze-ah*). See *Acroesthesia*.

Acroanesthesia (*ah-ro-an-es-the'-ze-ah*) [*ἀκρον*, extremity; *ἀναισθησία*, want of feeling]. Anesthesia of the extremities.

Acroasphyxia (*ah-ro-as-fiks'-e-ah*) [*ἀκρον*, extremity; *â* priv.; *σφίξις*, pulse]. Asphyxia of the extremities, the so-called phenomena of Raynaud.

Acrocephalia (*ah-ro-sef'-a'-le-ah*) [*ἀκρον*, a point; *κεφαλή*, the head]. A deformity of the head in which the vertical diameter is increased and the top is more or less pointed.

Acrocephalic, Acrocephalous (*ah-ro-sef'-al-ik*, *ah-ro-sef'-al-us*) [*ἀκρον*, a point; *κεφαλή*, the head]. Characterized by acrocephalia; having the top of the head unusually high.

Acrocinesis (*ah-ro-sin'-e'-sis*) [*ἀκρος*, extreme; *κίνησις*, movement]. Excessive motility; abnormal freedom of movement, as seen in certain cases of hysteria.

Acrocinetie (*ah-ro-sin'-e'-ik*) [*ἀκρος*, extreme; *κίνησις*, movement]. Characterized by acrocinesis.

Acrodynia (*ah-ro-din'-e-ah*) [*ἀκρος*, an extremity; *δύνη*, pain]. Epidemic erythema; a disease closely allied to pellagra. It is characterized principally by pricking pains in the palm and soles, hyperesthesia followed by anesthesia of these parts, and an erythematous eruption preceded by bullæ, chiefly on the hands and feet. This is followed by exfoliation and dark-brown or black pigmentation.

Acroesthesia (*ah-ro-es-the'-ze-ah*) [*ἀκρος*, extreme; *αἴσθησις*, sensation]. Exaggerated sensitiveness, or sensibility.

Acrolein (*ah-ro'-le-in*) [*acer*, sharp; *oleum*, oil], $\text{C}_3\text{H}_4\text{O}$. Acrylic aldehyd. A volatile liquid derived from the decomposition of glycerol.

Acromania (*ah-ro-ma'-ne-ah*) [*ἀκρος*, extreme; *μανία*, madness]. Incurable or extreme insanity.

Acromastitis (*ah-ro-mas-ti'-tis*) [*ἀκρος*, extreme; *μαστός*, nipple; *itis*, inflammation]. Inflammation of the nipple.

Acromegalia, Akromegaly (*ah-ro meg'-a'-le-ah*, *ah-ro-meg'-al-e*) [*ἀκρος*; *μεγάλη*, large]. A disease characterized by an overgrowth of the extremities and of the face, including the bony as well as the soft parts. The etiology is unknown. In a number of cases the pituitary body has been enlarged; disease of the thyroid gland has also been found in some instances.

Acromial (*ah-ro'-me-al*) [*ἀκρος*, the summit; *ὤμος*, the shoulder]. Relating to the acromion.

Acromio-clavicular (*ah-ro'-me-o-kla-vik'-u-lar*) [*ἀκρος*, the summit; *ὤμος*, the shoulder; *clavus*, a key]. Relating to the acromion and the clavicle.

Acromio-humeral (*ak-ro'-me-o-hu'-mer-al*) [*ἀκρον*, the summit; *ὤμος*, the shoulder; *humerus*]. Relating to the acromion and the humerus. **A. Muscle**, the deltoid.

Acromion (*ak-ro'-me-on*) [*ἀκρον*, the summit; *ὤμος*, the shoulder]. The triangular-shaped process at the summit of the scapula.

Acromio-thoracic (*ak-ro'-me-o-tho-ra'-sik*) [*acromion*; *θώραξ*, thorax]. Relating to the shoulder and thorax.

Acromphalus (*ak-rom'-fal-us*) [*ἀκρον*, extremity; *ὀμφαλός*, the navel]. 1. The center of the umbilicus, to which the cord is attached. 2. The first stage of umbilical hernia, marked by a pouting of the navel.

Acronarcotic (*ak-ro-nar-kot'-ik*) [*acer*, sharp; *ναρκῶν*, to benumb]. Both acrid and narcotic.

Acroneurosis (*ak-ro-nu-ro'-sis*) [*ἀκρον*, an extremity; *νεῦρον*, a nerve]. Any neurosis manifesting itself in the extremities.

Acronyx (*ak'-ro-nix*) [*ἀκρον*, an extremity; *ὄνυξ*, a nail]. The ingrowing of the nail.

Acroparalysis (*ak-ro-par-al'-is-is*) [*ἀκρον*, an extreme; *παρά*, by; *λύειν*, to loose]. Paralysis of the extremities.

Acroparesthesia (*ak-ro-par-es-the'-se-ah*) [*ἀκρον*, extremity; *παρά*, around; *αἰσθήσις*, sensation]. Abnormal or perverted sensation in the extremities.

Acropathology (*ak-ro-path-ol'-o-je*) [*ἀκρον*, an extremity; *πάθος*, disease; *λόγος*, treatise]. The pathology of the extremities.

Acropathy (*ak-ro-p'-a-the*) [*ἀκρόν*, an extremity; *πάθος*, suffering]. Any disease of the extremities.

Acrophobia (*ak-ro-f'-be-ah*) [*ἀκρον*, a height; *φόβος*, fear]. Morbid dread of being at a great height.

Acrotarsium (*ak-ro-tar'-se-um*) [*ἀκρον*, the summit; *ταρσός*, the tarsus]. The instep.

Acrotic (*ak-rof'-ik*) [*ἀ* priv.; *κρότος*, a striking]. Relating to acrotism.

Acrotism (*ak'-krot-izm*) [*ἀ* priv.; *κρότος*, a striking]. Any defective beating of the pulse; failure of the pulse.

Acrylaldehyd (*ak-ril-al'-de-hid*) [*acer*, sharp; *aldehyd*], C_2H_2O . *Acrolein*, the aldehyd of the allyl series; a colorless, mobile liquid, of a pungent odor.

Actea, or **Actæa** (*ak-te'-ah*) [*ἄκτῃ*, the elder]. A genus of ranunculaceous plants having active medicinal qualities. **A. alba**, the white cohosh, has much the same qualities as **A. spicata**. **A. cimicifuga** is more important. See *Cimicifuga*. **A. racemosa**. See *Cimicifuga*. **A. rubra**, red cohosh, and **A. spicata** are purgative and emetic.

Actinic (*ak-tin'-ik*) [*ἄκτις*, a ray]. Referring to those rays of the spectrum capable of producing chemic changes; they occur in the violet and ultra-violet.

Actinism (*ak'-tin-izm*) [*ἄκτις*, a ray]. The chemic quality of light.

Actinomyces (*ak-tin-o-mi'-sez*) [*ἄκτις*, a ray; *μύκης*, fungus]. A vegetable parasite, the cause of the disease actinomycosis. It is also called the *Ray Fungus*. It probably belongs to the Cladothrix group of Schizomycetes. As seen in tissues it presents itself in the form of a rosette of fine filaments clubbed at their outer ends; in the center are numerous coccus-like bodies, the spores of the organism.

Actinomycosis (*ak-tin-o-mi-co'-sis*) [*ἄκτις*, a ray; *μύκης*, a fungus]. A parasitic, infectious, inoculable disease first observed in cattle, and also occurring in man, and characterized by the manifestations of chronic inflammation, with or without suppuration, often resulting in the formation of granulation tumors, especially about the jaws. The disease is due to the presence of a parasite, the *ray fungus*, or *actinomyces*. The disease is often called *lumpy-jaw*, *holdfast*, or *wooden tongue*.

Actinomycotic (*ak-tin-o-mi-kot'-ik*) [*ἄκτις*, ray; *μύκης*, fungus]. Pertaining to or affected with actinomycosis.

Action (*ak'-shun*) [*agere*, to do, or perform]. A doing, a working; especially the performance of a function. **A., Reflex**, an involuntary movement of part of the body resulting from an impression carried by a sensory or afferent nerve to a center, and then sent back by an efferent nerve to the part, usually at or near the source of irritation.

Active (*ak'-tiv*) [*agere*, to do or perform]. 1. Energetic; decisive; as **A. treatment**. 2. Due to an intrinsic force as distinguished from passive, as **A. hyperemia**.

Actual (*ak'-chu-al*) [*agere*, to do]. Real, effective. **A. Caution**. See *Caution*.

Actuation (*ak'-chu-a'-shun*) [*agere*, to do or perform]. The mental function that is exercised between the impulse of volition and its performance.

Acuity (*ak-u'-it-e*) [*acuitas*; *acuere*, to sharpen]. Acuteness or clearness, as acuity of vision.

Acumeter (*ak-oo'-met-er*) [*ἀκούειν*, to hear; *μέτρον*, a measure]. An instrument for testing hearing.

Acuminate (*ak'-u'-min-at*) [*acuminatus*, pointed, acute]. Sharp-pointed.

Acupressure (*ak'-u-presh-ūr*) [*acus*, a needle; *premere*, to press]. An operation to stop hemorrhage by compressing the artery with a needle inserted into the tissues upon either side.

Acupuncture (*ak'-u-punk-chūr*) [*acus*, a needle; *pungere*, to prick]. Puncture of the skin or tissue by one or more needles for the relief of pain, the exit of fluid, the coagulation of blood in an aneurysm, etc.

Acute (*ak-ūt'*) [*acutus*, sharp]. Sharp, sharp-pointed, keen; of diseases, having a rapid onset, a short course, and pronounced symptoms and termination.

Acuteness (*ak-ūt'-nes*) [*acutus*, sharp]. The quality of being acute.

Acutorsion (*ak-u-tor'-shun*) [*acus*, a needle; *torsion*]. The twisting of an artery with a needle as a means of controlling hemorrhage.

Acyesis (*ah-si-e'-sis*) [*ā* priv.; *κῆσις*, pregnancy]. 1. Sterility of the female. 2. The absence of pregnancy.

Ad [*ad*, to]. A Latin preposition, signifying *to*, *toward*, *at*, etc. *Ad deliquium*, to fainting. *Ad libitum*, at pleasure, or according to discretion.

Adactylism (*ah-dak'-til-izm*) [*ā* priv.; *δάκτυλος*, a finger]. The absence of the digits.

Adactylous (*ah-dak'-til-us*) [*ā* priv.; *δάκτυλος*, finger]. Without fingers or toes.

Adam's Apple (*ad'-amz ap'-pl'*). See *Fo-mum Adami*.

Adansonia digitata (*ad-an-so'-ne-ah dij-it-a'-tah*). The Baobab tree, a native of Africa. The bark is used in the form of an infusion, $\frac{3}{4}$ j to Oj, as a remedy for intermittent fever. Unof.

Addison's Disease. Melasma suprarenale, dermatomelasma suprarenale, or cutis area ("bronzed skin"), a disease of the suprarenal capsules, first described by Addison, and characterized by tuberculous infiltration of the capsules, discoloration of the skin, progressive anemia, and asthenia, ending in death from exhaustion. Bronzed skin can occur without disease of the suprarenal capsules, and the latter have been the seat of morbid processes without an accompanying change in the skin. See *Diseases, Table of*.

A. Keloid. See *Diseases, Table of*.

Adducens (*ad-du'-senz*) [*adducere*, to bring toward]. An adductor, a term applied to certain muscles. **A. Oculi**, the internal rectus muscle of the eye.

Adducent (*ad-du'-sent*) [*adducere*, to bring toward]. Performing adduction.

Adduction (*ad-dukt'-shun*) [*adducere*, to bring toward]. Any movement whereby a part is brought toward another or toward the median line of the body.

Adductor (*ad-dukt'-tor*) [*adducere*, to bring toward]. Any muscle effecting adduction.

Adelomorphous (*ad-el-o-mor'-fus*) [*ἀδελός*, not evident; *μορφή*, form]. Not clearly defined; not having a determinate form (a term applied to certain cells in the gastric glands).

Adenia (*ad-e'-ne-ah*) [*ἀδην*, gland]. A hyperplasia of the tissue of lymphatic glands leading to the formation of tumors. **A.**, Leukemic, adenia associated with a leukemic

condition of the blood. **A.**, Simple, a synonym of *Hodgkin's Disease*.

Adeniform (*ad-en'-e-form*) [*ἀδην*, gland; *forma*, resemblance]. Of the shape of a gland; gland-like.

Adenin (*ad'-en-in*) [*ἀδην*, a gland—first discovered in the pancreas], $C_8H_8N_6$. The simplest member of the uric acid group of leukomains, apparently formed by polymerization of hydrocyanic acid. It occurs, with other bases, as a decomposition product of nuclein, and may be obtained from all animal and vegetable tissues rich in nucleated cells. It crystallizes in leaflets with pearly luster. It exists abundantly in the liver and urine of leukocythemic patients. Adenin is not poisonous.

Adenitis (*ad-en-i'-tis*) [*ἀδην*, gland; *τις*, inflammation]. Inflammation of a gland.

Adeno- [*ἀδην*, gland]. A prefix denoting relation to glands.

Adenocarcinoma (*ad-en-o-kar-sin-o'-mah*) [*ἀδην*, gland; *καρκίνος*, a cancer]. Adenoma blended with carcinoma.

Adenocoele (*ad'-en-o-sel*, or *ad-en-o-se'-le*) [*ἀδην*, gland; *κύλη*, a tumor]. A cystic tumor containing adenomatous elements.

Adenocystoma (*ad-en-o-sis-to'-mah*) [*ἀδην*, gland; *κύστις*, tumor; *μα*, tumor]. A cystic adenoma.

Adenofibroma (*ad-en-o-fi-bro'-mah*) [*ἀδην*, gland; *fibra*, fiber]. A combination of adenoma and fibroma.

Adenography (*ad-en-og'-ra-fe*) [*ἀδην*, a gland; *γράφειν*, to write]. A treatise on the glandular system.

Adenoid (*ad'-en-oid*) [*ἀδην*, gland; *ειδός*, resemblance]. Resembling a gland. **A.**

Acne. See *Lupus, Disseminated Follicular*.

A. Body, the prostate gland. **A. Disease**.

Synonym of *Hodgkin's Disease*. **A. Tis-**

sue. Lymphadenoid tissue. **A. Tumor**.

See *Adenoma*. **A. Vegetations**, a term

applied to an hypertrophy of the adenoid

tissue that normally exists in the naso-phar-

ynx.

Adenology (*ad-en-ol'-o-je*) [*ἀδην*, gland; *λόγος*, a discourse]. The science of the glandular system.

Adenoma (*ad-en-o'-mah*) [*ἀδην*, gland; *μα*, tumor]. An epithelial tumor constructed after the type of a secreting gland. **A. des-**

truens, a destructive form of adenoma.

A., Malignant, an adenomatous carcinoma.

A., Racemose, an adenoma after the type

of a racemose gland. **A. Simplex**, a tumor-

like hyperplasia of a gland. **A.**, Tubular,

an adenoma after the type of a tubular

gland.

Adenomalacia (*ad'-en-o-mal-a'-she-ah*)

[*ἀδην*, a gland; *μαλαξία*, softening]. Ab-

normal softening of a gland.

Adenomyoma (*ad-en-o-mi-o'-mah*) [*ἀδὴν*, a gland; *μῦς*, a muscle; *oma*, tumor]. A tumor composed of glandular and muscular tissues.

Adenomyxoma (*ad-en-o-miks-o'-mah*) [*ἀδὴν*, a gland; *μύξα*, mucus; *oma*, tumor]. A growth having the characters of adenoma and myxoma.

Adenopathy (*ad-en-op'-a-the*) [*ἀδὴν*, a gland; *πάθος*, disease]. Any disease of a gland.

Adenopharyngitis (*ad-en-o-far-in-jit'-tis*) [*ἀδὴν*, a gland; *φάρυγξ*, the throat; *τις*, inflammation]. Inflammation of the tonsils and pharynx.

Adenophthalmia (*ad-en-off-thal'-me-ah*) [*ἀδὴν*, gland; *ὀφθαλμός*, the eye]. Inflammation of the Meibomian glands.

Adenosarcoma (*ad-en-o-sar-ko'-mah*) [*ἀδὴν*, a gland; *σάρξ*, flesh; *oma*, a tumor]. A tumor with the characters of adenoma and sarcoma.

Adeps (*ad'-eps*) [L., *gen.*, *Adipis*]. Lard. The fat obtained from the abdomen of the hog, composed of 38 per cent. stearin and margarin, and 62 per cent. olein. It forms 70 per cent. of ceratum and 80 of unguentum.

A. anserinus, goose grease. **A. benzoïnatus**, contains two per cent. of benzoin.

A. lanæ, lanolin. **A. lanæ hydrosus**, hydrous wool fat, the purified fat of the wool of the sheep. **A. ovillus**, mutton suet.

Adipis, Oleum, a fixed oil expressed from lard. **A. preparatus** (B. P.), purified fat of the hog. **Unguentum Simplex**, contains white wax, benzoated lard, and almond oil. **A. suillus**, hog's lard; adeps.

Adermia (*ah-der'-me-ah*) [*ἀ* priv.; *δέρμα*, skin]. Absence of the skin.

Adhesion (*ad-he'-zhun*) [*adharere*, to stick to]. 1. The attractive force between two dissimilar bodies that are in contact. 2. Abnormal union of two surfaces as a result of inflammation, etc. **A.**, Primary, called also *Healing by First Intention* and by *Immediate Union*, a method of healing of wounds by the production of lymph, followed by the vascularization and cicatrization of the exudate. **A.**, Secondary, or *Healing by Second Intention*, or by *Granulation*, is that mode of healing attended by the production of pus and the formation of granulations. **A.**, Plaster. See *Emplastrum*.

Adhesive (*ad-he'-siv*) [*adharere*, to adhere]. Sticky, tenacious. **A.** Inflammation, inflammation accompanied by plastic exudation, and tending to the union of apposed surfaces. **A.** Plaster. See *Resin* and *Emplastrum*.

Adiantum (*ad-e-an'-tum*) [*ἀ* priv.; *διαντός*, wettable]. A genus of ferns; the maiden-hair. **A. capillus Veneris** and **A. pedatum**, of N. America, are serviceable in coughs and as demulcents.

Adipic (*ad-ip'-ik*) [*adepts*, lard]. Of or belonging to fat. **A. Acid**. See *Acid*.

Adipocere (*ad'-ip-o-ser*) [*adepts*, fat; *cera*, wax]. A wax-like substance formed by the exposure of fleshy tissue to moisture, with the exclusion of air, *i. e.*, in the earth or under water. It consists of the fatty acids in combination with the alkaline earths and ammonium. Human bodies in moist burial places often undergo this change.

Adipoma (*ad-ip-o'-mah*) [*adepts*, fat; *oma*, a tumor; *pl.*, *Adipomata*]. A fatty tumor; lipoma.

Adipose (*ad'-ip-ös*) [*adepts*, fat]. Fatty. **A.** Tissue, fatty tissue, which is distributed extensively through the body. It consists of areolar connective tissue, the cells of which contain fat globules.

Adiposis (*ad-ip-o'-sis*) [*adepts*, fat]. Corpulence; fatty infiltration. **A. dolorosa**, a disease characterized by the formation of soft nodules throughout the connective tissue of the body, accompanied by neuralgic pains. **A. hepatica**, fatty degeneration or infiltration of the liver.

Adipositas (*ad-ip-os'-it-as*) [L.]. Fatness; corpulency.

Adiposity (*ad-ip-os'-it-e*). Fatness; corpulency.

Adipsia (*ah-dip'-se-ah*) [*a* priv.; *διψα*, thirst]. Absence of thirst.

Aditus (*ad'-it-us*) [*ad*, to; *ire*, to go]. In anatomy, an entrance. **A. laryngis**, the entrance to the larynx.

Adjuster (*ad'-jus'-ter*) [Fr., *adjuster*, to adjust]. 1. He who or that which adjusts. 2. A device formerly used for the forcible reduction of dislocations.

Adjustment, Coarse. The rack and pinion for raising or lowering the tube of a microscope a considerable distance. **A.**, Fine, the micrometer screw generally at the top of the column of a microscope for raising or lowering the tube slowly through a short distance.

Adjuvant (*aj'-u-vant*) [*adjuvare*, to assist]. A medicine that assists the action of another to which it is added.

Adnexa (*ad-neks'-ah*) [*ad*, to; *nectere*, to join]. Adjunct parts, as the adnexa of the uterus. **A. bulbi**, the appendages of the bulb of the eye.

Adolescence (*ad-o-les'-ens*) [*adolescere*, to grow]. Youth, or the period between puberty and maturity, usually reckoned as extending in males from about 14 to 25 years, and in females from 12 to 21 years.

Adonidin (*ad-on'-id-in*) [*Adonis*]. A glucosid derived from *Adonis vernalis*, a plant indigenous in Europe and Asia. It is recommended in cardiac dropsy. Dose gr. $\frac{1}{8}$ – $\frac{1}{4}$ (0.0075–0.015). Unof.

Adonis æstivalis (*ad-o'-nis æ-tiv-a'-lis*). A plant much used in Italy as a cardiac tonic.

Similar properties are possessed by *A. ver-nalis*. Dose of fld. ext. m-jj (0.06-0.13).

A., Tinctura, dose m-xxx (0.65-2.0).

Adrenal (*ad-rè'-nal'*) [*ad*, near to; *ren*, the kidney]. 1. Adjacent to the kidney. 2. The suprarenal capsule.

Adrue (*ad-rù'-e*). Antiemetic root. The root of *Cyperus articulatus*; it is anthelmintic, aromatic, stomachic. Dose of the fld. ext. gtt. xx-xxx (1.3-2.) Unof.

Advancement (*ad-vans'-ment*) [Fr. *avancer*, to advance]. The act of bringing or going forward. Specifically, an operation to remedy strabismus, generally in conjunction with tenotomy, whereby the opposite tendon from the over-acting one, having been cut, is brought forward, so that, growing fast in a more advanced position, it shall have more power to act upon the globe of the eye.

Adventitia (*ad-ven-tish'-e-ah*) [*adventitius*, foreign]. The external coat of a blood-vessel.

Adynamia (*ah-din-a'-me-ah*) [*à priv.*; *dynamis*, power]. Deficiency or loss of vital or muscular power; prostration.

Adynamic (*ah-din-am'-ik*) [*à priv.*; *dynamis*, power]. Pertaining to or characterized by adynamia.

Æ-. See *E-*.

Ægophony (*e-goff'-o-ne*). See *Egophony*.

Æquator. See *Equator*.

Aer (*a'-er*). See *Air*.

Aerated (*a'-er-a-ted*) [*âhp*, the atmosphere]. Impregnated or charged with carbon dioxide or air.

Aeration (*a'-er-a'-shun*) [*âhp*, air]. The process of supplying or charging with air or with some gas, such as carbon dioxide; the state of being supplied with air or gas.

Aerial (*a'-er-e-al*). Pertaining to the air; conveyed by the air, as *A.* conduction of sound waves.

Aeriform (*a'-er-re-form*) [*âhp*, air; *forma*, form]. Resembling air or gas.

Aerobic (*a'-er-oh'-ik*) [*âhp*, air; *βιος*, life]. Requiring oxygen (air) in order to live. A term applied to those bacteria requiring free oxygen. Those that do not grow in its presence are called *anaerobic*. Between these extremes there are forms that are able to grow without oxygen under favorable conditions, though they make use of it when present; others that may grow in its presence, though flourishing best in its absence; these are called respectively *facultative aerobic* or *facultative anaerobic*, in distinction from those first mentioned, which are called *obligatory aerobic* or *obligatory anaerobic*.

Aerobiotic (*a'-er-o-hi-oh'-ik*) [*âhp*, air; *βιωτικός*, pertaining to life]. Thriving only in the presence of air.

Aerodynamics (*a'-er-o-di-nam'-ics*) [*âhp*, air; *δυναμις*, power]. The branch of physics that deals with gases in motion.

Aerography (*a'-er-og'-ra-fe*) [*âhp*, air; *γραφη*, a writing]. The description of the air and its qualities.

Aerology (*a'-er-ol'-o-je*) [*âhp*, air; *λόγος*, treatise]. The science of the air and its qualities.

Aerometer (*a'-er-om'-et-er*) [*âhp*, air; *μέτρον*, a measure]. An instrument for ascertaining the density of gases.

Aerophobia (*a'-er-o-fò'-be-ah*) [*âhp*, air; *φόβος*, fear]. Dread of a current of air.

Aerophone (*a'-er-o-fon*) [*âhp*, air; *φωνή*, sound]. An instrument for increasing the amplitude of sound-waves.

Aerophore (*a'-er-o-fôr*) [*âhp*, air; *φορεῖν*, to carry]. 1. A device for the inflation of the lungs of a still-born child with air. 2. A breathing apparatus to be used by firemen and others, to prevent the inhalation of noxious gases.

Aeroplethysmograph (*a'-er-o-pleth-iz'-mo-graf*) [*âhp*, air; *πλήθος*, fulness; *γράφειν*, to write]. An apparatus for registering graphically the expired air; the latter raises a very light and carefully equipoised box placed over water, and this moves a writing-style.

Aeroscope (*a'-er-o-skôp*) [*âhp*, air; *σκοπεῖν*, to observe]. An instrument for estimating the purity of the air.

Aeroscopy (*a'-er-oh'-ko-pe*) [*âhp*, air; *σκοπεῖν*, to observe]. The investigation of atmospheric conditions.

Aerostatics (*a'-er-o-stat'-iks*) [*âhp*, air; *στατικός*, standing]. The branch of physics that treats of the properties of gases at rest.

Aerotherapeutics, Aerotherapy (*a'-er-ther-a-pu'-tiks, a'-er-o-ther'-a-pe*) [*âhp*, air; *θεραπεύειν*, to heal]. A mode of treating disease by compressed or rarefied air or by other gases.

Aerothorax (*a'-er-o-tho'-raks*) [*âhp*, air; *θώραξ*, the chest]. See *Pneumothorax*.

Aerotonometer (*a'-er-o-ton-om'-et-er*) [*âhp*, air; *τόνος*, tension; *μέτρον*, measure]. An instrument for estimating the tension of gases in the blood.

Aerourethroscopy (*a'-er-o-u-re-thros'-ko-pe*) [*âhp*, air; *οὐρήθρα*, urethra; *σκοπεῖν*, to examine]. Urethroscopy conjoined with inflation of the urethra with air.

Aerteriversion (*a'-er-ter-iv-er'-shun*) [*âhp*, air; *τηρεῖν*, to hold; *vertere*, to turn]. A method of arresting hemorrhage by everting the cut end of an artery, invaginating the vessel in itself, and fixing the parts by a needle.

Aerteriverter (*a'-er-ter-iv-er'-ter*) [*âhp*, air; *τηρεῖν*, to hold; *vertere*, to turn]. An instrument used in performing aerteriversion.

Ærugo (*c-ru'-go*) [L., *gen. Æruginis*]. 1. Rust of a metal. 2. Subacetate of copper.
Æ. Ferri, the subcarbonate of iron. **Æ. Plumbi**, lead carbonate, or subcarbonate.

Æsculin (*es'-ku-lin*). See *Esculin*.

Æsculus hippocastanum (*es'-ku-lus hip-o-kas-ta'-num*) [L.]. Horse-chestnut. The fruit of the common "buckeye," or horse-chestnut. It is recommended in hemorrhoids. Dose gr. iij (0.20). Unof.

Æstas (*es'-tas*) [L., *pl.*]. Freckles, or sunburn.

Æsthesio- (*es-the'-se-o*). For all compounds of this word see *Esthesio-*.

Æsthesia (*es-the'-sis*). See *Esthesia*.

Æsthesodic (*es-the-sod'-ik*). See *Esthesodic*.

Æstival (*es'-tiv-al*). See *Estival*.

Æstivation (*es-tiv-a'-shun*). See *Estivation*.

Ætas (*e'-tas*) [L.]. Age; a period of life. See *Age*.

Æther (*e'-ther*). See *Ether*.

Æthyl (*eth'-il*). See *Ethyl*.

Ætiology (*e-te-ol'-o-je*). See *Etiology*.

Afebrile (*ah-feb'-ril*) [*à priv.*; *febrilis*, feverish]. Without fever.

Affection (*af-ek'-shun*) [*afficere*, to affect]. Disease.

Affective (*af-ek'-tiv*) [*afficere*, to affect]. Exciting emotion. **A. Faculties**, the emotions and propensities—especially those peculiar to man. **A. Insanity**, emotional or impulsive insanity.

Afferent (*af'-er-ent*) [*afferens*, carrying to]. Carrying toward the center. Of *nerves*: conveying impulses toward the central nervous system; sensory, centripetal. Of *blood-vessels*: those, as the arteries, conveying blood to the tissues. Of *lymphatics*: those conveying lymph to a lymphatic gland.

Afferentia, or **Vasa afferentia** (*va'-sah af-er-ent'-she-ah*) [L.]. 1. The lymphatic vessels. 2. Any vessels carrying blood or lymph to an organ.

Affiliation (*af-il-e-a'-shun*) [*ad*, to; *filius*, son]. In medical jurisprudence, the act of imputing or affixing the paternity of a child in order to provide for its maintenance.

Affinity (*af-in'-it-e*) [*affinis*, akin to]. Relationship. Attraction. **A., Chemic**, the force, exerted at inappreciable distances, that unites atoms of different substances. **A., Elective**, the preference of one substance for another.

Afflux (*af'-lux*) [*affluere*, *affluxus*, to flow toward]. The flow of the blood or other liquid to a part.

Affusion (*af-u'-zhun*) [*affundere*, to pour upon]. The pouring of water upon an object, as upon the body in fever, to reduce temperature and calm nervous symptoms.

African (*af'-rik-an*) [*Africa*]. Pertaining to Africa. **A. Arrow poison**. See *Stro-*

phanthus. **A. Fever**. Synonym of *Dengue*. **A. Gum**, gum arabic. **A. Lethargy**, or **Nelavan**, a "sleeping sickness" affecting negroes of the West African coast. Increasing somnolence is the characteristic symptom. It is very fatal. Death from exhaustion follows in from three to six months.

After (*af'-ter*) [A. S., *after*, back]. 1. The anus. The buttocks. 2. Next in succession.

A.-Action, the negative variation in an electric current continuing for a short time in a tetanized muscle.

A.-Birth, the popular designation of the placenta, cord, and membranes.

A.-Brain. See *Hindbrain* and *Metencephalon*.

A.-Cataract, *Cataracta Secundaria*; an opacity of the media of the eye after operation for cataract due to opacification of the capsule or to non-absorption of the remains of the lens-substance.

A.-Hearing, a condition in which sounds are heard after the wave-motion that produces them has ceased.

A.-Images, continued retinal impressions after the stimulus of the light or image has ceased to act.

A *positive after-image* is a simple prolongation of the sensation; a *negative after-image* is the appearance of the image in complementary colors.

A.-Pains. See *Pains*.

A.-Perception, the perception of a sensation after the stimulus has passed away.

A.-Sensation, a sensation lasting longer than the stimulus producing it.

A.-Taste, a gustatory sensation produced some time after the stimulus has been removed.

Agalactia (*ah-gal-ak'-te-ah*) [*à priv.*; *γάλα*, milk]. Failure of secretion of the milk after childbirth.

Agar-agar (*a'-gar-a'-gar*) [Ceylon]. A kind of glue made from certain sea-weeds, such as *Gracilaria lichenoides* and *Gigartina speciosa*, used in medicine to make suppositories, and in bacteriologic studies in the preparation of culture media.

Agaric (*ag-ar'-ik*). Touchwood; Spunk; Tinder; the product of different species of *Boletus*, a genus of mushrooms. *Boletus laricis*, *Polyporus officinalis*—is the *white* or *purging agaric*. *Agaricin* is an impure alcoholic extract, which in doses of 1-3 grains (0.065-0.19) [U. S. Dispensatory] gr. $\frac{1}{2}$ - $\frac{1}{4}$ (0.005-0.01) [Brunton] is valuable in the night-sweats of pulmonary tuberculosis.

Agaric or *Agaricinic acid*, in doses of gr. $\frac{1}{8}$ - $\frac{1}{4}$ (0.0038-0.02) is also useful in night-sweats.

A., Ext. Dose gr. iij-vj (0.19-0.39).

A., Tinct. Dose m iij-xx (0.19-1.3).

A. chirurgorum, *Boletus chirurgorum*, Surgeon's *Agaric*, a parasitic fungus formerly used for *Moxa*. Soaked in solution of potassium nitrate it forms *spunk*.

A. muscarius, Fly Agaric, a poisonous mushroom, contains an alkaloid, *Muscarin*. Dose of

the alkaloid gr. $\frac{1}{8}$ -ij (0.008-0.13). *Muscarin Nitrate*, used hypodermatically. Dose gr. $\frac{1}{16}$ - $\frac{1}{4}$ (0.0065-0.048).

Agaricin (*ag-ar'-is-in*) [see *Agaric*], $C_{16}H_{30}O_5 + H_2O$. A white crystalline substance, the active principle of *Agaricus albus*. It has proved useful in the night sweats of pulmonary tuberculosis. Dose gr. $\frac{1}{16}$ - $\frac{1}{8}$ (0.0032-0.0065). Unof.

Agaster (*ah-gas'-ter*) [*â* priv.; *γαστήρ*, the stomach]. One without a stomach.

Agastric (*ah-gas'-tric*) [*a* priv.; *γαστήρ*, the stomach]. Without an intestinal canal, as the tape-worms.

Agathin (*ag'-ath-in*) [*ἀγαθός*, good], $C_8H_8(OH).CH.N.N.(CH_3).C_6H_5$. A greenish-white, crystalline substance, obtained by the interaction of salicylic aldehyd and alpha-methyl-phenyl-hydrazin. It is used as an antineuralgic in doses of eight grains (0.52) two or three times daily. Its action is cumulative. Unof.

Agave (*a-ga'-ve*) [*ἀγανή*, noble]. A large genus of amaryllidaceous plants, natives of North America. *A. americana*, American Aloe, the leaves of a plant growing in North America. It is diuretic and antispythilitic. Dose of the fld. ext. $\frac{3}{4}$ ss-j (2.0-4.0). The fresh juice is also similarly employed. The fermented juice, called *pulque*, is a moderately stimulant drink, very popular in Mexico. Unof.

Agnesia, Agnesis (*ah-jen'-e'-se-ah, ah-jen'-es-is*) [*â* priv.; *γένεσις*, generation]. Absent development.

Agnosomus (*ah-jen-o-so'-mus*) [*â* priv.; *γενῶν*, to beget; *σῶμα*, body]. A variety of single autositic monsters, of the species *Celosoma*, in which there is a lateral or median eventration occupying principally the lower portion of the abdomen, while the genital and urinary organs are either absent or very rudimentary.

Ageusia (*ah-gu'-se-ah*) [*â* priv.; *γεῦσις*, taste]. Abolition of the sense of taste.

Agglomerate (*ag-lom'-er-ât*) [*agglomerare*, to wind into a ball]. Grouped or clustered.

Agglutinant (*ag-lu'-tin-ant*) [*agglutinare*, to paste to]. See *Agglutinative*.

Agglutinate (*ag-lu'-tin-ât*) [*agglutinare*, to paste to]. To glue together; to unite by adhesion.

Agglutination (*ag-lu-tin-a'-shun*) [*agglutinare*, to glue to]. A joining together.

Agglutinative (*ag-lu'-tin-a-tiv*) [*agglutinare*, to glue upon or to]. 1. Favoring agglutination; adhesive. 2. Any substance with adhesive properties, fitted to retain the edges of wounds in apposition.

Aggregate (*ag'-re-gât*) [*ad*, to; *gregare*, to collect into a flock]. Grouped into a mass.

Aglobulia (*ah-glo-bu'-le-ah*) [*â* priv.; *globu-*

lus, a globule]. A decrease in the quantity of red blood corpuscles.

Aglossia (*ah-glos'-e-ah*) [*â* priv.; *γλῶσσα*, the tongue]. Absence of the tongue.

Aglossus (*ah-glos'-us*) [*â* priv.; *γλῶσσα*, the tongue]. A person without a tongue.

Agmatology (*ag-mat-ol'-o-je*) [*ἀγμός*, a fracture; *λόγος*, discourse]. The science or study of fractures.

Agminate (*ag'-min-at*) [*agmen*, a multitude]. Aggregated, clustered. A. Glands. See *Gland*, *Peyer's*.

Agnail (*ag'-nâl*) [A. S., *angnagl*]. 1. Hangnail. 2. A whitlow. 3. A corn.

Agnathia (*ag-na'-the-ah*) [*â* priv.; *γνάθος*, a jaw]. Absence or defective development of the jaws.

Agnin (*ag'-nin*) [*agnus*, a lamb]. A fatty substance derived from sheep's wool.

Agomphious (*ah-gom'-fe-us*) [*â* priv.; *γομφίος*, a tooth]. Without teeth.

Agomphosis (*ah-gom-fol'-sis*) [*â* priv.; *γομφίος*, a tooth]. 1. Absence of the teeth. 2. A loosening of the teeth.

Agony (*ag'-o-ne*) [*ἀγωνία*, a contest or struggle]. Violent pain; extreme anguish; the death-struggle.

Agoraphobia (*ag-o-ra-fol'-be-ah*) [*ἀγορά*, a market-place, assembly; *φόβος*, fear]. A morbid fear of open places or spaces.

Agrammatism (*ah-gram'-at-izm*) [*â* priv.; *γράμμα*, a word]. A phenomenon of aphasia, consisting in the inability to form words grammatically, or the suppression of certain words of a phrase; a form of aphasia.

Agraphia (*ah-gra'-fe-ah*) [*â* priv.; *γράφειν*, to write]. Inability to express ideas by writing. In some cases not a single letter can be formed (*A.*, *Absolute or Literal*); in others, letters are strung together in a manner resembling words, but without meaning (*A.*, *Verbal*).

Agraphic (*ah-gra'-fik*) [*â* priv.; *γράφειν*, to write]. Affected with agraphia.

Agria (*ag'-re-ah*) [*ἀγριος*, wild]. A pustular eruption; malignant pustule; herpes.

Agrimony (*ag'-rim-o-ne*) [*ἀγρός*, a field; *μόνος*, alone]. The root of *Agrimonia eupatoria*, a mild astringent. Dose of fld. ext. $\frac{3}{4}$ ss-ij (2.0-8.0). Unof.

Agriothymia (*ag-re-o-thi'-me-ah*) [*ἀγριός*, wild; *θυμός*, mind, will]. Maniacal fury.

Agrippa (*ah-grip'-ah*) [L.]. A person born with the feet foremost.

Agromania (*ag-ro-ma'-ne-ah*) [*ἀγρός*, a field; *μανία*, madness]. A mania for living in the country.

Agrypnia (*ah-grip'-ne-ah*) [*â* priv.; *ὑπνος*, sleep]. Loss of sleep, insomnia.

Agrypnotic (*ah-grip-not'-ik*) [*â* priv.; *ὑπνος*, sleep]. 1. Preventing sleep. 2. A medicine that prevents sleep.

Ague (*ah'-gu*) [*acutus*, sharp, acute; Fr., *aigu*]. Malarial or intermittent fever, characterized by paroxysms, consisting of chill, fever, and sweating, at regularly recurring times, and followed by an interval or intermission the length of which determines the epithets, quotidian, tertian, etc. In some cases there is a double paroxysm, and hence called double quotidian, double tertian, etc. The duration of each paroxysm varies from two to twelve hours. See *Malarial Fever*. **A.**, **Aden**. See *Dengue*. **A.**, **Brass-founders'**, a disease common among brass-founders, characterized by symptoms somewhat resembling an imperfect attack of intermittent fever, the recurrence of the paroxysms, however, being irregular. The direct cause is generally thought to be the inhalation of the fumes of deflagrating zinc or "spelter." **A.**, **Brow**, intermittent neuralgia of the brow. **A. Cake**, chronic enlargement of the spleen in diseases of malarial origin. **A.**, **Catenating**, ague associated with other diseases. **A. Drop**. See *Fowler's Solution*. **A.**, **Dumb**, latent ague; masked ague; ague without well-marked chill, and with at most only partial or slight periodicity. **A.**, **Latent**. See *A.*, *Dumb*. **A.**, **Masked**. See *A.*, *Dumb*. **A. Tree**, common Sassafras. **A. Weed**. 1. See *Grindelia*. 2. The *Eupatorium perforiatum*, or thorough-wort.

Ailanthus (*a-lan'-thus*) [Malacca, *Ailanto*, "tree of heaven"]. The bark of *A. glandulosa*, commonly known as Tree of Heaven. Its properties are due to an oleoresin and a volatile oil. It is a nauseant and drastic purgative, and an excellent anthelmintic against tape-worm. **A.**, **Ext. Fld.** Dose $\text{m}\bar{x}-3j$ (0.65-4.0). **A.**, **Tinct.** Dose $\text{m}\bar{x}-3ij$ (0.65-8.0).

Ailment (*ah'-ment*) [M.E., *cyle*]. A disease; sickness; complaint.

Ainhum (*in'-hum*) [Negro word, meaning *to shrivel*]. A disease of Guinea and Hindostan, peculiar to negroes, in which the little toes are slowly and spontaneously amputated at about the digitoplar fold. The process is very slow, is unaccompanied by any constitutional symptoms, and its cause is unknown. It sometimes attacks the great toe.

Air [*ahp*, from *aeu*, to blow, or breathe]. The atmosphere. *Atmospheric air* consists of a mixture of 77 parts by weight, or 79.19 by volume, of nitrogen, and 23 parts by weight, or 20.81 by volume, of oxygen, with 3.7 to 6.2 parts by volume of CO₂ in 10,000 parts. It also contains traces of ammonia, argon, nitrites, and organic matter. By virtue of its oxygen it is able to sustain respiration. 100 cubic inches weigh 30,935 grains. The pressure of the air at sea-level is about 14 $\frac{3}{4}$ pounds upon the square inch. **A.**, **Alkaline**, free or

volatile ammonia. **A.**, **Azotic**, nitrogen. **A. Bag**. See *A. Cushion*. **A. Bath**, therapeutic exposure to air which may be heated, condensed, or variously medicated. **A.-bed**, an air-tight, rubber mattress, inflated with air, employed in conditions requiring long confinement to bed. **A.-bladder**. See *A.-vesicle*. **A.-cell**, an air-sac; an air-vesicle of the lung. **A.**, **Complemental**, the amount of air that can still be inhaled after an ordinary inspiration. **A. Conduction**, a method of testing the hearing-power by means of a watch held at varying distances from the ear, or by the employment of a number of tuning forks of varying pitch. **A.-cure**, the therapeutic employment of air. **A.-cushion**, a cushion filled with air, and usually made of soft india-rubber. **A.**, **Dephlogisticated**, an old name for oxygen. **A.-douche**, the inflation of the middle ear through the nose. **A.-embolism**, the entrance of free air into the blood-vessels during life. **A.**, **Expired**, that driven from the lungs in expiration. **A.**, **Fixed**, an old name for carbon dioxide. **A.**, **Inspired**, that taken into the lungs on inspiration. **A.**, **Mephitic**, carbon dioxide. **A. Passages**, the nares, mouth, larynx, trachea, and bronchial tubes. **A.-pump**, an apparatus for exhausting or compressing air. **A.**, **Reserve or Supplemental**, the air that can still be exhaled after an ordinary expiration. **A.**, **Residual**, that remaining in the lungs after the most complete expiration possible. **A.-sac**. See *Air-vesicle*. **A.**, **Stationary**, that remaining in the lungs during normal respiration. **A.**, **Supplemental**. See *A.*, *Reserve*. **A.-tester**, an instrument for testing the purity of the air. **A.**, **Tidal**, that taken in and given out at each respiration. **A.-vesicles**, the alveoli of the lung, the ultimate division of the air-passages. **A.**, **Vital**, an old name for Oxygen.

Akanthion (*a-kan'-the-on*). See *Acanthion*. **Akatamathesia** (*ah-kat-am-ath-e'-ze-ah*) [*â* priv.; *καταμάθησις*, understanding]. Inability to understand.

Akinesia, Akinesis (*ah-kin-e'-se-ah, ah-kin-e'-sis*) [*â* priv.; *κίνησις*, motion]. Absent or imperfect motion. **A. Algeria**, an affection characterized by abstinence from voluntary movement on account of pain, which any active muscular effort causes. The condition is probably a form of neurasthenia.

Akromegaly (*ak-ro-meg'-a-le*). See *Acromegalia*.

Al. 1. The Arabic definite article, *the*, prefixed to many words to designate pre-eminence, etc., as alkali, alcohol. 2. A chemical suffix denoting similarity to or derivation from an aldehyd, as chloral, butyral, etc.

Ala (*al'-lah*) [L., a wing; *pl.*, *Alæ*]. A wing. **A. Auris**, the pinna of the ear. **A. cinerea**, a triangular space of gray matter in the fourth ventricle of the brain, probably giving origin to the pneumogastric nerves. **A. majores**. 1. The greater wings of the sphenoid. 2. The external labia pudendi. **A. minores**. 1. The lesser wings of the sphenoid. 2. The labia minora pudendi. **A. Nasi**, the lateral cartilage of the nose. **A. parvæ**, the lesser wings of the sphenoid. **A. Pontis**. See *Ponticulus*. **A. Pulmonum**, the lobes of the lung. **A. of Sacrum**, the flat triangular surface of bone extending outward from the base of the sacrum, supporting the psoas magnus muscle. **A. Ves-pertilionis**, the broad ligament of the uterus. **A. Vulvæ**, the labia of the pudendum.

Alalia (*al-a'-le-ah*) [*â* priv.; *λάλει*, talk]. Impairment of articulation from paralysis of the muscles of speech or from local laryngeal disease.

Alantin (*al-an'-tin*). Same as *Inulin*.

Alanthol (*al-an'-thol*), $C_{10}H_{12}O$. Inulol. An aromatic liquid obtained from elecampane; used like creosote in pulmonary tuberculosis.

Alar (*al'-lar*) [*ala*, wing]. Winglike. **A. Ligaments**, lateral synovial folds of the ligament of the knee-joint. **A. Odontoid Ligaments**, lateral ligaments of the odontoid process.

Alate (*al'-lāt*) [*ala*, a wing]. Winged.

Albedo (*al-be'-dō*) [L., "whiteness"]. Whiteness. **A. retinæ**, retinal edema. **A. unguis**, the lunula of the nail.

Albescent (*al-bes'-ent*) [*albescere*, to become white]. Whitish.

Albicans (*al'-be-kans*) [*albicare*, to grow white]. White; whitish.

Albican-tia (*al-be-kan'-she-ah*) [L.]. Plural of *Albicans*.

Albinism, **Albinismus** (*al'-bin-izm*, *al-bin-iz'-mus*) [*albus*, white]. *Congenital Achro-ma*; *Congenital Leukoderma*. That condition of the skin in which there is a congenital absence of pigment involving its entire surface, including the hair and the choroid coats and irides of the eyes. It is usually associated with nystagmus, photophobia, and astigmatism.

Albino (*al-bi'-nō*) [Sp.]. A person affected with albinism.

Albinotic (*al-bin-ot'-ik*) [Sp.]. Affected with albinism.

Albolene (*al'-bo-len*) [*albus*, white; *oleum*, oil]. A hydrocarbon oil, colorless, tasteless, odorless, used as an application to inflamed surfaces.

Albuginea (*al-bu-jin'-e-ah*) [*albus*, white]. White, or whitish. **A. oculi**, the sclerotic coat of the eye. **A. ovarii**, and **testis**, the tunica albuginea of the ovary and testicle.

Albugineous (*al-bu-jin'-e-us*) [*albus*, white]. Whitish.

Albumen (*al-bu'-men*) [*albus*, white]. The white of an egg. See *Albumin*.

Albumimeter (*al-bu-mim'-et-er*) [*albumen*; *μέτρον*, measure]. An instrument for the quantitative estimation of albumin in urine.

Albumin (*al-bu'-min*) [*albus*, white]. A proteid substance, the chief constituent of the animal tissues. Its molecule is highly complex. It is soluble in water and coagulable by heat. It contains the following elements: Carbon 51.5 to 54.5; hydrogen 6.9 to 7.3; nitrogen 15.2 to 17.0; oxygen 20.9 to 23.5; sulphur 0.3 to 2.0. Its approximate formula is $C_{73}H_{113}N_{16}O_{22}S$. **Albumen**, white of egg, often called albumin, is largely composed of it. Other varieties are called after their sources or characteristic reactions, as acid-albumin, alkali-albumin, muscle-, serum-, ovum-, vegetable-albumin, etc. **A., Blood**. See *Serum-albumin*. **A., Caseiform**, that variety not coagulated by heat, but precipitated by acids. **A., Circulating**, that found in the fluids of the body. **A., Derived**, a modification of albumin resulting from the action of certain chemicals upon native albumin. **A., Muscle-**, a variety found in muscle-juice. **A., Native**, any albumin occurring normally in the tissues. **A., Organic**, that forming an integral part of the tissue. **A., Serum-**. See *Serum-albumin*. **A., Vegetable**, that found in various vegetable juices.

Albuminate (*al-bu'-min-āt*) [*albus*, white]. A compound of albumin and certain bases, as albuminate of iron.

Albuminiferous (*al-bu-min-īf'-er-us*) [*albumin*; *ferre*, to bear]. Yielding albumin.

Albuminimeter (*al-bu-min-im'-et-er*). See *Albumimeter*.

Albuminogenous (*al-bu-min-ōj'-en-us*) [*albumin*; *γεννᾶν*, to produce]. Producing albumin.

Albuminoid (*al-bu'-min-oid*) [*albumin*; *είδος*, form]. 1. Resembling albumin. Applied to certain compounds having many of the characteristics of albumin. 2. Any nitrogenous principle of the class of which normal albumin may be regarded as the type. **A. Disease**. See *Amyloid Degeneration*.

Albuminometer (*al-bu-min-om'-et-er*). See *Albumimeter*.

Albuminone (*al-bu'-min-ōn*) [*albumin*]. A principle derived from certain albuminoids; it is soluble in alcohol, and is not coagulable by heat.

Albuminorrhæa (*al-bu-min-or-e'-ah*) [*albumin*; *ροια*, a flow]. Excessive discharge of albumins.

Albuminose (*al-bu'-min-ās*) [*albumin*]. 1. A product of the digestion of fibrin or any

albuminoid in very dilute hydrochloric acid; acid albumin. 2. Albumose, or one of the products of the digestion of albumin by the gastric juice.

Albuminous (*al-bu'-min-us*) [*albumin*]. Containing, or of the nature of, albumin.

Albuminuria (*al-bu-min-u'-re-ah*) [*albumin*; *oïpov*, urine]. The presence in the urine of albumin, usually serum-albumin. Albumin in the urine may result from disease of the kidneys or from the admixture of blood or pus to the urine. Its presence is sometimes not accounted for by either of these causes. See *A.*, *Cyclic*. **A. of Adolescence.** See *A.*, *Cyclic*. **A. Adventitious.** See *A.*, *Pseudo*. **A. Cardiac**, that due to chronic valvular disease. **A. Cyclic**, a condition also known as physiologic, simple, functional, or transient albuminuria, or the albuminuria of adolescence, in which a small quantity of albumin appears in the urine, especially of the young, at stated times of the day; hence the term "cyclic." The nature of this phenomenon is not positively known, but it is probably the result of some changes in the renal circulation. **A.**, *Dietetic*, that due to the ingestion of certain forms of food. **A.**, *False*, a mixture of albumin with the urine during its transit through the urinary passages, where it may be derived from blood, pus, or special secretions that contain albumin. **A.**, *Febrile*, that due to fever, or associated with acute infectious diseases, slight changes occurring in the glomeruli without organic lesion. **A.**, *Functional*. See *A.*, *Cyclic*. **A.**, *Gouty*, albumin in the urine of elderly persons, who secrete a rather dense urine containing an excess of urea. **A.**, *Intrinsic*. See *A.*, *True*. **A.**, *Mixed*, the presence of a true with a pseudo-albuminuria. **A.**, *Nephrogenous*, that due to renal disease. **A.**, *Neurotic*, that due to nervous disorders. **A.**, *Paroxysmal*. See *A.*, *Cyclic*. **A.**, *Physiologic*, the presence of albumin in normal urine, without appreciable coexisting renal lesion or diseased condition of the system. **A.**, *Pseudo*-. *A.*, *Adventitious*; albuminuria dependent upon the presence of such fluids as blood, pus, lymph, spermatic fluid, or the contents of an abscess cavity, in the urine. **A.**, *Simple*. See *A.*, *Cyclic*. **A.**, *Temporary*, that lasting but for a short time. **A.**, *Transient*. See *A.*, *Cyclic*. **A.**, *True*, *A.*, *Intrinsic*; that due to the excretion of a portion of the albuminous constituents of the blood with the water and salts of the urine.

Albuminuric (*al-bu-min-u'-rik*) [*albumin*; *oïpov*, urine]. Associated with, of the nature of, or affected by, albuminuria.

Albumose (*al'-bu-môs*) [*albumin*]. Any

albuminoid substance ranking among the first products of the splitting up of proteids by enzymes, and intermediate between the food-albumins and the typical peptones. According to Kühne there are at least two albumoses, *antialbumose* and *hemialbumose*. The latter yields the following: *Protalbumose*, *deuteroalbumose*, *heteroalbumose*, and *dysalbumose*.

Albumosuria (*al-bu-môs-u'-re-ah*) [*albumose*; *oïpov*, urine]. The presence of albumose in the urine.

Alchemy (*al'-kem-e*) [Arab., of doubtful derivation]. The supposed art of the transmutation of metals (into gold) and of finding a remedy for all diseases.

Alcohol (*al'-ko-hol*) [Arab., *al-koh'l*, the fine powder for staining eyelids]. 1. Any compound of an organic hydrocarbon radicle with hydroxyl. Alcohols are classed as *monacid* (monatomic), *diacid* (diatomic), and *triacid* (triatomic), according to the number of hydroxyl radicles present in the molecules. 2. Ethyl alcohol, C_2H_5O . A liquid obtained by the distillation of fermented grain or starchy substance. It is inflammable, colorless, and possesses a pungent odor and burning taste. Internally, it is a cerebral excitant and cardiac stimulant; in large doses a depressant, narcotic poison, producing muscular incoordination, delirium, and coma. It exists in wine, whisky, brandy, beer, etc., and gives to them their stimulant properties. Commercial alcohol contains 91 per cent. of absolute alcohol, with 9 per cent. of water. It is valuable as a cardiac stimulant in acute failure of the heart's action and in adynamic conditions. **A.**, *Benzyl*, C_7H_5O , obtained from benzaldehyd by the action of sodium amalgam. **A.** *deodoratum*, deodorized alcohol, containing 92.5 per cent. by weight of ethyl alcohol and 7.5 per cent. of water. **A.**, *Dilute*, composed of 41 per cent., by weight, of alcohol. **A.**, *Ethyl*. See *Alcohol*. **A.**, *Methyl*, CH_3O , commonly known as "Wood Spirit." **A.**, *Phenic*. See *Acid*, *Carbolic*. **A.**, *Primary*, **A.**, *Secondary*, **A.**, *Tertiary*, an alcohol produced by the replacement of one, two, or three hydrogen atoms in carbinol by alkyls. **A.**, *Unsaturated*, those derived from the unsaturated alkylens in the same manner as the normal alcohols are obtained from their hydrocarbons. In addition to the general character of alcohols, they are also capable of directly binding two additional affinities.

Alcoholate (*al'-ko-hol-ât*) [see *Alcohol*]. 1. A chemic compound, as a salt, into which an alcohol enters as a definite constituent. 2. A preparation made with alcohol.

Alcoholature (*al'-ko-hol-at-chur*) [Fr. *alcoholature*]. An alcoholic tincture.

Alcoholic Radicle, the name applied to the hydrocarbon radicle that unites with hydrogen and oxygen to form an alcohol.

Alcoholica (*al-ko-hol'-ik-ah*) [see *Alcohol*]. In pharmacy, alcoholic preparations.

Alcoholimeter (*al-ko-hol-im'-it-er*). See *Alcoholometer*.

Alcoholism (*al'-ko-hol-izm*) [see *Alcohol*]. The morbid results of excessive or prolonged use of alcoholic liquors. The term **acute A.** has been used as a synonym for inebriety. The **chronic** form is associated with severe disturbances of the digestive and nervous systems.

Alcoholization (*al-ko-hol-iz-a'-shun*) [see *Alcohol*]. The art or process of alcoholizing; the state of being alcoholized; the product of the process of alcoholizing.

Alcoholize (*al'-ko-hol-iz*) [see *Alcohol*]. 1. To impregnate with alcohol. 2. To convert into an alcohol.

Alcoholometer (*al-ko-hol-om'-et-er*) [*alcohol*; *μέτρον*, a measure]. An hydrometer or other instrument used in determining the percentage of alcohol in any liquid.

Alcoholophilia (*al-ko-hol-o-fil'-e-ah*) [*alcohol*; *φιλέω*, to love]. The appetite for strong drink; a craving for intoxicants.

Aldehyd (*al'-de-hid*) [*al*, the first syllable of alcohol; *dehyd*, from *dehydratus*]. 1. A class of compounds intermediate between alcohols and acids, derived from their corresponding primary alcohols by the oxidation and removal of two atoms of hydrogen, and converted into acids by the addition of an atom of oxygen. They contain the group COH. 2. C_2H_4O . Alcohol deprived of two atoms of hydrogen, or *acetic aldehyd*. It is a colorless, limpid liquid, with a characteristic odor. **A.**, Anisic, $C_9H_8O_2$, results on oxidizing various essential oils (anise, fennel, etc.) with dilute HNO_3 . **A.**, Benzoic, C_7H_6O . *Benzaldehyd*, the oil of bitter almonds. **A.**, Cinnamic, C_9H_8O , the chief ingredient of the essential oil of cinnamon and cassia. **A.**, Formic, CH_2O , *Formaldehyd*, is said to be microbicidal and antiseptic. See *Formaldehyd*.

Ale (*al'*) [*A. S.*, *eolw*]. An alcoholic beverage brewed from malt and hops. It contains from three to seven per cent. of alcohol.

Alecithal (*ah-les'-ith-al*) [*ἀ* priv.; *λεκιθος*, yolk]. A term applied to certain ova having the food-yolk absent, or present only in very small quantity.

Alembic (*al'-em'-bik*) [*Arab.*, *al*, the; *ἀμβίξ*, a cup]. A vessel used for distillation.

Alembroth (*al'-em'-broth*) [origin unknown]. An old name for a compound of the chlorid of ammonium and mercury. Its solution has been used as an antiseptic.

Aleppo Boil, Button, Evil, Pustule, or Ulcer. See *Furunculus orientalis*.

Aletris (*al'-et-ris*). Star Grass, Unicorn Root, Starwort, Colic Root. The root of *A. farinosa*. It is tonic, diuretic, and anhelminthic, and was formerly a popular domestic remedy in colic, dropsy, and chronic rheumatism. Dose of fld. ext. $\mathfrak{m}\text{x}-\text{xxx}$ (0.65-2.0); of tincture (1 in 8 proof spirit) \mathfrak{z} i-ij (4.0-8.0); of *aletrin*, the extractive, gr. \mathfrak{z} i-iv (0.016-0.26). **A. Cordial**. Dose \mathfrak{z} j (4.0) three times daily. Unof.

Aleukemia (*ah-lu-ke'-me-ah*) [*ἀ* priv.; *λευκός*, white; *αἷμα*, blood]. Deficiency in the proportion of white cells in the blood.

Aleurometer (*al-u-rom'-et-er*) [*ἀλετρον*, flour; *μέτρον*, a measure]. An instrument used for the examination of crude gluten as to its power of distending under the influence of heat, as a means of judging of the value of a flour for bread-making.

Aleuronat (*al-u'-ro-nat*) [*ἀλετρον*, flour]. A vegetable albumin used as a substitute for bread in cases of diabetes.

Aleuroscope (*al-u'-ro-skop*) [*ἀλετρον*, flour; *σκοπεῖν*, to view]. See *Aleurometer*.

Alexanderism (*al-eks-an'-der-izm*) [*Alexander*, the Macedonian King]. The insanity of conquest.

Alexia (*al-eks'-e-ah*) [*ἀ* priv.; *λέξις*, word]. Word-blindness. A form of aphasia, in which the patient is unable to recognize written or printed characters. **A.**, Musical, musical blindness; loss of the ability to read music.

Alexin (*al-eks'-in*) [*ἀλέξις*, help]. A defensive proteid existing normally in the blood.

Alexipharmac, or **Alexipharmic** (*al-eks-e-far'-mak*, or *-mik*) [*ἀλέξιν*, to repel; *φάρμακον*, a poison]. A medicine neutralizing a poison.

Alexipharmacon (*al-ekt-e-far'-mak-on*) [*ἀλέξιν*, to repel; *φάρμακον*, a poison]. Any alexipharmac medicine.

Algæ (*al'-je*) [*alga*, a seaweed]. A group of acotyledonous plants, living mostly in the water.

Algaroth (*al'-gar-oth*) [*It.*, *Algarotti*, the Veronese physician]. Oxychlorid of antimony.

Algesia (*al-je'-se-ah*) [*ἀλγος*, pain]. Pain; suffering.

Algesimeter (*al-jes-im'-et-er*) [*ἀλγος*, pain; *μέτρον*, a measure]. An instrument for determining the acuteness of the sense of pain. **A.**, Boas', an instrument consisting of a pad and spring, used to determine the relative sensitiveness over the epigastrium. The normal tolerance is 9 to 10 kilograms. In cases of gastric ulcer, 1 to 2 kilograms.

Algesthesis, or **Algæsthesis** (*al-jes-the'-sis*) [*ἀλγος*, pain; *αἰσθησις*, feeling]. The perception of pain; painful disease.

Algid (*al'-jid*) [*algidus*, cold]. Cold, chilly.

A. Fever, a pernicious intermittent fever, with great coldness of the surface of the body.

A. Cholera, the cold stage of Asiatic cholera.

Algotometer (*al-gom'-et-er*) [*άλγος*, pain; *μέτρον*, a measure]. An instrument for testing the sensibility of a part to pain.

Algor (*al'-gor*) [L.]. A sense of chilliness or coldness.

Alible (*al'-i-bl*) [*alibilis*, nutritive]. Nutritive; absorbable and assimilable.

Alienation (*al-yen-a'-shun*) [*alienus*, strange]. Mental derangement.

Alienist (*al'-yen-ist*) [*alienus*, strange]. One who treats mental diseases.

Aliform (*al'-if-orm*) [*ala*, wing; *forma*, shape]. Wing-shaped. **A. Process**, the wing of the sphenoid.

Aliment (*al'-im-ent*) [*alimentum*, from *alimentare*, to nourish]. Nourishment, food.

Alimentary (*al-im-en'-ta-re*) [*alimentum*, from *alimentare*, to nourish]. Nourishing.

A. Bolus, the food after mastication and just prior to swallowing. **A. Canal, Duct, System**, or **Tube**, the digestive tube, from the lips to the anus, with its accessory glands.

Alimentation (*al-im-en-ta'-shun*) [*alimentare*, to nourish]. The act of supplying with food. **A., Rectal**, the nourishing of a patient by the administration of small quantities of concentrated food through the rectum. There are many formulæ for these nutritive enemata, the most important of which are the following: 1. *Leube's Pancreatic-meat Emulsion*.—"For rectal alimentation: chop five ounces of finely-scraped meat still finer, add to it one and a half ounces of finely-chopped pancreas, free from fat, and then three ounces of lukewarm water; stir to the consistence of a thick pulp. Give at one time, care being taken to wash out the rectum with water about an hour before." 2. *Mayet's Formula*.—"Fresh ox-pancreas 150 to 200 grams; lean meat 400 to 500 grams. Bruise the pancreas in a mortar with water at a temperature of 37° C., and strain through a cloth. Chop the meat and mix thoroughly with the strained fluid, after separating all the fat and tendinous portions. Add the yolk of one egg, let stand for two hours, and administer at the same temperature, after having cleansed the rectum with an injection of oil. This quantity is sufficient for 24 hours' nourishment, and should be administered in two doses." 3. *Rennie's Formula*.—"Add to a bowl of good beef-tea one-half pound of lean, raw beefsteak pulled into shreds. At 99° F. add one dram of fresh pepsin and one-half dram of dilute HCl. Place the mixture before the fire and let it remain for four hours, stirring frequently. The heat must not be too great, or the artificial digestive process will be

stopped altogether. It is better to have the mixture too cold than too hot. If alcohol is to be given it should be added at the last moment. Eggs may also be added, but should be previously well beaten." 4. *Peaslee's Formula*.—"Crush or grind a pound of beef-muscle fine; then add one pint of cold water; allow it to macerate 40 minutes, and then gradually raise it to the boiling point; and then it to boil two minutes—no more—and then strain." 5. *Flin's Mixture*.—"Milk $\frac{3}{4}$ ij, whisky $\frac{3}{4}$ ss, to which add half an egg."

Alinasal (*al-in-a'-sal*) [*ala*, a wing; *nasus*, the nose]. Pertaining to the *ala nasi*, or wing of the nose.

Alisphenoid (*al-is-fe'-noid*) [*ala*, a wing; *σφήν*, a wedge; *ειδος*, form]. 1. Pertaining to the greater wing of the sphenoid bone. 2. The bone that in adult life forms the main portion of the greater wing of the sphenoid.

Alkalinescence (*al-ka-les'-ens*) [Arab., *al-qalt*, soda-ash]. Slight or commencing alkalinity.

Alkalinescent (*al-ka-les'-ent*) [Arab., *al-qalt*, soda-ash]. Somewhat alkaline.

Alkali (*al'-ka-li*) [Arab., *al-qalt*, soda-ash]. The term includes the hydrated oxides of the alkali metals; these are electro-positive, are strong bases, uniting with acids to form salts, turn red litmus blue, and saponify fats. **A., Caustic**, the solid hydroxid of potassium or sodium. **A., Fixed**, potassium or sodium hydrate. **A. Metals**, sodium, potassium, lithium, cesium, and rubidium. **A., Volatile**, ammonium hydrate, which is decomposed by heat with the evolution of ammonia.

Alkali-albumin (*al'-ka-li-al-bu'-min*). A derived albumin; a proteid that has been acted upon by dilute alkalies and yields an alkaline reaction.

Alkalimeter (*al-ka-lim'-et-er*) [*alkali*; *μέτρον*, a measure]. An instrument for estimating the alkali in a substance.

Alkalimetry (*al-ka-lim'-et-re*) [*alkali*; *μέτρον*, a measure]. The measurement of the amount of an alkali in a substance.

Alkaline (*al'-ka-lin*) [*alkali*]. Having the qualities of or pertaining to an alkali. **A. Earths**, the oxides of calcium, barium, strontium, and magnesium.

Alkalinity (*al-ka-lin'-i-te*) [*alkali*]. The quality of being alkaline.

Alkalinuria (*al-ka-lin-u'-re-ah*) [*alkali*; *ούρον*, urine]. Alkalinity of the urine.

Alkalization (*al-ka-li-za'-shun*) [*alkali*]. The act of rendering a thing alkaline; the state or quality of being rendered alkaline.

Alkaloid (*al'-ka-loid*) [*alkali*; *ειδος*, likeness]. Any one of the nitrogeaneous compounds occurring in plants and resembling ammonia in being basic and capable of forming salts with acids. They are believed

to be substituted ammonias. Several of the important alkaloids are related to piperidin and to quinolin. Alkaloids are, as a rule, the most active parts of plants; many are used in medicine. **A.**, **Animal**, substances chemically like alkaloids, formed in the decomposition of animal tissues. **A.**, **Cadaveric**, or **Putrefactive**. See *Ptomains*.

Alkanet (*al'-kan-et*) [Sp., dim. of *alkanna*, henna]. The root of the herb, *Alkanna* (*Achusa*) *tinctoria*, yielding a red dye that is used in staining wood, coloring adulterated wines, and in pharmacy to give a red color to salves, etc.

Alkanna Red. See *Alkannin*.

Alkannin (*al'-kan-in*) [Sp., dim. of *alkanna*, henna]. *Alkanna* red; a valuable coloring matter obtained from alkanet.

Alkapton (*al-kap'-ton*) a yellowish, resinous, nitrogenous body occasionally found in urine.

Alkaptonuria (*al-kap-ton-u'-re-ah*) [*alkap-ton*; *oûr*, urine]. The presence of alkapton in the urine. It has been found in cases of pulmonary tuberculosis and in other instances in which there were no local lesions or general disease. Urine containing alkapton turns dark on standing or on the addition of an alkali.

Alkarsin (*al-kar'-sin*) [*alcohol*; *arsenic*]. Cadet's Fuming Liquid; an extremely poisonous liquid containing cacodyl. It is of a brown color, and on exposure to the air ignites spontaneously.

Alkyls (*al'-kilz*) [*alkali*]. The name applied to the univalent alcohol radicles, C_nH_{2n+1} ; methyl, ethyl, etc., are alkyls.

Alkyl-sulphids (*al'-kil-sul'-fids*). Thioethers. Sulphur analogues of the ethers. They are colorless liquids, generally insoluble in water, and possessing a disagreeable odor resembling that of garlic.

Allantiasis (*al-an-ti'-as-ís*) [*állās*, a sausage]. Sausage poisoning, due to the ingestion of sausages in which putrefactive changes have taken place.

Allantoic (*al-an-to'-ik*) [*állās*, a sausage; *eîdos*, resemblance]. Pertaining to the allantoids.

Allantoid (*al-an'-to'id*) [*állās*, sausage; *eîdos*, resemblance]. 1. Resembling a sausage. 2. Relating to the allantoids.

Allantoin (*al-an'-to-in*) [*állās*, a sausage; *eîdos*, resemblance], $C_4H_6N_2O_3$. A crystalline substance occurring in traces in normal urine, and prepared from uric acid by oxidation. Also the characteristic constituent of the allantoid fluid, and likewise found in fetal urine and amniotic fluid.

Allantois (*al-an'-to-ís*) [*állās*, a sausage; *eîdos*, resemblance]. One of the fetal membranes, derived from the mesoblastic and

hypoblastic layers. Its function is to convey the blood-vessels to the chorion. The lower part finally becomes the bladder, the upper, the urachus.

Allantotoxicon (*al-an-to-toks'-ik-on*) [*állās*, a sausage; *τοξικόν*, a poison]. A poisonous substance, probably a ptomain, that develops during the putrefactive fermentation of sausage.

Allesthesia (*al-es-the'-se-ah*) [*állas*, other; *αἴσθησις*, feeling]. Same as *Allochiria*.

Alleviator (*al-e'-ve-a-tor*) [*allevare*, to lighten]. A device for raising or lifting a sick person from the bed.

Alliaceous (*al-e-a'-shus*) [*allium*, garlic]. Resembling garlic, or pertaining to the same.

Alliteration (*al-it-er-a'-shun*) [*ad*, to; *littera*, letter]. A form of dysphasia in which the patient arranges his words according to the sound.

Allium (*al'-e-um*) [L.]. Garlic. The bulb of *A. sativum*. It contains a pungent volatile oil (allyl sulphid), that is found also in the leek and the onion. The tubers are used both as a food and as a condiment, and are stimulants to digestion. **A. cepa**, the common onion, and **A. porrum**, the leek, have similar qualities. **A.**, **Syr.**, contains, fresh garlic 20 gm., sugar 80 gm., dil. acetic acid, a sufficient quantity to make 100 c.c. Dose 3j-5 iv (4.0-16.0).

Allocheiria, or **Allochiria** (*al-o-ki'-re-ah*) [*állas*, other; *χείρ*, hand]. A peculiar disturbance of sensation in which, if one extremity be pricked, the patient locates the sensation in the corresponding part on the other side.

Allochesesthesia, **Allochæsthesia** (*al-ok-the'-se-ah*). Same as *Allochiria*.

Allolalia (*al-o-la'-le-ah*) [*állas*, other; *λαλεῖν*, to speak]. Any perversion of the faculty of speech.

Allopathy (*al-op'-a-the*) [*állas*, other; *πάθος*, affection]. According to Hahnemann, the inventor of the term, that method of the treatment of disease consisting in the use of medicines the action of which upon the body in health produces morbid phenomena different from those of the disease treated. Opposed to homeopathy.

Allorhythmia (*al-o-rith'-me-ah*) [*állas*, other; *ῥυθμός*, rhythm]. Variation in intervals of the pulse.

Allotoxin (*al-o-toks'-in*) [*állas*, other; *τοξικόν*, poison]. Any substance, produced by tissue metamorphosis within the organism, that tends to shield the body by destroying microorganisms or toxins that are inimical to it.

Allotriodontia (*al-at-re-o-don'-the-ah*) [*ἀλλότριος*, strange; *ὀδούς*, tooth]. 1. The transplanting of teeth from one person to another.

2. The existence of teeth in abnormal situations, as in tumors.

Allotriogeustia (*al-ot-re-o-gūs'-te-ah*) [ἀλλότριος, strange; γεῖναι, taste]. Perversion of the sense of taste; abnormality of the appetite.

Allotriophagy (*al-ot-re-off'-a-je*) [ἀλλότριος, strange; φαγεῖν, to eat]. Depraved or unnatural appetite.

Allotropic (*al-o-trop'-ic*) [ἄλλος, different; τρέπος, turn]. Characterized by allotropism.

Allotropism (*al-ot'-ro-pism*) [ἄλλος, other; τρέπος, turn]. 1. The term expresses the fact of certain elements existing in two or more conditions with differences of physical properties. Thus carbon illustrates allotropism by existing in the forms of charcoal, plumbago, and the diamond. 2. Appearance in an unusual or abnormal form.

Alloxan (*al-oks'-an*) [*allantoin*; *oxalic*], $C_4H_2N_2O_6$. A crystalline substance produced by the oxidation of uric acid.

Alloxantin (*al-oks-an'-tin*) [*allantoin*; *oxalic*], $C_8H_4N_4O_8 + 3H_2O$. A substance obtained by reducing alloxan with $SnCl_2$, zinc, and HCl, or H_2S in the cold. It occurs in small, hard, colorless prisms that turn red when treated with ammonia.

Alloy (*al-oi'*) [from the French word *aloi*, a contraction of *à la loi*]. 1. A compound of two or more metals by fusion. 2. The least valuable of two or more metals that are fused together.

Allspice (*awl'-spis*). See *Pimenta*, and *Florida Allspice*.

Allyl (*al'-il*) [*allium*, garlic], C_3H_5 . A univalent alcohol-radicle. **A.** Alcohol, C_3H_5HO , a colorless, inflammable liquid, with pungent odor, boiling at $97^\circ C$. **A.** Aldehyde, C_3H_4O , a syn. of *Acrolein*. **A.** Cyanid. See *Sinamin*. **A.** Mustard Oil, $CS.N.C_3H_5$, Allyl thiocarbimid. The principal constituent of ordinary mustard oil. **A.** sulphid, $(C_3H_5)_2S$, the essential oil of garlic. **A.** thiocarbimid. See *A. Mustard Oil*. **A.** tribromid, $C_3H_5Br_3$, a colorless liquid used as an antispasmodic. Dose, five drops (0.15). Unof.

Allylamin (*al-il'-am-in*) [*allium*, garlic; *ammonia*], $NH_2(C_3H_5)$. Ammonia in which a hydrogen atom is replaced by allyl. It is a caustic liquid.

Almond (*am'-und*) [ME., *almonde*]. See *Amygdala*. **A.** Bread, a variety of bread made from almond flour, for use in diabetes as a substitute for ordinary bread. **A.-eyed**, applied to the Mongolian race on account of the peculiar elliptical form and slanting position of the eyes. **A. of the Ear**, the tonsil. **A.** Mixture. See *Amygdalæ Mixture*. **A.** Oil, oleum amygdalæ, the fixed oil

obtained from both the sweet and the bitter almond. It is odorless, agreeable to the taste, and of a yellow color; sp. gr. .919 at $15^\circ C$. **A.** Oil, Bitter, oleum amygdalæ amaræ. See *Amygdala*. **A.** Paste, a magma of bitter almonds, alcohol, white of egg, and rose-water, used to soften the skin and prevent the hands and lips from chapping. **A. of the Throat**, the tonsil.

Alnus (*al'-nus*) [L.]. Alder Bark. The bark of the American Alder. *A. serrulata*, contains tannic acid. The decoction of bark and leaves is astringent, and used as a gargle and as a lotion for wounds and ulcers. Dose of powdered bark gr. x (0.65).

Alochia (*ah-lo'-ke-ah*) [*ā* priv.; λoχία, the lochia]. Absence of the lochia.

Aloe (*al'-o*) [ἀλόη, the aloe]. The inspissated juice of several species of aloe, of which the *A. socotrina*, *A. barbadensis*, and *A. capensis* are most commonly used. Its properties are due to a glucosid, *aloin*, $C_{17}H_{18}O_8$. It is a tonic astringent, useful in amenorrhea, chronic constipation, and atonic dyspepsia. It is also an emmenagogue and anthelmintic. Dose gr. j-v (0.065-0.32). **A. americana**. See *Agave*. **A.** Decoct., Comp. (B. P.). Dose gr. ss-ij (0.032-0.13). **A.** Enema (B. P.), aloes, carbonate of potassium, and mucilage of starch. **A. et Asafoetida**, Pil., aloes and asafoetida aa gr. $1\frac{1}{2}$ (0.1). **A. et Canella**, Pulv., contains powdered bark of Canella alba. Dose gr. v-xx (0.32-1.3). **A. et Ferri**, Pil., contains gr. j each of aloes, ferrous sulphate, and aromatic powder, incorporated with confection of roses. **A. et Mastich**, Pil., "Lady Webster's" pill, contains aloes gr. ij (0.13), mastich and red rose gr. ss (0.032). **A. et Myrrh.**, Pil., contains each aloes gr. ij (0.13), myrrh gr. j (0.065), aromatic powder gr. ss (0.032), mixed with syrup. **A. et Myrrh.**, Tinct., aloes 10, myrrh 10, alcohol 100 parts. Dose ʒss-ʒij (2.0-8.0). **A.** Ext., Aquosum, prepared by mixing aloes 1 part with 10 parts boiling water, straining and evaporating. Dose gr. ss-v (0.032-0.32). **A.** Hepatic, dark liver-colored aloes, mostly Socotrine or Barbadian. **A.** Pil., aloes and soap aa gr. ij (0.13). **A. Purificata**, the common drug purified by solution in alcohol and evaporation. Dose gr. j-v (0.065-0.32). **A. Socotrin.**, Pil. (B. P.), contains socotrine aloes, hard soap, oil of nutmeg, and confection of roses. Dose gr. v-x (0.32-0.65). **A.** Tinct., consists of aloes 10, glycyrrhiza 10, dil. alcohol 100 parts. Dose ʒss-ʒij (2.0-8.0). **A.** Vin., has aloes 6, cardamon 1, ginger 1, white wine 100 parts. Dose ʒj-ʒiv (4.0-16.0). **Aloeretin** (*al-o-e-re'-tin*) [ἀλόη, aloes; ῥητινῇ, resin]. A brown resin contained in aloes.

Aloetic (*al-o-et'-ik*) [*ἀλόη*, the aloe]. Containing or pertaining to aloes.

Aloetin (*al-a-et'-in*). 1. Aloeretin. 2. A yellow crystalline principle obtainable from aloes.

Aloin (*al'-o-in*) [*ἀλόν*, the aloe]. A bitter principle found in aloes. It forms fine needles, possesses a very bitter taste, and acts as a strong purgative. Several glucosids of this name are described; as *Barbaloin*, *Nataloin*, *Zamaloïn*, *Socaloin*. Dose gr. ss-ij (0.032-0.13).

Alopecia (*al-o-pe'-she-ah*) [*ἀλωπηξ*, a fox]. Deficiency of hair; baldness. It may be universal or partial, congenital or acquired. It follows a large number of systemic affections. **A. adnata**. See *A. Congenital*. **A. areata**; *Area celsi*; *Tinea decalvans*; *Porrigio decalvans*; *Alopecia circumscripta*; that condition in which suddenly or slowly one or several, usually asymmetrically distributed, patches of baldness appear upon the hairy regions of the body, more often upon the scalp and parts covered by the beard.

A. circumscripta. See *A. areata*. **A. Congenital**, a rare form, seldom complete, due to absence of hair-bulbs. **A. furfuracea**; *Alopecia pityroides capillitii*; *Pityriasis capitis*; *Seborrhœa capillitii*; *Pityriasis simplex*; *Dandruff*; a form of baldness associated with a disorder of the scalp, marked by hyperemia, itching, and exfoliation of dry or fatty scales from its surface. It may be acute or chronic, and produce a dryness, brittleness, and lack of luster in the hair. **A. localis**; *A. neuritica*; that form occurring in one or more patches at the site of an injury, or in the course of a nerve. **A. neuritica**. See *A. localis*. **A. neurotica**, a name given to baldness of trophoneurotic origin. **A. orbicularis**. Same as *A. circumscripta*. **A. pityroides capillitii**. See *A. furfuracea*. **A. pityroides universalis**, a rapid and general denudation of hair occurring in debilitated states, preceded by abundant desquamation of fatty scales. **A. senilis**, that occurring in old age. **A. simplex**, the idiopathic premature baldness of young adults. It is most common in males, and is often associated with premature grayness. **A. syphilitica**, that due to syphilis. **A. universalis**, that in which there is a general falling out of the hairs of the body.

Aloxanthin (*al-oks-an'-thin*), $C_{15}H_{10}O_8$. A yellow substance obtained from barbaloin and socaloin by the action of bichromate of potassium.

Alpha (*al'-fah*) [*ἀλφα*, the first letter of the alphabet]. The Greek letter *α*, used in combination with many chemist terms to indicate the first of a series, as alpha-naphthol.

Alphos (*al'-fos*) [*ἀλφός*, vitiligo]. 1. An old name for leprosy. 2. Psoriasis.

Alphosis (*al'-fo'-sis*) [*ἀλφός*, vitiligo]. Albinism; leukoderma.

Alpinia (*al'-pin'-e-ah*) [*Alpinus*, an Italian botanist of the sixteenth century]. A genus of zingiberaceous tropical plants. *A. officinarum*, *A. chinensis*, and other species furnish Galangal.

Alsace Gum (*al'-sas'-gum*). See *Dextrin*.

Alstonidin (*al'-ston'-id-in*) [*Alston*, a Scotch botanist]. An amorphous substance contained in a variety of Dita bark.

Alstonin (*al'-sto'-nin*) [*Alston*, a Scotch botanist]. 1. An amorphous substance contained in a variety of Dita bark. 2. A crystalline alkaloid, $C_{21}H_{20}N_2O_4$, obtained from *Alstonia Constricta*.

Alter (*awel'-ter*). To castrate or spay.

Alterant (*awel'-ter-ant*). Same as *Alterative*.

Alterative (*awel'-ter-a-tiv*) [*alter*, another]. A medicine that alters the processes of nutrition, restoring, in some unknown way, the normal functions of an organ, or of the system. The most important alteratives are arsenic, iodine, and the iodids, mercury, and gold.

Alternate (*awel'-ter-nāt*) [*alternare*, to do by turns]. Occurring successively in space or time. **A. Hemiplegia**. See *Hemiplegia*.

Alternating (*awel'-ter-na-ting*) [*alternare*, to do by turns]. Occurring successively. **A. Currents**, electric currents, the direction of which is constantly changing. **A. Insanity**, a form of insanity in which there are regular cycles of exaltation and depression.

Alternation (*awel'-ter-na'-shun*) [*alternare*, to do by turns]. Repeated transition from one state to another.

Althaus's Oil. It is made as follows: Metallic mercury 1 part, pure lanolin 4 parts, 2 per cent. carbolic oil 5 parts. It is used in the treatment of syphilis in injections of m.v (0.32) at a dose.

Althea, or **Althæa** (*al-the'-ah*) [L.]. Marsh-mallow. The peeled root of *A. officinalis*, a plant of the mallow family. It consists of about one-third of vegetable mucus and starch, together with the alkaloids *Asparagin* and *Althein* (latterly regarded as identical). Its decoction is employed as a mucilaginous drink. **A. Syrupus**, contains four per cent. althea. Dose indefinite. *Asparagin* possesses sedative and diuretic properties, and is useful in ascites and gout. Dose gr. ij-ijj (0.13-0.19).

Altitude (*al'-ti-tud*) [*altitudo*, height]. The elevation of a place above the sea-level.

Alum (*al'-um*) [*alumen*, alum]. Any one of a class of double sulphates formed by the union of one of the sulphates of certain non-alkaline metals with a sulphate of some alkaline metal. The standard (or common

commercial) alum, the official *Alumen*, U. S. P., is the aluminium-potassium sulphate, $\text{Al}_2(\text{SO}_4)_3 + \text{K}_2\text{SO}_4 + 24\text{H}_2\text{O}$. It is a powerful astringent and styptic, and is also extensively used in the arts. **A.**, *Ammonia*, is the same as the above, except that the potassium is replaced by ammonium. It is official in Great Britain, and is extensively used on account of its cheapness. What is known as *concentrated* or *patent* alum, is the normal aluminium sulphate (*aluminii sulphas*), which is not a true alum. **A.**, *Ammonio-ferric* (*Ferri et Ammonii sulphas*), U. S. P., is strongly styptic, and is useful in leukorrhea. Dose gr. v-x (0.32-0.65). **A.**, *Potassio-ferric*, is similar to the **A.**, ammonio-ferric. **A.**, *Soda*, double sulphate of sodium and aluminium, is too soluble for ordinary uses. **A.**, *Whey*, a preparation obtained by boiling 3 ij of alum in a pint of milk, and straining. It is used as an astringent and internal hemostatic, in wineglassful doses.

Alumen (*al-u'-men*) [L., *gen.*, *Aluminis*]. See *Alum*. **A.** *exsiccatum*, burnt or dehydrated alum.

Alumina (*al-u'-min-ah*) [L.], Al_2O_3 . Aluminium oxid; the principal ingredient of clay, and of many stones, earths, and minerals.

Aluminate (*al-u'-min-at*) [*alumina*]. A basic salt of alumina.

Aluminated Copper. See *Lapis divinus*.

Aluminiferous (*al-u-min-iff'-er-us*) [*alumen*, alum; *ferre*, to form]. Yielding alum.

Aluminium (*al-u-min'-e-um*) [L.]. $\text{Al} = 27$. Quantivalence II, IV. A silver white metal distinguished by its low sp. gr.,—about 2.6. It is largely used in the arts and for certain surgical instruments. **A.** *hydrate*, $\text{Al}_2(\text{HO})_6$, a tasteless white powder, feebly astringent. Dose gr. iij-xx (0.2-1.3). **A.** and *Potassium sulphate*, $\text{K}_2\text{Al}_2(\text{SO}_4)_4$ (*Alum*), a valuable astringent used in catarrh, leukorrhea, gonorrhea. Dose gr. x-xx (0.65-1.3). In teaspoonful doses it is an emetic. **A.** *sulphate*, $\text{Al}_2(\text{SO}_4)_3$, an antiseptic and astringent used as a lotion in 5 per cent. solution. **A.** *acetate*, a disinfectant. Unof.

Aluminol, **Alumnol** (*al-um-i'-nol*, *al-um'-nol*) [*aluminium*]. An astringent and antiseptic sulphur compound of aluminium, used in gonorrhea, endometritis, and diseases of the ear, nose, skin, etc. Unof.

Alum Root. The root of *Heuchera americana*. Its properties are due to gallic and tannic acids. It is very astringent. Dose of the fld. ext., gtt. x-xx (0.65-1.3). Also the root of *Geranium maculatum*, a mild astringent.

Alvegniat's Pump. See *Alvergniat's Pump*.

Alveolar (*al-ve'-o-lar*) [*alveolus*, a small hollow]. Pertaining to an alveolus. **A.** *Abscess*, a gum-boil. **A.** *Arch*, the alveolar surface of the jaw. **A.** *Artery*, a branch of the internal maxillary artery. **A.** *Border*, the margin of the jaws. **A.** *Index*, in craniometry, the gnathic index; the ratio of the distance between the basion and alveolar point, to the distance between the basion and the nasal point, multiplied by 100. (Sometimes the basilar index is called the alveolar index). **A.** *Passages*. The ultimate division of the bronchi, emptying into the infundibula. **A.** *Points*. See *Craniometric Points*. **A.** *Process*, the border of the superior maxilla in which the alveoli are placed. **A.** *Sarcoma*. See *Sarcoma*.

Alveolitis (*al-ve-o-li'-tis*) [*alveolus*, a hollow; *itis*, inflammation]. Inflammation of the alveolus of a tooth.

Alveolodental (*al-ve'-o-lo-den'-tal*) [*alveolus*, a hollow; *dens*, a tooth]. Pertaining to the teeth and their sockets.

Alveolus (*al-ve'-o-lus*) [L.]. 1. The bony socket of a tooth. 2. An air cell of the lung. **A.** of *Glands*, the ultimate sacs of a racemose gland. **A.** of the *Stomach*, one of the honeycomb-like depressions found in the stomach.

Alvergniat's Pump. A form of mercurial air-pump, used in estimating the gaseous constituents of the blood.

Alvine (*al'-vin* or *al'-vân*) [*alvus*, belly]. Pertaining to the belly. **A.** *Concretion*, an intestinal calculus. **A.** *Dejections*, **A.** *Discharges*, the feces. **A.** *Obstruction*. Constipation.

Alvus (*al'-vus*) [L.]. The belly, or its contained viscera.

Alymphia (*ah-limf'-e-ah*) [*à priv.*; *lymph*, lymph]. A deficiency of lymph.

Amadou (*am'-a-doo*) [Fr., *amadouer*, to coax]. German tinder or touchwood; *Boletus ignarius*, a fungus found on old tree-trunks, used to stanch local hemorrhage, and as a dressing of wounds, etc.

Amalgam (*am-al'-gam*) [*μάλαγμα*, a soft mass]. A combination of mercury with any other metal.

Amalgamation (*am-al-gam-a'-shun*) [*μάλαγμα*, a soft mass]. In metallurgy, the process of combining mercury with some other metal, as practised in separating silver and gold from ores.

Amanitin (*am-an'-it-in*) [*ἀμανίται*, a kind of fungus]. 1. A principle identical with cholin, obtained from the fly agaric. 2. A poisonous glucosid obtainable from various species of agaric.

Amara (*am-a'-ra*) [*amarus*, bitter]. Bitters. **Amasesis** (*ah-mas-e'-sis*) [*à priv.*; *μάσησις*, chewing]. Inability to chew the food.

Amastia (*ah-mas'-te-ah*) [*à* priv.; *μαστός*, breast]. Absence of the mammae or nipples.

Amativeness (*am'-at-iv-nes*) [*amare*, to love]. The sexual passion.

Amaurosis (*am-aw-ro'-sis*) [*ἀμαυρόειν*, to darken]. Blindness, especially that occurring without demonstrable lesion of the eye. **A.**,

Albuminuric, that due to renal disease. **A.**,

Cerebral, that due to disease of the brain. **A.**,

Congenital, that form existing from birth. **A.**,

Diabetic, that associated with diabetes. **A.**,

Hysteric, that associated with hysteria. **A.**,

Reflex, that resulting from a reflex action upon the optic nerve from some remote source of irritation. **A.**,

Saburral, sudden temporary blindness occurring in an attack of acute gastritis. **A.**,

Uremic, that due to uremia.

Amaurotic (*am-aw-ro'-tik*) [*ἀμαυρόειν*, to darken]. Relating to, or affected with amaurosis. **A.** **Cats-eye**, a light-reflex through the pupil in suppurative choroiditis.

Amazia (*ah-ma'-ze-ah*) [*à* priv.; *μαστός*, the breast]. Absence of the mammary gland.

Amber (*am'-ber*). See *Succinum*.

Ambergris (*am'-ber-gris*) [*amber*; Fr., *gris*, gray]. A substance excreted by the sperm whale, *Physeter macrocephalus*. It is not known whether it is a pathologic product or not. It exhales a fragrant, musky odor when warmed, and is used in adynamic fevers, chronic catarrh, and nervous diseases.

Dose, gr. j-ij (0.065-0.19). Unof.

Ambidexter (*am-be-dek'-ter*) [*ambo*, both; *dexter*, the right hand]. An ambidextrous person.

Ambidextrous (*am-be-dek'-trus*) [*ambo*, both; *dexter*, the right hand]. Able to use both hands equally well.

Amblyopia (*am-ble'-o'-pe-ah*) [*ἀμβλῆς*, dulled; *ὤψ*, eye]. Dimness of vision, especially that not due to refractive errors or organic disease of the eye. It may be *congenital* or *acquired*, being due, in the latter case, to the use of tobacco (*A. nicotinic*), or alcohol, or to other toxic influences; to *traumatism*; or it may be *hysteric*. *Nyctalopia* and *hemeralopia* are other forms; it may arise from *entoptic phenomena*, such as *musca volitans*, *micropsia*, *megalopsia*, *metamorphopsia*, etc. It may take the form of *contracted fields of vision*, of *color-blindness*, or *anesthesia of the retina*. **A.** **exanopsia**, amblyopia from disuse, or from non-use.

Ambrosia (*am-bro'-she-ah*) [*ἀμβροσία*, the food of the gods]. A genus of composite-flowered herbs. *A. artemisiaefolia*, common hog-weed of North America; stimulant, tonic, antiperiodic, and astringent. *A. trifida* has similar properties. The pollen of these plants is by some regarded as a cause of hay-fever.

Ambulance (*am'-bu-lans*) [*ambulare*, to walk about]. In Europe the term is applied to the surgical staff and arrangements of an army in service. In the United States the word is restricted to a vehicle for the transference of the sick or wounded from one place to another.

Ambulant, **Ambulating**, **Ambulatory** (*am'-bu-lat-o-ry*) [*ambulare*, to walk about]. Relating to walking, or changing location; not confined to bed. **A.** **Blister**, a blister that changes its location. **A.** **Clinic**, a clinic for patients that can walk. **A.** **Erysipelas**, erysipelas that shifts from place to place. **A.** **Tumor**, a pseudo-tumor. **A.** **Typhoid**, walking typhoid; enteric fever in which the patient does not, or will not, take to his bed.

Ambustion (*am-bus'-chun*) [*ambustio*, a burn]. A burn or scald.

Ameba, **Amœba** (*am.-e'-bah*) [*ἀμοιβή*, a change]. A colorless, single-celled, jelly-like, protoplasmic organism found in sea and fresh waters, constantly undergoing changes of form and nourishing itself by englobing surrounding objects. **A.** **coli**, the ameba of dysentery. This is a protoplasmic mass, resembling the water ameba, 20 to 30 μ in diameter, and composed of a nucleus and a highly granular protoplasm containing vacuoles. It is found in large numbers in the stools of certain forms of dysentery, in the intestinal mucous membrane, and at times in the so-called dysenteric abscess of the liver. Whether it is the real cause of the disease is not definitely established.

Amebic (*am.-e'-bik*) [*ἀμοιβή*, a change]. Pertaining to or characterized by amebæ.

A. **Dysentery**, dysentery associated with the presence in the bowel of the amœba coli.

Ameboid (*am.-e'-boid*) [*ἀμοιβή*, a change; *εἶδος*, resemblance]. Resembling an ameba in form or in movement, as the white blood-cells.

Amelia (*ah-me'-le-ah*) [*à* priv.; *μέλος*, a limb]. Absence of the limbs.

Amelioration (*am-él-yo-ra'-shun*) [*ad*, to; *melior*, better]. Improvement.

Ameloblast (*am.-el'-o-blast*) [*Anglo-French*, *amel*, enamel; *βλαστός*, a germ]. An enamel-cell, one of the cylindric cells covering the papilla of the enamel organ of the teeth, and forming a beautifully regular epithelial layer that produces the enamel.

Amelus (*am.-el-us*) [*à* priv.; *μέλος*, a limb]. A monstrosity without limbs.

Amenia (*ah-me'-ne-ah*). See *Amenorrhœa*.

Amenorrhœa (*ah-men-or.-e'-ah*) [*à* priv.; *μήν*, month; *ῥέω*, to flow]. Abnormal absence of menstruation. **A.**, **Primitive**, is a term applied to those cases in which the catamenia have not appeared at the proper time, and **A.**, **Secondary**, to those in which the dis-

charge has been arrested after it has existed during the reproductive period.

Amenorrheal (*ah-men-or-e'-al*) [*à* priv.; *μήν*, month; *ῥέειν*, to flow]. Pertaining to amenorrhea.

Amentia (*ah-men'-she-ah*) [*à* priv.; *mens*, mind]. Defective intellect; idiocy.

Ametria (*ah-met'-re-ah*) [*à* priv.; *μήτρα*, womb]. Absence of the uterus.

Ametrometer (*ah-met-rom'-et-er*) [*à* priv.; *μέτρον*, a measure]. An instrument for measuring ametropia.

Ametropia (*ah-met-ro'-pe-ah*) [*à* priv.; *μέτρον*, a measure; *ὄψις*, sight]. Ametropia exists when an imperfect image is formed upon the retina, due to defective refractive power of the media, or to abnormalities of form of the eye. In *myopia* the antero-posterior diameter is too great, or the power of the refractive media is too great; *hypermetropia* (or *hyperopia*) is the exact reverse; *astigmatism* is due to imperfect curvature of the cornea, or of the retina, or to inequality of refracting power in different parts of the lens; *presbyopia* is due to inelasticity of the lens, producing insufficient accommodation; *aphakia*, or absence of the lens, produces both insufficient refracting power and loss of accommodation.

Ametropic (*ah-met-rof'-ik*) [*à* priv.; *μέτρον*, a measure; *ὄψις*, sight]. Affected with or pertaining to ametropia.

Amid (*am'-id*) [*ammonia*]. A chemic compound produced by the substitution of an acid radicle for one or more of the hydrogen atoms of ammonia. The amids are primary, secondary, or tertiary, according as one, two, or three hydrogen atoms have been so replaced. They are white crystalline solids, often capable of combining with both acids and bases. **A. Bases.** See *Amins*, *Primary*.

Amidin (*am'-id-in*) [*Fr.*, *amidon*, starch]. Starch altered by heat into a horny, transparent mass; soluble starch; the part of starch that is soluble in water.

Amidins (*am'-id-ins*) [*ammonia*]. Mono-acid bases produced from the nitrites by heating with ammonium chlorid. In the free condition they are quite unstable. They contain the group C.NH.NH₂.

Amido-acetic Acid (*am'-id-o-as-e'-tik as'-id*). See *Glycocol* and *Glycin*.

Amido-acid (*am'-id-o-as'-id*) [*ammonia*; *acetum*, vinegar]. An acid containing the amido-group, NH₂.

Amidogen (*am'-id-o-jen*) [*amid*; *γεννᾶν*, to produce]. The hypothetical radicle, NH₂, regarded as an essential part of all amids. See *Amid*.

Amido-succinamic Acid (*am'-id-o-suk-sin-am'-ik as'-id*). Same as *Asparagin*.

Amidulin (*am-id'-u-lin*) [*Fr.*, *amidon*, starch]. Soluble starch; prepared by the action of H₂SO₄ on starch, thus removing the starch-cellulose.

Amimia (*ah-mim'-e-ah*) [*à* priv.; *μίμος*, a mimic]. Loss of the power of imitation or of making gestures.

Amin (*am'-in*) [*ammonia*]. The *Amins* are chemic compounds produced by the substitution of a basic atom or radicle for one or more of the hydrogen atoms of ammonia; or basic derivatives of carbon, containing nitrogen and viewed as ammonia derivatives. In deportment they bear some similarity to ammonia. The lower members are gases with ammoniacal odor and are readily soluble in water; they differ from ammonia in their combustibility. The highest members are liquids soluble in water; only the highest are sparingly soluble. They are called *Monamins*, *Diamins*, *Triamins*, etc., according to the number of amidogen molecules, NH₂, substituted for H. **A.**, **Primary**, an amin in which one hydrogen atom is replaced by a univalent alkyl. **A.**, **Secondary**, an amin in which two hydrogen atoms are replaced by univalent alkyls. **A.**, **Tertiary**, an amin in which three hydrogen atoms are replaced by univalent alkyls.

Aminol (*am'-in-ol*) [*amin*]. A gaseous substance derived from the methylamin of herring-brine mixed with milk of lime. It is disinfectant, and it has been used in the purification of sewage.

Amitosis (*ah-mit-o'-sis*) [*à* priv.; *μίτρος*, a thread]. Cell multiplication by direct division, or simple cleavage.

Amitotic (*ah-mit-o'-ik*) [*à* priv.; *μίτρος*, thread]. Of the nature of, or characterized by, amitosis. **A.** **Cell Division**, direct cell division, as distinguished from karyokinesis.

Ammonemia (*am-o-ne'-me-ah*). The supposed presence of ammonium carbonate in the blood.

Ammonia (*am-o'-ne-ah*) [from the name of Jupiter *Ammon*, from the neighborhood of whose temple in Libya, ammonium chlorid was obtained]. A colorless, pungent gas, NH₃, very soluble in water. The preparations of ammonium are used as antacids and as gastric and cardiac stimulants, in headache, hysteria, etc. It is a stimulant to the heart, and, in its elimination through the lungs, stimulates and liquefies the bronchial secretion. **Ammonia**, **Aqua**, water of ammonia, a solution containing ten per cent. of the gas in water. Dose m̄v-ʒss (0.32-2.0), well diluted. **A.**, **Aqua Fortior**, contains 28 per cent. of the gas in solution. **A.**, **Linimentum**, aqua ammonia 35, cottonseed oil 60, alcohol 5 per cent. **A.**, **Spiritus**,

a ten per cent. solution of aqua ammonia in alcohol. Dose $\text{m}\bar{x}$ - 3j (0.65-4.0), diluted.

A., Spiritus, Aromaticus, aromatic spirit of ammonia, an alcoholic solution of ammonium carbonate, flavored with lemon, lavender, and pimenta. Dose 3 ss - ij (2.0-8.0).

Ammoniac (*am-o'-ne-ak*). See *Ammoniacum*.

Ammoniaccal (*am-o-ni'-ak-al*) [*ammonia*]. Containing or relating to ammonia.

Ammoniacum (*am-o-ni'-ak-um*) [*ammonia*].

Ammoniac. A gum obtained from a Persian plant, *Dorema ammoniacum*. It is a stimulating expectorant and laxative, resembling asafetida, employed in chronic bronchial affections. Dose gr. x -xxx (0.65-2.0). **A. cum Hydrargyro, Emplastrum**, ammoniac 72, mercury 18 per cent., with sulphur, acetic acid, and oil, *q. s.* **A., Emplastrum**, 100 parts of ammoniac, digested with 140 parts of acetic acid, diluted, strained, and evaporated. **A., Emulsum**, a four per cent. emulsion in water. Dose 3 ss - j (16.0-32.0).

Ammoniameter (*am-o-ne-am'-et-er*) [*ammonia*; $\mu\bar{\epsilon}\tau\rho\omega\nu$, a measure]. An instrument for testing the strength of ammonia solutions.

Ammoniated (*am-o-ne-a'-ted*) [*ammonia*]. Combined with ammonia.

Ammonium (*am-o'-ne-um*) [*ammonia*]. A hypothetical univalent alkaline base, having the composition NH_4 . It exists only in combination. **Ammonii acetatis, Liq.**, Spirit of Mindererus, dilute acetic acid neutralized with ammonia. Dose 3 j - 3 j (4.0-32.0).

A. benzoate, $\text{NH}_4\text{C}_6\text{H}_5\text{O}_2$. Dose gr. v-xv (0.32-1.0). **A. bromid**, NH_4Br , used in epilepsy, cough, and rheumatism. Dose gr. x - 5 ss (0.65-2.0). **A. carbonate**, NH_4HCO_3 , $\text{NH}_4\text{NH}_2\text{CO}_3$, a mixture of carbonate and dicarbonate. It is a stimulant expectorant and cardiac stimulant. Dose gr. v-x (0.32-0.65).

A. chlorid, NH_4Cl , sal ammoniac, is used in bronchitis, rheumatism, and liver disease. Dose gr. j-xx (0.065-1.3).

A. chloridi, Trochisci, each lozenge contains gr. ij of the salt. **A. fluorid**, used in enlargement of the spleen. Dose gr. $\frac{1}{4}$ - $\frac{1}{2}$ (0.0027-0.032). **A. glycyrrhizas**, Unof.

An expectorant. **A. iodid**, NH_4I . Dose gr. ij-x (0.13-0.65). **A. nitrate**, NH_4NO_3 , used in preparing nitrous oxide. **A. phosphate**, $(\text{NH}_4)_2\text{HPO}_4$. Dose gr. v-xx (0.32-1.3).

A. picras, $\text{C}_6\text{H}_7(\text{NH}_4)(\text{NO}_2)_3\text{O}$, a salt in yellow needles, of bitter taste; like other picrates it is explosive, and must be handled with care. It is antipyretic and antiperiodic, and tends to correct gastric disturbances. Dose gr. v (0.32) in 24 hours. **A. sulphate**, $(\text{NH}_4)_2\text{SO}_4$, used in the preparation of other ammonium salts. **A. urate**, occurs in alkaline urine and at times in urinary cal-

culi. **A. valerianate**, $\text{NH}_4\text{C}_8\text{H}_9\text{O}_2$, is used as a sedative in hysteria. Dose gr. j-v (0.065-0.32).

Ammotherapy (*am-o-ther'-a-pe*) [$\bar{\alpha}\mu\mu\omega\varsigma$, sand; $\theta\epsilon\rho\alpha\pi\epsilon\iota\tau\epsilon\iota\nu$, to heal]. The use of sand-baths in the treatment of disease.

Amnesia (*am-ne'-se-ah*) [$\bar{\alpha}\mu\eta\sigma\iota\alpha$, forgetfulness]. Loss of memory, especially of the ideas represented by words. **A. Auditory**, word-deafness. **A., Visual**, word-blindness.

Amnesic (*am-ne'-sik*). Relating to amnesia.

A. Aphasia. See *Amnesia*.

Amnio-chorial (*am-ne-o-ko'-re-al*) [$\bar{\alpha}\mu\eta\omega\nu$, the amnion; $\chi\omicron\rho\iota\omega\nu$, a membrane]. Pertaining to both amnion and chorion.

Amnion (*am'-ne-on*) [$\bar{\alpha}\mu\eta\omega\nu$, a young lamb]. The innermost of the fetal membranes; it is continuous with the fetal epidermis at the umbilicus, forming a complete sheath for the umbilical cord and a sac or bag in which the fetus is enclosed. It contains one or two pints of *liquor amnii*. It is a double, non-vascular membrane, the inner layer or *sac* derived from the epiblast, the outer from the mesoblast. The cavity of the inner folds is called the *true amnion*, that of the outer, the *false*. **A., Dropsy of**, excessive secretion of liquor amnii.

Amnionic (*am-ne-on'-ik*) [$\bar{\alpha}\mu\eta\omega\nu$, a young lamb]. Relating to the amnion.

Amniorrhea (*am-ne-o-re'-ah*) [*amnion*, $\rho\eta\iota\alpha$, a flow]. The discharge of the liquor amnii.

Amniotic (*am-ne-ot'-ik*) [$\bar{\alpha}\mu\eta\omega\nu$, a young lamb]. Relating to the amnion. **A. Fluid**. The liquor amnii. See *Amnion*.

Amœba (*am-e'-bah*). See *Ameba*.

Amœboid (*am-e'-boid*). See *Ameboid*.

Amomum (*am-o'-mum*) [$\bar{\alpha}\mu\alpha\mu\omega\nu$, an Eastern spice plant]. A genus of scitamineaceous plants to which the cardamon (*A. cardamomum*) and Grains of Paradise (*A. granum paradisi*) belong.

Amorphia (*ah-mor'-fe-ah*) [$\bar{\alpha}$ priv.; $\mu\omicron\rho\phi\eta$, form]. Shapeless condition.

Amorphism (*ah-mor'-fiz-m*) [$\bar{\alpha}$ priv.; $\mu\omicron\rho\phi\eta$, a form]. The state of being amorphous or without shape; want of crystalline structure.

Amorphous (*ah-mor'-fus*) [$\bar{\alpha}$ priv.; $\mu\omicron\rho\phi\eta$, a form]. Formless, shapeless, not crystalline.

Ampelotherapy (*am-pel-o-ther'-a-pe*) [$\bar{\alpha}\mu\pi\epsilon\lambda\omega\varsigma$, a grape vine; $\theta\epsilon\rho\alpha\pi\epsilon\iota\tau\epsilon\iota\nu$, to heal]. The *Grape-cure*.

Ampereage (*am-pâr'-ahj*) [*Ampère*, a French physicist]. The number of amperes passing in a given circuit.

Ampere (*am-pâr')* [*Ampère*, a French physicist]. A unit of measurement of an electric current.

Ampere-meter (*am-pâr'-mē-ter*) [*Ampère*; $\mu\bar{\epsilon}\tau\rho\omega\nu$, to measure]. An instrument for estimating the current of an electric circuit in amperes.

Amphiarkyochrome (*am-fe-ar'-ke-o-krom*) [*ἀμφί*, both; *ἀρκυς*, net; *χρῶμα*, color]. A term applied by Nissl to a nerve cell the stainable portion of whose cell body is in the form of a pale network, the nodal points of which are joined by an intensely staining network.

Amphiarthrosis (*am-fe-ar-thro'-sis*) [*ἀμφί*, around; *ἄρθρον*, a joint]. A form of mixed articulation in which the surfaces of the bones are connected by broad discs of fibro-cartilage, or else are covered with fibro-cartilage and connected by external ligaments. It is distinguished by limited flexion in every direction, as, e.g., between the vertebrae.

Amphiasier (*am'-fe-as-ter*) [*ἀμφί*, around; *ἀστὴρ*, a star]. The figure formed in indirect cell-division by the achromatin threads and chromatin granules united to form the so-called nuclear spindle, together with the threads of cell-protoplasm radiating from a rounded clear space at each end of the spindle, known as the stars or suns.

Amphibia (*am-fib'-e-ah*) [*ἀμφί*, both; *βίος*, life]. A class of the *Vertebrata*, living both in the water and upon the land, as the frog, newt, etc.

Amphibious (*am-fib'-e-us*) [*ἀμφί*, both; *βίος*, life]. Living both on land and water.

Amphiblastic (*am-fe-blas'-tik*) [*ἀμφί*, on both sides; *βλαστός*, a germ]. Pertaining to that form of complete segmentation that gives rise to an amphiblastula.

Amphiblastula (*am-fe-blas'-tu-lah*) [*ἀμφί*, on both sides; dim. of *βλαστός*, a germ]. The mulberry-mass or morula-stage in the development of a holoblastic egg. It follows the stage known as amphimorula.

Amphibolia (*am-fe-bo'-le-ah*) [*ἀμφιβολία*, uncertainty]. The vacillating period of a fever or disease.

Amphibolic (*am-fe-bol'-ik*) [*ἀμφίβολος*, uncertain]. Uncertain; doubtful. Applied to a period in the febrile process occurring between the fastigium and the defervescence, and marked by exacerbations and remissions.

Amphicrania (*am-fe-kra'-ne-ah*) [*ἀμφί*, both; *κρανίον*, the skull]. Headache affecting both sides of the head.

Amphicreatin (*am-fe-kre'-at-in*) [*ἀμφί*, around; *κρέας*, flesh], $C_9H_{10}N_7O_4$. One of the muscle-leukomains. It crystallizes in brilliant oblique prisms of a yellowish-white color, and is faintly basic.

Amphicreatinin (*am-fe-kre-at'-in-in*) [*ἀμφί*, around; *κρέας*, flesh], $C_9H_{10}N_7O_4$. A member of the creatinin group of leukomains derived from muscle.

Amphidiarthrosis (*am-fe-di-ar-thro'-sis*) [*ἀμφί*, both; *διάρθρωσις*, articulation]. The articulation of the lower jaw, as it partakes of the nature both of ginglymus and arthrodia.

Amphigony (*am-fig'-o-ne*) [*ἀμφί*, on both sides; *γόνος*, offspring]. The sexual process in its broadest sense; gamogenesis.

Amphimixis (*am-fi-miks'-is*) [*ἀμφί*, on both sides; *μῖξις*, mixing]. The mingling of two individuals or their germs; sexual reproduction.

Amphimorula (*am-fe-mor'-u-lah*) [*ἀμφί*, on both sides; *morula*, a mulberry]. The morula, or globular mass of cleavage cells resulting from unequal segmentation, the cells of the hemispheres being unlike in size.

Amphistoma (*am-fis'-to-mah*) [*ἀμφί*, double; *στόμα*, mouth]. A genus of trematode worms, named from the mouth-like apparatus at either end. One species, *A. hominis*, has been found in the large intestine of man.

Amphodiplopia (*am-fo-dip-lo'-pe-ah*) [*ἀμφω*, both; *διπλός*, double; *ὤψ*, eye]. Double vision affecting each of the eyes.

Amphophile, **Amphophilous** (*am'-fo-fil*, *am-fof'-il-us*) [*ἀμφω*, both; *φιλεῖν*, to love]. Readily stainable alike with acid and basic dyes.

Amphoric (*am-for'-ik*) [*amphora*, a vase with two handles]. Resembling the sound produced by blowing across the mouth of a bottle. **A. Breathing**, breath sounds with musical quality heard in diseased conditions of the lung, especially in pulmonary tuberculosis with cavity-formation. **A. Resonance**, in auscultation, a metallic sound like that of blowing into a bottle, caused by the reverberation of sound in a cavity of the lung. **A. Respiration**. See *A. Breathing*.

Amphorophony (*am-for-of'-o-ne*) [*amphora*, a vase with two handles; *φωνή*, a sound]. An amphoric resonance or sound.

Amphoteric, **Amphoteros** (*am-fo-ter'-ik*, *am-fof'-er-us*) [*ἀμφοτερος*, both of two]. Double-sided; having the power of altering the color of both red and blue litmus test paper; a condition sometimes presented by the urine.

Amplification (*am-plif-ik-a'-shun*) [*amplificare*, to enlarge]. Increase of the visual area, in microscopy.

Amplifier (*am'-ple-fi-er*) [*amplificare*, to enlarge]. An apparatus used in microscopy for increasing the magnification. It consists of a diverging lens or combination placed between the objective and the ocular, and gives to the image-forming rays from the objective an increased divergence.

Amplitude (*am'-ple-tūd*) [*amplus*, broad]. The range or extent, as of vibrations and undulations, the pulse wave, etc.

Ampulla (*am-pul'-ah*) [L., a Roman wine jug]. The trumpet-mouthed or dilated extremity of a canal, as of the lachrymal canal, the receptaculum chyli, the Fallopian tubes, mammary ducts, semicircular canals, vas de-

ferens, etc. **A. Chyli**, the receptaculum chyli. **A., Lieberkühn's**, the cecal terminus of one of the lacteals in the villi of the intestines. **A. of Rectum**, the portion above the perineal flexure. **A. of Vater**, the site of entrance of the common bile-duct and pancreatic duct into the duodenum.

Amputation (*am-pu-ta'-shun*) [*amputare*, to cut away]. The removal of a limb or any projecting part of the body. Amputation may be by the knife, ligature, or other means, or it may be the result of pathologic processes, as gangrene, constriction (*e. g.*, of the cord in the fetus). **A., Accidental**, the separation of a limb by some form of accident. **A., Bloodless**, one in which there is but slight loss of blood, on account of the circulation being controlled by mechanical means. **A., Central**, one in which the scar is situated at or near the center of the stump. **A., Circular**, that performed by making a single flap, by circular sweeps of a long knife, through skin and muscles, in a direction vertical to the long axis of the limb. **A., Circular Skin Flap**, a modification of the circular, in which the skin flap is dissected up, and the muscles divided at a higher level. **A., Coat-sleeve**, a modification of the circular, in which the cutaneous flap is made very long, the end being closed by being gathered together by means of a tape. **A., Congenital**, amputation of fetal portions, due to constriction by bands of lymph. **A. in the Contiguity**, amputation at a joint. **A. in the Continuity**, amputation of a limb elsewhere than at a joint. **A., Consecutive**, an amputation during the period of suppuration or later. **A., Cutaneous**, one in which the flaps are composed exclusively of the integuments. **A., Dielastie**, one in which the bone is broken with an osteoclast, and the soft tissues divided by means of an *écraseur*. Its object is to avoid hemorrhage and purulent infection. **A., Double Flap**, one in which two flaps are formed from the soft tissues. **A., Dry**. See *A., Bloodless*. **A., Eccentric**, one in which the scar is situated away from the center of the stump. **A., Elliptical**, one that may be performed by a single sweep, as in the circular method; the wound, however, having an elliptical outline, on account of the oblique direction of the incision. **A. of Expediency**, one performed for cosmetic effect. **A., Flap**, one in which one or more flaps are made from the soft tissues, the division being made obliquely. **A., Flapless**, one in which, on account of destruction of the soft parts, flaps cannot be formed, the wound healing by granulation. **A., Galvano-caustic**, one in which the soft parts are divided with the galvano-cautery followed by division of the bone by

the saw. **A., Immediate**, one done within twelve hours after the injury, during the period of shock. **A., Intermediary**, or **Intermediate**, one performed during the period of reaction, and before suppuration. **A., Intrauterine**. See *A., Congenital*. **A., Major**, amputation of an extremity above the wrist or ankle joint. **A., Mediate**. See *A., Intermediary*. **A., Minor**, amputation of a small part, as a finger. **A., Mixed**, a combination of the circular and flap methods. **A., Multiple**, amputation of two or more members at the same time. **A., Musculo-cutaneous**, one in which the flaps consist of skin and muscle. **A., Musculo-tegumentary**. See *A., Musculo-cutaneous*. **A., Natural**. See *A., Congenital*. **A., Oblique**. See *A., Oval*. **A., Osteoplastic**, one in which there is section and apposition of portions of bone, in addition to the amputation. **A., Oval**, a modification of the elliptical, in which the incision consists of two reversed spirals, instead of the one oblique. **A., Partial**, 1. One in which but a portion of the extremity is removed. 2. An incomplete congenital amputation. **A., Pathologic**, one done for tumor or other diseased condition. **A., Primary**, one done after the period of shock and before the occurrence of inflammation. **A., Racket**, a variety of the oval amputation, in which there is a single longitudinal incision continuous below with a spiral incision on either side of the limb. **A., Secondary**, one performed during the period of suppuration. **A., Spontaneous**. See *A., Congenital*. It also occurs in the disease, *ainhum*. **A., Subperiosteal**, one in the continuity, the cut end of the bone being covered by periosteal flaps. **A., Synchronous**. See *A., Multiple*. **A. by Transfixion**, one done by thrusting a long knife completely through a limb, and cutting the flaps from within out.

Amusia (*ah-mu'-se-ah*) [*à priv.*; *musa*, *μουσα*, muse]. Loss of the ability to produce or comprehend music or musical sounds; an abnormality as regards music analogous to aphasia as regards the faculty of speech. **Motor A.**, the music is understood, but there is loss of the power of singing or of otherwise reproducing music. **Paramusia**, faulty and imperfect reproduction of music. **Sensory A.**, musical deafness, or the loss of the power of comprehension of musical sounds.

Amyelencephalia (*ah-mi-el-en-sef-a'-le-ah*) [*à priv.*; *μυελός*, marrow; *κεφαλή*, the head]. Absence of both brain and spinal cord.

Amyelia (*ah-mi-el'-le-ah*) [*à priv.*; *μυελός*, marrow]. Absence of the spinal cord.

Amyelinic (*ah-mi-el-in'-ik*) [*à priv.*; *μυελός*, marrow]. Without myelin.

Amyelotrophy (*ah-mi-el-ot'-ro-fe*) [*â* priv.; *μυελός*, marrow; *τροφή*, nourishment]. Atrophy of the spinal cord.

Amyelus (*ah-mi'-el-us*) [*â* priv.; *μυελός*, marrow]. A fetal monstrosity with partial or complete absence of the spinal cord.

Amygdala (*am-ig'-dal-ah*) [*ἀμυγδάλη*, almond]. 1. The tonsil. 2. A small lobe on the lower surface of each cerebellar hemisphere, projecting into the fourth ventricle. 3. Almond. The seeds of *A. amara* and *A. dulcis*, containing the principle *Emulsin*. The former contains *Amygdalin*. The expressed oil of the sweet almond is a demulcent, and is useful in skin affections; in doses of $\frac{3}{4}$ – $\frac{1}{2}$ (4.0–8.0) a mild laxative. That of *A. amara* is used in cosmetics. **A. Amaræ, Aqua**, a 1:1000 solution of the oil in water. Dose indefinite. **A. Amaræ, Oleum**, a bitter volatile oil, containing 3–14 per cent. of hydrocyanic acid and having similar uses. Dose $\frac{1}{4}$ – $\frac{1}{2}$ (0.016–0.065). **A., Oleum expressum**, oil of sweet almonds. **A., Emulsum**, oil of sweet almonds 6 per cent., sugar, water, and acacia q. s. Dose $\frac{3}{4}$ – $\frac{3}{4}$ ss. (4.0–16.0). **A., Pulv. Comp.** (B. P.), containing sweet almonds, refined sugar, and gum acacia. It is used to make *A., Emulsum*. **A., Syrupus**, contains oil of bitter almonds 3, sweet almonds 10, syrup 87 per cent. It is used as a vehicle.

Amygdalin (*am-ig'-dal-in*) [*ἀμυγδάλη*, almond], $C_{20}H_{27}NO_{11} + 3H_2O$. A glucosid formed in bitter almonds, in various plants, and in the leaves of the cherry laurel. Under the influence of emulsin, contained in the almond, it splits up into glucose and hydrocyanic acid.

Amygdaline (*am-ig'-dal-in*) [*ἀμυγδάλη*, almond]. 1. Almond-like. 2. Pertaining to the tonsil.

Amygdalitis (*am-ig-dal-i'-tis*) [*ἀμυγδάλη*, almond; *ιτις*, inflammation]. Tonsillitis.

Amygdaloid (*am-ig'-dal-oid*) [*ἀμυγδάλη*, almond; *ειδός*, form]. Resembling an almond. **A. Fossa**, the depression for the lodgment of the tonsil. **A. Tubercle**, a projection of gray matter at the end of the descending cornu of the lateral ventricle of the brain. It is attached to the temporal lobe, and appears to be nearly isolated by white substance.

Amygdalolith (*am-ig-dal'-o-lith*) [*ἀμυγδάλη*, an almond; *λίθος*, a stone]. A concretion or calculus found in the tonsil.

Amygdalopathy (*am-ig-dal-op'-ath-e*) [*ἀμυγδάλη*, an almond; *πάθος*, a disease]. Any disease of the tonsils.

Amygdalotome (*am-ig'-dal-o-tōm*) [*ἀμυγδάλη*, tonsil; *τέμνειν*, to cut]. An instrument used in cutting the tonsils.

Amygdalotomy (*am-ig-dal-ot'-o-mē*) [*ἀμυγ-*

δάλη, a tonsil; *τέμνειν*, to cut]. Tonsillotomy.

Amyl (*am'-il*) [*ἀμυλον*, starch]. The radicle, C_5H_{11} , of amyl alcohol, the fifth member of the series of alcohol radicles, C_nH_{2n+1} . **A. Alcohol**. See *Amylic Alcohol*. **A. Hydrate**. See *Amylic Alcohol*. **A. Nitrite**, nitrite of amyl, $C_5H_{11}NO_2$, a clear, yellowish, volatile liquid, of a penetrating odor. It produces vascular dilatation and stimulates the heart's action, and is useful in angina pectoris, respiratory neuroses, etc. Dose, internally, $\frac{1}{4}$ – $\frac{1}{2}$ (0.016–0.065) dissolved in alcohol; by inhalation, $\frac{1}{2}$ – $\frac{1}{2}$ (0.13–0.32). **A. Valerianate**. See *Apple Oil*.

Amylaceous (*am-il-a'-se-us*) [*ἀμυλον*, starch]. Containing starch; starch-like.

Amylene (*am'-il-en*) [*ἀμυλον*, starch], C_5H_{10} . A liquid hydrocarbon, having anesthetic properties. See *Anesthetic*. **A. Hydrate**, $C_5H_{12}O$, a tertiary alcohol used as an hypnotic. Dose $\frac{1}{4}$ – $\frac{1}{2}$ (2.0–4.0). Unof.

Amylic (*am-il'-ik*) [*ἀμυλον*, starch]. Pertaining to amyl. **A. Alcohol**, Fusel Oil, Potato-starch Alcohol, Amyl Hydrate. An alcohol having the composition $C_5H_{12}O$, produced in the continued distillation of fermented grain. It was formerly used to adulterate whisky. It is employed as a solvent and reagent.

Amylin (*am'-il-in*) [*ἀμυλον*, starch]. The insoluble wall of the starch grain.

Amylogenic (*am-il-o-jen'-ik*) [*ἀμυλον*, starch; *γέννᾶν*, to produce]. Starch-producing.

Amyloid (*am'-il-oid*) [*ἀμυλον*, starch; *ειδός*, form]. Starch-like. **A. Bodies**, bodies resembling starch grains, found in the nervous system, the prostate, etc. They are the result of a localized amyloid degeneration. **A. Degeneration**, waxy, or lardaceous degeneration. A degeneration characterized by the formation of an albuminous substance, resembling starch in its chemic reactions. The process affects primarily the connective tissue of the blood-vessels of various organs, and is connected with or due to chronic suppuration in the body. Amyloid substance gives a brown color with iodine, a red color with gentian-violet, and turns blue on being treated with iodine and sulphuric acid. **A. Kidney**. See *Bright's Disease*.

Amylolysis (*am-il-ol'-is-is*) [*ἀμυλον*, starch; *λύσις*, solution]. The digestion of starch, or its conversion into sugar.

Amyolytic (*am-il-o-lit'-ik*) [*ἀμυλον*, starch; *λύσις*, solution]. Pertaining to or effecting the digestion of starch, as the ferments in the saliva and pancreatic juice, that convert starch into sugar.

Amylopsin (*am-il-op'-sin*) [*ἀμυλον*, starch; *δψις*, appearance]. A ferment found in the

pancreatic juice, which changes starch into sugar.

Amylose (*am'-il-as*) [*ἀμύλου*, starch]. Any one of the group of carbohydrates, comprising starch, glycogen, dextrin, inulin, gum, cellulose, and tunicin.

Amylum (*am'-il-um*) [*L.*], $C_6H_{10}O_5$. Starch. **A.**, **Glyceritum**, contains starch 10, water 10, glycerin 80 per cent.; used for external application. **A.**, **Mucilago** (*B. P.*), used in making enemata. **A. iodatum**, contains starch 95, iodine 5 per cent., triturated with distilled water and dried. Dose \mathfrak{z} j- \mathfrak{z} ss (4.0-16.0).

Amyocardia (*am-i-o-kar'-de-ah*) [*ἀ* priv.; *μῦς*, muscle; *καρδία*, the heart]. Lack of muscular power in the heart's contractions.

Amyostasia (*am-i-os-ta'-ze-ah*) [*ἀ* priv.; *μῦς*, muscle; *στάσις*, standing]. An abnormal trembling of the muscles while in use, often seen in locomotor ataxia.

Amyosthenia (*am-i-os-the'-ne-ah*) [*ἀ* priv.; *μῦς*, muscle; *σθένος*, force]. Deficient muscular power.

Amyotrophia (*am-i-o-tro'-fe-ah*) [*ἀ* priv.; *μῦς*, muscle; *τροφή*, nourishment]. Atrophy of a muscle.

Amyotrophic (*am-i-o-trof'-ik*) [*ἀ* priv.; *μῦς*, muscle; *τροφή*, nourishment]. Characterized by muscular atrophy. **A. Lateral Sclerosis**, lateral sclerosis combined with muscular atrophy. The lesion is in the pyramidal tracts and in the ganglion cells of the anterior gray horns of the spinal cord. The disease has a marked tendency to involve the medulla. **A. Paralysis**, that due to muscular atrophy.

Anabasis (*an-ab'-as-is*) [*ἀναβαίνειν*, to go up]. The increasing stage of acute disease.

Anabatic (*an-ab-at'-ik*) [*ἀναβατικός*, ascending]. Increasing; growing more intense; as the anabatic stage of a fever.

Anabiosis (*an-ab-i'-sis*) [*ἀναβίειν*, to come to life again]. The reappearance of vitality in an apparently lifeless organism.

Anaboly (*an-ab-ol'-er-je*) [*ἀναβάλλειν*, to throw up; *ἔργον*, work]. The force expended or work performed in anabolism, or in anabolic processes.

Anabolic (*an-ab-ol'-ik*) [*ἀναβάλλειν*, to throw up]. Pertaining to or characterized by anabolism.

Anabolin (*an-ab'-o-lin*) [*ἀναβάλλειν*, to throw up]. Any substance formed during the anabolic process.

Anabolism (*an-ab'-o-lizm*) [*ἀναβάλλειν*, to throw up]. Synthetic or constructive metabolism. Activity and repair of function; opposed to katabolism.

Anacamptic (*an-ab-amp'-tik*) [*ἀνακάμπειν*, to bend back]. Reflected, as sound or light; pertaining to or causing a reflection.

Anacardium (*an-ak-ar'-de-um*) [*ανά*, up; *καρδία*, the heart, from its heart-shaped seeds].

1. A genus of tropical trees. *A. occidentale* yields cashew gum and the cashew nut. 2. The oil of the pericarp of the cashew nut, known as cardol, and used as an escharotic. It is said to be of value in leprosy. **A.**, **Tinct.**, 1 to 10 of rectified spirit. Dose, \mathfrak{m} ij-x (0.13-0.65). **A.**, **Unguentum**, one part of the tar to eight of lard or vaselin, used as a blistering ointment.

Anaclasis (*an-ak'-las-is*) [*ἀνάκλασις*, a breaking off, or back]. 1. Reflection or refraction of light or sound. 2. A fracture; forcible flexion of a stiff joint.

Anacrotic (*an-ak-ro'-tik*) [*ἀνά*, upward; *κρότος*, a stroke]. Relating to or characterized by anacrotism.

Anacrotism (*an-ak'-ro-tizm*) [*ἀνά*, upward; *κρότος*, a stroke]. The condition in which there is one or more notches on the ascending limb of the pulse curve.

Anacusia (*an-ak-ot'-se-ah*) [*ἀν* priv.; *ἀκοῦειν*, to hear]. Complete deafness.

Anadenia (*an-ad-e'-ne-ah*) [*ἀν* priv.; *ἀδην*, gland]. Insufficiency of glandular function.

Anadicrotic (*an-ah-di-krot'-ik*) [*ἀνά*, upward; *δῖς*, twice; *κρότος*, a stroke]. Characterized by anadicrotism.

Anadicrotism (*an-ah-di'-krot-izm*) [*ἀνά*, upward; *δῖς*, twice; *κρότος*, a stroke]. Di-crotism of the pulse-wave occurring in the upward stroke.

Anadidymous (*an-ad-id'-im-us*) [*ἀνά*, up; *δίδυμος*, a twin]. Cleft upward into two, while single below—said of certain joined twins.

Anadidymus (*an-ad-id'-im-us*) [*ἀνά*, up; *δίδυμος*, a twin]. An anadidymous monster.

Anadipsia (*an-ah-dip'-se-ah*) [*ἀνά*, intensive; *διψα*, thirst]. Intense thirst.

Anæmia (*an-e'-me-ah*). See *Anemia*.

Anaerobia (*an-a-er-o'-be-ah*) [*ἀν* priv.; *ἀήρ*, air; *βίος*, life]. Microorganisms having the power of living without free oxygen. **A.**, **Facultative**, applied to organisms normally or usually living in the presence of oxygen, but capable of becoming anaerobic.

Anaerobic (*an-a-er-o'-bik*) [*ἀν* priv.; *ἀήρ*, air; *βίος*, life]. Living in the absence of free oxygen or air. See *Aerobic*.

Anaerobiosis (*an-a-er-o-bi-o'-sis*) [*ἀν* priv.; *ἀήρ*, air; *βίος*, life]. Life sustained in the absence of free oxygen; the power of living where there is no free oxygen.

Anaerobiotic, **Anaerobious** (*an-a-er-o-bi-ot'-ik*, *an-a-er-o'-be-us*) [*ἀν* priv.; *ἀήρ*, air; *βίος*, life]. Capable of existing without free oxygen.

Anæsthesia (*an-es-the'-ze-ah*). See *Anesthesia*.

Anæsthetic (*an-es-the'-tik*). See *Anesthetic*.

Anagraph (*an'-a-graf*) [*αναγραφή*, a writing out]. A physician's prescription or recipe.

Anakusis (*an-ak-oo'-sis*). See *Anacusia*.

Anal (*a'-nal*) [*anus*, the fundament]. Pertaining to the anus.

Analepsia, Analepsis (*an-al-ep'-se-ah*, *an-al-ep'-sis*) [*ἀνάληψις*, a taking up]. Recovery of strength after disease.

Analeptic (*an-al-ep'-tik*) [*ἀναληπτικός*, restorative]. 1. Restorative. 2. Any agent restoring health after illness.

Analgen (*an-al'-jen*) [*ἀν* priv.; *ἄλγος*, pain], $C_{10}H_{14}N_2O_4$. A white, tasteless, crystalline powder, almost insoluble in water, soluble with difficulty in cold alcohol, but more readily in hot alcohol, and dilute acids. It melts at 406.4° F. It is employed as an analgesic, antineuralgic, and antipyretic. Dose gr. x-xxx (0.65-2.0).

Analgesia (*an-al'-je'-ze-ah*) [*ἀν* priv.; *ἄλγος*, pain]. Insensibility to or absence of pain.

Analgesic (*an-al'-je'-sik*) [*ἀν* priv.; *ἄλγος*, pain]. 1. Anodyne; relieving pain. 2. Affected with analgesia. 3. A remedy that relieves pain.

Analgesin (*an-al'-je'-sin*). See *Antipyrrin*.

Analgin (*an-al'-jik*) [*ἀν* priv.; *ἄλγος*, pain]. Analgesic.

Analgine. Synonym of *Creolin*.

Analogous (*an-al'-o-gus*) [*ἀνάλογος*, conformable]. Conforming to, proportionate, answering to.

Analogue (*an'-al-og*) [*ἀνάλογος*, conformable]. A part or organ having the same function as another, but with a difference of structure. The correlative term, *homologue*, denotes identity of structure with difference of function. The wing of the butterfly and that of the bird are *analogous*, but the wing of a bird and the arm of a man are *homologous*.

Analysis (*an-al'-is-is*) [*ἀναλίσκειν*, to unloose]. The resolution of a compound body into its constituent parts. **A.**, **Gasometric**, the determination of the constituents of gaseous compounds, especially the determination of the amount of oxygen in samples of atmospheric air. **A.**, **Gravimetric**, the quantitative determination by weight of the elements of a body. **A.**, **Organic**, the determination of the elements of matter formed under the influence of life. The analysis of animal and vegetable tissues. **A.**, **Proximate**, the determination of the simpler compound into which a substance may be resolved. **A.**, **Qualitative**, the determination of the nature of the elements that compose a body. **A.**, **Quantitative**, the determination of the proportionate parts of the various elements of a compound. **A.**, **Spectral**, the determination of the composition of a body by means of the spectroscope. **A.**, **Ultimate**, the resolution of a compound into its ultimate ele-

ments. **A.**, **Volumetric**, the quantitative determination of a constituent by volume.

Analyzer (*an'-al-i-zer*) [*ἀναλίσκειν*, to unloose]. In microscopy, the Nicol prism, which exhibits the properties of light after polarization.

Anamirta (*an-am-er'-tu*). A genus of Menispermaceae. **A.** *paniculata*, or *Menispermum cocculus*, is the source of *Cocculus indicus*.

Anamnesis (*an-am-ne'-sis*) [*ἀνάμνησις*, a recalling to mind]. 1. The faculty of memory; recollection. 2. That which is recollected: information gained regarding the past history of a case from the patient.

Anamnestic (*an-am-ne'-sik*) [*ἀνάμνησις*, a recalling to mind]. 1. Recalling to mind. 2. Restorative of the memory.

Anamniotic (*an-am-ne-of'-ik*) [*ἀν* priv.; *ἄμνιον*, amnion]. Without an amnion.

Anandria (*an-an'-dre-ah*) [*ἀν* priv.; *ἀνήρ*, man]. Lack of virility; impotence.

Anapeiratic (*an-ap-i-rat'-ik*) [*ἀναπειράσθαι*, to do again]. Due to excessive use of certain parts, or of certain muscles, as in writers' and telegraphers' cramp.

Anaphases (*an-af-a'-sez*) [*ἀνά*, up; *φάσις*, a phase]. The phenomena of karyokinesis immediately preceding the formation of the daughter stars, and up to the formation of the resting daughter nuclei.

Anaphia (*an-a'-fe-ah*) [*ἀν* priv.; *ἄφή*, touch]. Defect in the sense of touch.

Anaphoresis (*an-af-or-e'-sis*) [*ἀν* priv.; *φορεῖν*, to carry]. A diminution in the activity of the sweat-glands.

Anaphrodisia (*an-af-ro-dis'-e-ah*) [*ἀν* priv.; *ἄφροδιτη*, Venus]. Absence or impairment of sexual appetite.

Anaphrodisiac (*an-af-ro-dis'-e-ak*) [*ἀν* priv.; *ἄφροδιτη*, Venus]. 1. Relating to, affected by, or causing anaphrodisia. 2. An agent that allays the sexual desire.

Anaplastic (*an-ap-las'-tik*) [*ἀναπλάσσειν*, to build up]. 1. Relating to anaplasty; restoring a lost or defective part. 2. An agent that facilitates repair.

Anaplasty (*an'-ap-las-te*) [*ἀναπλάσσειν*, to build up]. An operation for the restoration of lost parts; plastic surgery.

Anapnograph (*an-ap'-no-graf*) [*ἀναπνοή*, respiration; *γράφειν*, to write]. An apparatus registering the movements of inspiration and expiration, together with the quantity of air inhaled.

Anapnoic (*an-ap-no'-ik*) [*ἀνά*, against; *ἀπνοια*, want of breath]. Relieving dyspnea.

Anapophysis (*an-ap-off'-is-is*) [*ἀνά*, back; *ἀπόφυσις*, an offshoot]. An accessory process of a lumbar or dorsal vertebra, corresponding to the inferior tubercle of the transverse process of a typical dorsal vertebra.

Anarthria (*an-ar'-thre-ah*) [*ἀν* priv.; *ἀρθρον*, articulation]. Defective articulation.

Anasarca (*an-ah-sar'-kah*) [*ἀνά*, through; *σάρξ*, the flesh]. An accumulation of serum in the subcutaneous areolar tissues of the body.

Anasarca (*an-ah-sar'-kus*) [*ἀνά*, through; *σάρξ*, the flesh]. Affected with anasarca.

Anaspadias (*an-as-pa'-de-as*) [*ἀνά*, up; *σπάειν*, to draw]. A urethral opening upon the upper surface of the penis.

Anastaltic (*an-as-tal'-tik*) [*ἀνασταλτικός*, checking, putting back]. 1. Strongly astringent. 2. Centripetal; afferent.

Anastasis (*an-as'-tas-is*) [*ἀνάστασις*, a setting up]. Recovery; convalescence.

Anastatic (*an-as-tat'-ik*) [*ἀνάστασις*, a setting up]. Tending to recovery; restorative.

Anastole (*an-as'-to-le*) [*ἀναστολή*, retracted]. Retraction; shrinking away, as of the lips of a wound.

Anastomosis (*an-as-to-mo'-sis*) [*ἀναστομῶν*, to bring to a mouth]. 1. The intercommunication of blood-vessels. 2. The establishment of a communication between two hollow parts, or between two distinct portions of the same organ. See *A.*, *Intestinal*. **A.**, *Crucial*, an arterial anastomosis in the upper part of the thigh, formed by the anastomotic branch of the sciatic, the first perforating, the internal circumflex, and the transverse branch of the external circumflex arteries. **A.**, *Intestinal*, an operation consisting in establishing a communication between two parts of the intestine.

Anastomotic (*an-as-to-mo'-ik*) [*ἀναστομῶν*, to bring to a mouth]. Pertaining to anastomosis.

Anastomotica magna. See *Arteries*, *Table of*.

Anatherapeusis (*an-ath-er-ap'-u'-ris*) [*ἀνά*, forward; *θεραπεύω*, medical treatment]. Treatment by increasing doses.

Anatomic (*an-at-om'-ik*) [*ἀνατομία*, anatomy]. Pertaining to anatomy. **A.** *Tubercle*. See *Verruca necrogenica*.

Anatomist (*an-at'-om-ist*) [*ἀνατομία*, anatomy]. One who is skilled in anatomy.

Anatomy (*an-at'-o-me*) [*ἀνατομία*]. The science of the structure of organs or of organic bodies. **A.**, *Applied*, anatomy as concerned in the diagnosis and treatment of pathologic conditions. **A.**, *Comparative*, the investigation and comparison of the anatomy of different orders of animals or of plants, one with another. **A.**, *Descriptive*, a study of the separate and individual portions of the body, apart from their relationship to surrounding parts. **A.**, *Gross*, anatomy dealing with the naked-eye appearance of tissues. **A.**, *Homologic*, the study of the correlations of the several parts of the body. **A.**, *Microscopic*, or *Minute*, that studied under the microscope. **A.**,

Morbid, or **Pathologic**, a study of diseased structures. **A.**, **Physiognomic**, the study of expressions depicted upon the exterior of the body, especially upon the face. **A.**, **Regional**, a study of limited parts or regions of the body, the divisions of which are collectively or peculiarly affected by disease, injury, operations, etc. **A.**, **Surgical**, the application of anatomy to surgery. **A.**, **Topographic**, the anatomy of a part in its relation to other parts. **A.**, **Veterinary**, the anatomy of domestic animals.

Anatriptic (*an-at-rip'-tik*) [*ἀνὰτριψς*, a rubbing]. A medicine to be applied by rubbing.

Anazoturia (*an-az-et-u'-re-ah*) [*ἀν* priv.; *azotum*, nitrogen; *οὔρον*, urine]. A condition of deficient excretion of nitrogen in the urine, the urea being chiefly diminished.

Anchusin (*ang'-ku-sin*) [*ἀγχουσα*, alkanet], $C_{25}H_{40}O_8$. The red coloring matter found in alkanet-root. See *Alkanet*.

Anchyloglossia (*ang'-kil-o-glos'-e-ah*) [*ἀγκύλη*, a loop; *δύσσω*, tongue]. Tongue-tie.

Anchylous (*ang'-kil-o'-sis*). See *Ankylosis*.

Anchylostomiasis (*ang'-kil-o-sto-mi'-as-is*) [*anchylostomium*]. The morbid condition depending upon the presence of anchylostom duodenale in the intestines.

Anchylostomum (*ang'-kil-o'-st-o-mum*) [*ἀγκύλος*, crooked; *στόμα*, mouth]. A genus of nematoid worms of which the species *A. duodenale* is sometimes found in the human intestine. It produces a condition analogous to pernicious anemia. See *Parasites*, *Animal*, *Table of*, in *Gould's Illustrated Dictionary*.

Ancipital (*an-sip'-it-al*) [*anceps*, double]. Two-edged.

Anconagra (*ang'-kon-a'-grah*) [*ἀγκών*, the elbow; *ἄγρα*, a seizure]. Arthritic pain at the elbow.

Anconal, **Anconéal** (*ang'-kon-al*, *ang'-ko-ne-al*) [*ἀγκών*, the elbow]. Pertaining to the elbow.

Anconeus (*ang'-ko-ne'-us*) [*ἀγκών*, the elbow]. See *Muscles*, *Table of*.

Ancyroid (*an'-sir'-oid*) [*ἄγκυρα*, anchor; *εἶδος*, form]. Shaped like an anchor.

Andersch's Ganglion. The inferior ganglion of the glosso-pharyngeal nerve.

Anderson's Pill. The compound gamboge pill.

Androgalactozemia (*an-dro-gal-ak-to-ze-me-ah*) [*ἄρρ*, man; *γάλα*, milk; *ζημία*, loss]. The presence of milk in the male mamma.

Androgyna (*an-droj'-in-ah*) [*ἄρρ*, man; *γυνή*, woman]. An hermaphrodite; a female in whom the genital organs are similar to those of the male.

Androgyneity (*an-droj'-in-e'-it-e*) [*ἄρρ*, man; *γυνή*, woman]. Hermaphroditism.

Androgynus (*an-droj'-in-us*) [*ἀνήρ*, man; *γυνή*, woman]. An hermaphrodite. A male with genital organs similar to those of the female.

Andrology (*an-drol'-o-je*) [*ἀνήρ*, man; *λόγος*, science]. 1. The science of man, especially of the male sex. 2. The science of the diseases of the male genito-urinary organs.

Andromania (*an-dro-ma'-ne-ah*) [*ἀνήρ*, a man; *μανία*, madness]. Nymphomania.

Androphobia (*an-dro-fo'-be-ah*) [*ἀνήρ*, man; *φόβος*, dread]. Fear or dislike of the male sex.

Anelectrode (*an-el-ek'-trōd*) [*ἀνά*, upward; *electrode*]. The positive pole of a galvanic battery.

Anelectrotonic (*an-el-ek-tro-ton'-ik*) [*ἀν* priv.; *ἤλεκτρον*, amber; *τόνος*, tension]. Relating to anelectrotonus.

Anelectrotonus (*an-el-ek-tro'-o-nus*) [*ἀν* priv.; *ἤλεκτρον*, electricity; *τόνος*, tension]. The decreased irritability that is present in a nerve in the neighborhood of the anode.

Anel's Probe, or Sound, a fine probe used in operations upon the lacrymal passages.

A. Syringe, a syringe used in injecting fluids into the lacrymal passages.

Anemia (*an-e'-me-ah*) [*ἀν* priv.; *αἷμα*, blood]. Deficiency of blood as a whole, or deficiency of the number of the red corpuscles, or of the hemoglobin. It may be *general* or *local*. Local anemia, or *ischemia*, is the result of mechanical interference with the circulation of the affected part. General anemia is either idiopathic or symptomatic. **A.**, **Cytogenic**. Synonym of **A.**, **Idiopathic**. **A.**, **Essential**. Synonym of **A.**, **Idiopathic**. **A.**, **Idiopathic**, one in which the lesion is in the blood or the blood-making organs. **A. lymphatica**. Synonym of *Hodgkin's Disease*. See *Lymphadenoma*. **A.**, **Primary**. See **A.**, **Idiopathic**. **A.**, **Secondary**, or **Symptomatic**, that due to a distinct cause, as hemorrhage, cancer, wasting discharges, poisons, etc.

Anemic (*an-em'-ik*) [*ἀν* priv.; *αἷμα*, blood]. Pertaining to anemia. **A. Infarct**, a wedge-shaped area of coagulation-necrosis occurring in organs possessing terminal arteries. It is the result of the sudden stopping of such an artery by a thrombus or an embolus. **A. Murmur**, a murmur heard in anemic conditions, soft and blowing in character, and disappearing with the anemia. It is generally heard over the base of the heart. **A. Necrosis**, the coagulation-necrosis of tissues resulting from the sudden stoppage of the supplying artery.

Anemometer (*an-em'-om'-et-er*) [*ἀνεμος*, wind; *μέτρον*, a measure]. An instrument for measuring the velocity of the wind.

Anemone (*an-em'-o-ne*) [*ἀνεμώνη*, wind

flower]. A genus of ranunculaceous herbs, most of which have active medicinal and poisonous qualities. See *Pulsatilla*.

Anemonin (*an-em'-o-nin*) [*ἀνεμώνη*, wind-flower], $C_{15}H_{12}O_8$. The active principle of the *Anemone*. It is given in bronchitis, asthma, and spasmodic cough. Dose gr. $\frac{1}{4}$ – $\frac{1}{2}$ (0.016–0.048) twice daily.

Anencephalia (*an-en-sef'-a'-le-ah*) [*ἀν* priv.; *ἐγκέφαλος*, brain]. Absence of the brain.

Anencephalic (*an-en-sef'-al'-ik*) [*ἀν* priv.; *ἐγκέφαλος*, brain]. Pertaining to or characterized by anencephalia.

Anencephalus (*an-en-sef'-al-us*) [*ἀν* priv.; *ἐγκέφαλος*, brain]. A species of single autostic monsters in which there is no trace of the brain.

Anergia (*an-er'-je-ah*) [*ἀν* priv.; *ἔργον*, work]. Sluggishness; inactivity.

Anergic (*an-er'-jik*) [*ἀν* priv.; *ἔργον*, work]. Characterized by sluggishness; as *anergic* dementia.

Aneroid (*an'-er-oid*) [*ἀν* priv.; *νηρός*, wet; *εἶδος*, form]. Working without a fluid. **A. Barometer**. See *Barometer*.

Anerythrospia (*an-er-ith-rof'-se-ah*) [*ἀν* priv.; *ἐρυθρός*, red; *ὄψις*, sight]. Impaired color perception of red.

Anesthesia, or Anæsthesia (*an-es-the'-ze-ah*) [*ἀναίσθησία*, want of feeling]. A condition of total or partial insensibility, particularly to touch. **A.**, **Central**, due to disease in the nerve centers. **A.**, **Crossed**, anesthesia on one side of the body, due to a central lesion of the other side. **A. dolorosa**, severe pain experienced after the occurrence of complete motor and sensory paralysis, a symptom observed in certain diseases of the spinal cord. **A.**, **Local**, that limited to a part of the body. **A.**, **Muscular**, loss of the muscular sense. **A.**, **Peripheral**, that depending upon changes in the peripheral nerves. **A.**, **Primary**, a temporary insensibility to slight pain occurring in the beginning of anesthesia and during which minor operations can be performed. **A.**, **Surgical**, that induced by the surgeon by means of anesthetics for the purpose of preventing pain, producing relaxation of muscles, or for diagnostic purposes.

Anesthetic, or Anæsthetic (*an-es-the'-ik*) [*ἀν* priv.; *αἰσθησία*, feeling]. 1. Without feeling; insensible to touch or pain. 2. A substance that produces insensibility to touch or to pain, diminished muscular action, and other phenomena. Anesthetics may be general, local, partial, and complete. **A.**, **General**; the following are the substances that have been used for general anesthesia: AMYLENE, C_5H_{10} (*Pental*), a thin, colorless, translucent liquid; action rapid, producing partial anesthesia. It should not be brought near a flame. CARBON TETRACHLORID,

not so irritating to the organs of respiration, but far more dangerous than chloroform. CHLORAL HYDRATE, action indirect and incomplete, and rarely, if ever, now used. CHLOROFORM, by inhalation. Largely employed in general surgery. It seems to have a selective action upon the nervous system, and also exercises a direct influence upon the muscular tissues of the heart. It paralyzes the vasomotor system, and death results from cardiac paralysis. Chloroform should always be administered freely mixed with air. The *Lister Method*, also known as the *Scotch or Open Method* of administering chloroform, consists in pouring a small amount of the anesthetic upon a common towel arranged in a square of six folds, and holding this as near to the face as can be borne without inconvenience. ETHYLENE CHLORID, formerly called *ethylene chlorid*, *Dutch liquid*, *chloric ether*, closely resembles chloroform, but is less depressant to the heart, and is considered safer than chloroform. Death results from paralysis of the respiratory centers. ETHER, by inhalation, is probably the safest known agent for the production of prolonged narcosis. Its action is directed largely to the nervous system, which becomes profoundly affected. It frequently causes spasmodic action and suspension of respiration. Death results from paralysis of respiration. It may be administered alone or in combination with nitrous oxid gas, in which case the patient is more rapidly narcotized. ETHIDENE CHLORID is similar in its action to chloroform. Patients take a longer time to recover consciousness than when chloroform is used, but they experience fewer after-effects. Anesthetization occurs in from 3 to 5 minutes. HYDROBROMIC ETHER (*bromid of ethyl*) produces unconsciousness and anesthesia in one minute, and complete muscular relaxation in two or three minutes. The heart's action is somewhat weakened. Return to consciousness after withdrawal of the ether is very prompt. Its use may be followed by vomiting. According to some observers, death results from cardiac failure, while others believe it kills by direct action upon the respiratory center. It is best administered by an Allis inhaler, and is of most service in minor surgery for short operations. NITROUS OXID, by inhalation, is much used by dentists as an anesthetic in the extraction of teeth. The symptoms resemble those of asphyxia, hence it is more important to watch the respiration than the pulse. It may be employed in minor surgical operations. There are but few after-effects, those most often observed being headache and malaise. **A., Local**, an anesthetic that, locally ap-

plied, produces absence of sensation in the organ or tissue so treated. ALCOHOL, locally, removes sensation to pain, while tactile sense persists. Cool the alcohol to about 10° below the freezing point, by placing it in ice and salt, and place the part to be numbed in it. CARBOLIC ACID painted over the skin—its action, however, is caustic. CHLORETHYL, in vapor form, is useful in minor and dental surgery. CHLORID OF METHYL, CH_3Cl , allowing the liquid chlorid to drop on the skin or mucous membrane. Unless kept carefully under control the vitality of the tissues may be affected by the substance. COCAIN is used in subcutaneous injections, by painting over mucous or cutaneous surfaces, or, in the case of the eye, by instillation. As a paint, a 20 per cent. solution is used, weaker preparations being of little value over cutaneous surfaces; several coats are necessary. A 10 per cent. solution should be used on mucous surfaces; for the eye a 2 to 4 per cent. solution will answer, and this solution is strong enough when cocain is employed as a spray. Hypodermically. Mij-v of a 10 to 20 per cent. solution are usually injected, and this may be repeated two or three times during the operation. DISUTPHID OF CARBON, by spray or irrigation, is a local anesthetic, but has a disgusting odor and is a powerful poison. ETHER, in spray, is also a local anesthetic. The anesthesia thus produced is confined to the skin, and is very transient. It may produce a slough from excessive freezing. RHIGOLENE, in spray, its use being similar to that of ether. **A. Mixtures** contain combinations of substances for producing anesthesia. The following are the most important: **A. C. E. MIXTURE**:—alcohol, sp. gr. .838, 1 part; chloroform, sp. gr. 1.497, 2 parts; ether, sp. gr. .735, 3 parts. **BILLROTH'S**:—chloroform 3 parts, alcohol and ether each 1 part. **LINHART'S**:—alcohol 1, chloroform 4. It is administered similarly to chloroform. **MARTINDALE'S**:—a volumetric mixture, the ingredients of which evaporate almost uniformly; it consists of absolute alcohol, sp. gr. .795, one volume; chloroform, sp. gr. 1.498, two volumes; pure ether, sp. gr. .720, three volumes. **MEDICO-CHIRURGICAL SOCIETY OF LONDON**:—ether 3, chloroform 2, alcohol 1. **METHYLENE**, or **METHYLENE BICHLORID**:—a mixture of methylic alcohol 30 per cent., and chloroform 70 per cent. The so-called "*Liquid of Rognaud*" consisted of 80 per cent. chloroform, 20 per cent. methylic alcohol. Methylene is not much used, several deaths from cardiac paralysis having occurred from its employment. **NUSSBAUM'S**:—ether 3, chloroform 1, alcohol 1. **RICHARDSON'S**:—alcohol 2, chloroform 2, ether 3. **SANFORD'S**:—the

so-called "CHLORAMYL," is a mixture of chloroform and amyl nitrate in the proportion of two drams of the nitrate to the pound of chloroform. It is a dangerous mixture, both drugs depressing the heart. VIENNA GENERAL HOSPITAL:—ether 9, chloroform 30, alcohol 9. VIENNA MIXTURE:—ether 3, chloroform 1. VON MERING'S:—chloroform one volume, dimethylacetal two volumes. It is said not to produce failure of respiration or heart, nor lowering of the blood pressure. WACHMUTH'S:—one-fifth part of oil of turpentine is added to the chloroform. It is said to prevent any danger of heart-failure.

Anesthetization (*an-es-thet-iz-a'-shun*) [*ἀναίσθησις*, insensible]. The act of placing under the influence of an anesthetic.

Anesthetize (*an-es'-thet-iz*) [*ἀναίσθητος*, insensible]. To put under the influence of an anesthetic.

Anesthetizer (*an-es'-thet-i-ser*) [*ἀναίσθητος*, insensible]. One who administers an anesthetic.

Anethol (*an'-eth-ol*) [*ἀνά*, up; *αἰθεῖν*, to burn; *oleum*, oil], $C_{10}H_{12}O$. The chief constituent of the essential oils of anise and fennel. It is employed in preparing the *Elixir anethi*, N. F., being more fragrant and agreeable than the anise oil.

Anethum (*an'-el'-thum*) [*ἀνῆθον*, anise]. Dill; the dried fruit of *Peucedanum graveolens*, indigenous to Southern Europe. It is aromatic, carminative, and stimulant. **A.**, Aqua (B. P.). Dose ʒj-ij (32.0-64.0). **A.**, Oleum (B. P.). Dose ʒj-iv (0.065-0.26).

Aneuria (*ah-nu'-re-ah*) [*ἀ* priv.; *νεῦρον*, a nerve]. Lack of nervous power.

Aneuric (*ah-nu'-rik*) [*ἀ* priv.; *νεῦρον*, a nerve]. Characterized by aneuria.

Aneurysm (*an'-u-rizm*) [*ἀνεύρυσμα*, a widening]. A circumscribed dilatation of the walls of an artery. The symptoms of aneurysm depend upon the location of the aneurysmal tumor. Expansile pulsation and a bruit are important; very significant are the so-called pressure-symptoms, which vary with the organ or part pressed upon. **A.**, Abdominal, one of the abdominal aorta. **A.**, Ampullary, a small saccular aneurysm; it is most common in the arteries of the brain. **A.** by Anastomosis, a dilatation of a large number of vessels—small arteries, veins, and capillaries—the whole forming a pulsating tumor under the skin. This form of aneurysm is especially seen upon the scalp. **A.**, Arterio-venous, the simultaneous rupture of an artery and a vein, the blood from both being poured out into the cellular tissue and forming a false aneurysm. A *Varicose Aneurysm* is produced by the rupture of an aneu-

rysm into a vein. An *Aneurysmal Varix* results from the establishment of a communication between an artery and a vein, the latter becoming dilated and pulsating. **A.**, Bérard's, a varicose aneurysm with the sac in the tissues immediately around the vein. **A.**, Cardiac, an aneurysm of the heart. **A.**, Cirroid, a tortuous lengthening and dilatation of a part of an artery. **A.**, Compound, one in which one or several of the coats of the artery are ruptured and the others merely dilated. **A.**, Consecutive, or Diffused, follows rupture of all the arterial coats with infiltration of surrounding tissues with blood. **A.**, Dissecting, one in which the blood forces its way between the coats of an artery. **A.**, Ectatic, an expansion of a portion of an artery due to yielding of all the coats. **A.**, Endogenous, one formed by disease of the vessel walls. **A.**, Exogenous, one due to traumatism. **A.**, False, or Spurious, one due to a rupture of all the coats of an artery, the effused blood being retained by the surrounding tissues. **A.**, Fusiform, a spindle-shaped dilatation of the artery. **A.**, Sacculated, a sac-like dilatation of an artery communicating with the main arterial trunk by an opening that is relatively small. **A.**, Park's, a variety of arterio-venous aneurysm in which the arterial dilatation communicates with two contiguous veins. **A.**, Pott's. Same as *Aneurysmal Varix*. **A.**, Rodrigues', a varicose aneurysm in which the sac is immediately contiguous to the artery. **A.**, Spurious. See *A.*, False. **A.**, Varicose. See *A.*, Arterio venous.

Aneurysmal (*an-u-riz'-mal*) [*ἀνεύρυσμα*, a widening]. Of the nature of or pertaining to an aneurysm. **A.**, Varix. See *Aneurysm*.

Angeo- (*an'-je-o*). See *Angio-*.

Angelica (*an-jel'-ik-ah*) [L.]. The seeds and root of *Angelica archangelica*. It is an aromatic stimulant and emmenagogue. Dose of the seeds or roots, gr. xxx-ʒj (2.0-4.0).

Angel's Wing (*an'-jels wing*). A deformity of the scapula in which it turns forward and then backward, giving the shoulder a peculiar dorsal bulge.

Angiectasis (*an-je-ek'-tas-is*) [*ἀγγειον*, a blood-vessel; *ἐκτασις*, dilatation]. Abnormal dilatation of a vessel.

Angiitis, or **Angieitis** (*an-je-i'-tis*) [*ἀγγειον*, a vessel; *ιτις*, inflammation]. Inflammation of a lymph-vessel or a blood-vessel.

Angina (*an'-jin-ah* or *an-jit'-nah*) [*angere*, to strangle]. Any disease attended by a sense of choking or suffocation, particularly an affection of the fauces or pharynx presenting such symptoms. **A. acuta** or **simplex**, simple sore-throat. **A. externa**. Synonym of *Mumps*. **A. laryngea**. Synonym of

Laryngitis. **A. lingualis**. Same as *Glossitis*. **A. Ludovici**, or **A.**, Ludwig's, acute suppurative inflammation of the connective tissue surrounding the submaxillary glands. **A. membranacea**. Synonym of *Diphtheria*. **A. parotidea**, the mumps, or Parotitis. **A. pectoris**, a paroxysmal neurosis with intense pain and oppression about the heart. It usually occurs in the male after forty years of age, and is generally associated with diseased conditions of the heart and aorta. There is a sense of impending death, and frequently there is a fatal termination. **A. pectoris vasomotoria**, a term given by Nothnagel and Landois to an angina associated with vasomotor disturbances, coldness of the surface, etc. **A.**, **Pseudo-**, a neurosis occurring in anemic females, characterized by a less grave set of symptoms and never resulting fatally. **A. simplex**. See *A. acuta*. **A. tonsillans**, Quinsy. **A. trachealis**, Croup. **A. ulceromembranus**. See *Tonsillitis*, *Herpetic*. **Anginoid** (*an'-jin-oid*) [*angere*, to strangle]. Resembling angina.

Anginose (*an'-jin'-ose*) [*angere*, to strangle]. Pertaining to angina; characterized by symptoms of suffocation.

Angiocholitis (*an-je-o-kol-i'-tis*) [*ἀγγειον*, a vessel; *κολή*, bile; *τις*, inflammation]. Inflammation of the biliary ducts.

Angioderma pigmentosum (*an-je-o-der'-mah pig-men-to'-sum*). See *Atrophoderma*.

Angiogenesis, **Angiogeny** (*an-je-o-jeu'-es-is*, *an-je-og'-en-e*) [*ἀγγειον*, a vessel; *γεννῶν*, to produce]. The development of the vessels.

Angioglioma (*an-je-o-gli-o'-mah*) [*ἀγγειον*, a vessel; *glioma*]. A glioma rich in blood-vessels.

Angiograph (*an'-je-o-graf*) [*ἀγγειον*, a vessel; *γράφειν*, to write]. A variety of sphygmograph.

Angiography (*an-je-og'-ra-fe*) [*ἀγγειον*, a vessel; *γράφω*, a writing]. A description of the vessels; angiology.

Angiokeratoma (*an-je-o-ker-at-o'-mah*) [*ἀγγειον*, a vessel; *κέρως*, horn; *δμα*, tumor]. Lymphangiectasis; Telangiectatic warts; a very rare disease of the extremities characterized by warty-looking growths that develop on dilated vessels in persons with chilblains, etc. Dark vascular spots the size of pins' points or pins' heads, develop as an attack of chilblains is subsiding. The disease is peculiar to childhood.

Angioleucitis (*an-je-o-lu-si'-tis*) [*ἀγγειον*, a vessel; *λευκός*, white; *τις*, inflammation]. Inflammation of the lymphatic vessels.

Angiolithic (*an-je-o-lith'-ik*) [*ἀγγειον*, a vessel; *λίθος*, a stone]. A term applied to neoplasms in which crystalline or mineral deposits take place, with hyaline degeneration of the coats of the vessels.

Angiology (*an-je-ol'-o-je*) [*ἀγγειον*, a vessel; *λόγος*, science]. The science of the blood-vessels and lymphatics.

Angiolymphitis (*an-je-o-limf-i'-tis*). Same as *Angioleucitis*.

Angiolymphoma (*an-je-o-limf-o'-mah*) [*ἀγγειον*, a vessel; *lymphā*, lymph; *δμα*, tumor]. A tumor formed of lymphatic vessels.

Angioma (*an-je-o'-mah*) [*ἀγγειον*, a vessel; *δμα*, a tumor]. A tumor formed of blood-vessels. **A.**, **Cavernous**, an angioma with communicating blood spaces, like the cavernous tissue of the penis. **A.**, **Telangiectatic**, an angioma composed of dilated blood-vessels.

Angiomalacia (*an-je-o-mal-a'-se-ah*) [*ἀγγειον*, a vessel; *μαλακία*, a softening]. Softening of the blood-vessels.

Angiometer (*an-je-om'-et-er*). See *Sphygmograph*.

Angioneurosis (*an-je-o-nu-ro'-sis*) [*ἀγγειον*, a vessel; *νεῖρον*, a nerve]. A neurosis of the blood-vessels; a disturbance of the vasomotor system, either of the nature of a spasm of the blood-vessels (*Angiospasm*) or of paralysis (*Angioparalysis*).

Angioneurotic (*an-je-o-nu-ro'-ik*) [*ἀγγειον*, a vessel; *νεῖρον*, a nerve]. Pertaining to angioneurosis. **A. Edema**, an acute circumscribed swelling of the subcutaneous or submucous tissues, probably due to vasomotor lesion. The disease often runs in families. It is at times periodic and is associated with colic and gastric disturbances.

Angioparalysis (*an-je-o-par-at'-is-iy*) [*ἀγγειον*, a vessel; *παράλυσις*, paralysis]. Vasomotor paralysis.

Angioparalytic (*an-je-o-par-at'-it'-ik*) [*ἀγγειον*, a vessel; *παράλυσις*, paralysis]. Relating to or characterized by angioparalysis.

Angioparesis (*an-je-o-par'-es-is*) [*ἀγγειον*, a vessel; *πάρεσις*, paresis]. Partial paralysis of the vasomotor apparatus.

Angiopathy (*an-je-op'-a-the*) [*ἀγγειον*, a vessel; *πάθος*, disease]. Any disease of the vascular system.

Angiorrhexis (*an-je-o-reks'-is*) [*ἀγγειον*, a vessel; *ρήξις*, a bursting]. Rupture of a blood-vessel.

Angiosarcoma (*an-je-o-sar-ko'-mah*) [*ἀγγειον*, a vessel; *σάρξ*, flesh; *δμα*, a tumor]. A vascular sarcoma.

Angiosialitis (*an-je-o-si-al-i'-tis*) [*ἀγγειον*, a vessel; *σάλων*, saliva; *τις*, inflammation]. Inflammation of the duct of a salivary gland.

Angiosis (*an-je-o'-tis*) [*ἀγγειον*, a vessel]. Any disease of blood-vessels or lymphatics.

Angiospasm (*an-je-o-spazm*) [*ἀγγειον*, a vessel; *σπασμός*, a spasm]. A vasomotor spasm.

Angiospastic (*an-je-o-spas'-tik*) [*ἀγγειον*, a vessel; *σπασμός*, spasm]. Characterized by or of the nature of angiospasm.

Angiostenosis (*an-je-o-sten-o'-sis*) [*ἀγγειον*, a vessel; *στένωσις*, a narrowing]. Narrowing of a vessel.

Angiectasia (*an-je-o-tel-ek-ta'-se-ah*) [*ἀγγειον*, a vessel; *τέλος*, end; *εκτασις*, dilatation]. Dilatation of the blood-vessels.

Angiotomy (*an-je-ot'-o-me*) [*ἀγγειον*, a vessel; *τομή*, a cutting]. Incision into a vessel.

Angle, Angulus (*ang'-gl* or *ang'-gu-lus*) [*angulus*, an angle]. 1. A corner. 2. The degree of divergence of two lines or planes that meet each other; the space between two such lines. **A. of Aberration.** See *A. of Deviation*.

A., Acromial, that formed between the head of the humerus and the clavicle. **A., Alpha.** In optics, that formed by the intersection of the visual line and optic axis. **A., Alveolar**, that formed between a line passing through a spot beneath the nasal spine and the most prominent point of the lower edge of the alveolar process of the superior maxilla and the cephalic horizontal line. **A. of Aperture**, in optics, that included between two lines joining the opposite points of the periphery of a lens and the focus. **A., Biorbital**, in optics, that formed by the intersection of the axes of the orbits. **A., Costal.** The angle formed by the meeting of ribs at the ensiform cartilage.

A., Critical, that made by a beam of light passing from a rarer to a denser medium, with the perpendicular, without being entirely reflected. **A. of Deviation.** 1. In magnetism, the angle traversed by the needle when disturbed by some magnetic force. 2. In optics, that formed by a refracted ray and the prolongation of the incident ray. **A. of Elevation**, in optics, that made by the visual plane with its primary position when moved upward or downward.

A. of Incidence, in optics, the angle at which a ray of light strikes a denser medium and undergoes reflexion or refraction. **A. of Inclination (of Pelvic Canal)**, in obstetrics, that formed by the anterior wall of the pelvis with the conjugate diameter. **A. of Inclination (of Pelvis)**, in obstetrics, that formed by the pelvis with the general line of the trunk, or that formed by the plane of the inferior strait with the horizon. **A. of Jaw**, the junction of the lower border of the ramus of the mandible with its posterior border. **A., Limiting.** See *A., Critical*.

A. of the Lips, those formed by the union of the lips at each extremity of the mouth. **A., Louis's**, that between the manubrium and gladiolus of the sternum. **A., Ludwig's.** See *A., Louis's*. **A., Meter-**, in optics, the degree of convergence of the eyes when centered on an object one meter distant from each. **A., Optic**, that included between lines joining the extremities

of an object and the nodal point. The smallest is about 30 seconds. **A. of Polarization**, in optics, the angle of reflection at which light is most completely polarized. **A. of the Pubes**, that formed by the junction of the pubic bones at the symphysis. **A. of Reflection**, in optics, that which a reflected ray of light makes with a line drawn perpendicular to the point of incidence. **A. of Refraction**, in optics, that which exists between a refracted ray of light and a line drawn perpendicular to the point of incidence. **A., Sacro-vertebral**, that which the sacrum forms with the last lumbar vertebra. **A., Sterno-clavicular**, that existing between the clavicle and the sternum. **A., Subcostal.** See *A., Costal*. **A., Subpubic**, that formed at the pubic arch. **A. Visual.** See *A. Optic*. **A., Xiphoid**, that formed by the sides of the xiphoid notch.

Anglesey Leg (*ang'-gl-se leg*) [so called after the Marquis of Anglesey]. An artificial limb formed from a solid piece of wood hollowed out to receive the stump and provided with a steel joint at the knee. The ankle-joint was made of wood, to which motion was communicated by strong cat-gut strings posteriorly and a spiral spring anteriorly.

Angophrasia (*ang-go-fra'-se-ah*) [*ἀγγειν*, to choke; *φράσις*, utterance]. A speech-defect consisting of a choking, drawing utterance, occurring in paralytic dementia.

Anguillula (*ang-gwil'-u-lah*) [dim. of *anguilla*, an eel]. A genus of parasitic round worms. **A., Stercoralis.** See *Thread-worms*.

Angular (*ang'-gu-lar*) [*angulus*, an angle]. Pertaining to an angle. **A. Artery**, the terminal branch of the facial artery. **A. Gyrus**, or **Convolution**, a convolution of the brain. See *Convolution*. **A. Movement**, the movement between two bones that may take place forward and backward, or inward and outward. **A. Processes**, the external and internal extremities of the orbital arch of the frontal bone.

Angulus (*ang'-gu-lus*). [L.]. See *Angle*.

Angustura (*ang-gus-tu'-rah*) [Sp., *Angostura*, a S. A. town]. Cusparia Bark. The bark of *Galipea cusparia*. It is a stimulant tonic and febrifuge used in malignant bilious fever, intermittent fever, and dysentery. In large doses it is emetic. Dose of fld. ext. ℥x-xxx (0.65-2.0); of the bark gr. x-xl (0.65-2.6). Unof. **Infus. Cuspariæ** (B. P.). Dose ʒj-ij (32.0-64.0).

Angusturin (*ang-gus-tu'-rin*) [Sp., *Angostura*, a S. A. town]. Synonym of *Brucin*.

Anhelation (*an-hel'-a'-shun*) [*anhelare*, to pant]. Shortness of breath; dyspnea.

Anhematosis (*an-hem-at-o'-sis*) [*ἀν* priv; *ἡματίζειν*, to make bloody]. Defective formation of the blood.

Anhidrosis (*an-hid-ro' sis*) [*ân* priv.; *îdros*, sweat]. Partial or complete absence of sweat secretion.

Anhidrotic (*an-hid-ro't-ik*) [*ân* priv.; *îdros*, sweat]. 1. Tending to check sweating. 2. An agent that checks sweating.

Anhydremia (*an-hi-dre'-me-ah*) [*ân* priv.; *îdros*, water; *aima*, blood]. The opposite of hydremia. A diminution of the watery constituents of the blood.

Anhydrid (*an-hi'-drîd*) [*ân* priv.; *îdros*, water]. A chemic compound, particularly an acid, formed by the withdrawal of a molecule of water. Carbon dioxide and sulphur dioxide are examples.

Anhydrous (*an-hi'-drous*) [*ân* priv.; *îdros*, water]. In chemistry, a term used to denote the absence of water.

Anideus (*an-id'-e-us*) [*ân* priv.; *eidros*, form]. The lowest form of omphalosite, in which the parasitic fetus is reduced to a shapeless mass of flesh covered with skin.

Anidrosis (*an-id-ro'sis*). See *Anhidrosis*.

Anidrotic (*an-id-ro't-ik*). See *Anhidrotic*.

Anilid (*an'-il-id*) [Arab., *al*, the; *nil*, dark blue]. Compounds formed by the action of acid chlorids or acid anhydrids upon the anilins. They are very stable derivatives.

Anilin (*an'-il-in*) [Arab., *al*, the; *nil*, dark blue], C_6H_5N . Amidobenzene; formed in the dry distillation of bituminous coal, bones, indigo, isatin, and other nitrogenous substances. It is made by reducing nitrobenzene. It is a colorless liquid with a faint, peculiar odor, boiling at 183° ; its sp. gr. at 0° is 1.036. When perfectly pure it solidifies on cooling, and melts at -8° . It is slightly soluble in water, but dissolves readily in alcohol and ether. Combined with chlorin, the chlorates and hypochlorites, it yields the various anilin dyes. It is used in chorea and epilepsy in one-half grain doses (0.03). Unof.

Anilism (*an'-il-izm*) [*anilin*]. An acute or chronic disease produced in workmen in anilin factories by the poisonous fumes. The symptoms are debility, vertigo, gastro-intestinal disturbance, and cyanosis.

Animal (*an'-im-al*) [*anima*, the spirit, breath, or life]. An organism capable of ingesting and digesting food. No sharp line of distinction exists between the lowest animals and certain vegetables. The higher animals are distinguished by the power of locomotion and the possession of a nervous system. **A.**

Charcoal, bone-black, ivory-black, etc., is the product of the calcining of bones in closed vessels. **A.** Chemistry, that concerning itself with the composition of animal bodies. **A.** Electricity, electricity generated in the body. **A.** Gum, $C_{12}H_{20}O_{16} + 2H_2O$. A substance prepared from mucin by Landwehr, and so named on account of its

resemblance to the gum of commerce. It occurs in many tissues of the body, is soluble in water, and in alkaline solution readily dissolves cupric oxide, the solution not being reduced on boiling. It yields no coloration with iodine, and is very feebly dextrorotatory. **A.** Heat, the normal temperature of the body in man—about $98.5^\circ F.$ ($37^\circ C.$). **A.** Magnetism, mesmerism, hypnotism. **A.** Starch. See *Glycogen*.

Animalcule (*an-im-al'-kûl*) [*animakulum*, a minute animal]. An animal organism so small as to require the microscope for its examination.

Anime (*an'-im-e*) [Fr., *anime*, origin doubtful]. A name of various resins, especially that of *Hymenaea courbaril*, a tree of tropical America; sometimes used in plaster, etc. Unof.

Anion (*an'-e-on*) [*âná*, up; *îdros*, going]. In electrolysis, an electro-negative element.

Aniridia (*an-i-rid'-e-ah*) [*ân*, priv.; *îros*, the rainbow]. Absence or defect of the iris.

Anisated (*an'-is-a-ted*) [*anisum*, anise]. Containing anise.

Anise (*an'-is*). See *Anisum*.

Anisic Acid, $C_8H_8O_3$. Methyl-para-oxybenzoic acid, an oxidation product of anethol. It is antiseptic and antipyretic, and is used in the treatment of wounds and acute articular rheumatism. Dose of the sodium salt 15 grains (1.0).

Anisin (*an'-is-in*) [*anisum*, anise], $C_{12}H_{24}N_2O_6$. A crystalline alkaloid, a derivative of anise.

Anisocoria (*an-is-o-ko'-re-ah*) [*ânisos*, unequal; *korê*, pupil]. Inequality of the diameter of the pupils.

Anisol (*an'-is-ol*) [*anisum*, anise], C_7H_8O . Methyl-phenyl ether, produced by heating phenol with potassium and methyl iodide or potassium methyl sulphate in alcoholic solution. It is an ethereal-smelling liquid, boiling at 152° ; its sp. gr. at 15° is 0.991.

Anisomelous (*an-is-om'-el-us*) [*ânisos*, unequal; *melos*, a limb]. Having limbs of unequal length.

Anisometropia (*an-is-o-met-ro'-pe-ah*) [*ânisos*, unequal; *metron*, a measure; *ôps*, the eye]. A difference in the refraction of the two eyes.

Anisometropic (*an-is-o-met-rof'-ik*) [*ânisos*, unequal; *metron*, a measure; *ôps*, the eye]. Affected with anisometropia.

Anisopia (*an-is-ô'-pe-ah*) [*ânisos*, unequal; *ôps*, eye]. Inequality of visual power in the two eyes.

Anisosthenic (*an-is-o-sthen'-ik*) [*ânisos*, unequal; *sthenos*, strength]. Not of equal power; used of pairs of muscles.

Anisotropal, **Anisotropic**, **Anisotropous** (*an-is-o-trop'-al*, *an-is-o-trop'-ik*, *an-is-ot'*

ro-pus) [ἀνισος, unequal; τρόπος, turning]. Not possessing the same light-refracting properties in all directions; a term applied to doubly refracting bodies. In biology, varying in irritability in different parts or organs.

Anisum (*an'-is-um*) [L.]. Anise. The fruit of *Anisum pimpinella*. Its properties are due to a volatile oil. It is slightly stimulant to the heart action. It is useful chiefly to liquefy bronchial secretions, and is therefore a favorite ingredient in cough mixtures. Dose gr. x-xx (0.65-1.32). **A.**, **Aqua**, oil of anise 1, water 500 parts. Dose indefinite. **A.**, **Essentia** (B. P.). Dose ℥x-xx (0.65-1.3). **A.**, **Oleum**, an ingredient in tinct. opii camph. Dose ℥j-v (0.065-0.32). **A.**, **Spiritus**, a ten per cent. solution of the oil in alcohol. Dose ʒj-ij (4.0-8.0).

Anitrogenous (*ah-ni-troj'-en-us*) [ā priv.; nitrogen]. Non-nitrogenous.

Ankle (*ang'-kl*) [ME., *ancle*]. The joint between the leg and the foot. It is a ginglymus joint, with three ligaments, the anterior, internal, and external. **A-bone**, the astragalus. **A. Clonus**, the succession of a number of rhythmic muscular contractions in the calf of the leg, when the foot is suddenly flexed by a pressure upon the sole. It is a symptom of various diseases of the spinal cord, especially those involving the lateral pyramidal tracts. **A.-jerk**. See *A. Clonus*. **A.-joint**. See *Ankle*. **A. Reflex**. See *A. Clonus*.

Ankyloblepharon (*ang-kil-o-blef'-ar-on*) [ἀγκύλη, a thong or loop; βλέφαρον, the eyelid]. The adhesion of the ciliary edges of the eyelids.

Ankylochilia (*ang-kil-o-ki'-le-ah*) [ἀγκύλη, loop; χείλος, lip]. Adhesion of the lips.

Ankylocolpos (*ang-kil-o-kol'-pōs*) [ἀγκύλη, a loop; κόλπος, the vagina]. Atresia of the vagina or vulva.

Ankyloglossia, Ankyloglossum (*ang-kil-o-glos'-e-ah, ang-kil-o-glos'-sum*) [ἀγκύλη, a loop; γλῶσσα, the tongue]. Tongue-tie.

Ankylose (*ang'-kil-ōs*) [ἀγκύλη, a loop]. To be, or to become, consolidated or firmly united.

Ankylosis (*ang-kil-o'-sis*) [ἀγκύλη, a loop]. Union of the bones forming an articulation, resulting in a stiff joint. **A.**, **Extracapsular**, that due to rigidity of the parts external to the joint. **A.**, **False**, or **Spurious**, is due to the rigidity of surrounding parts. **A.**, **Intracapsular**, that due to rigidity of the structures within a joint. **A.**, **True**, or **Bony**, when the connecting material is bone. **A.**, **Ligamentous**, when the medium is fibrous.

Ankylostoma, Ankylostomum (*ang-kil-os'-to-mah, ang-kil-os'-to-mum*) [ἀγκύλος, crooked; στόμα, a mouth]. A genus of

nematoid worms, one species of which, *A. duodenale*, is found in the human intestine.

Ankylostomiasis (*ang-kil-os-to-mi'-as-is*) [ἀγκύλος, crooked; στόμα, a mouth]. A form of pernicious anemia produced by the presence of the parasite *Ankylostoma duodenale* in the human intestine. It is especially prevalent among brickmakers and other workmen in Europe. It is called also *Dochmiasis*, *Brickmakers' Anemia*, *Tunnel Anemia*, *Miners' Cachexia*, *Egyptian Chlorosis*, etc. Male fern and thymol expel the worm.

Ankylotia (*ang-kil-o'-she-ah*) [ἀγκύλη, a noose; οὖς, ear]. Union of the walls of the meatus auditorius.

Anatto (*an-at'-o*). See *Annotto*.

Annectant (*an-ek'-tant*) [*ad*, to; *nectere*, to bind]. Linking or binding together. **A. Convolutions**. See *Convolution*.

Annidalin (*an-id'-al-in*). 1. Dithymol triiodid. A substitute for iodoform and aristol. 2. See *Aristol*.

Annotto (*an-ot'-o*) [native American]. Annatto, annatto. A coloring matter obtained from the pellicles of the seeds of *Bixa orellana*. It is used to color plasters, and as an artificial color for butter.

Annuens (*an'-u-ens*) [*annuere*, to nod]. The rectus capitis anticus minor muscle.

Annular (*an'-u-lar*) [*annulus*, a ring]. Ring-like. **A. Cartilage**, the cricoid cartilage. **A. Finger**, the ring finger. **A. Ligament**, the ligament surrounding the wrist and the ankle. **A. Muscle of Müller**, the circular fibers of the ciliary muscle. **A. Process**, or *Protuberance*, the pons varolii. **A. Reflex** (of macula), a ring-like reflection sometimes seen with the ophthalmoscope around the macula.

Annulus (*an'-u-lus*) [*annulus*, a ring]. A ring-shaped or circular opening. **A. abdominalis**, the external or internal abdominal rings. **A. abdominis**, the inguinal ring. **A. ciliaris**, the boundary between the iris and choroid. **A. fibrosus**, the external part of the intervertebral discs. **A. membranæ tympani**, an incomplete bony ring that forms the fetal auditory process of the temporal bone. **A. osseus**, the tympanic plate. **A. ovalis**, the rounded or oval margin of the *Foramen ovale*. **A. tracheæ**, any tracheal ring. **A. umbilicus**, the umbilical ring.

Anococcygeal (*a-no-kok-sij'-e-al*) [*anus*, the fundament; κόκκυξ, the coccyx]. Pertaining to the anus and the coccyx. **A. Ligament**, a ligament that connects the tip of the coccyx with the external sphincter ani muscle.

Anodal (*an'-o-dal*) [ἀνά, up; ὁδός, a way]. Relating to the anode; electro-positive. **A. Opening Contraction**. See *Contraction*.

Anode (*an'-ôd'*) [*âná*, up; *ôdós*, a way]. The positive pole of a galvanic battery.

Anodontia (*an-o-don'-she-ah*) [*ân* priv.; *ôdôn*, tooth]. Absence of the teeth.

Anodyne (*an'-o-dîn*) [*ân* priv.; *ôdînê*, pain]. A medicine that gives relief from pain. **A.**, Hoffmann's. See *Ether*.

Anoesia (*an-o-ê'-ze-ah*) [*ânôesia*, a want of sense]. Want of understanding.

Anoia (*an-oi'-ah*) [*ânôia*, idiocy]. Synonym of *Idiocy*.

Anomalous (*an-om'-al-us*) [*ânómalos*, not ordinary]. Irregular; characterized by deviation from the common or normal order.

Anomaly (*an-om'-al-é*) [*ânómalia*, irregularity]. A marked deviation from the normal; an abnormal thing or occurrence.

Anonyma (*an-on'-im-ah*) [*ân* priv.; *ônyma*, name]. The innominate artery.

Anonymous (*an-on'-im-us*) [*ân* priv.; *ônyma*, a name]. Nameless. **A. Bone**. See *Innominate*.

Anophthalmia (*an-off-thal'-me-ah*) [*ân* priv.; *ôphthalmos*, eye]. Congenital absence of the eyes. **A. cyclopica**, a congenital malformation in which the eye-socket is very ill-developed and the orbit rudimentary or altogether absent.

Anophthalmus (*an-off-thal'-mus*) [*ân* priv.; *ôphthalmos*, eye]. 1. Congenital absence of the eyes. 2. A person born without eyes.

Anopia (*an-o'-pe-ah*) [*ân* priv.; *ôps*, the eye]. Absence of sight, especially that due to defect of the eyes.

Anorchia (*an-or'-ke-ah*). See *Anorchism*.

Anorchism (*an'-or-kism*) [*ân* priv.; *ôrches*, the testicle]. Absence of the testicles.

Anorexia (*an-or-ek'-se-ah*) [*ân* priv.; *ôrêxis*, appetite]. Absence of appetite. **A. nervosa**, an hysterical affection occurring chiefly in young neurotic females, and characterized by a great aversion to food.

Anorthopia (*an-or-tho'-pe-ah*) [*ân* priv.; *ôrthós*, straight; *ôps*, vision]. 1. A defect in vision in which straight lines do not seem straight, and parallelism or symmetry is not properly perceived. 2. Squinting; obliquity of vision.

Anorthosis (*an-or-tho'-sis*) [*ân* priv.; *ôrthosis*, a making straight]. Absence or defect of erectility.

Anosmia (*an-oi'-me-ah*) [*ân* priv.; *ôsmê*, smell]. Absence of the sense of smell.

Anosphrasia (*an-os-fra'-ze-ah*) [*ân* priv.; *ôsphrasia*, smell]. Absence of the sense of smell.

Anospinal (*a-no-spi'-nal*) [*anus*, the fundament; *spina*, the backbone]. Relating to the anus and the spinal cord. **A. Center**, a center that controls the anal sphincters. It is situated in the lumbar portion of the spinal cord.

Anotus (*an-o'-tus*) [*ân* priv.; *ôûs*, the ear]. An earless monstrosity.

Anovesical (*a-no-ves'-ik-al*) [*anus*, the fundament; *vesica*, the bladder]. Pertaining conjointly to the anus and urinary bladder.

Ansa (*an'-sah*) [*La*, a "handle."]. A loop. **A. capitis**, the zygomatic arch. **A.**, Haller's, the loop formed by the nerve joining the facial and glosso-pharyngeal nerves. **A.**, Henle's. See *Henle's Loop*. **A. hypoglossi**, a loop formed at the side of the neck by the junction of the descendens noni nerve with branches of the second and third cervical nerves. **A. sacralis**, a loop joining the ganglion impar with the sympathetic trunks of the two sides. **A. of Vieussens**, a loop extending from the third cervical ganglion and surrounding the subclavian artery. **A. of Wrisberg**, the nerve joining the right great splanchnic and right pneumogastric nerves.

Anserine (*an'-ser-in*) [*anser*, a goose]. Resembling a goose. **A. Disease**, a wasting of the muscles of the hands, rendering the tendons unduly prominent, and suggesting the appearance of a goose's foot. **A. Skin**, goose skin.

Antacid (*ant-as'-id*) [*anti*, against; *acidus*, acid]. 1. Neutralizing acidity. 2. A substance counteracting or neutralizing acidity.

Antacidin (*ant-as'-id-in*). Saccharate of lime.

Antagonism (*an-tag'-on-izm*) [*ántagonisthai*, to struggle against]. Opposition; opposed action, as of two sets of muscles or of two remedial agents.

Antagonist (*an-tag'-o-nist*) [*ántagonistês*, counteracting]. A term applied to a drug that neutralizes the therapeutic effects of another. In anatomy, a muscle that acts in opposition to another. **A.**, **Associated**, a name given to that muscle of a healthy eye that turns the globe in the same direction as the affected muscle of the opposite eye would, if normal, turn the eye to which it belongs.

Antagonistic (*an-tag-o-nis'-tik*) [*ántagonistês*, counteracting]. Opposing.

Antalgic (*ant-al'-jik*) [*ân*ti, against; *âlgos*, pain]. 1. Relieving pain. 2. A remedy that relieves pain.

Antalkaline (*ant-al'-kal-in*) [*ân*ti, against; *alkali*]. 1. Neutralizing alkalies. 2. An agent neutralizing alkalies, as acids.

Antaphrodisiac (*ant-af-ro-dis'-e-ah*) [*ân*ti, against; *âphrodisia*, sexual desire]. 1. Lessening venereal desire. 2. An agent that lessens the venereal impulse; an aphrodisiac.

Antapoplectic (*ant-ap-o-plek'-tik*) [*ân*ti, against; *âποπληξία*, apoplexy]. Efficient in preventing or treating apoplexy.

Antarthritic (*ant-ar-thrit'-ik*) [*ân*ti, against;

- ἀσθητικός**, gouty]. 1. Relieving gout. 2. A medicine for the relief of gout.
- Antasthenic** (*ant-as-then'-ik*) [*ἀντί*, against; *ἀσθένεια*, weakness]. Tending to correct debility and restore the strength.
- Antasthmatic** (*ant-as-mat'-ik*) [*ἀντί*, against; *ἀσθμα*, short-drawn breath]. 1. Relieving asthma. 2. A medicine serving for the relief of asthma.
- Antebrachium** (*an-te-bra'-ke-um*) [*ante*, before; *brachium*, the arm]. The forearm.
- Antecurvature** (*an-te-ker'-vat-chûr*) [*ante*, forward; *curvatus*, bent]. A forward curvature.
- Anteflexion** (*an-te-flek'-shun*) [*ante*, before; *flectere*, to bend]. A bending forward. **A. of Uterus**, a condition in which the fundus of the uterus is bent forward.
- Antemetetic** (*ant-em-et'-ik*). See *Antiemetic*.
- Ante mortem** (*an'-te mor'-tem*) [L.]. Before death.
- Antenna** (*an-ten'-ah*) [*ἀντεινεῖν*, to stretch out]. In biology, one of the paired, articulated, sensory appendages of the head of an arthropod.
- Ante partum** (*an'-te par'-tum*) [L.]. Before delivery.
- Antephialtic** (*ant-ef-e-al'-tik*) [*ἀντί*, against; *ἐπιμάτης*, nightmare]. Preventive of nightmare.
- Antepileptic** (*ant-ep-il-ep'-tik*) [*ἀντί*, against; *ἐπιληψία*, epilepsy]. Relieving epilepsy.
- Anteprostatic** (*an-te-pros-tat'-ik*) [*ante*, before; *προστάτης*, one who stands before]. Situated before the prostate. **A. Glands**, certain small accessory glands sometimes found between Cowper's gland and the prostate.
- Anterior** (*an-te'-re-or*) [*anterior*, forward]. Situated before or in front of; pertaining to the part of organ situated toward the ventral aspect of the body. **A. Poliomyelitis**, inflammation of the anterior horns of the spinal cord, giving rise to a characteristic paralysis, common in children. **A. Rotation**, the forward turning of the presenting part in labor.
- Antero-** (*an'-te-ro*) [*anterior*, before]. A prefix signifying position in front.
- Antero-inferior** (*an-te-ro-in-fe'-re-or*) [*anterior*, forward; *inferior*, lower]. Situated in front and below.
- Antero-lateral** (*an-te-ro-lat'-er-al*) [*anterior*, before; *latus*, a side]. In front and to or on one side; from the front to one side.
- Antero-parietal** (*an'-te-ro-par-i'-et-al*) [*anterior*, before; *paries*, a wall]. Anterior and also parietal. **A. Area**, the anterior part of the parietal area of the cranium.
- Antero-posterior** (*an'-te-ro-pos-te'-re-or*) [*anterior*, forward; *posterior*, backward]. Extending from before backward.
- Antero-superior** (*an'-te-ro-su-pe'-re-or*) [*anterior*, forward; *superior*, upper]. Situated in front and above.
- Anteversion** (*an-te-ver'-shun*) [*ante*, before; *vertere*, to turn]. A turning forward. **A. of Uterus**, a tilting forward of the uterus.
- Anthelix** (*ant'-he-lik*s or *an'-the-lik*s) [*ανθήλιξ*, the inner curvature of the ear]. The ridge surrounding the concha of the external ear posteriorly.
- Anthelmintic** (*an-thel-min'-tik*) [*ἀντί*, against; *ἐλμυς*, a worm]. 1. Efficacious against worms. 2. A vermicide.
- Anthemis** (*an'-them-is*) [*ἀνθεμῖς*, a flower]. Camomile. The flower-heads of *A. nobilis*, the properties of which are due to a volatile oil, a camphor, and a bitter principle. It is useful in coughs and spasmodic infantile complaints, and is an excellent stomachic tonic. Infusion of \mathfrak{z} iv to Oj, given in doses of \mathfrak{z} j-ij (32.0-64.0). No official preparations. **A., Extract.** (B. P.), dose gr. ij-x (0.13-0.65). **A., Infus.** (B. P.), dose \mathfrak{z} j-iv (32.0-128.0). **A., Oleum**, the volatile oil of camomile. Dose \mathfrak{m} ij-x (0.13-0.65).
- Anthemorrhagic** (*ant-hem-or-aj'-ik*) [*ἀντί*, against; *αἷμα*, blood; *παύω*, a bursting]. Checking or preventing hemorrhage.
- Anthony's Fire**, St. A popular name for erysipelas.
- Anthracemia** (*an-thras-e'-me-ah*) [*ἀνθραξ*, a coal; *αἷμα*, blood]. Wool-sorter's disease; splenic fever of animals; a disease due to the presence in the blood of the *Bacillus anthracis*.
- Anthracene** (*an'-thra-sen*) [*ἀνθραξ*, a coal], $C_{14}H_{10}$. A hydrocarbon formed from many carbon compounds when they are exposed to a high heat; also from coal-tar. It crystallizes in colorless, monoclinic tables, showing a beautiful blue fluorescence; dissolves with difficulty in alcohol and ether, but easily in hot benzene; melts at 213°. It is the base from which artificial alizarin is prepared.
- Anthracia** (*an-thra'-se-ah*) [*ἀνθραξ*, a coal]. A name for diseases characterized by the formation of carbuncles. **A. pestis**, the plague. **A. rubula**. Synonym of *Frambesia*.
- Anthracic** (*an'-thras-ik*) [*ἀνθραξ*, a coal]. Pertaining to or of the nature of anthrax.
- Anthracin** (*an'-thras-in*) [*ἀνθραξ*, a coal]. A toxic ptomain derived from pure cultures of the bacillus of anthrax.
- Anthracoid** (*an'-thrak-oid*) [*ἀνθραξ*, coal; *ειδός*, form]. Resembling anthrax.
- Anthracometer** (*an-thrak-om'-et-er*) [*ἀνθραξ*, coal; *μέτρον*, a measure]. An instrument for estimating the amount of carbon dioxide in the air.
- Anthraconecrosis** (*an-thrak-o-ne-kro'-sis*) [*ἀνθραξ*, coal; *νέκρωσις*, death]. The ne-

crotic transformation of a tissue into a black mass, as in dry gangrene.

Anthracosis (*an-thrak-ō-sis*) [*ἀνθραξ*, carbon; *νόσος*, disease]. "Miner's Lung." A diseased condition of the lung produced by the inhalation of coal dust. It is a form of pneumonokoniosis.

Anthraquinone (*an-thra-kwin-ōn'*) [*ἀνθραξ*, coal; Sp., *quina*, bark], $C_{14}H_{10}O_2$. A substance produced by oxidizing anthracene with HNO_3 . It sublimes in yellow needles, melting at 277° , and is soluble in hot benzene and HNO_3 .

Anthrarobin (*an-thra-ro'-bin*), $C_{14}H_{10}O_2$. A derivative of alizarin, similar to chrysarobin. It is a yellowish-white powder, insoluble in water, but soluble in alcohol and dilute alkaline solutions. It is useful in psoriasis, herpes, pityriasis versicolor. Unof.

Anthrax (*an'-thraks*) [*ἀνθραξ*, a coal or carbuncle]. 1. A carbuncle. 2. An acute infectious disease due to the bacillus anthracis. It is most common in cattle and sheep, but may occur in man. Infection in man may occur (a) through a wound, the resulting lesion being a boil or carbuncle, whence the disease becomes disseminated through the blood. (b) Through the intestinal tract. In this case there are marked local lesions, either such as resemble the dermal boils, or a hemorrhagic infiltration of the mucous membrane. (c) Through the respiratory tract. The results are inflammatory processes in the lungs and pleura. Synonyms: *Miltbrand*, *Charbon*, *Woolsorter's Disease*. A., Symptomatic. See *Black-leg*.

Anthropogenesis (*an-thro-po-jen'-es-is*) [*ἄνθρωπος*, man; *γένεσις*, generation]. The development of man, as a race (*phylogenesis*) and as an individual (*ontogenesis*).

Anthropoglot (*an'-thro-po-glot*) [*ἄνθρωπος*, man; *γλῶσσα*, the tongue]. Human-tongued, as a parrot.

Anthropography (*an-thro-pog'-ra-fee*) [*ἄνθρωπος*, man; *γράφειν*, to write]. A treatise upon the human structure or organism.

Anthropoid (*an'-thro-poid*) [*ἄνθρωπος*, a man; *εἶδος*, form]. Man-like.

Anthropology (*an-thro-pol'-o-je*) [*ἄνθρωπος*, man; *λόγος*, discourse]. The science of man.

Anthropometalism (*an-thro-po-met'-al-izm*) [*ἄνθρωπος*, man; M. E., *metal*]. Hypnotism or the like condition, induced by looking at a metallic disc.

Anthropometry (*an-thro-pom'-et-re*) [*ἄνθρωπος*, man; *μέτρον*, a measure]. The determination of the measurement, weight, strength, and proportions of the parts of the human body.

Anthropomorphic (*an-thro-po-mor'-fik*) [*ἄνθρωπος*, man; *μορφή*, form]. Man-like.

Anthropophagy (*an-thro-poff'-a-je*) [*ἄνθρωπος*, man; *φαγεῖν*, to devour]. 1. Cannibalism. 2. Sexual perversion leading to rape, mutilation, and cannibalism.

Anthropophobia (*an-thro-po-fo'-be-ah*) [*ἄνθρωπος*, man; *φόβος*, fear]. A symptom of mental disease consisting in fear of society.

Anthropotoxin (*an-thro-po-toks'-in*) [*ἄνθρωπος*, man; *τοξικόν*, poison]. The toxic substance supposed to be excreted by the lungs of human beings.

Anthydropic (*ant-hi-drop'-ik*) [*ἀντι*, against; *ἵδωρ*, water]. Effective against dropsy.

Anthyptotic (*ant-hip-not'-ik*) [*ἀντι*, against; *ἕπνοσ*, sleep]. 1. Preventive of sleep. 2. An agent that tends to induce wakefulness.

Anthysteric (*ant-his-ter'-ik*) [*ἀντι*, against; *ἱστέρα*, the womb]. Overcoming hysteria.

Antialbumate (*an-te-al'-bu-mat*) [*ἀντι*, against; *albumen*, white of egg]. Parapeptone; a product of the imperfect digestion of albumin. It is changed by the pancreatic ferment into antipeptone.

Antialbumin (*an-te-al'-bu'-min*) [*ἀντι*, against; *albumen*, white of egg]. One of the products of the action of the digestion of albumin; it is probably one of the preformed substances existing in the proteid molecule.

Antialbumose (*an-te-al'-bu'-mās*) [*ἀντι*, against; *albumen*, white of egg]. One of the albumoses produced by the action of pancreatic juice on albumin. It resembles syntonin or acid-albumin, and is convertible into antipeptone.

Antiafrodisiac (*an-te-af-ro-dis'-e-ak*). See *Antiafrodisiac*.

Antiarin (*an'-te-ar-in*) [Javanese, *antiar* or *antschar*], $C_{14}H_{10}O_2$ + $2H_2O$. The active principle of *Antiaris toxicaria* or *U'pas antiar*, the Javanese poison-tree. It is intensely poisonous and is used as an arrow poison. It is a cardiac depressant. Dose gr. 1/10 (0.00065). Unof.

Antiarthritic (*an-te-ar-thrit'-ik*). Synonym of *Antiarthritic*.

Antiasthmatic (*an-te-as-mat'-ik*). Synonym of *Antasthmatic*.

Antibacterial (*an-te-bak-te'-re-al*) [*ἀντι*, against; *βακτηριον*, a staff, stick]. Acting against bacteria.

Antibiotic (*an-te-bi-ot'-ik*) [*ἀντι*, against; *βίος*, life]. Tending to destroy life.

Antiblenorrhagic (*an-te-ble-nor-aj'-ik*) [*ἀντι*, against; *βλιννος*, mucus; *πηγνύμαι*, to burst]. Efficient in preventing or curing gonorrhea.

Antibrachial (*an-te-bra'-ke-al*) [*ἀντι*, against; *βραχίον*, the arm]. Pertaining to the forearm.

Antibrachium (*an-te-bra'-ke-um*) [*ἀντι*, against; *βραχίον*, the arm]. The forearm; antebrachium.

Anticardium (*an-te-kar'-de-um*) [*ἀντι*, ever

against; *καρδία*, the heart]. The scrobiculus cordis, or pit of the stomach; the infrasternal depression.

Antichirotonus (*an-te-ki-ro't-o-nus*) [*ávri*, against; *χείρ*, hand; *τόνος*, tension]. Forcible and steady inflection of the thumb, seen at times in or before attacks of epilepsy.

Anticholerin (*an-te-kol'-er-in*) [*ávri*, against; *χολέρα*, cholera]. A product isolated by Klebs from cultures of cholera-bacilli. Immunizing and curative properties have been ascribed to it.

Anticipating (*an-tis'-ip-a-ting*) [*anticipare*, to take before]. Occurring before the regular or expected time, as an anticipating intermittent fever, one in which the paroxysms occur earlier on successive days.

Anticlinal (*an-te-kli'-nal*) [*ávri*, against; *κλίνειν*, to slope]. Sloping in opposite directions. **A.** Vertebra, in man, the tenth thoracic vertebra, where the thoracic vertebrae begin to assume the characters of the lumbar.

Anticnemion (*an-tik-ne'-me-on*) [*ávri*, against; *κνήμη*, leg]. The shin or front of the leg.

Anticus (*an-ti'-kus*) [*anticus*, that in front]. Anterior, in front of.

Anticylic Acid. A white, fragrant powder, with a pleasant taste. It is said to be antipyretic. Dose $\frac{1}{16}$ grain (0.0006). Unof.

Antidinic (*an-te-din'-ik*) [*ávri*, against; *δίνω*, a whirl]. Curing or preventing vertigo.

Antidotal (*an-te-do'-tal*) [*ávri*, against; *δοῖός*, given]. Having the nature of an antidote.

Antidote (*an'-te-dôt*) [*ávri*, against; *δοῖός*, given]. An agent preventing or counteracting the action of a poison. **A.**, **Arsenical**, G. Ph., is prepared by dissolving 100 parts of the hydrated sulphate of iron in 250 parts of water, to which 15 parts of burnt magnesia and 250 parts of water are added. **A.**, **Chemic**, one that changes the chemic nature of the poison so that it becomes insoluble or harmless. **A.**, **Mechanical**, one that prevents absorption by holding the poison in mechanical suspension or by coating the stomach. **A.**, **Physiologic**, one that counteracts the physiologic effects of a poison. **A.**, **Universal**, a mixture of one part of dissolved iron sulphate in two parts of magnesia water.

Antidynamic (*an-te-di-nam'-ik*) [*ávri*, against; *δύναμις*, force]. Weakening.

Antidysenteric (*an-te-dis-in-ter'-ik*) [*ávri*, against; *δυσεντερία*, dysentery]. 1. Serviceable against dysentery. 2. A remedy for dysentery.

Antiemetic (*an-te-em-et'-ik*) [*ávri*, against; *εμετικός*, causing vomit]. Preventing emesis; relieving nausea.

5

Antifebrin (*an-te-feb'-rin*) [*ávri*, against; *febris*, a fever], $C_6H_5.C_2H_5O.NH$. The proprietary name of acetanilid or phenylacetamid. A white, crystalline powder insoluble in water, freely soluble in alcohol, ether, and chloroform. It is antipyretic and analgesic. The drug's official name is acetanilidum. Dose gr. v-x (0.3-0.6).

Antiferment (*an-te-fer'-ment*) [*ávri*, against; *fermentum*, leaven]. An agent that prevents fermentation.

Antifermentative (*an-te-fer-men'-ta-tive*) [*ávri*, against; *fermentum*, leaven]. Preventing fermentation.

Antigalactic (*an-te-gal-ak'-tik*) [*ávri*, against; *γάλα*, milk]. 1. Lessening the secretion of milk. 2. A drug that lessens the secretion of milk.

Antihelix (*an-te-he'-lik*). See *Anthelix*.

Antihidrotic (*an-te-hi-drot'-ik*) [*ávri*, against; *ιδρώς*, sweat]. 1. Diminishing the secretion of sweat. 2. An agent lessening the secretion of sweat.

Antihydropin (*an-te-hi'-dro-pin*) [*ávri*, against; *ὕδωρ*, water]. A crystalline principle obtainable from the common cockroach, *Blatta (Periplaneta) orientalis*, and said to be diuretic. Dose gr. x-xx (0.6-1.3).

Antikamnia (*an-te-kam'-ne-ak*) [*ávri*, against; *κάμνειν*, to suffer pain]. A proprietary remedy said to be composed of sodium bicarbonate, acetanilid, and caffeine. It is used as an analgesic in doses of gr. v-x (0.32-0.65).

Antikol (*an-te-tik-ol*). A proprietary antipyretic mixture said to contain acetanilid, sodium bicarbonate, and tartaric acid.

Antilemic, or **Antilœmic**, **Antiloimic** (*an-te-le'-mik*, or *an-te-loi'-mik*) [*ávri*, against; *λοιμός*, the plague]. Efficacious against the plague or other pestilence.

Antileptic (*an-ti-lep'-tik*) [*ἀντιληψις*, a receiving in return]. Revulsive.

Antilithic (*an-te-lith'-ik*) [*ávri*, against; *λίθος*, a stone]. 1. Efficacious against calculus. 2. An agent preventing the deposit of urinary sediment.

Antilœmic. See *Antilemic*.

Antiluetic (*an-te-lu-et'-ik*) [*ávri*, against; *λues*, the plague; syphilis]. Efficacious against syphilis.

Antilyssic (*an-te-lis'-ik*) [*ávri*, against; *λύσσα*, rabies]. 1. Tending to cure rabies. 2. A remedy for rabies.

Antimiasmatic (*an-te-mi-as-mat'-ik*) [*ávri*, against; *μίασμα*, exhalation]. Preventive of malaria.

Antimonial (*an-te-mo'-ne-al*) [*antimonium*, antimony]. Containing antimony.

Antimonic (*an-te-mon'-ik*) [*antimonium*, antimony]. A term applied to those compounds of antimony that correspond to its higher oxid.

Antimonious (*an-te-mo'-ne-us*) [*antimonium*, antimony]. A term denoting those compounds of antimony that correspond to its lower oxid.

Antimony (*an'-te-mo-ne*) [*antimonium*]. Sb = 122; quantivalence III and V. A metallic, crystalline element possessing a bluish-white luster. The symbol Sb is derived from the old name, *stibium*. Antimony is found native, as the sulphid, Sb_2S_3 , as the oxid, and is a constituent of many minerals. It is used commercially chiefly for making alloys. Type-metal, Britannia-metal, and Babbitt's anti-friction metal are alloys of antimony. In medicine antimonium salts are used less frequently than formerly. The salts are cardiac and arterial depressants, diaphoretic and emetic, and in large doses powerful gastro-intestinal irritants, producing symptoms resembling Asiatic cholera. Antimony has been used as an antiphlogistic in sthenic inflammation, as a diaphoretic and expectorant, and as an emetic. **A. chlorid**, $SbCl_3$, the "butter" of antimony; a strong caustic. Unof. **A.**, *Pilulæ Compositæ*, Plummer's pills, which contain calomel and sulphuretted antimony, aa gr. ss (0.032). **A. tri-oxid**, antimonious acid, Sb_2O_3 ; soluble in hydrochloric and tartaric acids. Dose gr. j-ij (0.065-0.13). It is an ingredient of James' powder. **A. oxychlorid**, the "powder of algaroth;" now little used. **A. pentoxid**, Sb_2O_5 , antimonic acid, combines with bases to form antimonates. **A. et Potassii tartas** (*A. tartaratum*) (B. P.), $2KsOOC_4H_4O_6 \cdot H_2O$, tartrate of antimony and potassium, "tartar emetic." Dose gr. $\frac{1}{16}$ - $\frac{1}{8}$ (0.004-0.016). **A.**, *Pulvis*, powder of antimony, James' powder, consists of antimonious oxid 33, and calcium phosphate 67 parts, and is diaphoretic; in large doses, emetic and cathartic. Dose gr. iij-vij (0.2-0.5). **A. sulphid**, Sb_2S_3 , black sulphid of antimony. Dose gr. $\frac{1}{4}$ -j (0.016-0.065). **A. sulphuratum**, the sulphid with a small but indefinite amount of the oxid. Dose gr. j-v (0.065-0.32). **Syr. Scillæ Comp.**, Cox's hive mixture, hive syrup. Each f $\frac{3}{4}$ j contains gr. $\frac{3}{4}$ of tartar emetic. Dose m $\frac{v}{f\frac{3}{4}j}$ (0.32-4.0) cautiously. **A.**, *Vinum*, wine of antimony. Boiling water 60, tartar emetic 4, stronger white wine 1000 parts. It contains about 2 grains of tartar emetic to the ounce. Dose m $\frac{v}{xv}$ (0.32-1.0).

Antimycotic (*an-te-mi-kot'-ik*) [*avri*, against; *μύκη*, a fungus]. Destructive to microorganisms.

Antinarcotic (*an-te-nar-kot'-ik*) [*avri*, against; *νάρκωσις*, a benumbing]. Preventing narcosis.

Antineuralgic (*an-te-nu-rat'-ik*) [*avri*, against; *νεῖρον*, a nerve; *ἄλγος*, pain]. Overcoming neuralgia.

Antiparasitic (*an-te-par-as-it'-ik*) [*avri*, against; *παράσιτος*, a parasite]. 1. Destroying parasites. 2. An agent destroying parasites.

Antipeptone (*an-te-pep'-tōn*) [*avri*, against; *πέπτειν*, to cook, digest]. A variety of peptone not acted upon by trypsin.

Antiperiodic (*an-te-pe-re-od'-ik*) [*avri*, against; *περίοδος*, a going round]. 1. Preventing periodic attacks of a disease. 2. A remedy against periodic disease. **A. Tincture**. See *Warburg's Tincture*.

Antiperistalsis (*an-te-per-is-tal'-sis*) [*avri*, against; *περί*, around; *στάσις*, compression]. Reversed peristalsis.

Antiperistaltic (*an-te-per-is-tal'-tik*) [*avri*, against; *περί*, around; *στάσις*, compression]. Relating to antiperistalsis.

Antiphlogistic (*an-te-flo-jis'-tik*) [*avri*, against; *φλόγωσις*, inflammatory heat]. 1. Counteracting fever. 2. An agent subduing or reducing inflammation or fever. **A. Treatment**, consists in bloodletting, the application of cold, the administration of antipyretics, etc.

Antiplastic (*an-te-plas'-tik*) [*avri*, against; *πλάσσειν*, to form]. 1. Unfavorable to granulation or to the healing process. 2. An agent impoverishing the blood.

Antipneumotoxin (*an-te-nu-mo-toks'-in*). See *Pneumotoxin*.

Antipodal Cells (*an-tip'-o-dal*) [*avri*, opposite; *πόδις*, foot]. A term applied to a group of four cells formed in the lower end of the embryo-sac opposite to the cells constituting the egg-apparatus.

Antipraxia (*an-te-prak'-e-ah*) [*avri*, against; *πράσσειν*, to do]. Antagonism of functions or of symptoms.

Antipruritic (*an-te-pru-rit'-ik*) [*avri*, against; *pruritus*, itching]. 1. Relieving the sensation of itching. 2. A drug that relieves the sensation of itching.

Antipsoric (*an-te-psy'-rik*) [*avri*, against; *ψώρα*, the itch]. Effective against itching or the itch.

Antipyresis (*an-te-pi-re'-sis*) [*avri*, against; *πυρεσσειν*, to be feverish]. The reduction of fever by means of antipyretics.

Antipyretic (*an-te-pi-rel'-ik*) [*avri*, against; *πυρετός*, fever]. 1. Cooling; lowering the temperature. 2. An agent reducing temperature. The most important antipyretic agents are cold, diaphoretics, and the newer remedies, many of which are coal-tar products, such as antipyrin, acetanilid, phenacetin, etc.

Antipyrin (*an-te-pi'-rin*) [*avri*, against; *πυρετός*, fever]. $C_{11}H_{12}N_2O$. Phenazon. The scientific name is dimethyl-oxy-phenic-phenyl-dimethylpyrazolon, or dehydrodimethylphenylpyrazin. An alkaloidal pro-

duct of the destructive distillation of coal-tar. It may be produced by heating acetoacetic ester with methyl-phenyl-hydrazin. It is a grayish or reddish-white crystalline powder, slightly bitter, soluble in water, alcohol, and chloroform, and crystallizes from ether in shining leaflets, melting at 113°. It reduces the temperature, causes sweating, at times vomiting, peculiar eruptions, pruritus, coryza, etc. Not rarely a cyanotic condition of the face and hands is produced. Antipyrin is incompatible with nitrous compounds. It is a powerful antipyretic and analgesic. Dose gr. v-xv (0.3-1.0). **A. Salicylate.** See *Salipyrin*.

Antirabic (*an-te-rab'-ik*) [*ávri*, against; *rabies*, madness]. Preventing or curing rabies.

Antirheumatic (*an-te-ru-mat'-ik*) [*ávri*, against; *ρευματικός*, of or pertaining to a flux]. Preventing or curing rheumatism.

Antirheumatin (*an-te-ru'-mat-in*). A combination of sodium salicylate and methylene blue. It occurs as blue prismatic crystals easily soluble in water and alcohol, and tasting very much like sodium salicylate. It colors the urine blue or green.

Antirrhinum (*an-te-ri'-num*) [L.]. A genus of scrophulariaceous plants. *A. linaria*, called also *Linaria vulgaris*, Toad-flax, Ramsted, Butter-and-Eggs; an herbaceous plant of Europe and North America; diuretic, cathartic, and irritant; used as a poultice and fomentation. Unof.

Antiscorbutic (*an-te-skor-bu'-tik*) [*ávri*, against; *scorbutus*, scurvy]. 1. Effective against scurvy. 2. A remedy useful in scurvy.

Antiseptin (*an-te-sep'-sin*) [*ávri*, against; *σήψις*, putrefaction], $C_6H_5BrNHC_2H_5O$. Asepsin; bromated acetanilid; soluble in alcohol and ether, insoluble in water. It is antipyretic, analgesic, and antiseptic. Dose gr. vj-vij (0.39-0.45).

Antisepsis (*an-te-sep'-sis*) [*ávri*, against; *σήψις*, putrefaction]. Exclusion of the germs that cause putrefaction.

Antiseptic (*an-te-sep'-tic*) [*ávri*, against; *σήψις*, putrefaction]. 1. Having power to prevent the growth of the bacteria upon which putrefaction depends. 2. A remedy or agent that prevents the development of bacteria. The principal antiseptics in use are corrosive sublimate, creolin, carbolic acid, iodoform, thymol, salicylic acid, boric acid, etc. **A. Gauze**, open cotton cloth charged with an antiseptic. **A. Ligature**, catgut or other material rendered aseptic by soaking in antiseptic solutions. **A. Treatment of Wounds** looks to thorough antisepsis as regards the wound, the instruments, the operator's hands, the dressings, etc.

Antisepticism (*an-te-sep'-tis-izm*) [*ávri*, against; *σήψις*, putrefaction]. The theory

or systematic employment of antiseptic methods.

Antisepticize (*an-te-sep'-tis-iz*) [*ávri*, against; *σήψις*, putrefaction]. To render antiseptic; to treat with antiseptics.

Antiseptin (*an-te-sep'-tin*) [*ávri*, against; *σήψις*, putrefaction]. Zinc boro-thymo-iodid. It consists of 85 parts zinc sulphate, 2.5 parts each of zinc iodid and thymol, and 10 parts boric acid. It is an antiseptic. Unof.

Antiseptol (*an-te-sep'-tol*) [*ávri*, against; *σήψις*, putrefaction]. Cinchonin iodosulphate, an odorless and fairly effective substitute for iodoform.

Antisialagogue (*an-te-si-al'-a-gog*) [*ávri*, against; *σάλων*, saliva; *ἀγωγός*, leading]. 1. Preventing or checking salivation. 2. A remedy that is effective against salivation.

Antisialic (*an-te-si-al'-ik*) [*ávri*, against; *σάλων*, saliva]. 1. Checking the flow of saliva. 2. An agent that checks the secretion of saliva.

Antispasmin (*an-te-spas'-min*). A whitish powder, consisting of a mixture of narcein-sodium and sodium salicylate. It is sedative and hypnotic, and has been used in laryngismus stridulus and whooping-cough. Dose gr. ¼-iv (0.01-0.26).

Antispasmodic (*an-te-spas-mod'-ik*) [*ávri*, against; *σπασμός*, a spasm]. 1. Tending to relieve spasm. 2. An agent relieving convulsions or spasmodic pains, as the narcotics, the nitrites, etc.

Antispastic (*an-te-spas'-tik*) [*ávri*, against; *σπαστικός*, drawing]. 1. Revulsive; counter-irritant. 2. Antispasmodic. 3. A revulsive agent.

Antisudoral (*an-te-su'-dor-al*) [*ávri*, against; *sudor*, sweat]. Checking the secretion of sweat.

Antisyphilitic (*an-te-sif-il-il'-ik*) [*ávri*, against; *syphilis*]. 1. Effective against syphilis. 2. A remedy used in the treatment of syphilis.

Antithenar (*an-tith'-en-ar*) [*ávri*, against; *θέναρ*, the hollow of the hand or foot]. 1. Opposite to thenar. 2. A muscle that extends the thumb, or opposes it to the hand: an antithenar muscle. **A. Eminence**, the border of the palm of the hand from the base of the little finger to the wrist. **A. Muscles**, of the toe and of the thumb; the abductor pollicis pedis and the flexor brevis pollicis manus; also, the first dorsal interosseous muscle.

Antithermic (*an-te-ther'-mik*) [*ávri*, against; *θερμη*, heat]. Cooling; antipyretic.

Antithermin (*an-te-ther'-min*) [*ávri*, against; *θερμη*, heat], $C_{11}H_{14}O_2N_2$. Phenylhydrazin-levalulinic acid, a coal-tar derivative used as an antipyretic, analgesic, and antiseptic. Dose gr. v (0.3). Unof.

Antitoxic (*an-te-toks'-ik*) [*ἀντί*, against; *τοξικόν*, poison]. Antidotal; counteracting poisons.

Antitoxin (*an-te-toks'-in*) [*ἀντί*, against; *τοξικόν*, poison]. A counter poison or antidote elaborated by the body to counteract the toxins of bacteria. According to some authorities, antitoxins are, like the toxins, bacterial products. Antitoxins are used in the treatment of certain infectious diseases and also to confer immunity against these diseases.

Antitragic (*an-te-traf'-ik*) [*ἀντί*, opposite to; *τράγος*, the tragus]. Pertaining to the antitragus. **A.** Muscle, a mere rudiment in man; it arises from the antitragus, and extends to the cauda of the helix.

Antitragus (*an-te-tra'-gus*) [*ἀντί*, opposite to; *τράγος*, the tragus]. An eminence of the external ear opposite the tragus.

Antitrope (*an'-tro-trop*) [*ἀντί*, against; *τροπή*, to turn]. Any organ set over against another to form a symmetric pair. Thus, the right eye is an antitrope to the left.

Antivenene (*an-te-ven'-en*) [*ἀντί*, against; *venenum*, poison]. A term given to the blood-serum of animals rendered immune against snake-poison, on account of its antidotal properties.

Antizymotic (*an-te-si-mot'-ik*) [*ἀντί*, against; *ζύμωσις*, fermentation]. 1. Preventing or checking fermentation. 2. An agent preventing the process of fermentation; an antiferment.

Antraccele (*an'-tra-sel'*) [*antrum*, a cavity; *κίλη*, a tumor]. Dropsy of the antrum; an accumulation of fluid in the maxillary sinus.

Antral (*an'-tral*) [*antrum*, a cave]. Relating to an antrum.

Antrectomy (*an-tek'-to-me*) [*ἀντρεξ*, antrum; *ἐκτομή*, excision]. Surgical removal of the walls of the antrum.

Antritis (*an-tri'-tis*) [*ἀντρεξ*, a cave; *itis*, inflammation]. Inflammation of an antrum.

Antrophore (*an'-tro-för*) [*ἀντρεξ*, a cavity; *φέρω*, to bear]. A soluble, medicated bougie.

Antrotympanic (*an-tro-tim-pan'-ik*) [*ἀντρεξ*, a cave; *τύμπανον*, a drum]. Relating to the cavity of the tympanum and to the tympanic antrum.

Antrum (*an'-trum*) [*L.*]. A cavity or hollow space, especially in a bone. **A.** *cardiacum*, a dilatation of the lower end of the esophagus. It has been noted in cases of rumination, and in other conditions. **A.** *ethmoidale*, the ethmoidal sinus. **A.** *highmorianum*, Antrum of Highmore, a cavity in the superior maxillary bone. **A.** *highmori testis*. See *Mediastinum testis*. **A.** *Mastoid*, the hollow space beneath the roof of the mastoid process. **A.** *Maxillary*.

See *A. highmorianum*. **A.** *pyloricum willisii*, the cavity of the pylorus.

Anuresis (*an-u-re'-sis*) [*ἀν* priv.; *ούρην*, urine]. Anuria.

Anuretic (*an-u-ret'-ik*) [*ἀν* priv.; *ούρην*, urine]. Pertaining to or affected with anuria.

Anuria (*an-u'-re-ah*) [*ἀν* priv.; *ούρην*, urine]. Suppression of the urine.

Anuric (*an-u'-rik*) [*ἀν* priv.; *ούρην*, urine]. Pertaining to anuria.

Anurous (*an-u'-rus*) [*ἀν* priv.; *οὐρά*, a tail]. Without a tail.

Anus (*a'-nus*) [*anus*, the fundament]. The extremity of the rectum; the lower opening of the alimentary canal. **A.** *Artificial*, an opening established from the bowel to the exterior at a point above the normal anus, most commonly from the colon, either in the lumbar or the iliac region. **A.** *Fissure* of, a slight tear in the mucous membrane at the anus, usually due to passage of hardened feces. It is very painful. **A.** *Fistula* of, fistula in ano, a sinus opening from the rectum into the connective tissue about the rectum, or discharging externally. **A.** *Imperforate*, absence of the anus, the natural opening being closed by a membranous septum.

Anvil (*an'-vil*). See *Incus*.

Anydremia, **Anydræmia** (*an-id-re'-me-ah*) [*ἀν* priv.; *ὕδωρ*, water; *αἷμα*, blood]. A deficiency of water in the blood.

Anypnia (*an-ip'-ne-ah*) [*ἀν* priv.; *ἕπνοα*, sleep]. Sleeplessness.

Aorta (*a-ort'-ah*) [*ἀορτή*, aorta]. The large vessel arising from the left ventricle and distributing by its branches arterial blood to every part of the body. It ends by bifurcating into the two common iliacs at the fourth lumbar vertebra. *The arch*, that extending from the heart to the third dorsal vertebra, is divided into an *ascending*, a *transverse*, and a *descending* part. The *thoracic* portion extends to the diaphragm; the *abdominal* to the bifurcation.

Aortal (*a-ort'-al*) [*ἀορτή*, the aorta]. Relating to the aorta.

Aortic (*a-ort'-ik*) [*ἀορτή*, the aorta]. Pertaining to the aorta. **A.** *Arch*. See *Aorta*. **A.** *Arches*. See *Arch*. **A.** *Foramen*. See *A. Opening of Diaphragm*. **A.** *Murmur*. A murmur produced by disease of the aortic valves. **A.** *Opening of Diaphragm*, the aperture in, or really behind, the diaphragm, through which the aorta passes. **A.** *Opening of Heart*, the opening between the heart and the aorta. **A.** *Plexus*, the plexus of sympathetic nerves, situated on the front and sides of the aorta, between the origins of the superior and inferior mesenteric arteries. **A.** *Sinus*, a deep depression between the leaflets of the aortic valve and the

aortic wall. **A. Valve**, the three semilunar valves closing the aortic opening during the cardiac diastole.

Aortitis (*a-ort-i'-tis*) [*ἀορτή*, the aorta; *τις*, inflammation]. Inflammation of the aorta.

Aortomalacia, or **Aortomalaxia** (*a-ort-o-mal-a'-se-ah*, or *-aks'-e-ah*) [*ἀορτή*, aorta; *μαλαξία*, softening]. Softening of the aorta.

Aortostenosis (*a-ort-o-sten-o'-sis*) [*ἀορτή*, aorta; *στενός*, narrow]. Stenosis or narrowing of the aorta.

Apatropin (*ap-at'-ro-pin*) [*ἀπό*, from; *ἀτροπος*, unchangeable], $C_{11}H_{11}NO_2$. A compound derived from atropin by the action of nitric acid. It is said to produce peculiar convulsions.

Ape (*ap*) [*ΜΕ.*, *ape*]. A man-like monkey.

A. Fissures, those fissures of the human brain that are also found in apes. **A.-hand**, a peculiar shape of the hand produced by the wasting of the thumb-muscles; it is seen in some cases of progressive muscular atrophy.

Apellous (*ah-pell'-us*) [*ἀ* priv.; *πέλλα*, skin]. Skinless.

Apepsia (*ah-pep'-se-ah*) [*ἀ* priv.; *πέπτειν*, to digest]. Cessation or absence of the digestive function. **A.**, Hysterical condition of apesia due to hysteria. It is also called *Hysterical Anorexia*. **A. nervosa**. See *Anorexia nervosa*.

Apeptic (*ah-pep'-tik*) [*ἀ* priv.; *πέπτειν*, to digest]. Affected with apesia.

Aperient (*ap-e'-re-ent*) [*aperire*, to open]. 1. Laxative; mildly purgative. 2. A mild purgative; a laxative.

Aperistalsis (*ah-per-is-tal'-sis*) [*ἀ* priv.; *περί*, around; *στάσις*, constriction]. Cessation of the peristaltic movements of the intestine.

Aperture (*ap'-er-chûr*) [*apertura*, an opening]. An opening. **A.**, Angular, in the microscope, the angle formed between a luminous point placed in focus and the most divergent rays that are capable of passing through the entire system of an objective. **A.**, Numeric, the capacity of an objective for admitting rays from the object and transmitting them to the image.

Apex (*a'-peks*) [*apex*, the extreme end of a thing]. The summit or top of anything; the point or extremity of a cone. **A. Beat**, the impulse of the heart felt in the fifth intercostal space, about $3\frac{1}{2}$ inches from the middle of the sternum. **A. Murmur**, a murmur heard over the apex of the heart. **A. of the Lung**, the upper extremity of the lung behind the border of the first rib.

Aphacia (*ah-fa'-se-ah*). Same as *Aphakia*.

Aphacic (*ah-fa'-sik*). Syn. of *Aphakic*.

Aphagia (*ah-fa'-je-ah*) [*ἀ* priv.; *φαγεῖν*, to eat]. Inability to eat or to swallow.

Aphakia (*ah-fa'-ke-ah*) [*ἀ* priv.; *φακός*, a

lens], the crystalline lens]. The condition of an eye without the lens.

Aphakic (*ah-fa'-kik*) [*ἀ* priv.; *φακός*, the lens]. Not possessing a crystalline lens.

Aphasia (*ah-fa'-se-ah*) [*ἀ* priv.; *φάσις*, speech]. Partial or complete loss of the power of expressing ideas by means of speech or writing. Aphasia may be either motor or sensory. **Motor or Ataxic Aphasia** consists in a loss of speech owing to inability to execute the various movements of the mouth necessary to speech, the muscles not being properly coordinated, owing to disease of the cortical center. It is usually associated with *agraphia*, "aphasia of the hand," inability to write, and right-sided hemiplegia. Some aphasics can write, but are unable to articulate words or sentences; this variety is variously named *aphemia*, *alalia*, or *anarthria*, according as the impairment of speech is more or less marked. Charcot supposes the center for articulate language divided into four sub-centers, a visual center for words, an auditory center for words, a motor center of articulate language, and a motor center of written language. Lesions of one or more of these centers produce the characteristic forms of aphasia, all of which have clinical exemplifications. **Sensory Aphasia**, or **Amnesia** is the loss of memory for words, and may exist alone or in association with motor aphasia. **Amnesia** appears clinically in three distinct forms: 1. Simple loss of memory of words. 2. *Word-deafness*, or inability to understand spoken words (there is usually some paraphasia connected with this form). 3. *Word-blindness*, or inability to understand written or printed words. **Paraphasia**, a form of aphasia in which there is inability to connect ideas with the proper words to express the ideas. **Agrammatism**, a phenomenon of aphasia consisting in the inability to form words grammatically, or the suppression of certain words of a phrase. **Ataxaphasia**, inability to arrange words synthetically into sentences. **Bradyphasia**, abnormal slowness of speech, from pathologic cause. **Tumultus sermonis**, a stuttering manner of reading, from pathologic cause. **A.**, *Conduction*, such as is due to defect in some commissural connection between centers. **A.**, *Gibberish*, aphasia in which the patient can utter many words, but uses them in such a way that they express no meaning.

Aphasiac (*ah-fa'-se-ak*) [*ἀ* priv.; *φάσις*, speech]. One who is aphasic.

Aphasic (*ah-fa'-zik*) [*ἀ* priv.; *φάσις*, speech]. Relating to or affected with aphasia.

Aphemia (*ah-fe'-me-ah*) [*α* priv.; *φήμη*, voice]. Motor aphasia; inability to articulate words or sentences, from centric and not from peripheral disease. See *Aphasia*.

Aphemic (*ah-fem'-ik*) [*á* priv.; *φῆμη*, voice]. Relating to or affected with aphemia.

Aphonia (*ah-fō'-ne-ah*) [*á* priv.; *φωνή*, voice]. Loss of speech due to some peripheral lesion, as in clergyman's sore throat; it may be due to hysteria.

Aphonic (*ah-fon'-ik*) [*á* priv.; *φωνή*, voice]. Speechless; voiceless.

Aphrasia (*ah-fra'-ze-ah*) [*á* priv.; *φράζειν*, to utter]. Absence of the power to utter connected phrases.

Aphrodisia (*af-ro-diz'-e-ah*) [*Ἀφροδίτη*, Venus]. Sexual desire, especially when morbid, or immoderate; sexual congress.

Aphrodisiac (*af-ro-diz'-e-ah*) [*Ἀφροδίτη*, Venus]. 1. Stimulating the sexual appetite; erotic. 2. An agent stimulating the sexual passion.

Aphtha (*af'-thah*) [*ἀφθα*, an eruption]. An eruption; an ulcer.

Aphthæ (*af'-the*) [*ἀφθα*, an eruption]. Also called *Aphthous stomatitis*. A form of stomatitis characterized by the presence of small white vesicles in the mouth, occurring chiefly in children under three years of age, and supposed to be due to a special microorganism. **A.**, Bednar's, two symmetrically placed ulcers seen at times on the hard palate of cachectic infants, one on either side of the mesial line. **A.**, Cachectic, aphthæ appearing beneath the tongue, and associated with grave constitutional symptoms; Riga's disease.

Aphthoid (*af'-thoid*) [*ἀφθα*, an eruption]. Resembling aphthæ.

Aphthongia (*af-thong'-ge-ah*) [*á* priv.; *φθόγγος*, a sound]. A peculiar form of aphasia due to spasm of the muscles supplied by the hypoglossal nerve.

Aphthous (*af'-thus*) [*ἀφθα*, an eruption]. Pertaining to or affected with aphthæ.

Apical (*á'-pik-al*) [*apex*, the top]. Pertaining to the apex.

Apices (*á'-pis-es*) [*L.*]. Plural of *Apex*.

Apinoid (*ap'-in-oid*) [*á* priv.; *πῖνος*, dirt; *εἶδος*, form]. Clean; not foul. **A.** Cancer, scirrhus; so called from its cleanly section.

Apiol (*ap'-e-ol*) [*apium*, parsley; *oleum*, oil], $C_{12}H_{14}O_4$. A principle occurring in parsley seeds; it crystallizes in long white needles, with a slight parsley odor; melts at 30° C. (86° F.), and boils at 294° C. (572° F.). It is used in dysmenorrhea and in malaria. In large doses it produces ringing in the ears and frontal headache. It is also called *Parsley-camphor*. Dose gr. x-xv (0.65-1.0). Unof.

Aplacental (*ah-plas-en'-tal*) [*á* priv.; *placenta*, a cake]. Destitute of placenta.

Aplanatic (*ah-plan-at'-ik*) [*á* priv.; *πλανάειν*, to wander]. Not wandering; rectilinear. **A.** Focus, that focus of a lens the rays

from which do not undergo spheric aberration in their passage through the lens. **A.** Lens, a lens corrected for aberration of light and color; a rectilinear lens.

Aplasia (*ah-pla'-se-ah*) [*á* priv.; *πλάσσειν*, to form]. Incomplete or defective development; also termed *agenesis*.

Aplastic (*ah-plas'-tik*) [*á* priv.; *πλάσσειν*, to form]. 1. Structureless, formless. 2. Incapable of forming new tissue. **A.** Lymph, a non-fibrinous material incapable of coagulation or organization.

Apnea (*ap-ne'-ah*) [*á* priv.; *πνέειν*, to breathe]. 1. A transient cessation of respiration from an over-abundance of oxygen, as, e. g., after forcible respiration. 2. Asphyxia.

Apneumatosis (*ap-nu-mat'-o'-sis*) [*á* priv.; *πνευμάτωσις*, inflation]. Collapse of the air cells.

Apnœa (*ap-ne'-ah*). See *Apnea*.

Apoaconitin (*ap-o-ak-en'-it-in*) [*ἀπό*, from; *aconitum*, aconite], $C_{23}H_{41}NO_{11}$. An alkaloid prepared from aconitin by dehydration.

Apoatropin (*ap-o-at'-ro-pin*) [*ἀπό*, from; *ἀτροπος*, unchangeable], $C_{17}H_{21}NO_3$. An alkaloid obtained by the action of HNO_3 on atropin.

Apochromatic (*ap-o-kro-mat'-ik*) [*ἀπό*, away; *χρώμα*, color]. Without color. **A.** Lens, a lens of a special variety of glass, corrected for spheric and chromatic aberration.

Apocodein (*ap-o-ko'-de-in*) [*ἀπό*, from; *κώδεια*, the head], $C_{18}H_{19}NO_2$. An alkaloid prepared from codein by dehydration. It is emetic and expectorant, with other qualities much like those of codein, and is recommended in chronic bronchitis. The hydrochlorate is generally used. Dose gr. iij-iv (0.2 to 0.25). Unof.

Apocynin (*ap-os'-in-in*) [*apocynon*, dogbane]. The precipitate from a tincture of *Apocynum cannabinum*; tonic, alterative, and cathartic. Dose gr. $\frac{1}{4}$ -j (0.016-0.065). Unof.

Apocynum (*ap-os'-in-um*) [*apocynon*, dogbane]. Canadian Hemp. The root of *A. cannabinum*, the properties of which are due to *apocynin*. It is a good expectorant; in full doses it is emetic and cathartic. Dose gr. v-xx (0.32-1.3). **A.**, Extractum fluidum. Dose \mathfrak{m} v-xx (0.32-1.3). **A.**, Tinct. Dose \mathfrak{m} v-xl (0.32-2.6). Unof. Another American species, **A. androsæmifolium**, has similar properties. Unof.

Apodia (*ah-po'-de-ah*) [*á* priv.; *πούς*, a foot]. Absence of feet.

Apolar (*ah-po'-lar*) [*á* priv.; *πόλος*, the end of an axis]. Not possessing a pole. **A.** Cells, nerve cells without processes.

Apollinaris Water (*ap-ol-in-a'-ris*). A German alkaline mineral water, highly charged

with carbonic acid, and largely used as a diluent in gout, rheumatism, etc.

Apomorphin (*ap-o-mor'-fin*) [*ἀπό*, from; *Morpheus*, the god of sleep] $C_{17}H_{17}NO_2$. An artificial alkaloid, derived from morphin by the abstraction of a molecule of water. The *hydrochlorate* is the salt used, and is a grayish crystalline powder. It acts as a centric emetic. Dose gr. $\frac{1}{16}$ – $\frac{1}{8}$ (0.003–0.0065), hypodermically, or gr. $\frac{1}{16}$ – $\frac{1}{8}$ (0.0065–0.001) by the mouth. It is expectorant in small doses. **A.**, *Injectio hypodermica* (B. P.). A two-per-cent. solution. **Apomorphinæ hydrochloratis**, Syrupus. Dose \mathfrak{z} ss–j (16.0–32.0).

Aponeurosis (*ap-on-u-ro'-sis*) [*ἀπό*, from; *νεῖρον*, a tendon]. A fibrous, membranous expansion of a tendon giving attachment to muscles or serving to enclose and bind down muscles. Important are the *infraspinous*, enclosing the *infraspinous* muscle; that of the *deltoid muscle*, of the *external oblique* or *abdominal*, of the *forearm*, of the *leg*, of the *transversalis*, of the *head*, etc., etc. **A.** of *Occipito-frontalis Muscle*, the aponeurosis that separates the two slips of the occipito-frontalis muscle. **A.** of *Soft Palate*, a thin but firm fibrous layer, attached above to the hard palate, and becoming thinner toward the free margin of the velum. **A.**, *Subscapular*, a thin membrane attached to the entire circumference of the subscapular fossa, and affording attachment by its inner surface to some of the fibers of the subscapularis muscle. **A.**, *Supraspinous*, a thick and dense membranous layer that completes the osseo-fibrous case in which the supraspinatus muscle is contained, affording attachment by its inner surface to some of the fibers of the muscle. **A.**, *Vertebral*, a thin aponeurotic lamina extending along the whole length of the back part of the thoracic region, serving to bind down the erector spinæ, and separating it from those muscles that connect the spine to the upper extremity.

Aponeurosis (*ap-on-u-ro-si'-tis*) [*ἀπονευρωσις*, an aponeurosis; *ιτις*, inflammation]. Inflammation of an aponeurosis.

Aponeurotic (*ap-on-u-ro'-ik*) [*ἀπονευρωσις*, aponeurosis]. Pertaining to an aponeurosis.

Apophysis (*ap-off'-is-is*) [*ἀπό*, from; *φύσις*, growth]. A process, outgrowth, or swelling of some part or organ, as of a bone. **A.** *lenticularis*, the orbicular process of the temporal bone. **A.** *raviana*, the process gracilis of the malleus.

Apoplectic (*ap-o-plek'-tik*) [*ἀποπληξία*, apoplexy]. Pertaining to or affected with apoplexy. **A.** *Equivalents*, a name given to the premonitory symptoms of apoplexy, indicating that the brain is subject to alterations in blood-pressure.

Apoplectiform (*ap-o-plek'-tif-orm*) [*ἀποπληξία*, apoplexy; *forma*, form]. Resembling apoplexy.

Apoplexy (*ap'-o-pleks-e*) [*ἀποπληξία*, a striking down]. The symptom-complex resulting from hemorrhage, or from the plugging of a vessel, in the brain or spinal cord. The term is sometimes also applied to the bursting of a vessel in the lungs, liver, etc. **A.**, *Capillary*, one resulting from rupture of capillaries. **A.**, *Ingravescent*, a term applied to a form of apoplexy in which there is a slowly progressive loss of consciousness due to a gradual leakage of blood from a ruptured vessel. **A.**, *Pulmonary*, escape of blood into the pulmonary parenchyma. **A.**, *Simple*, the name given to those cases of death from coma in which no cerebral lesion is found. **A.**, *Spinal*, rupture of a blood-vessel of the spinal cord.

Aporetin (*ap-o-re'-tin*) [*ἀπό*, from; *ρητινη*, a resin]. A resinous substance found in rhubarb.

Apositia (*ap-o-sit'-e-ah*) [*ἀπό*, from; *σῆτος*, food]. Aversion to or loathing of food.

Apostasis (*ap-os'-tas-is*) [*ἀπόστασις*, a standing away from]. 1. An abscess. 2. The end or the crisis of an attack of disease; termination by crisis. 3. An exfoliation.

Aposthia (*ah-pos'-the-ah*) [*ἀ* priv.; *πύσθη*, penis, prepuce]. Absence of the penis or of the prepuce.

Apostoli's Method. The use of strong electrolytic or chemic galvano-caustic currents in the treatment of diseases of the female generative organs, especially uterine fibroids.

Apothecaries' Weight. A system of weights and measures used in compounding medicines. The Troy pound of 5760 grains is the standard. It is subdivided into 12 ounces. The ounce is subdivided into eight drams, the dram into three scruples, and the scruple into 20 grains. For fluid measure the quart of 32 fluidounces is subdivided into two pints, the pint into 16 fluidounces, the ounce into eight fluidrams, and the fluidram into 60 minims. The following abbreviations are used:—

℥, <i>minim</i> .	℥, <i>uncia</i> , an ounce (480 grains).
ʒ, <i>scrupulus</i> , a scruple (20 grains).	℔, <i>libra</i> , a pound.
ʒ, <i>drachma</i> , a dram (60 grains).	℥, <i>octarius</i> , a pint.
	gr., <i>granum</i> , a grain.
	ss., <i>semissis</i> , one-half.

See *Weights and Measures*.

Apothem, or **Apothema** (*ap'-o-them* or *ap-oth'-em-ah*) [*ἀπό*, from; *θέμα*, a deposit]. A brown powder deposited from vegetable infusions or decoctions exposed to the air.

Apothesis (*ap-oth'-es-is*) [*ἀποthesis*, a putting back]. The reduction of a fracture or luxation.

Apotheter (*ap-ath'-et-er*). A navel-string repositr devised by Braun, consisting of a staff with a sling attached in which the prolapsed funis is placed and carried up into the uterine cavity.

Apozem, or Apozema (*ap'-o-zem* or *ap-oz'-em-ah*) [*ἀπό, away; ζέω, to boil*]. A decoction, especially one to which medicines are admixed.

Apparatus (*ap-ar'-tus*) [*apparatus, prepared*]. 1. A collection of instruments or devices used for a special purpose. 2. Anatomically the word is used to designate collectively the organs performing a certain function. **A.** ligamentosus colli, the *occipito-axoid ligament*, a broad band at the front surface of the spinal canal that covers the odontoid process.

Appendage (*ap-en'-daj*) [*appendere, to weigh; hang*]. Anything appended, usually something of minor importance. **A.**, Auricular, the projecting part of the cardiac auricle. **A.**, Epiploic. See *Appendix*. **A.** of the Eye, the eye lashes, eye-brows, lacrymal gland, lacrymal sac and ducts, and conjunctiva. **A.**, Ovarian, the parovarium. **A.** of the Skin, the nails, hairs, sebaceous and sweat glands. **A.**, Uterine, the uterine ligaments, the Fallopian tubes, and the ovaries.

Appendectomy (*ap-en-dis-ek'-to-me*) [*appendix; εκτομή, excision*]. Excision of the appendix.

Appendicitis (*ap-en-dis-it'-tis*) [*appendix; ιτις, inflammation*]. Inflammation of the appendix vermiformis. **A.** obliterans, an inflammation characterized by the progressive obliteration of the lumen of the appendix, by the disappearance of the epithelial lining and glandular structure. The symptoms are acute attacks of brief duration, moderate swelling at the seat of disease, and persistence of tenderness in the region of the appendix during the intermissions.

Appendicular (*ap-en-dik'-u-lar*) [*appendicula, a small appendix*]. Pertaining to the appendix vermiformis. **A.** Colic, a spasmodic colicky pain originating in the appendix.

Appendix (*ap-en'-diks*) [*appendere, to hang upon or to; pl., Appendices*]. An appendage. **A.** auricularis, the auricular appendage, a process of the auricles of the heart. **A.** cerebri, the pituitary body. **A.**, Ensiform. See *Xiphoid*. **A.** epiploicæ, fatty projections of the peritoneum of the large intestine. **A.** vermiformis, the small, blind gut projecting from the cecum. **A.**, Xiphoid. See *Xiphoid*.

Apperception (*ap-er-sip'-shun*) [*appercepere, to receive*]. The conscious reception or perception of a sensory impression; the

power of receiving and appreciating sensory impressions.

Appetence (*ap'-e-tens*) [*appetentia, appetite*]. An appetite or desire; the attraction of a living tissue for those materials that are appropriate for its nutrition.

Appetite (*ap'-e-tit*) [*appetere, to desire*]. The desire for food; also any natural desire; lust. **A.**, Perverted, that for unnatural and undigestible things, frequent in disease and in pregnancy.

Applanate (*ap'-lan-at*) [*ad, to; planus, flat*]. Horizontally flattened.

Applanatio, Applanation (*ap-lan-at'-she-o, ap-lan-at'-shun*) [*L.*]. A flattening. **A.** corneæ, flattening of the entire surface of the cornea from disease.

Apple (*ap'-l*) [*AS., æppel, an apple*]. The fruit of the tree, *Pyrus malus*. **A.**, Adam's. See *Pomum adam*. **A.**-brandy, an alcoholic spirit distilled from cider; cider-brandy. **A.** Extract. See *Extractum ferri pomatum*. **A.**-eye, a synonym of *Exophthalmos*. **A.**-oil, amyl valerianate.

Applicator (*ap'-lik-a-tor*) [*L.*]. An instrument used in making applications.

Apposition (*ap-o-zish'-un*) [*appondere, to apply to*]. The act of fitting together; the state of being fitted together.

Apraxia (*ah-praks'-e-ah*) [*ἀ priv.; πρᾶσσειν, to do*]. Soul-blindness; mind-blindness; object blindness; an affection in which the memory for the uses of things is lost, as well as the understanding for the signs by which the things are expressed.

Aproctia (*ah-prok'-te-ah*) [*ἀ priv.; πρωκτός, anus*]. Absence of the anus.

Apron (*ap'-prun*) [*ME., apren*]. 1. A cloth or rubber covering to prevent the clothing from becoming soiled. 2. The omentum. **A.**, Hottentot, artificially elongated labia minora. **A.** of Succor, a canvas appliance borne by two men, and used for the transportation of wounded persons.

Aprosexia (*ap-ro-seks'-e-ah*) [*ἀ priv.; προσέχειν, to give heed*]. A mental disturbance consisting in inability to fix the attention upon a subject. An inability to think clearly and to comprehend readily what is read or heard; a condition sometimes observed in the course of chronic catarrh of the nose or of the nose and pharynx.

Aprosopia (*ap-ro-so'-pe-ah*) [*ἀ priv.; πρόσωπον, the face*]. A form of fetal monstrosity with partial or complete absence of the face.

Apselaphesia (*ap-sel-af'-e-ah*) [*ἀ priv.; ψηλάφειν, touch*]. Loss of the tactile sense.

Apsithyria, or Apsithurea (*ap-sith'-v'-re-ah, or -u'-re-ah*) [*ἀ priv.; ψιθυρίζειν, to whisper*]. Hysterical aphonia, in which the patient not only loses the voice, but is also unable to whisper.

Apsychia (*ap-sik'-e-ah*) [*ā* priv.; *ψυχή*, spirit]. Unconsciousness; a faint or swoon.

Aptyalia, Aptyalism (*ap-ti-a'-le-ah*, *ap-ti'-al-ism*) [*ā* priv.; *πτύαλιζειν*, to spit]. Deficiency or absence of saliva.

Apyonin (*ah-pi'-on-in*) [*ā* priv.; *πύον*, pus]. A remedy introduced as a substitute for pyoktanin in ophthalmic practice. It is said to be identical with yellow pyoktanin. Unof.

Apyretic (*ah-pi-ret'-ik*) [*ā* priv.; *πυρετός*, fever]. Without fever.

Apyrexia (*ah-pi-reks'-e-ah*) [*ā* priv.; *πυρετός*, feverish]. Absence of fever.

Apyrexial (*ah-pi-reks'-e-al*) [*ā* priv.; *πυρετός*, feverish]. Pertaining to, of the nature of, or characterized by, apyrexia.

Aqua (*a'-kwah*) [L.]. Water. An oxid of hydrogen, having the composition H_2O . It is a solid below 32° , a liquid between 32° and 212° , vaporizes at 212° at the sea level (bar. 760 mm.), giving off vapor of tension equal to that of the air. It covers four-fifths of the surface of the earth, but is never pure in nature, containing from a trace of soluble matter, in rain water, to 26 per cent. of soluble mineral salts, in the Dead Sea. Water is an essential constituent of all animal and vegetable tissues. In the human body it forms two per cent. of the enamel of the teeth, 77 per cent. of the ligaments, 78 per cent. of the blood, and 93 per cent. of the urine. Water is a valuable antipyretic; internally, it is diuretic. It is the most useful of all the solvents. **A. bulliens**, boiling water. **A. communis**, common water. **A. destillata**, distilled water. **A. fervens**, hot water. **A. fluvialis**, river-water. **A. fontana**, well- or spring-water. **A. marina**, sea-water. **A. pluvialis**, rain-water. **Aqua**, in pharmacy, designates various medicated waters, as: **A. ammoniæ**. Dose \mathfrak{m} x-xxx (0.65-2.0). **A. amonizæ fortior**, used externally. **A. amygdalæ amaræ**. Dose $\mathfrak{f}\mathfrak{z}$ ij (8.0). **A. anethi**, dill-water (Br. Ph.). Dose $\mathfrak{f}\mathfrak{z}$ ss- $\mathfrak{f}\mathfrak{z}$ ij (16.0-64.0). **A. anisi**. Dose $\mathfrak{f}\mathfrak{z}$ ss- $\mathfrak{f}\mathfrak{z}$ ij (16.0-64.0). **A. aurantii florum**. Dose $\mathfrak{f}\mathfrak{z}$ ss- $\mathfrak{f}\mathfrak{z}$ ij (16.0-64.0). **A. aurantii florum fortior**, triple orange flower water. **A. camphoræ**. Dose $\mathfrak{f}\mathfrak{z}$ ss- $\mathfrak{f}\mathfrak{z}$ ij (16.0-32.0). **A. chlori**. Dose $\mathfrak{f}\mathfrak{z}$ j- $\mathfrak{f}\mathfrak{z}$ iv (4.0-16.0). **A. chloroformi**. Dose $\mathfrak{f}\mathfrak{z}$ ss- $\mathfrak{f}\mathfrak{z}$ ij (16.0-64.0). **A. cinnamomi**. Dose $\mathfrak{f}\mathfrak{z}$ ss- $\mathfrak{f}\mathfrak{z}$ ij (16.0-64.0). **A. creosoti**. Dose $\mathfrak{f}\mathfrak{z}$ j- \mathfrak{z} iv (4.0-16.0). **A. destillata**, distilled water. **A. fœniculi**. Dose $\mathfrak{f}\mathfrak{z}$ j- $\mathfrak{f}\mathfrak{z}$ ij (32.0-64.0). **A. fortis**. See *Acid, Nitric*. **A. hydrogenii dioxidi**, used chiefly locally. **A. lauro-cerasi**, cherry-laurel water (Br. Ph.). Dose \mathfrak{m} v-xxx (0.32-2.0). **A. menthæ piperitæ**. Dose $\mathfrak{f}\mathfrak{z}$ j- $\mathfrak{f}\mathfrak{z}$ ij (32.0-64.0). **A. menthæ viridis**. Dose $\mathfrak{f}\mathfrak{z}$ j- $\mathfrak{f}\mathfrak{z}$ ij (32.0-64.0). **A. pimentæ** (Br. Ph.). Dose $\mathfrak{f}\mathfrak{z}$ ss- $\mathfrak{f}\mathfrak{z}$ ij (16.0-64.0). **A. regia**. See *A. Nuro-*

hydrochloric. **A. rosæ**. Dose $\mathfrak{f}\mathfrak{z}$ j- $\mathfrak{f}\mathfrak{z}$ ij (32.0-64.0). **A. rosæ fortior**, used for making rose-water. **A. vitæ**, brandy or spirit. In anatomy, **A. labyrinthi**, the clear fluid existing in the labyrinth of the ear. **A. oculi**, aqueous humor.

Aquacapsulitis (*a-kwah-kap-su-li'-tis*) [*aqua*, water, *capsula*; *itis*, inflammation]. Inflammation of the membrane of Descemet; serous iritis.

Aquæductus (*ak-we-duk'-tus*). See *Aqueduct*.

Aquapuncture (*ak-wah-pungk'-chûr*) [*aqua*, water; *punctura*, a puncture]. Counter-irritation by means of a very fine jet of water impinging upon the skin; it is useful in neuralgic disorders.

Aquatic (*a-kwah'-ik*) [*aqua*, water]. Pertaining to water. **A. Cancer**. Synonym of *Cancrum oris*.

Aqueduct, or Aquæductus (*ak'-we-duk't*, or *ak-we-duk'-tus*) [*aqua*, water; *ductus*, a leading]. A canal for the passage of fluid; any canal. **A. cerebri**, the infundibulum. **A. cochleæ**, aqueduct of the cochlea. **A. of Cotunnus**, the aqueduct of the vestibule, extending from the utricle to the posterior wall of the pyramid in the brain. **A. Fallopii**, the canal in the petrous portion of the temporal bone through which the facial nerve passes. **A. Sylvii**, the aqueduct of Sylvius, the passageway from the third to the fourth ventricle, the *iter a tertio ad quartum ventriculûm*. **A. vestibuli**, the aqueduct of the vestibule of the ear.

Aqueous (*a'-kwe-us*) [*aqua*, water]. Watery. **A. Chamber of the Eye**, the space between the cornea and the lens; the iris divides it into an anterior and a posterior chamber. **A. Extract**, a solid preparation of a drug made by evaporation of its aqueous solution. **A. Humor**, the fluid filling the anterior chamber of the eye.

Aquocapsulitis (*a-kwo-kap-su-li'-tis*) [*aqua*, water; *capsula*, a small box]. Serous iritis.

Aquosity (*a-kwo'-it-e*) [*aquositas*, watery]. The state or condition of being watery.

Arabic (*ar'-ab-ik*) [*Ἀραβικός*, Arabic]. Of or pertaining to Arabia. **A. Acid**. See *Arabin*. **A. Gum**. See *Acacia*.

Arabin (*ar'-ab-in*) [*Ἀραβικός*, Arabic], ($C_6H_{10}O_5$)₂ + H_2O . Arabic Acid. A transparent, glassy, amorphous mass, an exudate from many plants. It is soluble in water, and is the principal constituent of gum arabic.

Arabinose (*ar'-ab-in-ô*) [*Ἀραβικός*, Arabic], $C_6H_{10}O_5$. One of the glucoses. made from gum arabic on boiling with dilute H_2SO_4 . It crystallizes in shining prisms that melt at 100° ; is slightly soluble in cold water, has a sweet taste, and reduces Fehling's solution, but is not fermented by yeast.

Arabite (*ar'-ab-it*) [*Ἀραβικός*, Arabic], $C_5H_{12}O_5$. A substance formed from arabinose by the action of sodium amalgam. It crystallizes from hot alcohol in shining needles, melting at 102° . It has a sweet taste, but does not reduce Fehling's solution.

Arachnitis (*ar-ak-ni'-tis*) [*ἀράχνη*, a spider's web; *itis*, inflammation]. Inflammation of the arachnoid membrane of the brain.

Arachnoid (*ar-ak'-noid*) [*ἀράχνη*, a spider's web; *oidos*, form]. Resembling a web. **A. Membrane**, the delicate membrane of the brain and cord between the dura and pia mater. It is separated from the latter by the subarachnoid space, and passes over the convolutions without dipping down into the fissures between them.

Arachnoidal (*ar-ak'-noid'-al*) [*ἀράχνη*, a spider's web; *oidos*, form]. Pertaining to the arachnoid.

Arack (*ar'-ak*) [*Ind.*]. A spirituous liquor distilled from rice, dates, palm-sap, etc.

Aræometer (*ar-e-om'-et-er*). See *Areometer*.

Aralia (*ar-ä'-le-ah*) [*L.*]. A genus of plants, order *Araliaceæ*, embracing several species, having aromatic, diaphoretic, and resolvent properties. Ginseng, wild sarsaparilla, petty morrel, and other plants esteemed in popular medicine belong here; few have active qualities of high value in any disease. Unof.

Aran-Duchenne's Disease. See *Diseases, Table of*.

Araneous (*ar-ä'-ne-us*) [*aranea*, a spider's web]. Full of webs; resembling a cobweb.

A. Membrane, the arachnoid.

Arantius, Bodies of. See *Ductus*. **A.**

Canal of. See *Canal*. **A.**, **Duct of**. See *Duct*. **A.**, **Nodules of**. See *Arantius, Bodies of*. **A.**, **Valves of**. See *Valvule Arantii*. **A.**, **Ventricle of**, a small cul-de-sac in the medulla oblongata, forming the lower termination of the fourth ventricle.

Araroba (*ar-ar-ä'-bah*) [*Brazil*]. *Goa Powder*. An oxidation product of the resin found deposited in the wood of the trunk of the *A. andira* of Brazil. Its active principle is *Chrysarobin* or *Chrysophanic acid*. It is largely used in skin affections.

Arbor (*ar'-bor*) [*L.*]. A tree. A name for the arbor vitæ of the cerebellum. **A. vitæ**, i. a term applied to the arborescent appearance of a section of the cerebellum, and also to a similar appearance of the folds of the interior of the cervix uteri. 2. The *Thuja occidentalis*.

Arborescent (*ar-bor-es'-ent*) [*arbor*, a tree]. Branching like a tree.

Arborization (*ar-bor-iz-a'-tion*) [*arbor*, a tree]. A form of nerve termination in which nerve-fiber is brought into contact with muscle fiber by means of an expansion.

Arbutin (*ar'-bu-tin*) [*arbutus*, the wild

strawberry tree], $(C_{12}H_{14}O_7)_2H_2O$. A bitter glucosid obtained from *Uva ursi*, or bearberry. It is neutral, crystalline, and resolvable into glucose and hydroquinon. It is diuretic. Dose gr. xv-xxx (1.0-2.0). See *Uva ursi*.

Arbutus (*ar-bu'-tus*) [*L.*]. A genus of ericaceous shrubs and trees. **A. menziesii**, the madroño of California, has an astringent bark, useful in diarrheas. Unof. **A. unedo**, the European arbutus; astringent and narcotic. **A.**, **Trailing**. See *Epigæa*.

Arc (*ark*) [*arcus*, a bow]. A part of the circumference of a circle; a more or less curved passage-way. **A.**, **Binauricular**, a measurement from the center of one auditory meatus to the other, directly upward across the top of the head. **A.**, **Bregmato-lambdoid**, a measurement along the sagittal suture. **A.**, **Naso-bregmatic**, a line measured from the root of the nose to the bregma. **A.**, **Naso-occipital**, measurement from the root of the nose to the lowest occipital protuberance. **A.**, **Reflex**, the pathway for a reflex act, comprising the center, the afferent and efferent nerve.

Arcanum (*ar-ka'-num*) [*arcantum*, a secret]. A secret medicine.

Arcate (*ar'-kāt*) [*arcatus*, bow-shaped]. Bow-shaped; curved; arcuate.

Arch (*arch*) [*arcus*, a bow]. A structure having a curved outline resembling that of an arc or a bow. **A.**, **Aortic**, five pairs of vascular arches existing in the fetus. **A.**, **Branchial**, the cartilaginous arches that support the gills of fishes. In the human fetus they are also present. **A.**, **Crural**, Poupart's ligament. **A.**, **Femoral**, same as *A.*, *Crural*. **A.**, **Palmar**, the arch formed by the radial artery and ulnar arteries in the palm of the hand; there are two, a superficial and a deep. **A.**, **Plantar**, the arch made by the external plantar artery. **A.**, **Postoral**, arches in the fetus, five in number, that develop into the lower jaw and throat. See *A.*, *Branchial*. **A.** of **Pubes**, that part of the pelvis formed by the convergence of the rami of the ischia and pubes on each side. **A.**, **Supraorbital**, the curved and prominent margin of the frontal bone that forms the upper boundary of the orbit. **A.** of a **Vertebra**, the part of a vertebra formed of two pedicles and two laminae and enclosing the spinal foramen. **A.**, **Zygomatic**, the arch formed by the malar and temporal bones.

Archameba (*ark-am-e'-bah*) [*ἀρχή*, primitive; *ἀμειβή*, change]. Haeckel's hypothetical progenitor of all amebæ and of all higher forms of life.

Archamphiaster, or **Archiamphiaster** (*ark-am-fe-as'-ter*, or *ar-ke-am-fe-as'-ter*) [*ἀρχή*,

first; *αμφί*, around; *ἀστήρ*, star]. In biology, those amphistars concerned in the production of the polar globules.

Archangelica (*ark-an-jel'-ik-ah*) [*ἀρχαγγελικός*, archangel]. A genus of umbelliferous plants. See *Angelica*.

Archebiosis (*ar-ke-bi-ō'-sis*) [*ἀρχή*, the beginning; *βίος*, life]. Spontaneous generation.

Archegogenesis (*ark-e-jen'-es-is*). The same as *Archebiosis*.

Archenteron (*ark-en'-ter-on*) [*ἀρχή*, a beginning; *έντερον*, intestine]. The embryonic alimentary cavity.

Archetype (*ar'-ke-tip*) [*ἀρχή*, a beginning; *τύπος*, a type]. In comparative anatomy, an ideal type or form with which the individuals or classes may be compared. A standard type.

Archiblast (*ar'-ke-blast*) [*ἀρχή*, first; *βλαστός*, germ]. In embryology, the granular areola surrounding the germinal vesicle. It is composed of three layers, the outer, or *epiblast*, from which the skin and its adnexa, the nervous system, and the terminal portions of the alimentary canal are formed; the middle, or *mesoblast*, from which the epithelium of the genito-urinary organs and the smooth and striated muscle-tissue are derived; and the inner, or *hypoblast*, for the development of the epithelium of the respiratory tract and of the alimentary canal. In pathology, the important tissues of the body as contrasted with the parblast, or connective tissues.

Archiblastic (*ar-ke-blas'-tik*) [*ἀρχή*, first; *βλαστός*, germ]. Derived from the archiblast. The parenchymatous tissues are regarded as archiblastic.

Archiblastoma (*ar-ke-blas-to'-mah*) [*ἀρχή*, first; *βλαστός*, germ; *δύμα*, a tumor]. A tumor composed of archiblastic tissue, such as myoma, neuroma, papilloma, adenoma, carcinoma, etc.

Archiblastula (*ar-ke-blas'-tu-lah*) [*ἀρχή*, first; *βλαστός*, a germ or bud]. In embryology, a ciliated, vesicular morula, resulting from complete and regular yelk-division and by invagination forming the archigastrula.

Archicytula (*ar-ke-sit'-u-lah*) [*ἀρχή*, a beginning; *κύτος*, a cell]. A fertilized egg-cell in which the nucleus is discernible.

Archigaster (*ar-ke-gas'-ter*) [*ἀρχή*, first; *γαστήρ*, belly]. The primitive, perfectly simple intestine; archenteron.

Archigastrula (*ar-ke-gas'-tru-lah*) [*ἀρχή*, beginning; *γαστήρ*, belly]. The gastrula as it is observed in the most primitive types of animal development; called also bell-gastrula from its shape.

Archigenesis (*ar-ke-jen'-es-is*). See *Archebiosis*.

Archil (*ar'-kil*) [ME., *orchell*]. A coloring

matter similar to litmus, chiefly obtained from the lichen, *Rocella tinctoria*; used for staining animal tissues.

Archimonerula (*ar-ke-mon-er'-u-lah*) [*ἀρχή*, first; *μονήρης*, single, solitary]. In embryology, a special name given by Haeckel to the monerula stage of an egg undergoing primitive and total cleavage.

Archimorula (*ar-ke-mor'-u-lah*) [*ἀρχή*, first; *μόρον*, a mulberry]. In embryology, the solid mass of cleavage cells, or mulberry mass, arising from the segmentation of an archicytula, and preceding the archiblastula and archigastrula.

Archineuron (*ar'-che-nu-ron*) [*ἀρχή*, beginning; *neuron*]. See *Neuron*, 2d definition.

Archinephron (*ar-ke-nef'-ron*) [*ἀρχή*, beginning; *νεφρός*, kidney]. The primitive or embryonic stage of the kidney or renal apparatus.

Archistome (*ar'-kis-tōm*) [*ἀρχή*, first; *στόμα*, mouth]. The gastrula mouth or opening of the archenteron; blastoporus.

Architis (*ar-ki'-tis*) [*ἀρχός*, anus]. Proctitis; inflammation of the anus.

Arciform (*ar'-se-form*) [*arcus*, bow; *arch*; *forma*, form]. Bow-shaped; especially used to designate certain sets of fibers in the medulla oblongata.

Arctation (*ark-ta'-shun*) [*arctatio*; *arctare*, to draw close together]. Contraction of an opening or canal.

Arctium (*ark'-she-um*) [L.]. Burdock. See *Lappa*.

Arcual (*ar'-ku-al*) [*arcualis*, arched]. Arched; bent or curved.

Arcuate (*ar'-ku-āt*) [*arcuatio*, a bowing]. Arched; curved; bow-shaped. **A. Fibers of the Cerebellum**, associating fibers connecting one lamina with another. **A. Fibers of the Cerebrum**, associating fibers connecting adjacent convolutions.

Arcuation (*ar-ku-a'-shun*) [*arcuatio*, a bowing]. Curvature, especially of a bone.

Arcus (*ar'-kus*) [*arcus*, a bow]. A bow or arch. **A. dentalis**, the dental arch. **A. senilis**, a ring of fatty degeneration seen about the periphery of the cornea in old persons. **A. senilis lentis**, an opaque ring in the equator of the crystalline lens; it sometimes occurs in the old. **A. zygomatiscus**, the zygomatic arch.

Ardor (*ar'-dor*) [*ardor*, heat]. Violent heat, burning. **A. urinæ**, burning pain in the inflamed urethra during micturition.

Area (*a'-re-ah*) [*area*, an open space]. A limited extent of surface. **A. Celsi**. See *Alopecia areata*. **A.**, Cohnheim's, certain clear-cut areas seen in a transverse section of muscle-fibers. **A.**, Crural, a space at the base of the brain included between the pons and chiasm. **A. germinativa**, or *embryonic*

spot, the oval germinating spot of the embryo.

A., Motor, the convolutions on either side of the rolandic fissure, containing the centers for voluntary motion. **A. opaca**, the opaque circle about the *A. pellucida*. **A. pellucida**, the light central portion of the *A. germinativa*. **A. vasculosa**, the vascular area of the *A. opaca*.

Areca Nut (*ar'-ek-ah*). See *Betel*.

Areane, Arekane (*ar'-ek-ān*) [Malay]. An oily and volatile basic substance obtainable from areca nut; said to be a purgative and sialagogue, and to slow the pulse. Unof.

Arecin (*ar'-es-in*) [Sp., *areca*], $C_{23}H_{26}N_2O$. 1. An organic base isomeric with brucin, derived from cinchona bark. 2. A brown-red coloring matter, obtained from areca nuts; areca red.

Areola (*ar'-el-o-lah*) [dim. of *area*, an open space; *pl.*, *Areolae*]. 1. The brownish space surrounding the nipple of the female breast. This is sometimes called *Areola papillaris*. A second areola, surrounding this, occurs during pregnancy. The pigmentation about the umbilicus is called the *umbilical areola*. 2. Any interstice or minute space in a tissue.

Areolar (*ar'-el-o-lar*) [*areola*, dim. of *area*, an open space]. Relating to or characterized by areolae. **A. Tissue**, cellular tissue; loose connective tissue.

Argentamin (*ar-jen-tam'-in*) [*argentum*, silver; *amin*]. A solution of silver phosphate in ethylenediamine. It is an antiseptic agent that does not coagulate albumin.

Argentation (*ar-jen-ta'-shun*) [*argentum*, silver]. Staining with a preparation of silver.

Argentite (*ar-jen'-tik*) [*argentum*, silver]. Containing silver.

Argentine (*ar'-jen-tin*) [*argentum*, silver]. Containing or resembling silver.

Argentum (*ar-jen'-tum*) [L.]. Silver. $Ag=108$; quantivalence, 1; specific gravity, 10.4 to 10.5. A malleable and ductile metal of brilliant white luster. It tarnishes only in the presence of free sulphur, sulphur gases, and phosphorus. The following salts are used in medicine: **A. cyanidum**, $AgCN$, used in the preparation of hydrocyanic acid. **A. iodidum**, AgI , sometimes used internally instead of *A. nitras*. Dose gr. $\frac{1}{4}$ - $\frac{1}{2}$ (0.016-0.065). **A. nitras**, $AgNO_3$, nitrate of silver, argentic nitrate, "lunar caustic," a powerful astringent and an escharotic of moderate strength. It stains skin and other tissues black. If too long administered, it leaves a slate-colored, insoluble deposit of silver under the skin (*Argyria*). It is used in gastric catarrh, in gastric ulcer, in intestinal ulceration, and as an alternative in scleroses of the nervous system. Dose gr. $\frac{1}{6}$ - $\frac{1}{2}$ (0.01-0.032). **A. nitras dilutum**, the mitigated caustic, or diluted stick, is fused with an

equal amount of potassium nitrate. **A. nitras fusum**, "stick caustic," contains four per cent. of silver chlorid. It is used locally. **A. oxidum**, Ag_2O , explosive when treated with ammonia. Dose gr. $\frac{1}{2}$ - $\frac{1}{2}$ (0.032-0.13). **A. vivum**, an old name for mercury or quicksilver.

Argilla (*ar'-jil'-ah*) [*ἀργίλλος*, potter's clay]. White or potter's clay; alumina.

Argol (*ar'-gol*) [*ἀργός*, white]. The impure tartar derived from wine.

Argon (*ar'-gon*) [*ἀργός*, inactive]. A new element discovered in the atmosphere in 1894 by Lord Rayleigh and Prof. Ramsey. Its symbol is A; the atomic weight has not been definitely determined, but is thought to be 19.7. It is soluble to the extent of 4 volumes in 100 of water, and condenses to a colorless liquid at -128.6° C. and under a pressure of 33 atmospheres, the liquid having a density of 1.5. Argon may be obtained by freeing air which has been deprived of its carbon dioxide and water, from oxygen by means of red-hot copper, and then absorbing the nitrogen by means of metallic magnesium. The residual gas, the passage of the gases being repeated a number of times, is argon. Chemically, it is the most inert element yet known.

Argyll Robertson Pupil. See *Signs and Symptoms*, *Table of*, and *Pupil*.

Argyria (*ar'-ji'-re-ah*) [*argentum*, silver]. A form of discoloration of the skin and mucous membranes produced by the prolonged administration of silver, the granules of silver being deposited in about the same position as those of the natural pigment of the skin.

Argyrosis (*ar'-ji-ro'-sis*). Same as *Argyria*, *q. v.*

Arhinia (*ah-rin'-e-ah*) [*ἀ priv.*; *ῥίς*, nose]. Congenital absence of the nose.

Arrhythmia (*ar'-ith'-me-ah*). See *Arrhythmia*.

Arica Bark. Calisaya bark exported from Arica, Peru. It contains the alkaloid, aricin.

Aricin (*ar'-is-in*) [*Arica*, in Peru], $C_{23}H_{26}N_2O_4$. An alkaloid obtained from several varieties of cinchona bark.

Aristol (*ar'-is-tol*) [*ἀριστος*, best], $(C_6H_5-CH_2O)_2$. Dithymol-iodid. It is also called annidalin, although this is dithymol-triiodid. An iodine compound used as a substitute for iodoform as an antiseptic dressing. It has the advantage of being odorless, and is used either in the powder form or as a five to ten per cent. ointment with vaselin or lanolin.

Aristolochia (*ar'-is-to-lo'-ke-ah*) [*ἀριστος*, best; *λοχία*, the lochia]. A genus of exogenous herbs, many species of which have

active medicinal qualities. **A. clematitis**, of Europe, has been used as a tonic, stimulant, and diaphoretic. **A. cymbifera**, of S. America, furnishes a part of the drug called *Guaco*, and is a good tonic and stimulant. **A. serpentaria** (see *Serpentaria*) is at present more used in medicine than any other species.

Aristolochin (*ar-is-to-lo'-kin*) [*ἀριστος*, best; *λοχία*, the lochia]. A bitter principle found in Virginia snake-root. See *Serpentaria*.

Aristotle's Experiment. The double feeling experienced by the fingers when a single pebble is placed between the crossed fingers of one hand.

Arithmomania (*ar-ith-mo-ma'-ne-ah*) [*ἀριθμός*, a number; *μανία*, madness]. An insane anxiety with regard to the number of things that fall under the observation. Sometimes it consists in constant or uncalled-for counting, either of objects or in mere repeating of consecutive numbers.

Arit's Ointment. An ointment containing grs. viiss of belladonna to 3j¼ of blue ointment.

Arm (*arm*) [*armare*, to arm]. 1. That part of the upper extremity from the shoulder to the wrist. 2. That portion of the stand connecting the body or tube of a microscope to the pillar. **A. Center**, the cortical center for the movement of the arm; it is situated in the middle third of the ascending frontal and ascending parietal convolutions.

Armature (*ar'-mat-châr*) [*armatura*, equipment]. A mass of soft iron at the extremity of a magnet. Also, the core of iron around which coils of insulated wire are wound.

Army Itch (*ar'-me itch*). A distressing, chronic form of itch prevalent in the United States at the close of the civil war. The itching was intense. The eruption was seen especially on the arms, forearms, chest, abdomen, and lower extremities, particularly on the ulnar side of the forearm and inner aspect of the thigh. It resembled prurigo associated with vesicles, pustules, and eczema.

Arnica (*ar'-nik-ah*) [L.]. A genus of composite-flowered plants. The name in medicine denotes the plant commonly known as "Leopard's bane," *A. montana*. Both flowers and roots are used in medicine. Its properties are probably due to an alkaloid, *trimethylamin*, C_3H_7N . In small doses it is a cardiac stimulant; in larger doses a depressant. It is a popular remedy, when locally applied, for sprains, bruises, and surface wounds. **A., Emplastrum**, contains extract of root 33, lead plaster 67 parts. **A. Radicis**, Ext. Dose gr. j-ij (0.065-0.2). **A. Rad.**, Ext. **Fld.** Dose ℥v-xx (0.32-1.3). **A., Infus-** flowers, 100 parts water. It

is superior to the tincture for local use. **A., Tinct.**, 20 per cent. Dose ℥v-xxx (0.32-2.0). **A. Rad.**, Tinct., 10 per cent. Dose ℥v-xxx (0.32-2.0). **Arnicae flores**, arnica flowers. **Arnicae radix**, the root of arnica. **Arnicin** (*ar'-nis-in*) [*arnica*], $C_{20}H_{30}O_6$. A brownish, bitter glucosid extracted from the flowers of *Arnica montana*.

Arnold's Nerve. The auricular branch of the pneumogastric nerve.

Aroma (*ar-o'-mah*) [*ἀρώμα*, spice]. The fragrance or odor emanating from certain vegetable substances, especially those used for food and drink.

Aromatic (*ar-o-mat'-ik*) [*ἀρώμα*, spice]. 1. Having a spicy odor. 2. A substance characterized by a fragrant, spicy taste and odor, as cinnamon, ginger, the essential oils, etc. 3. A qualification applied to any carbon compound originating from benzin, C_6H_6 . Their stability is relatively great as compared with that of the fatty bodies. **A. Acids**, those derived from the benzin group of hydrocarbons. **A. Compound**, any benzyl derivative. **A. Fluid Extract**, aromatic powder 100, alcohol sufficient to make 100 c.c. **A. Group**, a series of hydrocarbons having the composition $C_{10}H_{16}$. **A. Powder**. See *Cinnamomum*. **A. Sulphuric Acid**. See *Acid, Sulphuric*. **A. Vinegar**, any mixture of aromatic oils in vinegar. **A. Wine**, a wine containing in each 100 parts, one part each of lavender, origanum, peppermint, rosemary, sage, and wormwood.

Aromin (*ar-o'-min*) [*ἀρώμα*, spice]. An alkaloid obtainable from urine. When heated it emits a fragrant odor.

Arrack (*ar'-ak*) [Ind.]. A liquor distilled from malted rice. Any alcoholic liquor is called arrack in the East.

Arrector (*ar-ek'-tor*) [L., an erector]. An erector. **A. pili**, a fan-like arrangement of a layer of smooth muscular fibers surrounding the hair follicle, the contraction of which erects the follicle and produces *cutis anserina*, or "goose-skin."

Arrhinia (*ah-rin'-e-ah*). Same as *Arhinia*.

Arrhythmia (*ah-rith'-me-ah*) [*ἄ priv.*; *ῥυθμός*, rhythm]. Absence of rhythm.

Arrhythmic (*ah-rith'-mik*) [*ἄ priv.*; *ῥυθμός*, rhythm]. Without rhythm; irregular.

Arrow-poison (*ar'-o-poi'-zun*). See *Curare*.

Arrowroot (*ar'-o-root*) [ME., *arow*; *route*].

A variety of starch derived from *Muranta arundinacea* of the West Indies, Southern States, etc. It is a popular remedy for diarrhea, and is widely used as a food. Many other starchy preparations are sold as arrowroot.

Arsenate, or **Arseniate** (*ar'-sen-at*, or *ar-se'-ne-at*) [*arsenum*]. Any salt of arsenic acid.

Arsenum (*ar-se'-num*). As = 75; quantivalence III, V. A brittle, crystalline metal, of a steel-gray color, tarnishing on exposure to the air. Sp. gr. 5.73. It sublimes at 180° C., and gives off a garlicky odor. It forms two oxids: arsenous oxid, As_2O_3 , and arsenic oxid, As_2O_5 . The sulphids are used as pigments in the arts, rarely in medicine. Scheele's green is copper arsenite; Schweinfurth and Paris-green likewise contain arsenic. Arsenic is also an ingredient of certain poisons used against rodents and insects. In medicine, arsenic is used as an alterative in anemias, chronic malaria, asthma, phthisis pulmonalis, as a gastric sedative, and in chorea.

Arsenous Acid, white arsenic, "ratsbane," As_2O_3 . Dose gr. $\frac{1}{2}$ – $\frac{1}{4}$ (0.002–0.006). **A. Acid**, **Liquor** *Liq. arsenici hydrochlorici*, B. P., a 1 per cent. solution of the oxid in hydrochloric acid and distilled water. Dose $\mathfrak{m}\mathfrak{j}$ –v (0.13–0.32). **Ferri Arsenias**, $Fe_3As_2O_8$. Dose gr. $\frac{1}{2}$ – $\frac{1}{4}$ (0.004–0.03). **Liq. arsenici hydrochlor.** (B. P.). Dose $\mathfrak{m}\mathfrak{j}$ –v (0.13–0.32). **Potassii arsenitis**, **Liq.** (*Liquor Arsenitalis*, B. P.). *Fowler's solution*. It contains A. acid 1, potass. bicarbonate 1, comp. tinct. lavender 3, and distilled water q. s. to make 100 parts. Dose $\mathfrak{m}\mathfrak{j}$ –x (0.13–0.65). **Sodii arsenias**, $Na_2HASO_4 \cdot 7H_2O$. Dose of the dried salt gr. $\frac{1}{2}$ – $\frac{1}{4}$ (0.0025–0.005). **Sodii arseniatis**, **Liq.**, Pearson's solution. Dose $\mathfrak{m}\mathfrak{j}$ –x (0.13–0.65). **A. iodidum**, AsI_3 . Dose gr. $\frac{1}{2}$ – $\frac{1}{4}$ (0.003–0.008). **A. et hydrargyri iod.**, **Liq.**, liquor of the iodid of arsenic and mercury. Donovan's solution. It contains A. iodid 1, mercuric iodid 1, distilled water 100 parts. Dose $\mathfrak{m}\mathfrak{j}$ –x (0.13–0.65). **A. bromidum**, $AsBr_3$, is used in diabetes. Dose gr. $\frac{1}{2}$ – $\frac{1}{4}$ (0.003–0.008). **A. bromidi**, **Liq.**, Clemens' solution, a one per cent. solution of the bromid of arsenic. Dose $\mathfrak{m}\mathfrak{j}$ –ix (0.265–0.26).

Arsenical (*ar-sen'-i-kal*) [*arsenium*, arsenic]. Pertaining to arsenic.

Arsenicalism, **Arsenism** (*ar-sen'-i-kal-izm*, *ar-sen'-izm*) [*arsenium*, arsenic]. Chronic arsenical poisoning.

Arsenite (*ar-sen'-it*) [*arsenium*, arsenic]. Any salt of arsenous acid.

Arsenous (*ar-sen'-us*) [*arsenium*, arsenic]. Pertaining to arsenic. **A. Acid**. See *Arsen.*

Arsins (*ar-sin')* [*arsenium*, arsenic]. Permanent volatile arsenical bases found by Selmi to be produced by the contact of arsenous acid and albuminous substances.

Arsonium (*ar-sen'-i-um*) [*arsenium*, arsenic; *ammonium*]. AsH_4^+ . A univalent radical in which arsenic replaces the nitrogen of ammonium.

Artefact (*ar'-te-fak'-tuls*, by art; *artum*, man). In microscopy and histology, a structure that has been produced by mechanic, chemical, or other artificial means; a struc-

ture or tissue that has been changed from its natural state.

Artemisia (*ar-tem-iz'-e-ak*) [*Ἀρτεμις*, the goddess Diana]. A genus of plants of the order *Compositae*. **A. abrotanum**, *Southern wood*, is stimulant, tonic, and vermifuge, and is popularly used as a vulnerary. It is similar in properties to wormwood. Dose of fld. ext. $\mathfrak{m}\mathfrak{x}$ –xx (0.65–1.3). Unof. **A. absinthium**. See *Absinthe*. **A. maritima** affords pure wormseed. **A. vulgaris**, mugwort, a popular remedy in various diseases.

Arteria (*ar-te'-re-ak*) [*ἀρτηρία*, the trachea]. A hollow tube. See *Artery*.

Arterial (*ar-te'-re-al*) [*ἀρτηρία*, the trachea]. Pertaining to an artery.

Arterialization (*ar-te'-re-al-iz-a'-shun*) [*ἀρτηρία*, the trachea]. The process of making or becoming arterial; the change from venous blood into arterial.

Arterin (*ar'-ter-in*) [*ἀρτηρία*, trachea]. Hoppe-Seyler's term for the arterial blood-pigment as contained in the red corpuscles.

Arteriocapillary (*ar-te'-re-o-kap'-il-a-re*) [*ἀρτηρία*, trachea; *capillus*, a hair]. Pertaining to arteries and capillaries. **A. Fibrosis**, a chronic inflammatory process characterized by an overgrowth of connective tissue in the walls of the blood-vessels. It is also known as arteriocapillary fibrosis of Gull and Sutton.

Arteriofibrosis (*ar-te-re-o-fi-bro'-sis*). See *Arteriocapillary Fibrosis*.

Arteriogram (*ar-te'-re-o-gram*). See *Sphygmogram*.

Arteriography (*ar-te-re-og'-ra-fe*) [*ἀρτηρία*, trachea; *γραφία*, a writing]. 1. A description of the arteries. 2. The graphic representation of the pulse waves.

Arteriole (*ar-te'-re-ol*) [*arteriola*]. A very small artery. **A.**, Straight, the small blood-vessels that supply the medullary pyramids of the kidneys.

Arteriology (*ar-te-re-ol'-a-je*) [*ἀρτηρία*, trachea; *λογία*, science]. The science of the arteries; the anatomy, physiology, and pathology of the arteries.

Arteriomalacia (*ar-te-re-o-mal-a'-se-ak*) [*ἀρτηρία*, trachea; *μαλακία*, softness]. Softening of the wall of an artery.

Arteriosclerosis (*ar-te-re-o-skle-ro'-sis*) [*ἀρτηρία*, trachea; *σκληρός*, hard]. A chronic inflammation of the arterial walls, especially of the intima.

Arteriosclerotic (*ar-te-re-o-skle-ro'-ik*) [*ἀρτηρία*, trachea; *σκληρός*, hard]. Pertaining to arteriosclerosis. **A. Kidney**, a kidney the seat of chronic interstitial inflammation affecting primarily the blood-vessels.

Arteriosis (*ar-te-re-o'-it-e*) [*ἀρτηρία*, trachea]. The quality of being arterial.

Arteriotomy (*ar-te-re-ol'-o-me*) [*ἀρτηρία*,

trachea; *téuveiv*, to cut]. The cutting or opening of an artery for the purpose of letting blood. The temporal artery is generally selected.

Arteriovenous (*ar-te-re-o-ve'-nus*) [*âprepia*, trachea; *vena*, vein]. Both arterial and venous; involving an artery and a vein, as an arteriovenous aneurysm.

Arterioversion (*ar-te-re-o-ver'-shun*) [*âprepia*, trachea; *verteo*, to turn]. Weber's method of arresting hemorrhage by turning vessels inside out by means of an instrument called the arterioverter.

Arterioverter. See *Arterioversion*.

Arteritis (*ar-te-ri'-tis*) [*âprepia*, trachea; *itis*, inflammation]. Inflammation of an artery. **A. deformans.** See *Endarteritis, Chronic*. **A. Obliterans.** See *Endarteritis obliterans*.

Artery (*ar'-ter-e*) [*arteria*; *âprepia*, trachea]. One of the tube-like vessels through which the blood is propelled by the heart to all parts of the body. Arteries end in arterioles and capillaries. They are composed of three coats: the outer, or *tunica adventitia*; the middle, or *tunica media*, the muscular coat; the internal, or *intima*, composed of endothelial cells, fibrous and elastic tissue. **A. Compressor, A. Constrictor**, an instrument for occluding an artery for the purpose of arresting or preventing hemorrhage. **A., End, or A., Terminal**, an artery that does not anastomose with other arteries by means of large branches; there is usually a capillary anastomosis. **A. Forceps**, a forceps for catching or twisting an artery. A table of the arteries is appended.

TABLE OF ARTERIES.

NAME.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Acromio-thoracic (thoracic axis).	Second branch of first part of axillary.	Shoulder, arm, upper anterior part of chest, and mammary gland.	Acromial, humeral, pectoral, clavicular.
Alar thoracic.	Second part of axillary.	Lymphatic glands in axilla.	
Anastomotic (of external plantar).	External plantar.	Outer border of foot.	Anastomoses with the tarsal and metatarsal branches of the dorsalis pedis.
Anastomotic (of internal plantar).	Internal plantar.	Inner side of foot.	Anastomoses with internal tarsal branch of the dorsalis pedis.
Anastomotica magna (of brachial).	Brachial.	Elbow.	Posterior and anterior.
Anastomotica magna (of superficial femoral).	Superficial femoral (in Hunter's canal).	Knee.	Superficial and deep.
Angular.	The termination of the facial.	Lacrimal sac and lower part of orbicularis palpebrarum.	Anastomoses with infraorbital.
Aorta, abdominal.	Thoracic aorta.	Two common iliacs.	Phrenic (right and left), celiac axis, suprarenal or capsular (right and left), superior mesenteric, lumbar (four pairs), renal (right and left), spermatic (right and left), inferior mesenteric, right and left common iliac, middle sacral.
Aorta, arch.	Left ventricle of heart.	Thoracic aorta.	Two coronary, innominate, left common carotid, left subclavian.
Aorta, primitive.	That portion from the origin to the point at which the first branch is given off.		
Aorta, thoracic.	Arch of aorta.	Abdominal aorta.	Two or three pericardiac, three bronchial, four or five esophageal, twenty intercostal, subcostal, for twelve dorsal, diaphragmatic, aberrant.

TABLE OF ARTERIES—*Continued.*

NAME.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Articular, middle (of knee).	Popliteal.	Crucial ligaments and joint.	
Articular, superior external (of knee).	Popliteal.	Crureus and knee.	
Articular, superior internal.	Popliteal.	Knee.	
Auricular, posterior.	Fifth branch of external carotid.	Back of auricle, scalp, and part of neck.	Parotid, muscular, stylo-mastoid, anterior terminal or auricular, and posterior terminal or mastoid.
Axillary.	Subclavian.	Brachial and seven branches.	Superior thoracic, acromio-thoracic, long thoracic, alar thoracic, subscapular, anterior and posterior circumflex.
Basilar.	By confluence of right and left vertebral.	Brain.	Transverse (or pontile), internal auditory, anterior cerebellar, superior cerebellar, two posterior cerebral.
Brachial.	Axillary.	Arm and forearm.	Superior and inferior profunda, anastomotica magna, nutrient, muscular, radial and ulnar.
Cardiac.	Gastric.	Cardiac end of stomach.	
Carotid, common.	<i>Right side</i> , innominate; <i>left side</i> , arch of aorta.	External and internal carotid.	External and internal carotid.
Carotid, external.	Common carotid.	Anterior part of neck, face, side of head, integuments and dura mater.	Ascending pharyngeal, superior thyroid, lingual, facial, occipital, posterior auricular, temporal, internal maxillary.
Carotid, internal.	Common carotid.	Greater part of brain, the orbit, internal ear, forehead, and nose.	Tympanic, vidian, arteria receptaculi, pituitary, gasserian, meningeal, ophthalmic, posterior communicating, anterior choroid, anterior cerebral, middle cerebral.
Carotid, primitive.	See <i>Carotid, common</i> .		
Celiac.	Abdominal aorta.	Stomach, duodenum, spleen, pancreas, liver, and gall-bladder.	Gastric, hepatic, splenic.
Central (of retina).	Ophthalmic.	Retina.	
Cerebellar, anterior.	Basilar.	Anterior inferior surface of cerebellum.	
Cerebellar, inferior.	Vertebral.	Vermiform process, and cortex of cerebellum.	Inferior vermiform and the hemispherical.
Cerebellar, superior.	Basilar.	Superior vermiform process and circumference of cerebellum.	Superior vermiform and hemispherical.
Cerebral, anterior.	Internal carotid.	Anterior portion of cerebrum.	Anterior communicating, ganglionic (or central), commissural, hemispherical (or cortical).
Cerebral, middle.	Internal carotid.	Middle portion of cerebrum.	Ganglionic (or central), hemispherical (or cortical).
Cerebral, posterior.	Basilar.	Temporo-sphenoidal and occipital lobes.	Ganglionic (or central), and hemispherical (or cortical).
Cervical.	Uterine.	Cervix uteri.	

TABLE OF ARTERIES—*Continued.*

NAME.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Cervical, ascending.	Inferior thyroid.	Deep muscles of neck and spinal canal.	Muscular, spinal, and phrenic.
Cervical, deep.	Superior intercostal.	Deep muscles of neck and spinal canal.	Muscular, anastomotic, vertebral (or spinal).
Cervical, superficial.	Transverse cervical.	Trapezius, levator anguli scapulae, splenius muscles, and posterior chain of lymphatic glands.	
Cervical, transverse (transversalis colli).	Thyroid axis.	Posterior cervical and scapular regions.	Posterior scapular and superficial cervical.
Circumflex, anterior (of axillary).	Axillary.	Pectoralis major, biceps, and shoulder-joint.	Bicipital and pectoral.
Circumflex, posterior (of axillary).	Axillary.	Deltoid, teres minor, triceps, and shoulder-joint.	Nutrient, articular, acromial, muscular.
Colic, left.	Inferior mesenteric.	Descending colon.	
Colic, middle.	Superior mesenteric.	Transverse colon.	
Colic, right.	Superior mesenteric.	Ascending colon.	
Colic, transverse.	Colic, middle.	Transverse colon.	
Comes nervi phrenici.	See <i>Phrenic</i> .	<i>superior.</i>	
Communicating.	Dorsalis pedis.	Enters into formation of plantar arch.	Two digital.
Communicating (or perforating).	Deep palmar arch.	Join proximal ends of metacarpal and second and third dorsal interosseous arteries.	
Communicating, anterior.	Anterior cerebral.	Assists in formation of anterior boundary of circle of Willis; sends branches to caudate nucleus.	
Communicating, posterior.	Posterior cerebral.	Enters into formation of circle of Willis; uncinat convolution and optic thalamus.	Uncinate, middle thalamic.
Coronary, inferior.	Facial.	Lower lip.	
Coronary, left.	Left anterior sinus of Valsalva.	Heart.	Left auricular, anterior interventricular, left marginal, terminal.
Coronary, right.	Right anterior sinus of Valsalva.	Heart.	Right auricular, preventricular, right marginal, posterior interventricular, transverse.
Coronary, superior.	Facial.	Upper lip.	
Diaphragmatic.	Thoracic aorta.	Diaphragm.	
Digital.	External plantar.	Outer side of the 2d and 3d, 4th, and 5th toes.	
Digital, palmar.	Superficial palmar arch.	Both sides of little, ring, and middle finger, and ulnar side of index finger.	
Dorsal (of penis).	Pudic.	Penis.	
Dorsalis hallucis.	Continuation of dorsalis pedis.	Great and second toe.	

TABLE OF ARTERIES—*Continued.*

NAME.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Dorsalis pedis.	Continuation of anterior tibial.	Assists to form plantar arch.	Tarsal, metatarsal, dorsalis hallucis, communicating.
Epigastric, deep (or inferior).	External iliac.	Abdominal wall.	Cremasteric, pubic, muscular, cutaneous, terminal.
Epigastric, superficial.	Common femoral.	Inguinal glands, skin, superficial fascia, and abdominal wall.	
Epigastric, superior.	Internal mammary.	Abdominal wall and diaphragm, liver and peritoneum.	Phrenic, xiphoid, cutaneous, muscular, hepatic, and peritoneal.
Epiploic.	Right and left gastro-epiploic.	Omentum.	
Esophageal.	Gastric.	Esophagus.	
Esophageal.	Inferior thyroid.	Esophagus.	
Esophageal.	Left phrenic.	Esophagus.	
Esophageal (4 or 5).	Thoracic aorta.	Esophagus.	
Esophageal, inferior.	Coronary (of stomach).	Esophagus.	
Facial.	Third branch external carotid.	Pharynx and face.	Ascending, or inferior palatine, tonsillar, glandular, muscular, submental, masseteric, buccal, inferior labial, inferior and superior coronary, lateralis nasi, angular.
Femoral, common.	Continuation of external iliac.	Lower part of abdominal wall, upper part of thigh and genitalia.	Superficial epigastric, superficial circumflex iliac, superficial external pudic, deep external pudic, profunda.
Femoral, deep.	See <i>Femoral, profunda</i> .		
Femoral, profunda.	Common femoral.	Muscles of thigh.	External circumflex, internal circumflex, and three perforating.
Femoral, superficial.	Continuation of common femoral.	Muscles of thigh and knee-joint.	Muscular, saphenous, anastomotica magna.
Gastric (or coronary).	Celiac axis.	Stomach, liver, and esophagus.	Esophageal, cardiac, gastric, and hepatic.
Gastro-duodenal.	Hepatic.	Stomach and duodenum.	Right gastro-epiploic and superior pancreatico-duodenal.
Gastro-hepatic.	See <i>Coronary, of stomach</i> .		
Helicine.		The arteries found in cavernous tissue, as in the testicle, uterus, ovary, etc.	
Hemorrhoidal, inferior (or external).	Pudic.	Sphincter muscle, levator ani.	
Hemorrhoidal, middle.	Internal iliac, anterior division.	Middle part of rectum.	
Hemorrhoidal, superior.	Inferior mesenteric.	Upper part of rectum.	
Hepatic.	Celiac axis.	Liver, pancreas, part of duodenum, and stomach.	Pancreatic, subpyloric, gastroduodenal, right and left terminal.
Iliac, common.	Terminal branch of abdominal aorta.	Peritoneum, subperitoneal fat, ureter, and terminates in external and internal iliac.	Peritoneal, subperitoneal, ureteric, external and internal iliac.

TABLE OF ARTERIES—*Continued.*

NAME.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Iliac, external.	Common iliac.	Lower limb.	Deep epigastric, deep circumflex iliac, muscular, and continues as femoral.
Iliac, internal.	Common iliac.	Pelvic and generative organs, and inner side of thigh.	Anterior and posterior trunk.
Iliac, internal, anterior trunk.	Internal iliac.	Pelvic and generative organs and thigh.	Hypogastric, superior, middle, and inferior vesical, middle hemorrhoidal, uterine, vaginal, obturator, sciatic, internal pudic.
Iliac, internal, posterior trunk.	Internal iliac.	Muscles of hip and sacrum.	Ilio-lumbar, lateral sacral, and gluteal.
Innominate.	Arch of aorta.	Right side of head and right arm.	Right common carotid, right subclavian, occasionally thyroidea ima.
Intercostal, anterior.	Internal mammary.	Intercostal muscles, ribs (upper 5 or 6), and pectoralis major.	
Intercostal, anterior.	Musculo-phrenic.	Lower 5 or 6 intercostal spaces.	
Intercostal, superior.	Subclavian.	Neck and upper part of thorax.	Deep cervical, first intercostal, arteria aberrans.
Interosseous, anterior.	Interosseous (common).	Muscles of forearm.	
Interosseous, common.	Ulnar.	Interosseous membrane and deep muscles of the forearm.	Anterior and posterior interosseous.
Interosseous, posterior.	Ulnar.	Muscles of forearm.	
Labial, superior.	See <i>Coronary, superior.</i>		
Laryngeal, superior.	Superior thyroid.	Intrinsic muscles and mucous membrane of larynx.	
Lenticulo-striate.	Middle cerebral.	Lenticular and caudate nuclei.	
Lingual.	External carotid.	Tongue.	Hyoid, dorsalis linguae, sublingual, ranine.
Mammary, external.	See <i>Thoracic, long.</i>		
Mammary, internal.	Subclavian.	Structures of thorax.	Superior phrenic, mediastinal (a. thymic), pericardiac, omental, anterior intercostal, perforating, lateral intercostal, superior epigastric, internal mammary.
Maxillary, external.	See <i>Facial.</i>		
Maxillary, internal (maxillary group).	External carotid.	Structures indicated by names of branches.	Deep auricular, tympanic, middle meningeal, mandibular, small meningeal.
Maxillary, internal (pterygoid group).	External carotid.	Structures indicated by names of branches.	Masseter, pterygia deep, temporal, internal and external pterygoid, buccal, mental deep, temporal.
Maxillary, internal (sphenomaxillary group).	External carotid.	Structures indicated by names of branches.	Pterygia anterior in alveolar, inferior orbital pterygia in cavernosus, palatine, sphenopterygoid, sphenomaxillary, in sphenoidal.
Median (arteria comes nervi mediani).	Anterior interosseous.	Median nerve and superficial palmar arch.	

TABLE OF ARTERIES—*Continued.*

NAME.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Mediastinal, anterior (or thymic).	Internal mammary.	Connective tissue, fat and lymphatics, in superior and anterior mediastina, thymus gland.	
Meningeal.	Ascending pharyngeal.	Membranes of brain.	
Meningeal.	Posterior ethmoidal.	Dura mater.	
Meningeal, anterior.	Internal carotid.	Dura mater.	
Meningeal, middle or great.	Internal maxillary.	Cranium and dura mater.	Anterior and posterior.
Meningeal, posterior.	Occipital.	Dura mater.	
Meningeal, posterior.	Vertebral.	Dura mater.	
Meningeal, small.	Internal maxillary.	Gasserian ganglion, walls of cavernous sinus, and dura mater.	
Mesenteric, inferior.	Abdominal aorta.	Lower half of large intestine.	Left colic, sigmoid, superior hemorrhoidal.
Mesenteric, superior.	Abdominal aorta.	Whole of small intestine and upper half of large.	Inferior pancreatico-duodenal, colica media, colica dextra, ileo-colic, vasa intestini tenuis.
Musculo-phrenic.	Internal mammary.	Diaphragm, fifth and sixth lower intercostal spaces, oblique muscles of abdomen.	Phrenic, anterior intercostals, muscular.
Nasal.	Ophthalmic.	Lacrymal sac and integuments of nose.	Lacrymal, and transverse nasal.
Obturator.	Anterior division, internal iliac.	Pelvis and thigh.	Iliac (or nutrient), vesical, pubic, external and internal pelvic.
Obturator, external.	Obturator.	Muscles about obturator foramen.	
Occipital.	Fourth branch of external carotid.	Muscles of neck, and scalp.	Sterno-mastoid, posterior meningeal, auricular, mastoid, princeps cervicis, communicating, muscular, terminal.
Omphalo-mesenteric.	Primitive aorta.	Subsequently becomes the umbilical.	
Ophthalmic.	Internal carotid.	The eye, adjacent structures, portion of face.	Lacrymal, supraorbital, central artery of retina, muscular, ciliary, posterior and anterior ethmoidal, palpebral, frontal, nasal.
Ovarian.	Abdominal aorta.	Ovary, ureter, Fallopian tube, uterus.	Ureteral, Fallopian, uterine, ligamentous.
Palatine.	Ascending pharyngeal.	Soft palate and its muscles.	
Palatine, ascending (or inferior).	First branch of facial.	Upper part of pharynx, palate and tonsils.	Palatine, tonsillar.
Palatine, descending.	Internal maxillary.	To soft and hard palate.	Anterior and posterior.
Palmar arch, deep.	Radial and communicating of ulna.	Palm and fingers.	Princeps pollicis, radial, palmar interosseous, carpal, posterior.

TABLE OF ARTERIES—*Continued.*

NAME.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Palmar arch, superficial.	Ulnar and superficialis volæ.	Palm and fingers.	Digital (4), muscular, cutaneous.
Pancreatic.	Hepatic.	Pancreas.	
Pancreatic.	Splenic.	Pancreas.	
Pancreatico-duodenal, inferior.	Superior mesenteric.	Pancreas and duodenum.	
Pancreatico-duodenal, superior.	Gastro-duodenal.	Duodenum and pancreas.	
Perforating (or posterior communicating), (3).	Deep palmar arch.	Interosseous spaces.	
Pharyngeal.	Pterygo-palatine.	Roof of pharynx.	
Pharyngeal.	Spheno-palatine.	Roof and contiguous portions of pharynx.	
Pharyngeal, ascending.	First branch external carotid.	Pharynx, soft palate, tympanum, posterior part of neck, and membranes of brain.	Prevertebral, pharyngeal, palatine, tympanic, meningeal.
Phrenic.	Ascending cervical.	Phrenic nerve.	
Phrenic, superior (or comes nervi phrenici).	Internal mammary.	Pleura, pericardium, and diaphragm.	
Plantar arch.	External plantar artery.	Anterior part of foot and toes.	Articular and plantar digital.
Plantar, deep.	Metatarsal.	Assists in formation of plantar arch.	
Plantar, external.	Posterior tibial.	Sole and toes.	Muscular, calcaneal, cutaneous, anastomotic, posterior perforating, plantar arch.
Plantar, internal.	Posterior tibial.	Inner side of foot.	Muscular, cutaneous, articular, anastomotic, superficial digital.
Popliteal.	Continuation of femoral.	Knee and leg.	Cutaneous, muscular (superior and inferior) or sural, articular, superior and inferior external, superior and inferior internal and sylvia, terminal (anterior and posterior tibial).
Profunda (or deep femoral).	Femoral.	Thigh.	External and internal circumflex, three perforating.
Profunda, inferior.	Brachial.	Triceps, elbow-joint.	
Profunda, superior.	Brachial.	Humerus, muscles and skin of arm.	Ascending, cutaneous, articular, nutrient muscular.
Pterygo-palatine (or pterygo-pharyngeal).	Internal maxillary.	Pharynx, Eustachian tubes, and sphenoidal cells.	Pharyngeal, Eustachian, sphenoidal.
Pudic, external, deep (or inferior).	Femoral, common.	Skin of scrotum (or labium in female).	
Pudic, external, superficial (or superior).	Common, femoral.	Integument above pubes and external genitalia.	
Pudic, internal.	Internal iliac, anterior division.	Generative organs.	External (in relation, haemorrhoidal, superficial perineal, muscular, arteries of penis, urethra, and ducts of glands of penis).

TABLE OF ARTERIES—*Continued.*

NAME.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Pulmonary.	Right ventricle.	Lunga.	Right and left.
Pyloric, inferior.	Gastro - duodenal or right gastro-epiploic.	Pyloric end of stomach.	
Pyloric, superior.	Hepatic.	Pyloric end of stomach.	
Radial.	Brachial.	Forearm, wrist, hand.	Radial recurrent, muscular, anterior and posterior carpal, superficial volar, metacarpal, dorsalis pollicis, dorsalis indicis, deep palmar arch.
Ranine.	Lingual.	Tongue and mucous membrane of mouth.	
Renal.	Abdominal aorta.	Kidney.	Inferior suprarenal, capsular, ureteral.
Sacra media. See	<i>Sacral, middle.</i>		
Sacral, middle.	Continuation of aorta.	Sacrum and coccyx.	
Scapular, dorsal.	Subscapular.	Muscles of infraspinous fossa.	Infrascapular.
Scapular, posterior.	Continuation of transverse cervical.	Muscles of scapular region.	Supra-spinous and infra-spinous, subscapular, muscular.
Sciatic.	Internal iliac, anterior division.	Pelvic muscles and viscera, and branches.	Coccygeal, inferior gluteal, muscular, anastomotic, articular cutaneous, comes nervi ischiadici, vesical, rectal, prostatic, etc.
Spermatic.	Abdominal aorta.	Scrotum and testis.	Ureteral, cremasteric, epididymal, testicular.
Spheno-palatine (or naso-palatine).	Internal maxillary.	Pharynx, nose, and sphenoidal cells.	Pharyngeal, sphenoidal, nasal, ascending septal.
Spinal.	Ascending cervical.	Spinal canal.	
Spinal.	Intercostals.	Spinal canal and spine.	
Spinal.	Lateral sacral.	Spinal membranes and muscles and skin over sacrum.	
Spinal, anterior.	Vertebral.	Spinal cord.	
Spinal, lateral.	Vertebral.	Vertebrae and spinal canal.	
Spinal, posterior.	Vertebral.	Spine.	
Splenic.	Celiac axis.	Spleen, pancreas, part of stomach, omentum.	Small and large pancreatic, left gastro-epiploic, vasa brevia, terminal.
Splenic.	Left phrenic.	Spleen.	
Subclavian.	<i>Right, innominate. Left, arch of aorta.</i>	Neck, thorax, arms, brain, meninges, etc.	Vertebral, thyroid axis, internal mammary, superior intercostal.
Subscapular.	Axillary.	Subscapularis, teres major, latissimus dorsi, serratus magnus, axillary glands.	Dorsal and infrascapular.
Suprascapular (or transversalis humeri).	Thyroid axis.	Muscles of shoulder.	Inferior sternomastoid, subclavian, nutrient, suprasternal, acromial, articular, subscapular, supraspinous and infraspinous.

TABLE OF ARTERIES—Continued

NAME.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Temporal.	External carotid.	Forehead, parietal gland, masseter muscle, etc.	Frontal, articular, masseteric, anterior auricular, transverse facial, middle, anterior and posterior temporal.
Temporal, deep, anterior.	Internal maxillary.	Anterior part of temporal fossa.	
Termatic.	Anterior communicating.	Lamina cerebri and corpus callosum.	
Thoracic, acromial.	Axillary.	Muscles of shoulder, arm, and chest.	Acromial, humeral, pectoral, circumflex.
Thoracic, alar.	Axillary.	Axillary glands.	
Thoracic, external.	See <i>Thoracic, long.</i>		
Thoracic, internal.	See <i>Mammary, in termal.</i>		
Thoracic, long (external mammary)	Axillary.	Pectoral muscles, serratus magnus, mammary and axillary glands.	
Thymic.	Internal mammary.	Connective tissue, fat and lymphatics of mediastinum, and thymus.	
Thyroid axis.	Subclavian.	Shoulder, neck, thorax, spine, mast.	Inferior thyroid, suprascapular and transverse cervical.
Thyroid, inferior.	Thyroid axis.	Larynx, esophagus, and muscles of neck.	Muscles ascending cervical, omohyoid, tracheal, and inferior thyroideus.
Thyroid, superior.	External carotid.	Glandular, sternohyoid, sternothyroid, thyroid gland.	Superior, sternohyoid, superior thyroideus, thyroideus.
Thyroidea ima.	Innominate (usually).	Thyroid body.	
Tibial, anterior.	Popliteal.	Leg.	Anterior and posterior tibial, peroneal, anterior tibial and lateral malleolus.
Tibial, posterior.	Popliteal.	Leg, heel, and foot.	Posterior tibial, anterior tibial, posterior tibial, posterior malleolus and lateral malleolus.
Tonsillar.	Ascending palatine.	Tonsil and Eustachian tube.	
Tonsillar.	Facial.	Tonsil and root of tongue.	
Transversalis colli.	Thyroid axis.	Neck and scapular region.	Transverse cervical and posterior auricular.
Ulnar.	Brachial.	Forearm, wrist, and hand.	Anterior and posterior ulnar, interosseous, ulnar, and posterior ulnar, posterior ulnar, and posterior ulnar.
Uterine.	Internal iliac, anterior branch.	Uterus.	Uterine, vaginal, ovary.
Uterine.	Ovarian.	Uterus.	
Vasa brevia.	Splenic.	Stomach.	
Vertebral.	Subclavian.	Neck and shoulders.	Deep cervical, superficial cervical, posterior auricular, and posterior auricular.

TABLE OF ARTERIES—Continued.

NAME.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Vesical, inferior.	Internal iliac, anterior division.	Bladder, prostate, seminal vesicles, and vagina (in female).	
Vesical, middle.	Superior vesical.	Bladder.	
Vesical, superior.	Internal iliac, anterior division.	Bladder.	Deferentia, ureteric, middle vesical (occasionally).
Vidian.	Internal maxillary.	Roof of pharynx, Eustachian tube, and tympanum.	Pharyngeal, Eustachian, tympanic.
Vitelline. See Om.	phalo-mesenteric.		

Arthragra (*ar-thra'-grah*) [*ἀρθρα*, a joint; *ἀρα*, seizure]. Gout.

Arthralgia (*ar-thral'-je'-ah*) [*ἀρθρα*, a joint; *ἄλγος*, pain]. Neuralgic pain in a joint.

Arthralgic (*ar-thral'-jik*) [*ἀρθρα*, a joint; *ἄλγος*, pain]. Relating to arthralgia.

Arthrectomy (*ar-threk'-to-me*) [*ἀρθρα*, a joint; *ἐκτομή*, a cutting out]. Excision of a joint.

Arthritic (*ar-thrit'-ik*) [*ἀρθρα*, a joint]. Relating to arthritis or to gout.

Arthritis (*ar-thrit'-tis*) [*ἀρθρα*, a joint; *τις*, inflammation]. Inflammation of a joint. **A. deformans**, chronic inflammation of a joint with deformity; rheumatoid arthritis. **A. fungosa**, tuberculous disease of the joints; white swelling. **A. Gouty**, that due to gout. **A. pauperum**. Synonym of *Rheumatoid Arthritis*. **A. Proliferating**. See *A. deformans*. **A. Rheumatoid**. Chronic rheumatoid arthritis; *Osteo-arthritis*; *Rheumatic gout*; *Nodular rheumatism*; *Arthritis deformans*; a chronic joint-affection characterized by inflammatory overgrowth of the articular cartilages and synovial membranes, with destruction of those parts of the cartilages subject to intraarticular pressure; there is progressive deformity. The nature of the disease is unknown; by some it is considered a trophic disorder consequent upon spinal changes. It occurs in the young, in whom it is often acute, and in the old, in whom it is of a more mild form. **A. Urethral**, gonorrheal rheumatism.

A. urtica, arthritis due to gout.

Arthritism (*ar'-thrit-izm*) [*ἀρθρα*, a joint]. Gout or the gouty diathesis.

Arthrobacterium (*ar-thro-bak'-te'-ri-um*). A bacterium forming arthrospheres.

Arthroclasia (*ar-thro-kla'-se-ah*) [*ἀρθρα*, a joint; *κλάειν*, to break]. The breaking down of ankyloses in order to produce free movement of a joint.

Arthrodosis (*ar-throd'-es-is*) [*ἀρθρα*, joint; *δῆσις*, a binding]. Fixation of paralyzed joints.

Arthrodia (*ar-throd'-de-ah*) [*ἀρθρωδία*, a kind of articulation]. A form of joint admitting of a gliding movement.

Arthrodynia (*ar-thro-din'-e-ah*) [*ἀρθρα*, a joint; *δύνη*, pain]. See *Arthralgia*.

Arthrography (*ar-throg'-ra-fe*) [*ἀρθρα*, a joint; *γράφειν*, to write]. A description of the joints.

Arthrogryposis (*ar-thro-grip'-o'-sis*) [*ἀρθρα*, a joint; *γρῦπος*, curved]. 1. Persistent contracture of a joint. 2. Tetany or tetanilla.

Arthroneuralgia (*ar-thro-nu-ral'-ge-ah*) [*ἀρθρα*, a joint; *νεῖρον*, nerve; *ἄλγος*, pain]. Neuralgic pain in a joint.

Arthropathy (*ar-throp'-a-the*) [*ἀρθρα*, a joint; *πάθος*, disease]. 1. Any joint-disease. 2. A peculiar trophic disease of the joints, sometimes occurring in locomotor ataxia and syringomyelia; rarely in general paralysis of the insane, and in disseminated sclerosis. It is also known as Charcot's joint.

Arthrophyma (*ar-thro-fī'-ma*) [*ἀρθρα*, a joint; *φῦμα*, a swelling]. Swelling of a joint.

Arthroplasty (*ar'-thro-plas-te*) [*ἀρθρα*, a joint; *πλάσσειν*, to form]. The making of an artificial joint.

Arthropodous (*ar-throp'-o-dus*) [*ἀρθρα*, a joint; *πούς*, a foot]. In biology, having jointed legs.

Arthropoysis (*ar-thro-pi'-o'-sis*) [*ἀρθρα*, a joint; *πύωσις*, suppuration]. Pus-formation in a joint.

Arthrosis (*ar-thro'-sis*) [*ἀρθρώσειν*, to fasten by a joint]. Articulation or jointing.

Arthrosphere (*ar'-thro-spōr*) [*ἀρθρα*, a joint; *σπῆρος*, a seed]. A permanent form assumed by some bacteria, chiefly the cocci. It is analogous to a spore.

Arthrotomy (*ar-throt'-o-me*) [*ἀρθρα*, a joint; *τείνειν*, to cut]. Incision of a joint.

Artiad (*ar'-te-ad*) [*ἀρτιος*, even]. In chemistry, a term used to designate atoms having an even valivalence, as *a. white* bivalent.

Articular (*ar-tik'-u-lar*) [*articularis*, of the joints]. Pertaining to an articulation or joint.

Articulate (*ar-tik'-u-lal*) [*articulare*, to divide in joints]. Divided into joints, distinct. **A. Speech**, the communication of ideas by spoken words.

Articulation (*ar-tik'-u-lal'-shun*) [*articulus*, a joint]. 1. A joint; a connection between two or more bones, whether allowing movement between them or not. The articulations are divided into: (1) *Synarthroses*, immovable, subdivided into *schindyleses*, or grooved joints, *gomphoses*, in sockets, as the teeth, and *suturæ*, as in the bones of the skull; (2) *Diarthroses*, or movable joints, subdivided into the *arthrodia*, or gliding joints, the *ginglymus*, or hinge-like, the *enarthroses*, or ball-and-socket joints; (3) *Amphiarthroses*, or those of a mixed type. 2. The enunciation of spoken speech.

Articulo mortis, In (*ar-tik'-u-lo-mor'-tis*) [l.]. At the moment of death. In the act of dying.

Artifact (*ar'-te-fakt*) [*arte*, by art; *factum*, made]. See *Artefact*.

Artificial (*ar-te-fish'-al*) [*artificialis*]. Made or imitated by art. **A. Anus**, an opening in the abdomen or loin to give exit to the feces. **A. Eye**, a film of glass, celluloid, rubber, etc., made in imitation of the front part of the globe of the eye and worn in the socket or over a blind eye for cosmetic reasons. **A. Feeding**, the feeding of an infant by other means than mother's milk. Various mixtures and foods are to be had, such as *Meigs's*, *Rotch's*, *Biedert's Mixtures*, etc. See *Table at end of this article*. **A. Leech**. See *Leech*. **A. Palate**, a mechanical contrivance for supplying the loss of the whole or a portion of the hard or soft palate, or both. **A. Pupil**, removal of a piece of the iris (*iridectomy*, *iridodialysis*, etc.), to allow the light to pass through the opening. **A. Respiration**, the aeration of the blood by artificial means. A method of inducing the normal function of respiration when from any cause it is temporarily in abeyance, as in asphyxia neonatorum, drowning, etc. *Bain's Method*: a modification of Sylvester's method, the axilla itself being seized so that the traction is made directly upon the pectoral muscles. *Byrd's (H. L.) Method*: the physician's hands are placed under the middle portion of the child's back with their ulnar borders in contact and at right angles to the spine. With the thumbs extended, the two extremities of the trunk are carried forward by gentle but firm pressure, so that they form an angle of 45 degrees with each other in the diaphragmatic region. Then the angle is reversed by carrying backward

the shoulders and the nates. *Dew's Method*: the infant is grasped in the left hand, allowing the neck to rest between the thumb and forefinger, the head falling far over backward. The upper portion of the back and scapulæ rest in the palm of the hand, the other three fingers being inserted in the axilla of the babe's left arm, raising it upward and outward. The right hand grasps the babe's knees, and the lower portion of the body is depressed to favor inspiration. The movement is reversed to favor expiration, the head, shoulders, and chest being brought forward and the thighs pressed upon the abdomen. *Forest's Method*: the child is placed on its face and quick, firm pressure is made on the back; then it is placed in a pail of hot water, and the hands carried upward until the child is suspended by its arms, and mouth-to-mouth insufflation is practised; the arms are then lowered and the body doubled forward; these movements are repeated at the rate of 40 per minute. *Hall's Method*: by turning the body alternately upon the side or face to compress the chest, and then upon the back to allow the lungs to expand. *Howard's Method*: by pressure upon the lower ribs every few seconds. *Pacini's Method*: for resuscitating asphyxiated infants. The child lying on its back, the operator stands at its head, and grasps the axillæ, pulling the shoulders forward and upward to compress the thorax, and allowing them to fall in order to expand the chest. *Satterthwaite's Method*: pressure upon the abdomen alternating with relaxation to allow descent of the diaphragm. *Schrader's Method*: the babe while in a bath is supported by the operator on the back, its head, arms, and pelvis being allowed to fall backward; a forceful expiration is then effected by bending up the babe over its belly, thereby compressing the thorax. *Schultze's Method*: the child is seized from behind with both hands, by the shoulders, in such a way that the right index finger of the operator is in the right axilla of the child from behind forward, and the left index finger in the left axilla, the thumbs hanging loosely over the clavicles. The other three fingers hang diagonally downward along the back of the thorax. The operator stands with his feet apart and holds the child as above, practically hanging on the index fingers in the first position, with the feet downward, the whole weight resting on the index fingers in the axillæ, the head being supported by the ulnar borders of the hands. This is the first inspiratory position. At once the operator swings the child gently forward and upward. When the operator's hands are somewhat above the horizontal the child is moved gently, so that the lower end of the body falls for-

ward toward its head. The body is not flung over, but moved gently until the lower end rests on the chest. In this position the chest and upper end of the abdomen are compressed tightly. The child's thorax rests on the tips of the thumbs of the operator. As a result of this forcible expiration the fluids usually pour out of the nose and mouth of the infant. The child is allowed to rest in this position (the first expiratory position) about one or two seconds. The operator gradually lowers his arms, the child's body bends back, and he again holds the infant hanging on his index fingers with its feet downward; this is the second inspiratory position. These movements are repeated 15 to 20 times in the minute. *Sylvester's Method* consists chiefly of movements of the arms. This method is valueless in asphyxia neonatorum, owing to non-development of the pectoral muscles.

ARTIFICIAL FEEDING OF INFANTS.

TABLE OF QUANTITY REQUIRED.

AGE.	INTERVAL.	NUMBER OF FEEDINGS IN 24 HOURS.	AMOUNT OF FOOD AT EACH FEEDING.	TOTAL AMOUNT IN 24 HOURS.
1st week, . . .	2 hours.	10	1 oz.	10025.
2d to 4th week, . .	2 "	9	1½ ozs.	13½ "
2d to 3d month, . .	3 "	6	3 "	18 "
3d to 4th month, . .	3 "	6	4 "	24 "
4th to 5th month, . .	3 "	6	4-4½ "	24-27 "
6th month, . . .	3 "	6	5 "	30 "
8th month, . . .	3 "	6	6 "	36 "
10th month, . . .	3 "	5	8 "	40 "

Aryepiglottic (*ar-ep-e-glōt'-ik*). Same as *Arytenoepiglottic*.

Arytenoepiglottic (*ar-i-ten-o-ep-e-glōt'-ik*) [*arytana*, a pitcher; *eidōs*, likeness; *ēpi*, upon; *glōt'tis*, glottis]. Relating to an arytenoid cartilage and to the epiglottis; as the arytenoepiglottic fold (or folds), consisting of a fold of mucous membrane that extends from each arytenoid cartilage to the epiglottis.

Arytenoid (*ar-i-tē-nōid'*) [*arytana*, a pitcher; *ēidōs*, likeness]. Resembling the mouth of a pitcher. **A. Cartilages**, two cartilages of the larynx regulating, by the means of the attached muscles, the tension of the vocal bands. **A. Glands**, muciparous glands found in large numbers along the posterior margin of the arytenoepiglottic fold in front

of the arytenoid cartilages. **A. Muscle**, a muscle arising from the posterior surface of one arytenoid cartilage and inserted into the corresponding parts of the other. It is composed of three planes of fibers, two oblique and one transverse. It draws the arytenoid cartilages together.

Arythmia (*ar-ith'-me-ah*). See *Arrhythmia*.

Arythmic (*ar-ith'-mik*). See *Arrhythmic*.

Asafetida, **Asafotida** (*as-a-fet'-id-ah*) [*asa*, gum; *fetida*, stinking]. A resinous gum obtained from the root of *Ferula narihex* and *F. scordosoma*. It is slightly soluble in alcohol and forms an emulsion with water. Its properties are due to *allyl sulphid*, C₆H₁₀S. It is a powerful antispasmodic, stimulant, and expectorant, and is used in hysteria and bronchial affections. Dose gr. v-xx (0.32-1.3). **A.**, Tinct., strength 20 per cent. Dose ʒss-ij (2.0-8.0). **A.**, Emplastrum, asafetida 35, lead plaster 35, galbanum 15, yellow wax 15, alcohol 120 parts. **A.**, Emulsum, a 4 per cent. emulsion. Dose ʒss-ʒij (16.0-64.0). **A.**, Enema (B. P.), asafetida and distilled water. **A. et Magnesium carb.** 5, tinct. asafetida 7, tinct. opii 1, sugar 10, distilled water sufficient to make 100 parts. Dose ʒss-ʒss (2.0-16.0). **A.**, Pilulæ, contain each gr. iij (0.19) of asafetida and gr. j (0.065) of soap. Dose j-iv. **A. et Aloes**, Pil., have gr. 1½ (0.087) of each ingredient. **A.**, Pil., Comp. (B. P.). Pil. galbani comp. It consists of asafetida, galbanum, myrrh, and treacle. Dose gr. v-xx (0.32-1.3).

Asaphia (*as-a' fe-ah*) [*ἀσάφεια*, indistinctness]. Indistinctness of utterance, especially that due to cleft palate.

Asaprol (*as'-ap-rol*), Ca(OH.C₁₀H₆SO₃)₂·3Aq. Calcium-β-naphthol-α-mono-sulphonate, a substance readily soluble in water and alcohol, and recommended in asthma, tonsillitis, and acute articular rheumatism, in doses of from 15 to 60 grains (1.0-4.0).

Asarol (*as'-ar-ol*) [*ἀσάρων*, asarabacca; *oleum*, oil], C₁₀H₁₄O. A camphor-like body derived from asarum.

Asarum (*as'-ar-um*) [*ἀσάρων*, asarabacca]. A genus of aristolochiaceae plants. **A. canadense**, called wild ginger, Canada snake-root, with other N. American species, is used chiefly in domestic practice. **A. canadense** is a fragrant, aromatic stimulant. Dose of fld. ext. ʒ xv-ʒss (1.0-2.0). **A. europæum** has diaphoretic, emetic, purgative, and diuretic qualities, but is now little used except in veterinary practice.

Ascariasis (*as-kar-i-as-iis*) [*ἀσκαρίς*, a species of intestinal worm]. The symptoms produced by the presence of ascariæ in the gastro-intestinal canal.

Ascaricide (*as-kar'-is-id'*) [*ἄσκαρις*, a species of intestinal worm; *cādere*, to kill]. A medicine that kills ascarides.

Ascaris (*as'-kar-is*) [*ἄσκαρις*, a species of intestinal worm: *pl.*, *Ascarides*]. A genus of parasitic worms inhabiting the intestines of most animals. *A. lumbricis* is found in the ox, hog, and man. It inhabits the small intestine, especially of children. *A. mystax*, the round-worm of the cat, and *A. alata* have rarely been found in man. *A. vermicularis*, a synonym of *Oxyuris vermicularis*.

Ascending (*as-end'-ing*) [*ascendere*, to rise]. Taking an upward course; rising (as parts of the aorta and colon, and as one of the *venæ cavæ*). *A. Current*, in electricity, one going from the periphery to a nerve-center. *A. Degeneration*, a degeneration of the nerve-fibers extending from the periphery to the center, or, in the spinal cord, from below upward toward the brain. *A. Paralysis*. See *Paralysis*. *A. Tracts*, the centripetal tracts of the spinal cord, carrying afferent impulses.

Ascherson's Vesicles. The peculiar small globules formed when oil and an albuminous fluid are agitated together, and formerly thought to be cells.

Ascia (*as'-se-ah* or *as'-ke-ah*) [*ἄ priv.*; *σκία*, shadow]. *Fascia spiralis*; *dolabra currens*; a spiral bandage applied without reverses, each turn of which overlaps the preceding for about a third of its width. *Dolabra repens*, is the same as the preceding, in which, however, the spirals are formed more obliquely, and do not overlap each other, but are separated by a greater or less interval.

Ascites (*as-i'-tiz*) [*ἀσκίτης*, a kind of dropsy; from *ἄσκος*, a bag]. An abnormal collection of serous fluid in the peritoneal cavity; dropsy of the peritoneum. It is either local in origin or a part of a general dropsy. The ascitic fluid is usually clear, yellow, and coagulates on standing. It may be turbid, blood-stained, and contain lymph-particles or shreds. There is a uniform enlargement of the abdomen, fluctuation, percussion-dulness. Its usual cause is cirrhosis of the liver. *A. adiposus*, ascites characterized by a fluid milky appearance due to the presence in it of numerous cells that have undergone fatty degeneration and solution. It is seen in certain cases of carcinoma, tuberculosis, and other chronic inflammations of the peritoneum. *A. chylosus*, the presence of chyle in the peritoneal cavity. It follows rupture of a chyle duct.

Ascitic (*as-i'-ik*) [*ἀσκίτης*, a kind of dropsy]. Pertaining to or affected with ascites.

Asclepiadin (*as-kle-pi'-ad-in*) [*Ἀσκληπιός*]. A bitter glucosid obtainable from various species of *Asclepias*. It is poisonous, and has emetic, purgative, and sudorific properties. Unof.

Asclepias (*as-kle'-pe-as*) [*Ἀσκληπιός*]. Pleurisy Root. The root of *Asclepias tuberosa*. A popular remedy in the Southern States for pleurisy. It is diaphoretic, emetic, and cathartic. The infusion recommended has a strength of $\frac{3}{4}$ of the powdered root to $\frac{3}{4}$ xxxij of water. Dose, a teacupful every three or four hours. Also, *A. curassavica*, blood flower, an herb common to tropical America; astringent, styptic, and anthelmintic against the tape-worm. *A.*, Ext. fld. Dose $\frac{1}{2}$ xx-f $\frac{3}{4}$ j (1.3-4.0).

Asclepin (*as-kle'-pin*) [*Ἀσκληπιός*]. 1. A poisonous principle obtainable from asclepiadin by the separation of glucose from the latter. 2. The precipitate from a tincture of *Asclepias tuberosa*; alterative, evacuant, tonic, sedative. Dose grs. ij-iv (0.13-0.27). Unof.

Ascococcus (*as-ko-kok'-us*) [*ἄσκος*, a leather bag; *κόκκος*, a kernel]. A genus of the family of Schizomycetes. The ascococci are microorganisms made up of round or ovoid cells united in massive colonies, and surrounded by tough, thick, gelatinous envelopes. *A. billrothii*, found in putrefied meat; its natural habitat is the air; it is probably not pathogenic.

Ascospore (*as'-ko-spōr*) [*ἄσκος*, a bag; *σπῆρος*, seed]. A spore produced by or in an ascus.

Ascus (*as'-kus*) [*ἄσκος*, a bag or bladder]. The characteristic spore case of some fungi and lichens, usually consisting of a single terminal cell containing eight spores.

Asemasia (*as-em-a'-ze-ah*) [*ἄ priv.*; *σημασία*, a signaling]. Absence of the power to communicate either by signs or language.

Asepsin (*ah-sep'-sin*). See *Antisepsin*.

Asepsis (*ah-sep'-sis*) [*ἄ priv.*; *σῆψις*, putrefaction]. Absence of pathogenic microorganisms.

Aseptic (*ah-sep'-tik*) [*ἄ priv.*; *σῆψις*, putrefaction]. Free from pathogenic bacteria, as aseptic wounds. *A. Surgery*, the mode of surgical practice in which everything that is used as well as the wound is in a germ-free condition.

Asepticism (*ah-sep'-tis-izm*) [*ἄ priv.*; *σηπτός*, septic]. The doctrine or principles of aseptic surgery.

Asepticize (*ah-sep'-tis-iz*) [*ἄ priv.*; *σηπτός*, septic]. To render aseptic.

Aseptin (*ah-sep'-tin*) [*ἄ priv.*; *σηπτός*, septic]. A secret preparation containing boric acid, used for preserving articles of food.

Aseptol (*ah-sep'-tol*) [*ἄ priv.*; *σηπτός*, septic], $C_6H_6SO_4$. Sulphocarbolic acid; sozolic acid. A reddish liquid, with an odor of carbolic acid, recommended as a disinfectant and antiseptic. It is used externally (1-10 per cent. solution), and internally in about the same dose as carbolic acid.

Asexual (*ah-seks'-u-al*) [*ἄ priv.*; *sexus*, sex]. Without sex; non-sexual.

Ash (*ash*) [M.E., *asch*]. 1. The incombustible mineral residue that remains when a substance is incinerated. 2. See *Fraxinus*. **A. Manna**. See *Fraxinus*. **A., Prickly**. See *Xanthoxylum*.

Asialia (*as-e-a'-le-ah*) [*â* priv.; *σίαλον*, spittle]. Defect or failure of the secretion of saliva.

Asiatic (*a-ze-at'-ik*) [*Asia*]. Pertaining to or belonging to Asia. **A. Cholera**. See *Cholera*. **A. Pill**, a pill composed of arsenous acid, black pepper, powdered licorice, and mucilage.

Asitia (*ah-sit'-e-ah*) [*â* priv.; *σιτος*, food]. The want of food. Also a loathing for food.

Asomata (*ah-so'-mat-ah*) [*â* priv.; *σῶμα*, body]. A species of omphalositic monster characterized by an absence of the trunk. The head is never well-formed, and the vessels run from it to the placenta in the membranes. Beneath the head is a sac in which rudiments of body-organs may be found. This is the rarest form of omphalositic.

Aspalasoma (*as-pal-as-o'-mah*) [*ἀσπάλασξ*, mole; *σῶμα*, body]. A variety of single autotomic monsters of the species celosoma, in which there is a lateral or median eventration occupying principally the lower portion of the abdomen, with the urinary apparatus, the genital apparatus, and the rectum opening externally by three distinct orifices.

Asparagin (*as-par'-ag-in*) [*ἀσπάραγος*, asparagus]. $C_4H_8N_2O_3 \cdot H_2O_3$. An alkaloid found in the seeds of many plants, in asparagus, beet root, peas, and beans. It forms shining, four-sided, rhombic prisms, readily soluble in hot water, but not in alcohol or ether. It is an amid of aspartic acid, and forms compounds with both acids and bases. It is diuretic. The hydrargyrate of asparagin has been used as an antisyphilitic, in doses of $\frac{1}{2}$ grain (0.01) hypodermically.

Asparaginic Acid, $C_4H_8NC_2$. A dibasic acid, a decomposition-product of asparagin and also of proteids and gelatin.

Asparagus (*as-par'-ag-us*) [*ἀσπάραγος*, asparagus]. The green root of *Asparagus officinalis*, a mild diuretic. Dose of fld. ext. $\overline{3}$ ss-j (2.0-4.0). Unof.

Aspartic Acid (*as-par'-tik*). See *Acid*.

Aspergillus (*as-per-jil'-us*) [*aspergere*, to scatter]. A genus of fungi. **A. auricularis**, a fungus found in the wax of the ear. **A. fumigatus**, found in the ear, nose, and lungs. **A. glaucus**, the bluish mold found upon dried fruit. **A. mucoroides**, a species found in tuberculous or gangrenous lung-tissue.

Aspergillusmycosis (*as-per-jil'-us-mi-ko'-sis*). See *Otomycosis*.

Aspermatic (*ah-sper-mat'-ik*) [*â* priv.; *σπέρ-*

μα, seed]. Affected with or relating to aspermaticism.

Aspermatism (*ah-sper'-mat-izm*) [*â* priv.; *σπέρμα*, seed]. Defective secretion of semen or lack of formation of spermatozoa.

Aspermous (*ah-sper'-mus*) [*â* priv.; *σπέρμα*, seed]. Without seed.

Aspersio (*as-per'-shun*) [*aspersio*, *aspergere*, to sprinkle]. Treatment of disease by sprinkling the body or the affected part with a medicinal agent.

Asphyctic, or **Asphyctous** (*as-fik'-tik*, *as-fik'-tus*) [*â* priv.; *σφίξις*, the pulse]. Affected with asphyxia.

Asphyxia (*as-fiks'-e-ah*) [*â* priv.; *σφίξις*, the pulse]. Suffocation; the suspension of vital phenomena resulting when the lungs are deprived of oxygen. The excess of carbon dioxid in the blood at first stimulates, then paralyzes the respiratory center of the medulla. Artificial respiration is therefore required in cases of asphyxia. **A., Local**, that stage of Raynaud's disease in which the affected parts are dusky red from intense congestion. **A. neonatorum**, the asphyxia of the new-born from any cause.

Asphyxiant (*as-fiks'-e-ant*) [*ἀσφύκτωρ*, pulseless]. 1. Producing asphyxia. 2. An agent capable of producing asphyxia.

Asphyxiate (*as-fiks'-e-āt*) [*â* priv.; *σφίξιν*, to pulsate]. To produce or cause asphyxia.

Aspidium (*as-pid'-e-um*) [L.; gen. *Aspidii*].

1. A genus of ferns, known as shield ferns. 2. The rhizome of *Dryopteris filix mas* and of *D. marginalis*, or male fern. Its properties are due to a resin containing filicilic acid. It is valuable chiefly against tape-worm. Dose $\overline{3}$ ss- $\overline{3}$ ss (2.0-16.0). **A., Oleoresina**, an ethereal extract. Dose $\overline{3}$ ss-j (2.0-4.0). **Extractum Filicis Liquidum** (B. P.). Dose \overline{m} xv- $\overline{3}$ j (1.0-4.0).

Aspidosamin (*as-pid-os'-am-in*). $C_{22}H_{38}N_2O_2$. A basic principle, from quebracho bark. It is emetic. Unof.

Aspidosperma (*as-pid-o-sper'-mah*) [*ἀσπίς*, a shield; *σπέρμα*, a seed]. A genus of apocynaceous trees, of which the *Quebracho* is the most important.

Aspidospermin (*as-pid-o-sper'-min*) [*ἀσπίς*, shield; *σπέρμα*, seed]. $C_{22}H_{38}N_2O_2$. An alkaloid extracted from Quebracho. (*Aspidosperma quebracho*). It has the general effects of quebracho. Unof.

Aspiration (*as-pir-a'-shun*) [*ad*, to; *spirare*, to breathe]. 1. The act of sucking up or sucking in; inspiration, imbibition. 2. The act of using the aspirator. 3. A method of withdrawing the fluids and gases from a cavity. **A. Pneumonia**. See *Pneumonia*.

Aspirator (*as'-pir-a-tor*) [*ad*, to; *spirare*, to breathe]. An apparatus for withdrawing liquids from cavities by means of suction.

Asporogenic (*ah-spôr-o-jen'-ik*) [*â* priv.; *σπόρος*, seed; *γενής*, producing]. Not reproducing by means of spores; not producing spores.

Assault (*as-awlt'*) [*assalire*, to assail]. An attack. **A.**, Criminal, in medical jurisprudence, the touching or attempting to touch, on the part of a male, any of the sexual organs (the breasts included) of a female against her will, even though they be covered by clothing.

Assideration (*as-id-er-a'-shun*) [*ad*, intensive; *sideratio*, an evil influence]. In forensic medicine, infanticide by immersing in ice-water.

Assimilable (*as-in'-il-a-bl'*) [*assimulare*, to make like]. Capable of being assimilated; nutritious.

Assimilation (*as-im-il-a'-shun*) [*assimulare*, to make like]. The process of transforming food into such a nutrient condition that it is taken up by the circulatory system, to form an integral part of the economy; synthetic or constructive metabolism; anabolism. **A.**, Mental, the mental reception of impressions and their assignment by the consciousness to their proper place. **A.**, Primary, that concerned in the conversion of food into chyle and blood. **A.**, Secondary, that relating to the formation of the organized tissues of the body.

Associated (*as-o'-se-a-ted*) [*associatus*, united]. Joined. **A. Movements**, coincident or consensual movements of other muscles than the leading one, and which by habit or unity of purpose are involuntarily connected with its action. Both eyeballs move alike in reading, though one be a blind eye. Movement of the normal arm will sometimes produce slight motion of the opposite paralyzed arm. Uniformity of innervation is usually the cause of these movements. **A. Paralysis**, **A. Spasm**, a common paralysis, or spasm, of associated muscles.

Assurin (*as'-u-rin*), $C_{44}H_{84}N_4P_2O_9$. A name given by Thudichum to a complex substance occurring in brain-tissue.

Astasia (*as-ta'-se-ah*) [*â* priv.; *στάσις*, standing]. Motor incoordination for standing. **A.-Abasia**, a symptom consisting in inability to stand or walk in a normal manner. The person affected seems to collapse when attempting to walk.

Asteatosis (*as-te-at-o'-sis*) [*â* priv.; *στεατ*, tallow; *ώδης*, fulness]. 1. A deficiency or absence of the sebaceous secretion. 2. Any skin disease (as xeroderma) characterized by scantiness or lack of the sebaceous secretion. **A. cutis**, a condition of diminished sebaceous secretion, as the result of which the skin becomes dry, scaly, and often fissured.

Aster (*as'-ter*) [*aster*, a star]. The stellate form assumed by the mitome of the nucleus when undergoing karyokinesis.

Asterion (*as-te'-re-on*) [*ἀστήρ*, star]. A point on the skull corresponding to the junction of the occipital, parietal, and temporal bones.

Asternal (*ah-ster'-nal*) [*â* priv.; *στέρνον*, the breast-bone]. Not connected with the sternum. **A. Ribs**, the five lower pairs, because not joined directly to the sternum.

Asternia (*ah-ster'-ne-ah*) [*â* priv.; *στέρνον*, the breast-bone]. Absence of the sternum.

Asteroid (*as'-ter-oid*) [*ἀστήρ*, a star; *εἶδος*, resemblance]. Shaped like a star.

Asthenia (*as-then-i'-ah*) [*ἀσθένεια*; *â* priv.; *σθένος*, strength]. Absence of strength; adynamia.

Asthenic (*as-then'-ik*) [*â* priv.; *σθένος*, strength]. Characterized by asthenia.

Asthenometer (*as-then-oni'-et-er*) [*ἀσθένεια*, want of strength; *μέτρον*, a measure]. An instrument for detecting and measuring asthenia; especially, a device for measuring muscular asthenopia.

Asthenopia (*as-then-o'-pe-ah*) [*â* priv.; *σθένος*, strength; *ὤψ*, eye]. Weakness of the ocular muscles or of visual power, due to errors of refraction, heterophoria, over-use, anemia, etc. **A.**, Accommodative, due to hyperopia, astigmatism, or a combination of the two, producing strain of the ciliary muscle. **A.**, Muscular, due to weakness, incoordination (heterophoria), or strain of the external ocular muscles. **A.**, Retinal, or Nervous, a rare variety, caused by retinal hyperesthesia, anesthesia, or other abnormality, or by general nervous affections.

Asthenopic (*as-then-op'-ik*) [*â* priv.; *σθένος*, strength; *ὤψ*, eye]. Characterized by asthenopia.

Asthma (*as'-mah*) [*ἄσθμα*, panting]. A paroxysmal affection of the bronchial tubes characterized by dyspnea, cough, and a feeling of constriction and suffocation. The disease is probably a neurosis, and is due to hyperemia and swelling of the bronchial mucous membrane, with a peculiar secretion of a mucin-like substance. The attacks may be caused by direct irritation of the bronchial mucous membrane or by indirect or reflex irritation, as from the nose, the stomach, the uterus. When dependent upon disease of the heart, the kidneys, stomach, thymus, etc., it has been designated, *cardiac*, *renal*, *peptic*, *thymic*, etc. **A.**, Bronchial. See *Asthma*. **A.**, Cardiac, paroxysmal dyspnea due to heart-disease. **A. convulsivum**. Synonym of *Bronchial Asthma*. **A. Crystals**, acicular crystals (Charcot-Leyden crystals) contained in the sputum of asthmatic patients. They are generally associated with eosinophile cells. **A. dyspepticum**, is due to nervous reflexes

through the vagus. **A., Grinders'.** Same as *Fibroid Phthisis*. **A., Hay.** See *Hay Fever*. **A., Heberden's,** angina pectoris. **A., Kopp's,** spasm of the glottis. **A., Marine.** See *Beriberi*. **A., Miller's.** See *Laryngismus stridulus*. **A., Miner's.** See *Anthraxosis*. **A. nervosum.** Synonym of *Bronchial Asthma*. **A., Renal,** a paroxysmal dyspnea sometimes occurring in the course of Bright's disease. **A., Spasmodic.** See *Asthma*. **A., Thymic.** Synonym of *Laryngismus stridulus*.

Asthmatic (*as-mat'-ik*) [*ἀσθμα*, panting]. Relating to or affected with asthma.

Astigmatic (*as-tig'-mat'-ik*) [*ἀ priv.; στίγμα*, a point]. Pertaining to or affected with astigmatism.

Astigmatism (*as-tig'-mat-izm*) [*ἀ priv.; στίγμα*, a point, because rays of light from a point are never brought to a point by the refractive media of the eye]. That condition of the eye in which rays of light do not converge to a point on the retina. It is usually due to inequality of curvature of the different meridians of the cornea (*corneal A.*), but may be caused by imperfections of the lens (*lenticular A.*), unequal contraction of the ciliary muscle, or may perhaps be due to retinal imperfection. It may be **acquired** or **congenital**, and may complicate hypermetropia or myopia, producing either **simple hypermetropic A.**, in which one principal meridian is enmetropic, the other hypermetropic; or **compound hypermetropic A.**, in which both meridians are hypermetropic, but one more so than the other. Complicating myopia, we may in the same way have **simple myopic** or **compound myopic A.** In **mixed A.**, one principal meridian is myopic, the other hypermetropic. **Regular A.** is when the two principal meridians are at right angles to each other. **Irregular A.**, when different parts of a meridian have different refracting powers.

Astigmatometer (*as-tig'-mat-om'-et-er*) [*ἀ priv.; στίγμα*, a point; μέτρον, a measure]. An instrument for measuring the degree of astigmatism.

Astigmometer (*as-tig'-mom'-et-er*). See *Astigmatometer*.

Astigmometry (*as-tig'-mom'-et-re*) [*ἀ priv.; στίγμα*, a point; μέτρον, a measure]. The measurement of astigmatism.

Astragalectomy (*as-trag'-al-ek'-to-me*) [*ἀσπράγαλος*, astragalus; ἐκτομή, excision]. Excision of the astragalus.

Astragaloscaphoid (*as-trag'-al-o-scaf'-oid*) [*ἀσπράγαλος*, astragalus; σκαφοειδής, like a boat]. Relating to the astragalus and the scaphoid bone.

Astragalus (*as-trag'-al-us*) [*ἀσπράγαλος*, a dice; the analogous bones of the sheep were

used by the ancients as dice]. 1. The ankle-bone, upon which the tibia rests. 2. A genus of leguminous plants from some varieties of which gum tragacanth is derived. **A. mollissimus**, Loco Plant. The active principle of this plant has mydriatic properties. Unof.

Astrapaphobia (*as-trap'-af-o'-be-ah*) [*ἀστραπή*, lightning; φόβος, fear]. Morbid fear of lightning.

Astrokinetic (*as-tro-kin-et'-ik*) [*ἀστρον*, a star; κινεῖν, to move]. Applied to the phenomena of motion as exhibited by the centrosomes of cells.

Astrostatic (*as-tro-stat'-ik*) [*ἀστρον*, a star; ἵστασθαι, to stand]. Applied to the resting condition of the centrosomes of cells.

Asylum (*as-il'-i-um*) [*asylum*, a place of refuge]. An institution for the support, safe-keeping, cure, or education of those incapable of caring for themselves, such as the insane, the blind, etc. **A. Ear.** See *Hemiatoma auris*.

Asymbolia (*ah-sim-bo'-le-ah*) [*ἀ priv.; σύμβολον*, symbol]. The loss of all power of communication, even by signs or symbols.

Asymmetry (*ah-sim'-et-re*) [*ἀ priv.; συμμετρία*, symmetry]. 1. Unlikeness of corresponding organs or parts of opposite sides of the body that are normally of the same size, etc., e. g., asymmetry of the two halves of the skull or brain. 2. The linking of carbon atoms to four different groups; the combination of carbon atoms with different atoms or atomic groups.

Asynergy (*ah-sin'-er-je*) [*ἀ priv.; συνεργία*, cooperation]. Faulty coordination of the different organs or muscles normally acting in unison.

Asynesia (*as-in-e'-ze-ah*) [*ἀσυνεσία*, stupidity]. Stupidity; loss or disorder of mental power.

Asynodia (*ah-sin-o'-de-ah*) [*ἀ priv.; συνωδία*, a traveling together]. Impotence.

Asystole (*ah-sis'-to-le*) [*ἀ priv.; συστολή*, a shortening]. Imperfect contraction of the ventricles of the heart.

Asystolic (*ah-sis-to'-ik*) [*ἀ priv.; συστολή*, a shortening]. Characterized by asystole.

Atactic (*at-ak'-tic*) [*ἀτακτος*, irregular]. Irregular, incoordinate. Pertaining to muscular incoordination, especially in aphasia.

Atavic (*at'-av-ik*) [*atavus*, a forefather]. Relating to or characterized by atavism.

Atavism (*at'-av-izm*) [*atavus*, a forefather]. A variety of heredity characterized by the reappearance of a peculiarity in an individual whose more or less remote progenitors possessed the same peculiarity but whose immediate ancestors did not present it.

Atavistic (*at-av-is'-tik*). Same as *Atavic*.

Ataxia (*at-aks'-e-ah*) [*ἀταξία*, want of order]. Incoordination of muscular action. **A.,**

Cerebellar, A., Cerebral, A., Spinal, that due to disease of the cerebellum, of the brain, or of the spinal cord. **A., Briquet's**, an hysteric state marked by loss of sensation in the skin and in the leg-muscles. **A. cordis**. See *Delirium cordis*. **A. Family**. See *A., Hereditary*. **A., Hereditary**. See *Friedreich's Disease*. **A., Locomotor**, a synonym of tabes dorsalis, or posterior spinal sclerosis, a disease of the posterior columns of the spinal cord, characterized by static and motor ataxia, by fulgurant pains, girdle-sensation, the Robertson pupil, disturbances of sensation and of the sphincters, and the loss of the patellar reflex. **A., Motor**, inability to coordinate the muscles in walking. **A., Static**, the failure of muscular coordination in standing still, or in fixed positions of the limbs. **A., Thermal**, peculiar large and irregular fluctuations of the body-temperature, due to a condition of incoordination, or a disordered or weakened thermotaxic mechanism. This may give rise to the so-called *paradoxical* or *hysteric temperatures*, rising occasionally to 108° or 110° F., without grave or permanent injury. **A., Vasomotor**. See *Vasomotor Ataxia*.

Ataxic (*at-aks'-ic*) [*ἀταξία*, want of order]. 1. Pertaining to or affected with ataxia. 2. A person affected with ataxia. **A. Aphasia**. See *Aphasia*. **A. Fever**. Synonym of Typhus Fever.

Ataxophemia (*at-aks-o-fe'-me-ah*) [*ἀταξία*, want of order; *φήμη*, to speak]. Lack of coordination in speech.

Ataxy (*at-aks'-e*). See *Ataxia*.

Atelectasis (*at-el-ek'-tas-is*) [*ἀτελής*, imperfect; *ἐκτασις*, expansion]. Imperfect expansion or collapse of the air-vesicles of the lung. It may be present at birth, or may be acquired from diseases of the bronchi or lungs.

Atelectatic (*at-el-ek'-tat'-ik*) [*ἀτελής*, imperfect; *ἐκτασις*, expansion]. Relating to or characterized by atelectasis.

Ateloccephalous (*at-el-e-o-sef'-al-us*) [*ἀτελής*, imperfect; *κεφαλή*, head]. Having the skull or head more or less incomplete.

Atelia (*at-el-i'-ah*) [*ἀτέλεια*, imperfection]. Imperfect development. The word is compounded with others to designate the part affected, as *atelocardia*, etc., imperfect development of the heart, etc. Cf. *Asthenia*.

Atelocardia (*at-el-o-kar'-de-ah*) [*ἀτελής*, imperfect; *καρδία*, heart]. An imperfect or undeveloped state of the heart.

Atelochilia (*at-el-o-ki'-le-ah*) [*ἀτελής*, imperfect; *χείλος*, lip]. Defective development of a lip.

Atelochiria (*at-el-o-ki'-re-ah*) [*ἀτελής*, imperfect; *χείρ*, hand]. Defective development of the hand.

Ateloencephalia (*at-el-o-en-sef-a'-le-ah*) [*ἀτελής*, incomplete; *ἐγκέφαλον*, brain]. Incomplete development of the brain.

Atelogliossia (*at-el-o-glos'-e-ah*) [*ἀτελής*, imperfect; *γλῶσσα*, tongue]. Congenital defect in the tongue.

Atelognathia (*at-el-og-na'-the-ah*) [*ἀτελής*, imperfect; *γνάθος*, jaw]. Imperfect development of a jaw, especially of the lower jaw.

Atelomyelia (*at-el-o-mi-e'-le-ah*) [*ἀτελής*, imperfect; *μυελός*, marrow]. Congenital defect of the spinal cord.

Atelopodia (*at-el-o-po'-de-ah*) [*ἀτελής*, imperfect; *πούς*, foot]. Defective development of the foot.

Ateloprosopia (*at-el-o-pro-so'-pe-ah*) [*ἀτελής*, incomplete; *πρόσωπον*, face]. Incomplete facial development.

Atelorrhachidia (*at-el-o-rak-id'-e-ah*) [*ἀτελής*, imperfect; *ράχης*, spine]. Imperfect development of the spinal column, as in spina bifida.

Atelostomia (*at-el-o-sto'-me-ah*) [*ἀτελής*, incomplete; *στόμα*, mouth]. Incomplete development of the mouth.

Atheroma (*ath-er-o'-mah*) [*ἀθήρη*, gruel; *δμα*, tumor]. 1. A sebaceous cyst containing a cheesy material. 2. The fatty degeneration of the walls of the arteries in arteriosclerosis; by common usage, the word is also applied to the whole process of arteriosclerosis. Arterial atheroma is also termed atherosclerosis.

Atheromasia (*ath-er-o-ma'-ze-ah*) [*ἀθήρη*, gruel; *δμα*, tumor]. Atheromatous degeneration; the condition of atheroma.

Atheromatous (*ath-er-o'-mat-us*) [*ἀθήρη*, gruel; *δμα*, tumor]. Characterized by or affected with atheroma. **A. Abscess**, is a collection of soft matter beneath the intima resulting from chronic arteritis, while an **A. Ulcer** is formed by the abscess breaking through the intima.

Atherosclerosis (*ath-er-o'-sis*) [*ἀθήρη*, gruel]. A synonym of atheroma (second definition).

Athetoid (*ath'-et-oid*) [*ἀθητος*, unfixed]. Pertaining to or affected with athetosis. **A. Spasm**, a spasm in which the affected member performs athetoid movements.

Athetosis (*ath-et-o'-sis*) [*ἀθητος*, unfixed; changeable]. A condition most frequently occurring in children, and characterized by continual slow change of position of the fingers and toes. It is usually due to a lesion of the brain. It is also called "post-hemiplegic chorea," from its occurrence after hemiplegia. **A., Double Congenital**. See *Paraplegia, Infantile Spasmodic*.

Athlete's Heart (*ath'-let*) [*ἀθλῆν*, to contend with]. A slight incompetency of the aortic valves, a condition sometimes found in athletes.

Athrepsia (*ah-threps'-e-ah*) [*á* priv.; *τρέφειν*, to nourish]. Malnutrition.

Athymia (*ah-thí'-me-ah*) [*á* priv.; *θύμος*, mind]. 1. Despondency. 2. Loss of consciousness. 3. Insanity.

Atlas (*at'-las*) [*ἀτλας*, able]. The first of the cervical vertebrae. It articulates with the occipital bone of the skull and with the axis.

Atlodymus (*at-lod'-im-us*) [*ἀτλας*, able; *διδυμος*, double]. A monosomic dual monstrosity with two heads and a single body.

Atmograph (*at'-mo-graf*) [*ἀτμός*, breath; *γράφειν*, to record]. A form of self-registering respirometer.

Atmometer, or Atmidometer (*at-mom'-et-er*, or *at-mid-om'-et-er*) [*ἀτμός*, vapor; *μέτρον*, a measure]. An instrument for measuring the amount of water exhaled by evaporation from a given surface in a given time, in order to determine the humidity of the atmosphere.

Atmosphere (*at'-mos-fer*) [*ἀτμός*, vapor; *σφαῖρα*, a sphere]. 1. The air; the mixture of gases, surrounding the earth, to the height of about 200 miles. 2. The pressure exerted by the earth's atmosphere at the level of the sea; it is about 15 pounds to the square inch, or 1 kilogram to the square centimeter.

Atmospheric (*at-mos-fer'-ik*) [*ἀτμός*, vapor; *σφαῖρα*, a sphere]. Pertaining to the atmosphere. **A. Moisture**, the vapor of water mingled with the atmosphere. It varies in quantity according to the temperature. **A.**

Tension, the pressure of the air per square inch on the surface of a body. Normally, at the sea level, it is about 15 lbs. per square inch, or equal to that of a column of mercury about 30 in. in height. It decreases about $\frac{1}{8}$ in. or $\frac{1}{8}$ lb. per square inch for every 90 feet of altitude. Above 10,000 feet, the rarity of the atmosphere is usually noticeable in quickened breathing and pulse-rate.

Atocia (*at-ot'-se-ah*) [*ἀτοκος*, barren]. Sterility of the female.

Atom (*at'-um*) [*á* priv.; *τέμνειν*, to cut]. The ultimate unit of an element; that part of a substance incapable of further division, or the smallest part capable of entering into the formation of a chemie compound, or uniting with another to form a molecule—which last is the smallest quantity of a substance that can exist free or uncombined.

Atomic (*at-om'-ik*) [*á* priv.; *τέμνειν*, to cut]. Pertaining to atoms. **A. Heat**, the specific heat of an atom of a chemie element as compared with that of an atom of hydrogen. **A.**

Theory, the theory of Dalton, that all matter is composed of atoms, the weight of each atom differing for the different elements. **A.**

Valence, Equivalence, or the Atomicity of an element, is the saturating power of its atom as compared with that of hydrogen. **A.**

Weight, the weight of an atom of an element

as compared with the weight of an atom of hydrogen.

Atomicity (*at-om-is'-i-te*) [*á* priv.; *τέμνειν*, to cut]. Chemie valence; quantitative.

Atomization (*at-om-i-z-a'-tion*) [*á* priv.; *τέμνειν*, to cut]. The mechanic process of breaking up a liquid into fine spray.

Atomizer (*at'-om-i-zer*) [*á* priv.; *τέμνειν*, to cut]. An instrument for transforming a liquid into a spray.

Atonic (*at-on'-ik*) [*ἀτονία*, want of tone]. Relating to or characterized by atony.

Atony (*at'-o-ne*) [*á* priv.; *τόνος*, tone]. Want of power, especially of muscular power.

Atrabiliary (*at-rah-bil'-e-a-re*) [*ater*, black; *bilis*, bile]. Pertaining to black bile. **A.** Capsules, an old name for the suprarenal capsules.

Atresia (*at-re'-ze-ah*) [*á* priv.; *τρέφειν*, perforation]. Imperforation of a normal opening or canal, as of the anus, vagina, meatus auditorius, pupil, etc.

Atresic (*at-re'-zik*) [*á* priv.; *τρέφειν*, perforation]. Characterized by atresia.

Atrichia (*ah-trik'-e-ah*) [*á* priv.; *τριξ*, hair]. Absence of the hair.

Atrichosis (*ah-trik-ot'-is*) [*á*, priv.; *τριξ*, hair]. A condition characterized by absence of hair.

Atrioventricular (*a-tre-o-ven-trik'-u-lar*) [*atrium*, hall; *ventriculus*, ventricle]. Relating both to the atrium (or auricle) and to the ventricle of the heart.

Atrium (*at'-tre-um*) [*atrium*, the fore-court or hall]. 1. That part of the auricle of the heart into which the venous blood is poured. 2. The part of the tympanic cavity of the ear below the head of the malleus. **A.** Infection, the point of entrance of the bacteria in an infectious disease.

Atropa (*at'-ro-pah*) [*Ἀτροπις*, "she who turns not," undeviating; one of the three Fates, who cut the thread of life,—in allusion to the poisonous effects of the plant]. A genus of the natural order *Solanaceae*. **A.** belladonna, the deadly nightshade, from which *atropin* is obtained. See *Belladonna*.

Atrophia (*at-ro'-fe-ah*) [*á* priv.; *τροφή*, nourishment]. Atrophy. Diminution in size of a part from degeneration or shrinkage of its component cells. **A. cordis**, atrophy of the heart, following senile changes, or occurring in cachectic conditions, or as a result of pressure exerted by mediastinal tumors, etc. **A. cutis**, Atrophoderma; a term applied to various changes in the skin characterized by the diminution or disappearance of certain of its elements, especially seen in advanced age. The skin becomes thin, loose, wrinkled, and discolored. **A. cutis senilis**, Senile Atrophoderma; an atrophy of the skin usually associated with general signs of senile

degeneration. **A. infantum.** Synonym of *Tabes mesenterica*. **A. mesenterica.** Synonym of *Tabes mesenterica*. **A. pilorum propria,** atrophy of the hair, either symptomatic or idiopathic in origin. **A. unguis,** atrophy of the nails.

Atrophic (*at-rof'fik*) [*â* priv.; τροφή, nourishment]. Pertaining to, or affected with atrophy.

Atrophoderma (*at-ro-fod'er-mah*) [*â* priv.; τροφή, nourishment; δέρμα, the skin]. Atrophy of the skin.

Atrophy (*at-ro-fē*) [*ârpoûia*, want of nourishment]. Diminution in the size of a tissue, organ, or part, the result of degeneration of the cells or a decrease in the size of the cells. **A., Acute Yellow.** See *Icterus gravis*. **A., Brown,** a form of atrophy in which the normal pigment of the organ is retained, and in which there is also frequently the addition of new pigment. It occurs most frequently in the heart, muscles, and liver, and is caused by chronic congestion; also called pigmented atrophy. **A., Correlated,** an atrophy of certain portions of the body following the removal or destruction of other portions. Thus amputation of an arm will be followed by an atrophy of the scapula; of a leg, by atrophy of the corresponding os innominatum. **A., Cruveilhier's.** See *Diseases, Table of*. **A., Degenerative,** that due to degeneration of the cells. **A., Gray,** a degenerative change in the optic disc in which the latter assumes a grayish color. **A., Idiopathic Muscular,** muscular wasting, beginning in various groups of muscles, usually progressive in character, and dependent on primary changes in the muscles themselves. There is a strong hereditary predisposition to the disease. **A., Muscular,** atrophy affecting muscles; it may be hereditary or acquired, idiopathic, myelopathic, myopathic, neuropathic, primary, secondary, simple, or progressive. **A., Pigmentary, A., Pigmented,** a form of atrophy so called from a deposit of pigment (yellow or yellowish-brown) in the atrophied cells. **A., Progressive Muscular;** Chronic Anterior Poliomyelitis; Wasting Palsy. A chronic disease characterized by progressive wasting of individual muscles or physiologic groups of muscles, and by an associated and proportional amount of paralysis. It is due to a degeneration and atrophy of the multipolar cells in the anterior gray horns of the cord, with consecutive degeneration of the anterior nerve-roots and muscles. The right hand is usually the part first attacked, and takes on a peculiar claw-like form (*main en griffe*). The disease is most frequent in males of adult life, and follows excessive muscular exertion. **A., Progressive Unilateral Facial,** a disease characterized by progressive wasting of

the skin, connective tissue, fat, bone, and more rarely the muscles of one side of the face. It is most common in females; its course is slow and generally progressive. **A., Red,** a form of atrophy due to chronic congestion, as seen in the liver in mitral and tricuspid valvular lesions. **A., Senile,** the physiologic atrophy of advanced life. It affects the lungs, the sexual and other organs. **A., Serous,** atrophy associated with an infiltration of fluid into the atrophic tissues. **A., Simple,** that due to a decrease in the size of individual cells. **A., Trophoneurotic,** that dependent upon abnormality of the nervous supply of an organ or tissue, best illustrated in muscular atrophy from disease of the anterior horns of the spinal cord.

Atropina, or Atropin (*at-ro-pi'-nah*, or *at'-ro-pin*) [*Ἄτροπος*, one of the Fates who cut the thread of life; *gen., Atropine*], $C_{17}H_{23}NO_3$. A crystalline alkaloid derived from *Atropa belladonna*. The sulphate is a white powder of bitter taste and neutral reaction, and is soluble in water. Atropin is a mydriatic, antispasmodic, and anodyne; in small doses a cardiac, respiratory, and spinal stimulant, in large doses a paralyzant of the cardiac and respiratory centers, the spinal cord, motor nerves, and involuntary and voluntary muscles. It lessens all the secretions except the urine. In full doses it produces dryness of the throat, flushing of the face, dilatation of the pupils, a rise of temperature and sometimes an erythematous rash. It is extensively used in ophthalmic practice to dilate the pupil, to paralyze accommodation, and also in various corneal, iritic, and other ocular diseases. Its therapeutic use in general medicine is also manifold; *e. g.*, in inflammatory affections and the pain of cerebral and spinal hyperemia, atonic constipation, cardiac failure, hypersecretions, especially of the sweat, to relieve local spasms, as in intestinal and biliary colic, in asthma, whooping cough, etc., and as a physiologic antagonist in opium poisoning. Dose of atropin sulphate gr. $\frac{1}{100}$ – $\frac{1}{50}$ (0.0005–0.001).

Atropinize (*at'-ro-pin-iz*) [*Ἄτροπος*]. To bring under the influence of, or to treat with atropin.

Attar (*at'-ar*) [Arab. *itr*, perfume]. A general name for any of the volatile oils. **A. of Rose, Oil of Rose.** The volatile oil distilled from the fresh flowers of the Damascene rose. It comes mainly from eastern Roumelia and is generally adulterated with other volatile oils. It is used as a perfume.

Attendant (*at-ten'-dant*) [*attendere*, to attend]. A non-professional attaché of an asylum or hospital, especially of an insane asylum.

Attenuant (*at-ten'-u-ant*) [*attenuare*, to

make thin]. A medicine or agent increasing the fluidity or thinness of the blood or other secretion.

Attenuating (*at-ten'-u-a-ting*) [*attenuare*, to make thin]. Making thin.

Attenuation (*at-ten-u-a'-shun*) [*attenuare*, to make thin]. The act of making thin; a thinning, narrowing, or reduction of the strength or size of a substance, especially the weakening of the pathogenic virulence of microorganisms by successive cultivation, by exposure to light, air, heat, or other agency, or by passing through certain animals, so that they may be used as a vaccine to confer immunity from future attacks of the disease. **A.**, Sanderson's Method of, the passing of the virus through the system of another animal (*e. g.*, the guinea-pig, in anthrax), so that it becomes modified in virulency.

Attic (*at'-ik*) [*ἄττικός*, Attic]. Part of the tympanic cavity situated above the atrium. **A. Disease**, chronic suppurative inflammation of the attic of the tympanum.

Attitude (*at'-e-tūd'*) [*aptitudo*, aptitude]. See *Posture*. **A.**, Crucifixion, in hystero-epilepsy, a rigid state of the body, the arms stretched out at right angles. **A.**, Frozen, a peculiar stiffness of the gait characteristic of disease of the spinal cord, especially of amyotrophic lateral sclerosis. **A.**, Passionate, the assumption of a dramatic or theatrical expression, a position assumed by some hysteric patients.

Attollens (*at-ol'-enz*) [*attollere*, to rise up]. Raising. **A. auris**, a muscle raising the external ear.

Attraction (*at-rak'-shun*) [*attrahere*, to draw to]. The tendency of one particle of matter to approach another. Affinity. As existing between masses, it is termed *gravitation*, while *molecular attraction* or *cohesion* expresses the force aggregating molecules. **A.**, Chemic, the attraction of affinity, relates to the attraction of atoms of one element to those of others, resulting in chemic compounds. **A.**, Capillary, the force that causes liquids to rise in fine tubes or between two closely approximated surfaces, or on the sides of the containing vessel. **A.**, Electric, the tendency of bodies toward each other when charged with opposite electricities. **A.**, Magnetic, the influence of a magnet upon certain metallic substances, chiefly iron.

Attrahens (*at'-rah-enz*) [*attrahens*, drawing]. Drawing forward, as **Attrahens aurem**, a muscle drawing the ear forward and upward.

Attrition (*at-rish'-un*) [*atterere*, to rub against]. Rubbing or friction.

Atypic (*ah-tip'-ik*) [*ἀ priv.; τύπος*, a type]. Irregular; not conformable to the type. **A. Fever**, an intermittent fever with irregularity of the paroxysm.

Atypical (*ah-tip'-ik-al*). See *Atypic*.

Audiometer (*aw-de-om'-et-er*) [*audire*, to hear; *μέτρον*, a measure]. An instrument for measuring the acuteness of hearing.

Audiphone (*aw'-dif-ōn*) [*audire*, to hear; *φωνή*, a sound]. An instrument for improving the power of hearing by conveying sounds through the bones of the head to the labyrinth.

Audition (*aw-dish'-un*) [*audire*, to hear]. The act of hearing. **A. colorée**, color-hearing, a peculiar association between the auditory and optic nerves, by which a certain sound or musical note will give rise to a subjective sensation of color, the same note in the same person being always associated with the same color.

Auditory (*aw'-dit-o-re*) [*audire*, to hear]. Pertaining to the act or the organs of hearing. **A. After-Sensations**, the sensations of sounds continuing or occurring after the cessation of the stimulus. **A. Amnesia**. See *Mind-deafness*. **A. Area**, the cerebral center for hearing, probably located in the temporo-sphenoidal lobe. **A. Aura**, an auditory sensation preceding an attack of epilepsy. **A. Center**, same as *A. Area*. **A. Hairs**, the processes of the crista acustica. **A. Meatus** (external and internal), the external and internal canals or openings of the ear. **A. Nerve**, the eighth cranial nerve, supplying the internal ear; formerly the *portio mollis* of the seventh pair of cranial nerves. **A. Ossicles**, the chain of small bones of the middle ear. **A. Pit**, the depression in the epiblast on both sides of the embryonic after-brain, destined to form the labyrinth of the ear. **A. Vertigo**, dizziness due to pathologic conditions of the ear. See *Ménière's Disease*.

Auerbach's Plexus. A nerve-plexus found between the circular and longitudinal muscular coats of the stomach and intestines, and consisting of a network of pale nerve-fibers, at the nodal points of which minute ganglia exist.

Augnathus (*awg-na'-thus*) [*αὐ*, besides; *γνάθος*, the jaw]. A monster with two lower jaws.

Aula (*aw'-lah*) [*αὐλή*, a hall or open court]. The common mesal cavity of the cerebrum, it being also the anterior portion of the third ventricle.

Aura (*aw'-rah*) [*ἀτφα*, a breath]. A breath of wind; a soft vapor. The phenomena preceding an attack of epilepsy. It may be motor, sensory, vasomotor, secretory, or psychic. It is also applied to the symptom preceding an attack of any disease or paroxysm, as the *aura hysterica*, *aura vertiginosa*, etc.

Aural (*aw'-ral*) [*auris*, the ear]. Relating

to the ear or to hearing. **A. Vertigo.** See *Mnière's Disease*.

Auramin (*aw'-ram-in*) [*aurum*, gold; *amin*]. Yellow Pyoktanin, a yellow anilin color used to some extent as an antiseptic. See *Pyoktanin*.

Aurantia (*aw-ran'-she-ah*) [*aurantium*, an orange]. An orange coal-tar dye; an ammonium salt of hexanitro-diphenylamin.

Aurantia (*aw-ran'-tin*). See *Heptane*.

Aurantium (*aw-ran'-she-um*) [L.: *gen.*, *Aurantii*]. Orange. The fruit of *Citrus vulgaris* and *C. aurantium*. Both the flowers and the rind of the fruit are employed in medicine. The volatile oil from the rind is aromatic and a mild tonic, but is used mainly as a flavor. **A. amari**, Ext. Fld., bitter orange peel, alcohol, and water. It is used as a flavor. Dose 3 ss-j (2.0-4.0). **A. amari**, Tinct., bitter orange peel 20, dilute alcohol, q. s. ad 100. Dose 3 j-ij (4.0-8.0). **A. corticis**, Oleum, the volatile oil expressed from the rind of the orange. Dose gtt. j-v (0.065-0.32). **A. dulcis**, Tinct., sweet orange peel 20, dilute alcohol, q. s. ad 100. Dose 3 j-ij (4.0-8.0). **A.**, Elixir, oil of orange 1, sugar 100, alcohol and water, q. s. ad 300. **A. florum**, Aqua, stronger orange water and distilled water, of each, one volume. **A. florum fortior**, Aqua. Water saturated with the volatile oil of fresh orange flowers. **A. florum**, Oleum, "oil of neroli," a volatile oil distilled from fresh orange flowers. Dose gtt. j-v (0.065-0.32). **A. florum**, Syr., sugar 85, orange flower water sufficient to make 100 parts. A common flavoring agent. **A.**, Infus. (B. P.). Dose 3 j-ij (32.0-64.0). **A.**, Infus. Comp. (B. P.). Dose 3 j-ij (32.0-64.0). **A. Spt.**, oil of orange 5, deodorized alcohol 95. Dose according to quantity of alcohol desired. **A.**, Spt. Comp., oil of orange peel 20, oil of lemon 5, oil of coriander 2, oil of anise 5, deodorized alcohol, sufficient to make 100 parts. **A.**, Syr., sweet orange peel 5, calcium phosphate precip. 1, sugar 70, alcohol and water, each a sufficient quantity to make 100 parts. **A.**, Tinct. (B. P.). Dose 3 j-ij (4.0-8.0). **A. recentis**, Tinct. (B. P.), tincture of fresh orange peel. Dose 3 j-ij (4.0-8.0). **A.**, Vinum (B. P.), contains 12 per cent. of alcohol.

Auric (*aw-rik*) [*aurum*, gold]. Pertaining to aurum or gold. **A. Acid.** See *Acid*.

Auricle (*aw'-rik-l'*) [*auricula*, the ear]. 1. The expanded portion or pinna of the ear.

2. One of the upper chambers of the heart receiving the blood from the lungs (left A.) or from the general circulation (right A.).

Auricular (*aw-rik'-u-lar*) [*auricula*, the ear]. 1. Relating to the auricle of the ear.

2. Pertaining to the auricles of the

heart, as *A. appendix*. 3. Relating to the auricular nerve, arteries, veins, etc. **A. Finger**, the little finger. **A. Point**, the central point of the external auricular meatus.

Auricularis (*aw-rik-u-la'-ris*) [*auricula*, the ear]. Auricular. **A. magnus**, a branch of the cervical plexus of nerves.

Auriculotemporal (*aw-rik'-u-lo-tem'-po-ral*) [*auricula*, the ear; *tempus*, the temple].

Relating to the auricle and to the temporal region. **A. Nerve**, a branch of the inferior maxillary, supplying superficial parts about the auricle and the temple.

Auriculoventricular (*aw-rik'-u-lo-ventrik'-u-lar*) [*auricula*, the ear; *ventriculus*, the ventricle].

Relating to an auricle and a ventricle of the heart. **A. Opening**, the opening between the auricles and the ventricles of the heart.

Auripuncture (*aw'-re-punkt-chūr*) [*auris*, ear; *punctura*, puncture]. Puncture of the membrana tympani.

Auris (*aw'-ris*) [L.]. The ear.

Auriscopes (*aw'-ris-kōp*) [*auris*, the ear; *σκοπεῖν*, to examine]. An instrument for examining the ear, and especially the Eustachian passage; an otoscope.

Aurist (*aw'-rist*) [*auris*, the ear]. A specialist in diseases of the ear.

Aurum (*aw'-rum*) [L.: *gen.*, *Auri*]. Gold. Au = 196.7; quantivalence, III. A brilliant yellow metal, having a specific gravity of 19.3. It is soluble in a mixture of nitric and hydrochloric acids. **A. bromidum**, AuBr₃, is used in epilepsy and migraine. Dose gr. 1/10-1/8 (0.003-0.01). **A. chloridum**. Dose gr. 1/10-1/8 (0.0012-0.002). Gold chlorid is also used as a stain for nerve-tissue. **A. et sodii chloridum**, the double chlorid of gold and sodium. Dose gr. 1/10-1/8 (0.002-0.006). It is used as an alternative in chronic inflammations, diabetes, in the treatment of the alcohol habit, etc.

Auscult, or **Auscultate** (*aws-kult'*, *aws'-kul-tat*) [*auscultare*, to listen to]. To perform or practise auscultation; to examine by auscultation.

Auscultation (*aws-kul-tat'-shun*) [*auscultare*, to listen to]. A method of investigation of the functions and condition of the respiratory, circulatory, digestive, and other organs by the sounds they themselves give out, or that are elicited by percussion. It is called *immediate*, when the ear is directly applied to the part, and *mediate*, if practised by the aid of the stethoscope. *Obstetric auscultation* is practised in pregnancy to detect or study the fetal heart-sounds or the placental murmur.

A.-tube, in otology, an instrument for listening to the forced passage of air through the ear of another.

Auscultatory (*aws-kul'-tat-o-re*) [*auscultare*,

to listen to]. Relating to auscultation. **A.** Percussion, the practice of listening with the stethoscope to the sounds produced by percussing a part.

Autechoscope (*aw-tek'-o-skop*) [*αὐτός*, self; *ἦχος*, sound; *σκοπεῖν*, to inspect]. A device for enabling a person to listen to sounds produced within his own body.

Autecious, or Autécious (*aw-te'-shus*) [*αὐτός*, self; *οἶκος*, dwelling]. Applied to parasitic fungi that pass through all the stages of their existence in the same host.

Autochthon (*aw-tok'-then*) [*αὐτόχθων*, sprung from the land]. An aboriginal inhabitant.

Autochthonous (*aw-to-k'-then-us*) [*αὐτόχθων*, sprung from the land]. Aboriginal; formed (as *e. g.*, a clot) in the place where it is found.

Autocinesis (*aw-to-sin-el'-sis*) [*αὐτός*, self; *κίνησις*, motion]. Motion that is voluntary.

Autocinetic (*aw-to-sin-el'-ik*) [*αὐτός*, self; *κίνησις*, motion]. Possessed of the power of spontaneous motion.

Autoclave (*aw-to-klaw*) [*αὐτός*, self; *clavis*, a key]. 1. Self-fastening; closing itself. 2. An apparatus for sterilizing objects by steam-heat at high pressure.

Autodidact (*aw'-to-di-dakt*) [*αὐτός*, self; *διδασκός*, taught]. One who is self-taught in his profession.

Autodigestion (*aw-to-di-jes'-chun*) [*αὐτός*, self; *digere*, to digest]. Digestion of an organ by its own secretion.

Autogenesis (*aw-to-jen'-es-is*) [*αὐτός*, self; *γένεσις*, production]. Spontaneous generation; self-production.

Autogenetic (*aw-to-jen-el'-ik*) [*αὐτός*, self; *γένεσις*, production]. Produced within the organism.

Autogenous (*aw-toj'-en-us*) [*αὐτός*, self; *γένεσις*, production]. Pertaining to diseases or conditions self-produced within the body and not derived from external sources; applied to poisons generated in the body by its inherent processes. **A.** Hemorrhage, hemorrhage due to causes residing within the body; not traumatic.

Autohypnotism (*aw-to-hip'-not-izm*) [*αὐτός*, self; *ἵπνος*, sleep]. Mental stupor induced by dwelling intensely upon some all-absorbing thought.

Autoinfection (*aw-to-in-fek'-shun*) [*αὐτός*, self; *infection*]. Infection by virus originating within the body or transferred from one part of the body to another.

Autoinoculable (*aw-to-in-ok'-u-la-bl*) [*αὐτός*, self; *inoculare*, to implant]. Capable of being inoculated upon the person already infected. Chancroid is autoinoculable.

Autoinoculation (*aw-to-in-ok-u-la'-shun*) [*αὐτός*, self; *inoculare*, to implant]. Inoculation in one part of the body by virus present in another part; self-inoculation.

Autointoxication (*aw-to in-toks-ik-a'-shun*) [*αὐτός*, self; *ροξικόν*, a poison]. Poisoning by faulty metabolic products elaborated within the body; autoinfection.

Autokinesis (*aw-to-kin-el'-sis*) [*αὐτός*, self; *κίνησις*, movement]. Voluntary movement.

Autokinetic (*aw-to-kin-el'-ik*) [*αὐτός*, self; *κίνησις*, movement]. Pertaining to, or of the nature of, autokinesis.

Automatic (*aw-to-mat'-ik*) [*αὐτοματίζω*, to act spontaneously]. Performed without the influence of the will.

Automatism (*aw-tom'-at-izm*) [*αὐτοματίζω*, to act spontaneously]. The performance of acts without apparent volition, as seen in certain somnambulists and in some hysteric and epileptic patients. **A., Epileptic.** See *Automatism*.

Automaton (*aw-tom'-at-on*) [*αὐτόματος*, spontaneous]. One who acts in an involuntary or mechanic manner.

Autonomy (*aw-ton'-o-me*) [*αὐτός*, self; *νόμος*, law]. Independence.

Autonomous (*aw-ton-om'-us*) [*αὐτός*, self; *νόμος*, law]. Self-ruled; independent.

Autopepsia (*aw-to-pep'-se-ah*) [*αὐτός*, self; *πέπτειν*, to digest]. Autodigestion.

Autophagia (*aw-to-pha'-je-ah*) [*αὐτός*, self; *φαγεῖν*, to eat]. Self-consumption; emaciation.

Autophagy (*aw-toff'-a-je*). See *Autophagia*.

Autophobia (*aw-to-fo'-be-ah*) [*αὐτός*, self; *φόβος*, fear]. A morbid dread of one's self, or of solitude.

Autophonous (*aw-toff'-on-us*) [*αὐτός*, self; *φωνή*, voice]. Having the character of autophony.

Autophony (*aw-toff'-o-ne*) [*αὐτός*, self; *φωνή*, voice]. 1. The auscultation of the physician's own voice through the patient's chest. 2. The condition in which one's own voice appears changed. It may be due to chronic inflammation of the ear or to other causes.

Autophthalmoscopy (*aw-toff-thal'-mas'-ko-pe*) [*αὐτός*, self; *ὀφθαλμός*, the eye; *σκοπεῖν*, to see]. Ophthalmoscopy performed upon one's own eye.

Autoplasty (*aw-to-plas-te*) [*αὐτός*, self; *πλάσσειν*, to form]. A method of repairing the effects of a wound or lesion involving loss of tissue by grafting or implanting fresh parts taken from other portions of the patient's body.

Autopsy (*aw-top-se*) [*αὐτός*, self; *ὄψις*, a seeing]. The post-mortem examination.

Autoscope (*aw-to-skop*) [*αὐτός*, self; *σκοπεῖν*, to see]. An instrument, *e. g.*, the ophthalmoscope, arranged for the examination of an organ by oneself.

Autoscopy (*aw-to'-ko-pe*) [*αὐτός*, self; *σκοπεῖν*, to see]. The examination of one's own organs by means of an autoscope.

Autosite (*aw'-to-sit'*) [*αὐτός*, self; *σῖτος*, food].

1. A monster capable of an independent existence after birth. 2. That member of a double fetal monstrosity that nourishes itself by its own organs and also the other member, which is called the parasite.

Autositic (*aw-to-sit'-ik*) [*αὐτός*, self; *σῖτος*, food]. Of the nature of an autosite.

Autostethoscope (*aw-to-steth'-o-skōp*) [*αὐτός*, self; *στήθος*, the breast; *σκοπεῖν*, to examine]. A stethoscope so arranged that by it one may listen to his own chest-sounds.

Autosuggestion (*aw-to-sug-jes'-chun*) [*αὐτός*, self; *suggestio*, an intimation]. A term suggested by Page for a peculiar mental condition, often developing after accidents, especially railway accidents; it is intimately associated with the hypnotic state. In both of these conditions the mental spontaneity, the will, or the judgment, is more or less suppressed or obscured, and suggestions become easy. Thus the slightest traumatic action directed to any member may become the occasion of a paralysis, of a contracture, or of an arthralgia. It is also called *traumatic suggestion*.

Autotherapy (*aw-to-ther'-a-pe*) [*αὐτός*, self; *θεραπεία*, a waiting on]. The spontaneous or self-cure of a disease.

Autotoxemia (*aw-to-toks'-e'-me-ah*) [*αὐτός*, self; *τοξικόν*, a poison; *αἷμα*, blood]. Toxemia from poisons derived from the organism itself.

Autotoxin (*aw-to-toks'-in*) [*αὐτός*, self; *τοξικόν*, a poison]. Any poisonous product of tissue-metamorphosis.

Autotransfusion (*aw-to-trans-fu'-shun*) [*αὐτός*, self; *transfusio*, a pouring out or forth]. The transfer of the blood to the brain and other central organs by elevating the hips and legs, and by the use of elastic bandages compressing the limbs.

Autovaccination (*aw-to-vaks'-in-a'-shun*) [*αὐτός*, self; *vaccinare*, to vaccinate]. The reinsertion of fresh-vaccine lymph upon the same person from whom it is taken.

Autumn Catarrh. Synonym of Hay-fever, since it occurs in August and the fall of the year.

Autumnal (*aw-tum'-nal*) [*autumnus*, autumn]. Pertaining to the fall of the year.

A. Fever. Synonym of Typhoid Fever.

Auxiliary (*awks-il'-e-a-re*) [*auxilium*, help]. Aiding.

Auxillium (*awks-il'-e-um*) [L., help]. A wheeled vehicle or ambulance with couch and mattresses, for use in the service of field military hospitals.

Ava Kava (*ah'-vah kah'-vah*). See *Kava-Kava*.

Avalanche Theory. Pflüger's theory that nerve-energy gathers intensity as it passes toward the muscles.

Avascular (*ah-vas'-cul-ar*) [*à* priv.; *vasa*, vessel]. Without blood; not possessing blood-vessels.

Avena (*av-e'-nah*) [L.]. A genus of plants. Oat. **A. farina**, oatmeal. **A. sativa**, the embryo of the seed of the common oat plant. It contains starch, gluten, a ferment called diastase, and a small amount of alkaline phosphates, and is a nutritious food. Dose of the concentrated tinct. or fld. ext. ℥x-ʒij (0.65-8.0). The pericarp contains an alkaloid possessed of slight narcotic powers. Unof.

Avenin (*av-e'-nin*) [*avena*, the oat]. 1. A precipitate made from a tincture of *avena sativa*, or the oat. It is a nerve-stimulant and tonic. Unof. 2. A nitrogenous principle obtained from the oat, and nearly identical with legumin; the gluten-casein of oats.

Avogadro, Law of. See *Law*.

Avoirdupois Weight (*av-or-du-pois'-wät*) [Fr., *avoirduois*, to have some weight]. See *Weights and Measures*.

Avulsion (*av-ul'-shun*) [*avulsio*; *avellere*, to tear away]. A tearing or wrenching away of a part, as a polypus, a limb, etc.

Axial (*aks'-e-äl*) [*axis*, axis]. Pertaining to or situated in an axis. **A. Current**, the column of red corpuscles which, by reason of the weight of the cells, occupies the center or axis of the blood-stream. **A. Hypermetropia**. See *Hypermetropia*. **A. Stream**. See *A. Current*.

Axilemma (*aks-il-em'-ah*) [*axis*, axis; *λέμμα*, husk, skin]. An elastic sheath composed of neurokeratin, enclosing the axis-cylinder of medullated nerve-fibers.

Axilla (*aks-il'-ah*) [L.: *pl.*, *Axillæ*]. The armpit.

Axillary (*aks'-il-a-re*) [*axilla*, armpit]. Pertaining to the axilla. **A. Artery**, the continuation of the subclavian artery, extending from the lower border of the first rib to the insertion of the pectoralis major muscle, where it becomes the brachial. See *Arteries*, *Table of*. **A. Glands**, the lymphatic glands in the axilla. **A. Plexus**, the brachial plexus, formed by the last three cervical and the first dorsal nerves. **A. Space**, the irregular conical space of the axilla. **A. Vein**, a continuation of the brachial vein, corresponding with the artery and terminating in the subclavian vein.

Axis (*aks'-is*) [L. for axletree]. 1. An imaginary line passing through the center of a body. 2. The second cervical vertebra. 3. A short artery which breaks up into several branches, e. g. A., Thyroid; A., Celiac. See *Arteries*, *Table of*. **A.**, Basicranial, in craniometry, a line drawn from the basion to the middle of the anterior border of the cerebral surface of the sphenoid

bone. **A.**, Basifacial, in craniometry, a line drawn from the anterior border of the cerebral surface of the sphenoid to the alveolar point. **A.**, Binauricular, in craniometry, the imaginary line joining the two auricular points. **A.**, Cerebrospinal, the central nervous system. **A.**, Frontal (of the eye), an imaginary line running through the eyeball from right to left, and corresponding with the movements of elevation and depression of the eyeball. **A.**, Optic. 1. The line from the center of the cornea to the macula lutea. 2. An imaginary line passing from the center of the eye-piece of a microscope through the body, objective, stage, and sub-stage, to the mirror. **A.**, Sagittal (of the eye), an imaginary line running through the eyeball from before backward, and coinciding with the line of vision. **A.**, Visual, the line from the object through the nodal point to the macula.

Axis-cylinder (*aks-is-sil'-in-der*) [*axis; cylinder*]. The conducting or essential part of a nerve. It is also called the axis-cylinder of Purkinje. **A.-cylinder Process**, that one of the protoplasmic processes of a nerve-cell which becomes an axis-cylinder.

Axis-traction (*aks-is-trak'-shun*) [*axis; trahere, to draw*]. Traction on the fetus in the axis of the pelvis. **A.-traction Forceps**, a forceps for performing axis-traction.

Axungia (*aks-un'-je-ah*) [*L.: gen., Axungia*]. Fat; lard; adeps.

Azalein (*az-ol'-le-in*). Same as *Rosanilin*.

Azedarach (*az-ed'-ar-ak*) [*Pers. âzâd, free; dirakht, a tree*]. Pride of China, the bark of *Melia azedarach*, an Asiatic tree naturalized in the southern U. S. It occurs in curved pieces or quills, having a sweetish taste. A decoction, $\frac{3}{4}$ ss to Oj, is used as an anthelmintic against the round-worm. Dose $\frac{3}{4}$ ss-j (16.0-32.0). **A.**, Ext. Fld. Dose $\frac{3}{4}$ j (4.0). Unof. **A.**, Tinctura, 1 to 8. Dose $\frac{3}{4}$ ss-ij (2.0-8.0). Unof.

Azerin (*az'-er-in*) [*â priv.; ξηρός, dry*]. A ferment analogous to ptyalin and found in the digestive secretions of *Drosera Nepenthes*, and probably all other insectivorous plants.

Azobenzene (*az-o-ben'-zen*) [*azote, nitrogen*;

benzoin], $C_{12}H_{10}N_2$. A compound formed by the action of sodium-amalgam upon the alcoholic solution of nitro-benzene. It forms orange-red, rhombic crystals, readily soluble in alcohol and ether, but sparingly soluble in water. It melts at 68°, and disils at 293°.

Azo-compounds. In chemistry, compounds intermediate between the nitro-compounds and the amido-compounds, and made from the former by partial reduction, or from the latter by partial oxidation.

Azo-dyes. A well defined group of the coal-tar colors, all containing the diatomic group $-N=N-$, bound on either side to a benzene radical. They may be prepared by reduction of the nitro-compounds in alkaline solutions, or by acting on diazo-compounds with phenols or amines of the aromatic series. The azo-dyes are the amido-derivatives of simple azo-compounds, and are to be distinguished as amidoazo-dyes and oxyazo-dyes.

Azoic (*az-ol'-ik*) [*â priv.; ζωή, life*]. Destitute of living organisms.

Azolitmin (*az-ol'-it'-min*) [*â priv.; ζωή, life; litmus*], $C_7H_7NO_4$. A deep blood-red coloring matter obtained from litmus.

Azoospermia (*ah-zo-o-sper'-me-ah*) [*â priv.; ζωή, life; σπέρμα, seed*]. Absence of, or deficient vitality of, the spermatozoa.

Azote (*az'-ât*) [*â priv.; ζωή, life*]. A synonym of nitrogen. *Azotic acid*, nitric acid.

Azotemia (*az-ol'-te-me-ah*) [*azote, nitrogen; αἷμα, blood*]. The presence of nitrogenous compounds in the blood; uremia.

Azotized (*az'-ot'-ized*) [*azote, nitrogen*]. Nitrogenized; containing nitrogen.

Azoturia (*az-ol'-tu'-re-ah*) [*azote, nitrogen; οὖρον, the urine*]. An increase of the urea and urates in the urine.

Azygos (*az'-ig-os*) [*â priv.; ζυγός, a yoke*]. Applied to parts that are single, not in pairs.

A. uvulæ, a small muscle of the uvula. **A. Vein**, a vein connecting the superior and inferior venæ cavæ.

Azygous (*az'-ig-us*) [*â priv.; ζυγός, a yoke*]. Not paired.

Azymia (*ah-zi'-me-ah*) [*â priv.; ζύμα, a ferment*]. Absence of ferment.

Azzle Teeth (*az'-l-teth*) [*E. dial., assal teeth*]. A name given to the molar teeth.

B

B. In chemic terminology the symbol of *Boron*.

Ba. The chemic symbol of *Barium*.

Babbitt Metal. See *Antimony*.

Bacca (*bak'-ka*) [L.]. *Berry*.

Bacilliform (*bas-il'-if-orm*) [*bacillum*, rod; *forma*, form]. Having the shape or appearance of a bacillus.

Bacilluria (*bas-il'-u'-re-ah*) [*bacillum*, a rod; *urine*, urine]. The presence of bacilli in the urine.

Bacillus (*bas-il'-us*) [dim. of *baculum*; a small staff; *pl.*, *Bacilli*]. 1. A genus of the *Schizomycetes* comprising the rod-shaped forms of bacteria. 2. An individual of the genus *Bacillus*. 3. A medicated rod or bougie. 4. Any rod-like body, or specifically, one of the retinal rods.

Bacony Infiltration (*ba'-kon-e in-fil-tra'-shun*). Same as *Anyloid Degeneration*.

Bacteremia (*bak-ter-e'-me-ah*) [*Bakteriuv*, a little rod; *alua*, blood]. The presence of bacteria in the blood.

Bacterium (*bak-te'-re-um*) [*Bakteriuv*, a little stick; *pl.*, *Bacteria*]. 1. A synonym of *Schizomyces* or microorganism. 2. The word was formerly restricted to a genus of schizomycetous fungi established by Ehrenberg (1838) and Dujardin (1841), characterized by short, linear, inflexible, rod-like forms, without tendency to unite into chains or filaments. Morphologically, bacteria are spheric (cocci); in the form of straight rods (bacilli); or of twisted rods (spirilla). Bacteria are either aerobic requiring free oxygen, or anaerobic, not requiring free oxygen. Again, certain forms appear to possess the ability to flourish in either condition, and are known as facultative aerobic or anaerobic. Bacteria are either motile or nonmotile; they may exist as saprophytes, facultative parasites, strict, obligate, or true parasites. Those that produce pigment are known as chromogenic; those that produce fermentation as zymogenic; those that affect adversely the health of plants or animals as pathogenic. See *Bacteria*, *Table of*, and special terms under appropriate headings.

TABLE OF BACTERIA.*

Actinobacter polymorphus. See *Bacillus butyricus*, Prazmowski. **Actinomyces bovis**, Harz. Syn. *Cladothrix bovis*, *Cladothrix canis*, Rabe (?). Found in

Actinomycosis. Occurs in grains the size of a poppy-seed; composed of radiating threads. Stains with anilin dyes; is not decolorized by Gram's method. Grows on egg-albumin, gelatin, potato, and bouillon. *liqf.*, *pg.*, *chg.* (Sulphur-yellow.) **Anthrax bacillus.** See *Bacillus anthracis*. **Arthrobacterium aceti**, De Bary. See *Bacillus aceti*. **Arthrobacterium chlorinum**, De Bary. See *Bacillus chlorinus*. **Arthrobacterium merismopædioides**, De Bary. See *Bacillus merismopædioides*. **Arthrobacterium pastorianum**, De Bary. See *Bacillus pasteurianus*. **Arthrobacterium viride**, De Bary. See *Bacillus viridis*. **Arthrobacterium zopfii**. See *Bacillus zopfii*. **Ascobacillus citreus**, Unna and Tommasoli. Syn. *Ascococcus citreus*. On the skin, in cases of eczema seborrhœicum. *aer.*, *liqf.*, *mot.*, *chg.* (lemon-yellow), *npg.* **Ascobacterium ulvina**, Van Tieghem. Short rods occurring in liquids containing decaying leguminous seeds. **Ascococcus billrothii**, Cohn. Lobulated masses, producing a viscous fermentation of saccharine fluids and evolving butyric acid in solution of ammonium tartrate. **Ascococcus johnei**, Cohn. See *Micrococcus botryogenus*. **Ascococcus mesenteroides**, Cienkowski. See *Leucomostoc mesenteroides*. **Ascococcus vibrans**, Van Tieghem. Upon water containing *Beggiatoa*; distinguished from *A. billrothii* by the whirling and oscillating of the cells. **Bacilli of Butyric Acid Fermentation.** See *Bacillus butyricus*, Hueppe and Prazmowski. *Bacillus butylicus*, Fitz. **Bacilli of Fermentation of Carbohydrates.** See *Bacillus acidi lactici*, Hueppe. *Bacillus butyricus*, Hueppe and Prazmowski. *Bacillus butylicus*, Fitz. *Bacillus dysodes*, Zopf. *Bacillus polymixa*, Prazmowski. **Bacilli of Guillebeau** (a, b, c), obtained from milk of cows with mastitis; give rise to a peculiar fermentation of cheese. (a) *aer.*, *facanaer.*, *mot.*, *nliqf.*; (b) *aer.*, *facanaer.*, *liqf.*, *mot.*; (c) *aer.*, *nliqf.* **Bacillus aceti**, Kützing, Sommer. Syn., *Arthrobacterium aceti*, De Bary. *Bacillus aceticus*, Flüggé; *Bacterium aceti*, Lanzi. *Micrococcus aceti*; *Mycoderma aceti*, Pasteur. *Ulvina aceti*, Kützing. Found in air and vinegar; common. Large cylindrical cells of varying size. Absorbs oxygen from air and oxidizes alcohol to acetic acid; produces vinegar; may, by further oxidation, convert this into carbon dioxide and water.

* Abbreviations.—*aer.* = aerobic. *anarr.* = anaerobic. *chg.* = chromogenic. *facanaer.* = facultative anaerobic. *liqf.* = liquefactive. *monom.* = monomorphic. *mot.* = motile. *nliqf.* = nonliquefactive. *nmot.* = nonmotile. *npg.* = nonpathogenic. *obl.* = obligate. *osci.* = oscillating. *pg.* = pathogenic. *phos.* = phosphorescent. *pleom.* = pleomorphic. *sap.* = saprophytic.

aer., *nlif.*, *mot.*, *npg.*, *pleom.* **Bacillus aceticus**, Flügge. See *Bacillus aceti*, Kütz-
ing. **Bacillus acidiformans**, Sternberg.
Obtained from the liver of a yellow fever
cadaver. *aer.*, *facanaer.*, *nlif.*, *nmot.*, *pg.*
Bacillus acidi lactici, Hueppe. Syn., *Bac-*
terium lactis, Lister. Found in the air and
in sour milk. *aer.*, *facanaer.*, *nlif.*, *nmot.*,
npg. **Bacillus actinobacter**, Duclaux.
Syn., *Actinobacter polymorphus*. A minute
capsulated bacillus, rendering milk gelatinous
and ropy. *aer.*, *nmot.* See *Bacillus buty-*
ricus, Prazmowski. **Bacillus aerogenes**,
I, II, III, Miller. Syn., *Bacterium aéro-*
genes, Miller. *Helicobacterium aerogenes*,
Miller. Three small bacilli from the alimen-
tary tract of healthy persons. *aer.*, *nlif.*,
mot., *npg.* See *Bacillus lactis aerogenes*,
Escherich. **Bacillus aerogenes capsula-**
tus, Welch and Nuttall. Found in the blood-
vessels in a case of thoracic aneurysm. *an-*
aer., *nlif.*, *nmot.*, *npg.* **Bacillus aéro-**
philus, Liborius. Obtained from the air.
aer., *lif.*, *nmot.*, *npg.* Syn., *Bacille aéro-*
phile, Fr. *Bacillus aeruginosus*. See *Ba-*
cillus pyocyaneus, Gessard. **Bacillus albu-**
minis, Bienstock. Found abundantly in
feces. Decomposes albumin, hence its name.
aer., *mot.*, *npg.* **Bacillus albus**, Becker,
Eisenberg. The white bacillus of water.
aer., *nlif.*, *mot.*, *npg.* **Bacillus albus an-**
aerobiscens, Vaughan. Found in water.
aer., *facanaer.*, *nlif.*, *nmot.*, *npg.* **Bacil-**
lus albus cadaveris, Strassmann and
Stricker. Found in blood of cadaver. *aer.*,
lif., *mot.*, *pg.* **Bacillus albus putridis**,
De Bary. Found in water. *aer.*, *lif.*, *mot.*,
npg. **Bacillus allantoides**, Klein. Ob-
tained from the air. *aer.*, *mot.*, *npg.* Not
sufficiently studied. **Bacillus allii**, Griffiths.
Syn., *Bacterium allii*. Found on surface of
decaying onions. *aer.*, *chg.* (green), *npg.*
Bacillus alvei, Cheshire and Cheyne. Syn.,
Bacillus preussii, Ciesielski. **Bacillus milit-**
tophorus, Cohn. Bacillus of foul brood.
Obtained from bee larvae infected with "foul
brood." *aer.*, *facanaer.*, *lif.*, *mot.*, *pg.* Also
pathogenic for mice and guinea pigs. Cf.
Bacillus of Canestrini. **Bacillus anaero-**
bicus liquefaciens, Sternberg. Obtained
from contents of intestine of yellow fever
cadaver. *anaer.*, *lif.*, *nmot.* Pathogenesis
not determined. **Bacillus anthracis**, Pol-
lender and Davaine. Syn., *Bacterium an-*
thraxicum, Bollinger. *Bacterium anthracis*,
Zopf. *Bacterium carbunculare*, Pollender,
Brauell, Delafond, Davaine. *Mettalacter*
anthracis, Trevisan. *Bacille du charbon*,
Fr.; *Milzbrand-bacillus*, Ger. Found in the
blood of animals and persons infected with
anthrax. *aer.*, *facanaer.*, *lif.*, *nmot.*, *pg.*
Bacillus aquaticus liquefaciens, Pokrows-

ky. Found in river water (Kura), and
hydrant water (Tiflis). **Bacillus aquatilis**,
Frankland. Found in well-water in the chalk-
formations of Kent, Eng. *aer.*, *lif.*, *mot.*,
npg. **Bacillus aquatilis sulcatus**, I, II,
III, IV, V, Weichselbaum. Found in hy-
drant water (Vienna). *aer.*, *facanaer.*, *nlif.*,
mot., *npg.* **Bacillus arborescens**, Frank-
land. Found in hydrant water (London).
Bacillus argenteo-phosphorescens, I, II,
III, Katz. Obtained respectively from sea-
water, phosphorescent pieces of fish, and
cuttle-fish (Sydney, N. S. W.). *aer.*, *nlif.*,
mot. (except II), *phos.*, *npg.* **Bacillus ar-**
genteo-phosphorescens liquefaciens,
Katz. Obtained from sea-water, near Syd-
ney, N. S. W. *aer.*, *facanaer.*, *lif.*, *mot.*,
phos., *npg.* **Bacillus aurantiacus**, Frank-
land. Found in well-water. *aer.*, *nlif.*,
mot., *chg.* (orange), *npg.* **Bacillus aureus**,
Adametz. Found in water; also on the skin
in cases of eczema seborrhœicum, Unna and
Tommasoli. *aer.*, *nlif.*, *mot.*, *chg.* (golden
yellow). **Bacillus b**, Vignal. See *Bacillus*
buccalis, Vignal. **Bacillus B.**, Hofmann.
Found in the larvæ of *Liparis manacha*. The
cause of *Flacherie*, or *Schlafsucht*. **Bacillus**
beribericus, Lacerda. Syn., *Micrococcus*
of beriberi, Lacerda; *Neisseria winkleri*.
Found by Lacerda, Pekelharing and Wink-
ler, in the blood of persons affected with beri-
beri. Negative results were obtained by
Eykmann and Sternberg. See *Beri-beri*.
Bacillus berolinensis indicus, Clâssen.
Found in water of the Spree. *aer.*, *nlif.*,
mot., *chg.* (indigo-blue), *npg.* **Bacillus bien-**
stockii, Bienstock. Found in human feces.
mot., *pg.* **Bacillus brassicæ**, Pommer.
Found in infusions of cabbage-leaves. *aer.*,
facanaer., *lif.*, *nmot.*, *npg.* **Bacillus**
brunneus, Adametz and Weichmann. Found
in water. *aer.*, *facanaer.*, *nlif.*, *nmot.*, *npg.*
Bacillus buccalis, Vignal. Syn., *Bacillus*
ulna, Vignal. *Bacillus a*, Vignal; *Bacillus*
b, Vignal; *Bacillus f*, Vignal; *Bacillus g*,
Vignal; *Bacillus j*, Vignal. *Bacillus buc-*
calis fortuitus, Vignal. *Bacillus buccalis mi-*
nutus, Vignal. *Leptothrix buccalis*, Vignal.
From the salivary secretions of healthy per-
sons. *aer.*, *lif.*, *npg.*, *chg.* (golden yellow).
Bacillus buccalis maximus, Miller.
Found in the mouth of man. Common.
Biologic characters undetermined. **Bacillus**
buccalis minutus, Vignal. See *Bacillus*
buccalis, Vignal. **Bacillus butylicus**, Fite.
Syn., *Bacille butylique*, Fr. See *Bacillus*
butyricus, Hueppe. **Bacillus butyricus**,
Botkin, Hueppe, Prazmowski. Obtained
from milk, hydrant water, and well water,
old cheese, garden earth, dust, etc. Syn.,
Actinobacter polymorphus, Duclaux. *Amy-*
lobacter clostridium, Trecul. *Bacillus actino-*

bacter, Duclaux. *Bacillus amylobacter*, Van Tieghem. *Bacillus butylicus*, Fitz. *Bacteridium navicula*, Reinke and Berthold. *Clostridium butyricum*, Prazmowski. *Micrococcus amylovorus*, Burrill. *Vibrione butyrique*, Pasteur. *Bacillus* of butyric acid fermentation; *Bacillus* of fire-blight; *Bacillus* of pear-blight; *Micrococcus* of apple-blight; *Micrococcus* of pear-blight. One of the most widely diffused forms, exceedingly important and varied in its powers of decomposition. It is typically anaerobic and the chief of all the ferments giving rise to butyric acid as a primary product. It plays an important part in the human economy and in the destruction of plant cellulose. Botkin regards the form observed by him as distinct. *aer.*, *anaer.*, *liqf.*, *mot.*, *npg.*, *pleom.* *Bacillus butyri fluorescens*, Lafar. Occurring in every sample of natural butter examined. See *Bacterium butyri colloideum*, Lafar. *Bacillus melochloros*, Winkler and Schröter. *Bacillus butyri viscosus*, Lafar. Found in butter. *aer.*, *nliqf.*, *np.* *Bacillus cadaveris*, Sternberg; obtained from yellow fever cadavers. *obl.*, *anaer.* (strict), *nmot.*, *pg.* *Bacillus canalis capsulatus*, Mori. Found in sewer water. *aer.*, *facanaer.*, *nliqf.*, *nmot.*, *pg.* (for mice; guinea pigs and rabbits immune). *Bacillus canalis parvus*, Mori. Found in sewer water. *aer.*, *nliqf.*, *nmot.*, *pg.* (for mice and guinea pigs). *Bacillus candicans*, Frankland. Found in soil. *aer.*, *nliqf.*, *nmot.*, *npg.* *Bacillus capsulatus*, Pfeiffer. Obtained from the blood of a guinea pig. *aer.*, *facanaer.*, *nliqf.*, *nmot.*, *pg.* (for white mice and house mice, guinea pigs, rabbits and pigeons (septicemia)). *Bacillus capsulatus mucosus*, Fäsching. Found in the nasal secretions of man in cases of influenza. *aer.*, *facanaer.*, *nmot.*, *nliqf.*, *pg.* (for white mice and field mice), *npg.* (for rabbits and pigeons). *Bacillus capsulatus smithii*, *a, b, c*, Theobald Smith. Syn., *Capsule-bacillus* of Smith. Found in the intestines of swine. *aer.*, *facanaer.*, *nliqf.*, *nmot.*, *npg.* According to Smith, possibly identical with *Bacillus lactis aerogenes*, Escherich. *Bacillus carabiformis*, Kaczynsky. Found in the stomach of meat-fed dogs. *aer.*, *liqf.*, *mot.*, *npg.* *Bacillus carotarum*, A. Koch. Obtained from cooked carrots and sugar beets. *aer.*, *liqf.*, *nmot.*, *npg.* *Bacillus catenula*, Duclaux. Syn., *Tyrophix catenula*. Found in cheese. *anaer.* (will, however grow in air), *nmot.*, *npg.* *Bacillus caucasicus*, Kern. Syn., *Bacterium caucasicum*. *Dispora caucasica*, Kern. Used in the preparation of the peculiar milk-wine, "Kefir." *aer.*, *facanaer.*, *nmot.*, *npg.* *Bacillus caviæ fortuitus*, Sternberg. Found in exudates of guinea pig inoculated with liver from yellow

fever cadaver. *aer.*, *facanaer.*, *nliqf.*, *mot.*, *npg.* *Bacillus cavidus*, Brieger. See *Bacillus cavidus*, Brieger. *Bacillus cavidus havaniensis*, Sternberg. Obtained from intestine of yellow fever cadaver. *aer.*, *facanaer.*, *nliqf.*, *mot.* *Bacillus cavidus*, Brieger. Syn., *Bacillus cavidus*. Brieger's bacillus. Obtained from human feces. *aer.*, *facanaer.*, *nliqf.*, *pg.* (not for rabbits or mice). Perhaps identical with *Bacillus neapolitanus*, Emmerich. *Bacillus chauvæi*, Bollinger and Feser, Arloing, Cornevin, and Thomas. Syn., *Bacillus* of symptomatic anthrax. *Bacille du charbon symptomatique*. Rauschbrandbacillus. *Clostridium* of symptomatic anthrax, Neelsen and Ehlers. Found in tissues of cattle suffering from symptomatic anthrax, "black leg," or "quarter evil." *anaer.*, *liqf.*, *mot.*, *pg.* *Bacillus chlorinus*, Engelmann. Syn., *Arthrobacterium chlorinum*, De Bary. *Bacterium chlorinum*, Engelmann. Found in infusions of decaying vegetable matter. *aer.*, *sap.*, *chg.* (green), *npg.* *Bacillus cholerae asiaticæ*, Koch. See *Spirillum cholerae asiaticæ*, Koch. *Bacillus cholerae gallinarum*, Flügge. See *Bacillus septiciemia hemorrhagica*, Sternberg. *Bacillus chromoaromaticus*, Galtier. Obtained from the carcass of a diseased pig. *aer.*, *facanaer.*, *liqf.*, *mot.*, *pg.* (for rabbits). *Bacillus circulans*, Jordan. Found in the water of Merrimac River. *aer.*, *facanaer.*, *liqf.*, *mot.*, *npg.* *Bacillus citreus*. See *Ascobacillus citreus*, Unna and Tommasoli. *Bacillus citreus cadaveris*, Strassmann. Found in cadaver. *aer.*, *liqf.*, *nmot.*, *chg.* (yellow). *Bacillus claviformis*, Duclaux. Syn., *Tyrophix claviformis*, Duclaux. Found in fermenting casein. *anaer.*, *pg.* *Bacillus cloacæ*, Jordan. One of the most common microbes in sewage. *aer.*, *facanaer.*, *liqf.*, *mot.*, *npg.* *Bacillus cœruleus*, Smith. Found in river water (Schuykill). *aer.*, *liqf.*, *chg.* (blue), *npg.* *Bacillus coli communis*. See *Bacillus neapolitanus*, Emmerich. *Bacillus coli similis*, Sternberg. Obtained from human liver. *aer.*, *facanaer.*, *nliqf.*, *nmot.*, *npg.* *Bacillus constrictus*, Zimmermann. Found in water. *aer.*, *facanaer.*, *nliqf.*, *mot.*, *chg.* (cadmium yellow), *npg.* *Bacillus coprogenes foetidus*, Schottelius. Obtained from intestinal contents of pigs dead of malignant erysipelas. *aer.*, *nliqf.*, *nmot.*, *pg.* *Bacillus coprogenes parvus*. Obtained from human feces. *aer.*, *nliqf.*, *nmot.*, *pg.* *Bacillus crassus*, Van Tieghem. The broadest known bacterium, 4 μ . *mot.*, *npg.* *Bacillus crassus sputigenus*, Kreibohm. Syn., *Bacillus sputigenus crassus*. From the sputum and tongue of man. *aer.*, *nliqf.*, *nmot.*, *pg.*

Bacillus crystallosus. *Bacillus cuneatus.* Syn., *Bacterium cuneatum*, Rivolta. Found in the blood and viscera of horses, cattle, and dogs dead of septic processes. *pg.* **Bacillus cuniculicidus**, Koch. See *Bacillus septicanisæ hæmorrhagiæ*, Sternberg. **Bacillus cuniculicida havaniensis**, Sternberg. Found in viscera of yellow fever cadavers. *aer., facanaer., nliqf., pg.* **Bacillus cuticularis**, Tils. Found in water. *aer., liqf., mot., chg.* (yellow), *npg.* **Bacillus cyaneophosphorescens**, Katz. Obtained from sea water (Australia). *aer., facanaer., liqf., mot., phos., chg.* (greenish), *npg.* **Bacillus cyanofuscus**, Beyerinck. Obtained from glue and Edam cheese. *aer., chg.* (green, changing to blue, brown, black), *liqf., mot., npg.* **Bacillus cyanogenus**, Fuchs. Hueppe, Neelsen. Syn., *Bacillus lactis cyanogenus*. *Bacillus syncyanum*. *Bacterium cyanogenum*, Fuchs. *Bacterium syncyanum*, Ehrenberg, Schröter. *Vibrio cyanogenus*, Fuchs. *Vibrio syncyanus*, Ehrenberg. **Bacillus of blue milk.** Found in milk. *aer., nliqf., mot., chg.* (grayish blue), *npg.* **Bacillus cystiformis**, Clado. Found in urine of patient with cystitis. *aer., nliqf., mot., npg.* **Bacillus delicatulus**, Jordan. Found in hydrant water (Lawrence, Mass.). *aer., liqf., mot., npg.* **Bacillus dentalis viridans**, Miller. Syn., Miller's bacillus. Found in carious dentine. *aer., facanaer., nliqf., pg.* **Bacillus denitrificans**, Giltay and Aberon. Obtained from the soil, air, and sewage. *aer., nliqf., mot., npg.* Cf. *Bacterium denitrificans*, Gayon and Dupetit. **Bacillus devorans**, Zimmermann. Found in well water. *aer., facanaer., liqf., mot., npg.* **Bacillus diffusus**, Frankland. Found in the soil. *aer., liqf., npg.* **Bacillus diphtheriæ**, Klebs and Löffler. Syn., *Bacillus diphtheriticus*, *Bacillus diphtheritidis*, *Micrococcus diphtheriticus*, Cohn. Found in diphtheritic false membranes. *aer., nliqf., nmot., facanaer., pg.* **Bacillus diphtheriæ columbarum**, Löffler. From the pseudomembranes in the mouths of pigeons infected with pigeon diphtheria. *aer., nliqf., nmot., pg.* **Bacillus diphtheriæ vitulorum**, Löffler. From the pseudomembranes in the mouths of calves infected with epidemic diphtheria; culture experiments unsuccessful. *pg.* Rabbits and guinea pigs immune. **Bacillus diphtheriticus**, *Bacillus diphtheridis*, *Bacillus diphtheritis*. See *Bacillus diphtheriæ*, Klebs and Löffler. **Bacillus distortus**, Duclaux. Syn., *Tyrophrix distortus*. Found in milk and cheese. *aer., mot., npg.* **Bacillus dysentericæ**, Chantemesse and Widal. Found in the intestinal contents and viscera of dysentery cadavers. *aer., nliqf., pg.* **Bacillus dysodes**, Zopf.

Syn., *Bacterium dysodes*. Found in yeast, inducing fermentative changes in bread, causing it to smell disagreeably and to become greasy and unfit for use. **Bacillus Emmerich.** See *Bacillus neapolitanus*, Emmerich. **Bacillus endocarditidis capsulatus**, Weichselbaum. From viscera of man who died of endocarditis with thrombi. *aer., nliqf., pg.* **Bacillus endocarditidis griseus**, Weichselbaum. From the heart in a case of endocarditis recurrens ulcerosa. *aer., nliqf., mot., pg.* **Bacillus enteritidis**, Gärtner. Obtained from the tissues of a cow killed on account of sickness attended with mucous diarrhea, and from the spleen of a man who died shortly after eating of the flesh of this cow. *aer., nliqf., mot., pg.* Dogs, cats, chickens, and sparrows are immune. **Bacillus epidermidis**, Bizzozzo. Syn., *Leptothrix epidermidis*, *Microsporum minutissimum*, Burchart. From the epidermis between the toes. *aer., nliqf., npg.* **Bacillus epidermidis**, Bordoni-Uffreduzzi. See *Bacillus of Scheurlen*. **Bacillus Epsilon**, von Miller. See *Spirillum*, Finkler-Prior. **Bacillus erysipelas suis**, Koch, Löffler, Schütz, Pasteur. Syn., *Bacillus erysipelas malignum*, *Bacillus minimus*, *Bacillus murisepticus*, Flügge. *Bacillus septicus*, Koch. *Bacillus of mouse septicæmia*. *Bacillus of hog erysipelas*. First obtained by Koch by injecting putrefying flesh beneath the skin of mice, afterward by Löffler and Schütz from carcasses of swine dead of "Rotlauf." *aer., facanaer., nliqf., pg.* **Bacillus erysipelas leporis**, Koch. *Bacillus of erysipelas in the rabbit*. Found in the erysipelous inflammation in a rabbit's ear caused by the injection of mouse's dung. **Bacillus erythrosporus**, Eidam, Cohn, and Miflet. Found in putrefying albuminous fluids, water, etc. *aer., nliqf., mot., chg.* (greenish yellow, fluorescent), *npg.* **Bacillus ethaceticus**, Frankland. Cause of fermentation of arabinose. **Bacillus ex penumo-enteritidis suis**, E. Klein, Schütz, Salmon, Cornil, Chantemesse, Selander. *Bacillus of hog cholera*. Found in the intestinal contents, blood, and viscera of swine dead of hog cholera. *aer., facanaer., nliqf., mot., pg.* **Bacillus f.** Vignal. See *Bacillus buccalis*, Vignal. **Bacillus facultatus**, Sadebeck and E. Fränkel. Found in non-malignant pharyngeal mycosis. **Bacillus fœtidus**, Liborius. Syn., *Bacterium fœtidum*, *Clostridium fœtidus*, Liborius. Obtained from soil. *anaer., liqf., mot.* **Bacillus fœtidus lactis**, Jensen. From milk. *aer., nliqf., npg.* **Bacillus fœtidus ozæniæ**, Hajek. Obtained from nasal secretions of patients with ozæna. *aer., facanaer., liqf., mot., pg.* **Bacillus figu-**

rana, Crookshank. See *Bacillus mesentericus vulgaris*. *Bacillus figuratus*, Vaughan. Found in water. *arr.*, *hgf.*, *mod.*, *app.* *Bacillus filiformis*, Duclaux. *Syn. Thiothrix filiformis*. Found in cheese and milk. *aer.*, *mod.*, *app.* *Bacillus filiformis*, Tills. Found in water. *arr.*, *hgf.*, *mod.*, *app.* *Bacillus filiformis havaniensis*, Sternberg. From the liver of a yellow fever cadaver. *anaer.*, *fac-aer.*, *slgt.*, *anaer.*, *app.* *Bacillus fitzianus*, Zopf. See *Bacillus mitchellii* Cohn. *Bacillus flavescens*, Poell. Found in swamp water. *aer.*, *slgt.*, *slgt.*, yellow. *Bacillus flavocoriaceus*, Esenb. *Syn. Sulphur yellow bacillus* (Adams). Found in water. *aer.*, *slgt.*, *mod.*, *slgt.*, sulphur yellow. *app.* *Bacillus flavus*, Max. Found in drinking water. *aer.*, *hgf.*, *slgt.*, golden yellow), *anaer.* *Bacillus fluorescens aureus*, Zimmermann. Found in hygienic water (Chemnitz). *aer.*, *slgt.*, *mod.*, *slgt.* (ocher yellow). *app.* *Bacillus fluorescens longus*, Zimmermann. Found in hygienic water (Chemnitz). *aer.*, *slgt.*, *mod.*, *slgt.* (grayish yellow). *app.* *Bacillus fluorescens liquefaciens*, Függe. Common in water and purifying effluents. *aer.*, *hgf.*, *mod.*, *slgt.*, greenish yellow. *app.* *Bacillus viscosus*, Frankland. *Bacillus fluorescens liquefaciens microaerophilus*, Unna and Tomaszewski. Found upon skin in cases of *eczema scabrosum*. *aer.*, *fac-aer.*, *hgf.*, *mod.*, *slgt.*, greenish yellow. *app.* Possibly identical with the preceding species. *Bacillus fluorescens nitralis*, Schmoelk. *Syn. Bacillus aralis* (Günther) bacillus. Found in snow and ice water from Norwegian glaciers. *aer.*, *hgf.*, *mod.*, *slgt.*, bluish green. *app.* *Bacillus fluorescens nonliquefaciens*, Schiller. *Bacillus fluorescens putidus*, Függe. Found in water. *aer.*, *slgt.*, *mod.*, *slgt.* (greenish). *app.* *Bacillus fluorescens tenuis*, Zimmermann. Found in hygienic water (Chemnitz). *aer.*, *slgt.*, *mod.*, *slgt.* (greenish yellow). *app.* *Bacillus fetidus*, Paster. *Syn. Bacterium fetidum*, Tur. *Clostridium fetidum*, Liborius. *Corynebacterium*. Found in cases of foot sweating feet, in the excretions of mice inoculated with garden earth and in cases of malignant edema. *aer.*, *hgf.*, *mod.*, *app.* *Bacillus fetidus lactis*, Jensen. Found in milk. *aer.*, *slgt.*, *app.* *Bacterium Bacillus aspidioticus*, Zimmermann. *Bacillus fulvus*, Zimmermann. Found in hygienic water (Chemnitz). *aer.*, *hgf.*, *mod.*, *slgt.*, gamboge yellow. *app.* *Bacillus faecalis*, (derived from a purifying effluent of waste, from the ex. etc. of *Bacterium faecale*, Schreiber) *Bacillus faecalis* Linbath, Scheidegger. (Derived from

[illegible]

liqf., mot. Pathogenesis not determined. *Bacillus hominis capsulatus*, Bordoni-Uffreduzzi. Syn., *Proteus capsulatus septicus*, Banti. *Proteus hominis capsulatus*, Bordoni-Uffreduzzi. Bacillus of rag-picker's disease. Obtained from persons dead from rag-picker's disease. *aer., nliqf., nmot., pg.* *Bacillus hyacinthi septicus*, Heinz. Found in diseased hyacinths. *aer., facanaer., nliqf., mot., npg.* Cf. *Bacterium hyacinthi*, Wakker. *Bacillus hyalinus*, Jordan. Found in hydrant water (Lawrence, Mass.). *aer., facanaer., liqf., mot., npg.* Reduces nitrates rapidly. *Bacillus hydrophilus fuscus*, Sanarelli. From the lymph of diseased frogs. *aer., liqf., mot., pg.* *Bacillus implexus*, Zimmermann. Found in hydrant water (Chemnitz). *aer., liqf., mot., npg.* *Bacillus incanus*, Pohl. Found in swamp water. *aer., liqf., mot.* Not sufficiently studied. *Bacillus indicus*, Koch. Syn., *Bacillus indicus ruber*, Koch. *Micrococcus indicus*, Koch. Found in the intestinal contents of a monkey. *aer., facanaer., liqf., mot., chg.* (brick-red), *pg.* (for rabbits). *Bacillus indigoferus*, Classen. Found in river water (Spree). *aer., nliqf., mot., chg.* (deep indigo blue), *npg.* *Bacillus indigogenus*, Alvarez. Found in an infusion of the leaves of the indigo plant. (*Indigofera tinctoria*, L.). *aer., mot., chg.* (indigo blue), *pg.* *Bacillus inflatus*, A. Koch. Obtained from the air. *aer., liqf., mot., npg.* *Bacillus intestinalis motilis*, Sternberg. From the intestinal contents of yellow fever cadavers. *aer., facanaer., nliqf., mot., npg.* *Bacillus inunctus*, Pohl. Found in swamp water. *aer., facanaer., nliqf., mot.* Pathogenesis not determined. *Bacillus invisibilis*, Vaughan. Found in water. *aer., facanaer., nliqf., mot., npg.* *Bacillus iodo-coccus vaginatus*, Miller. See *Iodococcus vaginatus*. *Bacillus iris*, Frick. *aer., nliqf., nmot.* (green), *npg.* *Bacillus janthinus*, Zopf. Syn., *Bacterium janthinus*, Zopf. Violet bacillus, found in sewage (Lawrence, Mass.). *aer., liqf., mot., chg.*, (bluish violet), *npg.* *Bacillus klebsii*. See *Bacillus syphilidis*, Lustgarten. *Bacillus lacticus*, Pasteur. See *Bacillus acidilactici*, Hueppe. *Bacillus lactis aerogenes*, Escherich and Abelous. Syn., *Bacillus aerogenes*, *Bacillus capsulatus*, Smith. *Bacterium lactis aerogenes*, Escherich. Found in the stomach and intestine of healthy adults. *aer., facanaer., nliqf., npg.* *Bacillus lactis albus*, Löffler. Found in milk. *aer., liqf., mot., npg.* *Bacillus lactis cyanogenus*. See *Bacillus cyanogenus*, Hueppe. *Bacillus lactis erythrogenes*, Hueppe and Baginsky. Syn., *Bacillus* of red milk. Found in milk and

the feces of a child. *aer., liqf., nmot., chg.* (yellow and red), *npg.* *Bacillus lactis pituitosi*, Löffler. Syn., *Bacillus lactis viscosus*, Adametz. Found in milk. *aer., nliqf., npg.* *Bacillus lactis viscosus*, Adametz. Found in ropy milk. *aer., nliqf., nmot., npg.* Capsulated. *Bacillus latericeus*, Eisenberg. Found in water. *aer., nliqf., nmot., chg.* (brick-red), *npg.* *Bacillus leioderms*, Löffler. Syn., *Bacillus leioderms*, Flügge. *Bacillus leporis lethalis*, Gibier and Sternberg. From the intestinal contents of yellow fever patients. *aer., liqf., mot., pg.* *Bacillus lepræ*, Armauer and Hansen. Found in leprosy tubercles. Cannot be cultivated in ordinary media. Specific pathogenesis settled by inoculation of a condemned criminal. *Bacillus leptosporus*, L. Klein. Obtained from the air. *aer., mot., npg.* *Bacillus lethalis*, Babes. Syn., *Proteus lethalis*. Obtained from spleen and lung of patient dead of septicemia. *aer., facanaer., liqf., mot., pg.* *Bacillus limbatus acidilactici*, Marpmann. Found in cow's milk. *aer., nliqf., nmot., npg.* *Bacillus limosus*, Russel. Obtained from mud (Bay of Naples). *aer., liqf., mot.* Pathogenesis not determined. *Bacillus lineola*. See *Bacterium lineola*, Cohn. *Bacillus lioderms*, Flügge. See *Bacillus leioderms*, Löffler. *Bacillus liquefaciens*, Eisenberg. Found in water. *aer., liqf., mot., npg.* *Bacillus liquefaciens bovis*, Arloing. Syn., *Pneumobacillus liquefaciens bovis*. From the lungs of a diseased ox. *aer., facanaer., liqf., nmot., pg.* *Bacillus liquefaciens communis*, Sternberg. Obtained from the feces of yellow fever patients. *aer., facanaer., liqf., mot., npg.* *Bacillus liquefaciens magnus*, Lüderitz. Found in the exudates of mice inoculated with garden earth. *anaer., liqf., mot., npg.* *Bacillus liquefaciens parvus*, Lüderitz. Source same as last. *anaer., liqf., nmot., npg.* *Bacillus liquidus*, Frankland. Found in river water (Thames). Common. *aer., liqf., mot., npg.* *Bacillus litoralis*, Russel. Obtained from mud (Bay of Naples). *aer., facanaer., liqf., mot.* Pathogenesis not determined. *Bacillus lividus*, Plagge and Proskauer. Found in hydrant water (Berlin). *aer., facanaer., liqf., mot., chg.* (brick-red), *npg.* *Bacillus lucens*, Van Tieghem. Syn., *Bacterium lucens*, Van Tieghem. Found on the surface of water. *aer., nmot.* Brilliant. *Bacillus luteus*, Flügge. Syn., *Bacterium luteum*. Found in superficial layers of gelatin plate culture. *aer., nmot., nliqf., chg.* (yellow), *npg.* *Bacillus luteus suis*, Salmon and Smith. Found in the perivisceral fluids of swine. *aer., liqf., mot.,*

chg. (yellow red). Smells like glue. *Bacillus lyssæ*, Pasteur. Syn., *Coccobacterium lyssa*, Rivolta. *Lissophyton suspectum*, Hallier. Cf. *Bacterium septicum sputigenum*, Fraenkel. Found in saliva of hydrophobic patients and animals. Cf. *Bacillus septicus sputigenus*, Flügge. *Bacillus mallei*, Cuboni. From macerated corn and the feces of pellagra patients. *aer.*, *liqf.*, *mot.*, *npg.* *Bacillus malandriæ*, Israel. See *Bacillus mallei*, Löffler. *Bacillus malarie*, Klebs and Tommasi-Crudeli. Bacillus of the blood. Obtained from the soil and air of the Campagna, from swamps, from the blood of malarial patients, etc. Pathogenesis disputed. *Bacillus mallei*, Löffler. Syn., *Bacillus malandria*, Israel. Found in the nasal discharges, nodules, etc., of animals with glanders. *aer.*, *facanaer.*, *mot.* (strictly parasitic), *pg.* (proven by inoculation). *Bacillus martinez*, Sternberg. Obtained from the liver of a yellow fever cadaver. *aer.*, *facanaer.*, *nlqf.*, *nmot.*, *npg.* *Bacillus megatherium*, De Bary. Found on the leaves of boiled cabbage. *aer.*, *liqf.*, *mot.*, *npg.* *Bacillus melittophorus*. See *Bacillus alvei*, Cheshire and Cheyne. *Bacillus melochloros*, Winkler and Schrötter. Syn., *Bacillus butyri fluorescens*, Lafar. Obtained from the dejections of the larva in a wormy apple. *aer.*, *liqf.*, *mot.*, *chg.* (emerald green), *pg.* (for rabbits). *Bacillus membranaceus amethystinus*, Eisenberg. Found in well water (Spalato). *aer.*, *liqf.*, *nmot.*, *chg.* (dark violet), *npg.* *Bacillus meningitidis purulentæ*, Naumann and Schäffer. Obtained from pus taken from beneath the pia mater in a person dead of purulent meningitis. *aer.*, *facanaer.*, *nlqf.*, *mot.*, *pg.* *Bacillus merismopædioides*, Zopf. Syn., *Arthrobacterium* or *Bacterium merismopædioides*, De Bary. Obtained from sewage mud; the type of the so-called "tablet-cocci." *Bacillus mesentericus fuscus*, Flügge. Potato bacillus. Obtained from the air, hay dust, water, etc. Common. *aer.*, *liqf.*, *mot.*, *npg.* *Bacillus mesentericus ruber*, Globig. Potato bacillus. Found upon potatoes. *aer.*, *liqf.*, *mot.*, *chg.* (reddish yellow or pink), *npg.* *Bacillus mesentericus vulgaris*, Flügge. Syn., *Bacillus figurans*, Crookshank. *Proteus vulgaris*. Potato bacillus. Common. Found in potatoes, water, milk, intestinal contents of man, etc. *aer.*, *liqf.*, *npg.* *Bacillus miniaceus*. See *Bacillus ruber*. *Bacillus minimus*, Klebs. See *Bacillus erysipelatos suis*, Koch, etc. *Bacillus mirabilis*, Hauser. Syn., *Proteus mirabilis*. Obtained from decaying animal matter. *aer.*, *facanaer.*, *liqf.*, *mot.*, *pg.* *Bacillus mollusci*, Domenici. Found in mol-

luscum contagiosum; resembles somewhat *Bacillus lepræ* and *Bacillus malarie*. Pathogenesis disputed, the disease being attributed by some to psorosperms. *Bacillus multipediculosus*, Flügge. Found in air and water. *aer.*, *nlqf.*, *nmot.*, *npg.* *Bacillus murisepticus*, Flügge. See *Bacillus erysipelatos suis*, Koch. *Bacillus murisepticus pleomorphus*, Karlinski. Found in purulent uterine discharges. *aer.*, *facanaer.*, *liqf.*, *mot.*, *pg.* Probably identical with *Proteus vulgaris*, Hauser. *Bacillus muscoides Liborius*, Flügge. Found in soil, old cheese, cow-dung, etc. *anaer.*, *nlqf.*, *mot.*, *npg.* *Bacillus mycoides*, Flügge. Common in soil and water. *aer.*, *nlqf.*, *mot.*, *npg.* *Bacillus mycoides roseus*, Scholl. Found in the soil. *aer.*, *liqf.*, *chg.* (red), *npg.* *Bacillus neapolitanus*, Emmerich. Syn., *Bacillus coli communis*, *Bacterium coli commune*, Escherich, *Bacterium neapolitanum*, Colon bacillus of Escherich. Bacillus of Booker, *a to n*. A normal inhabitant of the intestine of man; resembles the bacillus of typhoid fever, differing from it in producing gas, coagulating milk, forming lactic acid in media containing lactose or glucose, in producing indol, in being nonmotile, and in giving a visible growth on potato. It is the cause of certain forms of suppuration in the peritoneum, liver, pelvis of kidney, urethra, etc. *aer.*, *facanaer.*, *nlqf.*, *nmot.*, *pg.* *Bacillus necrophorus*, Löffler. Obtained from rabbits after inoculation in the anterior chamber of the eye with portions of condyloma. *aer.*, *facanaer.*, *nlqf.*, *pleom.*, *pg.* *Bacillus nodosus parvus*, Lustgarten. From the healthy urethra of man. *aer.*, *facanaer.*, *nlqf.*, *nmot.*, *npg.* *Bacillus œdematis aerobius*. A new bacillus of malignant edema, Klein. Found in the exudates of guinea pigs after inoculating with garden earth. *aer.*, *facanaer.*, *nlqf.*, *mot.*, *pg.* *Bacillus œdematis maligni*, Koch. Syn., *Bacillus œdematis*, Koch; *Bacillus septicus*, Pasteur. Bacillus of gangrene, Bacillus of gangrenous septicemia. Pink bacillus of spreading edema, A. B. Harris. A widely scattered species, obtained from surface soil, dust, putrefying matter, foul water, etc., and from the exudates produced by inoculating animals with garden earth. *anaer.*, *liqf.*, *mot.*, *pg.* *Bacillus of Afanassiew*. See *Bacillus tussis convulsivæ*. *Bacillus of Albuminous Decomposition*. See *Bacillus putrificus coli*. *Bacillus of Allantiasis*, Müller, Hoppe-Seyler. See also *Sarcina botulina*, Van den Corput. Cf. Bacillus of choleraic diarrhea from meat-poisoning, Klein. Found in poisonous sausages, particularly in "Blunzen." *aer.*, *liqf.*, *pg.* *Bacillus of Alopecia areata*, Kasauli. Found at-

tached to the hairs from the diseased patches; easily cultivated. See *Micrococcus of Alopecia areata*, and *Bacterium decalvans*. **Bacillus of Babes and Oprescu.** Obtained from a case presenting symptoms of typhus fever. *aer.*, *facanaer.*, *nlqf.*, *mot.*, *pg.* **Bacillus of Belfanti and Pascarola.** Syn., *Impftetanusbacillus*, Ger. From the pus of wounds in a person dead of tetanus. *aer.*, *facanaer.*, *nlqf.*, *nmot.*, *pg.* **Bacillus of Beri-beri**, Eykman. See *Bacillus beribericus*, Lacerda. **Bacillus of Blue Milk.** See *Bacillus cyanogenus*, Hueppe. **Bacillus of Blue or Green Pus.** See *Bacillus pyocyaneus*, Gessard. **Bacillus of Booker**, a to n. Found in alvine discharges of children suffering from cholera infantum; probably varieties of *Bacillus neapolitanus*, Emmerich. **Bacillus of Bo-vet.** Obtained from the intestines of a woman dead of acute choleraic enteritis. *aer.*, *nlqf.*, *mot.*, *pg.* **Bacillus of Braxy.** See *Bacillus anthracis*, Pollender and Davaine. **Bacillus of Bronchitis**, Lum-nitzer. Syn., *Bacillus of Lumnitzer*. From the bronchial secretions in cases of putrid bronchitis. *aer.*, *mot.* Cf. *Micrococcus of bronchitis*, Picchini. **Bacillus of Butyric Acid Fermentation.** See *Bacillus butyricus*, Prazmowski. **Bacillus of Carcinoma**, Rappin and Scheurlen and Domingo-Freire. See *Bacillus of Scheurlen*. **Bacillus of Canestrini.** Found in larvae and bees suffering from a malady common among bees in certain parts of Italy. *aer.*, *liqf.*, *mot.*, *chg.* (pink), *pg.* **Bacillus of Canon and Pielicke.** See *Bacillus of measles*. **Bacillus of Cattle Plague**, Metschnikoff. See *Bacillus septicaemia hæmorrhagica*, Sternberg. **Bacillus of Cazal and Vaillard.** Obtained from cheesy nodules upon the peritoneum and in the pancreas. *aer.*, *facanaer.*, *liqf.*, *mot.*, *pg.* **Bacillus of Chancroid**, Ducrey. Found in pustules developed upon the arm from the inoculation of chancroidal virus; does not grow in artificial cultures. Cf. *Micrococcus ulceris mollis*. **Bacillus of Cheese.** See *Spirillum tyrogenum*, Denecke. **Bacillus of Chicken Cholera.** See *Bacillus septicaemia hæmorrhagica*, Sternberg. **Bacillus of Cholera.** See *Spirillum cholerae asiatica*, Koch. **Bacillus of Choleraic Diarrhea from Meat Poisoning**, Klein. Found in the blood and feces of persons poisoned with tainted meat. Cf. *Bacillus of allantiasis*, Müller and Hoppe-Seyler. **Bacillus of Cholera in Ducks**, Cornil and Toupet. Obtained from the blood of ducks that had died of an epidemic disease characterized by choleraic symptoms. *aer.*, *nlqf.*, *nmot.*, *pg.* (for ducks, but not for chickens or pigeons). **Bacillus of Chyl-**

uria, Wilson. Found in the coagulum of chylous urine. *aer.*, *facanaer.*, *mot.* **Bacillus of Colomiatti.** Obtained from cases of conjunctivitis and xerotic masses in the eye. *aer.*, *nlqf.*, *nmot.*, *npg.* Cf. *Bacillus of xerosis epithelialis conjunctivæ*, Leber. **Bacillus of Conjunctival Catarrh**, Koch. **Bacillus of conjunctivitis**, *Bacillus of pink eye*, Weeks. *aer.*, *nlqf.*, *pg.* Resembles *Bacillus xerosis*. **Bacillus of Dantec.** Syn., *Bacille du rouge de morue*, Fr. Obtained from salted codfish to which it gives a red color. *aer.*, *liqf.*, *mot.*, *chg.* (red), *npg.* **Bacillus of Davaine's Septicemia.** See *Bacillus septicaemia hæmorrhagica*, Sternberg. **Bacillus of Demme.** Obtained from the contents of tumors and pustules and from the blood of patients suffering from erythema nodosum. *aer.*, *facanaer.*, (?) *nlqf.*, *pg.* (to guinea pigs; rabbits, dogs, and goats refractory). **Bacillus of Diphtheria in Calves.** See *Bacillus diphtherie vitulorum*, Löffler. **Bacillus of Diphtheria in Doves.** See *Bacillus diphtherie columbarum*, Löffler. **Bacillus of Diphtheria in Man.** See *Bacillus diphtherie*, Klebs and Löffler. **Bacillus of Döderlein.** See *Bacillus vaginalis*, Döderlein. **Bacillus of Endocarditis.** See *Bacillus pyogenes fatidus*, Fränkel and Saenger. **Bacillus of Egyptian Catarrhal Conjunctivitis.** See *Bacillus of Kartulis*. **Bacillus of Erysipelas in the Rabbit**, Koch. See *Bacillus erysipelatos leporis*, Koch. **Bacillus of False Hog Cholera.** See *Bacillus parvus ovatus*, Löffler. **Bacillus of False Tuberculosis in Rabbits.** See *Bacillus of pseudotuberculosis* in rabbits, Malassez and Vignal. **Bacillus of FioCCA.** Found in the saliva of cats and dogs. *aer.*, *facanaer.*, *nlqf.*, *nmot.*, *pg.* (for rabbits, guinea pigs, rats, mice). **Bacillus of Foot Sweat.** See *Bacillus fatidus*, *Bacillus saprogenes* No. II. and *Bacillus graveolens*. **Bacillus of Foul Brood of Bees.** See *Bacillus alvei*, Cheshire and Cheyne. **Bacillus of Fowl Cholera.** See *Bacillus septicaemia hæmorrhagica*, Sternberg. **Bacillus of Friedländer.** See *Bacillus pneumoniae friedländeri*. **Bacillus of Fulles**, III. **Bacillus of Gangrene.** See *Bacillus oedematis maligni*, Koch, *Bacillus saprogenes*, III, Rosenbach, and *Bacillus of senile gangrene*. **Bacillus of Gangrenous Septicemia.** See *Bacillus oedematis maligni*, Koch. **Bacillus of Gessner.** See *Bacterium tholoidum*, Gessner. **Bacillus of Glanders.** See *Bacillus mallei*, Löffler. **Bacillus of Green Diarrhea in Children**, Lesage. See *Bacillus of Lesage*. **Bacillus of Green Pus.** See *Bacillus pyocyaneus*, Gessard. **Bacillus of Grouse Disease**, Klein. Obtained

from the lungs and liver of grouse that had died of an epidemic disease. *aer.*, *nliqf.*, *nmot.*, *pg.* (for mice, guinea-pigs, linnets, green finches, sparrows; chickens, pigeons, and rabbits immune). *Bacillus* of Hay Infusions. See *Bacillus subtilis*, Ehrenberg. *Bacillus heminecrobiphilus*, Arloing. Obtained from the callous lymphatic glands of a guinea pig. *aer.*, *facanaer.*, *nliqf.*, *mot.*, *pleom.*, *pg.* *Bacillus* of Hog Cholera, Salmon and Smith. See *Bacillus ex pneumoenteritide suis*, Klein. *Bacillus* of Hog Erysipelas. See *Bacillus erysipelatos suis*, Koch, etc. *Bacillus* of Horse Pox, Dieckhoff and Grawitz. Syn., *Bacillus* of acne contagiosa of horses. Obtained from the pustules of horses suffering from acne contagiosa. *aer.*, *nliqf.*, *pg.* *Bacillus* of Hydrophobia. See *Bacillus lyssæ*, Pasteur. *Bacillus* of Icterus, Karlinsky and Ducamp. Found in the blood during an attack of infectious icterus. Did not thrive on ordinary culture media. *Bacillus* of Influenza, Pfeiffer, Kitasato, and Canon. Influenza bacillus. Found in the purulent bronchial secretions and in the blood of persons suffering from epidemic influenza. *aer.*, *nmot.*, *pg.* Cf. *Micrococcus influenzae*, Letzerich. *Bacillus* of Intestinal Diphtheria in Rabbits, Ribbert. Obtained from the viscera of rabbits dead of a disease characterized by a diphtheritic inflammation of the intestinal mucous membrane. *aer.*, *nliqf.*, *nmot.*, (?) *pg.* *Bacillus* of Jeffries. A group (A, G, J, K, P, S, Z), of bacilli, resembling *Bacillus neapolitanus*, Emmerich, and *Bacillus lactis aerogenes*, Escherich. Found in the alvine discharges of children suffering from summer diarrhea. *anaer.*, *npg.* *Bacillus* of Jequirity Ophthalmia, DeWecker and Sattler. Found in cases of jequirity ophthalmia and in infusions of jequirity seeds (*Abrus precatorius*), the supposed source of Abrin. *aer.*, *liqf.*, *mot.*, *npg.* The disease is now held to be caused by a soluble poison, *jequiritin*. *Bacillus* of Karlinski. See *Bacillus murisepticus pleomorphus*, Karlinski. *Bacillus* of Kartulis, Koch and Kartulis. Obtained from the conjunctiva in cases of Egyptian catarrhal conjunctivitis. *aer.* Pathogenesis not well determined. *Bacillus* of Koubasoff. Obtained from carcinomatous growth in the human stomach. *aer.*, *facanaer.*, *nliqf.*, *mot.*, *pg.* (for guinea pigs and rabbits). Cf. *Bacillus* of Scheuerlen. *Bacillus* of Lactic Acid Fermentation. See *Bacillus acidilactici*, Hueppe. *Bacillus* of Laser. Obtained from mice infected with an epidemic disease. *aer.*, *facanaer.*, *nliqf.*, *mot.*, *pg.* (for field-mice, guinea-pigs, rabbits and pigeons). *Bacillus* of Leprosy. See

Bacillus lepra, Hansen. *Bacillus* of Lesage. *Bacillus* of green diarrhea in children. Obtained in the green alvine discharges of infants suffering from green diarrhea. *aer.*, *nliqf.*, *mot.*, *pg.* According to Baumgarten identical with *Bacillus fluorescens non-liquefaciens*, Schiller. *Bacillus* of Letzerich. Obtained from the urine of children suffering from nephritis interstitialis primaria. *aer.*, *liqf.*, *pg.* *Bacillus* of Liborius. See *Bacillus amylobacter*, Prazmowski. *Bacillus* of Lichen Ruber, Laser. Found in the lymph passages in cases of lichen ruber. *Bacillus* of Lucet. Obtained from chickens and turkeys suffering from an infectious form of septicemia characterized by dysenteric discharges. *aer.*, *facanaer.*, *nliqf.*, *nmot.*, *pg.* *Bacillus* of Lumnitzer. See *Bacillus* of Bronchitis. *Bacillus* of Lupus. See *Bacillus tuberculosis*, Koch. *Bacillus* of Lustgarten. See *Bacillus syphilidis*, Lustgarten. *Bacillus* of Lymph in Fishes, Oliver and Richet. Observed in the lymph of certain fishes. *aer.*, *mot.*, *npg.* *Bacillus* of Malaria. See *Bacillus malariae*, Klebs and Tommasi-Crudeli. *Bacillus* of Malignant Edema. See *Bacillus adematidis maligni*, Koch. *Bacillus* of Measles, Canon and Pielicke. Syn., *Bacillus* of Canon and Pielicke. Found in the blood and in the secretions of the nose and conjunctiva of persons with measles. Cf. *Micrococcus* of measles, Klebs and Keating. *Bacillus* of Mouse Septicemia. See *Bacillus erysipelatos suis*, Pasteur. *Bacillus* of Necrosis of the Liver in Badgers, Eberth. Found in the necrotic liver of a badger. *Bacillus* of Necrosis of the Liver in Guinea Pigs, Eberth. Found in the necrotic liver of a guinea-pig. *Bacillus* of Nephritis, Letzerich. See *Bacillus* of Letzerich. *Bacillus* of Nocard. Found in the superficial abscesses of cattle suffering from farcy. *aer.*, *nmot.*, *pg.* *Bacillus* of Okada. Obtained from the dust between the boards of a floor. *aer.*, *facanaer.*, *nliqf.*, *nmot.*, *pg.* *Bacillus* of Osteomyelitis, Kraske and Becker. Found in cases of osteomyelitis. *aer.*, *liqf.* (or in some forms *nliqf.*), *mot.*, *pleom.*, *pg.* Held by Pasteur as identical with the micrococcus of furuncle. *Bacillus* of Phthisis. See *Bacillus tuberculosis*, Koch. *Bacillus* of Pink Eye, Weeks. See *Bacillus of Conjunctival Catarrh*, Koch. *Bacillus* of Potato Rot, Kramer. Obtained from potatoes affected with wet rot. *aer.*, *liqf.*, *mot.*, *npg.* Putrefies the albuminous substances of potatoes. *Bacillus* of Pseudotuberculosis in Rabbits, Eberth, Malassez, and Vignal. Syn., *Bacille* de la tuberculose zoogléique, *Bacille* de la pseudo-tuberculose du lapin,

Fr.; *Bacillus der Pseudo-tuberculose der Kaninchen*, Ger. Found in the nodular deposits characteristic of this disease. *Bacillus of Puerperal Fever*. See *Bacillus puerperalis*, Engel and Spillmann. *Bacillus of Purpura Hæmorrhagica of Babes*. Obtained from the viscera of a patient dead of purpura hæmorrhagica. *aer.*, *facanaer.*, *liqf.*, *nmot.*, *pg.* Resembles the next two forms. *Bacillus of Purpura Hæmorrhagica of Kolb*. Obtained from the viscera of patients dead of purpura. *aer.*, *facanaer.*, *liqf.*, *nmot.*, *pg.* *Bacillus of Purpura Hæmorrhagica of Tizzoni and Giovannini*. Obtained from the blood of two children dead of purpura hæmorrhagica. *aer.*, *facanaer.*, *nlif.*, *nmot.*, *pg.* *Bacillus of Pyæmia*, Beltzow. Two forms found in pyæmia, one of which resembles *Bacillus œdematis maligni*, Koch. *Bacillus of Rabbit Septicæmia*. See *Bacillus septicæmia hæmorrhagica*, Sternberg. *Bacillus of Rabies*. See *Bacillus lyssa*, Pasteur. *Bacillus of Rag-pickers' Disease*. See *Bacillus hominis capsulatus*, Bordoni-Uffreduzzi. *Bacillus of Red Bread*. See *Bacillus prodigiatus*, Ehrenberg. *Bacillus of Red Milk*. See *Bacillus lactis erythrogenes*, Hueppe; and *Bacillus prodigiatus*, Ehrenberg. *Bacillus of Rheumatism*. See *Bacillus rheumathritidis*, Kussmaul. *Bacillus of Rhinoscleroma*. See *Bacillus rhinoscleromatis*, Cornil and Alvarez. *Bacillus of Roth*, I, II. Obtained from old rags. *aer.*, *facanaer.*, *nlif.*, *nmot.*, *pg.* Resembles *Bacillus neapolitanus*, Emmerich; and *Bacillus cavicidus*, Burger. *Bacillus of Rouget*. See *Bacillus erysipelatos suis*, Koch, etc. *Bacillus of Scarlet Fever*, Crooke. Found in the nasal mucus and tissues of the throat in a case of anginose scarlet fever. *mot.* Resembles *Bacillus œdematis maligni*, Koch, and the *Proteus* of Hauser. According to Dehle the parasite of this disease is a protozoon. Cf. *Micrococcus scarlatinae*. *Bacillus of Scheuerlen*, Scheuerlen, Rapin, and Domingo Freire. Syn., *Bacillus epidermidis*, Bordoni Uffreduzzi. From cancerous tissues and from mammary epithelium of healthy persons. According to Macé possibly identical with a variety of *Bacillus mesentericus vulgaris*, Flügge. *aer.*, *liqf.*, *mot.*, *npg.* *Bacillus of Schimmelbusch*. Obtained from the necrotic tissues of a case of cancrum oris or noma. *aer.*, *nlif.*, *pg.* *Bacillus of Schou*. See *Bacillus pneumonicus agilis*, Schou. *Bacillus of Senile Gangrene*, Tricomi. *Bacillus of gangrene*, *Bacillus of Tricomi*. Found in the blood, in the tissues along the line of demarcation, in the subcutaneous tissue and the lymph-spaces of the skin, and in the ichor-

ous discharge. *aer.*, *liqf.*, *nmot.*, *pg.* Cf. *Bacillus œdematis maligni*, Koch and *Bacillus saprogenes*, III, Rosenbach. *Bacillus of Septicæmia in Man*, Klein. *Bacillus of Southern Cattle Plague*, F. S. Billings. Syn., *Bacillus of Texas fever in cattle*. Found in the blood, viscera, and manure of cattle infected with cattle plague or Texas fever, and in their discharges; also in the ticks (*Boophilus bovis*, Curtice) infecting the cattle. The spread of the disease is largely due to the manure of the cattle and also indirectly to the ticks. *aer.*, *nlif.*, *mot.*, *pleom.*, *pg.* Smith and Kilborn now attribute this disease to the psorosperm *Pyrosoma bigeminum*. *Bacillus of Spontaneous Rabbit Septicæmia*, Eberth. See *Bacillus of swine plague*, Marseilles. *Bacillus of Symptomatic Anthrax*. See *Bacillus chauvoui*, Bollinger and Feser. *Bacillus of Swine Pest*, Selander. See *Bacillus ex pneumoenteritide suis*, Klein. *Bacillus of Swine Plague*, Marseilles, Reitsch, and Jobert. Syn., *Bacillus suis*, Detmers. *Bacillus of spontaneous rabbit septicæmia*; *Bacillus of swine plague*, Detmers and Billings; *Bacillus of ferret disease*. First obtained from swine attacked by a fatal epidemic disease in Marseilles. *aer.*, *nlif.*, *mot.*, *pg.* Caneva and others identify this with the *Bacillus of ferret disease* and with that of American swine plague; also with the *Bacillus of Southern cattle plague*, Billings; but this latter seems to be a distinct species. See *Bacillus tardigradus*. *Bacillus of Syphilis*. See *Bacillus syphilidis*, Lustgarten. *Bacillus of Tetanus*. See *Bacillus tetani*, Nicolaier. *Bacillus of Texas Cattle Fever*, F. S. Billings. See *Bacillus of Southern Cattle Plague*. *Bacillus of the Blood*. *Bacillus of the Lungs of Cattle*, Lüstig. Found in the lungs of healthy and diseased cattle. *aer.*, *liqf.* *Bacillus of the Meconium*, Escherich. Found in the meconium. *Bacillus of the Pneumoenteritis of the Pig*. See *Bacillus ex pneumoenteritide suis*, Klein. *Bacillus of the Smegma*. Found in the smegma of the male and female genitals. *Bacillus of Tommasoli*. Found on the hair in a case of sycosis. *Bacillus of Tricomi*. See *Bacillus of senile gangrene*. *Bacillus of Tuberculosis of Vines*. Regarded by Corvo as the cause of the diseased condition of vines usually attributed to *Phylloxera vitatrix*, Planchon. *Bacillus of Ulcerative Stomatitis of Cattle*, Lingard and Batt. Found on the skin, the mucous membranes, and in the lungs of young cattle. Cf. *Streptococcus of eczema epizootica*, Schottelius. *Bacillus of Ulcus Molle*, Kräfting. Identical with that described by Ducrey as the

Bacillus of Chancroid. *Bacillus* of Utpadel, Utpadel and Gessner. Obtained from the stuffing of the bed quilts in a military hospital (Augsburg), and from the intestinal contents of man. *aer.*, *nlif.*, *mot.*, *pg.* *Bacillus* of Vagus Pneumonia. See *Bacillus pneumonicus agilis*, Schou. *Bacillus* of Verruga Peruana, Izquierdo. Found in the intercellular spaces and blood-vessels in the nodules characteristic of Peruvian wart. *Bacillus* of Vignal. See *Bacillus buccalis minutus*, Vignal. *Bacillus* of Whooping Cough. See *Bacillus tussis convulsiva*, Afanassiew. *Bacillus* of Xerosis Epithelialis Conjunctivæ, Leber, Kuschbert, Fränkel, and Neisser. Syn., *Bacillus xerosis*, Fränkel; *Bacillus* of Colomiatti, *Bacillus* of xerosis, Schreiber. Found in the white, fatty scales of the conjunctivæ in cases of xerophthalmia. *nmot.* Does not grow on gelatin or potato. Pathogenesis not fully determined. *Bacillus* of Yellow Fever. Found in the mucous membrane of the small intestine in two cases of yellow fever. Sternberg has found many forms in the intestinal contents and viscera of yellow fever cadavers, the most frequent and abundant being the *Bacillus neapolitanus*, Emmerich; *Bacillus cadaveris*, Sternberg; and *Bacillus X*, Sternberg. The last is held by Sternberg as being possibly concerned in the etiology of yellow fever. *Bacillus* of Yellow Milk. See *Bacillus synanthus*, Ehrenberg and Schröter. *Bacillus* *ovatus* minutissimus, Unna. Found upon the skin in cases of eczema seborrheicum. *aer.*, *facanaer.*, *nlif.*, *npg.* *Bacillus oxytocus* perniciosus, Wyssokowitsch. Found in milk left standing a long time. *aer.*, *nlif.*, *pg.* *Bacillus panificans*, Laurent. Claimed as the active agent in the raising of bread. Found especially in black bread. Dunnenberger considers it a mere impurity and not concerned in the leavening processes. *Bacillus parvus* *ovatus*, Löffler. *Bacillus* of false hog cholera. Found in the carcass of a pig dead of a disease resembling rouget. It resembles *Bacillus septiciemia hemorrhagica*, Sternberg. *Bacillus pasteurianus*, Hansen. Syn., *Bacterium pasteurianum*, *Micrococcus pasteurianus*. Found in beer-wort and beer, especially that poor in alcohol. Resembles *Bacillus acetii*, Kützing and Sommer. *Bacillus pestifer*, Frankland. Found in the air. *aer.*, *liqf.*, *mot.*, *npg.* *Bacillus phosphorescens*, Fischer. Syn., *Bacillus phosphorescens indicus*, Fischer; *Photobacterium indicum*, Fischer. Found in sea water (Gulf of Mexico). *aer.*, *liqf.*, *mot.*, *phos.*, *npg.* *Bacillus phosphorescens gelidus*, Forster. See *Bacillus phosphoreus*, Cohn. *Bacillus phosphorescens indicus*, Fischer. See

Bacillus phosphorescens, Fischer. *Bacillus phosphorescens indigenus*, Fischer. Found in sea water (harbor of Kiel) and upon herring. *aer.*, *liqf.*, *mot.*, *phos.*, *npg.* *Bacillus phosphoreus*, Cohn. Syn., *Micrococcus phosphoreus*, Cohn; *Bacterium phosphorescens*, Hermes; *Bacillus phosphorescens gelidus*, Forster. Found on sea fish, raw and cooked (cooked salmon, Cohn). *aer.*, *nlif.*, *phos.*, *npg.* *Bacillus plicatus*, Zimmerman. Found in hydrant water (Chemnitz). *aer.*, *liqf.*, *nmot.*, *chg.* (grayish yellow) *npg.* *Bacillus pneumoniae*, Flügge. See *Bacillus pneumoniae friedländeri*. *Bacillus pneumoniae friedländeri*. Syn., *Bacillus pneumoniae*, Flügge; *Diplococcus pneumoniae fibrinosa*, Friedländer; *Micrococcus pneumoniae infectiosa*, Friedländer. Occasionally obtained from the exudates in the pulmonary alveoli in cases of croupous pneumonia. *aer.*, *facanaer.*, *nlif.*, *nmot.*, *pg.* *Bacillus pneumonicus agilis*, Schou. Syn., *Bacillus* of Schou; *Bacillus* of Vagus pneumonia. Obtained from rabbits suffering from pneumonia induced by section of the vagi. *aer.*, *liqf.*, *mot.*, *pg.* *Bacillus pneumosepticus*, Babes. Obtained from the blood and tissues of a person dead of septic pneumonia. *aer.*, *facanaer.*, *nlif.*, *nmot.*, *pg.* *Bacillus polymyxa*, Prazmowski. Syn., *Claustidium polymyxa*, Prazmowski. Found in infusion of potatoes, lupin seeds, etc. *aer.*, *facanaer.*, *npg.* *Bacillus polypiformis*, Liborius. Found in cow dung and in the exudates of mice inoculated with garden earth. *anaer.*, *nlif.*, *mot.*, *npg.* *Bacillus preussii*, Ciesielski. See *Bacillus alvei*, Cheshire and Cheyne. *Bacillus prodigiosus*, Ehrenberg. Syn., *Bacteridium prodigiosum*, Schröter; *Bacterium prodigiosum*, *Micrococcus prodigiosus*, Cohn; *Micrococcus inetrophus*, Trevisan; *Monas prodigiosa*, Ehrenberg; *Palmella prodigiosa*, Mont; *Zoogalactina inetropha*, Sette. Frequently found upon food-stuffs, boiled potatoes, hard-boiled eggs, moist bread, etc. *aer.*, *facanaer.*, *liqf.*, *mot.*, *chg.* (red), *npg.* Acts as a protective to rabbits against anthrax. *Bacillus pseudodema*, Liborius. Syn., *Bacillus pseudosepticus*, Macé; *Pseudo-adembacillus*, Ger. Obtained from the exudates of mice, after inoculation with garden earth. *anaer.*, *liqf.*, *pg.* Resembles *Bacillus adematidis maligni*, Koch. *Bacillus pseudopneumonicus*, Passet. Syn., *Bacterium pseudopneumonicum*. Found in pus. *aer.*, *nlif.*, *nmot.*, *pg.* Resembles *Micrococcus pneumoniae crouposa*, Fränkel. *Bacillus pseudosepticus*, Macé. See *Bacillus pseudodema*, Liborius. *Bacillus pseudosepticus* of Mice, Bienstock. Syn., *Bacillus* of *pseudosepticemia* of

mice. Found in feces. *aer.*, *nlif.*, *nmot.*, *fg.* Found after inoculation, mostly in the edematous fluid and not in the blood. **Bacillus pseudotuberculosis**, Pfeiffer. Obtained from the viscera of a horse killed on suspicion of having glanders. *aer.*, *nlif.*, *nmot.*, *fg.* **Bacillus puerperalis**, Engel and Spillmann. Leptothrix of puerperal fever; *Bacillus* of puerperal fever. Found in two cases of puerperal sepsis. *fg.* (for mice and rabbits). Considered by Pasteur to be identical with *Bacillus anthracis*. **Bacillus pulpæ pyogenes**, Miller. Obtained from gangrenous tooth-pulp. *aer.*, *facanaer.*, *liqf.*, *fg.* **Bacillus punctatus**, Zimmermann. Found in hydrant water (Chemnitz). *aer.*, *liqf.*, *npg.* **Bacillus putrificus coli**, Bienstock. Found in human feces. *aer.*, *facanaer.*, *nlif.*, *mot.*, *npg.* **Bacillus pyocyaneus** α , Gessard. Syn., *Bacillus aeruginosis*, *Bacterium aeruginosum*, Schröter; *Micrococcus pyocyaneus*, Gessard. *Bacillus* of blue or green pus. A widely distributed form, found in purulent and serous wounds, in perspiration, and in the viscera of human cadavers. *aer.*, *facanaer.*, *liqf.*, *mot.*, *chg.*, *fg.* Two pigments, one fluorescent green, the other blue, pyocyanin. **Bacillus pyocyaneus** β , Ernst. Found in pus from bandages stained green. *aer.*, *liqf.*, *mot.*, *chg.* (yellowish green), *npg.* **Bacillus pyogenes foetidus**, Passet. Syn., *bacille pyogène fétide*, Fr. Obtained from an ischio-rectal abscess. *aer.*, *nlif.*, *mot.*, *fg.* **Bacillus pyogenes soli**, Bolton. Obtained from the exudates of a rat after inoculation with garden earth. *aer.*, *facanaer.*, *nlif.*, *nmot.*, *npg.* **Bacillus radiatus**, Lüderitz. Obtained from the exudates of mice and guinea-pigs after inoculation with garden earth. *anaer.*, *liqf.*, *mot.*, *npg.* **Bacillus radiatus aquatilis**, Zimmermann. Found in hydrant water (Chemnitz). *aer.*, *liqf.*, *mot.*, *npg.* **Bacillus ramosus**, Eisenberg. Syn., *Wurzelbacillus*. Common in soil and water. *aer.*, *liqf.*, *npg.* **Bacillus ramosus liquefaciens**. Obtained from the air. *aer.*, *liqf.*, *mot.*, *npg.* **Bacillus reticularis**, Jordan. Found in hydrant water (Lawrence, Mass.). *aer.*, *liqf.*, *mot.*, *npg.* **Bacillus rheumarthritidis**, Kussmaul. Syn., *Bacillus rheumarthritidis*, *Micrococcus rheumarthritidis*, Leyden. Found in the joint effusions in articular rheumatism. **Bacillus rhinoscleromatis**, Cornil and Alvarez, Von Frisch, Paltauf, Von Eiselsberg, Dittich, Stepanow, etc. Found in the newly-formed tubercles of rhinoscleroma. *aer.*, *facanaer.*, *nlif.*, usually *nmot.* (becomes motile on cultivation). *fg.* Etiologic relations not established. Considered by many as identical with *Bacillus pneumoniae*

friedländeri. It is less virulent, gelatin cultures more transparent, and the capsules more persistent. **Bacillus rosaceum metalloides**, Dowdeswell. **Bacillus rubefaciens**, Zimmermann. Found in hydrant water (Chemnitz). *aer.*, *nlif.*, *mot.*, *chg.* (pale pink), *npg.* **Bacillus rubellus**, Okada. Found in the exudates of guinea-pigs after inoculation with street dust. *anaer.*, *liqf.*, *mot.*, *chg.* (red), *npg.* **Bacillus ruber**, Frank, Cohn, and Becker. Syn., *Bacillus miniacus*, Zimmermann. *Bacillus* (or *Bacterium*) *rosaceum metalloides*, Dowdeswell. The red bacillus of water. Found in water and on boiled rice. *aer.*, *liqf.*, *mot.*, *chg.* (magenta red), *npg.* **Bacillus ruber indicus**. See *Bacillus indicus*, Koch. **Bacillus rubescens**, Jordan. Found in sewage (Lawrence, Mass.). *aer.*, *nlif.*, *mot.*, *chg.* (pale pink), *npg.* **Bacillus rubidus**, Eisenberg. Found in water. *aer.*, *liqf.*, *mot.*, *chg.* (brownish red), *npg.* (Lustig describes a red bacillus from river water which he claims to be different from this). **Bacillus salivarius septicus**, Biondi. See *Bacillus septicus sputigenus*, Flügge. **Bacillus sanguinis typhi**, Brannan and Cheesman. Obtained from the blood of typhus-fever patients. *aer.*, *facanaer.*, *nmot.* **Bacillus saprogenes**, I, II, III, Rosenbach. Found in putrefying matter on fetid feet, the white plugs of the pharyngeal follicles, etc. *aer.*, *facanaer.*, *fg.* **Bacillus scaber**, Duclaux. Syn., *Tyrophthrix scaber*. Found in cheese. *aer.*, *nlif.*, *mot.*, (in early stages becoming non-motile), *npg.* **Bacillus schafferi**, Freudenreich. Obtained from cheese and fermenting potato infusion. *aer.*, *facanaer.*, *nlif.*, *mot.*, *npg.* Closely resembles *Bacillus neapolitanus*, Emmerich. **Bacillus scissus**, Frankland. Found in the soil. *aer.*, *nlif.*, *nmot.*, *npg.* **Bacillus septicæmiæ hæmorrhagicæ**, Sternberg. *Bacillus cholera gallinarum*, Flügge. *Bacillus cuniculicida*, Flügge. *Bacillus cuniculicidus*, Koch. *Bacterium septicæmiæ*, Koch. *Bacterium morbilli*, Lanzi. *Coccobacteria septicæ*, Billroth. *Micrococcus septicus*, Cohn. *Micrococcus cholera gallinarum*, *Micrococcus gallicidus*, *Microsporon septicus*, Klebs. **Bacillus septicus**, Koch. See *Bacillus erysipelatos suis*, Koch. **Bacillus septicus**, Pasteur. See *Bacillus adenatis maligni*, Koch. Cf. *Proteus septicus*. **Bacillus septicus acuminatus**, Babes. Obtained from the umbilical stump, blood and viscera of a five days old child, dead of septic infection. *aer.*, no growth on gelatin, *fg.* (for rabbits and guinea-pigs, not for mice). **Bacillus septicus agrigenus**, Nicolaier. Obtained from manured garden soil. *aer.*, *nlif.* (?) Resembles *Bacillus septicæmiæ hæmorrhagicæ*, Sternberg. **Bacillus septicus** from

Sputum. See *Bacillus septicus sputi*, I, II, Kreibohm. *Bacillus septicus keratomalaciae*, Babes. Obtained from the corneal tissues and viscera of a child that died of septicemia following keratomalacia. aer., facanaer., nliqf., pg. (for rabbits and mice, slightly for birds, not for guinea-pigs). *Bacillus septicus sputi*, I, II, Kreibohm. Obtained from human buccal secretions. Does not grow in any known culture-medium. Resembles *Bacillus septicaemiae haemorrhagica*, Sternberg. *Bacillus septicus sputigenus*, Flügge. Syn., *Bacterium septicum sputigenum*, Fränkel. *Micrococcus pneumoniae crouposa*, Fränkel. *Micrococcus pasteurii*, Sternberg. *Diplococcus pneumoniae*, Weichselbaum. *Streptococcus lanceolatus pasteurii*, Gamaleja. *Micrococcus* of sputum septicemia, Fränkel. Lancet-shaped micrococcus, Talamon. *Pneumococcus* of Fränkel. Found both in healthy and in pneumonic sputum, in the fibrinous exudate of croupous pneumonia, and in the pus of meningitis, in middle-ear disease, in certain forms of pleurisy, joint-disease, and endocarditis. aer., facanaer., nmot., nliqf., pg. Held to be the cause of croupous pneumonia in man, and by some to be identical with *Bacillus lysae*, Pasteur. *Bacillus septicus ulceris gangrenosi*, Babes. Found in the blood and viscera of a boy dead of septicemia following gangrene of the skin. aer., liqf., mot., pg. *Bacillus septicus vesicae*, Clado. Found in the urine of a patient with cystitis. aer., facanaer., nliqf., mot., pg. *Bacillus aessilis*, L. Klein. Found in the blood of a cow supposed to have died of anthrax. aer., mot. (convulsive jerking), npg. *Bacillus similis*, Bienstock. See *Bacillus subtilis simulans*, I, II, Bienstock. *Bacillus smaragdinus foetidus*, Reimann. Found in the nasal secretions in a case of ozena. aer., facanaer., liqf., pg. *Bacillus smaragdino-phosphorescens*, Katz. Obtained from a herring in fish market (Sidney, N. S. W.). aer., nliqf., nmot., phos. (emerald green) npg. Resembles *Photobacterium phosphorescens*, Cohn, and *Photobacterium pfügeri*, Ludwig. *Bacillus solidus*, Lüderitz. Found in the exudates of mice inoculated with garden earth. aer., nliqf., mot., npg. *Bacillus spiniferus*, Unna. Found upon the skin in cases of eczema seborrhaicum. aer., nliqf., chg. (grayish yellow), npg. *Bacillus sputigenus crassus*. See *Bacillus crassus sputigenus*, Kreibohm. *Bacillus stolonatus*, Adametz. Found in water. aer., nliqf., mot., npg. *Bacillus stoloniferus*, Pohl. Found in swamp water. aer., liqf., mot. Pathogenesis not determined. *Bacillus striatus albus*, Von Besser. Found in the nasal secretions of healthy persons. aer., nliqf.,

npg. *Bacillus striatus flavus*, Von Besser. Occasionally found in nasal mucus. aer., nliqf., chg. (sulphur yellow), npg. *Bacillus subflavus*, Zimmermann. Found in hydrant water (Chemnitz). aer., nliqf., mot., chg. (pale yellow), npg. *Bacillus subtilis*, Ehrenberg, Cohn, Brefeld, Prazmowski, and Fitz. Syn. *Bacillus glycerina*, Fitz; *Bacillus fitzianus*, Zopf; *Bacterium subtile*, Zopf; *Metallacter subtile*, Pertz; *Vibrio subtilis*, Ehrenberg. DeBary confines *B. subtilis* to the form described by Brefeld and Prazmowski. The relation of the starch fermentation species of Fitz remains doubtful. A common form found in the air, water, soil, and on plants. aer., liqf., oscl., npg. *Bacillus subtiliformis*, Bienstock. Syn., *Bacillus subtilis simulans*, I, II, Bienstock. *Bacillus similis*, Bienstock. Found in human feces. aer., nliqf., nmot., npg. *Bacillus suis*, Detmers. See *Bacillus tardi gradus*. *Bacillus sulphhydrogenus*, Miquel. Found in water. aer., mot. Decomposes albumin with evolution of H_2S . *Bacillus sulfureum*, Holschewnikoff and Rosenheim. Two forms found in urine and in mud. *Bacillus superficialis*, Jordan. Common in sewage. aer., liqf., mot., npg. *Bacillus synxanthus*, Ehrenberg and Schröter. Syn., *Bacillus xanthogenus*, Fuchs and Ehrenberg. *Bacterium xanthinum*, Schröter. *Vibrio synxanthus*, Ehrenberg. *Vibrio xanthogenus*, Fuchs. Bacillus of yellow milk. Found in milk that has been boiled. aer., mot., chg. (golden yellow), npg. Precipitates the casein; renders the milk alkaline. *Bacillus syphilidis*, Lustgarten. Syn., *Bacillus syphilitis*, Lustgarten. Bacillus of Lustgarten. Found in syphilitic new growths and secretions; does not develop in ordinary culture media. In staining, reaction seems to be identical with that of the Bacillus of the Sinergma. Douhle describes parasitic infusoria as the causative agents of syphilis. *Bacillus tardi gradus*. Syn., *Bacillus* var., Detmers. *Micrococcus suis*, Burriell. Found in dew, water, and in the fluids of pigs affected with swine plague. aer., mot., npg. Cf. *Bacillus of swine plague*, Marcellus, Pouch and Jolbert. *Bacillus tenuis*, Dublanz. Syn., *Tyrophix tenuis*, Dublanz. Found in the decomposing albuminous matter, cheese, etc. aer., facanaer., nliqf., mot., npg. Used in ripening certain French cheeses. *Bacillus tenuis sputigenus*, Puzos. Observed from sputum. aer., nliqf., nmot., pg. *Bacillus termo*, Dupathé and Flandre. Syn., *Bacterium termo*, Cohn. *Bacillus termophilus*, Flandre. Syn., *Bacillus termo termophilus*, Rosenbach. *Bacterium termo*. Found in garden earth and in the exudate of pneumonia.

suffering from tetanus. Spores form at one end and give bacillus a drum-stick appearance. *anaer.*, *liqf.*, *mot.*, *pg.* *Bacillus tetani traumatici*. See the preceding. *Bacillus thalassophilus*, Russel. Obtained from mud (Bay of Naples). *anaer.*, *liqf.*, *mot.* Pathogenesis not determined. *Bacillus thermophilus*, Miquel. Found in the intestinal tract of man and animals, in sewage, and in the soil. *aer.*, *nmot.*, *npg.* *Bacillus tremelloides*, Schottelius. Found in hydrant water (Freiburg). *aer.*, *liqf.*, *chg.* (golden yellow), *npg.* *Bacillus tremulus*, Koch. Syn., *Metabacter tremulus*, Trevisan. Found as a surface pellicle on decomposing plant infusions. Resembles *Bacillus subtilis*, Ehrenberg. *Bacillus tuberculosis*, Koch. Syn., *Bacterium tuberculosis*, Zopf; *Bacterium tuberculosis*, *Bacillus tuberculi*, *Bacillus tuberculosis*. Found in the sputum, in tuberculous organs, and elsewhere in persons and animals affected with tuberculosis. Parasitic. *aer.*, *facanaer.*, *nmot.*, *pg.* *Bacillus tuberculosis gallinarum*, Maffucci. Obtained from cases of spontaneous tuberculosis in fowls. This seems to be a distinct species, characterized by its staining reactions, etc., its growth in pure cultures, and its pathogenic qualities, Koch. *Bacillus tumescens*, Zopf. Syn., *Bacterium tumescens*, Zopf. Found upon beets. *aer.*, *liqf.*, *mot.*, *npg.* *Bacillus turgidus*, Duclaux. Syn., *Tyrophrix turgidus*, Duclaux. Found in cheese. *aer.* Produces alkaline reaction (carbonate and butyrate of ammonium) in milk. *Bacillus tussis convulsivæ*, Afanassiev. Syn., *Bacillus of Afanassiev*. Found in the sputum of persons affected with pertussis. *aer.*, *nliqf.*, *mot.*, *pg.* *Bacillus typhi abdominalis*, Eberth, Gaffky, Klebs, Espinger, Brautlecht. Syn., *Bacillus typhosus*, *Bacterium typhosum*. Found in water and in milk, and in the feces and organs in typhoid fever. *aer.*, *facanaer.*, *nliqf.*, *mot.* *Bacillus typhi murium*, Löffler. Obtained from mice affected with an epidemic disease. *aer.*, *facanaer.*, *nliqf.*, *mot.* *Bacillus typhosus*. See *Bacillus typhi abdominalis*, Eberth, etc. *Bacillus ubiquitus*, Jordan. Found in air, water, sewage (Lawrence, Mass.). *aer.*, *facanaer.*, *nliqf.*, *nmot.*, *npg.* Resembles *Bacillus candicans*, Frankland. *Bacillus ulna*, Cohn, and Prazmowski. Syn., *Vibrio b*, Ehrenberg. *Bacterium ulna*, Miller. *Vibrio bacillus*, Müller, Ehrenberg. Found in egg albumin. *aer.*, *mot.*, *npg.* *Bacillus ulna* of Vignal. Found in saliva of healthy persons; perhaps identical with the preceding form. *aer.*, *liqf.*, *npg.* *Bacillus ureæ*, Miquel. Found in the air. *aer.*, *facanaer.*, *nliqf.*, *npg.* Perhaps identical with the preceding form. *Bacillus ureæ*, Leube.

Three varieties found in old urine. *aer.*, *nliqf.*, *npg.* Converts urea into ammonium carbonate. *Bacillus urocephalus*, Duclaux. Syn., *Tyrophrix urocephalus*, Duclaux. One of the commonest forms associated with putrefaction of animal matter. *aer.*, *facanaer.*, *npg.* *Bacillus vasculosis*, Sternberg. Obtained from the intestine and stomach of yellow-fever cadavers. *aer.*, *liqf.*, *mot.*, *npg.* *Bacillus vaginalis*, Döderlein. Constantly found in the normal vaginal secretions of adults. *aer.*, *facanaer.*, *nmot.*, (no growth in gelatin), *npg.* *Bacillus varicosus conjunctivæ*, Gombert. Found in the healthy conjunctival sac of man. *aer.*, *facanaer.*, *nliqf.*, *mot.*, *pg.* *Bacillus venenosus*, Vaughan. Found in water. *aer.*, *facanaer.*, *nliqf.*, *mot.*, *pg.* *Bacillus venenosus brevis*, Vaughan. Found in water. *aer.*, *facanaer.*, *nliqf.*, *mot.*, *pg.* *Bacillus venenosus invisibilis*, Vaughan. Found in water. *aer.*, *facanaer.*, *nliqf.*, *mot.*, *pg.* *Bacillus venenosus liquefaciens*, Vaughan. Found in water. *aer.*, *facanaer.*, *liqf.*, *mot.*, *pg.* *Bacillus ventriculi*, Raczynsky. Obtained from the stomach of meat-fed dogs. *aer.*, *facanaer.*, *nliqf.*, *mot.*, *npg.* *Bacillus vermicularis*, Frankland. Found in river water. *aer.*, *liqf.*, *oscl.*, *npg.* *Bacillus vermiculosus*, Zimmermann. Found in water. *aer.*, *liqf.*, *oscl.*, *npg.* *Bacillus violaceus*, Becker. Found in river water. *aer.*, *facanaer.*, *liqf.*, *mot.*, *chg.* (dark violet). *Bacillus violaceus laurentius*, Jordan. Found in hydrant water (Lawrence, Mass.). *aer.*, *facanaer.*, *liqf.*, *mot.*, *npg.* *Bacillus virens*, Van Tieghem. Syn., *Sporonema gracile*. Found in stagnant water. *aer.*, *mot.*, *chg.* (green, held by some to be chlorophyll, but not sufficiently studied), *npg.* *Bacillus virescens*, Frick. Found in green sputum. *aer.*, *nliqf.*, *mot.*, *chg.* (green), *npg.* *Bacillus virgula*, Duclaux. Syn., *Tyrophrix virgula*, Duclaux. Found in albuminous solutions. *aer.*, *nmot.*, *npg.* *Bacillus virgula*. See *Spirillum cholerae asiaticæ*, Koch. *Bacillus viridis*, Van Tieghem. Syn., *Bacterium viride*, Van Tieghem, *Arthrobacterium viride*, De Bary. Found in water collected in the hollow of a polyporous fungus. *aer.*, *chg.* (green). Not thoroughly investigated. *Bacillus viridis flavus*, Frick. Syn., *Bacillus viridis pallescens*, Frick. *aer.*, *nliqf.*, *mot.*, *chg.* (yellowish-green), *npg.* Resembles *Bacillus virescens*, Frick. *Bacillus viridis pallescens*, Frick. See *Bacillus viridis flavus*, Frick. *Bacillus viscosus*, Frankland. Found in river water. *aer.*, *liqf.*, *mot.*, *chg.* (fluorescent green). Perhaps identical with *Bacillus fluorescens*, Flügge. *Bacillus viscosus*, I, II, Van

Laer. Found in ropy milk. *aer.*, *nlqf.*, *npg.* *Bacillus viscosus cerevisiæ*, Van Laer. Found in viscous beer and in milk. *aer.*, *liqf.*, *npg.* *Bacillus viscosus sacchari*, Kramer. Found in viscous saccharine fluids. *aer.*, *liqf.*, *npg.* *Bacillus vulgaris*, Hauser; Syn. *Proteus vulgaris*. *Proteus sulfureus*, Holschenikoff. *Proteus* of Hauser. Commonly associated with putrefaction. *aer.*, *faeciaer.*, *liqf.*, *mot.*, *pg.* This species is probably one of those formerly included under *Bacterium termo*. *Bacillus X*, Sternberg. Found in yellow-fever cadavers. Sternberg thinks it possible that this form is concerned in the etiology of the fever. *pg.* (for rabbits). *Bacillus xanthogenus*, Fuchs and Ehrenberg. See *Bacillus synxanthus*, Ehrenberg. *Bacillus xerosis*, Fränkel. Syn. *Xerose bacillus*. See *Bacillus* of xerosis epithelialis conjunctivæ, Leber. *Bacillus zenkeri*, Hauser. Syn., *Proteus zenkeri*. Bacille de Hauser, Fr. Found in putrefying animal matter. *aer.*, *faeciaer.*, *nlqf.*, *mot.*, *pg.* *Bacillus zoffii*. Syn., *Bacterium zoffii*, Kurth. *Arthrobacterium zoffii*, De Bary. Found in the intestinal tract of fowls. *aer.*, *nlqf.*, *mot.*, *npg.* *Bacillus zurianum*, List. Found in water. *aer.*, *nlqf.*, *nmot.*, *npg.* *Bacteridium prodigiosum*, Schröter. See *Bacillus prodigiosus*, Ehrenberg. *Bacterium aceti*, Lanzi. See *Bacillus aceti*, Kützing. *Bacterium aerogenes*, Miller. See *Bacillus aerogenes*, II, Miller. *Bacterium aeruginosum*, Schröter. See *Bacillus pyocyaneus* a, Gessard. *Bacterium allii*. See *Bacillus allii*, Griffiths. *Bacterium anthracis*, Zopf. See *Bacillus anthracis*, Pollender and Davaine. *Bacterium articulatum*, Ehrenberg. See *Bacterium lineola*, Cohn. *Bacterium aurantiacum*, Trelease; a chromogenic form. *Bacterium bacillus*, Pasteur. See *Bacterium termo*, Davaine. *Bacterium brunneum*, Schröter. Found in a putrid infusion of Indian corn, *chg.* (brown). Perhaps identical with *Bacillus fuscus*, Zimmerman. *Bacterium butyri colloideum*, Lafar. Found in every sample of natural butter examined. Cf. *Bacillus butyri fluorescens*, Lafar. *Bacterium candidum*, Trelease. A chromogenous form, identical with *Bacillus merentericus fuscus*, Flügge. *Bacterium capitatum*, Davaine. Found in an infusion of albuminous matter. *aer.*, *mot.* *Bacterium carbuncolare*, Pollender, Brauell, Delafond, and Davaine. See *Bacillus anthracis*, Pollender and Davaine. *Bacterium carlsbergense*, Hansen. Found in the air. Cf. *Bacterium kochii*, Hansen. *Bacterium catenula*, Dujardin. Has been observed especially in putrid wine or putrefying blood, generally in albuminous liquids

in contact with air. Coze and Feltz found it in typhoid fever in man. It resembles *Bacillus acidi lactici*, Davaine, Hueppe. *Bacterium caucasicum*. See *Bacillus caucasicus*, Kern. *Bacterium chlorinum*, Engelmann. Found in water. *aer.*, *chg.* (green). Engelmann holds this to be chlorophyll, as a small amount of oxygen is given off, indicating the assimilation of carbon. *Bacterium coli commune*, Escherich. See *Bacillus neapolitanus*. *Bacterium commune*, Pasteur. See *Bacterium termo*, Dujardin. *Bacterium cuneatum*, Rivolta. See *Bacillus cuneatus*. *Bacterium cyanogenum*, Fuchs. See *Bacillus cyanogenus*, Hueppe. *Bacterium denitrificans*. See *Bacillus denitrificans*. *Bacterium dysodes*, Zopf. See *Bacillus dysodes*. *Bacterium enchelys*, Ehrenberg. Found in river water. *Bacterium farinaceum*, Wigand. Found in sour sponge or dough. *Bacterium fitzianum*, Zopf. See *Bacillus subtilis*, Ehrenberg. *Bacterium fætidum*, Thin. See *Bacillus fætidus*, Passet. *Bacterium fusiforme*, Warming. Found in sea water. *Bacterium gingivæ pyogenes*, Miller. See *Bacillus gingivæ pyogenes*. *Bacterium gliscrogenum*, Malerba. Syn. *Bacillus gliscrogenus*. Found in viscid acid urine. *aer.*, *nlqf.*, *mot.*, *npg.* *Bacterium griseum*, Warming. Syn. *Micrococcus griseus*, Winter, a doubtful form. *Bacterium gummi*, Commes. The cause of the gumous disease of the fig, almond, and orange trees, and the *mal nero* of vines. Cf. *Bacterium putredinis*, Davaine. *Bacterium gummosum*, Ritsert. Found to produce the viscosity of infusions of digitalis. *aer.*, *liqf.*, *pleom.* *Bacterium hessii*, Guillebeau. One of the two forms found by Guillebeau to cause milk to become ropy. Thrives best in pasteurized milk. *aer.*, *liqf.*, *mot.*, *npg.* Cf. *Micrococcus freudenreichii*. *Bacterium hyacinthi*, Wakker. Found in the bulbs and leaves of the hyacinth and causing the "yellows" of hyacinths. Cf. *Bacillus hyacinthi septicus*, Heinz. *Bacterium janthinum*, Zopf. See *Bacillus janthinus*, Zopf. *Bacterium kochii*, Hansen. Found in the air of Carlsberg. Cf. *Bacterium carlsbergense*, Hansen. *Bacterium lactis*, Lister. See *Bacillus acidi lactici*, Hueppe. *Bacterium lactis aerogenes*, Escherich. See *Bacillus lactis aerogenes*, Abeloos. *Bacterium lineola*, Müller, Cohn. Syn. *Bacterium punctum*, Rivolta. *Bacterium tremulans*, Trevisan. *Bacterium trilobulare*, Ehrenberg. *Bacterium articulatum*, Ehrenberg. *Vibrio lineola* (Müller), Ehrenberg. *Vibrio tremulans*, Ehrenberg. *Bacillus lineola*. Found in well and other water, in infusions, in soil on vegetables, etc. *aer.*, *mot.*

(trembling and darting back and forth). *pleom.*, *npg.* Probably represents several species, or a developmental form of a spirillum. **Bacterium litoreum**, Warming. Found only in sea water. *aer.*, *mot.*, *npg.* **Bacterium lucens**, Van Tieghem. See *Bacillus lucens*, Van T. **Bacterium luteum**, Trelease. See *Bacillus luteus*, Flügge. **Bacterium maidis**, Cuboni. See *Bacillus maidis*, Cuboni. **Bacterium merismopædioides**, Zopf. See *Bacillus merismopædioides*, Zopf. **Bacterium morbilli**, Lanzi. Found in the urine of persons with measles. Cf. *Bacillus septiciemise hæmorrhagica*, Sternberg. **Bacterium navicula**, Reinke and Berthold. See *Bacillus butyricus*, Prazmowski. **Bacterium neapolitanum**. See *Bacillus neapolitanus*, Emmerich. **Bacterium oblongum**, BOUTROUX. Syn., *Micrococcus oblongus*, BOUTROUX. Found in beer; the agent of gluconic fermentation. **Bacterium of Butyric Acid Fermentation**. See *Bacillus butyricus*, Prazmowski. **Bacterium of Diphtheria**, Emmerich. Found in cases of diphtheria in man and doves. **Bacterium of Fire Blight**. See *Bacillus butyricus*, Prazmowski. **Bacterium of Infectious Pneumonia in the Horse**. See *Streptococcus corysæ contagiosæ equorum*, Schütz. **Bacterium of Kefir**. See *Bacillus caucasicus*, Kern. **Bacterium of Pear Blight**. Cf. *Bacillus butyricus*, Prazmowski, and *Micrococcus amylocorus*, Burrill. **Bacterium of Sheep Pox**. Found in cases of sheep pox, or "Schafblattern." **Bacterium of Variola**, Cose and Feltz. Found in the vesicles of smallpox; pathogenic for rabbits. **Bacterium oleæ**, Arcangeli. Said to cause the so-called tuberculosis ("Maladie de la loupe;" "Rogner") of olives. **Bacterium ovatum**, Lebert. See *Nosema bombycis*, Nägeli. **Bacterium pasteurianum**, or **Bacterium pastorianum**. See *Bacillus pasteurianus*, Hansen. **Bacterium periplanetæ**, Tichomirow. Found in the common cockroach (*Periplaneta orientalis*), and producing in it a specific disease. **Bacterium pflügeri**, Ludwig. See *Photobacterium pflügeri*, Ludwig. **Bacterium phosphorescens**, Hermes. **Bacterium phosphorescens**, Cohn. See *Photobacterium phosphorescens*. **Bacterium photometricum**, Engelmann. Found in water; motion dependent on light; possibly not a bacterium at all. **Bacterium pneumoniae**, **Bacterium pneumoniae crouposæ**, Friedländer. See *Bacillus pneumoniae friedländeri*. **Bacterium porri**, Tommasi-Crudeli. Found in warts. **Bacterium prodigiosum**. See *Bacillus prodigiosus*,

Ehrenberg. **Bacterium pseudopneumonicum**. See *Bacillus pseudopneumonicus*, Flügge. **Bacterium punctum**, Rivolta. See *Bacterium lineola* (Müller), Cohn. **Bacterium putredinis**, Davaine. Found in rapidly-decaying plants. Cf. *Bacterium gummi*, Comes. **Bacterium pyri-forme**, Hansen. Found in the air. **Bacterium radicola**, Prazmowski. Syn., *Bacillus radicola*, Beyernick; *Rhizobium leguminosarum*, Frank. Found in the soil, particularly in the roots of the Leguminosæ, where it is held to stimulate the cells to unusual growth, affect nitrification, constitute the "bacteroids" of Woronin, and form the so-called "bacteroid tissue." **Bacterium rosaceum metalloides**, Dowdeswell. See *Bacillus ruber*, Frank, Cohn. **Bacterium rubescens**, Lankester. See *Beggiatoa roseopersicina*, Zopf. **Bacterium septiciemise**, Koch. See *Bacillus septiciemise hæmorrhagica*, Sternberg. **Bacterium septicum sputigenum**. See *Bacillus septicus sputigenus*, Flügge. **Bacterium subtile**, Buchner. See *Bacillus subtilis*, Ehrenberg. **Bacterium sulfuratum**; **Bacterium sulphuratum**, Warming. See *Beggiatoa roseopersicina*, Zopf. **Bacterium sulfureum**, Rosenheim. Found in wine. *nlqf.* Evolves H₂S. **Bacterium syncyanum**, Ehrenberg, Schröter. See *Bacillus cyanogenus*, Hueppe. **Bacterium synxanthum**, Schröter. See *Bacillus synxanthus*. **Bacterium termo**, Dujardin and Ehrenberg. Syn., *Bacillus termo*, Davaine. **Bacterium bacillus**, Pasteur. *Cryptococcus natans*, Kützing. *Cryptococcus nebulosus*, Kützing. *Micrococcus crepusculum*, Cohn. *Zoogloa termo*, Monas *termo* (Müller). A collective species. **Bacterium termo**, Vignal. Found in the saliva of healthy persons. *aer.*, *liqf.*, *mot.*, *chg.* (yellowish gray), *npg.* **Bacterium tholoideum**, Gessner. Syn., *Bacillus of Gessner*. Found in the intestinal contents of healthy persons. Resembles *Bacillus lactis aerogenes*, Escherich. *pg.* (for mice and guinea-pigs). **Bacterium tremulans**, Trevisan. See *Bacterium lineola*, Cohn. **Bacterium triloculare**, Ehrenberg. See *Bacterium lineola*, Cohn. **Bacterium tuberculosis**, Zopf. **Bacterium tuberculosis**, Koch. See *Bacillus tuberculosis*, Koch. **Bacterium tumescens**, Zopf. See *Bacillus tumescens*, Zopf. **Bacterium typhosum**. See *Bacillus typhi abdominalis*, Eberth and Gaffky. **Bacterium ulna**. See *Bacillus ulna*, Cohn. **Bacterium ulna**, Cohn. See *Micrococcus ureæ*. **Bacterium ureæ**, Jaksch. Found in ammoniacal urine. *aer.*, *fasanaer.*, *nlqf.* Not sufficiently investigated. **Bacterium vermiforme**, Ward. Found in fermented ginger-beer. **Bacterium viola-**

ceum, Bergonzini. Syn., *Chromobacterium violaceum*, Bergonzini. Found in putrefying solutions of egg-albumen. *chg.* (violet). *Bacterium viride*. See *Bacillus viridis*, Van Tieghem. *Bacterium xanthinum*. See *Bacillus synxanthus*, Ehrenberg and Schröter. *Bacterium xylinum*, A. J. Brown. Found in solutions of carbohydrates, giving rise to acetic acid, and converting dextrose into gluconic acid, and mannitol into levulose. *Bacterium zopfii*, Kurth. See *Bacillus zopfii*. *Bacteroides*, Woronin. Microorganisms forming tubercles in the roots of leguminous plants, and assisting in the fixing of nitrogen. *Beggiatoa alba*, Vaucher. Syn., *Beggiatoa punctata*, Trevisan. *Beggiatoa leptomitiformis*, Meneghi. *Oscillaria alba*, Vaucher. *Hygrocrocis vandelli*, Meneghi. Next to *Cladothrix* the most common bacterium of water, forming a superficial layer in the mud. *Beggiatoa alba*, var. *marina*, Cohn. Syn., *Beggiatoa ærstedtii*, Rabenhorst. See *Beggiatoa marina*, Cohn. *Beggiatoa arachnoidæ*, Agardh. Syn., *Oscillaria arachnoidæ*, Agardh. *Oscillaria versatilis*, Kütz. Found in swamps and sulphur springs. *Beggiatoa leptomitiformis*, Meneghi and Trevisan. Syn., *Oscillaria leptomitiformis*, Meneghi. Found in sulphur springs. *Beggiatoa marina*, Cohn. Syn., *Beggiatoa alba*, var. *marina*, Cohn. *Beggiatoa ærstedtii*, Rabenhorst. Found in a salt water aquarium, forming a whitish slimy layer on dead fish. *Beggiatoa minima*, Warming. Found in sea water; gray. *Beggiatoa mirabilis*, Cohn. Found in a salt water aquarium; snow white. *Beggiatoa multisepta*. See *Pragmidiothrix multisepta*, Engler. Held by Zopf to be identical with *Beggiatoa alba*, Vaucher. *Beggiatoa nivea*, Rabenhorst. Syn., *Leptomenia niveum*, *Symphiothrix nivea*, Bruggen. Found in sulphur springs. *Beggiatoa nodosa*, Van Tieghem. A doubtful form characterized by nodes in the filaments, which differ from other members of the genus in the absence of sulphur grains and of motion. *Beggiatoa ærstedtii*, Rabenhorst. See *Beggiatoa alba*, var. *marina*, Cohn. *Beggiatoa pellucida*, Cohn. Found in a marine aquarium; sulphur grains few. *Beggiatoa punctata*, Trevisan. See *Beggiatoa alba*, Cohn. *Beggiatoa roseopersicina*, Zopf. Syn., *Bacterium rubescens*, Lankester. *Bacterium sulfuratum* or *Bacterium sulphuratum*, Warming. *Clathrocystis roseopersicina*, Cohn. *Cohnia roseopersicina*, Winter, Kützing. *Microhaloa rosea*, Kütz. *Ophidiomonas sanguinea*, Ehrenberg. *Pleurococcus roseopersicina*, Rabenhorst. Peach-colored bacterium. Morphologically identical with *Beggiatoa alba*, Vaucher. Forms rose-colored to violet pellicles on putrid matter. Produces

bacteriopurpurin. *Beggiatoa tigrina*, Rabenhorst. Syn., *Oscillaria tigrina*, Roemer. Found in swamps on submerged wood. Forms a thin white layer. *Bienstock's Putrefactive Bacillus*. *Bienstock's Putrefactive Bacillus* from the Feces. See *Bacillus putreficus coli*, Bienstock. *Brieger's Bacillus*. See *Bacillus cavidus*, Brieger. *Buff Bacillus of Spreading Edema*, A. B. Harris. *nlif.*, *aer.*, *chg.* (buff), *pleom.* *Capsule Bacillus* of Loeb. Obtained from a case of keratomalacia infantum. *aer.*, *facanaer.*, *nlif.*, *nmot.*, *pg.* Resembles *Bacillus capsulatus*, Pfeiffer. *Capsule Bacillus of Ozena*, Marano. Found in the nasal secretions in ten cases of ozena. Resembles *Bacillus pneumoniae friedländeri*. Cf. *Micrococcus lawenbergii*. *Capsule Bacillus* of Smith. See *Bacillus capsulatus smithii*. *Cheese Spirillum*. See *Spirillum tyrogenum*, Denecke. *Chromobacterium violaceum*, Bergonzini. See *Bacterium violaceum*, Bergonzini. *Cladothrix bovis*. See *Actinomyces bovis*, Harz. *Cladothrix canis*. Held by Rabe to be very like *Actinomyces bovis*. *Cladothrix dichotoma*, Cohn and Zopf. See *Leptothrix ochracea*, Kützing. *Myconostoc gregarium*, Cohn. *Cladothrix fæsteri*, Cohn. Syn., *Crenothrix fæsteri*, *Streptothrix fæsteri*, Cohn. Forms yellowish or grayish concretions in the human lacrymal ducts. *Cladothrix intricata*, Russel. Obtained from sea mud (bay of Naples). *aer.*, *liqf.*, *mot.* *Clathrocystis roseopersicina*, Cohn. See *Beggiatoa roseopersicina*, Zopf. *Clostridium butyricum*, Prazmowski. See *Bacillus butyricus*, Prazmowski. *Clostridium fœtidum*, Liborius. See *Bacillus fœtidus*, Passet. *Clostridium of Symptomatic Anthrax* (or *Charbon*), Neelsen and Ehlers. See *Bacillus chauvæi*, Bollinger and Feser. *Clostridium polymyxa*, Prazmowski. See *Bacillus polymyxa*, Prazmowski. *Coccobacillus pyogenes ureæ*, Rösing. See *Diplococcus pyogenes ureæ*. *Coccobacteria septica*, Billroth. The single term under which Billroth includes all putrefactive bacteria. See *Bacillus septiciemia hæmorrhagica*. *Coccobacteria ureæ*, Rösing. See *Diplococcus pyogenes ureæ*. *Coccobacterium lyssæ*, Rivolta. See *Bacillus lyssæ*. *Cohnia roseopersicina*, Winter and Kützing. See *Beggiatoa roseopersicina*, Zopf. *Colon Bacillus* of Escherich. See *Bacillus neapolitanus*. *Comma Bacillus*, *Comma Bacillus* of Koch, *Comma-shaped Bacillus*, Waten. See *Spirillum cholerae asiaticæ*, Koch. *Corpuscles brillants*. See *Bacillus fœtidus*, Passet. *Coryza diplococci*. See *Micrococcus cereus aureus*, Schröter and Winckler. *Crenothrix fæsteri*, Cohn. See

Cladothrix farsteri. *Crenothrix kühniana*, Saccardo and Rabenhorst. Syn., *Crenothrix polyspora*, Cohn. *Hypoethrix kühniana*, *Leptothrix kühniana*. Found in drinking water, to which it imparts a disagreeable taste. *Crenothrix polyspora*, Cohn. See *Crenothrix kühniana*, Saccardo and Rabenhorst. *Cryptococcus alveareus*. See *Bacillus alvei*, Cheshire and Cheyne. *Cryptococcus clava*, Wedl. Found in the stomachs of ruminants. *Cryptococcus cerevisiæ*, Robin. *Cryptococcus fermentatum*, Kützing. See *Saccharomyces cerevisiæ*, Reess and Robin. *Cryptococcus glutinis*, Fersen. See *Saccharomyces glutinis*. *Cryptococcus guttulatus*, Robin. See *Saccharomyces guttulatus*. *Cryptococcus natans*, *Cryptococcus nebulosus*, Kützing. See *Bacterium termo*, Dujardin and Davaine. *Cryptococcus of Yellow Fever*. See *C. xanthogenicus*, Freire. *Cryptococcus xanthogenicus*, Domingo Freire. Considered by D. Freire to be the cause of yellow fever. Cf. *Bacillus of Yellow Fever*. Denecke's Cheese Bacillus. See *Spirillum tyrogenum*. Dengue Micrococci. See *Scheuteia laughlini*. *Detionella ochracea*, Saccardo. See *Leptothrix ochracea*, Kützing. *Diplococcus albicans amplius*, Bumm. Grayish white micrococcus. Found in mucus from the healthy vagina. aer., facanaer., liqf., nfg. *Diplococcus albicans tardus*, Unna and Tommasoli. Found in eczema. aer., nliqf., nmot. *Diplococcus albicans tardissimus*. Morphologically identical with the gonococcus; but more adherent, forming small masses. aer., nliqf., nfg. *Diplococcus citreus conglomeratus*, Bumm. Obtained from the air, in dust, and from gonorrheal pus. aer., facanaer., liqf., chg. (lemon-yellow), nfg. *Diplococcus citreus liquefaciens*, Unna and Tommasoli. Found on the skin in cases of eczema seborrhœicum. aer., liqf., nmot., chg. (lemon-yellow), nfg. *Diplococcus coryzæ*, Klebs and Hajek. Obtained from the nasal secretions in cases of acute nasal catarrh. aer., nliqf., nfg. *Diplococcus flavus liquefaciens tardus*, Unna and Tommasoli. Found on the skin in cases of eczema seborrhœicum. aer., facanaer., liqf., chg. (chrome-yellow), nfg. *Diplococcus fluorescens fœtidus*, Klamann. Obtained from the posterior nares. aer., facanaer., liqf., chg. (grass-green to violet), nfg. *Diplococcus intercellularis meningitidis*, Goldschmidt and Weichselbaum. Found within the cells of the exudate in cerebro-spinal meningitis. aer., nliqf., pg. *Diplococcus lacteus faviformis*, Bumm. Found frequently in the vaginal secretions. aer., nliqf., nfg. *Diplococcus luteus*, Ada-

metz. Found in water. aer., liqf., mot., chg. (yellow), nfg. *Diplococcus of Gonorrhea*. See *Micrococcus gonorrhœæ*, Neisser. *Diplococcus of Orchitis*, Hugouneng and Eraud. Syn., *Orchiococcus*. Frequently found in gonorrheal pus during the first few days. aer., nliqf., pg. *Diplococcus of Ozena*, Loewenberg. Found in the secretions in cases of ozena. *Diplococcus of Pemphigus*, *Diplococcus of Pemphigus Acutus*, Demme. Syn., *Micrococcus of Demme*. Found in the bullæ of pemphigus. aer., nliqf., pg. Resembles the gonococcus. *Diplococcus of Pneumonia in Horses*, Schütz. Obtained from the lungs of a horse affected with acute pneumonia. aer., nliqf., pg. *Diplococcus pneumoniae fibrinosæ*, Friedländer. See *Bacillus pneumoniae friedländeri*. *Diplococcus pneumoniae*, Weichselbaum. Syn., *Pneumococcus of Fränkel*. See *Bacillus septicus sputigenus*, Flügge. *Diplococcus pyogenes ureæ*, Rörising. Syn., *Diplococcus ureæ*, Rörising. *Coccobacillus pyogenes ureæ*, Rörising. *Coccobacterium ureæ*, Rörising. Found in purulent urine. Cf. *Diplococcus ureæ trifolius*, Rörising. *Micrococcus pyogenes ureæ flavus*, Rörising. *Diplococcus roseus*, Bumm. Found in the air. aer., facanaer., liqf., chg. (pink), nfg. *Diplococcus ureæ*, Rörising. See *Diplococcus pyogenes ureæ*, Rörising. *Diplococcus ureæ trifolius*, Rörising. Found in purulent urine. Cf. *Diplococcus pyogenes ureæ*, Rörising. *Micrococcus pyogenes ureæ flavus*, Rörising. *Dispora caucasica*, Kern. See *Bacillus caucasicus*, Kern. *Drumstick Bacillus*. See *Bacillus putreficiens coli*. *Epsilon Bacillus*. See *Spirillum finkleri*. *Erythroconus litoralis*, Christedt. See *Sarcina litoralis*. *Feet-sweat Bacillus*. See *Bacillus fœtidus*, *Bacillus saprogenes*, II, and *Bacillus graveolens*. *Finkler-Prior's Bacillus*. See *Spirillum Finkler-Prior*. *Flesh-colored Bacillus*, Tils. aer., liqf., mot., chg. (pink), nfg. *Flowers of Wine*. See *Saccharomyces mycoderma*, Reess. *Fränkel's Pneumobacillus*. See *Bacillus septicus sputigenus*, Flügge. *Frog-spawn Bacterium of Sugar Factories*, *Frog-spawn fungus*. See *Leuconostoc mesenteroides*. *Gaffky tetragenus*, Saccardo. See *Micrococcus tetragenus*, Gaffky. *Gas Forming Bacillus*. See *Bacillus gasiformans*, Eisenberg. *Ginger Beer Plant*, Marshall Ward. Forms jelly-like, semi-transparent, yellowish-white, brain-like masses at the bottom of the fermentation; resembles Caucasian kephir grains; it is a symbiotic association of *Saccharomyces pyeiformis* and of *Bacterium vermiforme*. *Glacier Bacillus*. See *Bacillus fluorescens nivialis*, Schmolck. *Gliococcus*, a micro-

coccus with a slimy capsule. *Golden Staphylococcus*. See *Staphylococcus pyogenes aureus*, Ogston, Becker. *Gray-white micrococcus*, Bumm. See *Diplococcus albicans amplius*, Bumm. *Hæmatococcus bovis*, Babes. Zimmermann. Obtained from the blood and viscera of cattle dead of an epidemic, infectious disease characterized by hemoglobinuria (in Roumania). *aer.*, *facanaer.*, *uliff.*, *pg.* *Hay bacillus*. See *Bacillus subtilis*, Ehrenberg. *Helicobacterium aerogenes*, Miller. See *Bacillus aerogenes*, III, Miller. *Helicobacterium klebsii*, Escherich. Found in the intestine of guinea pigs. *pleom.* *Helicomonas*, Klebs. Syphilis fungus. Cf. *Bacillus syphilidis*, Lustgarten. *Hormiscium cerevisiæ*, Bonard. *Hormiscium vini*. *Hormiscium vini et cerevisiæ*, Bonard. See *Saccharomyces mycoderma*. Hueppe's *Bacillus of Butyric Acid Fermentation*. See *Bacillus butyricus*, Hueppe. *Hygrocrocis vandelli*, Meneghi. See *Beggiatoa alba*, Trevisan. *Hypæothrix kühniana*. See *Crenothrix kühniana*. *Influenza-bacillus*. See *Bacillus of Influenza*, Pfeiffer. *Iodococcus vaginatus*, Miller. Found in unclean mouths; gives peculiar reaction with iodine, the sheath being stained yellow, the cocci dark blue. *Koch's Bacillus*. *Koch's Comma Bacillus*, *Commabacillus*. See *Spirillum cholera asiaticæ*, Koch. *Lanceolate Coccus*, *Lancet-shaped Micrococcus*, Talamon. See *Bacillus septicus eptigenus*, Flügge. *Lepra-bacillus*. See *Bacillus lepra*, Hansen. *Leptonema niveum*. See *Bacillus nivea*, Rabenhorst. *Leptothrix buccalis*, Robin. Syn., *Leptothrix gigantea*, Miller. *Leptothrix pulmonalis*, *Rasmussenia buccalis*, Saccardo. Found in the mouth of man and animals. Falsely considered the cause of dental caries. *Leptothrix buccalis of Vignal*. See *Bacillus buccalis (a)*, Vignal. *Leptothrix epidermidis*. Syn., *Bacillus epidermidis*, Bizzozero. *Microsporon minutissimum*, Burchart. Found on the epidermis between the toes, and held by Bizzozero to be nonpathogenic, but by Boeck as the cause of erythrasma. *Leptothrix gigantea*, Miller. Found on the teeth of dogs, sheep, cattle, and other animals affected with *Pyorrhæa alveolaris*. Some consider it identical with *Leptothrix buccalis*, Robin. *Leptothrix hyalina*, Reinsch. Found on marine algae. *Leptothrix innominata*, Miller. See *Leptothrix buccalis*, Robin. *Leptothrix kühniana*. See *Crenothrix kühniana*, Saccardo. *Leptothrix lacrimalis*. See *Cladotrix fursteri*, Cohn. *Leptothrix lanugo*, Kützing. Found on marine algae near Heligoland. *Leptothrix maxima buccalis*, Miller. See *Bacillus buccalis*

maximus, Miller. *Leptothrix muralis*, Kützing. See *Leptothrix parasitica*. *Leptothrix natans*. See *Sphaerotilus natans*, Saccardo. *Leptothrix ochracea*, Kützing. Syn., *Detionella ochracea*, Saccardo. *Cladotrix dichotoma*, Zopf. According to Zopf, this represents filaments of *Cladotrix* stained with oxid of iron. *Leptothrix of Puerperal Fever*, *Leptothrix puerperalis*, Fetz. See *Bacillus puerperalis*, Engel and Spillman. *Leptothrix parasitica*, Kützing. Parasitic on fresh-water algae. *Leptothrix pulmonalis*. See *Leptothrix buccalis*, Robin. *Leptothrix pusilla*, Rabenhorst. Found on fresh-water algae, also in the mouth, Klebs. *Leptothrix symplacoides*, Dickie. Found on marine algae. *Leptothrix vaginalis*. Found in the vagina of animals and women. *Leptothrix variabilis*, Saccardo. Found in saliva of healthy persons. *Leuconostoc mesenteroides*, Van Tieghem. Syn., *Asiococcus mesenteroides*, Cienkowski. Frog spawn bacterium of sugar factories, Frog spawn fungus. A source of great loss to the manufacturers of beet sugar, frequently and rapidly converting large quantities of the beet juice into a mucilagino-gelatinous mass. *Liborius's Bacillus of Butyric Acid Fermentation*, Liborius. See *Bacillus butyricus*, Prazmowski. *Lisophyton suspectum*, Hallier. See *Bacillus lysæ*, Pasteur. *Melunella flexuosa*, Borg. See *Spirillum rugula*, Müller. *Merismopedia chondroidea*, With, *Merismopedia glauca*. See *Sarcina renii*, Hepworth. *Merismopedia goodsirii*. See *Sarcina ventriculi*, Goodsir. *Merismopedia hyalina*, Kützing. *Sarcina hyalina*, Kützing. *Merismopedia litoralis*, Rabenhorst. See *Sarcina litoralis*, (Erstedt). *Merismopedia reitenbachii*, Caspary. See *Sarcina reitenbachii*, Caspary. *Merismopedia urinæ*, Rabenhorst. See *Sarcina urina*, Welcker. *Merismopedia ventriculi*, Husem. See *Sarcina ventriculi*, Goodsir. *Merismopedia violacea*, Kützing. See *Sarcina violacea*, Kützing. *Metallactar anthracis*, Trevisan. See *Bacillus anthracis*, Pollender and Davaine. *Metallactar tremulus*, Trevisan. See *Bacillus tremulus*, Koch. *Metallactar ulna*, Trevisan. See *Bacillus ulna*, Cohn. *Micoderma cerevisiæ*, Trecul. See *Saccharomyces mycoderma*, Reess. *Micrococcus aceti*. See *Bacillus aceti*, Kützing. *Micrococcus acidii lactici*, Marpmann. Found in milk; resembles in its action *Bacillus acidii lactici*, Hueppe. *aer.*, *uliff.*, *nmst.*, *npg.* Cf. *Sphaerococcus acidii lactici*, Marpmann. *Micrococcus acidii lactici liquefaciens*, Krüger. Found in butter and cheese. *aer.*, *facanaer.*, *liqf.*, *nmst.* *Micrococcus aerogenes*, Miller. Found in the intestinal

tract of man. *aer.*, *liqf.*, *nmot.*, *npg.*
Micrococcus agilis, Ali-Cohen. Found in drinking water. *aer.*, *liqf.*, *mot.*, *npg.*
 A motile coccus, provided with a flagellum; described by Menge. **Micrococcus agilis citreus**, Menge. **Micrococcus (Diplococcus) albicans amplius**, Bumm. Syn. *Neisseria albicans*. Frequently found in the vaginal secretions. *aer.*, *liqf.*, *npg.* **Micrococcus (Diplococcus) albicans tardus**, Unna and Tommasoli. Found in eczema. *aer.*, *nliqf.*, *nmot.* **Micrococcus albicans tardissimus**, Eisenberg. Bumm. Found in the vaginal secretions of puerperal women. *aer.*, *facanaer.*, *nliqf.*, *npg.* **Micrococcus liquefaciens**, Von Besser. Found in the nasal secretions of healthy persons. *aer.*, *facanaer.*, *liqf.*, *npg.* **Micrococcus albus urinæ**. See *Micrococcus urinæ*. **Micrococcus amylovorus**, Burrill and Arthur. Syn., *Micrococcus* of apple blight. *Bacterium* of pear blight. Produces fermentation in saccharine fluids. *aer.*, *nliqf.*, *mot.*, *fg.* (for apple and pear trees). Cf. *Bacillus butyricus*, Prazmowski. **Micrococcus aquatilis**, Bolton. Found frequently in water. *aer.*, *nliqf.*, *npg.* **Micrococcus arei celsii**, Buchner and Sehlen. Syn., *Bacterium decalvans*, Thin. Found on the diseased hairs in alopecia areata. Cf. *Bacillus of alopecia areata*, Kasauli, and *Micrococcus of alopecia areata*, Robinson. **Micrococcus ascoformans**, Johnne. See *Micrococcus botryogenus*, Rabe. **Micrococcus aurantiacus**, Cohn. Syn., *Pedococcus aurantius*. Found in water. *aer.*, *nliqf.*, *chg.* (orange-yellow). **Micrococcus biskræ**. See *Micrococcus of Heydenreich*. **Micrococcus blennorrhæus**. See *Micrococcus gonorrhææ*, Neisser. **Micrococcus bombycis**, Cohn, Bechamp. Syn., *Streptococcus bombycis*, Bechamp. Found in the intestinal tract of silkworms and other Lepidoptera dead of the epidemic disease known in France as *la flacherie*; etiologic relation demonstrated by Pasteur. **Micrococcus botryogenus**, Bollinger, Rabe. Syn., *Bollingeria equi*, Saccardo; *Ascooccus johnei*, Cohn; *Micrococcus ascoformans*, Johnne; *Micrococcus* of myko-desmoids of the horse. Found in colonies in the connective tissue of horses affected with "mykodesmoids." *aer.*, *liqf.*, *fg.* **Micrococcus candicans**, Flügge. Found abundantly in air and water. *aer.*, *nliqf.*, *npg.* **Micrococcus candidus**, Cohn. Found in water. *aer.*, *nliqf.*, *npg.* **Micrococcus capillorum**. Found on the scalp and said to alter the color of the hair. *aer.*, *chg.* (reddish-yellow). **Micrococcus carneus**, Zimmermann. Found in water. *aer.*, *nliqf.*, *chg.* (red to violet), *npg.* **Micrococcus cerasinus siccus**, List. *aer.*, *nliqf.*, *chg.*

(cherry-red), *npg.* **Micrococcus cereus albus**, Passet. Syn., *Staphylococcus cereus albus*, Passet. Found in the pus of acute abscesses, and by Tils in hydrant water (Freiburg). *aer.*, *nliqf.*, *npg.* **Micrococcus cereus aureus**, Schröter and Winkler. Syn., *Staphylococcus cereus aureus*, S. and W. Found in the thin secretions of coryza. *aer.*, *nliqf.*, *chg.* (orange-red), *npg.* **Micrococcus cereus flavus**, Passet. Syn., *Staphylococcus cereus flavus*, Passet. *Staphylococcus passetii*. Found in an acute abscess. *aer.*, *nliqf.*, *chg.* (lemon-yellow), *npg.* **Micrococcus chlorinus**. Occurs in zoogloea masses. *aer.*, *chg.* (greenish-yellow), *npg.* **Micrococcus cholerae gallinarum**, Pasteur. See *Bacillus septicaemia hemorrhagica*, Sternberg. **Micrococcus cinnabareus**, Flügge. Syn., *Streptococcus cinnabareus*, Flügge. Found in air and water. *aer.*, *nliqf.*, *chg.* (reddish-brown), *npg.* **Micrococcus citreus**, List. Syn., *Staphylococcus citreus*, List. Found in water, also by Lannelongue and Achard in the seat of an old osteomyelitis. *aer.*, *nliqf.*, *chg.* (pale yellow), *npg.* **Micrococcus citreus conglomeratus**, Bumm. See *Diplococcus citreus conglomeratus*, Bumm; *Micrococcus concentricus*, Zimmermann. Found in water. *aer.*, *nliqf.*, *sap.*, *npg.* **Micrococcus conglomeratus**, Weichselbaum. Syn., *Neisseria conglomerata*, W. **Micrococcus coronatus**, Flügge. Syn., *Streptococcus coronatus*, Flügge. Found in the air. *aer.*, *liqf.*, *npg.* **Micrococcus cremoides**, Zimmermann. Found in water. *aer.*, *liqf.*, *chg.* (yellowish-white), *npg.* **Micrococcus crepusculum**, Cohn. See *Bacillus cremeus*, Ehrenberg and Dujardin. **Micrococcus cumulus tenuis**, Von Besser. Found in abundance in the nasal mucus of man. *aer.*, *facanaer.*, *nliqf.*, *npg.* **Micrococcus cyaneus**, Schröter. Obtained from the air, on boiled potatoes. *aer.*, *chg.* (cobalt blue, altered to carmine by acids, restored by alkalis). **Micrococcus decalvans**, Schröter. Syn., *Bacterium decalvans*, Saccardo. Cf. *Bacillus of alopecia areata*, Kasauli, *Micrococcus arei celsii*, Buchner and Sehlen; *Micrococcus of alopecia areata*, Robinson. **Micrococcus diffuens**, Schröter. Found in the air, dust, and in excrement; common. *aer.*, *nliqf.*, *chg.* (yellow, with a greenish fluorescence; not modified by acids). Resembles *Bacillus fluorescens putidus*, Flügge. **Micrococcus diphtheriticus**, Oertel, Cohn, Löffler. Syn., *Streptococcus articularum*, Löffler. Found in the false membranes, and in the tissues of diphtheria patients; Cornil identifies this with *Micrococcus pyogenes*, Rosenbach. *aer.*, *nliqf.*, *fg.* **Micrococcus doyenii**. See *Micrococcus urinæ albus clarus*, Doyen. **Micrococcus endocarditicus**, Syn.,

Streptococcus endocarditidis. **Micrococcus** endocarditidis rugatus, Weichselbaum. Found in a case of ulcerative endocarditis. *aer., nliqf., pg.* **Micrococcus erysipelatis**, **Micrococcus erysipelatos**, **Micrococcus erysipelatosus**, Fehleisen. See *Micrococcus pyogenes*, Rosenbach. **Micrococcus fermenti**. See *Micrococcus viniperda*. **Micrococcus fervidus**, Adametz. Found in water. *aer., liqf., npg.* **Micrococcus fickii**. Found in healthy and diseased eyes. **Micrococcus finlayensis**, Sternberg. Obtained from the viscera of a yellow-fever cadaver. *aer., liqf., chg.* (pale-yellow), *npg.* **Micrococcus flavus**. See *Micrococcus flavus liquefaciens*, Flügge. **Micrococcus flavus desidens**, Flügge. Found in air and water. *aer., liqf., chg.* (yellowish-brown), *npg.* **Micrococcus flavus liquefaciens**, Flügge. Syn., *Micrococcus flavus, Micrococcus liquefaciens*. **Micrococcus flavus tardigradus**, Flügge. Syn., *Micrococcus tardigradus*. Found in the air and water. *aer., nliqf., chg.* (dark olive-green), *npg.* **Micrococcus foetidus**, Klamann, Rosenbach. Found in the posterior nares of man. *aer., liqf., npg.* **Micrococcus fervidus**, Adametz. Found in water. *aer., nliqf., npg.* **Micrococcus freudenreichii**, Guillebeau. One of the two forms found by Guillebeau to produce ropiness in milk. *aer., liqf., npg.* Cf. *Bacterium hessii*, Guillebeau. **Micrococcus friedländeri**. See *Bacillus pneumoniae friedländeri*. **Micrococcus fulvus**, Cohn. Syn., *Staphylococcus fulvus*. Found in horse and rabbit dung. *aer. nmot., nliqf., chg.* (rose), *npg.* **Micrococcus fuscus**, Maschek. Found in water. *aer., nmot., liqf., chg.* (sepia-brown), *npg.* **Micrococcus gallicidus**. See *Bacillus septicaemia hamorrhagica*, Sternberg. **Micrococcus gelatinosus**. Found in milk, which it causes to coagulate at about 22° C. **Micrococcus gingivæ pyogenes**, Miller. Found in the mouth of a man affected with alveolar abscess. *aer., facanaer., nliqf., pg.* **Micrococcus gonorrhææ**, Neisser. *Micrococcus blennorrhæus*, Neissera gonorrhææ, *Micrococcus gonococcus*. Found in gonorrhæal pus. *aer.* No growth on ordinary gelatin; has been grown on acid gelatin and human blood-serum. *pg.* **Micrococcus griseus**, Winter. See *Bacterium griseum*, Warming. **Micrococcus hæmatodes**, Babes. Syn., *Micrococcus of red sweat*. Found in foul sweat from the axilla. *aer., chg.* (red), *npg.* **Micrococcus havaniensis**, Sternberg. Syn., *Bacillus havaniensis*, Sternberg. Obtained by Sternberg in his investigation of yellow fever. *aer., nliqf., chg.* (blood-red). Only found in the presence of free oxygen. **Micrococcus of**

hepatitis enzootica of hogs. See *Micrococcus porcellorum*. **Micrococcus imetrophus**, Trevisan. See *Bacillus prodigiosus*, Ehrenberg. **Micrococcus indicus**, Koch. See *Bacillus indicus*, Kozin. **Micrococcus influenzae**, Letzenich. Syn. *Streptococcus seiferti*, *Micrococcus influenzae*, Seifert. See *Bacillus influenzae*, Pfeiffer. **Micrococcus insectorum**, Burri. *Streptococcus insectum*. Found in the oral appendages or pancreatic organs of the Chinese bug (*Blissus leucostictus* Say), and the higher Hemiptera. Especially recommended as a remedy for the Chinese bug and for *Haltina minutus*, so damaging to the peanut. **Micrococcus intercellularis meningitidis**, Neumann and Schäffer. Syn. *Micrococcus meningitidis*, Neumann. *Neisseria Weichselbaumii*. Found in the tissues of the brain and spinal cord in cases of cerebrospinal meningitis. **Micrococcus lacteus flaviformis**, Burri. Syn. *Neisseria lactea*. Frequently found in normal vaginal mucus. **Micrococcus lactis viscosus**, Conn. Syn. *Micrococcus* of bitter milk, Conn. The cause of the bitter taste of milk and cream. **Micrococcus liquefaciens**. See *Micrococcus fluxus liquefaciens*, Löffler. **Micrococcus lewenbergii**. Syn. *Micrococcus* of ozena. Found in the secretions in cases of ozena. **Micrococcus luteus**, Conn. Found in water, the superficial soil, dust of the rooms, food, etc. **Micrococcus lysæ**. *Neisseria lysæ*, Neumann. *Neisseria lysæ*, Neumann. *Neisseria* (Syn. *Micrococcus*) of lymphoma. **Micrococcus of rabbit**, Fol. Found in the spinal cord of rabbit, atanas. Conn. and Babes throw doubt on the identity of specimens of Fol. **Micrococcus major**, Dreyer. See *Micrococcus ureæ major* Dreyer. **Micrococcus manifestus**. Syn. *Streptococcus manifestus*. *Micrococcus* of granular granuloma formation. Found in the pattern of two cases of pyoderma in the skin. **Micrococcus mastitidis**, Nozard. *Micrococcus* of mastitis in cows. *Streptococcus* of mastitis in cows. *Micrococcus* of contagious mastitis. *Micrococcus* of gangrenous mastitis in sheep. Nozard. Obtained from the milk of cows and sheep affected with mastitis. **Micrococcus meningitidis**, Neumann. See *Micrococcus intercellularis meningitidis*, Neumann and Schäffer. **Micro-**

coccus meningitidis purulentæ, Heydenreich. *Micrococcus morbillosus*. Syn., *Streptococcus morbillosus*. *Micrococcus nasalis*, Hack. Found in the nasopharyngeal space. *aer.*, *mot.*, *nlif.*, *npg.* *Micrococcus nitrificans*, Van Tieghem. Found in water. *aer.*, *npg.* *Micrococcus*, II, of Fischel. Found in the blood of two cases of influenza. *aer.*, *facanaer.*, *liqf.*, *pg.* *Micrococcus oblongus*. See *Bacterium oblongum*, Boutroux. *Micrococcus ochroleucus*, Prove and Legrain. Syn., *Streptococcus ochroleucus*. Found in urine. *aer.*, *nlif.*, *chg.* (sulphur yellow), *npg.* *Micrococcus of Acute Infectious Osteomyelitis*. See *Micrococcus pyogenes aureus*, Rosenbach. *Micrococcus of Alopecia areata*, Von Sehlen, Robinson, Vaillard, and Vincent. Found on the hairs from the diseased patches. *aer.*, *liqf.*, *pg.* Cf. *Bacillus of Alopecia areata*, and *Micrococcus decalvans*. Schröter. *Micrococcus of Ammoniacal Urine*. See *Micrococcus ureæ*, Cohn, and *Micrococcus ureæ liquefaciens*, Flügge. *Micrococcus of Apple Blight*. See *Bacillus butyriscus*, Botkin, Prazmowski, etc. *Micrococcus of Beriberi*, Lacerda. See *Bacillus beribericus*, Lacerda. *Micrococcus of Biskra Button*. See *Micrococcus of Heydenreich*. *Micrococcus of Bitter Milk*. See *Micrococcus lactis viscosus*, Conn. *Micrococcus of Bronchitis*, Pichini. Three distinct forms. *aer.*, *liqf.* Cf. *Bacillus of Luminizer*. *Micrococcus of Bovine Mastitis*, Kitt. See *Micrococcus mastobius*, Nocard, Mollereau, and Kitt. *Micrococcus of Bovine Pneumonia*, Poels and Nolen. *Micrococcus of cattle pneumonia*. Found in the lungs of cattle suffering from infectious pleuropneumonia. *aer.*, *nlif.*, *pg.* *Micrococcus of Cerebrospinal Meningitis*. See *Micrococcus intercellularis meningitidis*. Neumann and Schäffer. *Micrococcus of Chicken Pox*. See *Staphylococcus viridis flavescens*, Guttman. *Micrococcus of Contagious Mammitis*. See *Micrococcus mastobius*, Nocard, Mollereau, and Kitt. *Micrococcus of Cow Pox*. See *Micrococcus variola et vaccinae*, Cohn. *Micrococcus of Dantec*. Obtained from salted codfish which had turned red. *aer.*, *nlif.*, *chg.* (yellow, changing to deep red), *npg.* *Micrococcus of Demme*. See *Diplococcus of pemphigus acutus*, Demme. *Micrococcus of Dental Caries*. See *Leptothrix vulgaris*, Robin; *Leptothrix gigantea*, Milles; *Miller's epsilon bacillus*, and *Spirillum dentium*, Arnt. *Micrococcus of Dengue*, McLaughlin. See *Schantzia laughlini*. *Micrococcus of Diphtheria*. *Streptococcus articolorum*, Löffler. *Micrococcus of Disse* and Taguchi. Obtained from the blood,

the secretions of broad condyloma, as well as the primary indurations of syphilis. Etiologic relations not confirmed. *Micrococcus of Drinking Water*. See *Micrococcus aquatilis*, Bolton. *Micrococcus of Egyptian Ophthalmia*. See *Micrococcus of trachoma*, Sattler and Michel. *Micrococcus of Endocarditis*. See *Streptococcus endocarditicus* and *Micrococcus endocarditicus rugatus*, Weichselbaum. *Micrococcus of Erysipelas*, Fehleisen. See *Micrococcus pyogenes*, Rosenbach. *Micrococcus of Foot and Mouth Disease*. Syn., *Streptococcus of erizema epizootica*, Schottelius, *Streptococcus aphthicola*, Hallier. Found in the vesicular eruptions about the mouth and feet of cattle, pigs, and sheep. Communicable to man. *aer.*, *facanaer.* Pathogenesis not fully settled. Cf. *Bacillus of ulcerative stomatitis of cattle*. *Micrococcus of Forbes*. Found in the bodies of larvae of the cabbage butterfly, *Pieris rapæ*, affected with an infectious disease. *Micrococcus of Fowl Cholera*. See *Bacillus septicemie hæmorrhagica*, Sternberg. *Micrococcus of Freire*. See *Cryptococcus xanthogenicus*. *Micrococcus of Furuncle*. See *Bacillus osteomyelitis*. *Micrococcus of Gangrenous Mastitis in Sheep*, *Micrococcus of Gangrenous Mammitis*. See *Micrococcus mastobius*, Nocard, Mollereau, and Kitt. *Micrococcus of Gonorrhæa*. See *Micrococcus gonorrhæa*, Neisser. *Micrococcus of Gray Parrot Disease*. See *Micrococcus psittaci*, Eberth and Wolff. *Micrococcus of Heydenreich*. Syn., *Streptococcus biskrae*. *Micrococcus of Biskra Button*. Found in the tumors and ulcers characteristic of Aleppo boil or Biskra button. *aer.*, *liqf.*, *pg.* *Micrococcus of Hydrophobia*. See *Micrococcus hyssæ*, Pasteur. *Micrococcus of Infectious Osteomyelitis*. See *Micrococcus pyogenes aureus*, Rosenbach, and *Micrococcus osteomyelitis*, Becker and Kraske. *Micrococcus of Influenza*, Seifert, *Streptococcus seiferti*. See *Micrococcus influenzae*, Letzerich. *Micrococcus of Kirchner*. Obtained from the sputum of patients suffering with influenza. *aer.*, *pg.* *Micrococcus of Manfredi*. See *Micrococcus manfredii*. *Micrococcus of Mastitis*, Kitt. See *Micrococcus mastobius*, Nocard and Mollereau. *Micrococcus of Measles*, Klebs and Keating. Syn., *Streptococcus morbillosus*, *Micrococcus morbillosus*. Found in the blood and exudates of persons afflicted with measles. Flügge considers it unimportant. *Micrococcus of Mykodesmoids of the Horse*. See *Micrococcus butyrigenes*, Rabe. *Micrococcus of Osteomyelitis*. See *Micrococcus pyogenes aureus*, Ogston. *Micrococcus of Ozena*. See *Micrococcus lawenberghii*. *Micrococcus of Pear Blight*,

Burrill. See *Bacillus butyricus*, Prazmowski. **Micrococcus of Pébrine.** See *Nosema bombycis*, Nägeli. **Micrococcus of Pneumonia.** See *Bacillus pneumoniae friedländerii* and *Micrococcus septicus*, Flügge. **Micrococcus of Progressive Abscess Formation in Rabbits,** Koch. Found in the exudates of rabbits inoculated with putrefying blood. *pg.* **Micrococcus of Progressive Granuloma Formation.** See *Micrococcus manfredii*. **Micrococcus of Progressive Tissue Necrosis in Mice,** Koch. Found in the exudates of mice inoculated with putrid blood. *pg.* **Micrococcus of Pyemia in Rabbits,** Koch. Found in the exudates of rabbits inoculated with putrefying flesh infusion. *pg.* **Micrococcus of Rabies,** Fol. See *Micrococcus lyssa*, Neisser. **Micrococcus of Red Sweat.** See *Micrococcus hematodes*, Babes. **Micrococcus of S. Le M. Moore.** See *Bacillus fatidus*, Liborius. **Micrococcus of Salvioli.** See *Bacillus septicus sputigenus*, Flügge. **Micrococcus of Scarlet Fever,** Pohl and Pincus. See *Micrococcus scarlatinae*. **Micrococcus of Septicemia in Rabbits,** Koch. Syn., *Streptococcus Kochii*. Found in the exudates of rabbits inoculated with putrefying flesh-infusion. *pg.* **Micrococcus of Smallpox.** See *Micrococcus variolæ et vaccinae*, Cohn. **Micrococcus of Sputum Septicemia.** See *Bacillus septicus sputigenus*, Flügge. **Micrococcus of Swine Plague or Hog Cholera.** See *Bacillus of swine plague*, Marseilles, Rietsch, and Jobert, and *Bacillus ex pneumo-enteritide suis*, Klein. **Micrococcus of Syphilis,** Haberkorn and Marcus. Found in the white blood corpuscles in cases of syphilis. *aer., osc., chg.* (red). **Micrococcus of Trachoma,** Sattler and Michel. Syn., *Neisseria rebellis*. **Micrococcus of Egyptian ophthalmia.** Found in the secretions and nodules of the conjunctiva in cases of trachoma. *aer., nliqf., pg.* Researches of Baumgarten and Kartulis go to show that this form has no etiologic relation to the diseases mentioned. **Micrococcus of Vaccinia.** See *Micrococcus variolæ et vaccinae*, Cohn. **Micrococcus of Whooping Cough,** Letzerich. Found in the sputum in cases of whooping cough. See *Bacillus tussis convulsivæ*, Afanassiew. **Micrococcus olearius.** See *Micrococcus urinae flavius olivarius*. **Micrococcus osteomyelitis,** Becker. Syn., *Micrococcus* of acute infectious osteomyelitis. See *Micrococcus pyogenes aureus*, Rosenbach. **Micrococcus ovalis,** Escherich. Found in the meconium and feces of infants. *aer., facanaer., nliqf., npg.* **Micrococcus ovatus.** See *Micrococcus of pébrine*. **Micrococcus pasteurii,** Sternberg. See *Bacillus*

septicus sputigenus, Flügge. **Micrococcus petechialis.** Syn., *Neisseria petechialis*. **Micrococcus pflügeri,** Ludwig. See *Photobacterium pflügeri*, Ludwig. **Micrococcus phosphoreus,** Cohn. See *Bacillus phosphoreus*, Cohn. **Micrococcus plumosus,** Bräutigam. Found in water. *aer., nliqf., npg.* **Micrococcus pneumoniae crouposæ,** Sternberg. See *Bacillus septicus sputigenus*, Flügge. **Micrococcus pneumoniae infectiosæ,** Friedländer. See *Bacillus pneumoniae friedländerii*. **Micrococcus porcellorum.** Syn., *Micrococcus* of Hepatitis enzootica of hogs. Found in hogs affected with hepatitis (Russia). *aer., liqf., pg.* **Micrococcus prodigiosus,** Cohn. See *Bacillus prodigiosus*, Ehrenberg. **Micrococcus pseudocyanus,** Cohn. Found in boiled potatoes exposed to air. *aer., chg.* (verdigris green), *npg.* **Micrococcus psittaci,** Wolf. Syn., *Streptococcus pernicius psittacorum*, Eberth and Wolf. *Micrococcus* of gray parrot disease. The cause of a disease of parrots resembling fowl cholera; fatal to many of the parrots imported from Guinea into Europe. **Micrococcus putridus.** A form emitting a fetid gas from gelatin when mixed with iodiform. **Micrococcus pyocyaneus,** Gessard. See *Bacillus pyocyaneus a*, Gessard. **Micrococcus pyogenes,** Rosenbach. Syn., *Streptococcus pyogenes*, *Streptococcus erysipelatos*, *Streptococcus longus*, Von Lingelsheim. *Micrococcus erysipelatis (tos) (tosus)*, *Streptococcus* of pus. *Micrococcus* of erysipelas, Fehleisen. Found in the pus of acute abscesses, and in the skin in cases of erysipelas. *aer., facanaer., nliqf., pg.* **Micrococcus pyogenes albus,** Rosenbach. Syn., *Staphylococcus pyogenes albus*, Passet. *Staphylococcus epidermidis albus*, Welch. Frequently found in company with *Micrococcus pyogenes aureus*. The cause of "stitch abscesses." *aer., liqf., pg.* **Micrococcus pyogenes aureus,** Rosenbach. Syn., *Staphylococcus pyogenes aureus*. *Micrococcus* of infectious osteomyelitis. The species most frequently found in pus. Obtained from various suppurations, boils, anthrax, osteomyelitis, ulcerative endocarditis, phlegmon, etc. *aer., facanaer., liqf., chg.* (brilliant golden yellow), *pg.* **Micrococcus pyogenes citreus,** Passet. Syn., *Staphylococcus pyogenes citreus*, Passet. *Micrococcus pyogenes flavus*. Found associated with the two preceding forms first described in acute abscesses. *aer., facanaer., liqf., chg.* (lemon yellow), *pg.* Held by some to be but a variety of *Micrococcus pyogenes aureus*. **Micrococcus pyogenes tenuis,** Rosenbach. Syn., *Micrococcus rosenbachii*. Found in pus. Culture thin, varnish-like; pathogenesis not settled. Cf. *Micrococcus septicus*,

Flügge. *Micrococcus pyogenes ureæ* flavus, Rörising. Found in purulent urine. Cf. *Diplococcus pyogenes ureæ*, Rörising; *Diplococcus ureæ trifolius*, Rörising. *Micrococcus pyosepticus*, Richet. Syn., *Staphylococcus pyosepticus*, Hericourt and Richet. Obtained from a carcinomatous tumor in a dog. aer., liqf., pg.; resembles *Micrococcus pyogenes albus*, Rosenbach. *Micrococcus radiatus*, Flügge. Syn., *Streptococcus radiatus*, Flügge. Found in air and water. aer., liqf., nfg. *Micrococcus rheumarthritidis*, Küssmaul. See *Bacillus rheumarthritidis*, Küssmaul. *Micrococcus rosenbachii*. See *Micrococcus pyogenes tenuis*, Rosenbach. *Micrococcus rosettaceus*, Zimmermann. Found in water. aer., nliqf., nfg. *Micrococcus roseus*, Eisenberg. Found in the sputum of a person suffering from influenza. aer., facanaer., liqf., chg. (pink), nfg. *Micrococcus roseus*, Bumm. See *Diplococcus roseus*, Bumm. *Micrococcus roseus*, Maggiora, Flügge. Syn., *Neisseria rosea*. Found on the skin. aer., nfg. *Micrococcus salivarius pyogenes*, Biondi. Syn., *Staphylococcus salivarius pyogenes*, Biondi. Found in an abscess produced in a guinea-pig by inoculation with saliva from a child affected with anginose scarlatina. aer., liqf., pg. *Micrococcus salivarius septicus*, Biondi. Obtained from animals after inoculation with saliva from a person with puerperal septicemia. aer., nliqf., pg. Cf. *Micrococcus sialosepticus*. *Micrococcus scarlatinæ*, *Micrococcus scarlatinus*, Coze and Feltz. Syn., *Perroncitoa scarlatina*. *Micrococcus* of scarlet fever, Pohl and Pincus. Found in the blood, on the desquamating skin, and on the palate of scarlet fever patients; resembles *Micrococcus septicus*, Flügge. Cf. *Streptococcus rubiginosus*, Edington, and *Bacillus* of scarlet fever, Crooke. *Micrococcus septicus*, Cohn. See *Bacillus septicæmia hæmorrhagica*, Sternberg. *Micrococcus septicus*, Flügge. Syn., *Streptococcus septicus*, Flügge, and *Streptococcus pyogenes*, Saccardo, *Streptococcus pneumoniae*, Weichselbaum. Found in soil. aer., nliqf., pg. *Micrococcus septopyæmicus*. Syn., *Streptococcus septopyæmicus*. Obtained from cases of phlegmonous angina and erysipelas of the larynx. aer., nliqf., pg. Resembles *Micrococcus septicus*, Flügge. *Micrococcus sialosepticus*. Found in the saliva in a case of puerperal septicemia. aer., nliqf., pg. Cf. *Micrococcus salivarius septicus*, Biondi. *Micrococcus subflavus*, Flügge, Neisser. Syn., *Neisseria subflava*. Yellowish-white micrococcus or diplococcus, Bumm. Obtained from the lochia of puerperal women, and from vaginal mucus. aer.,

liqf., pg. *Micrococcus suis*, Burrill. See *Bacillus tardigradus*. *Micrococcus tardigradus*. See *Micrococcus flavus tardigradus*, Flügge. *Micrococcus tetragenus*, *Micrococcus tetragenus*, Gaffky. Syn., *Gaffkyia tetragenus*, Saccardo. Obtained from a cavity in the lung of a person with pulmonary tuberculosis. aer., facanaer., nliqf., pg. *Micrococcus tetragenus*, Marotta. Constantly found in small-pox vesicles and pustules. aer., liqf., pg. Produces vaccinia in calves. *Micrococcus tetragenus febris flavæ*, Finlay. Syn., *Micrococcus tetragenus versatilis*, Sternberg. Obtained from the excrements of mosquitoes which had drawn blood from yellow fever patients; also from the surface of the body in the same disease (Sternberg). aer., liqf., chg. (lemon-yellow), nfg. *Micrococcus tetragenus mobilis ventriculi*, Mendosa. Found in the contents of the stomach of man. aer., nliqf., mot., nfg. *Micrococcus tetragenus subflavus*, Von Besser. Found in nasal mucus. aer., facanaer. (no growth in gelatin), nfg. *Micrococcus tetragenus versatilis*, Sternberg. See *Micrococcus tetragenus febris flavæ*, Finlay. *Micrococcus toxicatus*, Burrill. Syn., *Micrococcus variole et vaccinae*, Cohn. *Streptococcus toxicatus*, Burrill. Found in the poison of the poison ivy, *Rhus toxicodendron*. *Micrococcus ulceris mollis*. See *Bacillus* of chancreoid, Ducrey. *Micrococcus ureæ*, Cohn. Syn., *Bacterium ureæ*, Cohn. *Streptococcus ureæ*. *Torula ureæ*, Pasteur. Found in urine, in which it produces ammoniacal fermentation. *Micrococcus ureæ*, Pasteur and Van Tieghem. Found in the air and in ammoniacal urine. aer., facanaer., nliqf., nfg. Possibly the last two forms and the *Bacterium ureæ*, Jaksch, are identical. *Micrococcus ureæ liquefaciens*, Flügge. Syn., *Streptococcus athebius*. Found in ammoniacal urine. aer., facanaer., liqf., nfg. *Micrococcus urinæ albus*. Syn., *Micrococcus albus urinæ*. Found in the urine in cases of pyelonephritis and cystitis. *Micrococcus urinæ albus olearius*, Doyen. Syn., *Micrococcus doyenii*. Found under the same circumstances as the last form. aer., liqf. *Micrococcus urinæ flavus olearius*, Doyen. Syn., *Micrococcus olearius*. Found in the urine of cystitis, with the two preceding forms. aer., liqf., chg. (golden-yellow). *Micrococcus urinæ major*, Doyen. Syn., *Micrococcus major*. Found in same circumstances as the three preceding forms. *Micrococcus versatilis*. Found on the skin of healthy persons, also in the liver, spleen and kidneys after death from yellow fever. Not sufficiently studied. *Micrococcus versicolor*, Flügge. Found in water. aer., nliqf., chg. (yellow), nfg.

Micrococcus viniperda. Syn., *Micrococcus fermenti*. Found in spoiled wine. **Micrococcus violaceus**, Cohn. Found in water. *aer.*, *nlif.*, *chg.* (violet), *npg.* **Micrococcus viridis flavescens**, Guttman. Found in the lymph of a varicella pustule. *aer.*, *nlif.*, *npg.* **Micrococcus viscosus**, Pasteur. Found in the disease of wine known as *la graisse*. **Micrococcus viticulosus**, Flügge, Katz. Found in air and water. *aer.*, *facanaer.*, *nlif.*, *npg.* **Micrococcus xanthogenicus**, *Cryptococcus xanthogenicus*, Freire. Found in cases of yellow fever. **Microhaloa rosea**, Kützing. See *Beggiatoa roseopersicina*, Zopf. **Microsporon minutissimum**, Burchardt. See *Bacillus epidermidis*, Bizzozero. **Microsporon septicum**, Klebs. See *Bacillus septicæmia hæmorrhagica*, Sternberg. **Microzyma bombycis**, Béchamp. See *Micrococcus bombycis*, Cohn, Beck. **Milkwhite Micrococcus**. Bumm. See *Micrococcus albicans tardissimus*, Eisenberg. **Milk Yeast**. See *Saccharomyces lactis*, Adametz. **Miller's Bacillus**. See *Bacillus dentatis viridans*, Miller, and *Spirillum* of Miller. **Miller's Bacillus of the Intestine**. See *Bacillus aerogenes*, I, II, III, Miller. **Miller's Epsilon Bacillus**. See *Epsilon Bacillus*, Miller. **Milzbrandbacillus**. See *Bacillus anthracis*, Pollender and Davaine. **Monas prodigiosa**, Ehrenberg. See *Bacillus prodigiosus*, Ehrenberg. **Monas termo**, Müller. See *Bacillus termo*, Dujardin. **Mycoderma aceti**. See *Bacillus aceti*, Kützing. **Mycoderma vini**, Pasteur. See *Saccharomyces mycoderma*, Reess. **Myconostoc gregarium**, Cohn. A spiral or crumpled condition of a filamentous schizomycete, perhaps *Cladotrix dichotoma*, Cohn. **Neisseria albicans**. See *Micrococcus albicans amplius*, Neisser. **Neisseria citrea**. See *Micrococcus citreus conglomeratus*, Bumm. **Neisseria conglomerata**. See *Micrococcus conglomeratus*, Weichselbaum. **Neisseria gonorrhœa**. See *Micrococcus gonorrhœa*, Neisser. **Neisseria lactea**. See *Micrococcus lacteus farviformis*, Neisser. **Neisseria lyssæ**. See *Micrococcus lyssæ*, Neisser. **Neisseria petechialis**. See *Micrococcus petechialis*, Neisser. **Neisseria rebellis**. See *Micrococcus of trachoma*, Sattler. **Neisseria rosea**. See *Micrococcus roseus*, Maggiora. **Neisseria subflava**. See *Micrococcus subflavus*, Flügge. **Neisseria tardissima**. See *Micrococcus albicans tardissimus*, Bumm. **Neisseria weichselbaumii**. See *Micrococcus intercellularis meningitidis*, Neumann and Schäffer. **Neisseria winckleri**. See *Bacillus heribericus*, Lacerda. **Nitrifying Bacillus of Winogradsky**, Nitromonas of Winogradsky. It is found in the soil; no growth on ordinary culture media; one

of the very few nitrifying ferments. **Nosema bombycis**, Nägeli. Syn. *Micrococcus ovatus*. It is found in the organs and the blood of silk-worms suffering from *pébrine*. **Ophidomonas jenensis**, *Ophidomonas sanguinea*, Ehrenberg. See *Beggiatoa roseopersicina*, Zopf. **Orchiococcus**. See *Diplococcus of Orchitis*, Hugouneng and Eraud. **Oscillaria alba**, Vaucher. See *Beggiatoa alba*, Trevisan. **Oscillaria arachnoidea**, Agardh. See *Beggiatoa arachnoidea*, Trevisan. **Oscillaria leptomitiformis**, Meneghi. See *Beggiatoa leptomitiformis*, Trevisan. **Oscillaria tigrina**. See *Beggiatoa tigrina*, Rabenhorst. **Oscillaria versatilis**, Kützing. See *Beggiatoa arachnoidea*, Trevisan. **Palmella infusionum**, Ehrenberg. See *Bacillus termo*, Dujardin. **Palmella prodigiosa**, Mont. See *Bacillus prodigiosus*, Ehrenberg. **Palmellina capillorum** (Radlkoffer), Bulh. Syn., *Zooglaea capillorum*, Bulh. Found in the epidermis of the head; supposed to affect the color of the hair. **Panhystophyton ovatum**. Syn. of *Nosema bombycis*. **Pas-set's Putrefying Bacillus of Pus**. See *Bacillus pyogenes fœtidus*, Passet. **Pea-colored Bacterium**, Lankester. See *Beggiatoa roseopersicina*, Zopf. **Pediococcus acidi lactici**, Lindner. Found in hay infusion and in mash from malt. *aer.*, *facanaer.*, *nlif.*, *npg.* **Pediococcus albus**, Lindner. Found in well water. *aer.*, *liqf.*, *npg.* **Pediococcus aurantiacus**, Cohn. See *Micrococcus aurantiacus*, Cohn. **Pediococcus cerevisiæ**, Balcke. Syn., *Sarcina cerevisiæ*, Lindner. Found in beer and in the air of breweries. *aer.*, *facanaer.*, *nlif.*, *npg.* **Peroncitoa scarlatinosæ**. See *Micrococcus scarlatina*, Pohl-Pincus. **Pfeiffer's Bacillus**. See *Bacillus capsulatus*, Pfeiffer. **Photobacterium balticum**, Fischer. Found in the waters of the Baltic. *aer.*, *liqf.*, *phos.*, *npg.* Cf. *Bacillus phosphorescens indigenus*, Fischer. **Photobacterium fischeri**, Katz. Found in the waters of the Baltic. *aer.*, *liqf.*, *phos.*, *npg.* **Photobacterium indicum**, Fischer. See *Bacillus phosphorescens*, Fischer. **Photobacterium luminosum**, Beyerinck. Found on many of the phosphorescent animals, crustaceous polyps, infusoria, of the North Sea. *aer.*, *liqf.*, *phos.*, *npg.* **Photobacterium pflügeri**, Ludwig. Syn., *Bacterium pflügeri*, *Micrococcus pflügeri*. Found upon butcher's meat and upon market fish. The most phosphorescent of all light-producing bacteria. *aer.*, *liqf.*, *phos.*, *npg.* **Photobacterium phosphorescens**, Fischer. See *Bacillus phosphoreus*, Cohn. **Photobacterium phosphorescens**, Cohn. Syn., *Bacterium phosphorescens*. **Pink Bacillus of Spreading Edema**, A. B. Harris. *aer.*, *nlif.*, *pleom.*, *chg.* (pink),

nfg. Cf. *Bacillus oedematis maligni*, Koch. *Pleurococcus roseopersicina*, Rabenhorst. See *Beggiatoa roseopersicina*, Zopf. *Pneumobacillus friedländeri*. See *Bacillus pneumoniae friedländeri*. *Pneumobacillus liquefaciens bovis*. See *Bacillus liquefaciens bovis*, Arloing. *Pneumococcus*, Friedländer's; *Pneumoniococcus* of Friedländer, Sternberg. See *Bacillus pneumoniae friedländeri*. *Potato Bacillus*. Applied to *Bacillus mesentericus fuscus*, Flügge. *Bacillus mesentericus vulgaris*, Flügge. *Bacterium mesentericum rubrum*, Globig. *Pragmidiothrix multisepta*, Engler. Syn., *Beggiatoa multisepta*. Found in sea-water upon crustaceans (Kieler Bucht). It is distinguished from *Beggiatoa* by the absence of sulphur grains. Prazmowski's *Bacillus* of Butyric Acid Fermentation. See *Bacillus butyricus*, Botkin, Prazmowski. *Proteus capsulatus septicus*, *Proteus hominis capsulatus*. See *Bacillus hominis capsulatus*, Banti, Bordoni-Uffreduzzi. *Proteus lethalis*. See *Bacillus lethalis*, Babes. *Proteus mirabilis*. See *Bacillus mirabilis*, Hauser. *Proteus* of Karlinski. See *Bacillus murisepticus pleomorphus*, Karlinski. *Proteus septicus*. See *Bacillus septicus*, Pasteur. *Proteus sulfureus*. See *Bacillus sulfureus*, Lindenborn. *Proteus sulfureus* of Holochenikoff, *Proteus vulgaris*, Hauser. See *Bacillus vulgaris*, Hauser. *Proteus zenkeri*. See *Bacillus zenkeri*, Hauser. *Pseudodiphtheria* *Bacillus*, Löffler, Von Hoffmann, Roux, Yersin, and Abbott. Found in the mucus from the pharynx and tonsils of children suffering from nondiphtheric throat affections. *Pseudodiplococcus pneumoniae*, Bonome. Obtained from the serofibrinous exudate of a person dead of cerebrospinal meningitis. *aer.*, *liqf.*, *fg.* Cf. *Micrococcus meningitidis*, Neumann, and *Bacillus meningitidis purulenter*, Neumann. *Rasmussenia buccalis*, Saccardo. See *Leptothrix buccalis*, Robin. *Rauschbrand-bacillus*. See *Bacillus chauvoei*, Bollinger and Feser. *Red Bacillus of Water*. See *Bacillus ruber*, Frank and Becker. *Saccharomyces albicans*, Robin. Syn., *Oidium albicans*, Robin; the cause of thrush. *Saccharomyces apiculatus*, Reess. Occurs in fermented wine and beer, and, in the hot seasons, on sweet succulent fruits; in winter, in the soil beneath the trees that bear these fruits. *Saccharomyces cerevisiae*, Reess and Robin. Syn., *Cryptococcus cerevisiae*, Kütz- ing, *Cryptococcus fermentatum*, torula *cerevisiae*, yeast plant. Under this heading are included a series of different yeasts. *Saccharomyces conglomeratus*, Reess. As the result of Hansen's investigations this

species has been dropped by recent writers. *Saccharomyces ellipsoideus* I, II, Hansen, a "wild" species of wine ferment. *Saccharomyces exiguus*, Hansen. Found in German yeast; acts toward sugar exactly as does *Saccharomyces marxianus*. *Saccharomyces glutinis*. Syn., *Cryptococcus glutinis*, Fersen. A pullulating yeast that forms beautiful rose-colored patches on cooked potatoes. *Saccharomyces guttulatus*, Robin. Syn., *Cryptococcus guttulatus*, Robin, Winter. Found in the esophagus, stomach, and intestines of mammals, birds, and reptiles. The feces of patients suffering from tunnel anemia, or anchylostomiasis, frequently contain this species of *saccharomyces*. *Saccharomyces lactis*, Adametz. Syn., *Milk-yeast*. Found in milk, of which it ferments the milk-sugar. *Saccharomyces marxianus*. Found in wine; acts vigorously on saccharose, inverting and fermenting it with great activity; it also acts on dextrose. *Saccharomyces membrani-faciens*. Forms a bright yellow, tough scum on beer-wort; liquefies gelatin; has no fermentative action on ordinary carbohydrates, and does not invert cane-sugar. *Saccharomyces minor*, Engel. According to Engel this is the cause of fermentation in bread. *Saccharomyces mycoderma*, Reess. Syn., *Mycoderma mesentericum*, Pers. *Mycoderma cerevisiae et vini*, Desm. *Hormiscium vini et cerevisiae*, Bonard. *Mycoderma vini*, Pasteur. *Mycoderma cerevisiae*, Trecul. Found on the surface of wine and beer, forming the so-called "flowers of wine." *Saccharomyces pasteurianus*, Reess. According to Hansen, this represents a group of three forms that cause changes in beer and wine. *Saccharomyces pyriformis*, Ward. Found in the fermentation of ginger-beer. A yeast which develops much CO₂ but little alcohol; it inverts cane-sugar and ferments the products; does not ferment milk-sugar. *aer.*, *facanaer.* *Saccharomyces ruber*, Demme. Found in milk and cheese, forming red spots in the latter, and red sediment in the former. *liqf.* *Saccharomyces vacciniæ*, Pfeiffer. Found in the lymph of a vaccinia pustule; nothing is known of its action. *Sarcina alba*, Eisenberg. Found in air and water. *aer.*, *liqf.*, *nfg.* *Sarcina aurantiaca*. Syn., Orange *Sarcina*, Koch. Found in air and water. *aer.*, *liqf.*, *chg.* (orange yellow), *nfg.* *Sarcina aurea*, Macé. Found in the pulmonary exudates of a person dead of a pneumonia complicated with purulent pleurisy. *aer.*, *liqf.*, *chg.* (brilliant golden yellow). *Sarcina botulina*, Van den Corput. Found in cases of poisoning with tainted sausage (*botulismus*) and believed to be the active

agent. Cf. *Bacillus of allantiasis*. *Sarcina candida*, Reinke. Found in the air of breweries. *aer.*, *liqf.*, *npg.* *Sarcina cerevisiae*, Lindner. Syn., *Pediococcus cerevisiae*, Balcke. Found in beer and the air of breweries. *aer.*, *facanaer.*, *nliqf.*, *npg.* *Sarcina flava*, De Bary. Found in beer. *aer.*, *liqf.*, *chg.* (yellow), *npg.* *Sarcina fuscescens*, Falkenheim. Found in the human stomach. *Sarcina hyalina*, Kützing. Syn., *Merismopedia hyalina*, Kützing. Found in swamps. *Sarcina intestinalis*, Zopf. Found in the intestines of chickens and turkeys, principally in the cecum. *Sarcina litoralis*, CErstedt. Syn., *Merismopedia litoralis*, Rabenhorst, *Erythroconis*, CErstedt. Found in sea-water; forms a thin reddish layer between decomposing algae on the shore. *Sarcina lutea*, Schröter. Found in the air and in fungus cultures. *aer.*, *liqf.*, *chg.* (yellow), *npg.* *Sarcina minuta*, De Bary. Occurred spontaneously in a culture of sour milk on a microscopic slide; resembles *Sarcina welckeri*. *Sarcina mobilis*, Maurea. Found in old ascitic fluid. *aer.*, *liqf.*, *mot.*, *chg.* (brick-red). *Sarcina morrhuae*. Obtained from codfish. *Sarcina* of the Lungs. See *Sarcina pulmonum*, Hauser. *Sarcina* of the Urine. See *Sarcina urinae*, Welcker. *Sarcina paludosa*, Schröter. Found in bog water. *Sarcina pulmonum*, Hauser. Syn., *Sarcina* of the lungs. Found in sputum. *aer.*, *nliqf.*, *npg.* *Sarcina reitenbachii*, Caspary. Syn., *Merismopedium reitenbachii*. Found in fresh water on decaying plants. *Sarcina renis*, Hepworth. Found in the lungs of persons dead of pulmonary tuberculosis. *Sarcina rosea*, Menge. Found in milk and other food-stuffs. Held to be distinct from the form described by Schröter. *aer.*, *liqf.*, *chg.* (rose-red), *npg.* *Sarcina rosea*, Schröter. Found in the air. *aer.*, *liqf.*, *chg.* (intense red, giving same reaction as the pigment of *Sarcina aurantiaca*), *npg.* *Sarcina urinæ*, Welcker. Syn., *Merismopedia urinæ*, Rabenhorst. Found in the bladder; appears to be destitute of zymotic action. *Sarcina ventriculi*, Goodsir. Common in the stomach of man and animals. *aer.*, *nliqf.*, *npg.* *Sarcina welckeri*, Rossmann. Repeatedly found in the human bladder and voided for months at a time in the urine, which is usually abnormally rich in phosphates, the patient retaining good health. *Scheutzia laughlini*. Syn., *Micrococcus* of Dengue, *Staphylococcus* of Dengue, *Dengue Micrococcus*. Found in the blood of patients suffering from dengue. Pathogenetic relations not determined. *Schweinerotlaufbacillus*. See *Bacillus erysipelatos suis*, Koch.

Smegma bacillus. See *Bacillus* of the smegma. *Sphærococcus acidi lactici*, Marpmann. Found in fresh cow's milk. *aer.*, *nliqf.*, *npg.* *Sphærotilus natans*, Kützing, Saccardo. Syn., *Leptothrix natans*. Found in running and stagnant water. Old portions yellowish, younger portions colorless. *Spirillum amyliiferum*, Van Tieghem. Found in water. Forms granulose at certain developmental stages; gives the blue reaction with iodine. Cf. *Bacillus butyricus*, Prazmowski. *Spirillum anserum*. Syn., *Spirochate anserina*, Sakharoff. Found in the blood of geese affected with a fatal form of septicemia due to this microbe. A disease prevalent in the swampy regions of Caucasia. *aer.*, *mot.*, *pg.* (not for chickens). *Spirillum* of Weibel. See *Spirillum saprophiles* α, Weibel. *Spirillum attenuatum*, Warming. Found in sea water. *Spirillum aureum*. Syn., *Vibrio aureus*, Weibel. Found in the air and slime of sewers. *aer.*, *nliqf.*, *nmot.*, *chg.* (golden yellow), *npg.* *Spirillum* β of Weibel. See *Spirillum saprophiles* β, Weibel. *Spirillum cholerae asiaticæ*, Koch. Syn., *Spirillum cholerae*. *Spirillum* of cholera, *Bacillus* of cholera, *Comma bacillus* of Koch. Found in the discharges of cholera patients and in the intestines of cholera cadavers. *aer.*, *facanaer.*, *liqf.*, *mot.*, *pg.* *Spirillum concentricum*, Kitasato. Found in putrefying blood. *aer.*, *nliqf.*, *mot.*, *npg.* *Spirillum denticola*, Miller; or, *Spirillum dentium*, Arnt. Syn., *Spirochate denticola*, *Spirochate dentium*. Found under the margins of the gums in the mouths of healthy persons; biologic characters unknown, as it does not thrive in culture fluids. *Spirillum endoparagogenicum*, Sorokin. Found in a glairy, liquid exuding from a poplar tree. *Spirillum finkleri*, *Spirillum Finkler-Prior*. Syn., *Vibrio proteus*. *Bacillus* of Finkler-Prior. Found in the feces of persons affected with cholera nostras, after allowing the discharge to stand for some time. *aer.*, *facanaer.*, *liqf.*, *mot.* Pathogenetic relation not confirmed. *Spirillum flavescens*. Syn., *Vibrio flavescens*, Weibel. Found in the slime of sewers. *aer.*, *nliqf.*, *nmot.*, *chg.* (yellowish-green), *npg.* *Spirillum flavum*. Syn., *Vibrio flavescens*, Weibel. Found in the slime of sewers. *aer.*, *nliqf.*, *nmot.*, *chg.* (ocher yellow), *npg.* *Spirillum jenensis*, Ehrenberg. Syn., *Ophidomonas jenensis*, Ehrenberg. A doubtful species. *Spirillum leukomelænum*, Perty. Found in water containing decaying algae. *Spirillum linguae*. Syn., *Vibrio lingualis*. Found in the deposit on the tongue of a mouse, after inoculation. *aer.*, *facanaer.*, *nliqf.*, *nmot.*, *npg.* (for mice). *Spirillum litorale*, Warm-

ing. Syn., *Vibrio spirillum*, Müller; *Melunella spirillum*, Borg. Found in bog water. **Spirillum metschnikovi**. Syn., *Vibrio metschnikovi*, Gameleia. Found in the intestinal contents of chickens affected with an epidemic infectious disease resembling fowl cholera. aer., facanaer. (?), liqf., mot., pg. **Spirillum nasale**. Syn., *Vibrio nasale*, Weibel. Found in nasal mucus. aer., facanaer., nliqf., nmot., npg. **Spirillum obermeieri**, Cohn. Syn., *Spirochæta obermeieri*. Spirillum of relapsing fever. Found in the blood of persons affected with relapsing fever. aer., mot., (efforts to cultivate thus far unsuccessful), pg. **Spirillum of Asiatic cholera**. Spirillum of cholera. See *Spirillum cholerae asiaticæ*, Koch. **Spirillum of Denecke**. See *Spirillum tyrogenum*, Denecke. **Spirillum of Finkler and Prior**. See *Spirillum finkleri*. **Spirillum of Müller**. Syn., Miller's bacillus. Found in carious teeth. aer., facanaer., liqf., mot., npg. **Spirillum of Relapsing Fever**. See *Spirillum obermeieri*, Cohn. **Spirillum of Smith**. Found in the intestines of swine. aer., nliqf., mot., npg. **Spirillum plicatile**, Dujardin. Syn., *Spirochæta plicatilis*, Ehrenberg; *Spirulina plicatilis*, Cohn. Found in water containing decomposing algæ. Biologic characters undetermined. **Spirillum rosaceum**. Observed upon excrement in water. aer., chg. (red), npg. **Spirillum rosenbergii**, Warming. Found in brackish water. Contains refractive sulphur grains. aer., mot. **Spirillum roseum**, Macé. Found in a tube culture of blennorrhagic pus. aer., nliqf., mot., chg. (rose-red), npg. **Spirillum rubrum**, von Esmarch. Obtained from the putrefying body of a mouse. aer., facanaer., nliqf., mot., chg. (pink), npg. **Spirillum rufum**, Perty. Found in cistern water. aer., mot., chg. (rose-red), npg. **Spirillum rugula**, Müller. Syn., *Vibrio rugula*. Found in swamp water and frequently in the alimentary canal, on the teeth, etc. aer., facanaer., mot., npg. According to Prazmowski this species gives rise to decomposition of cellulose. **Spirillum sanguineum**. See *Beggiatoa roseopersicina*, Zopf. **Spirillum saprophiles a** of Weibel. Syn., *Vibrio saprophiles a* of Weibel. Found in rotting hay infusion and in the slime of sewers. aer., nliqf., mot., npg. **Spirillum saprophiles β** of Weibel. Syn., *Vibrio saprophiles β*, Weibel. Found in rotting hay infusion. aer., nliqf., mot., npg. **Spirillum saprophiles γ** of Weibel. Syn., *Vibrio saprophiles γ*, Weibel. Found in the slime of sewers. aer., nliqf., mot., npg. **Spirillum serpens**, Müller. Syn., *Vibrio serpens*, Müller. Found in stagnant water

containing decaying plants, etc. Biologic characters not determined. **Spirillum sputigenum**, Miller. Common in the neglected mouths of healthy persons. Does not grow on culture media. **Spirillum tenue**, Ehrenberg. Found in putrefying plant infusions. Biologic characters undetermined. **Spirillum tyrogenum**, Denecke. Spirillum of Denecke, Bacillus of cheese, Denecke's cheese spirillum. Found in old cheese. aer., facanaer., liqf., mot., pg. (for guinea pigs). **Spirillum undula**, Cohn, Müller. Syn., *Vibrio undula*, Müller; *Vibrio prolifer*, Ehrenberg. Found in zooglea masses in swamp water, vegetable infusions, etc. aer., mot., npg. **Spirillum violaceum**, Warming. Found in brackish water; cell contents violet, with a few sulphur grains. **Spirillum volutans**, Ehrenberg. Syn., *Vibrio spirillum*, Müller; *Melunella spirillum*, Borg. Found in infusions and in bog water. Biologic characters undetermined. **Spirochæta anserina**, Sakharoff. See *Spirillum anserum*. **Spirochæta denticola**, Miller, or *Spirochæta dentium*, Arnt. See *Spirillum dentium*. **Spirochæta obermeieri**. See *Spirillum obermeieri*, Cohn. **Spirochæta plicatilis**, Ehrenberg. See *Spirillum plicatile*, Dujardin. **Spiromonas cohnii**, Warming. Found in foul water. **Spiromonas volubilis**, Pertz. Found in putrefying infusions and in bog water. **Sporonema gracile**. See *Bacillus virens*, Van Tieghem. **Staphylococcus albus liquefaciens**, Escherich. Syn., White liquefying staphylococcus. Found in the alvine discharges of healthy infants. aer., liqf., npg. **Staphylococcus biskrae**. See *Micrococcus of Heydenreich*. **Staphylococcus cereus albus**, Passet. See *Micrococcus cereus albus*, Passet. **Staphylococcus cereus aureus**, Schröter and Winkler. See *Micrococcus cereus aureus*, S. and W. **Staphylococcus cereus flavus**, Passet. See *Micrococcus cereus flavus*, Passet. **Staphylococcus citreus**. See *Micrococcus citreus*, List. **Staphylococcus epidermidis albus**, Welch. See *Micrococcus pyogenes albus*, Rosenbach. **Staphylococcus of Dengue**, McLaughlin. See *Scheutisia laughlini*. **Staphylococcus of pemphigus**, De Michele. Obtained from the skin, kidney, and spleen of a case of pemphigus chronica. **Staphylococcus passetti**. See *Micrococcus cereus flavus*, Passet. **Staphylococcus pyogenes albus**, Rosenbach. See *Micrococcus pyogenes albus*, Rosenbach. **Staphylococcus pyogenes aureus**, Ogston, Becker. See *Micrococcus pyogenes aureus*, Rosenbach. **Staphylococcus pyogenes citreus**, Passet. See *Micrococcus pyogenes citreus*, Passet.

Staphylococcus pyosepticus, Hericourt and Richet. See *Micrococcus pyosepticus*, Richet. **Staphylococcus salivarius pyogenes**, Biondi. See *Micrococcus salivarius pyogenes*, Biondi. **Staphylococcus viridis flavescens**, Guttman. Syn., *Micrococcus* of chicken-pox. Found in the vesicles of varicella. *aer.*, *nliqf.*, *chg.* (greenish-yellow), *npg.* **Streptococcus acidilactici**, Grotenfeldt. Found in coagulated milk in Finland. *anaer.* (not strictly), *nliqf.*, *npg.* **Streptococcus aethiobius**. See *Micrococcus ureæ liquefaciens*, Flüge. **Streptococcus albus**, Maschek. Found in hydrant water (Freiburg). *aer.*, *liqf.*, *npg.* **Streptococcus aphicola**, Hallier. See *Streptococcus* of eczema epizootica, Schottelius. **Streptococcus articulorum**, Löfller. Syn., *Micrococcus* of diphtheria. Obtained from the affected mucous membrane in cases of diphtheria. Possibly a variety of *Micrococcus pyogenes*. *pg.* (for mice). **Streptococcus bombycis**. See *Micrococcus bombycis*, Cohn. **Streptococcus brevis**, Von Lingelsheim. Obtained from normal human saliva. *aer.*, *facanaer.*, *nliqf.*, *npg.* **Streptococcus cadaveris**, Sternberg. Found in the liver of a yellow-fever cadaver. *aer.*, *facanaer.*, *nliqf.*, *npg.* Probably identical with the preceding species. **Streptococcus cinnabareus**. See *Micrococcus cinnabareus*, Flüge. **Streptococcus coli gracilis**, Escherich. Found in the feces of healthy children living on a meat diet. *aer.*, *facanaer.*, *liqf.*, *npg.* **Streptococcus conglomeratus**, Kunth. Obtained from cases of scarlet fever. *pg.* (for mice). **Streptococcus coronatus**, Flüge. See *Micrococcus coronatus*, Flüge. **Streptococcus coryzæ contagiosæ equorum**, Schütz. Found in the pus from affected lymphatic glands in horses suffering with the disease known in Germany as *Drüse des Pferdes*. *aer.*, *facanaer.*, *pg.* (for horses and mice), *npg.* (for rabbits, guinea pigs, and pigeons). **Streptococcus endocarditicus**. See *Micrococcus endocarditicus rugatus*, Weichselbaum. **Streptococcus erysipelatis** (*erysipelatos*), Fehleisen. See *Micrococcus pyogenes*, Rosenbach. **Streptococcus fermenti**. See *Micrococcus viniperda*. **Streptococcus giganteus urethræ**, Lustgarten and Manneberg. Found in the healthy human urethra. *aer.*, *npg.* **Streptococcus havaniensis**, Sternberg. Found in the acid vomit of a yellow-fever patient. Biologic characters undetermined. **Streptococcus insectorum**, Burrill. See *Micrococcus insectorum*, Burrill. **Streptococcus kochii**. See *Micrococcus* of septicemia in rabbits, Koch. **Streptococcus lanceolatus pasteurii**, Gamaléia. See *Bacillus septicus sputigenus*, Flüge. **Streptococcus liquefaciens**,

Sternberg. Obtained from the intestines and livers of yellow-fever cadavers. *aer.*, *facanaer.*, *liqf.*, *npg.* **Streptococcus longus**, von Lingelsheim. See *Micrococcus pyogenes*, Rosenbach. **Streptococcus lucæ**. See *Micrococcus ulceris mollis*. **Streptococcus manfredii**. See *Micrococcus manfredii*. **Streptococcus meningitidis**. See *Micrococcus intercellularis meningitidis*, Naumann and Schäffer. **Streptococcus monomorphus**, Buijwid, Heryng. Found in benign pharyngeal ulcers. *npg.* **Streptococcus morbillosus**. See *Micrococcus* of measles. **Streptococcus nocardii**. See *Micrococcus mastobius*, Nocard. **Streptococcus ochroleukus**. See *Micrococcus ochroleukus*, Prove and Legrain. **Streptococcus of Bonome**. Obtained from the exudations in the cerebrospinal meninges and from hemorrhagic extravasations in the lungs, in cases of epidemic cerebrospinal meningitis. *pg.* (for white mice and rabbits). **Streptococcus of granuloma fungoides**, Rindfleisch and Auspetz. See *Streptococcus* of mycosis fungoides. **Streptococcus of Manneberg**. Obtained from the urine in cases of acute nephritis. *aer.*, *facanaer.*, *nliqf.*, *pg.* (for dogs and rabbits). **Streptococcus of Mastitis in Cows**. See *Micrococcus mastobius*, Nocard and Mollereau. **Streptococcus of Mycosis fungoides**. *Streptococcus* of granuloma fungoides, Rindfleisch and Auspetz. Found in the capillaries and subcutaneous tissue in the parts affected by this disease. Not sufficiently investigated. **Streptococcus of Pus**. See *Micrococcus septicus*, Flüge. **Streptococcus perniciosus psittacorum**, Eberth and Wolf. See *Micrococcus psittaci*, Wolf. **Streptococcus pneumoniae**, Weichselbaum. See *Micrococcus pyogenes*, Flüge. **Streptococcus putrefaciens**. *Micrococcus* of putrefaction. Refers to any one of the many micrococci found in putrefying substances. **Streptococcus pyogenes**, Rosenbach. See *Micrococcus pyogenes*, Rosenbach. **Streptococcus pyogenes**, Saccardo. See *Micrococcus septicus*, Flüge. **Streptococcus pyogenes maligni**, Krause and Flüge. Found in necrotic masses in a leukemic spleen. *pg.* (for mice and rabbits). **Streptococcus pyogenes ureæ**, Rösing. Found in purulent urine. Cf. *Micrococcus pyogenes ureæ flavus*, Rösing. **Streptococcus radiatus**. See *Micrococcus radiatus*, Flüge. **Streptococcus rubiginosus**, Edington. Found in cases of scarlatina. *npg.* Does not appear till late in the disease. **Streptococcus seiferti**. See *Micrococcus influenzae*, Letzerich. **Streptococcus septicus**, Flüge. See *Micrococcus septicus*, Flüge. **Streptococcus septicus liquefaciens**, Babes. Obtained from the blood and viscera

of a child that died of septicemia following scarlatina. *aer.*, *liqf.*, *fg.* (for mice and rabbits). **Streptococcus septopyæmicus**. See *Micrococcus septopyæmicus*, Biondi. **Streptococcus toxicatus**. See *Micrococcus toxicatus*, Burrill. **Streptococcus ureæ**. See *Micrococcus ureæ*, Cohn. **Streptococcus ureæ rugosus**, Rösing. Found in urine. **Streptococcus vermiformis**, Masehek. Found in hydrant water (Freiburg). *aer. liqf.*, *nfg.* **Streptocytus of Eczema epizootica**, Schottelius. Syn., *Streptococcus aphthicola*, Hallier. Micrococcus of foot-and-mouth disease. Found in vesicular eruptions about the mouth and feet of cattle, pigs, and sheep. Communicable to man. *aer.*, *facanaer.* Pathogenesis not fully settled. Cf. *Bacillus* of ulcerative stomatitis of cattle. **Streptothrix foersteri**, Cohn. See *Cladothrix foersteri*, Cohn. **Swine Plague Schizophyte**, Detmers. See *Bacillus* of swine plague, Marseilles, Rietsch, and Jobert. **Syphilis Bacillus**. See *Bacillus syphilidis*, Lustgarten. **Syphilis Bacillus of Eve and Lingard**. Obtained from the blood and diseased tissues of syphilitic persons who have not undergone mercurial treatment. *aer.*, *chg.* (pale yellow or brown). Pathogenetic relations not settled. Not stained by Lustgarten's method; capable of development on blood-serum. **Tetanus Bacillus**. See *Bacillus tetani*, Nicolaier. **Thiothrix nivea**. See *Beggiatoa nivea*, Rabenhorst. **Thiothrix tenuis**, Winogradsky. Found in sulphur springs. **Thiothrix tenuissima**, Winogradsky. Found in sulphur springs. **Torula cerevisiæ**. See *Saccharomyces cerevisiæ*, Reess and Robin. **Torula ureæ**, Pasteur. See *Micrococcus ureæ*, Cohn. **Tubercle bacillus**. See *Bacillus tuberculosis*, Koch. **Typhus bacillus**. See *Bacillus typhi abdominalis*, Eberth. **Tyrophthrix catenula**. **T. claviformis**. **T. distortus**. **T. filiformis**. **T. geniculatus**. **T. scaber**. **T. tenuis**. **T. turgidus**. **T. urocephalus**. **T. virgula**. See the corresponding terms under *Bacillus*. **Ulvina aceti**, Kützing. See *Bacillus aceti*, Kützing. **Urobacillus duclauxi**, Miquel. Found in sewage and river water. *aer.*, *facanaer.*, *mot.*, *liqf.* Pathogenesis not determined. **Urobacillus freudenreichii**, Miquel. Obtained from air, dust, sewage. *aer.*, *liqf.*, *mot.* Pathogenesis not determined. **Urobacillus maddoxi**, Miquel. Obtained from sewage and river water. *aer.*, *liqf.*, *mot.* Pathogenesis not determined. **Urobacillus pasteurii**, Miquel. Obtained from decomposing urine. *aer.*, *liqf.*, *mot.* Pathogenesis not determined. **Urobacillus schutzenbergii**, Miquel. Obtained from sewage and river water. *aer.*, *liqf.*, *mot.* Pathogenesis not determined.

Vibrio aureus, Weibel. See *Spirillum aureum*. **Vibrio bacillus**, Müller, Ehrenberg. See *Bacillus ulna*, Cohn. **Vibrio cholerae**. See *Spirillum cholerae asiatica*, Koch. **Vibrio cyanogenus**, Ehrenberg. See *Bacillus cyanogenus*, Fuchs. **Vibrio flavescens**, Weibel. See *Spirillum flavescens*, Weibel. **Vibrio flavus**, Weibel. See *Spirillum flavum*, Weibel. **Vibrio lineola**, Müller and Ehrenberg. See *Bacterium lineola*, Cohn. **Vibrio lingualis**. See *Spirillum linguar*, Weibel. **Vibrio metchnikovi**, Gamaleia. See *Spirillum metchnikovi*, Gamaleia. **Vibrio nasalis**, Weibel. See *Spirillum nasale*, Weibel. **Vibrio proteus**. See *Spirillum finkleri*. **Vibrio rugula**, Müller. See *Spirillum rugula*, Müller. **Vibrio saprophiles a**, Weibel. See *Spirillum saprophiles a*, Weibel. **Vibrio saprophiles β**, Weibel. See *Spirillum saprophiles β*, Weibel. **Vibrio saprophiles γ**, Weibel. See *Spirillum saprophiles γ*, Weibel. **Vibrio serpens**, Müller. See *Spirillum serpens*, Müller. **Vibrio syncyanus**, Ehrenberg. See *Bacillus cyanogenus*, Fuchs. **Vibrio synxanthus**, Ehrenberg. See *Bacillus synxanthus*, Ehrenberg and Schröter. **Vibrio tremulans**, Ehrenberg. See *Bacterium lineola*, Cohn. **Vibrio undula**, Müller. See *Spirillum undula*, Müller. **Vibrio Xanthogenus**, Fuchs. *Bacillus synxanthus*, Ehrenberg and Schröter. **Violet Bacillus**. See *Bacillus janthinus*, Zopf. **Zooglæa capillorum**, Buhl. See *Palmellina capillorum*, Radlkofer. **Zooglæa termo**, Müller, Cohn. See *Bacillus termo*, Dujardin. **Zooglæa ramigera**. A tree-like variety of *Cladothrix dichotoma*. **Zoogalactina imetropa**, Sette. See *Bacillus prodigiosus*, Ehrenberg.

Bacteriaceæ (*bak-te-re-ä'-se-e*) [*βακτηριον*, a little stick]. The Schizomycetes.

Bacterial (*bak-te'-re-äl*) [*βακτηριον*, a little stick]. Resembling, of the nature of, or derived from bacteria.

Bactericidal (*bak-te-ris-i'-dal*) [*βακτηριον*, a little stick; *cädere*, to kill]. Destructive to bacteria.

Bactericide (*bak-te'-ris-id*) [*βακτηριον*, a little stick; *cädere*, to kill]. 1. Destructive to bacteria. 2. An agent that destroys bacteria.

Bacteridium (*bak-ter-id'-e-nm*) [*βακτηριον*, a little stick]. According to Davaine, a genus of *Bacteriaceæ* characterized by immobility of the elements at all periods of their existence. The distinction does not now obtain.

Bacterioid (*bak-te'-re-oid*) [*βακτηριον*, a little stick; *oidos*, form]. Applied to certain microorganisms flourishing in and around the roots of leguminous plants, assisting in the accumulation of nitrogen, and giving rise to

tubercles. They may be cultivated in artificial media like ordinary bacteria.

Bacteriologist (*bak-te-re-ol'-o-jist*) [*βακτηριολογία*, a little stick; *λόγος*, science]. One who makes a special study of bacteriology.

Bacteriology (*bak-te-re-ol'-o-je*) [*βακτηριολογία*, a little stick; *λόγος*, science]. The science concerned with the study of bacteria.

Bacterioprotein (*bak-te-re-o-pro'-te-in*) [*βακτηριοπρωτεΐν*, a little stick; *πρωτός*, first]. A protein contained in bacteria.

Bacteriopurpurin (*bak-te-re-o-per'-pu-rin*) [*βακτηριοπυρρίν*, a little stick; *purpura*, purple]. A peach-colored pigment found by Lankester in the protoplasm of *Beggiatoa roseopersicina*. It is insoluble in water, alcohol, chloroform, ammonia, acetic and sulphuric acids.

Bacterioscopic (*bak-te-re-o-skop'-ik*) [*βακτηριοσκοπικός*, a little stick; *σκοπεῖν*, to view]. Pertaining to bacterioscopy.

Bacterioscopy (*bak-te-re-o-skop'-e*) [*βακτηριοσκοπία*, a little stick; *σκοπεῖν*, to view]. The microscopic study of bacteria.

Bacteriotherapy (*bak-te-re-o-ther'-ap-e*) [*βακτηριοθεραπεία*, a little stick; *θεραπεία*, treatment]. The treatment of disease by the introduction of bacteria into the system.

Bacterium (*bak-te'-re-um*) [*βακτήριον*, a little stick; dim. of *βακτηρία*, a stick, staff: *pl.*, *Bacteria*]. 1. A unicellular vegetable microorganism. The word is a generic term and includes the groups bacillus, coccus, and spirillum. 2. Formerly, a genus of fungi characterized by inflexible linear filaments.

Bacteroid (*bak'-ter-oid*) [*βακτηριοειδής*, a little stick; *εἶδος*, form]. Resembling bacteria.

Bael (*ba'-el*). See *Bela*.

Bag (*bag*) [A.S., *bæc*]. 1. A sac. 2. The scrotum. **B.**, Barnes's. A rubber bag shaped somewhat like a lyre, used for dilating the cervical canal. **B.**, Politzer's, a soft rubber bag for inflating the middle ear. **B. of Waters**, the fetal membranes enclosing the liquor amnii and projecting through the *os uteri* early in labor. The sac usually ruptures when the cervix is dilated.

Bagnio (*ban'-yo*) [It., *bagno*]. 1. A house of prostitution. 2. A bath-house.

Bakers' Itch. An eczematous affection of the hands, caused by the irritation of the yeast.

B. Leg, knock-knee, or genu valgum. **B. Salt**, a synonym of smelling salts, or the carbonate of ammonium; it is sometimes used by bakers in leavening cakes.

Balance (*bal'-ans*) [*bilanx*, having two scales]. 1. An instrument for weighing.

2. The harmonious adjustment of related parts.

Balanic (*bal'-an-ik*) [*βάλανος*, the glans penis]. Pertaining to the glans of the penis or of the clitoris.

Balanitis (*bal-an-i'-tis*) [*βάλανος*, the glans

penis; *τις*, inflammation]. Inflammation of the glans penis.

Balanoblenorrhoea (*bal-an-o-blen-o-re'-ah*) [*βάλανος*, the glans; *βλέννα*, mucus; *ῥοία*, a flow]. Gonorrheal balanitis.

Balanoplasty (*bal-an'-o-plas-te*) [*βάλανος*, the glans penis; *πλασσειν*, to form]. Plastic surgery of the glans penis.

Balanoposthitis (*bal-an-o-pos-thi'-tis*) [*βάλανος*, the glans penis; *πόσθη*, prepuce; *τις*, inflammation]. Inflammation of the glans penis and of the prepuce.

Balantidium (*bal-an-tid'-e-um*) [*βιλαντιδίων*, dim. of *βιλάντιον*, a bag]. A genus of protozoa. **B. coli**, a protozoan parasite that inhabits the intestine of the pig.

Balbuties (*bal-bu'-she-ēz*) [*balbutire*, to stammer]. Stammering.

Baldness (*bal-d'-nes*) [ME., *balde*, bald]. Loss of hair; alopecia.

Ball (*baul*) [ME., *bal*]. An object having a round or spheric shape. **B.-and-Socket Joint**. See *Diarthrosis* and *Euarthrosis*.

B. Thrombus. See *Thrombus*.

Ballottement (*bal-ot'-men-t*) [Fr., from *ballotte*, a ball]. A method of diagnosing pregnancy from the fourth to the eighth month. A push is given the uterus by the finger inserted into the vagina, and if the fetus be present, it will rise and fall again like a heavy body in water.

Balm (*balm*) [*balsamum*, a balsam]. 1. A popular synonym of *balsamum*. 2. Any soothing application or ointment. **B. of Gilead**. See *Balsam*, *Abies*, and *Melissa*.

Balneography (*bal-ne-og'-ra-fe*) [*balneum*, a bath; *γραφω*, a writing]. A treatise on bathing and baths.

Balneology (*bal-ne-ol'-o-je*) [*balneum*, a bath; *λόγος*, science]. The science of baths and their effects upon the system.

Balneotherapy (*bal-ne-o-ther'-ap-e*) [*balneum*, a bath; *θεραπεία*, treatment]. Systematic bathing for therapeutic purposes, or the treatment of disease by baths.

Balneum (*bal'-ne-um*) [L.]. A bath. See *Bath*. **B. arenae**, a sand-bath. See *Am-motherapy*. **B. lacteum**, a milk-bath. **B. luteum**, a mud-bath. **B. pneumaticum**, an air-bath.

Balsam (*ba-sam*) [*βάλσαμον*, the resin of the balsam tree]. The resinous, volatile, aromatic substance, liquid or solid, obtained from certain trees by natural exudation or by artificial extraction. Balsams are divided into two classes, those with, and those without benzoic and cinnamic acids. In general they are mixtures of various essential oils, resins, and acids. **B. Apple**, the plant *Momordica balsamina*, and its warty, gourd-like fruit. It is purgative, but its tincture is chiefly used in domestic medicine as a vul-

nerary. Unof. **B. Bog**, a singular stone-like, woody, umbelliferous plant, *Bolax*, or *Asorella glebaria*, of the Falkland Islands and Patagonia. Its aromatic gum is locally prized as a vulnerary, desiccative, and antigonorrheal remedy. Unof. **B., Canada**, a turpentine gathered from the natural blisters of the bark of *Abies balsamum*. It is used as a mounting-medium by microscopists. **B. of Copaiba**. See *Copaiba*. **B. of Fir**. Same as *Canada Balsam*. **B., Friar's**. See *Benoin*. **B. of Gilead**, the balm of the Old Testament; an oleoresin obtained from the *Balsamodendron gileadense*. **B. of Peru**, the balsam obtained from *Myroxylon pereiræ*; antiseptic, stimulant to the circulation, and sedative to the nervous system, tonic, and expectorant. Applied locally, it is useful in chronic inflammatory skin-diseases. Dose of the emulsion $\text{m}\bar{x}$ -xxv (0.65-1.6). **B. Root**, a popular name for certain composite-flowered plants of the genus *Balsamorhiza*. *B. hookerii*, *B. macrophylla*, *B. sagittata* are common in the Pacific States and abound in a resinous balsam. Unof. **B. of Tolutan**, or of *Tolu*, obtained from *Myroxylon toluifera*. Its properties are due to a volatile oil, *toluene*. It possesses an agreeable odor, and is a basis for many cough-mixtures. It is expectorant. *Tinct. tolutana* contains 10 per cent. of the balsam in 90 per cent. of alcohol. Dose $\text{m}\bar{x}$ -xxx (0.65-2.0). *Syr. tolutanus*. Dose $\text{f}\bar{5}\text{j}$ -ij (4.0-8.0).

Balsamation (*bawel-sam-a'-shun*) [$\beta\alpha\lambda\sigma\alpha\mu\omega\nu$, the resin of the balsam tree]. Embalment with balsamic or aromatic spices.

Balsamic (*bawel-sam'-ic*) [$\beta\alpha\lambda\sigma\alpha\mu\omega\nu$, the resin of the balsam tree]. Having the nature or qualities of a balsam.

Balsamodendron (*bawel-sam-o-den'-dron*). See *Myrrh* and *Bdellium*.

Balsamum (*bal'-sam-um*) [$\beta\alpha\lambda\sigma\alpha\mu\omega\nu$, balsam]. A balsam. **B. dipterocarpi**. See *Gurjun Balsam*.

Balser's Fat-Necrosis. See *Diseases, Table of*.

Bamberger's Fluid. An albuminous mercuric compound used in the treatment of syphilis. It is made as follows: To 100 c.c. of a filtered solution of white of egg (containing 40 c.c. of albumin and 60 c.c. of water) there are added 60 c.c. of a solution of mercuric chlorid (containing 5 per cent., or 3 grams, of bichlorid of mercury) and 60 c.c. of a solution of sodium chlorid (containing 20 per cent.); finally, 80 grams of distilled water are added, which brings the bulk of the solution up to 300, containing 0.010 mercuric chlorid in every cubic centimeter.

Bamboo (*bam-boo'*) [*E. Ind., bambu*]. A

popular name for many tree-like, woody-stemmed grasses, especially those of the genus *Bambusa*. *B. arundinacea* is employed as an alterative, anthelmintic, and depurative. Unof. **B. Brier**, the root of *Smilax sarsaparilla*, habitat, Southern United States. Its properties are identical with those of sarsaparilla. Dose of the fld. ext. $\text{f}\bar{3}\text{ss}$ -ij (2.0-8.0). Unof.

Banana (*ban-an'-ah*) [*Sp.*]. The fruit of the common banana, *Musa sapientum*, said to be a valuable alterative, and useful in strumous affections. Dose of the fld. ext. $\text{m}\bar{x}$ -xxx (0.65-2.0). Unof.

Band (*band'*) [*ME., bande*]. That which binds. A stripe. A ligament. **B., Axis**, the primitive streak. **B., Belly**, a flannel band wound around the abdomen. **B., Head**, a strap for securing a mirror to the forehead. **B., Horny**, of *Tarinus*, the fore part of the tania semicircularis. **B., Moderator**. See *B., Reil's*. **B., Reil's**, a fibromuscular fillet that frequently extends across the right ventricle of the heart; it is also called the moderator band. **B. of Remak**. See *Fiber, axial*; also *Purkinje, Axis Cylinder of*. **B. of Tarinus**. See *B., Horny*.

Bandage (*band'-aj*) [*Fr., bande, a strip*].

Bandages are usually strips of muslin or other material, of varying widths and lengths, used in surgery for the purpose of protecting, compressing, etc., a part, or for the retention of dressings and applications. A *simple bandage* or *roller* consists of one piece; a *compound*, of two or more pieces. *Starch, plaster-of-Paris, silica, dextrin, tripolite*, etc., etc., are used for making stiff and *immovable dressings or bandages*. According to their direction bandages are classed as: 1. *Circular*, circular turns about the part. 2. *Figure-of-8*, the turns crossing each other like the figure 8. 3. *Oblique*, covering the part by oblique turns. 4. *Recurrent*, the turns returning successively to the point of origin. 5. *Spica*, the turns resembling the arrangement of the husks of an ear of corn. 6. *Spiral*, each turn covering one-half of the preceding. 7. *Spiral reverse*, the bandage is reversed in order better to adapt it to the part. Bandages are also classed according to the part to which they are applied. **B., Genga's**. See *B., Theden's*. **B., Heliodorus's**, the T-bandage. **B., Hueter's**, a spica bandage for the perineum. **B., Langier's**, a many-tailed paper bandage. **B., Larrey's**, a form of many-tailed bandage, the edges being glued together. **B., Maisonneuve's**, a variety of plaster-of-Paris bandage, formed from cloths folded, these being supported by other bandages. **B., Recurrent**, is used after amputations, to support the flaps of the stump. **B., Richet's**, a form of plaster-

of-Paris bandage to which a small amount of gelatin has been added. **B. of Scultetus**, a compound bandage, similar to a spiral reverse in appearance and action, used in compound fractures, so that the short pieces of which it is composed may be removed without motion of the limb. **T.-Bandages** bandages shaped like the letter T. **B., The-den's**, a form of roller bandage applied from below upward over a graduated compress, to control hemorrhage from a limb. (For a more complete description of bandages, with illustrations, see *Gould's Illustrated Dictionary*.)

Baneberry (*bān'-ber-e*). See *Actea*.

Bang, Bhang, or Bangue (*bang*). See *Cannabis indica*.

Banian (*bau'-yan*) [*Ar.*, *banyan*, a trader]. A South Asiatic fig-tree, *Ficus bengalensis*, remarkable for the development of roots and secondary trunks from its branches. The bark and seeds are tonic, diuretic, and antipyretic. Unof.

Bantingism (*bant'-ting-izm*) [from *Banting*, the name of the inventor]. A method for the treatment of obesity, consisting in the abstinence from saccharine and farinaceous foods.

Banyan (*bau'-yan*). See *Banian*.

Baobab Tree. See *Adansonia digitata*.

Baptin (*bap'-tin*) [*βάπτισμα*, a dipping]. A purgative glucosid, obtainable from *Baptisia tinctoria*.

Baptisia (*bap-tiz'-e-ah*) [*βάπτισμα*, a dipping]. Wild Indigo. The root bark of *B. tinctoria* the properties of which are due to an impure glucosid, the so-called *Baptisin*. It is laxative and stimulant in moderate doses; emetic and cathartic in large doses, and is used in amenorrhea, typhus and typhoid fevers, and as a local application to indolent ulcers and gangrenous sores. **B., Extract**. Dose gr. j-x (0.065-0.65). **B., Ext. Fld.** Dose ℥ij-xx (0.13-1.3). **B., Tinct.** Dose ℥v-xxx (0.32-2.0). Dose of the resin gr. j-v (0.065-0.32). All unof.

Baptisin (*bap'-tiz-in*) [*βάπτισμα*, a dipping]. 1. A precipitate from the tincture of *Baptisia tinctoria*; antiseptic, purgative, ebolic, solvent. Dose gr. j-ij (0.065-0.2). Unof. 2. A bitter glucosid obtainable from the plant, *Baptisia tinctoria*; it has little medicinal activity.

Baraquet (*bar-ah-kwet'*) [*Fr.*]. Synonym of *Influenza*.

Barba (*bar'-bah*) [*L.*]. The beard.

Barbadoes Leg (*bar-ba'-dōs leg*). See *Elephantiasis Arabum*. **B. Aloes**. See *Aloes*. **B. Nut**. See *Purging Nut*.

Barbaloin (*bar-bal'-o-in*) [*Barbadoes*; *ἀλβή, aloē*], $C_{13}H_{20}O_7$. The aloin derived from Barbadoes Aloes.

Barber's Itch (*bar'-bers ich*). *Tinea barbae*.

Barberry (*bar'-ber-e*). See *Berberis*. **B. Gum**, a commercial name of certain varieties of gum-arabic.

Bardana (*bar-da'-nah*) [*L.*]. The burdock. See *Lappa*.

Baregin (*bar-a'-zhin*). See *Glaizin*.

Baresthesiometer (*bar-es-the-ze-om'-et-er*) [*βάρος*, weight; *αἴσθησις*, perception; *μέτρον*, a measure]. An instrument for estimating the sense of weight or pressure.

Baric (*ba'-rik*) [*βάρος*, weight]. Pertaining to or containing barium.

Barilla (*bar-il'-ah*) [*Fr.*, *barille*, impure soda]. The ashes obtained by burning various chenopodiaceous plants of the genera *Salicornia* and *Salsola*. They contain about 30 per cent. of sodium carbonate.

Barium (*ba'-re-um*) [*βάρος*, weight]. $Ba = 136.8$; quantivalence 11. A metal of the group of alkaline earths, of pale, yellow color, characterized by a strong affinity for oxygen. The salts are poisonous. **B. Carbonate**, $BaCO_3$, used in the preparation of the chlorid. **B. Chlorid**, $BaCl_2 \cdot 2H_2O$, soluble, used as a test for sulphates, which it precipitates as barium sulphate, and also as a cardiac and vasomotor stimulant. Dose gr. ss-v (0.032-0.32). **B. Hydrate**, $Ba(OH)_2$, caustic baryta; a crystalline substance, soluble in water, and used as a test for sulphates, which it precipitates as barium sulphate. **B. Iodid**, formerly used as an alternative. Dose gr. $\frac{1}{4}$ (0.008). It is employed in ointments. Unof. **B. Oxid**, BaO , baryta. **B. Sulphate**, $BaSO_4$.

Bark (*bark*) [*ME.*, *barkē*]. 1. The cortex or covering surrounding the wood of exogenous trees. 2. A synonym of cinchona or calisaya bark. **B., Peruvian**. See *Cinchona*. (For other barks, see the names of the plants or trees that produce them.)

Barley (*bar'-le*) [*ME.*, *barly*]. A cereal belonging to the order *Gramineæ*; the most common variety, *Hordeum distichon*, is used as a food, and also in the preparation of malt. **B. Decoct.** (*B. P.*). Dose $\frac{1}{3}$ j-iv (32.0-128.0). **B., Pearl**, is the decorticated grain, rounded and polished. **B. Water**, *decoctum hordei*, a decoction consisting of two ounces of Pearl B. boiled in one and one-half pints of water and afterward strained. It is used as a demulcent and food in the diarrheas of children.

Barnes's Curve [*Barnes*, an English obstetrician]. In obstetrics, the segment of a circle, having for its center the sacral promontory, its concavity looking backward. **B.'s Dilators**, graduated rubber bags used for dilating the cervix uteri in the induction of abortion or premature labor.

Barograph (*bar'-o-graf*) [*βάρος*, weight; *γράφειν*, to record]. A self-registering barometer.

Barometer (*bar-om'et-er*) [*βάρος*, weight; *μέτρον*, a measure]. An instrument for determining the weight and tension of the atmosphere. It consists essentially of a glass tube about 36 inches long, closed at one end, filled with mercury, and inverted in a basin of mercury. The mercury will sink in the tube until it rests at a height of about 30 inches (760 mm.) at the sea-level, the height varying as the atmospheric pressure increases or diminishes. **B.**, **Aneroid**, a metallic box from which the air has been exhausted, the tension being indicated by the collapsing or bulging of the thin corrugated cover, which is connected with a movable index. **B.-maker's Disease**, a form of chronic mercurial poisoning among the workmen who make barometers. It is due to the inhalation of the fumes of mercury.

Barometrograph (*bar-o-met'-ro-graf*). See *Barograph*.

Baroscope (*bar'-o-sköp*) [*βάρος*, weight; *σκοπεῖν*, to observe]. An instrument used for determining the loss of weight of a body in air, compared with its weight in a vacuum. A form of baroscope has been invented by Esbach for the quantitative determination of urea.

Barosma (*bar-oz'-mah*) [*βάρος*, weight; *βάρη*, smell]. A plant of the order *Rutaceæ*, native to the Cape of Good Hope and vicinity, several species of which yield the *Buchu* of commerce.

Barosmin (*bar-oz'-min*) [*βάρος*, weight; *βάρη*, smell]. A precipitate from the tincture of *Barosma crenata*. Dose grs. ij-iii (0.13-0.2). Unof.

Bartholin, Duct of (*bar'-tho-lin*). An excretory duct of the sublingual gland emptying into Wharton's duct. **B.**, **Glands of**, the vulvovaginal glands, a pair of glands situated at the entrance of the vagina, one on each side, and corresponding to Cowper's glands in the male.

Barton's Fracture. See *Diseases, Table of*.

Baruria (*bar-ur'-re-ah*) [*βάρος*, weight; *οὖρον*, urine]. The passage of urine having a high specific gravity; also the condition of the body associated therewith.

Baryencephalia (*bar-e-en-sef-ah'-le-ah*) [*βαρύς*, heavy; *ἐγκέφαλος*, brain]. Dulness of intellect.

Baryglossia (*bar-e-glos'-e-ah*) [*βαρύς*, heavy; *γλῶσσα*, a tongue]. Thick, slow utterance.

Barylalia (*bar-e-lah'-le-ah*) [*βαρύς*, heavy; *λαλῶ*, speech]. Thickness of speech.

Baryta, or **Barytes** (*bar-ih'-tah*, or *bar-ih'-tes*) [*βαρύς*, heavy]. Oxid of barium. See *Barium*.

Barythymia (*bar-e-thi'-me-ah*) [*βαρύς*, heavy; *θυμός*, mind]. A melancholy, gloomy, or sullen state of mind.

Basal (*ba'-sal*) [*βάσις*, a foundation]. Pertaining to or located at the base. **B. Ganglia**, the optic thalamus and corpus striatum of the brain.

Bascule Movement (*bas'-kül moov'-ment*) [*Fr.*, *bascule*, a swing]. The recoil of the heart in its systolic motion.

Base (*bäs*) [*βάσις*, a foundation]. The lowest part of a body or the foundation upon which anything rests. In chemistry, an element or radicle that combines with an acid to form a salt. The electropositive molecule or radicle of a compound. In pharmacy, the most important part of a prescription.

Basedow's Disease. Exophthalmic Goiter, or Graves' Disease. See *Goiter* and *Diseases, Table of*.

Basement (*bäs'-ment*) [*Fr.*, *bassement*]. The fundamental portion. **B. Membrane**. See *Membrane*.

Bas-fond (*bah fon(f)*) [*Fr.*]. The floor or lowest portion of the urinary bladder.

Basham's Mixture. Liquor ferri et ammonii acetatis.

Basial (*ba'-er-ah*) [*βάσις*, a foundation]. Relating to a base, or to the basion.

Basic (*ba'-sik*) [*βάσις*, a foundation]. Having properties of a base; *i. e.*, capable of neutralizing acids.

Basicity (*bas-is'-ih-e*) [*βάσις*, foundation]. 1. The quality of being basic. 2. The combining power of an acid.

Basiscranial (*bas-e-kra'-ne-ah*) [*βάσις*, foundation; *κράνιον*, the skull]. Relating to the base of the skull. **B. Axis**, a line running from a point midway between the occipital condyles through the median plane to the junction of the ethmoid and presphenoid.

Basifacial (*bas-e-fah'-shah*) [*βάσις*, foundation; *facies*, face]. Pertaining to the lower portion of the face. **B. Axis**, in craniometry, a line from the anterior point of the premaxilla to the anterior point of the basiscranial axis.

Basihyal (*bas-e-hi'-ah*) [*βάσις*, foundation; *ὑαλῆς*, from *υ*, *upsilon*; *ῥῖδος*, resemblance]. Either one of the two bones, one on each side, that form the principal part of the body of the hyoid arch.

Basihyoid (*bas-e-hi'-viah*). See *Basihyal*.

Basilar (*bas'-il-ar*) [*βάσις*, foundation]. Pertaining to the base. **B. Artery**. See *Arteries, Table of*. **B. Membrane**, a membranous division-wall separating the scala vestibuli from the scala tympani, extending from the base to the apex of the cochlea, and supporting the organ of Corti. **B. Process**, a strong quadrilateral plate of bone in front of the foramen magnum. **B. Suture**, the suture formed by the junction of the basilar process of the occipital bone with the posterior surface of the body of the sphenoid.

Basilateral (*bas-e-lat'-er-al*) [*βάσις*, foundation; *latus*, side]. Both basilar and lateral.

Basilic (*bas-il'-ik*) [*βασιλικός*, royal]. Illustrous; eminent. **B. Vein**, a large vein of the arm on the inner side of the biceps.

Basilicon Ointment (*bas-il'-ik-on oint'-ment*). Ceratum resinæ. It consists of resin 35 grams, yellow wax 15 grams, lard 50 grams.

Basilysis (*bas-il'-is-is*) [*βάσις*, foundation; *λίσις*, a loosening]. The breaking up of the fetal skull in craniotomy.

Basilyst (*bas'-il-ist*) [*βάσις*, foundation; *λίσις*, a loosening]. An instrument for use in perforating the cranial vault and breaking up the base of the skull.

Basin (*ba'-sin*) [M.E., *basin*]. 1. The third ventricle of the brain. 2. The pelvis. **B.-Trap**, a trap or seat in the outlet of the basin of a water-closet, placed there to prevent the escape into the apartment of noxious and offensive vapors and gases.

Basinasal (*bas-e-na'-sal*) [*βάσις*, foundation; *nasus*, the nose]. Relating to the basion and the nasion.

Basioccipital (*bas-e-ok-sip'-it-al*) [*βάσις*, foundation; *occiput*, the back of the head]. A bone, separate in many of the lower vertebrate animals, forming the central axis of the skull. In adult human life, it is the basilar process of the occipital bone.

Basioglossus (*bas-e-o-glos'-us*) [*βάσις*, foundation; *γλῶσσα*, the tongue]. That part of the hyoglossus muscle that is attached to the base of the hyoid bone.

Basion (*ba'-se-on*) [*βάσις*, foundation]. A point located at the middle of the anterior margin of the foramen magnum.

Basiotribe (*bas'-e-o-trib*) [*βάσις*, foundation; *τριβεῖν*, to grind or crush]. An instrument used for perforating or crushing the fetal head.

Basiotripsy (*bas-e-ot'-rip-se*) [*βάσις*, foundation; *τριβεῖν*, to grind or crush]. The operation of crushing the fetal head.

Basirrhinal (*bas-e-rin'-al*) [*βάσις*, foundation; *ῥίς*, nose]. Relating to the base of the brain and to the nose. Applied to a cerebral fissure located at the base of the olfactory lobe.

Basis (*ba'-sis*) [*βάσις*, foundation]. Base.

Basisphenoid (*bas-e-sfe'-noid*) [*βάσις*, foundation; *σφήν*, wedge; *ειδός*, form]. The lower part of the sphenoid bone.

Basisylvian (*bas-e-sil'-ve-an*) [*βάσις*, foundation; *Sylvius*, an anatomist]. Applied to the transverse basilar portion or stem of the Sylvian fissure.

Basitemporal (*bas-e-tem'-po-ral*) [*βάσις*, foundation; *tempora*, the temples]. Relating to the base or lower part of the temporal bone.

Basivertebral (*bas-e-ver'-te-bral*) [*βάσις*, foundation; *vertebra*, a joint]. Relating to the basis or centrum of a vertebra.

Basophile (*bas'-e-phil*). See *Basophilus*.

Basophilic (*bas-o-phil'-ic*) [*βάσις*, foundation; *φιλεῖν*, to love]. Combining readily with bases; stainable by means of basic dyes.

Basophilous (*bas-off'-il-us*) [*βάσις*, foundation; *φιλεῖν*, to love]. Stained by basic rather than by acid dyes.

Bass-deafness (*bās-def'-nes*) [Fr., *basse*, fem. of *bas*, low; M.E., *deaf*]. Deafness to certain bass-notes, the perception of the higher notes being retained.

Bassorin (*bas'-o-rin*) [*Bassora*, an Asiatic town] $C_6H_{10}O_5$, or $2C_6H_{10}O_5$. A tasteless, odorless, vegetable mucilage, insoluble in cold water, but rendered soluble by alkalis; it is found in gum-tragacanth (of Bassora) and in cherry and plum gums.

Bast (*bast*) [AS., *bast*, a lime-tree]. The inner bark of exogenous plants. The fibrous parts of the bark are used in making cordage, and have a limited use in surgery.

Bastard (*bast'-lard*) [OF., *bastard*, son of a bast, or pack-saddle]. 1. An illegitimate child. 2. Illegitimate.

Bastards (*bast'-lards*) [see *bastard*]. The name given to an impure sugar procured by concentrating molasses and allowing it to crystallize slowly in molds.

Basyl (*ba'-sil*) [*βάσις*, foundation]. The electropositive constituent of a compound.

Bateman's Drops. The *tinctura pectoralis*, N. F.; a weak tincture of opium, camphor, and catechu; a popular remedy in coughs.

Bath (*bath*) [AS., *bath*, a bath]. 1. A bathing-place or room. 2. The medium in which the body is wholly or partly immersed. As therapeutic agents, baths are classified according as water, vapor, air, etc., is used; according to the temperature, as hot, temperate, cold, etc.; according to the end desired, as nutritional, medicinal, stimulant, etc. Special forms of bath are the moor, peat, mud-slime, pine-leaf, herb (hay, gentian, camomile, juniper, marjoram, etc.), brine, sand, tan, bran, malt, glue, soup, milk, whey, blood, wine, guano, starch, soap, acid, iron, sulphur, carbonic acid, compressed air, mustard, electric, etc. **B., Acid**, add 1½ ounces of nitric acid and 1 to 3 ounces of hydrochloric acid to 30 gallons of warm water in a wooden or earthenware vessel, and immerse the patient in this for from 10 to 20 minutes. For a foot-bath, add ½ ounce of nitric acid and 1 ounce of hydrochloric acid to 4 gallons of warm water. This is said to be useful in cases of dyspepsia, with sluggish liver and constipation. **B., Air**, a bath in which but little water is employed, the body being exposed freely to the air. It is employed in those cases in which there is a tendency to catch cold on slight exposure. **B., Alcohol**, one in dilute alcohol, used to

reduce temperature in fever. **B., Alkaline**, add 3 ounces of potassium carbonate, or 6 of sodium carbonate, to 25 or 30 gallons of hot water. It is used in chronic squamous skin-diseases, chronic rheumatism, and lithemia. It should be taken in a wooden, earthenware, or enameled tub. **B., Bog**, a bath made by mixing bog-earth (produced by the decomposition of plants in the presence of water and found at iron and sulphur springs) with warm water to form a pulpy or mushy consistence. This is used as a mud-bath. **B., Borax**, borax, 4 ounces; glycerol, 3 ounces; water, 30 gallons. It is used in the same class of cases as the bran-bath. **B., Bran**, boil 1 pound of bran in 1 gallon of water, strain, and add 30 gallons of water. This is a soothing and emollient bath, and is of service in squamous and irritable conditions of the skin. **B., Chemic**, in chemistry, an apparatus for regulating the temperature of chemic processes by surrounding the substance with water, sand, oil, or mercury, through which the heat is communicated. **B., Cold**, a bath of cold water, the temperature of the latter varying from 32°-70° F. (0°-21° C.). It is used to reduce fever and as a general stimulant. **B., Foot**, a bath for the feet, used as a derivative agent in cases of cold, etc. **B., Full**, one in which the patient's body is entirely covered by water, so that his chin just clears it. **B., Graduated**, one in which the temperature of the water is gradually lowered by the addition of cold or ice-water. **B., Herb**; this is made by using the extract of pine-needles, or of some aromatic herbs, and is used as a tonic. **B., Hip**. See *B., Sitz*. **B., Hot**, one in which the temperature of the water ranges from 104° to 110° F. (40°-43.3° C.). It acts upon the skin, producing free perspiration, and accelerates the pulse and respiration. **B., Hot-air**; Turkish bath. A bath in hot air. It is useful as a diaphoretic, and in catarrhal, neuralgic, and rheumatic conditions, but is contraindicated in fatty degeneration of the heart. **B., Medicated**, a bath in which medicinal substances, as mineral salt, sulphur, etc., are dissolved or held in suspension. **B., Mercurial**, for the treatment of syphilis. A bath in the vapor of mercury, usually prepared by vaporizing calomel over an alcohol lamp. **B., Milk**, a bath in milk, for nutritive purposes. **B., Moor**, a bath in water mixed with the earth of moors. **B., Mud**, a bath prepared by mixing well-seasoned earths, containing more or less mineral matter, with water containing the same substances. It is useful in chronic rheumatism. **B., Mustard**, made by enclosing from two to four ounces of ordinary mustard in a piece

of muslin or thin linen and hanging it in about four gallons of hot water until the latter becomes yellow, or simply by adding mustard to water. It is used as a general bath for infants in collapse, convulsions, or severe bronchitis, the child being left in until the skin becomes distinctly reddened. It is also used as the foot or sitz-bath in amenorrhea. **B., Pack**, or **Sheet**, one in which the body is wrapped in wet cloths. **B., Pine**, prepared by adding a decoction of pine-needles, or some pine-extract, to hot water. It is mildly stimulating, and is employed in hysteria, gout, and rheumatism. **B., Rain**, consists of from four to six three-fourths circles of pipes secured together at a distance of two to three inches. Each pipe has three lines of fine perforations, from which the stream issues under pressure, striking the body of the patient at all points with considerable force. **B., Russian**, a vapor-bath, the vapor being generated by throwing water upon heated mineral or metallic substances. *Syn.* *B., Vapor*. **B., Sand**, the body of the patient is placed in a layer of dry sand that has been heated. In chemistry, the immersion of a crucible, etc., containing a chemic compound, in a vessel containing fine sand, the latter being heated gradually to a high temperature. **B., Sheet**, the application of cold water to the body through the medium of a wet sheet or towel of fine or coarse texture, placed upon the skin. It is used to reduce temperature. **B., Shower**, a douche in which the water is delivered against the body from a nozzle with numerous perforations. **B., Sitz**, one in which only the buttocks and hips are immersed in water. It is useful in pelvic inflammations, amenorrhea, and retention of urine. **B., Slime**, a bath in water mixed with the slimy deposit of organic matter found in rivers or ponds. **B., Sponge**, one in which the patient's body is rubbed with a wet sponge. It is used to reduce temperature. **B., Sulphur**, potassium sulphid 4 to 8 ounces in 30 gallons of water; a little sulphuric acid may be added. It is used in certain skin diseases, scabies, lead colic, and lead palsy. **B., Sun**, the exposure of the naked body to the sun's rays. **B., Tepid**, the temperature of the water ranges from 85° to 95° F. (29.4°-35° C.). It acts as a sedative, cleansing, and detergent agent; the pulse, respiration, excretion, secretion, and temperature are practically unaffected. **B., Turkish**, one in which the bather is placed successively in rooms of higher temperature, then shampooed or rubbed, and finally stimulated by a douche of cold water. See *Bath*, *Hot-air*. **B., Vapor**. See *B., Russian*. **B., Warm**, a bath in water having a tem-

perature of from 90° to 104° F. (32.2°–40° C.). It is used to calm the nervous system, produce sleep, and allay reflex irritability. **B.**, Water, a bath of water. In chemistry, a bath of water for immersing vessels containing substances that must not be heated above the boiling-point of water.

Batophobia (*bat-o-fə-be-ah*) [*βαρός*, a height; *φόβος*, fear]. 1. Acrophobia; dread of being at a great height. 2. Dread of high objects; fear of passing near a high building, or of going through a deep valley.

Battarism (*bat'-ar-izm*) [*βαρρατισμός*, a stuttering]. Stuttering or stammering.

Battery (*bat'-er-e*) [*batteria*, a beating; bat-

tery]. A series of two or more pieces of apparatus arranged to produce increased effect, as battery of boilers, prisms, lenses, galvanic cells. It is usually applied to a series of cells producing electricity (voltaic or galvanic battery); also, frequently to a single cell. **B.**, Faradic, one giving a faradic current. **B.**, Galvanic, one giving a galvanic or chemic current. **B.**, Primary, the combination of a number of primary cells so as to form a single battery. **B.**, Secondary, the combination of a number of storage-cells to form a single electric source. The following table shows the materials used in the more important batteries, and the electromotive force (E. M. F.) in volts.

NAMES, ELEMENTS, FLUIDS, ETC., OF THE PRINCIPAL BATTERIES.

NAME OF CELL.	POSITIVE ELEMENT.	NEGATIVE ELEMENT.	EXCITING AGENT.	DEPOLARIZING AGENT.	E. M. F. IN VOLTS.
Bunsen.	Zinc.	Carbon.	Sulphuric acid, dilute.	Nitric acid.	1.75 to 1.96
Beetz.	Zinc.	Copper.	Zinc sulphate in plaster-of-Paris.	Copper sulphate, in plaster-of-Paris.	1.04
Callaud.	Zinc.	Copper.	Solution of zinc sulphate.	Cupric sulphate.	1.0 to 1.14
Daniell.	Zinc.	Copper.	Zinc sulphate.	Cupric sulphate.	1. to 1.14
Edison-La-lande.	Zinc.	Cupric oxid and carbon.	Potassium hydroxid.	Cupric oxid.	0.75
Fuller.	Zinc.	Carbon.	Zinc chlorid.	Potassium dichromate and hydrochloric acid.	1.5
Gravity. See	<i>Callaud.</i>				
Gaiffe.	Zinc.	Silver.	Zinc chlorid.	Silver chlorid.	1.02
Grenet.	Zinc.	Carbon.	Sulphuric acid and potassium dichromate.	Chromic acid.	2.0
Grove.	Zinc.	Platinum.	Sulphuric acid, dilute.	Nitric acid.	1.04 to 1.97
Latimer Clark.	Zinc.	Mercury.	Mercuric sulphate.	Mercuric sulphate.	1.436
Leclanché.	Zinc.	Carbon.	Ammonium chlorid.	Manganese dioxid.	1.42
Lelande Chap-erone.	Zinc.	Copper or iron.	Potassium hydroxid.	Cupric oxid.	0.98
Maische.	Zinc scraps in a bath of mercury.	Platinized carbon.	Common salt solution.	None separate.	1.25
Marié-Davy.	Zinc.	Carbon.	Sulphuric acid, dilute.	Mercuric sulphate.	1.52
Maynooth.	Zinc.	Iron.	Sulphuric acid, dilute.	Nitric acid (strong).	
Niaudet.	Zinc.	Carbon.	Common salt solution.	Chlorated lime.	1.63
Poggendorf.	Zinc.	Carbon.	Potassium dichromate and sulphuric acid.	Chromic acid.	1.98
Schanschieff.	Zinc.	Carbon.	Mercurial solution.	None separate.	1.56
Scrivanoff.	Zinc.	Silver.	Potassium hydroxid.	Silver chlorid.	1.5
Smee.	Zinc.	Platinized silver.	Sulphuric acid, dilute.	None.	0.5 to 1.0
Walker.	Zinc.	Platinized carbon.	Sulphuric acid, dilute.	None.	0.66
Warren de la Rue.	Zinc.	Silver.	Ammonium chlorid.	Silver chlorid.	1.05

Battledore (*bat'-l-dor*) [ME., *batyldoure*, a bat for beating clothes]. An instrument shaped like a racket. **B. Placenta**, one in which there is a marginal insertion of the cord.

Baudelocque's Diameter (*bo'-del-oks*). In obstetrics, the external conjugate diameter of the pelvis.

Bauhin, Valve of (*bo'-hin*). The ileocecal valve.

Baunscheidtism (*bozon'-shit-izm*) [named from *Baunscheidt*, the inventor]. A mode of treating rheumatism and chronic neuralgias by counterirritation, the latter being produced by pricking the exterior of the part affected with fine needles dipped in oil of mustard, formic acid, or other irritant.

Bavarian (*bah'-va'-re-an*) [*Bavaria*]. Pertaining to Bavaria. **B. Dressing**. See *B. Splint*. **B. Splint**, a variety of immovable dressing in which the plaster is applied between two flannel cloths.

Bayberry (*ba'-ber-i*) [ME., *bay*; *bery*]. 1. The berry of the *Laurus nobilis*, bay, or noble laurel. 2. The wax-myrtle, *Myrica cerifera*, and its fruit. See *Myrica*. 3. The pimento, or allspice. **B. Rum** and **B., Oil of**. See *Myrica*. **B. Tree**, the *Laurus nobilis*; also *Prunus lauraceranus*; commonly called the laurel and the cherry-laurel.

Bazin (*ba'-zin*). Synonym of *Molluscum contagiosum*.

Bazin's Disease. See *Diseases, Table of*.

Bdellium (*del'-e-um*) [Heb., *b'dolach*]. A resinous gum exuding from various species of *Balaamodendron*. It resembles myrrh. **B., Indian**, has been recommended as an emmenagogue.

Beads, Rhachitic. "Rhachitic rosary;" the so-called "beading of the ribs" in rickets; a succession of visible and palpable swellings at the points where the ribs join their cartilages.

Beak (*bek*) [ME., *becke*]. 1. The mandibular portion of a forceps. 2. The lower end of the calamus scriptorius. 3. The pad or splenium of the corpus callosum.

Beaker (*be'-ker*) [ME., *biker*]. A wide-mouthed glass vessel used in chemist laboratories. **B.-cell**, the goblet-cell found in mucous membranes.

Bean (*ben*) [ME., *bené*]. The seeds of several plants, mostly *Leguminosa*, especially that of the common bean, *Faba vulgaris*. **B. of St. Ignatius**. See *Ignatia*.

Bearberry. See *Uva ursi*.

Bear's-foot. Leaf-cup. A composite-flowered plant, *Polymnia uvedalia* of N. America. A popular remedy for enlargement of the spleen, or the "ague-cake" of malarious regions.

Beat (*bet*) [ME., *beten*]. The pulsation of the arteries, or the impulse of the heart.

Beauperthuy's Method (*be'-per-thues*). A method of treating leprosy by bathing with olive oil, the internal administration of mercuric chlorid, abstinence from salted meats, and good hygiene.

Beberin (*be'-ber-in*). See *Nectandra*.

Bechic (*bek'-ik*) [*βήξ*, a cough]. Relieving cough; a remedy against cough.

Bechterew's Nucleus (*bek'-ter-är*). The accessory auditory nucleus.

Béclard's Hernia. See *Diseases, Table of*.

Becquerel's Pills (*bek'-a-rels*). Pills containing quinin, extract of digitalis, and colchicum seeds, for use in gout.

Bed (*bed*) [AS., *bedd*]. The couch or support on which the body may rest in sleep; usually a mattress of straw, hair, or similar substance. **B., Air**, a mattress of rubber or leather that can be inflated with air. **B.-bug**, an apterous insect, *Cimex lectularius*, that infests filthy bedsteads, and at times is parasitic upon the human body. **B.-pan**, a large shallow vessel for receiving the alvine discharges of bedridden patients. **B.-rest**, an apparatus for propping up patients in bed. **B.-ridden**, confined to bed. **B.-sore**, a sore produced on any part of the body by prolonged pressure against the bed, or by trophic changes in paralyzed parts.

B., Water, a rubber mattress containing water; it is used to prevent the formation of bed-sores.

Bedlam (*bed'-lam*) [ME., *bedlem*, a corruption of *Bethlehem*]. An insane asylum.

Bedlamism (*bed'-lam-izm*) [see *Bedlam*]. Insanity.

Bedlamite (*bed'-lam-it*) [see *Bedlam*]. A madman; an insane person.

Bednar's Aphthæ. See *Aphthæ*, and *Diseases, Table of*.

Beef (*bef*) [ME., *beſe*]. The flesh of domestic cattle. Good beef should be of red color, possess firm texture, and be free from unpleasant smell. Beef consists of water 73, fibrin 15, gelatin 4, albumin 3, fat and other substances 5 per cent. **B.-extract**, the soluble fibrin of lean meat partly desiccated. **B.-measles**. See *Cysticercus bovis*. **B.-tea**, the soluble extractive matter of beef, made by steeping finely-cut lean beef with its weight of water, and straining.

Beer (*ber*). See *Malt-liquors*.

Beer's Cataract Knife. A knife with a triangular-shaped blade, for making section of the cornea in the removal of the crystalline lens.

Beeswax (*bee'-waks*). *Cera flava*; wax secreted by bees, of which their cells are constructed. It is used in making candles, ointments, and pomades.

Beet (*bet*). See *Beta*.

Beggiatoa (*bej'-e-at-o'-ah*) [after the Italian

botanist, *J. Beggiato*]. A genus of bacteria founded by Trevisan, consisting of swinging or gliding, milk-white, gray, rosy, or violet threads. They decompose sulphur compounds, and store up sulphur granules in their protoplasm. They are found in stagnant, fresh, or salt water, particularly in that contaminated with sewage or factory-waste. See *Bacteria*, Table of.

Behen-nut (*be'-hen nut*). The seed of *Moringa pterygosperma*, and *M. aptera*, trees of tropical countries. They afford Oil of Behen, or Oil of Ben.

Bela (*be'-lah*) [Hind.]. The dried, half-ripe fruit of *Aegle marmelos*, or Bengal quince. It is recommended as a remedy for chronic diarrhea and dysentery. The ripe fruit is slightly laxative. Dose 3 ss-j (2.0-4.0). **B.**, Ext., Liq. (B.P.). Dose f3j-ij (4.0-8.0). Unof.

Belching (*belch'-ing*) [ME., *belchen*]. The expulsion of gas from the stomach; eructation. **Belemnoid** (*bel-em'-noid*) [*Βέλεμνον*, a dart; *ειδος*, form]. 1. Dart-shaped; styloid. 2. The styloid process of the ulna or of the temporal bone.

Bell (*bel*) [ME., *bel*]. A hollow, metallic, sounding instrument. **B.** Gastrula. See *Archigastula*. **B.** Metal, an alloy of copper, zinc, tin, and antimony. **B.**, Nerves of, the external and internal respiratory nerves, *i. e.*, the posterior thoracic and phrenic nerves. **B.** Sound, **B.** Tympany, the sound produced in pneumothorax by striking a coin, placed flat upon the chest, with another coin. It can be heard through the stethoscope placed over the affected side.

Belladonna (*bel-ah-don'-ah*) [It., *belladonna*, beautiful lady]. Deadly night-shade. A perennial plant of the order *Solanaceæ*, indigenous to Southern Europe and Asia, and cultivated in the U. S. Its properties are due to two alkaloids, *atropin* and *belladonnin*, the latter thought to be identical with hyoscyamin. Both leaves and root are employed. It is used as an antispasmodic, as a cardiac and respiratory stimulant, and to check secretion, as that of the sweat and milk. **B.**, Folia, the leaves of belladonna. **B.**, Radix, the root of belladonna. The preparations of the leaves are: **B.**, Extractum, Alcoholicum. Dose gr. $\frac{1}{8}$ - $\frac{1}{4}$ (0.0065-0.032). **B.**, Tinct., 15 per cent. Dose m-j-xxx (0.065-2.0). **B.**, Unguent., contains extract 10, dilute alcohol 6, benzoated lard 84 parts. From the root are prepared: **B.**, Abstrac-tum. Dose gr. $\frac{1}{8}$ -j (0.0065-0.065). **B.**, Emplastrum, made with resin-plaster. **B.**, Extractum, Fluidum. Dose m-j-v (0.065-0.32). **B.**, Linimentum, fluid extract 95, camphor 5. **B.**, Succus (B. P.). Dose

m-v-xv (0.32-1.0). **Atropin Sulphate**, the most frequently used preparation. Dose gr. $\frac{1}{160}$ - $\frac{1}{80}$ (0.00036-0.0008). **Lamellæ Atropinæ** (B. P.), each containing of atropin $\frac{1}{37500}$ grain (0.000013). Liq. **Atropinæ Sulphatis** (B. P.). Dose m-j-vj (0.065-0.40).

Belladonnin (*bel-ah-don'-in*) [see *Belladonna*], $C_{17}H_{23}NO_3$. An alkaloid found in *Belladonna*. It resembles atropin, hyoscyamin, and hyoscin. It occurs with atropin, and is likewise decomposed into tropic acid and oxytropin, $C_8H_{15}NO_3$. See *Belladonna*. **Bellocc's Canula** (*bel'-oks*). An instrument used in plugging the nares.

Bell's Disease. See *Periencephalitis and Diseases*, Table of. **B.** Mania. See *Periencephalitis and Diseases*, Table of. **B.** Palsy or Paralysis. See *Diseases*, Table of.

Belly (*bel'-e*) [ME., *bely*]. See *Abdomen*.

B.-ache. See *Colic*. **B.-bound**, a vulgar term for constipated. **B.-button**, the navel.

Ben, or **Benne Oil** (*ben*, or *ben'-e*). *Oleum balatinum*, is obtained by expression from the seeds of the several species of *Moringia*. It is a colorless, odorless oil, not readily turning rancid. It is used for extracting odors, and for lubricating clocks and light machinery.

Benedikt, Syndrome of. Paralysis of the muscles supplied by the oculomotor nerve of one side and tremor and paresis of the upper extremity of the other side.

Bengalin (*ben'-gaul-in*) [Hind., *Bengāl*]. A blue pigment derived from the benzene colors.

Benign, or **Benignant** (*be-nin'*, or *be-nig'-nant*) [*benignus*, kind]. Not endangering health or life; applied to certain tumors.

Benzaldehyd (*ben-zal'-de-hid*) [*benzoin*; *aldehyd*], C_7H_6O . Bitter-almond oil; a compound that results from the oxidation of benzyl alcohol.

Benzanalgen, $C_9H_7OC_2H_5.NHCOC_6H_5.N$. A derivative of chinolin. It is antiseptic, antipyretic, and antineuralgic, and has the power of dissolving uric acid. It is used in rheumatism, tabes dorsalis, and chronic gout. Dose gr. viiss-xlv (0.5-3.0).

Benzanilid (*ben-zan'-il-id*) [*benzoin*; *anilin*], $C_{15}H_{11}NO$. Benzoyl anilid, prepared by the action of benzoyl chlorid on anilin. It is used as an antipyretic in children, in doses of gr. iij-viiij (0.20-0.50). Unof.

Benzene (*ben'-zen*) [*benzoin*], C_6H_6 . A hydrocarbon contained in coal-tar. It is formed by the dry distillation of all benzene acids. It is a mobile, ethereal-smelling liquid, of specific gravity 0.899 at 0°. It solidifies about 0°, melts at + 6°, boils at 80.5°, and burns with a luminous flame. It readily

dissolves resins, fats, sulphur, iodine, and phosphorus. Aniline and the aniline-colors are derived from it. **B.-sulphonic Acid**, $C_6H_5 \cdot SO_3H$, is prepared by boiling together equal parts of benzene and H_2SO_4 . It occurs in small plates, readily soluble in alcohol and water, and which deliquesce in the air.

Benzimid (*ben'-zim-id*) [*benzoin*; *amid*], $C_{13}H_{15}N_3O_2$. A compound formed by the action of hydrocyanic acid on hydride of benzoyl. It occurs also in the resinous residue of the rectification of the oil of bitter almonds.

Benzine (*ben'-sin*). See *Benzinum*.

Benzinum (*ben-si'-num*) [L.]. Benzene, Petroleum ether. A purified distillate from American petroleum, having a specific gravity of .77 to .79, boiling at 80° to 90° C., colorless, of ethereal odor, and a slightly peppermint-like taste. It is a valuable solvent for oils, fats, resins, caoutchouc, and some alkaloids. It has been used against tapeworm. It should be distinguished from *Benzol*, which is called *Benzene* by English chemists. Dose grt. v-x (0.33-0.66), on sugar or in mucilage.

Benzoate (*ben'-zo-at*) [*benzoin*]. Any salt of benzoic acid. **B. of Naphthol**. See *Benzonaphthol*.

Benzoated (*ben'-zo-a-ted*) [*benzoin*]. Impregnated with benzoin or with benzoic acid.

Benzoic (*ben'-oi-ik*) [*benzoin*]. Pertaining to or derived from benzoin. **B. Acid**. See *Acid*.

Benzoin (*ben'-zo-in* or *-oin*) [origin obscure]. A resin obtained from *Syrax benzoin*, a tree native to Sumatra and Siam. It is a ketone alcohol, $C_{14}H_{12}O_2$, and may be produced by oxidizing hydrobenzoin with concentrated HNO_3 . It is sparingly soluble in water, cold alcohol, and ether, and crystallizes in shining prisms, melting at 134° . It yields benzoic and cinnamic acids, is antiseptic and disinfectant, and is used mainly as a stimulant expectorant in chronic bronchitis. **Adeps benzoïnatus**, benzoïnated lard, contains 2 per cent. of benzoin. **B., Tinct.**, 20 per cent. of the resin in alcohol. Dose fʒ ss-j (2.0-4.0). **B., Tinct. Comp.**, Friar's Balsam, benzoin 12, aloes 2, storax 8, balsam of Tolu 4, alcohol, sufficient to make 100 parts. Dose fʒ ss-ij (2.0-8.0).

Benzol (*ben'-zol*) [*benzoin*], C_6H_6 . A hydrocarbon formed by the dry distillation of organic substances, but in commerce chiefly derived from coal-tar. It is inflammable, volatile, and a solvent for fats. It is employed externally as a parasiticide; internally, as a pulmonary antiseptic, in influenza, and in trichiniasis. Its vapor is used in whooping-cough. Dose grt. v-x (0.32-0.65).

Benzonaphthol (*ben'-zo-naf'-tol*) [*benzoin*; *vâpha*, a kind of asphalt], $C_{10}H_8O$ (C_6H_5O). The benzoate of beta-naphthol, used as an intestinal antiseptic in typhoid fever and other intestinal diseases. Dose gr. ij-vij (0.13-0.52).

Benzonitrile (*ben'-zo-ni'-tril*) [*benzoin*; *nitrum*, niter], C_7H_5N . A substance obtained from benzene-sulphonic acid by distillation with potassium cyanide. It is an oil with an odor resembling that of oil of bitter almonds, and boils at 191° ; its specific gravity is 1.023 at 0° .

Benzosol (*ben'-zo-sol*) [*benzoin*], $C_{15}H_{17}O_2$. Benzoyl-guaïacol, a substitute for creasote in tuberculous affections. It also has antipyretic properties. Dose gr. iv-x (0.20-0.65).

B.-eugenol, $C_{17}H_{16}O_3$, a derivative of eugenol, used in tuberculosis. **B.-guaïacol**, $C_{14}H_{12}O_3$, benzosol. It is the benzoate of guaïacol and contains 54 per cent. of guaïacol. Its chief uses are as an intestinal antiseptic, and in pulmonary tuberculosis. Dose gr. iij-xij (0.2-0.75).

Benzoyl (*ben'-zo-il*) [*benzoin*], C_7H_5O .

The radicle of benzoic acid, of oil of bitter almonds, and of an extensive series of compounds derived from this oil, or connected with it by certain relations. **B.-ecgonin**, $C_{16}H_{16}NO_4H$, a substance intermediate in composition between cocaine and ecgonin. **B.-glycocoll**. See *Acid, Hippuric*.

Benzyl (*ben'-zil*) [*benzoin*], C_7H_7 . A univalent hydrocarbon radicle that does not exist in the free state, but in combination forms a considerable number of compounds.

B. Alcohol. See *Alcohol*.

Benzylidene (*ben'-zil'-id-en*) [*benzoin*], C_8H_6 . A bivalent hydrocarbon radicle derived from benzoin compounds.

Bérard's Aneurysm. See *Aneurysm and Diseases, Table of*.

Béraud, Ligament of (*ba-roh'*). The ligament that suspends the pericardium. **B., Valve of**, a valve supposed to exist at the bottom of the lacrymal sac over the entrance to the nasal duct.

Berberin (*ber'-ber-in*) [*berberina*], $C_{20}H_{17}NO_4$. An alkaloid found in the bark of *Berberis*, and in many other plants. It is recommended as a tonic and antiperiodic, and is an ingredient of various lotions for mucous membranes; it is useful in nasal catarrh, etc. Dose gr. j-x (0.065-0.65). Unof.

Berberis (*ber'-ber-is*) [L.]. Barberry. The root of *Berberis vulgaris*, or common barberry. Its properties are due to an alkaloid, *berberin*, $C_{20}H_{17}NO_4$. It is an astringent, bitter tonic; in large doses a cathartic. It has been used locally in conjunctivitis, and internally in malarial and typhoid fevers. **B., Ext., Fld.** Dose m v-xxx (0.32-2.0). **B., Tinct.**, contains 20 per cent. of the root.

Dose η x-f ζ j (0.65-4.0). **Berberin**, the alkaloid. Dose gr. j-x (0.065-0.65). **B. Hydrochlorate**, is used locally in gonorrhea. All unof. *B. asiatica*, *B. aquifolium* (or Oregon grape), and *B. lycium* are all medicinally active, and have similar properties to those of the common barberry.

Bergamot, Oil of (*ber'-gam-of*) [*Bergamo*, a town in Italy], $C_{10}H_{16}$. A volatile oil derived from the rind of the *Citrus bergamia*. It is used mainly as a perfume, and as a clearing-agent in histologic work.

Bergeron's Disease. See *Diseases, Table of*.

Beriberi (*ber'-e-ber'-e*) [Cingalese, *beri*, weakness]. Synonyms: *Kakke*, *Panneuritis epidemica*. An infectious disease, endemic in various countries of Asia (Ceylon, India, China, Japan), Africa, and Australia, and presenting the features of a multiple neuritis. Bad food and defective hygienic conditions are predisposing causes; the true etiologic factor is probably a microorganism. Various types of the disease have been described, particularly an anasarctous form and an atrophic form. The main symptoms are anæsthesia, paralysis, wasting of the muscles, edema, changes in the electric reactions, dyspnea, and palpitation. The mortality is variable; often it is very great. True beriberi has been met with in the seaport towns of the United States, introduced there by East Indian sailors, and an affection resembling it seems to occur among certain fishermen of New England.

Bernard's Granular Layer. The inner zone of the cells lining the acini of the pancreas. It is granular in appearance and stains but slightly with carmin.

Berry (*ber'-e*) [ME., *bery*]. An indehiscent fruit with a pericarp that is succulent throughout, as the grape and gooseberry.

Bertillonage (*ber-til-lon-a(h)j*) [*Bertillon*, a French criminologist]. A system of carefully-recorded measurements and descriptions of criminals, for the purpose of future identification, introduced into France by Bertillon and adopted by the police of many large cities of the U. S.

Bertin, Bones of. The sphenoidal turbinated bones, partly closing the sphenoidal sinuses. **B., Column of**, a cortical column of the kidney; the part separating the medullary pyramids. **B., Ligament of**, the iliofemoral ligament.

Berthollet's Law. See *Law*.

Beryllium (*ber-il'-e-um*) [*βήρυλλος*, *beryl*]. A bivalent metal obtained from the beryl, whence its name; it is called also glucinum. See *Elements, Table of*.

Bestiality (*bes-te-al'-it-e*) [*bestia*, a beast]. Unnatural intercourse with an animal.

Bestucheff's Mixture or Tincture. The

etheral tincture of chlorid of iron, used in erysipelas. It is made as follows: \mathcal{R} . Tinct. ferri chloridi, 1 part; sp. æther. nitrosi, 4 parts. Mix and expose to the rays of the sun in well-closed bottles until the brownish color disappears. The dose is from one to two teaspoonfuls every three hours.

Beta (*ba'-tah*) [L.]. 1. The beet. 2. The second letter of the Greek alphabet, used in chemic nomenclature, to indicate the second of two isomeric compounds. **B.-naphtol**. See *Naphtol*.

Betacism (*ba'-tas-izm*) [*beta*, β τα, the second letter of the Greek alphabet]. The too-frequent use of the *b*-sound in speech, or the conversion of other sounds into it.

Betain (*be-ta'-in*) [*beta*, a beet], $C_5H_{11}NO_2$, or $C_{10}H_{22}N_2O_4$. Called also oxyneurin and lycin. A basic substance obtained by the careful oxidation of cholin. It occurs already formed in the sugar-beet (*Beta vulgaris*) and crystallizes from alcohol with one molecule of water in shining crystals that deliquesce in the air; it has an alkaline reaction and a sweetish taste.

Beta-naphtol Bismuth. A preparation containing 80 per cent. of bismuth oxid and 20 per cent. of beta-naphtol. It is a brown powder, insoluble in water and decomposed into its component parts in the intestine, the beta-naphtol being absorbed and discharged with the urine, while the bismuth is evacuated with the stools. It is an intestinal antiseptic. Dose gr. xv-xlv (1.0-3.0).

Betel (*be'-tel*) [E. Ind.]. A masticatory used in the East. A few grains of the nut of the Catechu palm, *Areca C.*, are rolled up with a small amount of quicklime in a leaf of *Piper betel*, and chewed. It is tonic, astringent, stimulant, and aphrodisiac, and seems to increase the powers of endurance. Dose of fld. ext. f ζ j-ijj (4.0-12.0). Unof.

Betol (*be'-tol*), $C_{10}H_7O.C_7H_5O_2$. Salinaphthol. Naphthalol. A salicylic ether of naphtol, used in rheumatism and cystitis. It resembles salicylic acid in its properties. Dose gr. x-xv (0.65-1.0).

Betula (*be'-u-lah*). See *Birch*.

Betulin (*be'-u-lin*) [*betula*, birch], $C_{30}H_{60}O_3$ (Hausmann). Birch-resin, or birch-camphor, derived from the bark of the white birch.

Bezoar (*be'-zo-ar*) [Pers., *pūd-zahr*, the bezoar-stone, a supposed antidote against poison]. A concretion found in the stomach or intestines of some animals (especially ruminants), formerly believed to be efficacious in preventing the fatal effects of poison.

Bhang (*bang*). See *Cannabis indica*.

Bhel (*bel*). See *Bela*.

Biauricular (*bi-au-rik'-u-lar*) [*bis*, twice; *auricula*, the ear]. Relating to two auricles, or to two corresponding auricular points.

Bibulous (*bi-b' - u - lus*) [*bibere*, to drink]. Having the property of absorbing moisture, as, *B. Paper*.

Bicarbonate (*bi-kar' - bon - at*) [*bis*, twice; *carbo*, a coal]. Any salt of carbonic acid that has two carbonic acid equivalents for one of the base.

Bicaudal, Bicaudate (*bi-kaw' - dal, bi-kaw' - dat*) [*bicaudalis*, two-tailed]. Having two tails or appendages.

Biceps (*bi' - seps*) [*bis*, twice; *caput*, the head]. Having two heads, a term applied to several muscles, as *B. brachii*, *B. extensor*, *B. flexor cruris*.

Bichat (*be' - shak*) Canal of. See Canal. **B.**, Fissure of. See Fissure. **B.**, Foramen of. See Foramina, Table of. **B.**, Tunic of, the intima of the blood-vessels.

Bichlorid (*bi-klo' - rid*) [*bi*, two; *chlorin*]. A salt containing two equivalents of chlorin.

Bichromate (*bi-kro' - mat*) [*bi*, two; *chromium*]. A salt containing two equivalents of chromic acid.

Bicipital (*bi-sip' - it - al*) [*bi*, two; *caput*, the head]. Pertaining to the biceps muscle.

Biconcave (*bi-kon' - kav*). See Lens.

Biconvex (*bi-kon' - veks*). See Lens.

Bicuspid (*bi-kus' - pid*) [*bi*, two; *cuspis*, the point of a spear]. Having two cusps, as *B. Teeth*. **B. Valve**, the mitral valve of the heart.

Bidder's Ganglia. See Ganglia, Table of.

Bidet (*be-del'*, or *-da*) [Fr. for pony]. A tub or basin with fixed attachments for the administering of injections; also for use as a sitz-bath or hip-bath.

Biedert's Cream Mixture. An infant's food made by mixing 4 oz. of cream with 12 oz. of warm water, and adding $\frac{1}{2}$ oz. of milk-sugar. It contains 1 per cent. of casein, 2.5 per cent. of fat, and 3.8 per cent. of sugar. As the child grows older a larger proportion of milk is added.

Biermer's Change of Note. See Signs and Symptoms, Table of.

Bifid (*bi' - fid*) [*bis*, twice; *findere*, to cleave]. Divided into two parts; cleft, as *bifid uvula*.

Bifocal (*bi-fo' - kal*) [*bi*, two; *focus*, a point]. Having a double focus. Applied to a system of lenses or spectacle-glasses with two foci, chiefly used for the correction of presbyopia, when there is at the same time an error of refraction for distant vision. The distance lens is above that for near work. These are sometimes called *Pantoscopic lenses*, and also *Franklin spectacles*, because the device was first made by Benjamin Franklin. They are also called *Cement lenses*, because now made by cementing the lower segment to the distance lens.

Bifurcate (*bi-fer' - kat*) [*bi*, two; *furca*, a fork]. Divided into two, like a fork.

Bifurcation (*bi-fer - ka' - shun*) [*bifurcatio*; from *bi*, two; *furca*, a fork]. Division into two branches, as of the trachea or of the aorta.

Bigelow's Ligament. The Y-ligament of the hip-joint; iliofemoral ligament. **B.'s Septum**, in anatomy, a layer of hard bony tissue in the neck of the thigh-bone.

Bilateral (*bi-lat' - er - al*) [*bi*, two; *latus*, a side]. Relating to two sides; pertaining to or affecting both sides of the body. **B. Symmetry**, the symmetry of right and left halves.

Bilateralism (*bi-lat' - er - al - izm*) [*bi*, two; *latus*, a side]. Bilateral symmetry.

Bile (*bil*) [*bilis*, the bile]. The substance secreted by the liver. It is mucilaginous, golden-brown, and is composed of biliary salts, cholesterin, mucus, and certain pigments. The principal salts are the sodium salts of taurocholic acid ($C_{26}H_{45}NSO_7$), and glycocholic acid ($C_{26}H_{43}NO_6$). The taste of bile is intensely bitter, its reaction feebly alkaline, and its density from 1026 to 1032. It exercises a diastatic action on starch, emulsifies fats, and precipitates soluble peptone; it is also the natural antiseptic of the intestines, and has an action on the intestinal mucous membrane through which absorption is favored. Its composition varies, but according to Hoppe-Seyler, in 100 parts bile there are of—

Water,	91.68 Parts.
Inorganic matter,	8.32 "
Organic matter,	1.39 "
Mucus,	0.87 "
Taurocholate of sodium,	3.03 "
Glycocholate of sodium,	1.39 "
Saponified fat,	0.35 "
Cholesterin,	0.53 "
Lecithin,	0.73 "
Fat,	

B. Pigments, the coloring-matters of the bile.

Bilharzia (*bil-har' - ze - ah*) [named after Theodor Bilharz, a physician of Cairo]. A genus of trematode worms, established by Cobbold, characterized by having the sexes separate.

B. hæmatobia. See *Distoma*.

Bilharziosis (*bil-har - ze - o' - sis*) [*Bilharzia*]. The group of symptoms produced by the presence in the intestine of worms of the genus *Bilharzia*.

Biliary (*bil' - e - a - re*) [*bilis*, bile]. Pertaining to the bile. **B. Acids**, glycocholic and taurocholic acids. **B. Colic**, colic produced by the passage of gall-stones. **B. Diabetes**, Hanot's disease, or hypertrophic cirrhosis of the liver with icterus. **B. Ducts**, the hepatic and cystic ducts, and the ductus communis choledochus, together with the small ducts in the liver itself.

Bilicyanin (*bil-e-si' - an - in*) [*bilis*, bile; *cyano*,

- veos*, blue]. Cholecyanin, choleverdin. A blue pigment obtained from bilirubin.
- Biliflavin** (*bil-e-fla'-vin*) [*bilis*, bile; *flavus*, yellow]. A yellow coloring-matter derivable from biliverdin.
- Bilifulvin** (*bil-e-ful'-vin*) [*bilis*, bile; *fulvus*, fulvous]. An impure form of bilirubin; also a yellow bile-color from ox-gall, not normally present in human bile.
- Bilifuscin** (*bil-e-fus'-in*) [*bilis*, bile; *fuscus*, brown], $C_{18}H_{20}N_2O_4$. A pigment occurring in bile and in gall-stones.
- Bilihumin** (*bil-e-hu'-min*) [*bilis*, bile; *humus*, earth]. An insoluble residue left after treating gall-stones with various solvents.
- Bilin** (*bi'-lin*) [*bilis*, bile]. A mixture of the taurocholate and glycocholate of sodium, forming a constituent of the bile.
- Bilineurin** (*bil-e-nu'-rin*). Cholin.
- Bilious** (*bil'-yus*) [*biliosus*, full of bile]. A term popularly applied to disorders supposed to arise from a too free secretion of bile. **B. Fever**, a remittent fever characterized by the vomiting of bile.
- Biliousness** (*bil'-yus-nes*) [*biliosus*, full of bile]. A popular name for a condition characterized by anorexia, constipation, coated tongue, lassitude, and headache, and supposed to be due to an excessive secretion of bile.
- Biliprasin** (*bil-e-pra'-sin*) [*bilis*, bile; *πράσινος*, leek; *πράσινος*, leek-green], $C_{16}H_{22}N_2O_6$. A pigment occurring in gall-stones, icteric urine, and bile.
- Bilirubin** (*bil-e-ru'-bin*) [*bilis*, bile; *ruber*, red], $C_{42}H_{54}N_4O_6$. A red coloring-matter, the chief pigment of the bile, and also found in the urine in jaundice. It is insoluble in water, and almost so in ether and alcohol, but it is readily soluble in alkaline solutions. It crystallizes in rhombic plates or prisms.
- Biliverdin** (*bil-e-ver'-din*) [*bilis*, bile; *viridis*, green], $C_{42}H_{54}N_4O_6$, or $C_8H_8NO_2$. A green pigment, the first product of the oxidation of bilirubin. It gives the characteristic color to the bile of herbivora, and occurs in the urine in jaundice, and in gall-stones.
- Billroth's Mixture**. See *Anesthetic*.
- Bimanual** (*bi-man'-u-al*) [*bi*, two; *manus*, a hand]. With both hands; two-handed. **B. Palpation**, palpation by means of both hands.
- Binary** (*bi'-nar-e*) [*binus*, a couple]. In chemistry, compounded of two elements. In anatomy, separating into two branches.
- Binaural** (*bin-aw'-ral*) [*bis*, twice; *auris*, ear]. Pertaining to or having two ears; used for both ears. **B. Stethoscope**, a stethoscope with two tubes, one for each ear.
- Binauricular** (*bin-aw-rik'-u-lar*) [*bis*, twice; *auricula*, the ear]. Pertaining or belonging to both ears.
- Binder** (*bind'-er*) [ME., *byndere*]. A wide bandage about the abdomen, worn by women after labor, or after celiotomy, to support the abdominal walls. **B., Mammary**, a sling or suspensory for the mamma.
- Binioidid** (*bin-i'-o-did*). Having two atoms of iodine in the molecule.
- Binocular** (*bin-ok'-u-lar*) [*bi*, two; *oculus*, an eye]. Pertaining to both eyes. In optics, an instrument with two eye-pieces for use with both eyes at once. **B. Vision**, the faculty of using both eyes synchronously and without diplopia.
- Binoxid** (*bin-oks'-id*) [*bi*, two; *oxid*]. See *Dioxid*.
- Bioblast** (*bi'-o-blast*) [*βίος*, life; *βλαστός*, a germ]. A plastidule or formative cell; a corpuscle that has not yet become a cell.
- Biochemics** (*bi-o-kem'-iks*) [*βίος*, life; *χημεία*, chemistry]. The chemistry of life. See *Biochemy*.
- Biochemistry** (*bi-o-kem'-is-tre*) [*βίος*, life; *χημεία*, chemistry]. The chemistry of the living body, or of life.
- Biochemy** (*bi'-o-kem-e*) [*βίος*, life; *χημεία*, chemistry]. Chemic force as exhibited in living organisms.
- Biodynamics** (*bi-o-di-nam'-iks*) [*βίος*, life; *δύναμις*, power]. The dynamics of life; dynamic biology.
- Biogenesis** (*bi-o-jen'-es-is*) [*βίος*, life; *γένεσις*, origin]. 1. The doctrine that living things are produced only from living things—the reverse of abiogenesis. 2. Reproduction from living parents.
- Biogenetic** (*bi-o-jen'-et'-ik*) [*βίος*, life; *γένεσις*, origin]. Pertaining to biogenesis. **B. Law**, the fact that a certain tendency directs the drift or trend of development of a being along a line parallel with that of the series of forms ancestral to it. The being in the course of its development briefly recapitulates that of the ancestral series to which it belongs. This is also called *Müller's Law*.
- Biogeny** (*bi-oi'-en-e*) [*βίος*, life; *γένεσις*, origin]. In biology, the evolution of organic forms, either considered individually (*ontogeny*) or tribally (*phylogeny*).
- Biologic** (*bi-o-loj'-ik*) [*βίος*, life; *λόγος*, science]. Pertaining or belonging to biology.
- Biologist** (*bi-ol'-o-jist*) [*βίος*, life; *λόγος*, science]. One who is a student of biology.
- Biologos** (*bi-o-log'-os*) [*βίος*, life; *λόγος*, a word]. A designation proposed for the intelligent living power displayed in cellular and organic action and reaction.
- Biology** (*bi-ol'-o-jē*) [*βίος*, life; *λόγος*, science]. The science embracing the structure, function, and organization of living forms.
- Biometry** (*bi-on'-et-re*) [*βίος*, life; *μέτρον*, a measure]. Life-measurement; the estima-

tion of the probable duration of any given life-form—in the past or future.

Biondi's Fluid. A staining medium used in histologic laboratories. It is a mixture of orange-G, methyl-green, and acid-fuchsin.

Bionergy (*bi-on'-er-jē*) [*βίος*, life; *ἔργον*, work]. Life-force; force exercised in the living organism.

Bionomy (*bi-on'-o-mē*) [*βίος*, life; *νόμος*, a law]. Dynamic biology; biodynamics; the science of the laws of life.

Biophagous (*bi-off'-ag-us*) [*βίος*, life; *φαγεῖν*, to eat]. Feeding upon living organisms, or upon living tissue, as insectivorous plants.

Biophysiology (*bi-o-fiz-e-ol'-o-jē*) [*βίος*, life; *φύσις*, nature; *γράφειν*, to write]. Descriptive or structural biology; organography, as distinguished from biophysiology.

Biophysiology (*bi-o-fiz-e-ol'-o-jē*) [*βίος*, life; *φύσις*, nature; *λόγος*, science]. The branch of biology including organogeny, morphology, and physiology.

Bioplasm (*bi'-o-plazm*) [*βίος*, life; *πλάσμα*, form]. Any living matter, but especially germinal or forming matter; matter possessing reproductive vitality.

Bioplast (*bi'-o-plast*) [*βίος*, life; *πλαστός*, formed]. A mass or cell of bioplasm that is a unit of living matter.

Biorbital (*bi-or'-bit-al*) [*bi*, two; *orbita*, a circle]. Relating to both orbits.

Bioscope (*bi'-o-skōp*) [*βίος*, life; *σκοπεῖν*, to view]. An instrument used in bioscopy.

Bioscopy (*bi-ol'-ko-pē*) [*βίος*, life; *σκοπεῖν*, to examine]. Examination of the body to ascertain whether life is present. **B.**, Electro-, examination by the aid of the electric current. The muscular reaction is lost to faradic stimulation in about two hours after death in the tongue; after three or four hours in the extremities; after five or six hours in the trunk. The reaction to galvanism persists somewhat longer.

Biosis (*bi-ol'-is*) [*βίος*, life]. Life; vitality.

Biostatics (*bi-o-stat'-iks*) [*βίος*, life; *στατικός*, causing to stand]. Static biology; the science of the determinate parts of biology, including anatomy, and the physics of the living body.

Biotic (*bi-ol'-ik*) [*βίος*, life]. Pertaining to life or to the laws of animal and vegetable progress and evolution.

Biotomy (*bi-ol'-o-mē*) [*βίος*, life; *τέμνειν*, to cut]. Vivisection.

Biparasitic (*bi-par-as-it'-ik*) [*bi*, two; *παράσιτος*, a parasite]. Parasitic upon a parasite.

Biparietal (*bi-par-i'-et-al*) [*bi*, two; *paries*, a wall]. Relating to both parietal bones.

B. Diameter, the distance from one parietal eminence of the cranium to the other.

Biparous (*bi-p'-ar-us*) [*bi*, double; *parere*, to bring forth]. Producing two at a birth.

Bipartite (*bi-par'-it*) [*bi*, two; *pars*, a part]. In biology, composed of two parts or divisions.

Biped (*bi'-ped*) [*bi*, two; *pes*, a foot]. 1. Having two feet. 2. An animal with two feet.

Bipolar (*bi-po'-lar*) [*bi*, two; *polus*, a pole]. Having two poles. **B. Nerve-cells**, nerve-cells that have two prolongations of the cell matter. They are found chiefly in the spinal ganglia.

Birch (*berch*) [AS., *birce*]. Any tree of the genus *Betula*. Birch-tar, or the tarry oil of *B. alba*, is useful in certain skin-diseases. The bark of *B. lenta*, the American black birch, yields a fragrant volatile oil, identical with that of *Gaultheria procumbens*, for which it is extensively substituted. **B. Camphor**. Same as *B. Resin*. **B. Resin**. See *Betulin*.

Bird-lime (*berd'-lm*). A viscous vegetable substance used in Japan as a local dressing for wounds.

Bird's Formula. The last two figures of the specific gravity of urine nearly represent the number of grains of solids to the ounce contained in the urine. The same two figures multiplied by 2 (*Trapp's Factor*) give the parts per 1000.

Birth (*berth*) [ME., *byrth*]. 1. The delivery of a child; parturition. 2. That which is brought forth in parturition. **B.-mark**. See *Nevus pigmentosus*. **B.-palsy**, any paralytic affection due to an injury received at birth; less correctly, a congenital paralytic affection due to a lesion that existed in the fetal state.

Bisacromial (*bis-ak-ro'-me-al*) [*bis*, two; *acromion*]. Relating to the two acromia.

Biscara Button (*bis'-kar-ah but'-un*). See *Furunculus orientalis*.

Bisexual (*bi-seks'-u-al*) [*bi*, two; *sexus*, sex]. Having the reproductive organs of both sexes; hermaphrodite.

Bisiliac (*bis-il'-e-ak*) [*bis*, two; *iliacus*]. Relating to the two most distant points of the two iliac crests.

Biskra Boil, or **Button**. See *Furunculus orientalis*.

Bismarck Brown. A brown, basic anilin dye, extensively used as a stain and counterstain in histology.

Bismuth, or **Bismuthum** (*biz'-muth, biz-mu'-thum*) [L.]. Bi = 210; quantivalence 1, III, V. A pinkish-white crystalline metal. Its commercial salts often contain arsenic. The insoluble salts of bismuth are feebly astringent. The derivatives of bismuth are chiefly employed as astringents and sedatives to mucous membranes and as gastrointestinal antiseptics. The soluble salts are irritant in large doses. **B. et Ammonii Citras**, soluble in water. Dose gr. j-v (0.065-0.32). **B. et Ammon. Citratis, Liqueur** (B. P.).

Dose f 3 ss-j (2.0-4.0). **B. Carbonas** (B. P.), $(\text{Bi}_2\text{O}_3\text{CO}_3)_2\cdot\text{H}_2\text{O}$. Dose gr. v-xx (0.32-1.3). **B. Citras**, $\text{Bi}_2\text{C}_4\text{H}_5\text{O}_7$, soluble in water of ammonia. Dose gr. ij-v (0.13-0.32). **B. Oxid** (B. P.), Bi_2O_3 . Dose gr. v-xv (0.32-1.0). **B. Oxychlorid**, pearl white. Unof. It is used as a cosmetic. **B., Pulvis, Comp., Ferrier's Snuff**, contains two grains of hydrochlorate of morphin in one ounce, with bismuth and acacia. It is used in the treatment of coryza. **B. Salicylas**, $\text{Bi}(\text{C}_7\text{H}_5\text{O}_3)_3\cdot\text{Bi}_2\text{O}_3$. Dose gr. v-x (0.32-0.65); an intestinal antiseptic. **B. Subcarbonas**, $(\text{BiO})_2\text{CO}_3\cdot\text{H}_2\text{O}$, insoluble. Dose gr. x-3j (0.65-4.0); feebly astringent and sedative. **B. Subgallas**. See *Dermatol.* **B. Subiodid**, BiOI , used as an antiseptic dusting-powder, like iodoform. **B. Subnitras**, $\text{BiONO}_3\cdot\text{H}_2\text{O}$, the salt chiefly used in medicine as a sedative astringent to the gastrointestinal mucous membrane. Dose gr. x-3j (0.65-4.0). **B., Trochisci** (B. P.), each contains gr. ij (0.13).

Bismuthosis (*bis-muth-o'-sis*). Chronic bismuth-poisoning; it may follow the use of the soluble salts.

Bistoury (*bis'-too-re*) [*Fr. bistouri*]. A long, slender knife, straight or curved, used in surgery.

Bisulphate (*bi-sul'-fat*) [*bi*, two; *sulphur*]. A sulphate in which the base replaces but one of the two hydrogen atoms of the acid. Any acid sulphate having twice the amount of acid contained in the corresponding normal sulphate.

Bitartrate (*bi-tar'-trat*) [*bi*, two; *τάρταρον*, tartar]. Any tartrate in which one replaceable hydrogen atom has been replaced by a base.

Bitemporal (*bi-tem'-por-al*) [*bi*, two; *tem-pora*, the temples]. Pertaining to the two temples.

Bitter (*bit'-er*) [*AS., bitan*, to bite]. A peculiar, well-known taste, of which that of quinin is an example; unpalatable. **B. Almond**, the nut of the *Amygdalum amarum*. It contains hydrocyanic acid. **B. Almond Oil**, oleum amygdalæ amaræ. See *Benzaldehyd.* **B. Apple**, the fruit of the colocynth, a purgative remedy. **B. Blain**, a West Indian herb, *Vandellia diffusa*, employed in fevers and in hepatic disorders. Unof. **B. Bugleweed**, the herb *Lycopus europeus*, alterative and tonic. Dose of fld. ext. f 3 ss-j (16.0-32.0). Unof. **B. Cucumber**, colocynth. **B. Cup**, a cup made of quassia wood. It is a tonic. **B. Purging Salt**, magnesium sulphate. **B. Root**, the root of *Gentiana catartica*, a tonic. **B. Tincture**, the *tinctura amara* (N. F.), prescribed also in the German pharmacy; it is a tincture of gentian, centaury, bitter orange-peel, orange-berries, and zedoary; it is known also as

Stomach-drops. **B. Wine of Iron**, a solution of white wine, syrup, citrate of iron, and quinin.

Bitters (*bit'-ers*) [*AS., bitan*, to bite]. Medicines characterized by a bitter taste. **B., Aromatic**, medicines that unite the properties of aromatics with those of simple bitters. **B., Simple**, medicines that stimulate the gastrointestinal tract without influencing the general system. **B., Styptic**, medicines that add styptic and astringent properties to those of bitterness.

Bittersweet (*bit'-er-sweet*). See *Dulcamara*.

Bitumen (*bit'-u'-men*) [*L., bitumen*]. Mineral pitch or oil, composed of various hydrocarbons. In the solid form it is usually called *asphalt*; in the liquid form, *petroleum*. An intermediate form is known as mineral tar or *maltha*. By distillation, bitumen yields benzol, naphtha, paraffin, and various other hydrocarbons, liquid and gaseous.

Biuret (*bi'-u-ret*) [*bi*, two; *οὐρον*, urine], $\text{C}_2\text{H}_5\text{N}_3\text{O}_2 + \text{H}_2\text{O}$. A compound formed by exposing urea to a high temperature for a long time. It is readily soluble in water and in alcohol; it crystallizes with one molecule of water in the form of warts and needles. When anhydrous, biuret melts at 190° with decomposition. **B. Reaction**, or **Test**, a reaction given by proteids and by urea when heated (when biuret is formed). It consists in the development of a violet color when a few drops of a solution of cupric sulphate and an excess of potassium or sodium hydrate are added to the substance.

Bivalence (*bi'-al-ens*) [*bi*, two; *valens*, having power]. In chemistry, a valence or saturating power double that of the hydrogen atom.

Bivalent (*bi'-al-ent*) [*bi*, two; *valens*, having power]. In chemistry, applied to an element of which an atom can replace two atoms of hydrogen or other univalent element, or to a radicle that has the same valence as such an element.

Bivalve (*bi'-valve*) [*bi*, two; *valva*, a valve]. Having two valves or shells, as a speculum.

Biventer (*bi-ven'-ter*) [*bi*, two; *venter*, a belly]. Having two bellies, as a muscle. **B. cervicis**, the inner portion of the complexus muscle. **B. maxillæ**, the digastricus muscle.

Biventral (*bi-ven'-tral*) [*bi*, two; *ventrum*, a stomach]. Having two stomachs; having two bellies, as the digastric muscle.

Bixin (*bik'-in*) [*Bixa*, a genus of plants], $\text{C}_{16}\text{H}_{24}\text{O}_2$. An orange-red coloring matter found in annatto.

Bizygomatic (*bi-zi-go-mat'-ik*) [*bi*, two; *ζυγωα*, the zygomatic arch]. Relating to the most prominent points on the two zygomatic arches.

Bizzozzero's Blood-plates. See *Blood-plates*.

Black (*black*) [A.S., *blac*]. Characterized by an absence of color. The appearance of an object from the surface of which none of the spectrum colors are reflected. **B. Alder.** See *Prinos*. **B. Antimony**, antimonium tersulphid, SbS_3 . **B. Ash**, the bark of *Fraxinus sambucifolia*, a mild tonic and astringent. Dose of fld. ext. f 3ss-j (2.0-4.0). Unof. **B. Blood**, venous blood. **B. Cohosh.** See *Cimicifuga*. **B.-damp**, carbon dioxid gas, which is found in greater or less quantity in all collieries, being given off by many coals, either mixed with fire-damp, or separately, or produced in various other ways, as by the exhalations of the miners, by fires, and by explosions of fire-damp. Also called choke-damp. **B. Death.** See *Plague*. **B. Disease**, a disease of malarial origin and pernicious course, characterized by extreme darkening of the skin, which may be brown or black in color. It occurs in the Garo Hills in Assam. **B. Draught.** See *Senna*. **B. Drop.** See *Opium*. **B. Erysipelas.** Synonym of *Anthrax*. **B. Eye**, livor (or sugillation) oculi; ecchymosis of the tissues about the eye, usually from injury. **B. Haw.** See *Viburnum*. **B.-head.** See *Comedo*. **B. Hellebore.** See *Hellebore*. **B. Jaundice**, an excessive jaundice arising from obstruction of the gall-ducts. The color of the skin is greenish-black. **B. Lead**, a form of carbon properly known as the mineral graphite. **B.-Leg**, a febrile, generally fatal disease chiefly affecting cattle and sheep, which is characterized by the rapid appearance of irregular nodules in the skin and muscular tissues, that are at first tense and very painful, but rapidly become painless and crepitating. It is caused by the *Bacillus chauvæi*, or *B. of Symptomatic Anthrax*. It is also called *Symptomatic Anthrax*. **B. Measles**, *rubeola nigra*; a grave or malignant form of measles. **B. Pepper.** See *Piper nigrum*. **B. Phthisis**, a synonym of *Miner's Phthisis*. **B. Sickness**, cerebrospinal fever. **B. Snake-root.** See *Cimicifuga*. **B. Spauld.** See *Black-Leg*. **B. Spit.** See *Miner's Phthisis*. **B. Tongue.** 1. A name given to a condition characterized by the formation upon the dorsum of the tongue of a hair-like deposit that passes through various stages of coloration from yellow to brown and finally black, ultimately disappearing by desquamation as gradually as it came. Repeated recurrence is the rule. It is probably an undue proliferation of the epithelium, the result of irritation. 2. A term applied to erysipelatous glossitis. **B. Vomit**, the coffee-ground vomit of yellow fever, etc. **B. Walnut**, the leaves of *Juglans nigra*, a

tonic, alterative, and deobstruent. Dose of fld. ext. m xx-xxx (1.3-2.0). Unof. **B. Wash.** See *Hydrargyrum*. **B. Willow**, the buds of *Salix nigra*, a bitter tonic with aphrodisiac properties. Dose of fld. ext. m xv-[3j (1.0-4.0). Unof.

Blackberry (*black'-ber-ry*) [A.S., *blac*; *berie*, berry]. See *Rubus*. **B. Cordial**, the *cordiale rubi fructus* (N.F.). Its formula is: blackberry juice, 3 pints; cinnamon, in coarse powder, 2 Troy oz.; cloves and nutmeg, in coarse powder, each $\frac{1}{2}$ Troy oz.; dilute alcohol, 2 pints; syrup, 3 pints.

Blackwater-fever. A very fatal infectious disease occurring especially on the low coasts of tropic Africa, but also in Sicily, Greece, Central and South America, Java, and New Guinea. The disease is almost exclusively confined to the white race and is characterized by a chill, an irregular intermittent or remittent fever, vomiting, dyspnea, jaundice, and hemoglobinuria. An almost invariable complication is nephritis. Studies of the blood have revealed the presence of a small, annular, non-pigmented, intracellular parasite; also forms having a rosette or crescent shape.

Bladder (*blad'-er*) [A.S., *bladre*, a blister]. 1. A membranous sac serving for the reception of fluids or gases. 2. The receptacle for the urine. **B., Atony of**, inability to expel the urine, from deficient muscular power. **B., Catarrh of.** See *Cystitis*. **B., Extrophy**, or *Extroversion of*, absence of the anterior wall of the bladder, and more or less deficiency of the corresponding part of the abdomen. **B., Gall-**, the receptacle for the bile, situated on the lower surface of the liver. **B., Irritable**, a condition characterized by constant desire to urinate. **B., Neck of**, the constricted portion continuous with the urethra in front. **B., Nervous**, a condition in which there is a frequent desire to pass water, with inability at the same time to perform the act perfectly, and consequent slight dribbling at its close. **B., Sacculated**, pouches formed between the hypertrophied muscular fibers. **B. Worm.** See *Worm*.

Blancard's Pills (*blang'-kardi*). Pills of the iodid of iron.

Bland (*bland*) [*blandus*, mild]. Mild; soothing.

Blastema (*blas'-te'-mah*) [*βλαστημα*, from *βλαστειν*, to germinate]. The formative lymph or rudimentary tissue, from which, by differentiation, tissues and organs are developed.

Blastochyle (*blas'-to-ki'*) [*βλαστός*, a germ; *χυλός*, juice]. The colorless fluid in the blastodermic vesicles.

Blastocyst (*blas'-to-sist*) [*βλαστός*, a germ; *κύστις*, a bladder]. The germinal vesicle.

Blastoderm (*blas'-to-derm*) [*βλαστός*, a germ; *δέρμα*, the skin]. In embryology, the germinal membrane formed by the cells of the morula, lying on the internal surface of the vitelline membrane of the impregnated ovum. The whole hollow sphere, with its surrounding cells, is called the *blastodermic vesicle*, and is formed about the tenth day. The *ectoderm* (or *epiblast*) and the *endoderm* (*entoderm* or *hypoblast*) layers are simply due to a proliferation of the blastodermic cells about the *germinal area*, whereby the blastoderm is doubled, thus forming these outer and inner layers. The *mesoblast* or middle layer is developed after the others, and probably from the hypoblast.

Blastodermic (*blas-to-der'-mik*) [*βλαστός*, a germ; *δέρμα*, the skin]. Relating to the blastoderm.

Blastodisc (*blas'-to-disk*) [*βλαστός*, a germ; *δίσκος*, disc]. A blastodermic disc; a mass or disc capping one pole of the yolk.

Blastogeny (*blas-toj'-en-e*) [*βλάστος*, a germ; *γένεσις*, generation]. Haeckel's term for the germ-history of an individual organism; a division of ontogeny.

Blastomere (*blas'-to-mēr*) [*βλαστός*, a germ; *μέρος*, a part]. Any one of the nucleated cells or segments into which the fecundated vitellus divides.

Blastoneuropore (*blas-to-nu'-ro-pōr*) [*βλαστός*, a germ; *νεῦρον*, a nerve; *πόρος*, a pore]. In biology, the temporary aperture in certain embryos formed by the coalescence of the *blastopore* and *neuropore*.

Blastophore (*blas'-to-fōr*) [*βλαστός*, a germ; *φέρειν*, to bear]. That part of a sperm-cell that does not become converted into spermatozoa.

Blastophyllum (*blas-to-fil'-um*) [*βλαστός*, a germ; *φύλλον*, a leaf]. The endoderm or ectoderm; a primitive germ-layer.

Blastophyly (*blas-toff'-il-e*) [*βλαστός*, a germ; *φυλή*, a tribe]. The tribal history of individual organisms.

Blastopore (*blas'-to-pōr*) [*βλαστός*, a germ; *πόρος*, passage, pore]. The small opening leading into the notochordal canal, or, after the canal has fused with the yolk-cavity, leading into the archenteron. It is situated at the hind end of the primitive axis, and is a small portion of the gastrula mouth.

Blastula (*blas'-tu-lah*) [dim. of *βλαστός*, a germ]. The blastodermic vesicle.

Blaud's Pill (*blohs*). A pill containing equal parts iron sulphate and potassium carbonate; for use in anemia, etc.

Bleach (*bleck*) [ME., *blæcan*, to make white]. To make white or pale. **Bleaching Powder**, chlorinated lime, a mixture of calcium chlorid and calcium hypochlorite, containing free chlorin-gas. It is used as a

disinfectant. **Bleaching Fluid**, *eau de Javelle*, a fluid obtained by passing chlorin-gas into an emulsion of calcium hydrate.

Bleacher's Eczema (*blech'-erz ek'-zem-ah*). Eczema of the hands of bleachers, due to the use of hot water and strong lye.

Bleary Eye (*bler'-i*). See *Blepharitis ulcerosa*.

Bleb (*bleb*). See *Bulla*.

Bleeders (*ble'-derz*) [AS., *bltdan*, to bleed]. A popular term for those who are subjects of the hemorrhagic diathesis. **B's Disease**. See *Hemophilia*.

Blennorrhagia (*blen-or-a'-je-ah*) [*βλέννα*, mucus; *ῥεγνίναί*, to burst forth]. 1. An excessive mucous discharge. 2. Gonorrhea.

Blennorrhagic (*blen-or-a'-jik*) [*βλέννα*, mucus; *ῥεγνίναί*, to burst forth]. Relating to blennorrhagia.

Blennorrhea (*blen-or-e'-ah*) [*βλέννα*, mucus; *ῥεῖν*, to flow]. Same as *Blennorrhagia*.

Blennorrhéal (*blen-or-e'-al*) [*βλέννα*, mucus; *ῥεῖν*, to flow]. Relating to blennorrhea.

Blepharadenitis (*blef-ar-ad-en-i'-tis*) [*βλέφαρον*, the eyelid; *ἀδὴν*, a gland; *ιτις*, inflammation]. Inflammation of the Meibomian glands.

Blepharitis (*blef-ar-i'-tis*) [*βλέφαρον*, the eyelid; *ιτις*, inflammation]. Inflammation of the eyelids. **B. ciliaris**, or **B. marginalis**, inflammation of the ciliary or marginal border of the lids. **B. ulcerosa**, an ulcerative inflammation of the eyelids.

Blepharoadenitis (*blef-ar-o-ad-en-i'-tis*). See *Blepharadenitis*.

Blepharoadenoma (*blef'-ar-o-ad-en-o'-mah*) [*βλέφαρον*, the eyelid; *ἀδὴν*, a gland; *δμα*, a tumor]. An adenoma of the eyelid.

Blepharoatheroma (*blef-ar-o-ath-er-o'-mah*) [*βλέφαρον*, an eyelid; *ἀθήρη*, gruel; *δμα*, tumor]. A sebaceous cyst of the eyelid.

Blepharochromidrosis (*blef-ar-o-kro-mid-ro'-sis*) [*βλέφαρον*, the eyelid; *χρῶμα*, color; *ιδρώς*, sweat]. Colored sweat of the eyelids, usually of a bluish tint.

Blepharophimosis (*blef-ar-o-fi-mo'-sis*) [*βλέφαρον*, the eyelid; *φίμωσις*, a shutting up]. Abnormal smallness of the palpebral aperture.

Blepharoplasty (*blef'-ar-o-plas-te*) [*βλέφαρον*, the eyelid; *πλάσσειν*, to form]. An operation for the restoration of any part of the eyelid.

Blepharoplegia (*blef-ar-o-ple'-je-ah*) [*βλέφαρον*, the eyelid; *πληγή*, a stroke]. Paralysis of an eyelid.

Blepharoptosis (*blef-ar-op-to'-sis*) [*βλέφαρον*, the eyelid; *πτῶσις*, a fall]. Drooping of the upper eyelid.

Blepharorrhaphy (*blef-ar-or'-a-fe*) [*βλέφαρον*, the eyelid; *ῥαφή*, a seam]. The stitching together of a portion of the edges of the lids.

Blepharospasm (*blef'-ar-o-spa-zm*) [*βλέφαρον*, the eyelid; *σπασμός*, a spasm]. Spasm of

the orbicularis palpebrarum muscle; excessive winking.

Blepharostat (*blef'-ar-o-stat'*) [*βλεφαρον*, the eyelid; *ιστάμι*, to cause to stand]. An instrument for holding the eyelids apart whilst performing operations upon the eyes or lids.

Blepharostenosis (*blef'-ar-o-ste-no'-sis*) [*βλεφαρον*, the eyelid; *στενός*, narrow]. Diminution of the space between the eyelids.

Blepharosynechia (*blef'-ar-o-sin-ek-i'-ah*) [*βλεφαρον*, the eyelid; *συνήχεια*, a holding together]. The adhesion or growing together of the eyelids.

Blepharotomy (*blef'-ar-ot'-o-me*) [*βλεφαρον*, the eyelid; *τέμνω*, to cut]. Incision into the eyelid.

Blessed Thistle (*bles'-ed this'-l*). See *Centaurea benedicta*.

Blind (*blind'*) [AS., *blind*]. Without sight; deprived of sight. **B. Spot**, that part of the fundus of the eye where the optic nerve enters.

Blindness (*blind'-nes*) [AS., *blind*]. Want of vision. **B., Color-**, imperfect color-perception. This condition is found in about 4 per cent. of persons, is more frequent in men than in women, and is probably due to non-exercise of the color-sense. **Complete Color-B.** is very rare, the different colors probably appearing as different intensities or shades of white light. In **Partial Color-B.** defective perception of red is the most frequent, green, blue, and yellow, respectively, being next in order. **Tests for Color-B.** usually consist in matching and classifying colored yarns. **B., Cortical**, blindness due to lesion of the cortical center of vision.

B., Day-. See *Nyctalopia*. **B., Mind-**. See *B., Psychic*. **B., Moon-**, a rare condition of retinal anesthesia said to be due to exposure of the eyes to the moon's rays in sleeping. **B., Night-**. See *Hemeralopia*. **B., Object-**. See *Apraxia*. **B., Psychic**, loss of conscious visual sensation from destruction of the cerebral visual center; there is sight but not recognition. **B., Snow-**, photophobia and conjunctivitis due to exposure of the eyes to the glare of sunlight upon snow. **B., Word-**. See *Aphasia*.

Blister (*blis'-ter*) [ME., *blister*, a vesicle]. A vesicle resulting from the exudation of serous fluid between the epidermis and true skin. Also the agent by which the blister is produced. **B., Fly-**, a blister of cantharides. See *Cantharis*. **B., Flying**, a blister that remains long enough to produce redness of the skin and not vesication.

Blistering (*blis'-ter-ing*) [ME., *blister*, a vesicle]. Forming a vesicle upon the skin. **B. Collodion**, a solution of cantharidin in collodion. **B. Liquid**. Same as *B. Collodion*. **B. Paper**, paper saturated with cantharides; used for producing vesication.

Blood (*blud'*) [AS., *blōd*]. The fluid that circulates through the heart, arteries, and veins, supplying nutritive material to all parts of the body. In the human being the blood of the arteries is bright-red and dichroic; that of the veins dark-red and monochroic. Blood consists of plasma and corpuscular elements, the latter being the red corpuscles, the white corpuscles, and the blood-plaques. In a cubic millimeter there are about 5,000,000 red and 10,000 white corpuscles. The red color of the blood is due to the hemoglobin contained in the red corpuscles. The total amount of blood is estimated at $\frac{1}{12}$ of the body-weight. When exposed to the air it coagulates, forming a red clot and a yellowish fluid called serum. Healthy blood consists of 78 per cent. of water, and 22 per cent. solids. **B.-Casts**, tube-casts to which red blood-corpuscles are attached. **B.-Corpuscles**, Red, circular, biconcave discs floating in the blood. Red corpuscles are circular in mammals (except the camel), and elliptical in birds and reptiles. In man they are about $\frac{1}{2500}$ inch (7μ) in diameter and $\frac{1}{1250}$ inch thick. *Colorless corpuscles* are about one-third larger in diameter, $\frac{1}{1500}$ inch (10μ). They exhibit movements similar to those of the amoeba. **B.-crasis**, the mixture of the constituents of the blood. When the blood-crasis is diseased or disordered, the condition is known as dyscrasia. **B.-crystals**, crystals of hematin. **B., Defibrinated**, blood from which the fibrin has been removed by agitating it with twigs. **B.-disease**, a synonym of *Dyscrasia*. **B.-fluke**. See *Bilharzia hematobia*. **B.-islands**, a term applied to the groups of corpuscles developed during the first days of embryonic life, within the large branched cells of the mesoblast. **B.-plaque**. See *B.-platelets*. **B.-plasma**, the *liquor sanguinis*, or fluid part of the blood. **B.-plate**. See *B.-platelets*. **B.-platelets**, circular or oval, light-gray bodies found in the blood. They are from 1 to 1.3μ in size, and number from 18,000 to 300,000 in the cubic millimeter. Their function is not definitely known; they are an important factor in thrombosis. **B.-poisoning**, a morbid state due to the circulation of bacteria or their products, or both, in the blood, as the result of a local infection. **B.-pressure**, the tension of the blood in the arteries. **B.-root**. See *Sanguinaria*. **B.-tumor**. See *Hematoma*.

Bloodless (*blud'-less*) [AS., *blōd*]. Without blood. **B. Operations**, surgical operations, such as amputations, in which the member is so bandaged by compresses and elastic bands that the blood is expelled and kept from the part to be operated upon.

Bloodletting (*blud-let'-ing*) [AS., *blōd*;

letan]. The artificial abstraction of blood from the body. **B.**, General, venesection or phlebotomy. **B.**, Local or Topic, the removal of a small quantity of blood by cupping, leeching, or scarification.

Blue (*bleu*) [M.E., blew]. One of the colors of the spectrum. **B.-baby**, a child with congenital cyanosis. **B.-bottle**. See *Centaurea*. **B.** Blindness, defective color-perception for blue. **B. Cohosh**. See *Caulophyllum*. **B. Disease**, cyanosis of the newborn, usually due to congenital disease of the heart. **B. Flag**. See *Iris*. **B. Gentian**, the root of *Gentiana catesbaei*, tonic and stomachic. Dose of fld. ext. $\mathfrak{m}\text{x}-\text{xl}$ (0.65-2.65). Unof. **B. Gum Tree**. See *Eucalyptus*. **B. Cardinal Flower**. Synonym of *Lobelia syphilitica*. **B. Edema**, a puffed and bluish appearance of the limb sometimes seen in hysteric paralysis. **B. Mass**. See *Hydrargyrum*. **B.**, Methylene. See *Methylene Blue*. **B. Ointment**. See *Hydrargyrum*. **B. Pill**. See *Hydrargyrum*. **B.**, Prussian, ferric ferrocyanid, $\text{Fe}_4\text{Fe}(\text{C}_6\text{N}_6)_2$. **B. Stick**. Same as *B. Stone*. **B. Stone**, copper sulphate. **B.**, Turnbull's, ferrous ferrocyanid, $\text{Fe}_2\text{Fe}_2(\text{C}_6\text{N}_6)_4$. **B. Vitriol**, copper sulphate.

Blunt-hook (*blunt'-hook*). An obstetric instrument, used mainly in embryotomy.

Boas's Reagent. A reagent for detecting the presence of hydrochloric acid in the gastric juice. It consists of resorcin 5, sugar 3, dilute alcohol 100 parts.

Bochdalek's Ganglion. See *Ganglia*, Table of.

Body (*bod'-e*) [A.S., *bodig*]. 1. The animal frame with its organs. Also a cadaver or corpse. 2. The important and largest part of an organ, as the body of the uterus. See also *Corpus* and *Corpora*. **B. Louse**. See *Pediculus*.

Boil (*boil*) [M.E., *boile*]. A furuncle; a localized inflammation of the skin and subcutaneous connective tissue attended by the formation of pus. See *Furunculus*. **B.**, Aleppo, or **B.**, Delhi, a peculiar ulcerative affection endemic in India, due to a specific microorganism.

Boiled Oil. Linseed oil that has been heated to a high temperature (130° C. and upward), while a current of air is passed through or over the oil, and the temperature increased until the oil begins to effervesce from evolution of products of decomposition.

Boiler-makers' Deafness. See *Deafness*.

Boiling (*boil'-ing*) [M.E., *boilen*, to stir]. The vaporization of a liquid when it gives off vapor having the same tension as the surrounding air.

Boldo (*bol'-do*). See *Boldus*. **B.-glucine**, an aromatic glucosid obtained from *Peumus*

boldo and other species. It is an hypnotic in doses of gr. $\text{xx}-3\text{j}$ (1.3-4.0). Unof.

Boldus (*bol'-dus*) [L.]. Boldo. The leaves and stems of an evergreen, *Peumus boldo*, native to Chili, sometimes used in anemia and general debility as a substitute for quinin. It contains a bitter alkaloid, *Boldin*, an hypnotic in doses of gr. ij (0.2). **B.**, Ext., Fld. Dose $\mathfrak{m}\text{j}-\text{iv}$ (0.065-0.26). **B.**, Tinct., contains 20 per cent. of the drug. Dose $\mathfrak{m}\text{v}-\text{viii}$ (0.32-0.52).

Bole (*bol*) [*βῶλος*, a clod of earth]. A translucent, soft variety of clay formerly much used in medicine, internally as an astringent, externally as an absorbent. Several varieties are used, as the *Armenian bole*, the *Lemnian*, and the *French bole*.

Boletus (*bo-le'-tus*) [*βολιτης*, a kind of mushroom]. A genus of fungi, some species of which are edible, while others are highly poisonous.

Bolus (*bol'-lus*) [L.]. 1. A large pill. 2. The rounded mass of food prepared by the mouth for swallowing. 3. See *Bole*.

Bone (*bon*) [A.S., *bān*, a bone]. A hard tissue that constitutes the framework or skeleton of the body. Bone usually consists of a compact outer mass covered with periosteum, surrounding a reticulated inner structure that encloses a central cavity filled with marrow. A transverse section of a long bone shows bone-tissue to be composed of a number of nearly circular zones, each having a central tube, the Haversian canal, through which the blood circulates. Surrounding the Haversian canal are concentrically arranged laminae, or layers of bone, between which are found irregular spaces called lacunae, containing bone corpuscles and communicating with the Haversian canal and each other by means of canaliculi, through which the nutrition is conveyed to all parts of the bone.

B. Ash, the calcic phosphate remaining after bones have been incinerated. **B. Cartilage**. See *Ostein*. **B.**, Collar, the clavicle. **B.-conduction**, the transmission of sound-waves to the auditory nerve by means of sonorous vibrations communicated to the bones of the skull. **B.-cyst**, a tumor distending and thinning bone, and filled with serum or bloody fluid; in rare cases bone-cysts contain hydatids. **B.**, Epipteric [*ἐπί, upon*; *πτερόν*, a wing], a small Wormian bone sometimes found between the great wing of the sphenoid and the anterior inferior angle of the parietal bone. **B.**, Flat, a bone more or less in the form of a plate. **B.-gelatin**. See *Gelatin*. **B.**, Long, one consisting of a narrow shaft and two expanded ends. **B.**, Rider's, an ossification of the lower tendon of the adductor longus or magnus from pressure. **B.**, Sesamoid.

See *Sesamoid*. **B.**, **Short**, one the three dimensions of which are approximately equal. **B.**, **Wormian**, small bones in the sutures of the skull.

Boneset. See *Eupatorium*.

Bonnet's Capsule. See *Tenon's Capsule*.

Boot, Junod's. See *Junod's Boot*.

Boracic Acid (*bo-ras'-ik as'-id*). See *Acid, Boric*, and *Boron*.

Borate (*bo'-rat*) [*Ar.*, *bûraq*, borax]. Any salt of boric acid.

Borax (*bo'-raks*). See *Boron*.

Borborygmus (*bor-bor-ig'-mus*) [*βορβορυγμός*, a rumbling]. A rumbling of the bowels.

Boric Acid (*bo'-rik as'-id*). See *Boron*.

Borneo Camphor (*bor'-ne-o kam'-for*). See *Borneol*.

Borneol (*bor'-ne-ol*) [*Borneo*], $C_{10}H_{18}O$. Borneo Camphor; a substance that occurs in *Dryobalanops camphora*, a tree growing in Borneo and Sumatra. It is artificially prepared by treating the alcoholic solution of common camphor with sodium.

Boroglycerid (*bo-ro-glis'-er-id*). A mixture of boric acid, 62 parts, with glycerol, 92 parts. It is antiseptic. *Unof.*

Boron (*bo'-ron*) [*Ar.*, *bûraq*, borax]. $B = 11$; quantivalence III, V. A non-metallic element occurring in two allotropic forms—as a powder and as a crystalline substance. It is the base of boric acid and of the mineral borax. **Boracic**, or, more properly, **Boric Acid**, H_3BO_3 , a crystalline substance, found native in the volcanic lagoons of Tuscany. It occurs in white, transparent crystals, soluble in water and alcohol; it is antiseptic. Dose gr. v-xx (0.32-1.3). *Unguent. Acidi Borici* (B. P.), contains hard paraffin 2, soft paraffin 4, boric acid 1. It is used as an antiseptic and in dermatology. **Borax**, $Na_2B_4O_7 \cdot 10H_2O$, sodium tetraborate, occurs as white, transparent crystals, soluble in water, alcohol, and glycerol. It is used as an antiseptic. Dose gr. v-xl (0.32-2.6). *Glycerinum Boracis* (B. P.), contains borax, glycerol, and distilled water; it is used as a local application. *Mel Boracis* (B. P.), borax honey, contains borax, clarified honey, and glycerol (about 1 in 7); it is used as a local application.

Bosselated (*bos'-el-a-ted*) [*ME.*, *bos*]. Covered with bosses or small nodules.

Bot (*bat*) [*botus*, a belly worm]. The larva of certain species of flies of the genus *Æstrus*, which are conveyed into the stomach of man, where they hatch. Also the thread-worm, *Oxyuris vermicularis*.

Botal, or Botallo, Duct of. See *Duct. B., Foramen of*. See *Foramina, Table of*.

Botulismus (*bot-ul-is'-mus*). See *Botulismus*.

Botany (*bot'-an-ē*) [*βοτάνη*, an herb]. The

science of plants—their classification and structure.

Bothriocephalus (*both-re-o-sef'-al-us*) [*Βοθριοψ*, a pit; *κεφαλή*, a head]. A genus of tapeworms. **B. latus**, the fish tapeworm, a common parasite of man in certain European localities. It may reach twenty-five feet in length, with a breadth of three-fourths of an inch. It is also called *Tenia lata* and *Dibothrium latum*.

Botryoid (*bot'-re-oid*) [*Βότρυς*, a cluster; *ειδος*, likeness]. Resembling in shape a bunch of grapes.

Böttcher's Cells. Small cells in the cochlea, occurring in a single layer on the basilar membrane. **B.'s Crystals**, peculiar microscopic crystals formed by adding a drop of a one per cent. solution of phosphate of ammonia to a drop of prostatic secretion. They are composed of a phosphate formed from a base that exists in prostatic fluid, and that is supposed to impart to it its peculiar odor.

Böttger's Test. A test for sugar in the urine. The urine is alkalinized with sodium hydrate, a little bismuth subnitrate is added, and the urine boiled. A black color, due to the reduction of the bismuth, will be produced if sugar is present. Albumin, if present, must previously be removed.

Botulism, Botulismus (*bot'-u-lizm*, *bot-uliz'-mus*) [*botulus*, a sausage]. Sausage-poisoning.

Bougie (*boo-zhe'*) [*Fr.*, a candle]. A slender, cylindric instrument made of waxed silk, catgut, etc., for introduction into the urethra or other passage, for the purpose of dilatation, exploration, etc. **B. à boule**. See *B., Bulbous*. **B., Armed**, a bougie with a piece of silver nitrate or other caustic attached to its extremity. **B., Bulbous**, a bougie with a bulbous tip. **B., Filiform**, a whalebone or other bougie of very small diameter.

Bouillon (*boo-e-yon(g)'*) [*Fr.*]. 1. A broth made by boiling meat, usually beef, in water. 2. A liquid nutritive medium for the culture of microorganisms prepared from finely chopped beef or beef-extract.

Bourdin's Paste. A mixture of nitric acid with flowers of sulphur; it is used as an escharotic.

Bourdonnement (*boor-dun-mon(g)'*) [*Fr.*, *bourdonner*, to buzz]. Any buzzing sound. The murmur that is heard when the stethoscope is applied to any part of the body. It is thought to result from contraction of the muscular fibrille.

Bovine (*bo'-vin*) [*bovinus*, of an ox]. Ox-like. **B. Heart**, the immensely hypertrophied heart of aortic valvular disease.

Bovinine (*bo'-vin-in*) [*bas* (*gen. bovis*), an ox]. A proprietary preparation of beef used as a food for invalids and convalescents.

Bovista (*bo-vis'-tah*) [L.]. 1. A genus of fungi closely allied to Lycoperdon; some of the species are edible. 2. The Lycoperdon bovista, a fungus or puff-ball. When dry it is a good styptic, and its tincture has been used for nervous diseases.

Bowels (*bow'-els*) [OF., *boel*, from L. *botellus*, a sausage]. The intestines.

Bow-leg [ME., *bowe*]. An arching outward of the lower limbs. See *Genu varum*.

Bowman's Capsule. The expanded portion forming the beginning of a uriniferous tubule.

B. Discs, the products of a breaking up of muscle-fibers in the direction of the transverse striations. **B. Glands**, glands found in the olfactory mucous membrane. **B. Membrane**, a thin, homogeneous membrane representing the uppermost layer of the stroma of the cornea, with which it is intimately connected. **B. Muscle**. See *Muscles*, *Table of*. **B. Probe**, a probe used in dilating strictures of the lacrymal duct.

Boyle's Law. See *Law*.

Bozeman's Catheter. A double-current catheter.

Brachia (*bra'-ke-ah*). Plural of *Brachium*.

Brachial (*bra'-ke-al*) [*brachium*, the arm].

Pertaining to the arm. **B. Artery**, the continuation of the axillary artery, extending along the inner side of the arm. See *Arteries*, *Table of*. **B. Glands**, the lymphatic glands of the arm. **B. Plexus**, the plexus of the fifth, sixth, seventh, and eighth cervical and the first dorsal nerves. **B. Veins**, the veins of the arm that accompany the brachial artery.

Brachialgia (*bra-ke-al'-je-ah*) [*βραχίον*, the arm; *ἀλγος*, pain]. Pain or neuralgia in the arm or in the brachial plexus.

Brachinin (*brak'-in-in*) [*βραχίς*, short]. A substance obtained from the bombardier beetle, *Brachinus crepitans*, of Europe. It is said to be efficacious against rheumatism. Unof.

Brachiocephalic (*bra-ke-o-sef-al'-ik*) [*βραχίον*, the arm; *κεφαλή*, head]. Pertaining to the arm and the head. **B. Artery** and **Vein**, the innominate artery and vein.

Brachiocubital (*bra-ke-o-ku'-bit-al*) [*βραχίον*, the arm; *cubitus*, forearm]. Relating to the arm and forearm; as the brachiocubital ligament.

Brachium (*bra'-ke-um*) [*brachium*, *βραχίον*, the arm]. The arm, especially the upper arm. **B. cerebri** or **B. of Optic Lobes**, the bands connecting the nates and testes with the optic thalamus. **B. cerebelli**, the peduncles of the cerebellum. **B. pontis**, the brachium of the pons, being also the middle peduncle of the cerebellum.

Brach-Romberg Symptom. See *Signs and Symptoms*, *Table of*.

Brachycardia (*brak-e-kar'-de-ah*). Same as *Bradycardia*.

Brachycephalic (*brak-e-sef-al'-ik*) [*βραχίς*, short; *κεφαλή*, a head]. 1. Applied to skulls of an egg-like shape, with the larger end behind. 2. Having a skull the transverse diameter of which is more than eight-tenths of the long diameter.

Brachycephalous (*brak-e-sef'-al-us*) [*βραχίς*, short; *κεφαλή*, a head]. Brachycephalic.

Brachymetropia (*brak-e-met-ro'-pe-ah*) [*βραχίς*, short; *μέτρον*, a measure; *ὤψ*, the eye]. See *Myopia*.

Bradycardia (*brad-e-kar'-de-ah*) [*βραδύς*, slow; *καρδία*, the heart]. Slowness of the heart-beat.

Bradyesthesia (*brad-e-es-the'-ze-ah*) [*βραδύς*, slow; *αἴσθησις*, perception]. Dulness of perception.

Bradylalia (*brad-e-la'-le-ah*) [*βραδύς*, slow; *λαλεῖν*, to talk]. A slowness of utterance.

Bradypepsia (*brad-e-pep'-se-ah*) [*βραδύς*, slow; *πέψις*, digestion]. Slow digestion.

Bradyphasia (*brad-e-fa'-ze-ah*) [*βραδύς*, slow; *φάσις*, speech]. Bradylalia.

Bradyphrasia (*brad-e-fra'-ze-ah*) [*βραδύς*, slow; *φράσις*, utterance]. Slowness of speech.

Braidism (*brād'-izm*) [*Braid*, the originator]. The hypnotic state produced by fixation of the eyes upon a shining object; described by James Braid in 1842.

Brain (*brān*) [AS., *bragen*]. That part of the central nervous system contained in the cranial cavity, and consisting of the cerebrum, the cerebellum, the pons, and the medulla oblongata. **B., Abdominal**, the solar plexus. **B., After-.** See *Metencephalon*. **B.-axis**, that portion of the brain-substance including the island of Reil, the basal ganglia, the crura, pons, medulla, and cerebellum. **B.-bladder**, a cerebral vesicle of the embryo. **B. Fever**. See *Meningitis*; also synonym of *Typhoid Fever*. **B., Fore-.** See *Prosencephalon*. **B., Hind-.** See *Opencephalon*. **B., Mid-.** See *Mesencephalon*. **B., Railway-.** a condition analogous to railway spine and characterized by cerebral disturbance. See *Erichsen's Disease*, in *Diseases*, *Table of*. **B. Sand**, a gritty mineral matter found in and about the pineal gland, consisting mainly of calcium and magnesium carbonates and phosphates. **B.-tire**, a condition of brain-exhaustion due to excessive functional activity.

Bran (*bran*) [Breton, *bran*]. The epidermis or outer covering of the seeds of most cereals.

B. Bath. See *Bath*. **B. Dressing**, a dressing formerly used for compound fracture of the leg. The leg was placed in a fracture box and surrounded with clean bran.

Branchiæ (*brang'-ke-e*) [*βράχια*, gills]. The gills of fishes.

Branchial (*brang'-ke-al*) [*βράχια*, gills]. Pertaining to the branchiæ. **B. Arches.** See *Arches*. **B. Openings.** See *Clefts*, *Visceral*.

Branchiogenous (*brang-ke-ōj'en-us*) [*βράχια*, gills; *γεννᾶν*, to produce]. Produced or developed from a branchial cleft.

Brandy (*bran'-de*). See *Spiritus*.

Brassica (*bras'-ik-ah*) [AS., *brassica*, cabbage]. A genus of plants, Nat. Order, *Crucifera*, including the common cabbage.

Brawn (*brawn*) [OF., *braon*, a piece of flesh]. The flesh of a muscle; well-developed muscles.

Brawny (*brawn'-e*) [OF., *braon*, a piece of flesh]. Fleishy; muscular. **B. Induration**, pathologic hardening and thickening of the tissues.

Brayera (*bra-ye'-rah*) [Fr., *brayer*]. Kouso, Cusso, Koso. The female flowers of *B. anthelmintica*, containing tannic acid, a volatile oil, and a crystalline principle, Koussin, $C_{31}H_{38}O_{10}$. It is valuable as an anthelmintic against tapeworm. In large doses it produces nausea and emesis. Dose $\frac{3}{4}$ ij- $\frac{3}{4}$ ss (8.0-16.0), in infusion of boiling water. **Cusso**, Ext., Fld., dose $\frac{3}{4}$ ij-f $\frac{3}{4}$ j (8.0-32.0). **Infusum Cusso** (B. P.). Dose f $\frac{3}{4}$ iv-viii (128.0-256.0).

Bread (*bred*) [AS., *bræd*]. A mixture of flour and water rendered porous by carbon dioxide, and baked. The flour may be of wheat, corn, oat, or rye. The carbon dioxide may be introduced by decomposing an alkaline carbonate (sodium or potassium), by an acid ("cream of tartar"), or by fermenting the starch

with yeast. **B., Brown**, a kind of bread made from a mixture of corn, rye, and wheat-flour. **B., Graham**, made from unbolted wheat-flour; it contains more gluten, diastase, and mineral phosphates than ordinary bread. **B. Poulitice**, a poultice made with bread-crumbs. **B., White**, bread made from bolted wheat-flour, and therefore deficient in diastase, gluten, and mineral phosphates. Other kinds, such as rye (or black), corn, bran, barley, etc., indicate their composition by their name. **B.-paste**, a culture-medium for bacteria and molds. Stale, coarse bread is dried, ground to powder and made into a paste with water.

Break (*brāk*) [AS., *brecan*]. In electricity, to open the circuit of a battery. **B.-bone Fever.** See *Dengue*. **B. Shock**, a term sometimes employed in electrotherapeutics for the physiologic shock produced on the opening or breaking of an electric circuit.

Breast (*bræst*) [ME., *bræst*, the breast]. 1. The anterior part of the chest. 2. The mamma.

B.-bone, the sternum. **B., Broken**, abscess of the mammary gland. **B., Chicken**, *Pectus carinatum*, a deformity marked by prominence of the sternal portion of the chest. **B., Gathered**, mammary abscess. **B. Pang**, angina pectoris. **B., Pigeon**. Same as *B., Chicken*. **B. Pump**, a suction-apparatus for removing the milk from the breast.

Breath (*breth*) [AS., *bræth*]. The air exhaled from the lungs. **B.-sounds**, the respiratory sounds heard upon auscultation. A *Table of Breath-sounds* is given.

TABLE OF BREATH-SOUNDS IN HEALTH AND DISEASE.

(Altered and enlarged from J. K. Fowler.)

VARIETY OF BREATHING.	PERIOD.	PITCH.	QUALITY.	INTERVAL.	DURATION.	INTENSITY.	WHERE HEARD.	CONDITION IN WHICH HEARD.
Vesicular.	1. Inspiration.	Low.	Vesicular.	None.	...	Variable.	Over the lungs, in health.	In health.
	2. Expiration.	Lower.	Blowing.	...	Shorter than inspiration or absent.	Faint or absent.		
Bronchial.	1. Inspiration.	High.	Tracheal.	Distinct.	...	Variable.	In health, in regions of seventh cervical spine. In disease, over areas of consolidation	In health; pulmonary tuberculosis, lobar pneumonia, large pleuritic effusions, thoracic aneurysm, mediastinal tumors.
	2. Expiration.	Higher.	Tracheal.	...	Equal to or longer than inspiration.	Greater.		

TABLE OF BREATH-SOUNDS IN HEALTH AND DISEASE.—*Continued.*

VARIETY OF BREATHING.	PERIOD.	PITCH.	QUALITY.	INTERVAL.	DURATION.	INTENSITY.	WHERE HEARD.	CONDITION IN WHICH HEARD.
Broncho-vesicular.	1. Inspiration.	Higher than in vesicular breathing	More or less tracheal.	Slight.	...	Variable.	In <i>health</i> , ant., over sternal portion of infraclavic. region; <i>post.</i> , upper part of interscapular region. In <i>disease</i> , over slight consolidation.	In health; pneumonia, pulmonary tuberculosis.
	2. Expiration.	Higher than in vesicular breathing	More or less tracheal.	...	About equal to inspiration.	Greater.		
Amporic.	1. Inspiration.	Low.	Hollow and metallic.	Distinct.	...	Variable.	Over a large cavity communicating with an open bronchus.	Pulmonary tuberculosis.
	2. Expiration.	Lower.	Both characters more marked.	...	Longer than inspiration.	Greater.		
Cavernous.	1. Inspiration.	Low.	Blowing and hollow.	Distinct.	...	Variable.	Over a cavity communicating with an open bronchus.	Pulmonary tuberculosis.
	2. Expiration.	Lower.	Both characters more marked.	...	Longer than inspiration.	Greater.		
Tubular.	1. Inspiration.	Higher than in bronchial breathing	Laryngeal or whiffling.	Distinct.	...	Variable.	Over consolidated areas.	Lobar pneumonia, pulmonary tuberculosis.
	2. Expiration.	Higher.	Laryngeal or whiffling.	...	Equal to or longer than inspiration.	Greater.		

Breathing (*bréth'-ing*) [AS., *brath*]. See *Respiration*. **B.**, Abdominal, breathing in which the abdominal walls move decidedly and in which the diaphragm is actively engaged. **B.**, Interrupted, Wavy, or Cog-wheel, a broken or interrupted inspiratory sound produced by nervousness, irregular contraction of the muscles of respiration, or irregular expansion of the lung from disease. **B.**, Mouth, habitual respiration through the mouth. **B.**, Puerile, the breathing normally heard in children, and heard in adults when the respiratory murmur is exaggerated. **B.**, Suppressed, entire absence of breath-sounds, as in pleuritic effusion and certain solid conditions of the lung. **B.**, Thoracic, respiration in which the thoracic walls are actively moved.

Breech (*bréch*) [ME., *breech*]. The buttocks. **B. Presentation**, presentation of the buttocks of the child at the os uteri during labor.

Bregenin (*breg'-en-in*) [Low Ger., *bregen*, brain], $C_{10}H_{17}NO_6$. A name given by Thudichum to a viscous principle, soluble in and crystallizable from alcohol, by means of which it has been extracted from brain-

tissue. It is fusible like a fat, but is miscible with water.

Bregma (*breg'-mah*) [*βρεγμα*, the sinciput]. The part of the skull corresponding to the anterior fontanel.

Bregmatic (*breg-mat'-ik*) [*βρεγμα*, the sinciput]. Relating to the bregma.

Brick-dust Deposit (*brik'-dust de-pos'-it*). A reddish sediment in the urine consisting of urates.

Bricklayers' Itch (*brik'-lá-erz itch*). Eczema due to irritation of lime-mortar.

Brickmakers' Anemia. See *Anchylostomiasis*.

Bridle (*brí'-dl*) [AS., *bridel*]. A band or filament stretching across the lumen of a passage, or from side to side of an ulcer, scar, abscess, etc. **B. Stricture**, a stricture due to the presence of a delicate band stretched across the lumen of the urethra.

Bright's Disease (*bríts dis-ez'*) [*Bright*, an English physician]. A term applied to any disease of the kidney associated with albuminuria. **Acute Bright's Disease**, an acute inflammation of the kidney; it may be parenchymatous, interstitial, or diffuse. **Chronic Bright's Disease**, a chronic inflammation

of the kidney, affecting the parenchyma, the connective tissue, or both. Amyloid degeneration is also considered a chronic form of Bright's disease.

Brim (*brim*) [ME., *brim*]. An edge or margin, as the brim of the pelvis.

Briquet's Ataxia. See *Diseases, Table of*.

Brisement (*bréz-mont-g'*) [Fr.]. A crushing; the forcible breaking up of structures causing ankylosis of a joint.

Bristle-cell [AS., *hyrst*, a bristle]. Any one of the ciliated cells at the terminations of the auditory nerve-filaments.

British (*brit'-ish*) [AS., *Bryttisc*, Britons, the original inhabitants of Britain]. Of or pertaining to Great Britain. **B. Gum.** See *Dextrin*.

Broad (*brad*) [AS., *brád*]. Wide, extensive.

B. Ligament. See *Ligament*. **B. Tape-worm.** See *Bothriocephalus latus*.

Broca's Area. The speech-center in the frontal convolution. **B. Auricular Point,** the center of the external auditory meatus.

B. Convolution, the third left frontal convolution; the center for voluntary speech. **B. Fissure.** See *Fissures, Table of*.

Brodie's Abscess. See *Diseases, Table of*.

B. Disease. Same as *Brodie's Knee*. **B. Joint.** See *Diseases, Table of*. **B. Knee.** See *Diseases, Table of*.

Brokaw Ring (*brok'-aw ring*). A ring used in intestinal anastomosis, made of segments of rubber drainage-tubing and threaded with catgut strands.

Bromal (*bro'-mal*) [*βρωμος*, a stench; *aldehyd*], CBr_3CHO . Tribromaldehyd, analogous to chloral, and produced by the action of bromin on alcohol. It is a colorless, oily fluid, of a penetrating odor, and sharp, burning taste, boiling at $172^\circ\text{--}173^\circ$; it has been used in medicine, having properties similar to those of chloral. **B. Hydrate**, $\text{CBr}_3\text{CHO} + \text{H}_2\text{O}$, a fluid of oily consistence, having a structure similar to that of chloral hydrate, but more irritating and narcotic than the latter. It is used as an hypnotic and in epilepsy. Dose gr. j-v (0.065-0.32). Unof.

Bromamid (*bro'-mam-id*) [*βρωμος*, a stench; *amid*]. A bromin compound of the anilin group with the formula, $\text{C}_6\text{H}_4\text{Br}_2\text{NH}_2\text{HBr}$. It contains 75 per cent. of bromin, and is used as an antipyretic in 10 or 15-grain doses (0.65-1.0).

Bromate (*bro'-mat*) [*βρωμος*, a stench]. A salt of bromic acid.

Bromid (*bro'-mid*) [*βρωμος*, a stench]. A salt of hydrobromic acid; those of calcium, iron, ammonium, potassium, and sodium are used in medicine. They allay nervous excitement, and are employed as sedatives. **B. of Ethyl.** See *Ethyl bromid*, under *Bromin*.

Bromidrosis (*bro'-mid-ro'-sis*) [*βρωμος*, a

stench; *ιδρως*, sweat]. Osmidrosis; an affection of the sweat-glands in which the sweat has an offensive odor.

Bromin, Bromum (*bro'-min*, or *-mum*) [*βρωμος*, a stench]. $\text{Br} = 80$; quantivalence 1. A reddish-brown liquid, which, at ordinary temperatures, gives off a heavy, suffocating vapor. It is a very active escharotic, and internally a violent poison. It is used as an escharotic and disinfectant. The salts of bromin are cerebrospinal and cardiac depressants, and are employed as sedatives, particularly in epilepsy, eclampsia, spasmodic affections, insomnia, hysteria, migraine, etc. The salts of the alkaline metals are those most commonly used. **Ammonii bromidum**, NH_4Br . Dose gr. v-xx (0.32-1.3). **Calcii bromidum**, CaBr_2 . Dose gr. v- $\overline{\text{ss}}$ j (0.32-4.0). **Ethyl bromid**, $\text{C}_2\text{H}_5\text{Br}$, useful in spasmodic coughs. Dose $\overline{\text{m}}$ x- $\overline{\text{ss}}$ j (0.65-4.0). **Ferri bromidi**, *Syr.*, contains ten per cent. of the salt. Dose $\overline{\text{ss}}$ ss-j (2.0-4.0). **Lithii bromidum**, LiBr . Dose gr. v-xx (0.32-1.3). **Nickel bromidum.** See *Nickel*. **Potassii bromidum**, KBr . Dose gr. v- $\overline{\text{ss}}$ j (0.32-4.0). **Sodii bromidum**, NaBr . Dose gr. v- $\overline{\text{ss}}$ j (0.32-4.0). **Strontii bromidum**, SrBr_2 . Dose gr. xv-xx (1.0-1.3). **Zinci bromidum**, ZnBr_2 . Dose gr. ss-ij (0.032-0.13).

Brominism, Bromism (*bro'-min-izm*, *bro'-mizm*) [*βρωμος*, a stench]. Certain peculiar phenomena produced by the prolonged administration of the bromids. The most marked symptoms are headache, coldness of the extremities, feebleness of the heart's action, somnolence, apathy, anesthesia of the soft palate and pharynx, pallor of the skin, and a peculiar eruption of acne which is one of the earliest and most constant symptoms. There is also anorexia and at times loss of sexual power and atrophy of the testes or mammae.

Bromoform (*bro'-mo-form*) [*βρωμος*, a stench; *forma*, form], CHBr_3 . A bromid having a structure like that of chloroform, CHCl_3 ; it is sedative and anesthetic and is used in whooping-cough, and in seasickness, in doses of gr. ij-v (0.13-0.32). Unof.

Bromum. See *Bromin*.

Bronchial (*brong'-ke-al*) [*βρόγχος*, the windpipe]. Relating to the bronchi. **B. Arteries.** See *Arteries, Table of*. **B. Fluke.** See *Distoma ringeri*. **B. Glands,** the chain of lymphatic glands running beside the bronchi.

Bronchiectasis (*brong'-ke-ek'-tas-is*) [*βρόγχος*, the windpipe; *εκτασις*, dilatation]. Dilatation of the walls of the bronchi. It occurs in chronic bronchitis, in fibroid pneumonia, and tuberculosis of the lung. It may involve a tube uniformly, producing the cylindrical form; or it may occur irregularly in sacs

or pockets,—the sacculated form. The characteristic symptom of bronchiectasis is paroxysmal coughing, with the expectoration of large quantities of mucopurulent, often fetid, matter. Cavertous breathing may be heard over the dilated tubes.

Bronchiole (*brong'-ke-ol'*) [dim. of *bronchus*]. One of the smallest subdivisions of the bronchi.

Bronchiolitis (*bron-ke-o-li'-tis*) [*bronchiolus*, a little bronchus; *itis*, inflammation]. Inflammation of the bronchioles. **B.** *exudativa*, an inflammation of the bronchioles, with exudation, a condition by some held to be the cause of bronchial asthma.

Bronchitic (*brong-kil'-ik*) [*βρόγχος*, the windpipe]. Relating to, of the nature of, or affected with, bronchitis.

Bronchitis (*brong-kil'-tis*) [*βρόγχος*, the windpipe; *itis*, inflammation]. Inflammation of the mucous membrane of the bronchial tubes. **B.**, *Acute*, is due to exposure to cold, to the inhalation of irritant vapors, to certain infectious agents, etc. It is characterized by fever, cough, substernal pain, and by dry râles in the early, and moist râles in the later stages. **B.**, *Capillary*, an acute bronchitis of the finer bronchioles; it is generally the result of a downward extension of an acute bronchitis. Children are most frequently affected. Dyspnea, nervous depression, and cyanosis are prominent symptoms. Catarrhal pneumonia is a common complication. **B.**, *Catarrhal*, a form attended with profuse mucopurulent discharges. **B.**, *Chronic*, a form of bronchitis usually occurring in middle or advanced life, characterized by cough and by dry and moist râles. It may be due to repeated attacks of acute bronchitis, to gout, rheumatism, or tuberculosis, or it may be secondary to cardiac and renal disease. **B.**, *Croupous*, *Fibrinous*, or *Plastic*, a rare variety attended with the expectoration of casts of the bronchial tubes, containing Charcot-Leyden crystals and eosinophile cells, after a paroxysm of dyspnea and violent coughing. **B.**, *Dry*, that unattended by expectoration. **B.**, *Mechanic*, or *Potter's*, a form caused by the inhalation of dust, etc. **B.**, *Potter's*. See *B.*, *Mechanic*. **B.**, *Putrid*, or *Fetid*, a variety of chronic bronchitis characterized by the discharge of a copious, half-liquid, extremely offensive sputum. **B.**, *Summer*, hay-fever.

Bronchocele (*brong'-ko-sel*, or *-se'-le*) [*βρόγχος*, the windpipe; *κήλη*, a tumor]. Really a tumor of a bronchus, but generally signifying goiter.

Bronchocephalitis (*brong'-ko-sef'-al-i'-tis*). Synonym of *Whooping-cough*.

Broncholith (*brong'-ko-lith*) [*βρόγχος*, the windpipe; *λίθος*, a stone]. A calculus or concretion formed in a bronchial tube.

Bronchomycosis (*brong-ko-mi-ko'-sis*) [*βρόγχος*, the windpipe; *μύκης*, a fungus]. The growth or presence of fungi in a bronchial tube.

Bronchopathy (*brong-kop'-a-the*) [*βρόγχος*, the windpipe; *πάθος*, disease]. Any disease of a bronchus.

Bronchophony (*brong-koff'-o-ne*) [*βρόγχος*, the windpipe; *φωνή*, the voice]. The resonance of the voice within the bronchi as heard on ausculting the chest. It is normally present over the lower cervical spines, in the upper interscapular region, and over the sternal portion of the infraclavicular regions. The most frequent pathologic cause is consolidation of the lung. **B.**, *Whispered*, bronchophony elicited by causing the patient to whisper.

Bronchoplasty (*brong'-ko-plas-te*) [*βρόγχος*, the trachea; *πλάσσειν*, to form]. The closure of a tracheal or bronchial fistula by operation.

Bronchopneumonia (*brong-ko-nu-mo'-ne-ah*) [*βρόγχος*, the windpipe; *πνεύμων*, the lung]. Lobular pneumonia, a term applied to inflammation of the lungs, which, beginning in the bronchi, finally involves the parenchyma of the lungs. This disease is most frequently encountered in children, but may occur in old age, and may be a simple catarrhal or a tuberculous process.

Bronchorrhagia (*brong-kor-a'-je-ah*) [*βρόγχος*, the bronchus; *ρήγνιναι*, to burst forth]. Hemorrhage from the bronchial tubes.

Bronchorrhea (*brong-kor-e'-ah*) [*βρόγχος*, the bronchus; *ρῑειν*, to flow]. A form of bronchitis attended with profuse expectoration.

Bronchotome (*brong'-ko-töm*) [*βρόγχος*, the windpipe; *τέμνειν*, to cut]. An instrument for cutting the larynx or trachea in the operation of bronchotomy.

Bronchotomy (*brong-kot'-o-me*) [*βρόγχος*, bronchus; *τέμνειν*, to cut]. Incision into the larynx, trachea, or bronchus.

Bronchotracheal (*brong-ko-tra'-ke-al*) [*βρόγχος*, the bronchus; *τραχεΐα*, the windpipe]. Relating to a bronchus (or to both bronchi) and to the trachea.

Bronchovesicular (*brong-ko-ves-ik'-u-lar*) [*βρόγχος*, the bronchus; *vesicula*, a vesicle]. Both bronchial and vesicular. See *Breath-Sounds*, *Table of*.

Bronchus (*brong'-kus*) [*βρόγχος*, bronchus].

One of the primary divisions of the trachea. **Bronzed** (*bronzd'*) [*bronzium*, bronze]. Tanned; of a bronzed color. **B.** *Skin*, a symptom of Addison's Disease. **B.** *skin Disease*. See *Addison's Disease*, *Diseases*, *Table of*.

Broom (*broom*). See *Scoparius*.

Broth (*broth*). See *Bouillon*.

Brow (*brow*) [*AS.*, *brú*]. The forehead;

the upper anterior portion of the head. **B.-Ague**, neuralgia of the first division of the fifth cranial nerve, generally due to malaria. **B.-pang**. Synonym of *Hemiterania*. **B. Presentation**, presentation of the fetal brow in labor.

Brown (*brown*) [A.S., *brún*]. Having a dark color inclining toward red or yellow. **B. Atrophy**, an atrophy of a tissue associated with a deposit therein of a brown or yellow pigment. **B., Bismarck**. See *Bismarck Brown*. **B. Induration of Lung**, a state of the lung due to long-continued congestion, usually arising from valvular heart disease. It is characterized by an increase in connective tissue and an excess of pigment. **B. Mixture**. *Mistura glycyrrhizæ composita*. **B. Ointment**, the unguentum fuscum, N.F.; called also unguentum matris, or "mother's salve." It is composed of "brown plaster" two parts, oil one part, suet one part. **B. Plaster**, the emplastrum fuscum camphoratum, N.F.; called also emplastrum matris camphoratum, or "camphorated mother's plaster;" official in German pharmacy. It is made of red oxid of lead thirty parts, oil of olives sixty parts, yellow wax fifteen parts, camphor one part.

Brownian Movement. An oscillatory movement observed under the microscope in very fine granules, drops, etc., when suspended in a liquid. The movement is not locomotion, and is to be distinguished from that of the self-motility of living microorganisms. Its cause is not definitely known, but it may be due to heat, light, electricity, osmosis, etc.

Brown-Sequard's Disease. See *Diseases, Table of*. **B. Paralysis**. See *Diseases, Table of*.

Brucin (*bru'-sin*) [*Brucia*, a genus of shrubs], $C_{23}H_{26}N_2O_6$. A poisonous alkaloid found in *Strychnos nux vomica* and in *Strychnos ignatiua*. It crystallizes in prisms containing $4H_2O$, and melts at 178° . Its taste is exceedingly bitter and acrid. Its action on the animal economy is similar to but much less powerful than that of strychnin.

Bruit (*bru-e'*) [Fr., a noise or report]. An adventitious sound heard on auscultation. **B., Aneurysmal**. The blowing murmur heard over an aneurysm. **B. d'airain**, the ringing note heard through the stethoscope applied to the chest wall when a coin is struck against another pressed against the surface of the chest on the opposite side. It is pathognomonic of a collection of gas in the pleural cavity. **B. de clapotement**, a splashing sound often heard in cases of well marked dilatation of the stomach when pressure is made upon the abdominal walls. **B. de cuir neuf**, the creaking sound, like that of new leather, sometimes heard in pericarditis. **B. de**

diabie, a humming, rushing sound heard in the veins in anemia. **B. de galop**, a cantering rhythm of the heart sounds, in which owing to a reduplication of the second sound, three sounds are heard. It occurs most frequently in mitral stenosis. **B. de pot fêlé**, The cracked-pot sound. **B. de rappel**, a sound resembling the double beat upon a drum. **B. de soufflet**, the bellows-murmur. **B., Placental**, the uterine souffle, a blowing sound heard over the uterus in pregnancy.

Brunner's Glands. See *Glands*.

Brunonian Movement. See *Brownian Movement*. **B. Theory**, Brownism; a doctrine, taught by Dr. J. Brown (1735-88), that both physiologic and pathologic phenomena are due to variations in a natural stimulus, its excess causing sthenic and its deficiency producing asthenic diseases.

Brush (*brush*) [OF., *brave*]. An instrument consisting of a collection of some flexible material fastened to a handle. In medicine, various forms of brush are employed, as the acid brush, of glass threads; the electric brush, an electrode in the form of a brush; the laryngeal brush; the nasal, pharyngeal, and stomach brush. **B.-burn**, the injury produced by violent friction, and the resulting heat; it often resembles a burn or scald.

Bryce's Test. A test to determine if systemic infection has taken place after vaccination. It consists in repeating the inoculation during the evolution of the vaccine disease. If systemic infection has occurred, the second inoculation will mature rapidly, so as to overtake the first.

Bryonia (*bri'-o-ne-ah*) [*βρυωνία*, bryony]. Bryony. The root of *B. alba* and *B. dioica*; indigenous to Europe. Its properties are due to an intensely bitter glucosid, bryonin, $C_{40}H_{60}O_{19}$, which is a strong irritant when applied to the skin or mucous membrane, often producing vesication. It is used in pleurisy, pleuropneumonia, rheumatic fever, and colds. Dose of the powdered root gr. x-xxx (0.65-2.0). **B., Infusum** (unof.), has a strength of 1 to 16. Dose (3 ss-ij (16.0-64.0). **B., Tinct.**, a 10 per cent. solution of the root in alcohol. Dose f 3 j-ij (4.0-8.0).

Bryonin (*bri'-o-nin*) [*βρυωνία*, bryony]. See *Bryonia*.

Bryony (*bri'-o-ne*). See *Bryonia*.

Bubo (*bu'-bo*) [*βουβών*, the groin]. Inflammation and swelling of a lymphatic gland, particularly of the groin, and usually following chancroid, gonorrhea, or syphilitic infection. **B., Indolent**, one which has no tendency to break down; a syphilitic bubo. **B., Parotid**. See *Parotitis*. **B., Pestilential**, that associated with the plague. **B., Primary**, a slight adenitis of the groin due to

mechanic irritation or other cause; formerly supposed to be due to syphilis without a chancre having preceded. **B.**, Sympathetic, one caused by irritation, friction, injury, etc., and not arising from an infectious disease. **B.**, Syphilitic, that due to syphilis. **B.**, Venereal, that due to venereal disease.

Bubon d'emblée [Fr.]. See *Bubo*, *Primary*.

Bubonic (*bu-bon'-ik*) [*βουβών*, the groin]. Relating to a bubo. **B. Plague**. Synonym of the *Plague*.

Bubonocoele (*bu-bon'-o-sel*) [*βουβών*, the groin; *κήλη*, tumor]. Inguinal hernia when the gut does not extend beyond the inguinal canal.

Buccal (*buk'-al*) [*bucca*, the cheek]. Pertaining to the cheek.

Buccinator (*buk'-sin-a-tor*) [L., a trumpeter]. The thin, flat muscle of the cheek.

Buccula (*buk'-u-lah*) [L., dim. of *bucca*, the cheek]. The fleshy fold seen beneath the chin, and forming what is called a double chin.

Buchu (*bu'-ku*) [native African]. The leaves of several species of *Barosma*, yielding a volatile oil, to which its properties are probably due, and a bitter extractive, barosmin. Dose gr. j-v (0.065-0.32). It causes a sensation of glowing warmth over the body, stimulates the appetite, and increases the circulation. It is useful in cystitis and other affections of the genitourinary mucous membrane. Dose of the leaves, gr. xv-xxx (1.0-2.0). **B.**, Ext., Fld. Dose ℥x-fʒj (0.65-4.0). **B.**, Infusum (unof.), 1-16. Dose fʒss-ij (16.0-64.0). **B.**, Infusum (B. P.). Dose fʒj-iv (32.0-128.0). **B.**, Tinctura (B. P.). Dose fʒj-ij (4.0-8.0).

Buck's Fascia. The superficial perineal fascia.

Buckbean (*buk'-ben*). Bog-bean. The rhizome of *Menyanthes trifoliata*, tonic, antiscorbutic, and emmenagogue. It has been recommended as a vermifuge, and has been used in functional amenorrhea. Dose of fld. ext., ℥v-xxx (0.32-2.0). Unof.

Buckeye Bark (*buk'-i bark*). The bark of *Æsculus glabra*, astringent and tonic, used in rectal irritation, prolapsus, and various uterine derangements. Dose of fld. ext. gtt. ij-v (0.2-0.32). Unof.

Buckthorn (*buk'-thorn*). See *Frangula* **B.**, *California*. See *Cascara sagrada*.

Budding (*bud'-ing*) [ME., *buiden*]. In biology, a form of reproduction or cell division, occurring among the polyps and infusorians, in which a bud is given off by the parent and comes to resemble the latter. The process is also called gemmation.

Buffy Coat (*buff'-e kat*). A grayish or buff-

colored crust or layer sometimes seen upon a blood-clot, after phlebotomy, and once looked upon as a sign of inflammation. It is caused by the partial subsidence of the red blood-corpuscles.

Buhl's Disease. See *Diseases*, *Table of*.

Bulb (*bulb*) [*βολβός*, a bulb]. 1. An oval or circular expansion of a cylinder or tube. 2. The medulla oblongata. **B.** of a *Hair*, the swelling at the root of a hair. **B.**, *Olfactory*, one of the two bulbs of the olfactory nerve situated on either side of the corpus spongiosum penis. **B.** of the *Vagina*, a small body of erectile tissue on each side of the vestibule of the vagina, homologous to the bulb of the urethra of the male. **B.** of *Vena Jugularis*, the dilatation at the termination of the external jugular vein.

Bulbar (*bul'-bar*) [*βολβός*, a bulb]. Bulbous. Pertaining to the medulla. **B.** *Disease*, or *Paralysis*, a term applied to the progressive and symmetric paralysis of the muscles of the mouth, tongue, pharynx, and sometimes those of the larynx. This paralysis is due to a disease of the motor nuclei in the medulla oblongata; an acute and a chronic form are met with. The acute form is due to hemorrhage or softening; the chronic to degeneration. The disease is also called *lateral glossolaryngeal paralysis*. There is also a *pseudobulbar paralysis*, due to symmetric lesions of the motor cerebral cortex.

Bulbi vestibuli (*bul'-bi ves-tib'-u-li*) [L.]. A name sometimes given to the glands of Bartholin.

Bulbocavernosus (*bul-bu kav-ern'-o-sus*) [*βολβός*, a bulb; *caverna*, a cavity]. The accelerator urine muscle, corresponding to the sphincter vagina of the female.

Bulbourethral (*bul-bu ure-th'-al*) [*βολβός*, a bulb; *urethra*, the urethra]. Pertaining to the bulb of the urethra.

Bulbous [*βολβός*, a bulb]. Bulbous. Pertaining to a bulb. **B.** *Urethra*. See *Bartholin*.

Bulesia (*bul'-e-si*) [*βουλή*, a wish]. A wish, or an act of the will.

Bulimia (*bul'-i-mi*) [*βουλή*, a wish; *ιμία*, hunger]. Excessive hunger, especially a voracious desire for food, and is a symptom of emaciation, and the use of certain medicines, as opium.

Bulimus (*bul'-i-mus*) [*βουλή*, a wish; *ιμία*, hunger]. Excessive hunger, especially a voracious desire for food, and is a symptom of emaciation, and the use of certain medicines, as opium.

Bulla (*bul'-la*) [*βουλή*, a wish; *ιμία*, hunger]. Excessive hunger, especially a voracious desire for food, and is a symptom of emaciation, and the use of certain medicines, as opium.

alis, a rounded projection into the middle meatus of the nose, due to an enlarged ethmoid cell. **B. ossea**, the inflated or dilated part of the bony external meatus of the ear.

Bullet Forceps. A forceps for extracting bullets.

Bullous (*bull'-us*) [*bul-la*, a blister]. Marked by bullae; of the nature of a bulla.

Bundle (*bun'-dl*) [AS., *bindan*, to bind]. In biology, a fascicular grouping of elementary tissues, as nerve-fibers or muscle-fibers.

Bunion (*bun'-yun*) [origin uncertain]. A swelling of a bursa of the foot, especially of the great toe.

Bunsen Burner. See *Burner*. **B. Cell**. See *Batteries*, *Table of*.

Buphthalmia (*boof-thall'-me-ah*), or **Buphthalmos** (*boof-thall'-mos*) [*βοῦς*, ox; *ὀφθαλμός*, eye]. See *Keratoglobus*.

Burdach, Column of. The posteroexternal column of the spinal cord.

Burdock (*ber'-dok*). See *Lappa*.

Buret, or **Burette** (*bu-ret'*) [Fr.]. A graduated tube designed for measuring small quantities of a reagent. It is usually held vertically in a stand and is provided with a stopcock.

Burgundy Pitch. See *Pix*.

Burn [ME., *bernen*, to burn]. 1. To become inflamed. 2. To be charred or scorched. 3. To have the sensation of heat. 4. An injury caused by fire or dry heat. 5. A disease in vegetables. 6. In chemistry, to oxygenize. 7. In surgery, to cauterize.

Burner [ME., *bernen*, to burn]. A common name for a lamp or heating apparatus used in laboratories for chemic and pharmaceutic purposes. **B.**, **Argand**, uses gas or oil, and contains an inner tube for supplying the flame with air. **B.**, **Bunsen**, a form in which, before ignition, the gas is mixed with a sufficient quantity of air to produce complete oxidation.

Burns, Ligament of. The falciform process of the fascia lata.

Burrowing (*bur'-o-ing*) [ME., *borugh*, a lurking place]. The term given to the passage of pus through the tissues, after the formation of an abscess.

Bursa (*bur'-sah*) [*bursa*, a purse]. A small sac interposed between parts that move upon one another. **B. Pharyngea**, a blind pouch projecting upward from the pharynx toward the occipital bone. **B.**, **Popliteal**, a bursa

situated in the popliteal space between the tendon of the semimembranosus and the tendon of the inner head of the gastrocnemius, where they rub against each other. **B.**, **Prepatellar**, a bursa situated over the patella and the upper part of the patellar ligament. **B.**, **Synovial**, found between tendons and bony surfaces.

Bursitis (*bur-si'-tis*) [*bursa*, a purse; *itis*, inflammation]. Inflammation of a bursa.

Butter of Cacao. See *Theobroma*.

Butterin (*but'-er-in*) [*butyrum*, butter]. An artificial substitute for butter, made principally of beef-fat.

Buttocks (*but'-uks*) [dim. of *butt*, an end]. The nates. The fleshy part of the body posterior to the hip-joints, formed by the masses of the glutei muscles.

Button (*but'-un*) [ME., *boton*]. See *Furunculus orientalis*. **B.**, **Amboyna**. See *Frambesia*. **B.**, **Belly**, the navel. **B.**, **Biskra**. See *Furunculus orientalis*. **B.-bush**, the bark of *Cephalanthus occidentalis*, a tonic, febrifuge, and diuretic. Dose of fld. ext. f3 ss-j (2.0-4.0). Unof. **B.**, **Corrigan's**, a steel button-shaped cautery-iron, introduced by Sir J. C. Corrigan (1802-80). **B.-hole Mitral**, an advanced degree of constriction of the mitral orifice of the heart. **B.-makers' Chorea**. See *Chorea*. **B.**, **Murphy**, a device used in gastroenterostomy or intestinal anastomosis. **B.-snakeroot**, the root of *Liatris spicata*, and of *Eryngium yuccifolium*; a stimulant, tonic, diuretic, and emmenagogue. Dose of fld. ext. f3 ss-j (2.0-4.0). Unof. **Butyl** (*bu'-til*) [*butyrum*, butter]. C_4H_9 . A hydrocarbon alcohol radicle. **B. Chloral**. See *Chloral butylicum*. **B.-chloral Hydrate**. See *Chloral butylicum*.

Butylamin (*bu'-til'-am-in*) [*butyrum*, butter; *amin*], $C_4H_9NH_2$. A substance contained in cod-liver oil, possessing diuretic and diaphoretic properties.

Butylene (*bu'-til'-en*) [*butyrum*, butter], C_4H_8 . A hydrocarbon belonging to the olefin series. It exists in three isomeric forms, all of which are gases at ordinary temperatures.

Butyrin (*bu'-tir-in*) [*butyrum*, butter], $C_5H_8(C_2H_3O_2)_4$. A constant constituent of butter, together with olein, stearin, and other glycerids. It is a neutral, yellowish, liquid fat, having a sharp, bitter taste.

Butyroid (*bu'-tir'-oid*) [*butyrum*, butter]. **Buttery**; having the consistence of butter.

C

C. 1. The chemic symbol of Carbon. 2. The abbreviation of Centigrade.

Caballine Aloes (*kab'-al-in al'-ās*). An inferior quality of aloes, known also as fetid, or horse aloes.

Cabbage (*kab'-aj*). See *Brassica*. **C. Rose**. See *Rosa centifolia*. **C., Skunk**, a fetid plant of N. America, *Symplocarpus foetidus*. Its tincture and fluid extract are prescribed as antispasmodic and antasthmatic.

Cacao (*kak'-a'-o*). See *Theobroma*. **C. Butter**, oleum theobromæ, is obtained from seeds or nibs of *Theobroma cacao*. It is a pure white fat, with a pleasant odor and taste; it fuses at 86° F. (30° C.); its specific gravity is from .945 to .952. It is used in cosmetics and for pharmaceutical preparations. See also *Theobroma*.

Cachectic (*kak-ek'-tic*) [*κακός*, bad; *ἔξις*, a habit]. Characterized by cachexia.

Cachet (*kash-a'*) [Fr.]. A pharmaceutical preparation consisting of two concave pieces of wafer, varying in size from $\frac{3}{4}$ to $1\frac{1}{4}$ inches in diameter, round or oblong in shape, in one of which the powder to be administered is placed, and the other, having previously been moistened, is then laid over the powder and the two margins are pressed together, when they adhere and completely enclose the powder.

Cachexia (*kak-eks'-e-ah*) [*κακός*, bad; *ἔξις*, a habit]. A depraved condition of general nutrition, due to some serious disease, as syphilis, tuberculosis, carcinoma, etc. **C., Lymphatic**. Synonym of *Hodgkin's Disease*. **C., Pachydermic**. See *Myxedema*. **C. strumipriva**, the condition allied to, if not identical with, myxedema, following the extirpation of the thyroid gland. **C. thyropriva**. See *Myxedema*. **C. virginum**. See *Chlorosis*.

Cacodyl (*kak'-o-dil*). See *Kakodyl*.

Cacoplastic (*kak-o-plas'-tik*) [*κακός*, bad; *πλαστικός*, formed]. Characterized by a low degree of organization.

Cacosmia, or **Kakosmia** (*kak-oz'-me-ah*) [*κακός*, foul; *ὀσμή*, smell]. A disgusting smell.

Cacotrophy (*kak'-ot'-ro-fe*) [*κακός*, bad; *τρέφειν*, to nourish]. Disordered or defective nutrition.

Cactina (*kak-ti'-nah*) [*κάκτος*, a prickly plant]. A proprietary preparation said to be a proximate principle derived from night-blooming cereus (*Cereus grandiflora* and *C. mexicana*). It is a cardiac stimulant, recommended as a substitute for digitalis. Unof.

Cactus Grandiflorus, *Cereus grandiflorus*,

Night-blooming Cereus. The preparations of cactus are stimulant to the spinal cord, the vasomotor center, and the cardiac ganglia. They have been used as substitutes for digitalis. **C. G., Tinct.** Dose $\mathfrak{m}\text{xv}-\text{xx}$ (1.0-1.3). **C. G., Ext., Fld.** Dose $\mathfrak{m}\text{v}-\text{x}$ (0.32-0.65).

Cacumen (*kak-u'-men*) [L.: *πρ.*, *Cacumina*]. 1. The top, as of a plant. 2. The culmen of the vermis superior of the cerebellum.

Cadaver (*kad-ar'-er*) [*cadere*, to fall]. The dead body, especially that of a human being.

Cadaveric (*kad-ar'-er ik*) [*cadere*, to fall]. Pertaining to the cadaver. **C. Alkaloids**, ptomaines. **C. Ecchymoses**, **C. Lividity**, certain postmortem stains, closely resembling in their general appearance the effects of bruises or contusions. They occur on the lowest and most dependent parts of the body. **C. Spasm**, the early, at times instantaneous, appearance of rigor mortis, seen after death from certain causes. It is also called instantaneous rigor and tetanic rigidity.

Cadaverin (*kad-ar'-er in*) [*cadere*, to fall], $\text{C}_2\text{H}_5\text{N}_3$. A ptomaine, occurring very frequently in decomposing animal tissues. It is obtained from human hearts, lungs, livers, etc., after three days' decomposition at ordinary temperature,—also from horseflesh, from putrid mussel, from herring, and haddock. It is a constant product of the growth of the comma-bacillus. It is a thick, clear, syrupy liquid, having an exceedingly unpleasant odor.

Cade (*kad*) [a Languedoc name]. See *Juniperus*. **C., Oil of** (*oleum cadinum*), a tarry oil from the wood of *Juniperus communis*; it is used in the treatment of skin diseases.

Cadmium (*kad'-me-um*) [*cadmia*, calamin]. Cd 112; quantivalence 11. Sp. gr. 8.65-8.69. A bluish-white metal resembling zinc in its general properties. In its physiologic action it is escharotic and astringent; internally, in large doses, it produces emesis and violent gastritis. **C. iodidum**, CdI_2 , used as an ointment, 1 to 8 of lard. **C. sulphas**, $\text{CdSO}_4 \cdot 4\text{H}_2\text{O}$, an astringent in gonorrhea and in corneal opacities; used in a lotion in strength of gr. $\frac{1}{2}$ or 4 to $\frac{3}{4}$ of water, or as an ointment 1 to 40 of fresh lard.

Cæcal (*se' kal*). See *Cæcal*.

Cæcitis (*se-sit' tis*). See *Cæcitis*.

Cæcum (*se' kum*). See *Cæcum*.

Cæsarean Operation (*se-sar'-ee-an*) See *Cæsarean*.

Cæsium (*se'-ze-um*). See *Cæsium*.

Caffea (*kaf'-ee-ah*) [L.] The seed of *C.*

arabica. The dried and roasted seeds are almost universally used in infusion as a beverage, forming a cerebral stimulant and stomachic tonic. They are valuable in promoting digestion and allaying hunger and fatigue. The properties are due to an alkaloid, **Caffein**, $C_8H_{10}N_4O_2 \cdot H_2O$, identical with *Thein*. See *Tea* and *Caffein*. Dose of the alkaloid gr. j-v (0.065-0.32). **C. citrate**. Dose gr. j-v (0.065-0.32). **C. valerianae**, for nervous vomiting in hysteria. Dose gr. j-iv (0.065-0.26). **Ext. Caffæ Viridis** Fld., is intended as a substitute for the Fluid Extract of Guarana. Unof. Dose fʒss-ij (2.0-8.0). See *Guarana*. **Injectio Caf. hypodermatica**, one grain of caffein in three minims. Dose mʒj-vj (0.065-0.39).

Caffeic Acid (*kaf'e-ik*), $C_8H_6O_4$. A crystalline acid substance found in coffee.

Caffein (*kaf'e-in*) [*caffæa*, coffee], $C_8H_{10}N_4O_2 \cdot H_2O$. An alkaloid found in the leaves and beans of the coffee-tree, in tea, in Paraguay tea, and in guarana, the roasted pulp of the fruit of *Paulinia sorbilis*. It occurs in long, silky needles, slightly soluble in cold water and alcohol, with a feebly bitter taste. It is a cerebriospinal, circulatory, and renal stimulant. See *Caffæa*. **C. citrate**. Dose gr. j-v (0.065-0.32). **Sodium and Caffein benzoate**, and **Sodium and Caffein salicylate** are used hypodermically. Dose gr. iij-v (0.2-0.32).

Caffeinism (*kaf'e-in-izm*) [*caffæa*, coffee]. Chronic coffee-poisoning; a train of morbid symptoms due to excess in the use of coffee.

Caisson Disease (*ka'son diz-ez*). Diver's disease or tunnel disease, a morbid condition due to increased atmospheric pressure, sometimes occurring in divers, caisson-workers, etc. Paraplegia, hemiplegia, anesthesia, or apoplectic attacks are common, coming on only after return to the normal atmosphere. The nature of the lesion is obscure.

Cajuput Oil (*kaj'e-put oil*) [Malay, "white wood"; "aleum, oil"]. A volatile oil distilled from the leaves of *Leukadendron cajuputi*. It resembles oil of turpentine. It is used in flatulent colic, hysteria, cutaneous disorders, and toothache. Dose mʒj-v (0.065-0.32). **C., Spiritus** (B.P.), contains 2 per cent. of the oil. Dose fʒss-j (2.0-4.0).

Cajuputol (*kaj'u-putol*) [Malay, *Cajuputi*, "white wood"]. The more limpid part of oil of cajuput; it is found also in some other fragrant volatile oils.

Calabar Bean (*kal'-ab-ar ben*). See *Physostigma*.

Calabarin (*kal'-ab'-ar-in*) [*Calabar*]. An alkaloid from Calabar bean.

Calamin (*kal'-am-in*) [*calamina*, a corruption of *cadmia*]. Native zinc carbonate. *Calamina preparata*, the prepared calamin,

washed and pulverized, is used mainly as an external exsiccant and astringent. *Unguentum calaminæ*, Turner's cerate. Unof.

Calamus (*kal'-am-us*) [L., a reed]. Sweet flag. The rhizome of *Acorus calamus*. It contains a volatile oil and acorin, a bitter nitrogenous principle. The root is an aromatic, stomachic tonic, and a common ingredient of many popular "bitters." **C., Draco**, a species of rattan-palm that affords a part of the so-called Dragon's Blood of commerce. **C., Ext., Fld.** Dose mʒxv-fʒj (1.0-4.0). **C. scriptorius** (a writing pen or reed). The groove on the floor of the fourth ventricle, at the end of which is the ventricle of Arantius.

Calcaneum (*kal'-ka'-ne-um*) [L., the heel]. The os calcis, or heel-bone.

Calcareous (*kal'-ka'-re-us*) [*calx*, limestone]. Pertaining to or having the nature of limestone. **C. Infiltration**. See *Infiltration*.

Calcarine (*kal'-kar-en*) [*calcar*, a spur]. Spur-shaped; relating to the hippocampus minor. **C. Fissure**. See *Fissure*.

Calcic (*kal'-sik*) [*calx*, lime]. Of or pertaining to lime.

Calcicosis (*kal'-sik-a'-sis*) [*calx*, lime]. Marble-cutter's phthisis; a chronic inflammation of the lung due to the inhalation of marble-dust.

Calcification (*kal'-sif-ik-a'-shun*) [*calx*, lime; *fiere*, to become]. The deposit of calcareous matter within the tissues of the body. **C., Metastatic**, that resulting from an excess of lime-salts in the blood, as occurs in the rapid breaking down of bones from osteomalacia.

Calcine (*kal'-sin* or *kal'-sin'*) [*calcinare*, to calcine]. To separate the inorganic elements of a substance by subjecting it to an intense heat.

Calcium (*kal'-se-um*) [*calx*, lime]. Ca = 40; quantivalence 11. A brilliant, silver-white metal, the basis of limestone, characterized by strong affinity for oxygen, and isolated with great difficulty. It is best known in the form of *calcium oxid*, quicklime; *C. hydrate*, slaked lime; and *C. carbonate*, limestone or chalk. **C. benzoas**, $Ca(C_7H_5O_2)_2$, used in nephritis and albuminuria of pregnancy. Dose gr. v-x (0.32-0.65). **C. bromidum**, used as a sedative. Dose gr. xx-xxx (1.3-2.0). **C. chloridum**, $CaCl_2$, soluble in water; used internally to increase the coagulability of the blood. Dose gr. x-xx (0.65-1.3). **C. chlorid**, Liq., (B.P.). Dose mʒxv-l (1.0-3.2). **Calcis, Liquor**, lime-water, contains about $1\frac{1}{2}$ parts of lime in 1000 of water. Dose fʒss-ij (16.0-64.0). **Calcis, Liq., Saccharatus** (B.P.). Dose mʒxv-fʒj (1.0-4.0). **Calcis, Syr.**, saccharated syrup of lime, contains 5 per cent. lime, 30 per cent. sugar, 65 per cent. water. It is the antidote

to poisoning by phenol or oxalic acid. Dose 3 ss-ij (2.0-8.0). **Calx**, CaO, calcium oxid, quicklime.

Calculus (*kal'-ku-lus*) [*calculus*, a stone]. Of the nature of a calculus.

Calculus (*kal'-ku-lus*) [dim. of *calx*, chalk]. A calcareous or stone-like concretion found in the body, particularly in cavities. **C.**, **Arthritic**, a gouty concretion. **C.**, **Biliary**, a gall-stone. **C.**, **Bronchial**, a concretion in an air-passage. **C.**, **Cutaneous**. See *Milium*. **C.**, **Dental**, tartar on the teeth or gums. **C.**, **Fusible**, a urinary calculus composed of phosphates of ammonium, calcium, and magnesium. **C.**, **Lacteal**, or **Mammary**, a calcareous nodule sometimes obstructing the lactiferous ducts. **C.**, **Mulberry**, the oxalate-of-lime variety, resembling a mulberry in shape and color. **C.**, **Nasal**, one found in the nasal cavities. **C.**, **Prostatic**, one in the prostate gland. **C.**, **Renal**, a calculus found in the kidney. **C.**, **Salivary**, one formed in the ducts of the salivary glands. **C.**, **Uterine**, an intrauterine concretion; a womb-stone; formed mainly by calcareous degeneration of a tumor. **C.**, **Vesical**, a calculus found in the urinary bladder.

Calendula (*kal-en'-du-lah*) [*calendæ*, the first day of the month]. Marigold. The flowering plant known as the garden-marigold. *C. officinalis*. **C.**, **Tinct.**, contains 20 per cent. of the leaves and stems. It is used as a local application to wounds, bruises, and ulcers, and has been vaunted as a cure for carcinoma.

Calendulin (*kal-en'-du-lin*) [*calendæ*, the first day of the month]. An amorphous principle obtainable from calendula.

Calenture (*kal'-en-tur*) [Sp., *calentura*, heat; L., *calere*, to be hot]. 1. A tropical remittent fever with delirium; formerly, a supposed fever of this kind that attacked mariners, leading them to leap into the sea. 2. Sunstroke.

Calf (*kalf*) [Ice., *kálf*]. The thick, fleshy part of the back of the leg, formed by the gastrocnemius and soleus muscles.

California Buckthorn. See *Cassia torrida*.

Calipers (*kál'-ip-ers*) [contraction of *calipers*]. Compasses with curved legs.

Calisaya (*kál'-i-say*) [Sp., *América*]. Cinchona bark, especially that of *Maritima calisaya*. See *Cinchona*.

Calisthenics, or **Calisthenics**. An artificial skin [*calist*, beautiful; *sthen*, strength]. The practice of various rhythmic movements of the body, intended to develop the muscles and produce gracefulness of carriage. Light gymnastics, especially designed for the use of girls and young women.

Calix (*kál'-iks*) [L.]. A cup, especially the

of the cup-like divisions of the pelvis of the kidney into which the pyramids project.

Callosal (*kal'-o'-sal*) [*callosus*, hard]. Pertaining to the corpus callosum.

Callositas (*kal'-o'-sit-as*). See *Callosity*.

Callosity (*kal'-o'-sit-e*) [*callus*, hardness]. **Callositas**, Tylosis, Tyloma, Keratoma. A hard, thickened patch on the skin produced by excessive accumulation of the horny layers.

Callosomarginal (*kal'-o'-so-mar'-jin-al*) [*callosus*, hard; *margo*, margin]. Relating to the callosal and marginal gyri of the brain.

Callosum (*kal'-o'-sum*). Same as *Corpus callosum*.

Callosus (*kal'-us*) [*callosus*, hard]. Hard; tough, like callus.

Callus (*kal'-us*) [L.]. 1. A callosity; hardened and thickened skin. 2. The new growth of incomplete osseous tissue that surrounds the ends of a fractured bone during the process of repair. **C.**, **Permanent**, the permanent bond of bony union after reabsorption of the **C. Provisional**, or cartilage-like, plastic material first thrown out. **C. of Skin**, induration and thickening of the skin.

Calmative (*kál'-ma-tiv*) [F., *calme*, still]. Calming; sedative.

Calomel (*kal'-o-mel*) [*καλός*, fair; *μύλος*, black]. See *Hydrargyrum*.

Calor (*kal'-lor*) [L.]. 1. Heat. 2. Moderate fever-heat; less than *fever* and *ardor*. **C. animalis**, animal heat. **C. febrilis**, fever-heat. **C. fervens**, boiling heat. **C. innatus**, natural or normal heat. **C. internus**, inward fever; fever not appreciable on the surface of the body. **C. mordax**, **C. mordicans**, biting or pungent heat. **C. natus**, native or animal heat; blood-heat; normal heat.

Calorie (*kal'-lor-ee*) [Fr.] See *Calory*.

Calorific (*kal'-or-if-ic*) [*calor*, heat; *facere*, to make]. Heat-producing; applied to certain foods.

Calorimeter (*kal'-or-im-eter*, *Calor*, heat; *metron*, a measure). An instrument for measuring the amount of heat that bodies produce or absorb.

Calorimetry (*kal'-or-im-et-ree*, *Calor*, heat; *metron*, a measure). The measure of the heat given off by a substance.

Calory (*kal'-or-ee*) [*calor*, heat; *metron*, a measure]. The amount of heat required to raise the temperature of one gram of water one degree Fahrenheit.

Calvaria (*kal'-varee*) [*calvaria*, skull; *calva*, empty]. The skull of a human being, especially the frontal bone, which is the largest and most prominent part of the skull. It is composed of the frontal, parietal, and occipital bones, and is covered by the scalp. The skull is the protective covering of the brain, and is the most important part of the human skeleton.

ful in atonic dyspepsia, and as a mild, appetizing tonic in convalescence. **C., Ext.** (B. P.). Dose gr. ij-x (0.13-0.65). **C., Ext. Fld.** Dose \mathfrak{m} v-xxx (0.32-2.0). **C., Infus.** (B. P.). Dose $\mathfrak{f}\overline{\text{ss}}$ -ij (32.0-64.0). **C., Tinct.**, contains ten per cent. of C. Dose $\mathfrak{f}\overline{\text{ss}}$ -ij (2.0-8.0).

Calvaria, Calvarium (*kal'-va'-re-ah, kal'-va'-re-um*) [*calva*, the scalp]. The upper part of the skull.

Calvities (*kal'-vish'-e-ēz*) [*calvus*, bald]. Baldness.

Calx (*kalks*) [L.]. 1. The heel. 2. The oxid of calcium, CaO. See *Calcium*. **C. chlorata** (U. S. P.), **C. chlorinata** (B. P.), chlorinated lime. **C. sulphurata** (U. S. P., B. P.), consists largely of calcium sulphid; used externally and internally in skin diseases. Dose $\frac{1}{8}$ grain (0.006).

Cambodia, Cambogia (*kam-bo'-je-ah*) [*Camboja*, or *Cambodia*, in Siam]. Gamboge. A resinous gum from *Carcinia hanburii*, a tree native to Southern Asia. Its properties are due to gambogic acid. It is a drastic, hydragogue cathartic, decidedly diuretic. **C., Pil. Comp.** (B. P.), contains gamboge, Barbadoes aloes, hard soap, compound powder of cinnamon, and syrup. Dose gr. v-x (0.32-0.65). It is also officially a constituent of **Pil. Cathartic. Comp.** See *Colocynth*. Dose gr. ij-v (0.13-0.32).

Camera (*kam'-er-ah*) [*καμάρα*, an arched roof or chamber]. A box or chamber. In optics, the chamber or dark-box of the apparatus used for photography. **C. lucida**, an optic device for superimposing or combining two fields of view in one eye, invented by the chemist, Wollaston.

Camisole (*kam-is-ol'*) [Fr.]. The straight-jacket, formerly used for the restraint of violently insane persons.

Camomile, or Chamomile (*kam'-o-mil*). See *Anthemis* and *Matricaria*.

Camp Fever. Synonym of *Typhus Fever*.

Camphenes (*kam'-fenz*) [*campher*]. The volatile oils or hydrocarbons having the general formula $C_{10}H_{16}$, isomeric with oil of turpentine. Many camphenes exist ready-formed in plants, as oil of cloves, etc. They are liquid at ordinary temperatures.

Campho-phenique (*kam-fo-fen-ēk'*) [Fr.]. A proprietary preparation combining camphor and phenol; it is recommended as an antiseptic and local stimulant. Unof.

Camphor (*kam'-for*) [*camphora*, camphor], $C_{10}H_{16}O$. A solid, volatile oil obtained from *Cinnamomum camphora*, a tree indigenous to Eastern Asia. It yields camphoric and camphonic acids, also *C. Cymol* when exposed to a high heat in close vessels. It is antispasmodic, anodyne, diaphoretic, and stimulant. Applied locally, it is rubefacient. It is

used in cholera, vomiting, the typhoid state, headache, diarrhea with pain, etc., cardiac depression, and affections requiring an antispasmodic. **C., Aqua**, consists of camphor 8, alcohol 5, distilled water sufficient to make 1000 parts. Dose $\mathfrak{f}\overline{\text{ss}}$ -iv (4.0-16.0). **C. Ball**, an English preparation used as an application to chapped skin. Its composition is spermaceti 4, white wax 12, oil of almonds 5; melt in a water bath, and add flowers of camphor 4. Dissolve and when nearly cold pour into boxes or mould in gallipots. **C., Borneo.** See *Borneol*. **C., Carbolated**, a mixture of 2½ parts of camphor with one each of phenol and alcohol; it is a good antiseptic dressing for wounds. **C., Ceratum**, consists of camphor liniment 3, olive oil 12, simple cerate 85. It is used for itching skin-affections. **C., Chloral**, a fluid prepared by mixing equal parts of camphor and chloral. It is an excellent solvent for many alkaloids, and is used externally as a sedative application. **C.-ice**, the *ceratum camphora compositum*, N. F. **C., Linimentum**, has camphor 20, cottonseed oil 80 parts. **C., Liniment. Comp.** (B. P.), contains camphor and oil of lavender dissolved in rectified spirit, and strong solution of ammonia added. **C., Monobromated**, $C_{10}H_{15}BrO$, camphor in which one atom of hydrogen has been replaced by an atom of bromine. It resembles the bromids in therapeutic action. Dose gr. j-x (0.065-0.65). **C. salicylate**, prepared by heating together 14 parts of camphor with 11 of salicylic acid. It is used as an ointment. **C., Spt.**, contains camphor 10, alcohol 70, water 20 parts. Dose \mathfrak{m} v-xx (0.32-1.3). **C., Tinct., Comp.** (B. P.), contains opium, benzoic acid, camphor, oil of anise, and proof spirit. Dose \mathfrak{m} xv- $\mathfrak{f}\overline{\text{ss}}$ (1.0-4.0). **C., Tinct., Rubini's**. Unof. A saturated solution of camphor in alcohol. Dose \mathfrak{m} ij-v (0.13-0.32).

Camphorated (*cam'-for-a-ted*) [*camphora*, camphor]. Containing camphor.

Camphoric Acid (*kam-for'-ik*). See *Acid*.

Campimeter (*kam-pim'-et-er*). See *Perimeter*.

Canada (*kan'-a-dah*) [Sp.]. A British possession in North America. **C. Balsam**, the oleoresin of *Abies balsamea* used as a mounting medium in microscopy. **C. Hemp**. See *Apocynum*.

Canadol (*kan'-ad-el*). A transparent volatile liquid resembling benzene in smell. It is a local anesthetic used in minor surgical operations. Unof.

Canal, or Canalis (*kan-al'*, or *kan-at'-lis*) [*cannalis*]. A tubular channel or passage. **C., Abdominal**. See *C., Inguinal*. **C., Alcock's**, a strong sheath of the obturator layer of the pelvic fascia containing the in-

ternal pudic artery. **C., Alimentary**, the whole digestive tube from the mouth to the anus. **C., Alisphenoid**, in comparative anatomy, a canal in the alisphenoid bone, opening anteriorly into the foramen rotundum, and transmitting the external carotid artery. **C., Alveolar, Anterior**, one located in the superior maxilla; it transmits the anterior superior dental nerve. **C., Alveolar, Inferior**, the inferior dental canal. **C., Alveolar, Median**, one located in the superior maxilla and transmitting the middle superior dental nerve. **C., Alveolar, Posterior**, one situated in the superior maxilla; it transmits the posterior superior dental nerve. **C., Alveolodental**. See *C., Dental*. **C., Arachnoid**, a space formed beneath the arachnoid membrane of the brain; it transmits the *venæ magnæ Galeni*. **C. of Arantius**, the ductus venosus. **C., Archinephric**, the duct of the archinephron or primitive kidney. **C., Arterial**. See *Ductus arteriosus*. **C., Atrial**, the cavity of the atrium. **C., Auditory, External**, that from the auricle to the tympanic membrane. **C., Auditory, Internal**, that beginning on the posterior surface of the petrous bone, and extending outward and backward for a distance of about four lines; it transmits the auditory and facial nerves, and the auditory artery. **C., Auricular**. 1. See *C., Auditory, External*. 2. The constriction between the auricular and ventricular portions of the fetal heart. **C., Avant**, the anterior portion of the male urethra. **C., Bartholin's**, the duct of Bartholin's gland. **C., Bernard's**, a supplementary duct of the pancreas. Also called Santorini's canal. **C., Bichat's**. See *C., Arachnoid*. **C., Biliary**. See *C., Hepatic*. **C., Blastoporic**. See *C., Neurenteric*. **C. of Bone**, a canaliculus of bone. **C., Braun's**. See *C., Neurenteric*. **C., Braune's**, the continuous passage formed by the uterine cavity and the vagina during labor, after full dilatation of the os uteri. **C., Breschet's**. See *C. of the Diploe*. **C., Bullular**. See *C. of Petit*. **C., Caroticotympanic**, two or three short canals extending from the carotid canal to the tympanum; they transmit branches of the carotid plexus. **C., Carotid**, one in the petrous portion of the temporal bone; it transmits the internal carotid artery. **C. of Cartilage**, the canals in ossifying cartilage, during its vascularization intended to receive prolongations of the osteogenetic layer of the periosteum. They radiate in all directions from the center of ossification. **C., Central (of the modiolus)**, a canal running from the base to the apex of the cochlea. **C., Central (of spinal cord)**, the small canal that extends through

the center of the spinal cord from the *conus medullaris* to the lower part of the fourth ventricle. It represents the embryonic ectodermal canal. **C., Cerebrospinal**. 1. The neural or craniovertebral canal formed by the skull and the spine, and containing the brain and spinal marrow. 2. The primitive continuous cavity of the brain and spinal cord, not infrequently more or less extensively obliterated in the latter, but in the former modified in the form of the several ventricles and other cavities. **C., Cervical**. See *C. of Cervix Uteri*. **C., Cervicouterine**. See *C., Uterine*. **C. of Cervix Uteri**, that portion of the uterine canal that extends between the internal and external os. **C. of Chorda Tympani**, a small canal in the temporal bone, between its squamous and petrous portions, parallel with the Glaserian fissure; it transmits the chorda tympani nerve. **C., Ciliary**. See *C. of Fontana*. **C. of Cloquet**. See *C., Hyaloid*. **C., Cochlear**, the spiral and snail-like cavity of the cochlea, 28 to 30 mm. long. The base is turned inward toward the internal auditory meatus, and the apex outward toward the tympanum. **C., Connecting**, the arched or coiled portion of a uriniferous tubule, joining with a collecting tubule. **C. of Corti**, the triangular canal formed by the pillars of Corti, the base of which corresponds to the *membrana basilaris*. It extends over the entire length of the *lamina spiralis*. **C. of Cotunnus**. See *Aqueduct of Cotunnus*. **C., Cranio-vertebral**. See *C., Cerebrospinal*, and *C., Neural*. **C., Crural**. See *C., Femoral*. **C. of Cuvier**, the ductus venosus. **C., Cystic**, the cystic duct. **C., Deferent**, the *vas deferens*. **C., Demicircular**. See *C., Semicircular*. **C., Dental, Anterior**, one extending into the facial portion of the superior maxilla; it transmits the anterior dental vessels and nerves. **C., Dental, Inferior**, the dental canal of the inferior maxilla; it transmits the inferior dental nerve and vessels. **C., Dental, Posterior**, two canals in the superior maxilla. They transmit the superior posterior dental vessels and nerves. **C., Dentinal**, the minute canals in dentine, extending approximately at right angles to the surface of a tooth from the pulp-cavity, into which they open, to the cementum and enamel. **C., Digestive**. See *C., Alimentary*. **C. of Diploe**, canals in the diploe of the cranium transmitting Breschet's veins. **C., Ejaculatory**. See *Duct, Ejaculatory*. **C. of Epididymis**, a convoluted tube, about 20 feet long when straightened, forming the epididymis and continuous with the *vas deferens*. **C., Ethmoidal, Anterior**, one between the ethmoid and frontal bones; it transmits the nasal branch of the ophthalmic nerve, and the an-

terior ethmoidal vessels. **C., Ethmoidal, Posterior.** See *C., Orbital, Posterior, Internal.* **C., Eustachian,** one in the petrous portion of the temporal bone, containing a portion of the Eustachian tube. **C., Facial,** the aqueduct of Fallopius; it transmits the facial nerve. **C., Fallopiian.** See *C. Facial.* **C., Femoral.** 1. The inner compartment of the sheath of the femoral vessels behind Poupert's ligament. 2. See *C., Hunter's.* **C. of Ferrein,** a triangular channel, supposed to exist between the free edges of the eyelids when they are closed, and to serve for conducting the tears toward the puncta lacrymalia during sleep. **C. of Fontana,** a series of small spaces formed by the interlacing of the connective-tissue fibers of the framework of the peripheral processes of the iris, situated in the angle of the anterior chamber, and serving as a medium for the transudation of the aqueous humor from the posterior to the anterior chamber of the eye. They are also called *C. of Hovius, C., Ciliary,* and *Fontana's Spaces.* **C., Galactophorous,** the lactiferous tubules of the mammary gland. **C. of Gärtner,** in the female, the remains of the main portion of the Wolffian duct of the embryo; it is a tube extending transversely along the broad ligament, and is the homologue of the vas deferens. **C., Genital,** in comparative anatomy, any canal designed for copulation or for the discharge of ova. **C. of Guidi.** See *C., Vidian.* **C. of Havers.** See *C., Haversian.* **C., Haversian,** canals in the compact substance of bone forming a wide-meshed network, and establishing communication between the medullary cavity and the surface of the bone. Their average diameter is $\frac{5}{16}$ inch, and they are lined by a continuation of the endosteum; they transmit blood-vessels, lymph-vessels, and nerves. **C., Hemal,** the ventral of the two canals of which, according to R. Owen, the vertebrate animal is composed. It contains the heart and the other viscera, while the neural canal encloses the central nervous system. **C. of Henle,** a portion of the uriniferous tubules. **C., Hepatic.** 1. The excretory duct of the liver. 2. The radicles of the hepatic duct. **C., Hernial,** one transmitting a hernia. **C., of Hovius.** See *C. of Fontana.* **C. of Huguier.** See *C. of the Chorda Tympani.* **C., Hunter's,** a triangular canal formed in the adductor magnus muscle of the thigh; it transmits the femoral artery and vein and internal saphenous nerve. **C. of Huschke,** one formed by the junction of the tubercles of the annulus tympanicus. This is generally obliterated after the fifth year, but may persist through life. **C., Hyaloid,** a canal running antero-posteriorly through the vitreous body, through which in the fetus the hyaloid artery passes,

to ramify on the posterior surface of the crystalline lens. **C., Incisor,** a canal that opens into the mouth by an aperture just behind the incisor teeth of the upper jaw; it is formed by a groove on the adjoining surfaces of the superior maxillæ, and has two branches that open into the nasal fossæ. **C., Infraorbital,** a small canal running obliquely through the bony floor of the orbit; it transmits the infraorbital artery and nerve. **C., Inguinal,** a canal about one and one-half inches long, running obliquely downward and inward from the internal to the external abdominal ring, and constituting the channel through which an inguinal hernia descends; it transmits the spermatic cord in the male, and the round ligament of the uterus in the female. **C., Intestinal,** that portion of the alimentary canal that is included between the pylorus and the anus. **C., Intralobular, Biliary,** the radicles of the bile-ducts, forming a fine network in and around the hepatic cells, and communicating with vacuoles in the cells. **C. of Jacobson.** See *C., Tympanic.* **C. of Kowalevsky.** See *C., Neuroenteric.* **C., Lacrymal.** 1. The bony canal that lodges the nasal duct. 2. One of the lacrymal canaliculi. **C. of Loewenberg,** that portion of the cochlear canal that is situated above the membrane of Corti. **C., Malar,** one in the malar bone transmitting the malar division of the temporomalar branch of the superior maxillary nerve. **C., Maxillary.** See *C., Dental.* **C., Median.** 1. The central canal of the spinal cord. 2. The aqueduct of Sylvius. **C., Medullary.** 1. The hollow cavity of a long bone, containing the marrow. 2. See *C., Vertebral.* 3. The central canal of the spinal cord. 4. An Haversian canal. 5. In embryology, the medullary tube. **C., Medullary, Cerebrospinal,** the central canal of the spinal cord. **C., Membranous, of the Cochlea,** a canal in the cochlea, following the turns of the lamina spiralis; it is bounded by the basilar membrane, the membrane of Reissner, and the wall of the cochlea. **C., Membranous, Semicircular.** See *C., Semicircular.* **C. of Modiolus,** the spiral canal of the cochlea. **C., Myelonal,** the central canal of the spinal cord. **C., Nasal.** 1. See *C., Lacrymal.* 2. An occasional canal found in the posterior portion of the nasal bone; it transmits the nasal nerves. **C., Nasolacrymal.** See *C., Lacrymal.* **C., Nasopalatine.** See *C., Incisor.* **C., Neural.** See *C., Vertebral.* **C., Neuroenteric,** of Kowalevsky, also called blastoporic canal, in the embryo, a passage leading from the posterior part of the medullary tube into the archenteron. **C., Neurocentral.** See *C., Vertebral.* **C. of Nuck,** a pouch of peritoneum which, in the female fetus, de-

scends for a short distance along the round ligament of the uterus into the inguinal canal; it is the analogue of the processus vaginalis in the male. It sometimes persists after birth. **C., Nutritive.** See *C., Haversian.* **C., Obstetric.** See *C., Parturient.* **C., Obturator,** a canal in the ilium transmitting the obturator nerve and vessels. **C., Olfactory,** in the embryo, the nasal fossæ at an early period of development. **C., Omphalomesenteric,** in the embryo, a canal that connects the cavity of the intestine with the umbilical vesicle. **C., Orbital, Anterior Internal.** See *C., Ethmoidal, Anterior.* **C., Orbital, Posterior Internal,** the posterior of two canals formed by the ethmoid bone and the orbital plate of the frontal bone. It transmits the posterior ethmoidal vessels. **C., Palatine, Accessory Posterior,** one or two canals in the horizontal plate of the palate bone, near the groove entering into the formation of the posterior palatine canal. **C., Palatine, Anterior,** formed by the union of the incisive canals; it opens on the palate behind the incisor teeth. **C., Palatine, Descending.** See *C., Palatomaxillary.* **C., Palatine, External, Small,** a small canal in the pyramidal process of the palate bone, close to its connection with the horizontal plate. It transmits the external palatine nerve. **C., Palatine, Posterior.** See *C., Palatomaxillary.* **C., Palatine, Smaller.** See *C., Palatine, Posterior.* **C., Palatine, Superior,** one formed by the palate bone and the superior maxilla, transmitting the large palatine nerve and blood-vessels. **C., Palatomaxillary,** one formed by the outer surface of the palate bone and the adjoining surface of the superior maxilla. It transmits the large palatine nerve and blood-vessel. **C., Parturient,** the channel through which the fetus passes in parturition, comprising the cavity formed by the uterus and vagina considered as a single canal. **C., Pelvic,** the canal of the pelvis, from the superior to the inferior strait. **C., Perivascular,** the lymph spaces about the blood-vessels. **C. of Petit,** a space, intersected by numerous fine interlacing fibers, existing between the anterior and posterior laminae of the suspensory ligament of the crystalline lens. It extends from the periphery of the lens, nearly to the apices of the ciliary processes, and transmits the secretion from the posterior chamber. **C., Petromastoid,** a small canal, not always present, situated at the angle of union between the mastoid and petrous bones. It transmits a small vein from the middle fossa of the skull to the transverse sinus. **C., Petrosal,** two canals on the upper surface of the petrous portion of the temporal bone, transmitting the large and small superficial petrosal nerves. **C., Plas-**

matic, an Haversian canal. **C., Pore.** See *C., Porous.* **C., Porous,** a canal in the ovule, supposed to serve for the entrance of the spermatozooids in fecundation. **C., Portal,** the space in the capsule of Glisson of the liver, in which the portal vein, hepatic artery, and bile-duct lie. **C., Primitive,** the neural canal of the embryo. **C., Pseudostomatous,** the processes of branched cells that extend from a subepithelial or endothelial layer to the free surface, their free ends forming the pseudostomata. **C., Pterygoid.** See *C., Vidian.* **C., Pterygopalatine,** one formed by the root of the internal pterygoid plate of the sphenoid bone and the sphenoidal process of the palate bone. It transmits the pterygopalatine vessels and nerve. **C., Pulmoaortic.** See *Ductus arteriosus.* **C. of Recklinghausen,** minute channels supposed to exist in all connective tissue, which are directly continuous with the lymphatic vessels, and hence may be said to form their origin. **C. of Reissner.** See *C., Membranous, of the Cochlea.* **C. of Rivinus,** the duct of the sublingual gland. **C. of Rosenthal.** See *C., Spiral, of the Modiolus.* **C., Sacculocochlear,** one connecting the sacculus and the cochlea. **C., Sacculoutricular,** one connecting the sacculus and the utricle. **C., Sacral,** the continuation of the vertebral canal in the sacrum. **C. of Schlemm,** an irregular space or plexiform series of spaces occupying the sclerocorneal region of the eye; it is regarded by some as a venous sinus, by others as a lymph-channel. **C., Semicircular,** bony canals of the labyrinth of the internal ear. They are three in number, the external, superior, and posterior, and contain the membranous semicircular canals. **C., Semicircular, Anterior.** See *C., Semicircular, Superior.* **C., Semicircular, Anterior Vertical.** See *C., Semicircular, Superior.* **C., Semicircular, External,** that one of the semicircular canals of the labyrinth having its plane horizontal and its convexity directed backward. **C., Semicircular, Frontal.** See *C., Semicircular, Superior.* **C., Semicircular, Horizontal.** See *C., Semicircular, External.* **C., Semicircular, Inferior.** See *C., Semicircular, Posterior.* **C., Semicircular, Inner.** See *C., Semicircular, Posterior.* **C., Semicircular, Internal.** See *C., Semicircular, Posterior.* **C., Semicircular, Lateral.** See *C., Semicircular, External.* **C., Semicircular, Osseous.** See *C., Semicircular.* **C., Semicircular, Posterior,** that one of the semicircular canals having its convexity directed backward, and its plane almost parallel to the posterior wall of the pyramid. **C., Semicircular, Posterior Vertical.** See

C., Semicircular, Posterior. *C.*, Semicircular, Sagittal. See *C.*, Semicircular, Posterior. *C.*, Semicircular, Superior, that one of the semicircular canals having its convexity directed toward the upper surface of the pyramid. *C.*, Seminal, the seminiferous tubules. *C.*, Serous, any minute canal connected with the lymph-vessels and supposed to be filled with lymph. *C.*, Sheathing, the communication between the cavity of the tunica vaginalis of the testicle and the general peritoneal cavity. It soon closes in man, leaving the tunica vaginalis a closed sac. *C.*, Spermatic. 1. The vas deferens. 2. The inguinal canal in the male. *C.*, Sphenopalatine. See *C.*, Pterygopalatine. *C.*, Spinal. See *C.*, Vertebral. *C.*, Spiral, of the Cochlea, one that runs spirally around the modiolus, taking two turns and a half, diminishing in size from the base to the apex, and terminating in the cupola. *C.*, Spiral, of the Modiolus, a small canal winding around the modiolus at the base of the lamina spiralis. *C.*, Spiroid, of the Temporal Bone. See *C.*, Fallopiian. *C.* of Steno, the duct of the parotid gland. *C.* of Stilling. See *C.*, Hyaloid, and *C.*, Central, of Spinal Cord. *C.*, Suborbital. See *C.*, Infraorbital. *C.*, Supraorbital, one at the upper margin of the orbit. It transmits the supraorbital artery and nerve. *C.*, Temporal. See *C.*, Zygomaticotemporal. *C.*, Temporomalar. See *C.*, Zygomaticotemporal. *C.*, Thoracic, the thoracic duct. *C.*, Tympanic, one that opens on the lower surface of the petrous bone, between the carotid canal and the groove for the internal jugular vein. It transmits Jacobson's nerve. *C.*, Uterine, the cavity of the uterus, including the body and neck. *C.*, Uterocervical, the cavity of the cervix uteri. *C.*, Uterovaginal. 1. The common canal formed by the uterus and vagina. 2. In embryology, the duct of Müller. *C.*, Vaginal, the canal of the vagina. *C.*, Vascular. See *C.*, Haversian. *C.*, Vector, the oviduct. *C.*, Venous, the ductus venosus. *C.*, Vertebral, the canal formed by the vertebrae. It contains the spinal cord and its membranes. *C.*, Vidian, a canal of the sphenoid bone at the base of the internal pterygoid plate, opening anteriorly into the sphenomaxillary fossa, and posteriorly into the foramen lacerum. It transmits the vidian nerve and vessels. *C.*, Vulvar, the vestibule of the vagina. *C.*, Vulvouterine, the vagina. *C.*, Vulvovaginal. 1. The vagina and the vulva considered as a single canal. 2. The orifice of the hymen. *C.* of Wirsung, the pancreatic duct. *C.* of Wolff. See *Wolffian Duct*. *C.*, Zygomatic. See *C.*, Zygomaticotemporal. *C.*, Zygomaticofacial. See *C.*, Malar. *C.*, Zygomatic-

cotemporal, the temporal canal of the malar bone, running from its orbital to its temporal surface. It transmits a branch of the superior maxillary nerve.

Canalicular (*kan-al-ik'-u-lar*) [*canalis*, a channel]. Canal-shaped.

Canaliculus (*kan-al-ik'-u-lus*) [*L.*]. 1. A small canal; especially that leading from the punctum to the lacrymal sac of the eye. 2. Any one of the minute canals opening into the lacunæ of bone.

Canalization (*kan-al-iz-a'-shun*) [*canalis*, a canal]. 1. The formation of canals, as in tissues, etc. 2. A system of wound-drainage without tubes.

Cancellous (*kan'-sel-us*) [*cancelli*, lattice-work]. Resembling lattice-work, as the tissue in the articular ends of long bones.

Cancer (*kan'-ser*). See *Carcinoma*. *C.*

aquaticus. Synonym of *Stomatitis*, *Gangrenous*. *C.*-bandage, a crab-shaped bandage; a split cloth of eight tails. *C.*-cell, an epithelial cell of peculiar distorted shape, found in the interior of cancer-nests. It is an ordinary epithelial cell altered in outline by pressure. *C.*, Clay-pipe. See *C.*, *Smokers'*. *C.*, Colloid, one containing colloid material. *C.*, Encephaloid. See *C.*, *Soft*. *C.* en cuirasse, disseminated cancer of the skin of the thorax. *C.*, Hard, one containing an excess of fibrous tissue. *C.*-juice, the milky fluid yielded by the cut surface of a cancer on scraping. *C.*, Medullary. See *C.*, *Soft*. *C.*, Melanotic, a pigmented form. *C.*, Scirrhus. See *C.*, *Hard*. *C.*, *Smokers'*, epithelioma of the lip due to the irritation of a pipe. *C.*, *Soft*, one in which the cells predominate, the connective tissue being very small in amount.

Cancerin (*kan'-ser-in*) [*cancer*, a crab]. The name given to a ptomain obtained from the urine in cases of carcinoma of the uterus. It is a white substance crystallizing in fine needles and soluble in alkaline solutions. Its formula is $C_8H_5NO_3$.

Cancerous (*kan'-ser-us*) [*cancer*, a crab]. Having the qualities of a cancer; malignant.

Canceromyces (*kan'-ser-o-mi'-ses*). See *Cladosporium cancerogenes*.

Cancroid (*kan'-kroid*) [*cancer*, a crab]. Cancer-like. *C.* Corpuscles, the pearly bodies of squamous epithelioma. *C.* Ulcer. See *Rodent Ulcer*.

Cancroin (*kan'-kro-in*) [*cancer*, a crab]. A substance (said to be identical with neurin), introduced by Adamkiewicz as a material for hypodermic injection in cases of malignant disease, it being regarded by him as an alexin destructive of cancer-tissue.

Cancrum (*kan'-kru-m*) [*can-ser*, a crab]. A cancer or rapidly-spreading ulcer. *C.* oris, canker of the mouth, gangrenous stomatitis,

noma, gangrenous ulceration of the mouth. It is a disease of childhood between the ages of one and five, characterized by the formation of foul, deep ulcers of the buccal surfaces of the cheeks or lips. There is but slight pain, but the prostration is great, and death usually results from exhaustion or blood-poisoning. The disease is bacterial, poor hygienic surroundings and a debilitated system being predisposing causes.

Cane Sugar. See *Saccharum*.

Canella (*kan-el'-ah*) [dim. of *Canna*, a reed]. The bark of *C. alba* deprived of its corky layer and dried. It is a native of the West Indies and is an aromatic tonic and bitter stomachic. Dose of the powdered bark, gr. xv-xxx (1.0-2.0). It is official in the B. P.

Canine (*ka'-nin*) [*canis*, a dog]. Partaking of the nature of, relating to, or resembling a dog, or the sharp tearing-teeth of mammals, located between the incisors and the molars. **C. Appetite**, bulimia. **C. Eminence**, a prominence on the outer side of the upper maxillary bone. **C. Fossa**. See *Fossa*. **C. Laugh**, a sardonic smile or grin. **C. Madness**, rabies, hydrophobia. **C. Muscle**, the levator anguli oris. **C. Teeth**, dentes canini; cynodontes; dentes lanarii; dents angulaires; cuspidati; conoides; eye-teeth. The cuspid teeth next to the lateral incisors; so called from their resemblance to a dog's teeth.

Canities (*kan-ish'-e-ty*) [Lat.]. Poliosis; hoariness; blanching of the hair.

Canker (*kan'-ker*) [*cancer*, a crab]. An ulceration of the mouth, or any ulcerous or gangrenous sore; cancrum oris; in farriery, a fetid abscess of the horse's foot. See *Cancrum oris*. **C.-rash**. Synonym of *Scarlet Fever*.

Canna (*kan'-ah*) [*kávva*, a cane]. A genus of large-leaved marantaceous plants. **C. indica**, has an acrid and stimulant root; it is alterative, diuretic, and diaphoretic. Unof. The rhizome of *C. speciosa* affords canna-starch, a substitute for arrow-root.

Cannabene (*kan'-ab-ēn*). See *Cannabis*.

Cannabin. See *Cannabis*.

Cannabinin (*kan'-ab'-in-in*) [*cannabis*, hemp]. A volatile alkaloid from cannabis indica.

Cannabinon, Cannabinone (*kan'-ab'-in-on*) [*cannabis*, hemp]. An amorphous bitter resinoid from Indian hemp, used as an hypnotic. Dose 1 to 3 grains (0.065-0.2). Unof.

Cannabis (*kan'-ab-is*) [L.]. Hemp. Indian hemp. The flowering tops of *C. sativa*, of which there are two varieties, *C. indica* and *C. americana*, the former being the more potent. They contain a resin, cannabin, and a volatile oil, from which are obtained cannabene, $C_{15}H_{22}$, a light hydrocarbon, and hybrid of cannabene, a crystalline body.

It is antispasmodic, narcotic, and aphrodisiac. In large doses it produces mental exaltation, intoxication, and a sensation of double consciousness. It is used in migraine, in paralysis agitans, in spasm of the bladder, in sexual impotence, in whooping-cough, in asthma, and in other spasmodic affections. *Bang*, *Bhang*, *Gunjah*, *Churrus*, and *Hashish* are the various Indian names by which the drug is known. **C. indica**, Ext., dose gr. $\frac{1}{6}$ -j (0.01-0.065). **C. ind.**, Ext., Fld., an alcoholic preparation. Dose m-j-v (0.065-0.32). **C. ind.**, Tinct., contains 20 per cent. of the drug. Dose m-xx-f 3j (1.3-4.0). **Cannabin Tannate**. Dose as an hypnotic gr. v-x (0.32-0.65).

Cannula (*kan'-u-lah*) [dim. of *canna*, a tube]. A tube used for withdrawing fluids from the body. It is generally fitted with a pointed rod for puncturing the integument.

Canquoin's Paste. A paste of flour, water, and zinc chlorid; it is a powerful escharotic.

Cantani's Diet. An exclusive meat-diet in diabetes.

Cantering Rhythm (*kan'-ter-ing rithm*). See *Bruit de Galop*.

Canthal (*kan'-thal*) [*καὶθός*, a canthus]. Relating to a canthus.

Cantharidal (*kan'-thar'-id-al*) [*καὶθαρίς*, a blistering fly]. Relating to or containing cantharides.

Cantharides (*kan'-thar'-id-ēz*). Plural of *Cantharis*.

Cantharidin (*kan'-thar'-id-in*) [*καὶθαρίς*, a blistering fly], $C_{10}H_{12}O_4$. The bitter principle contained in Spanish flies and other insects; it crystallizes in prisms or leaflets, and melts at 218° . It has an extremely bitter taste, and produces blisters on the skin. See *Cantharis*.

Cantharis (*kan'-thar-is*) [*καὶθαρίς*, a blistering fly; *pl.*, *Cantharides*]. Spanish fly. The dried body of a species of beetle, *C. vesicatoria* (nat. ord. *Coleoptera*). It contains a powerful poisonous principle, cantharidin, $C_{10}H_{12}O_4$. Locally applied, cantharis is a rubefacient and vesicant; internally it is an irritant, causing pain and vomiting. In toxic doses it produces severe gastroenteritis, strangury, and priapism. It is used as an external counterirritant in the form of "blisters." Internally it is employed as a stimulant to the genitourinary mucous membrane, especially in cystitis, atony of the bladder, amenorrhea, etc.; also in skin diseases. **C., Acetum** (U. P.), of the strength of 1 to 8. **C., Ceratum**, cantharides 35, yellow wax 20, resin 20, lard 35 parts, alcohol q. s. **C., Ext.**, **Ceratum**, cantharides 30, resin 15, yellow wax 15, lard 35, alcohol q. s. **C., Charta**, cantharides 1, Canada turpentine 1, olive oil 4, spermaceti 3, white wax 2 atq, spread on

paper. *Charta epispastica* (B. P.), blistering-paper (white-wax, spermaceti, olive oil, resin, Canada balsam, cantharides, distilled water). **C. cum collodio**, cantharides 60, flexible collodion 85, chloroform q. s. **Emplastrum picis cum cantharide**, warming plaster, Burgundy pitch 92, cerat. canthar. 8 parts. **Emplastrum calefaciens** (B. P.), cantharides, expressed oil of nutmeg, yellow wax, resin, resin-plaster, soap-plaster, boiling water. **C., Emplastrum** (B. P.), cantharides, yellow wax, prepared suet, prepared lard, resin. **C., Liniment.**, cantharides 15 parts, oil of turpentine q. s. ad 100. **C., Tinct.**, contains 5 per cent. of the drug. Dose 3-10 drops (0.09-0.3). **C., Unguent.** (B. P.), cantharides, yellow wax, olive oil.

Canthectomy (*kan-thek'to-me*) [*κανθός*, canthus; *ἐκτομή*, a cutting out]. Excision of a canthus.

Canthitis (*kan-thi'tis*) [*κανθός*, canthus; *itis*, inflammation]. Inflammation of a canthus.

Canthoplasty (*kan'tho-plas-te*) [*κανθός*, canthus; *πλαστική*, to form]. An operation for increasing the size of the palpebral fissure by cutting the outer canthus.

Canthorraphy (*kan-thor'a-fe*) [*κανθός*, canthus; *ράφή*, a seam]. An operation to reduce the size of the palpebral fissure by suture of the canthus.

Canthotomy (*kan-thot'o-mé*) [*κανθός*, canthus; *τομή*, a cutting]. Surgical division of a canthus.

Canthus (*kan'thus*) [*κανθός*, canthus]. The angle formed by the junction of the eyelids.

Canula (*kan'-u-lah*). See *Cannula*.

Caoutchouc (*kao'-chuk*) [S. American]. India-rubber. The chief substance contained in the milky juice that exudes upon incision of a number of tropic trees belonging to the natural orders *Euphorbiaceae*, *Artocarpaceae*, and *Apocynaceae*. The juice is a vegetable emulsion, the caoutchouc being suspended in it in the form of minute transparent globules. When pure, caoutchouc is nearly white, soft, elastic, and glutinous; it swells up in water without dissolving; the best solvents are carbon disulphid and chloroform. It melts at about 150° C., and decomposes at 200° C.

Capacity (*kap-as'-it-é*) [*capacitas*, capacity]. The power of holding or containing; mental or physical ability. **C., Vital**, the total amount of air that can be expelled by the most forcible expiration after the deepest inspiration.

Capillarity (*kap-il-ar'-it-é*) [*capillus*, a hair]. Capillary attraction: the force that causes fluids to rise in fine tubes or bores.

Capillary (*kap'il-a-ré*) [*capillus*, a hair]. 1. Hair-like: relating to a hair, or to a hair-like

filament, or to a tube with a hair-like bore.

2. A minute blood-vessel connecting the smallest ramifications of the arteries with those of the veins. **C. Attraction**. See *Capillarity*. **C. Bronchitis**. See *Bronchitis*. **C. Fissure**, **C. Fracture**, a linear fracture, without displacement. **C. Nevus**. See *Nevus vascularis*. **C. Pulse**, pulsation of the capillaries sometimes seen in aortic regurgitation. **C. Vessels**, the capillaries. **Capillaries**, Meigs's, branching capillaries discovered by A. V. Meigs in the human heart.

Capital (*kap'-it-al*) [*caput*, the head]. 1. Pertaining to the head, or to the summit of a body or object. 2. Of great importance, as a capital operation in surgery.

Capitellum (*kap-it-el'-um*) [dim. of *caput*]. The rounded, external surface of the lower end of the humerus.

Capitulum (*kap-it'-u-lum*) [*capitulum*, a small head]. A little head. **C. of Santorini**, a small elevation on the apex of the arytenoid cartilage corresponding in position to the posterior extremity of the vocal band.

Capric Acid (*kap'-rik as'id*). See *Acid*.

Caprin (*kap'-rin*) [*capra*, a goat]. An oily and flavoring constituent of butter; glycerol caprate.

Caproic Acid (*ka-pro'-ik as'id*). See *Acid*. **Caprone** (*kap'-ron*) [*capra*, a goat], $C_{11}H_{22}O_2$. Caproic ketone; a clear, volatile oil found in butter, and forming the larger part of the oil of rûe.

Caproylamin (*kap-ro-il'-am-in*) [*capra*, a goat; *amin*], $C_6H_{13}N$. Hexylamin. A ptomain formed in the putrefaction of yeast.

Caprylic Acid (*kap-ri'l'-ic as'id*). See *Acid*. **Capsicin** (*kap'-sis-in*), $C_9H_{11}O_3$. The active principle of Cayenne pepper. It is a thick, yellowish-red substance.

Capsicol (*kap'-sik-al*) [*capsicum*; *oleum*, oil]. A red oil obtainable from the oleoresin of capsicum.

Capsicum (*kap'-sik-um*) [*capsa*, a box]. Cayenne pepper. The fruit of *C. fastigiatum*, native to tropic Africa and America. Its odor and hot taste are due to a volatile oil, capsin, $C_9H_{11}O_3$, which is irritant to the skin and mucous membranes. Internally, it is a stomachic, tonic, diuretic, and aphrodisiac. It is useful in atonic dyspepsia, flatulent colic, and in intermittent fever. **C., Emplastrum**, prepared from the oleoresin and resin plaster. **C., Ext., Fld.** Dose $\mathfrak{m}\jmath$ -f 5j (0.32-4.0). **C., Linimentum**, 1 in 10, for chest-affections, rheumatism, etc. **C., Oleoresin, Ethereal**. Dose $\mathfrak{m}\jmath$ -v (0.065-0.32). **C., Tinct.**, contains 5 per cent. of capsicum. Dose $\mathfrak{m}\jmath$ -f 5j (0.65-4.0).

Capsitis (*kap-si'tis*). Same as *Capsulitis*.

Capsular (*kap'-su-lar*) [*capsula*, a small

box]. Pertaining to a capsule. **C. Cataract**, an opacity of the capsule of the crystalline lens. **C. Hemiplegia**, a hemiplegia due to a lesion in the internal capsule.

Capsule (*kaf'-sül*) [dim. of *capsa*, a chest]. A receptacle or bag. In pharmacy, a small, spheroidal shell composed of glycerol and gelatin, divided so that the parts fit together like a box and cover. It is used for the administration of nauseous medicines. **C.**, **Atrabiliary**. See **C.**, **Suprarenal**. **C.**, **Bonnet's**, the posterior portion of the sheath of the eyeball. **C.**, **Bowman's**, the covering of the tuft of vessels of a renal glomerulus. It is the beginning of the uriniferous tubule. **C.**, **External**, a layer of white nerve-fibers forming part of the external boundary of the lenticular nucleus. **C.** of **Gliason**, the connective-tissue sheath of the hepatic artery, portal vein, and bile-duct. **C.**, **Internal**, a layer of nerve-fibers on the outer side of the optic thalamus and caudate nucleus, which it separates from the lenticular nucleus, and containing the continuation upward of the crus cerebri. **C.** of the **Lens**, a transparent, structureless membrane enclosing the lens of the eye. **C.**, **Malpighian**, the commencement of the uriniferous tubules. See **C.**, **Bowman's**. **C.**, **Mueller's**. See **C.**, **Bowman's**. **C.**, **Suprarenal**, the ductless, glandular body at the apex of each kidney. **C.** of **Tenon**, the tunica vaginalis of the eye.

Capsulitis (*kaf-su-lit'-tis*) [*capsula*, a small box; *itis*, inflammation]. Inflammation of the capsule of the lens, or of the fibrous capsule of the eyeball.

Capsulolenticular (*kaf-su-lo-len-tik'-u-lar*) [*capsula*, a little box; *lenticula*, a lentil]. Relating to the lens and to its capsule.

Capsulotome (*kaf'-su-lo-töm*). See **Cystotome**.

Capsulotomy (*kaf-su-lot'-o-me*) [*capsula*, a capsule; *teuein*, to cut]. The operation of rupturing the capsule of the crystalline lens in cataract-operations.

Captation (*kaf-tis'-shun*) [*captare*, to desire]. The first or opening stage of the hypnotic trance.

Capuron, **Cardinal Points** of. See **Pelvis**.

Caput (*kaf'-ut*) [L.]. The head; also the chief part or beginning of an organ. **C. coli**, the head of the colon. **C. cornu posterioris**. Same as **C. gelatinosum**. **C. galinaginis**. See **Verumontanum**. **C. gelatinosum**, the name given to the translucent gray matter covering the dorsomesal periphery of the dorsal horn of the spinal cord. It is a peculiar, striated substance composed of numerous closely crowded cellular elements, in part connective-tissue cells, in part nerve-cells. **C. medusæ**, the peculiar plexus of

veins surrounding the umbilicus in periportal cirrhosis of the liver. It represents collateral paths for the return of the venous blood from the abdominal viscera. **C. obstipum**. Synonym of *Wry-neck*. **C. quadratum**, the rectangular head of rickets, flattened upon the top and at the sides, with projecting occiput and prominent frontal bosses. **C. succedaneum**, a tumor composed of a serosanguineous infiltration of the connective tissue situated upon the presenting part of the fetus.

Caramel (*kar'-am-el*) [Fr., burnt sugar]. Cane-sugar deprived of two molecules of water. It is a viscid, brown-colored liquid.

Caraghen (*ka'-ra-ghen*). Irish moss.

Caraway (*kar'-ah-wa*). See **Carum**.

Carbamic Acid (*kar-bam'-ik as'-id*). See **Acid**.

Carbamid (*kar'-bam-id*) [*carbo*, a coal; *amid*]. $\text{CH}_3\text{N}_2\text{O}$. Urea.

Carbinol (*kar'-bin-el*) [*carbo*, a coal]. Methyl alcohol, CH_3OH . Also a generic term for the alcohols formed by substituting hydrocarbon radicles for the hydrogen in the methyl radicle of carbinol.

Carbazotic Acid (*kar-baz-ot'-ik*). See **Acid**, **Picric**.

Carbo (*kar'-bo*) [L.]. A coal; charcoal.

C. animalis, animal-charcoal; bone-black; it is used in pharmacy and in manufacturing chemistry largely as a decolorizing agent and as a filter. **C. ligni**, wood-charcoal; an absorbent, disinfectant, and deodorizer, used in poulticing wounds and dressing ulcers. It is used internally in gastrointestinal irritation.

Carbohydrate (*kar-bo-hi'-drät*) [*carbo*, coal; *hydr*, water]. An organic substance containing six carbon atoms or some multiple of six, and hydrogen and oxygen in the proportion in which they form water; that is, twice as many hydrogen as oxygen atoms. The carbohydrates form a large class of organic compounds, and may be arranged into three groups: the glucoses (*monoses*); the disaccharids, or sugars; and the polysaccharids. The glucoses are the aldehyd derivatives or ketone-derivatives of the hexahydric alcohols, into which they may be converted by the absorption of two hydrogen atoms. They are mostly crystalline substances, very soluble in water, but dissolving with difficulty in alcohol. They possess a sweet taste. The disaccharids and polysaccharids are ethereal anhydrids of the glucoses. All of them may be converted into the glucoses by hydrolytic decomposition. The disaccharids are ether-like anhydrids of the hexoses.

Carbolic (*kar-bol'-ik*) [*carbo*, a coal; *chem*, oil]. Containing or derived from coal tar oil. **C. Acid**. See **Acid**, **Carbolic**.

Carbolize (*kar'-bol-iz*) [*carbo*, a coal; *chem*, oil]. To impregnate with carbolic acid.

Carbon (*kar'-bon'*) [*carbo*, charcoal]. Charcoal. $C \equiv 12$; quantivalence 11, IV. A nonmetallic element occurring in the various forms of diamond, graphite or "black lead," charcoal, and lamp-black. It is the central or characteristic element of organic compounds.

Carbo animalis purificat., purified animal charcoal. Dose gr. xx-3j (1.3-4.0). **C.**, **Cataplasma** (B. P.), charcoal-poultice; wood-charcoal 1, bread-crumbs 1, linseed-meal 3, boiling water 20 parts. **C. dioxid**, the acid, gaseous product, having the composition of CO_2 , commonly known as "carbonic acid gas," or carbonic acid. It is a colorless gas, having a sp. gr. of 1.52, soluble in cold water, and possessing a pungent smell and an acid taste. Inhaled, it destroys animal life by asphyxiation. **C. disulphid**, bisulphid of carbon, CS_2 , a colorless, transparent liquid of offensive odor, highly inflammable, very poisonous. It is used as a solvent for caoutchouc and as a reagent. **C. monoxid**, CO, carbonic oxid, a colorless, tasteless, and inodorous gas, one of the products of imperfect combustion. It is actively poisonous. **C. nitrid**, CN, cyanogen. **C. tetrachlorid**, CCl_4 , anesthetic, used in asthma by inhalation. Unof. **C. oxysulphid**, a body, COS, formed by conducting sulphur-vapor and carbon monoxid through red-hot tubes; it is a colorless gas, with a faint and peculiar odor. It unites readily with air, forming an explosive mixture, and is soluble in an equal volume of water. It is present in the waters of some mineral springs.

Carbonate [*carbo*, a coal]. A salt of carbon dioxid.

Carbonemia (*kar-bon-e'-me-ah*) [*carbon*, charcoal; *aima*, the blood]. An accumulation of carbon dioxid in the blood.

Carbonic Acid (*kar-bon'-ik as'-id*). See *Carbon dioxid*.

Carbonization (*kar-bon-iz-a'-shun*) [*carbo*, coal]. The process of decomposing organic substances by heat without air, until the volatile products are driven off and the carbon remains.

Carbonyl (*kar'-bon-il*) [*carbo*, charcoal]. A hypothetical organic radicle having the formula CO.

Carboxyhemoglobin (*kar-boks-y-hem-o-glo'-bin*) [*carbo*, coal; *ōxēs*, sharp; *aima*, blood]. The compound of carbon monoxid and hemoglobin, formed when CO is present in the blood. The carbon monoxid displaces the oxygen and checks the respiratory function of the red corpuscles.

Carboxyl (*kar-boks'-il*) [*carbo*, charcoal; *ōxēs*, sharp]. The group, CO.OH, characteristic of the organic acids. The hydrogen of this can be replaced by metals, forming salts.

Carbuncle (*kar'-bung-kli*) [*carbo*, a live coal]. A hard, circumscribed, deep seated, painful, suppurative inflammation of the subcutaneous tissue. It differs from a boil in having greater size, a flat top, and several points of suppuration. It is erroneously called anthrax.

Carcinoma (*kar'-sin-o'-mah*) [*καρκίνωμα*; *καρκίνος*, a crab; *ōma*, tumor]. A malignant epithelial tumor composed of a connective-tissue stroma, surrounding groups or nests of epithelial cells. Three varieties are generally described—the squamous, the cylindric, and the glandular. **C.**, **Colloid**, affects chiefly the alimentary canal, uterus, etc.; the delicate connective-tissue stroma is filled with colloid matter, the result of a colloid degeneration of the epithelial cells. In some cases the degeneration is mucoid, instead of colloid. **C.**, **Cylindric**, one in which the cells tend to assume a cylindric or columnar shape. This shape is best seen in the cells nearest the periphery of the nests. **C.**, **Encephaloid**, is of rapid growth, with a small amount of stroma, large alveoli, and greater amount of cells and blood-vessels. **C.**, **Glandular**, a carcinoma in which the cells are of the glandular or secreting type. **C.**, **Squamous**, one derived from squamous epithelium; the cells are cuboidal in shape.

Carcinomatous (*kar-sin-o'-mat-us*) [*καρκίνος*, a crab]. Relating to or affected with carcinoma.

Carcinosis (*kar-sin-o'-sis*) [*καρκίνος*, a crab]. 1. A carcinomatous cachexia; a tendency to the development of malignant disease. 2. A form of carcinoma, usually fatal, beginning generally in the uterus or the stomach and spreading to the peritoneum. **C.**, **Acute**, rapidly fatal carcinosis. **C.**, **Miliary**, one in which there are many secondary nodules the size of miliary tubercles.

Cardamomum (*kar-dam-o'-mum*) [L.]. *Cardamom*. The fruit of *C. ellettaria*, cultivated in Malabar. Its properties are due to a volatile oil, $C_{10}H_{16}$. It is an aromatic, carminative stomachic, used as an ingredient of several "bitters." It is useful when combined with purgatives to prevent griping. **C.**, **Infusum**. Unof. Dose f3j (64.0). **C.**, **Tinct.**, 15 per cent. strength. Dose f3j-ij (4.0-8.0). **C.**, **Tinct., Comp.**, cardamom 20, cinnamon 20, caraway 10, cochineal 5, glycerol 60, dilute alcohol, q. s. ad 1000 parts. Dose f3j-ij (4.0-8.0).

Cardia (*kar'-de-ah*) [*καρδία*, the heart]. The heart. **C. of the Stomach**, the esophageal orifice of the stomach.

Cardiac (*kar'-de-ah*) [*καρδία*, the heart]. Pertaining to the heart or to the cardia of the stomach. **C. Cycle**, the period included between the beginning of one heart-beat and

the beginning of another. **C. Dropsy**, a dropsical effusion due to heart-disease with loss of compensation. **C. Ganglia**, ganglia lying in the grooves and substance of the heart—the principal ones are Remak's and Bidder's, the first on the surface of the sinus venosus, and the latter (two) at the auriculo-ventricular groove. **C. Impulse**, the elevation caused by the movement of the heart, usually seen in the fifth left intercostal space. **C. Murmurs**. See *Murmurs*. **C. Orifice**, of the stomach, the esophageal orifice. **C. Passion**, pain in the heart or the stomach. **C. Plexus**. See *Plexus*. **C. Rhythm**, the term given to the normal regularity in the force and volume of the individual heart-beats.

Cardialgia (*kar-de-al'-je-ah*) [*καρδία*, heart; *ἄλγος*, pain]. Pain in the region of the heart, usually due to gaseous distention of the stomach; heart-burn.

Cardinal (*kar'-din-al'*) [*cardo*, a hinge]. Important; preeminent. **C. Flower**, a common name for several species of *Lobelia*, chiefly *L. cardinalis*. **C. Points of Capuron**. See *Pelvis*. **C. Veins**, the venous trunks which, in the embryonic stage, form the primitive jugular veins.

Cardiocele (*kar'-de-o-sel*) [*καρδία*, heart; *κήλη*, tumor]. Hernial protrusion of the heart.

Cardiocentesis (*kar-de-o-sen-te'-sis*) [*καρδία*, heart; *κέντησις*, puncture]. Puncture of one of the chambers of the heart to relieve engorgement.

Cardiodynia (*kar-de-o-din'-e-ah*) [*καρδία*, heart; *δύνη*, pain]. Pain in or about the heart.

Cardiogram (*kar'-de-o-gram*) [*καρδία*, heart; *γράμμα*, a writing]. The tracing made of the cardiac impulse by the cardiograph.

Cardiograph (*kar'-de-o-graf*) [*καρδία*, heart; *γράφειν*, to write]. An instrument for registering graphically the modifications of the pulsations of the heart.

Cardioinhibitory (*kar-de-o-in-hib'-it-o-re*) [*καρδία*, the heart; *inhibere*, to restrain]. Inhibiting or diminishing the heart's action. The C. fibers pass to the heart through the pneumogastric nerves.

Cardiomalacia (*kar-de-o-mal-a'-se-ah*) [*καρδία*, heart; *μαλαξία*, softness]. Softening of the substance of the heart.

Cardiopathy (*kar-de-op'-a-the*) [*καρδία*, heart; *πάθος*, disease]. Any disease of the heart.

Cardiopericarditis (*kar-de-o-per-e-kar-di'-tis*) [*καρδία*, heart; *περί*, around; *καρδία*, the heart; *τις*, inflammation]. Associated carditis and pericarditis; inflammation of the heart-tissues and of the pericardium.

Cardioplegia (*kar-de-o-ple'-je-ah*) [*καρδία*,

heart; *πληγή*, a stroke]. Paralysis of the heart.

Cardiopneumatic (*kar-de-o-nu-mat'-ik*) [*καρδία*, heart; *πνεῦμα*, the breath]. Pertaining to the heart and respiration. **C. Movements**, those movements of the air in the lungs that are caused by the pulsations of the heart and larger vessels.

Cardiopneumograph (*kar-de-o-nu'-mo-graf*) [*καρδία*, heart; *πνεῦμα*, breath; *γράφειν*, to write]. An instrument designed for graphically recording cardiopneumatic movements.

Cardiopuncture (*kar-de-o-punk'-chūr*) [*καρδία*, heart; *punctura*, a puncture]. Cardiocentesis, especially on lower animals.

Cardiovascular (*kar-de-o-vas'-ku-lar*) [*καρδία*, heart; *vasculum*, a small vessel]. Pertaining to the heart and the blood-vessels.

Carditis (*kar-di'-tis*) [*καρδία*, the heart; *τις*, inflammation]. Inflammation of the heart.

C., Internal. Synonym of *Endocarditis*.

Cardol (*kar'-dol*). See *Anacardium*.

Carduus (*kar'-du-us*) [L., a thistle]. The seeds of *C. marianus*, St. Mary's thistle, and *C. benedictus*, blessed thistle. A decoction of the former, $\frac{3}{4}$ ij ad Oj, constitutes an old and popular remedy for hemoptysis. The latter is also a popular cure-all, used mainly as a tonic bitter. Dose of decoction f $\frac{3}{4}$ j-f $\frac{3}{4}$ ss (4.0-16.0); of a tincture, $\frac{m}{x}$ -xx (0.65-1.3).

Carica Papaya (*kar'-ik-ah pap-a'-yah*). See *Papain*.

Caricin (*kar'-is-in*). See *Papain*.

Caries (*kar'-re-ē*) [*caries*, rottenness]. A molecular death of bone, corresponding to ulceration in the soft tissues. **C., Necrotic**, when portions of the bone lie in a suppurating cavity. **C. sicca**, a form of tuberculous caries characterized by absence of suppuration, obliteration of the cavity of the joint, and sclerosis and concentric atrophy of the articular extremity of the bone. **C. of Spine**, or **Pott's Disease**, tuberculous osteitis of the bodies of the vertebrae and intervertebral fibrocartilage, producing curvature of the spine. **C. of Teeth**, a chemico decomposition of the earthy part or any portion of a tooth, accompanied by a partial or complete disorganization of the animal framework of the affected part.

Carious (*kar'-re-us*) [*caries*, rottenness]. Pertaining to or affected with caries.

Carmalum, Mayer's. A stain consisting of carminic acid 1, alum 10, water 200 parts. It is well adapted for sections cut on the freezing microtome.

Carmin (*kar'-min*), $C_{17}H_{16}O_{10}$. A coloring-matter extracted from cochineal.

Carnal (*kar'-nal*) [*carnalis*, fleshly]. Pertaining to flesh. **C. Knowledge**, sexual intercourse.

Carnauba (*kar-na-oo'-bah*) [Braz.]. The root of *Corypha cerifera*, a wax-producing palm-tree of tropic America. It is used in Brazil as an alterative and resembles sarsaparilla in its properties. Dose of the fld. ext. $\text{m xxx-f } \overline{3}$ (2.0-4.0). Unof.

Carnification (*kar-nif-ik-a'-shun*) [*caro*, flesh; *fieri*, to become]. A term indicating the alteration of tissue, especially the lung, to a dense fleshy appearance.

Carnin (*kar'-nin*) [*caro*, flesh], $\text{C}_2\text{H}_5\text{N}_2\text{O}_3$. A leukomycin isolated from American meat-extract, but not from muscle-tissue itself; also obtained from yeast and wine.

Carnivorous (*kar-niv'-o-rus*) [*caro*, flesh; *vorare*, to devour]. Flesh-eating.

Caroba (*kar-a'-bah*) [L.]. The leaflets of *Jacaranda procera* and of *Cyathoxanthus antipathicus*. It is a popular Brazilian remedy as an emetocathartic, alterative, and tonic in syphilis and in yaws. Dose of the fld. ext. $\text{m xv-f } \overline{3}$ (1.0-4.0). Unof.

Carotid (*kar-ot'id*) [*καρωτις*; *καρβειν*, to produce sleep]. 1. The carotid artery, the principal large artery on each side of the neck. See *Arteries*, Table of. 2. Of or relating to the carotid artery.

Carpal (*kar'-al*) [*καρπός*, the wrist]. Pertaining to the wrist.

Carphologia (*kar-fo-l'-je-ah*). See *Carphology*.

Carphology (*kar-fol'-o-je*) [*καρπος*, chaff; *λεγειν*, to collect]. The aimless picking at the bedclothes, seen in grave fevers, particularly in the so-called typhoid state.

Carpometacarpal (*kar-po-met-a-kar'-pal*) [*καρπός*, carpus; *μετά*, between, among; *καρπός*, carpus]. Relating to the carpus and to the metacarpus.

Carpopedal (*kar-po-pet'-dal*) [*καρπός*, carpus; *pes*, *pedis*, a foot]. Affecting the wrists or the hands and feet. **C. Spasm**, a spasm of the hands and feet, or of the thumbs and great toes, associated with the laryngismus stridulus of children. **C. Contraction**. See *Contraction*.

Carpozima (*kar-po-si'-mah*) [*καρπός*, fruit; *ζυμν*, ferment]. A genus of microorganisms producing fermentation.

Carpus (*kar'-us*) [*καρπός*, the wrist]. The eight bones collectively forming the wrist.

Carrageen, or **Carrageen** (*kar'-ag-en*) [*Carrageen* in Ireland]. Irish moss. See *Chondrus*.

Carron Oil (*kar'-on oil*). See *Linum*.

Carswell's Grapes. Pulmonary tubercles when they occur in a racemose distribution at the extremities of several adjacent bronchioles.

Cartilage (*kar'-til-aj*) [*cartilago*, gristle]. Gristle; a white, semi-opaque, nonvascular connective tissue composed of a matrix containing nucleated cells which lie in cavities or

lacunae of the matrix. When boiled, cartilage yields a substance called chondrin.

C., **Articular**, that lining the articular surfaces of bones. **C.**, **Costal**, that occupying the interval between the true ribs and the sternum or adjacent cartilages. **C.**, **Ensi-**

form, the third piece of the sternum. It is also called the xiphoid appendix. **C.**, **Ep-**

actal, small cartilaginous nodules on the upper edge of the alar cartilages of the nose.

C., **Hyaline**, is distinguished by a granular or homogeneous matrix. **C.**, **Fibro-**, dis-

tinguished by a fibrous matrix. **C. of Meckel**, a process from the first branchial arch, from

which the internal lateral ligament of the lower jaw is formed. **C.**, **Reticular**, a

peculiar cartilage found in the auricle of the ear, the epiglottis, and Eustachian tubes. Its

peculiarity consists in a network of yellow elastic fibers pervading the matrix in all

directions. **C. of Santorini**, a nodule at the apex of each arytenoid cartilage, the

corniculum laryngis. **C. of Weitbrecht**, the interarticular fibrocartilage that exists in

the acromioclavicular articulation. **C. of**

Wrisberg, the cuneiform cartilages, one on each side of the fold of membrane stretching

from the arytenoid cartilage to the epiglottis. **C.**, **Yellow**. See *C. Reticular*.

Cartilaginous (*kar-til-aj'-in-us*) [*cartilago*, cartilage]. Made up of or resembling cartilage.

Carum (*kar'-rum*) [*καρον*, caraway]. Caraway. The seeds of *C. Carui*, indigenous to

Europe and an allied species native to the Pacific coast of America. Its odor and

taste are due to a volatile oil. It is used chiefly as a flavor. **C.**, **Aqua** (B. P.).

Dose $\text{f } \overline{3}$ j-ij (32.0-64.0). **C.**, **Infusum**, unof., $\overline{3}$ j-ij, ad Oj. Dose $\text{f } \overline{3}$ ss-ij (16.0-

64.0). **C.**, **Oleum**. Dose m j-v (0.005-0.32).

Caruncle (*kar'-ung-kl*) [*caruncula*, dim. of *caro*, flesh]. A small, fleshy growth. **C.**,

Lacrymal, one upon the conjunctiva near the inner canthus. **C.**, **Urethral**, a small,

bright-red growth situated on the posterior lip of the meatus urinarius; a frequent

condition in women. The caruncle varies in size from a hempseed to a filbert; it is very

painful, especially during micturition and coitus, and bleeds readily.

Carunculæ (*kar-ung'-ku-le*) [*caruncula*, a caruncle]. The nymphæ. **C. myrtiformes**,

the projections of membrane near the orifice of the vagina, thought to be the remains of

the hymen after its rupture.

Carvacrol (*kar'-ak-rol*) [Ital., *carv*, caraway; *ἀσπες*, sharp], $\text{C}_{10}\text{H}_{18}\text{O}$. A liquid

body occurring in the oil of certain varieties of satureia.

Caryocinesis (*kar-e-o-sin-e'-sis*) [*καρυον*, a

nut (nucleus); *κίνησις*, movement]. See *Karyokinesis*.

Caryolysis (*kar-e-ol'-is-is*) [*κάρων*, nucleus; *λύειν*, to loose]. The process of the dissolution of cell-nuclei.

Caryomitosi. See *Karyomitosi*.

Caryophyllin (*kar-e-o-fil'-in*) [*κάρων*, a nut; *φύλλον*, a leaf], $C_{10}H_{16}O$ or $C_{20}H_{32}O_2$. The neutral crystalline principle of cloves.

Caryophyllus (*kar-e-o-fil'-us*) [*κάρων*, a nut; *φύλλον*, a leaf]. Clove. The unexpanded flowers of *Eugenia caryophyllata*, distinguished by their pungent, spicy taste. Its properties are due to a volatile oil, which is antiseptic, stimulant, and irritant. It also contains a crystalline body, eugenin, $C_{10}H_{12}O_2$, and a camphor, caryophyllin, $C_{10}H_{16}O$. It is useful as a stomachic, and to prevent "griping" when combined with purgatives. **C.**, Infusum (B. P.). A strength of 1 to 40 is recommended. Dose f 3j-ij (32.0-64.0). **C.**, Oleum, contains an acid and a phenol compound. Dose m j-iv (0.065-0.26). It is used also by microscopists to clarify preparations and tissues for mounting.

Caryoplasm (*kar'-e-o-plazm*) [*κάρων*, a nucleus; *πλάσμα*, that which is formed]. The plasm of a cell-nucleus; nuclear plasm.

Casca Bark (*kas'-kah bark*). Sassy bark; Ordeal bark. The bark of *Erythrophloeum guineense*, a tree native to Africa. Its properties are due to an alkaloid. It is valuable in intermittent fevers, and as a heart-tonic; in over-doses it produces nausea and vomiting. **Erythrophlein**, the active alkaloid, unof., is a local anesthetic. **C.**, Ext., Aq. Dose gr. j (0.065). **C.**, Ext., Fld. Dose m v-xv (0.32-1.0). **C.**, Tinct., 25 per cent. in strength. Dose m x (0.65). Unof.

Cascara (*kas-kar'-ah*) [Sp., "bark"]. The bark. **C. amarga**, Honduras bark. The bark of a tree native to Mexico, much used as an alterative tonic in syphilis and skin affections. **C. Cordial**, a trade preparation. Dose m xv-f 3j (1.0-8.0). **C.**, Ext., Fld. Dose f 3ss-j; (2.0-4.0); of the powdered extract gr. v-x (0.32-0.65). Unof. **C. sagrada**, Chittam bark, Sacred bark. The bark of *Rhamnus purshiana*, or California Buckthorn. Its properties are due to a volatile oil. It is useful in the treatment of chronic constipation. **Casc. sagrad.**, Ext. (B. P.). Dose gr. ij-viii (0.13-0.52). **Casc. sagrad.**, Ext., Liquidum (B. P.). Dose m v-f 3j (0.32-4.0).

Cascarilla (*kas-kar-il'-ah*) [Sp., dim. of *casca*, bark]. The bark of *Croton eluteria*, native to the Bahama Islands, an aromatic bitter, increasing the natural secretions of the digestive organs. **C.**, Infus. (B. P.). Dose f 3j-ij (32.0-64.0). **C.**, Tinctura (B. P.). Dose f 3ss-ij (2.0-8.0).

Cascarillin (*kas-kar-il'-in*) [*cascarilla*, dim. of *casca*, bark], $C_6H_8O_2$. The active principle of cascarilla-bark; a white, crystalline, bitter substance, scarcely soluble in water.

Caseation (*ka-ze-a'-shun*) [*caseum*, cheese]. The precipitation of casein during the coagulation of milk. Also a form of degeneration in which the structure is converted into a soft, cheese-like substance.

Casein (*ka'-ze-in*) [*caseum*, cheese]. A derived albumin, the chief proteid of milk, precipitated by acids and by rennet. It is closely allied to alkali-albumin, but contains more nitrogen and a large amount of phosphorus. It constitutes most of the curd of milk. **C. Saccharid**, a compound of dry casein one part, cane sugar nine parts, and sodium bicarbonate enough to render it slightly alkaline. It is useful in preparing emulsions of oils, balsams, terpenes, resins, or gum-resins. **C.**, Vegetable, two varieties have been described.—**Legumin**, in peas, beans, etc., and **Conglutin**, in hops and almonds.

Caseous (*ka'-ze-us*) [*caseum*, cheese]. Having the nature or consistency of cheese.

Cassava (*kas-a'-vah*) [Sp., *casabe*]. The fecula of manioc, *Japipha manihot*, separated from the juice. Unof. See *Tapioca*.

Casserian (*kas-e'-re-an*). See *Gasserian*.

Cassia (*cash'-e-ah*) [*kasia*, a perfume]. 1.

A genus of leguminous plants, several species of which afford senna. 2. An old name, still used commercially, for the coarser varieties of cinnamon. See *Cinnamon*.

C. Bark, Cassia lignea. See *Cinnamon*.

C. Buds, the immature fruit of Chinese cinnamon; used chiefly as a spice.

C. marilandica, of N. America, produces the leaves called American senna, which are less active as a cathartic than the true senna.

C. Oil, a variety of oil of cinnamon, used in pharmacy and in perfumery. The pulp, **C.**,

Pulpa (B. P.), is a mild laxative. Dose 3j-ij (4.0-8.0).

C., **Purgin**, is the fruit of a tree, *C. fistula*, growing in tropic regions.

Cast (*kast*) [ME., *casten*, to throw]. 1. A

mass of fibrous or plastic material that has

taken the form of some cavity in which it has

been moulded. From their source, casts may

be classified as bronchial, intestinal, nasal,

esophageal, renal, tracheal, urethral, vaginal,

etc. Of these, the renal casts are the most

important, by reason of their significance in

diseases of the kidney. Classed according

to their constitution, casts are epithelial, fatty,

fibrinous, granular, hyaline, mucous, san-

guineous, waxy, etc. See *Tube casts*. 2.

Strabismus.

Castanea (*kas-ta'-ne-ah*) [L.]. Chestnut.

The leaves of *C. vesca*. They contain tannic

and gallic acids and other principle. the

value of which is not known. They are used in infusion or decoction as a remedy for whooping-cough. **C.**, **Ext.**, **Fld.** Dose π v-lx (0.32-2.6).

Castor (*kas'-tor*). See *Castoreum*. **C. Bean** and **C. Oil**. See *Ricinus*.

Castoreum (*kas-to'-re-um*) [*κάστωρ*, the beaver]. The dried preputial follicles and their secretion, obtained from the beaver, *Castor fiber*. It is a reddish-brown substance with a strong odor. It is antispasmodic and stimulant, its action resembling that of musk. **C.**, **Tinct.** Dose $\frac{1}{2}$ ss-j (2.0-4.0).

Castration (*kas-tra'-shun*) [*castrare*, to cut]. Orchiectomy; the excision of one or both testicles. **C.**, **Female**, removal of the ovaries; oophorectomy; spaying.

Catabolergy, or **Katabolergy** (*kat-ab'-ol-er-je*) [*κατά*, down; *βάλλειν*, to throw; *ἔργον*, work]. Energy expended in katabolic processes.

Catabolic (*kat-ab'-ol'-ik*) [*κατά*, down; *βάλλειν*, to throw]. Of the nature of, or pertaining to, catabolism.

Catabolism (*kat-ab'-o-lizm*) [*κατά*, down; *βάλλειν*, to throw]. Destructive metamorphosis; dissimulation; physiologic disintegration.

Catacrotic (*kat-ah-krot'-ik*) [*κατά*, down; *κρότος*, a striking]. Interrupting the line of descent in a sphygmogram.

Catalepsy (*kat'-al-ep-se*) [*κατά*, down; *λαμβάνειν*, to seize]. A condition of morbid sleep, associated with a loss of voluntary motion and a peculiar plastic rigidity of the muscles, by reason of which they take any position in which they are placed and preserve it for an indefinite time. The condition is associated with hysteria, with forms of insanity, and is a stage of the hypnotic sleep. **C.**, **Local**, that affecting a single organ or group of muscles.

Cataleptic (*kat-al-ep'-tik*) [*κατά*, down; *λαμβάνειν*, to seize]. Relating to, affected with, or of the nature of, catalepsy.

Cataleptiform (*kat-al-ep'-tif-orm*) [*κατά*, down; *λαμβάνειν*, to seize; *forma*, form]. Resembling catalepsy.

Catalysis (*kat-al'-is-is*) [*καταλύειν*, to dissolve]. In chemistry, a reaction that appears to take place owing to the mere presence of another body that apparently undergoes no change.

Catalytic (*kat-al-it'-ik*) [*καταλύειν*, to dissolve]. 1. Of the nature of, or characterized by, catalysis. 2. Any medicine that is supposed to break down, destroy, or counteract morbid agencies existing within the economy.

Catamenia (*kat-am-e'-ne-ah*) [*κατά*, concerning, according to; *μήν*, month]. The recurrent monthly discharge of blood during

sexual life from the genital canal of the female.

Catamenial (*kat-am-e'-ne-ah*) [*κατά*, concerning; *μήν*, a month]. Pertaining to the catamenia.

Cataphoresis (*kat-af-or-e'-sis*) [*κατά*, down; *φορμειν*, to carry]. The introduction of drugs into the system through the skin, by means of ointments or solutions applied by the electrode of a battery.

Cataphoric (*kat-af-or'-ik*) [*κατά*, down; *φορμειν*, to carry]. Passing or causing to pass from the anode to the kathode, through a diaphragm or septum.

Cataplastm (*kat'-ap-lazm*) [*κατάπλασμα*, a poultice]. See *Poultice*.

Cataplasma (*kat-ap-las'-mah*). A poultice. **C. fermenti** (B. P.), a mixture of beer yeast, wheaten flour, water at 100° F. It is a stimulant and antiseptic for indolent ulcers.

Cataract (*kat'-ar-akt*) [*καταρράκτης*, a waterfall]. Partial or complete opacity of the crystalline lens or its capsule. **C.**, **Capsular**, cataract due to opacity of the capsule. **C.**, **Cortical**, one due to loss of transparency of the outer layers of the lens. **C.**, **Diabetic**, is associated with diabetes. **C.**, **Dissection of**, an operation preliminary to absorption, or extraction by suction, consisting in rupturing the capsule, so that the aqueous humor gains access to the lens. **C.**, **Extraction of**, removal of the cataractous lens by surgical operation. **C.**, **Fluid**, the breaking up of an opaque lens into a milky fluid. **C.**, **Green**, a name given to a grayish-green reflex seen in glaucoma; it is also seen when the pupil is dilated and the media are not completely transparent. **C.**, **Hard**. See *C.*, *Senile*. **C.**, **Immature**, only a part of the lens-substance is cataractous. **C.**, **Incipient**, one still in its early stages. **C.**, **Lacteal**. See *C.*, *Fluid*. **C.**, **Lamellar**, one due to opacity of certain layers between the cortex and nucleus, the remaining layers being transparent. **C.**, **Lenticular**, one occurring in the lens proper. **C.**, **Mature**, one in which the whole lens-substance is cataractous. **C.**, **Morgagnian**, when an over-ripe *C.* shrinks and leaves a nucleus floating in the dissolved outer layers. **C.**, **Polar** (anterior or posterior), the opacity is confined to one pole of the lens. **C.**, **Pyramidal**, the opacity is at the anterior pole and is conoid, the apex extending forward. **C.**, **Recurrent Capsular**, or **Secondary**, capsular cataract, appearing after the extraction of the lens. **C.**, **Ripe**. See *C.*, *Mature*. **C.**, **Senile**, the cataract of old persons, the most frequent form, and that understood when not specified as *congenital*, *juvenile*, *traumatic*, *soft*, etc. **C.**, **Soft**, is especially that of the young; the lens matter is of soft consistency

and milky appearance. **C.**, Unripe. See *C.*, Immature. **C.**, Zonular. See *C.*, Lamellar.

Cataractous (*kat-ar-ak'-tus*) [*καταράκτης*, a waterfall]. Of the nature of or affected with cataract.

Catarrh (*kat-ahr'*) [*καταρρέειν*, to flow down]. Inflammation of a mucous membrane. The term is also applied to certain inflammations of the tubules of the kidney and the air-vesicles of the lung. **C.**, Epidemic. See *Influenza*. **C.**, Gastric, gastritis. **C.**, Intestinal, enteritis. **C.**, Nasal, coryza. **C.**, Pulmonary, bronchitis. **C.**, Suffocative. Synonym of *Bronchitis*, *Capillary*. **C.**, Uterine, endometritis. **C.**, Venereal. Synonym of *Gonorrhea*. **C.**, Vesical, cystitis.

Catarrhal (*kat-ahr'-al*) [*καταρρέειν*, to flow down]. Of the nature of, affected with, or relating to catarrh. **C.** Fever. Synonym of *Influenza*. **C.** Inflammation, an inflammation of an archiblastic surface characterized by proliferation and desquamation of the epithelium.

Catastate or Katastate (*kat'-as-tat*) [*κατάστατος*, settling down]. Any one of a series of successive catabolic substances or conditions, each one of which is less complex, more stable, and exhibits less functional activity than its predecessor.

Catatonía, or Katatonía (*kat-at-ó'-ne-ah*) [*κατά*, down; *τόνος*, tension]. A form of insanity characterized by a pathetic emotional state and verberation, combined with a condition of motor tension.

Catechin (*kat'-e-chin*) [*catechu*], $C_{15}H_{10}O_6 + 5H_2O$. Catechinic acid, the active principle of catechu. It crystallizes in shining needles of a snow-white, silky appearance.

Catechu (*kat'-e-choo*) [E. Ind.]. An extract prepared from the wood of *Acacia catechu*, a native of the East Indies. It contains 50 per cent. of tannic acid, and hence is a powerful astringent. It is used for the diarrhea of children and as a gargle and mouth-wash. Dose of the powdered extract gr. x-3ss (0.65-2.0). **C.**, Infusum (B. P.). Dose f3j-iss (32.0-48.0). **C.**, Pulv., Comp. (B. P.). Contains catechu, kino, and rhatany. Dose gr. xx-xl (1.3-2.6). **C.**, Tinct., Composita, contains catechu 12, cinnamon 8, dilute alcohol q. s. ad 100. Dose ℥x-f3j (0.65-4.0). **C.**, Trochisci, each contains one grain of catechu with sugar, tragacanth, and orange-flower water.

Catelectrotonus (*kat-el-ek-tro'-o-nus*) [*κατά*, down; *ηλεκτρον*, amber; *τόνος*, tension]. The state of increased irritability of a nerve near the cathode. See *Anelectrotonus*.

Catgut (*kat'-gut*). The intestines of a sheep

treated to make ligatures. **C.**, Carbolized, catgut rendered aseptic by soaking in a solution of phenol. **C.**, Chromicized, treated with chromic acid. **C.** Plate, an appliance for uniting the intestinal edges in intestinal anastomosis. It is made of a solid catgut sheet; is thin, large, and flat, and resembles the Senn decalcified bone-plates.

Catharsis (*kath-ar'-sis*) [*καθαίρειν*, to purge]. Purgation.

Cathartic (*kath-ar'-tik*) [*καθαρτικός*, purging]. 1. Purgative. 2. A medicine used to produce evacuations of the bowels. A purgative. **C.** Acid. See *Senna*. **C.** Pill, Compound. See *Colocynth*.

Catheter (*kath'-et-er*) [*καθετήρ*, a thing put down]. A hollow tube for introduction into a cavity through a narrow canal. **C.**, Bozeman's, a double-current uterine catheter. **C.**, Eustachian, an instrument for examining the E. tube, distending, or making applications to it. **C.** Fever, systemic disturbance with fever, following the introduction of the catheter into the urethra. **C.**, Gouley's, a solid, curved-steel instrument grooved on its inferior aspect, for passing over a guide, through a stricture into the bladder. **C.**, Lung, a soft-rubber tube that may be passed down the trachea. **C.**, Schrötter's, instruments of hard rubber and of varying caliber, somewhat triangular on section, used for the dilatation of laryngeal strictures. **C.**, Self-retaining, one that will hold itself within the bladder without other appliances to assist it.

Catheterism, or Catheterization (*kath'-et-er-izm*, or *kath-et-er-iz-a'-shun*) [*καθετήρ*, a thing put down]. The introduction of a catheter.

Cathodal (*kath'-o-dal*) [*κατά*, down; *ὁδός*, way]. Relating to the cathode.

Cathode (*kath'-od*) [*κατά*, down; *ὁδός*, way]. The negative electrode or pole of an electric circuit.

Cathodic (*kath-od'-ik*) [*κατά*, down; *ὁδός*, way]. Relating to the cathode.

Cation (*kat'-e-on*) [*κατά*, downward; *ίέναι*, to go]. An electropositive element; one that accumulates at the cathode in electrolysis. See *Ion*.

Catling, or Catlin (*kat'-ling*, or *kat'-lin*) [dim. of *cat*]. A long, pointed, two-edged knife used in amputation.

Catnep, or Catnip (*kat'-nep*, or *-nip*) [corruption of catmint]. The leaves and tops of the herb *Nepeta cataria*, a stimulant and tonic; a popular remedy for chlorosis, hysteria, etc. Dose of fld. ext. f3j-ij (4.0-8.0). Unof.

Catoptric Test. The diagnosis of cataract by means of the reflection of images from the cornea and lens-capsules.

Catoptrics (*kat-op'-triks*) [*κατόπτρον*, a mirror]. The laws of the reflection of light.

Cauda (*kaw'-dah*) [L.]. Tail. **C. cerebelli**, the vermiform process. **C. epididymidis**, the inferior part of the epididymis. **C. equina**, a term applied collectively to the roots of the sacral and coccygeal nerves, from their resemblance to a horse's tail.

Caudad (*kaw'-dad*) [*cauda*, the tail; *ad*, to]. Toward the tail or cauda; opposed to cephalad; in man, downward.

Caudal (*kaw'-dal*) [*cauda*, the tail]. Pertaining to a cauda, or tail.

Caudate (*kaw'-dat*) [*cauda*, a tail]. Having or resembling a tail. **C. Lobe of Liver**, a small lobe of the liver. **C. Nucleus**, the intraventricular portion of the corpus striatum.

Caul (*kawl*) [ME., *calle*, a hood]. 1. A portion or all of the fetal membranes covering the head and carried out in advance of it in labor. 2. The great omentum.

Cauliflower Excrescence. A tumor with an irregular surface resembling the cauliflower.

Caulophyllum (*kaw-lo-fil'-um*) [*καυλός*, stalk; *φύλλον*, leaf]. Blue cohosh, "Squaw-root;" the rhizome and rootlets of *C. thalictroides*, growing in Canada and the northern U. S. It contains a glucosid, saponin, and two resins, one of which is caulophyllin. It produces intermittent contractions of the gravid uterus, and possesses diuretic, emmenagogue, and antispasmodic powers. There are no official preparations. Dose of the powdered drug gr. v-xx (0.32-1.3); of Caulophyllin, gr. ij-v (0.13-0.32).

Causalgia (*kaw-zal'-je-ah*) [*καυσός*, a burning; *άλγος*, pain]. The burning pain that is sometimes present in injuries of the nerves.

Caustic (*kaw'-tik*) [*καίω*, to burn]. 1. Very irritant; burning; capable of destroying tissue. 2. A substance that destroys tissue. **C. Alkali**, a pure alkaline hydrate or oxid. **C. Lunar**, argentic nitrate. **C. Mitigated**, argentic nitrate made less active by fusion with potassium nitrate or argentic chlorid. **C. Potash**, potassium hydrate. **C. Soda**, sodium hydrate.

Cauterant (*kaw'-ter-ant*) [*καυτήρ*, a burner]. 1. Caustic; escharotic. 2. A caustic substance.

Cauterism (*kaw'-ter-ism*). See *Cauterization*.

Cauterization (*kaw-ter-iz-a'-shun*) [*καυτηρίζω*, to cauterize]. The application of a cautery; the effect of such an application.

Cauterize (*kaw-ter-iz*) [*καυτηρίζω*, to cauterize]. To sear or burn with a cautery or a caustic.

Cautery (*kaw'-ter-e*) [*καυτήριον*; a branding iron]. A metal instrument heated by the electric current or in a flame, used to destroy tissue, or for producing counterirritation. **C.**, **Actual**, the white-hot iron. **C.**, **Button**, an

iron heated in hot water. **C.**, **Galvanic**, a platinum-wire heated by electricity. **C.**, **Paquelin's**, or **C.**, **Thermo-**, a hollow platinum-point kept at a uniform temperature by a current of benzene vapor. **C.**, **Potential**, or **C.**, **Virtual**, the application of caustic substances.

Cava (*ka'-vah*) [L.]. A vena cava.

Caval (*ka'-val*) [*cava*, a hollow]. Relating to a vena cava.

Cavalry Bone (*kaw'-al-re bon*). A bony deposit in the adductor muscles of the thigh.

Cavernoma (*kav-ern-o'-mah*) [*caverna*, a cavern; *μα*, a tumor]. A cavernous tumor; a cavernous angioma.

Cavernous (*kaw'-er-nus*) [*caverna*, a cave]. Having hollow spaces. **C. Bodies**, the corpora cavernosa of the penis. **C. Breathing**, the breath-sounds heard over a pulmonary cavity. **C. Groove**, the carotid groove. **C. Plexus**. See *Plexus*. **C. Sinus**, a venous sinus situated at the side of the body of the sphenoid. **C. Tissue**, erectile tissue. **C. Tumor**, a cavernous angioma.

Cavitary (*kaw'-it-a-re*) [*cavitaris*, hollow]. Hollow; having or forming cavities.

Cayenne Pepper (*ki'-en pep'-er*). See *Cap-sicum*.

Cazenave's Lupus. See *Diseases, Table of*.

Cc., an abbreviation for cubic centimeter.

Cd., symbol for Cadmium.

Ce., symbol for Cerium.

Cebocephalus (*se-bo-sef'-al-us*) [*κέβος*, a monkey; *κεφαλή*, head]. A variety of single autositic monsters of the species cyclocephalus, in which there is entire absence of the nose, with, however, two orbital cavities and two eyes, the interocular region being narrow and perfectly flat.

Cedar (*se'-dar*) [*cedrus*, cedar]. One of the genus of coniferous trees, *Cedrus*. **C.**, **Oil of**, a transparent oil obtained from *Juniperus virginiana*, and used as a clearing agent in histology and for oil-immersion lenses.

-cele (*sel*) [*κέλη*, a tumor]. A suffix denoting a tumor.

Celia (*se'-le-ah*) [*κοιλία*, the belly]. 1. The belly; the stomach. 2. A ventricle of the brain.

Celiac (*se'-le-ah*) [*celiacus*; *κοιλία*, belly]. Abdominal; pertaining to the belly. **C. Artery**. Same as *C. Axis*. See *Arteries, Table of*. **C. Axis**, a branch of the abdominal aorta; it divides into the gastric, hepatic, and splenic arteries. See *Arteries, Table of*. **C. Ganglion**. See *Ganglion, Semilunar*. **C. Plexus**. A sympathetic nerve-plexus situated about the origin of the celiac axis.

Celialgia (*se-le-al'-je-ah*) [*κοιλία*, belly; *άλγος*, pain]. Pain in the abdomen.

Celian, **Celine** (*se'-le-an*, *se'-lin*.) Same as *Celiac*.

Celiotomy (*se-le-ot'-o-me*) [*κοιλία*, belly; *rouq*, a cutting]. Surgical opening of the abdominal cavity.

Celiitis, Cœliitis (*se-le-i'-tis*) [*κοιλία*, belly; *itis*, inflammation]. Inflammation of the abdominal organs.

Cell (*sel*) [*cella*, a small, hollow cavity]. 1. A granular mass of protoplasm containing a nucleus. The typical adult cell consists of protoplasm or cell-contents, a nucleus, and within the latter, one or more nucleoli. The cell may or may not have a cell-wall. The protoplasm consists of two parts, the spongioplasm and the hyaloplasm. The nucleus is made up of a nuclear membrane, nuclear fibrils (chromatin), and nuclear matrix (achromatin). The nucleolus is a highly refracting body the function of which is not known. 2. A galvanic element, or single member of a galvanic battery without the connecting wire between the metals. C., Air. See *Air-cell*. C., Ameboid, a cell capable of changing its form and of moving about like an ameba. C., Apolar, a nerve-cell without processes. C., Beaker. See C., *Goblet*. C.-capsule, a thick or unusually strong cell-wall. C., Ciliated, one provided with cilia. C.-cones, the cancer-nests of a squamous epithelioma—so called from their cone-shape. C. of Corti, any one of the hair-cells on the outer surface of the organ of Corti. C., Cylindric, a variety of epithelial cell shaped like a miniature cylinder. C., Daughter, a cell originating from the division of the protoplasm of a mother-cell. C. of Deiters, one of the cells with fine processes resting on the basilar membrane of the cochlea, beneath the air-cells; also a neuroglia-cell. C. Demilune, Giannuzzi's cells, granular protoplasmic cells found in mucous glands, lying between the mucous cells and the basement-membrane. C.-doctrine, the theory that the cell is the unit of organic structure, and that cell-formation is the essential process of life and its phenomena. C., Endothelial, flat cells found on the inner surface of vessels and spaces that do not communicate directly with the external air. C., Epidermic, or Epithelial. See *Epithelium*. C., Ganglion, a large nerve-cell, especially that found in the spinal ganglia. C. of Giannuzzi. See C., *Demilune*. C., Giant, large multinuclear cells occurring in tuberculosis and other infectious granulomata, in bone, in giant-cell tumors, etc. C.-globulins, Halliburton's name for forms of globulin that occur in lymph-corpuscles and can be extracted from them by solutions of sodium chlorid. C., Goblet, an epithelial cell that has been bulged out like a goblet by the presence of mucin. C.-islets, the centers of most active growth in young cellular tissues.

They contain the stores of nutriment that are gradually dissolved and digested. C., Lymphoid, a small, round, connective-tissue cell containing a relatively large nucleus. C., Mother, a cell that divides its protoplasm and gives each part a new cell-wall. C.-multiplication, Cytogenesis, a name given to the process of reproduction of cells. It may be *direct*, as when a cell constricts and cuts off a part of itself, or *indirect*, when the division is preceded by the cycle of nuclear changes known as karyokinesis. C.-nests, a collection of epithelial cells closely packed together and surrounded by a connective-tissue stroma. Cell-nests are found in carcinomata. C., Squamous, a variety of epithelial cells found on the surface of the skin and certain mucous membranes and characterized by their scale-like flatness. C., Vasoformative, or Vasoformative, a cell that in embryonic or very early life, and probably in certain neoplasms, anastomoses with other similar cells so as to form blood-vessels.

Celloidin (*sel-oid'-in*) [*cella*; *eidos*, form]. A concentrated form of collodion for use in imbedding objects for histologic purposes.

Cellular (*sel'-u-lar*) [*cella*, cell]. Relating to or composed of cells. C. Pathology. See *Pathology*.

Cellule (*sel'-ül*) [*cellula*, a small cell]. A small cell.

Cellulitis (*sel-u-li'-tis*) [*cellula*, a small cell; *itis*, inflammation]. A diffuse inflammation of cellular tissue. C., Pelvic. See *Parametritis*.

Cellulose (*sel'-u-lös*) [*cellula*, a little cell], $C_{12}H_{10}O_{10}$. Wood-fiber; lignose, the principal ingredient of the cell-membranes of all plants. It is a white, amorphous mass, insoluble in most of the usual solvents.

Celosoma (*se-lo-so'-mah*) [*κήλη*, hernia; *σώμα*, a body]. A species of single auto-sitic monsters characterized by more or less extensive body-cleft with eventration, associated with various anomalies of the extremities, of the genitourinary apparatus, of the intestinal tract, and even of the whole trunk.

Celotomy (*se-lot'-o-me*) [*κήλη*, hernia; *τεμνω*, to cut]. The operation for strangulated hernia by incision of the stricture.

Cement (*se-ment'*) [*cimentum*, a rough stone]. Any plastic material capable of becoming hard and of binding together the objects that are contiguous to it; filling material for the teeth; the crusta petrosa of the teeth. C. Substance, the substance holding together the endothelial cells of the intima of blood-vessels.

Cenesthesia (*sen-es-the'-tis*) [*κοινῆ*, common; *αἴσθησις*, feeling]. The prevailing condition of feeling, either of depression or of exaltation, which is the resultant of the acts

conscious perceptions from the whole organism.

Centaurea (*sen-taw'-re-ah*) [*κένταυρος*, centaur]. A genus of composite-flowered herbs.

C. benedicta. See *Carduus*.

Centaur (*sen'-taw-re*) [*centauria*]. A popular name for various plants of the genera *Centaurea*, *Erythraea*, *Subbattia*, *Chlora*, etc. It is used as a simple, bitter tonic. Dose $\frac{3}{4}$ ss-j in decoction several times a day. Unof.

Center (*sen'-ter*) [*κέντρον*, the center]. 1.

The middle point of any surface or of a body.

2. The ganglion or plexus whence issue the nerves controlling a function. **C., Accelerating**, a center in the medulla sending accelerating fibers to the heart. These leave the cord through the branches of communication of the lower cervical and upper six dorsal nerves, passing thence into the sympathetic. **C., Auditory**, a center in the first temporosphenoidal convolution, upon each side. **C., Broca's**. See *C., Speech*.

C., Cardioinhibitory, in the medulla, efferent impulses being carried by the vagus. **C., Ciliospinal**, connected with the dilatation of the pupil; it is in the lower cervical part of the cord, and extends downward to from the 1st to the 3d dorsal. **C., Convulsional**, a hypothetic center said to lie in the floor of the 4th ventricle. **C., Coughing**, in the medulla, above the inspiratory center. **C., Diabetic**, in the posterior part of the anterior half of the floor of the 4th ventricle, in the median line. **C., Erection**, is in the lumbar region of the spinal cord, but is controlled from the medulla. **C. for Mastication and Sucking**, in the medulla. **C. for Secretion of Saliva**, on the floor of the 4th ventricle. **C., Olfactory**, probably in the hippocampal region of the temporal lobe. **C., Heat-regulating or Temperature**, the center for the control of body-temperature. See *Thermotaxis*. **C., Leg**, in the upper portion of the ascending frontal convolution. **C. of Ossification**, the place in bones at which ossification begins. **C., Parturition**, in the spinal cord, at the level of the 1st and 2d lumbar vertebrae. **C., Respiratory**, in the medulla, between the nuclei of the vagus and accessorius. **C., Sneezing**, in the medulla. **C., Spasm**, in the medulla, at its junction with the pons. **C., Speech**, in the third left frontal convolution in right-handed people; probably the island of Reil has some influence also. **C., Swallowing**, on the floor of the 4th ventricle. **C., Sweat**, the dominating center is in the medulla, with subordinate centers in the spinal cord. **C., Upper, for Dilator Pupillæ**, in the medulla. **C., Vasodilator**, in the medulla. **C., Vaso-motor**, in the medulla. **C., Visual**, in the occipital lobe, especially in the cuneus.

Centering (*sen'-ter-ing*) [*κέντρον*, a center].

In microscopy, the arrangement of an object or an accessory so that its center coincides with the optic axis of the microscope. In optics, having the pupil and the optic center of the refracting lens in the same axis.

Centesimal (*sen'-tes'-im-al*) [*centum*, hundred]. In the proportion of 1 to 100.

Centigrade (*sen'-te-grad*) [*centum*, a hundred; *gradus*, a step]. Abbreviation, *C.* Having 100 divisions or degrees. **C. Thermometer**, a thermometer with zero as the freezing-point and 100° as the boiling point of water. See *Thermometer*.

Centigram (*sen'-te-gram*) [*centum*, a hundred; *γράμμα*, a small weight]. The hundredth part of a gram, equal to 0.15433 Troy grain.

Centiliter (*sen'-til-e-ter*) [*centum*, a hundred; *λίτρα*, a pound]. The hundredth part of a liter, equal to 0.6102 of a cubic inch.

Centimeter (*sen'-tim-e-ter*) [*centum*, a hundred; *μέτρον*, a measure]. The hundredth part of a meter, equal to 0.39371 (or about $\frac{1}{2}$) of an inch.

Centinormal (*sen'-te-nor'-mal*) [*centum*, a hundred; *norma*, normal]. The $\frac{1}{100}$ of the normal; applied to a solution the $\frac{1}{100}$ of the strength of a normal solution.

Centrad (*sen'-trad*) [*centrum*, the center; *ad*, to]. Toward the center, or toward the median line.

Central (*sen'-tral*) [*centrum*, the center]. Relating to the center; passing through the center. **C. Artery**, an artery in the optic nerve and retina; it passes to the optic papilla and then divides. See *Arteries*, *Table of*. **C. Ganglia**, the corpora striata and thalami optici. **C. Ligament**, the terminal filum of the spinal cord. **C. Lobe**, the island of Reil.

Centraphose (*sen'-traf-ös*). See *Phase*.

Centric (*sen'-trik*) [*centrum*, a center]. Relating to a center, especially to a nerve-center.

Centrifugal (*sen'-trif'-u-gal*) [*centrum*, center; *fugere*, to fly]. Receding from the center to the periphery. **C. Machine**, one by which tubes of liquid are rapidly revolved for the purpose of driving particles floating in the liquid to the distal ends of the tubes.

Centrifuge (*sen'-trif-aj*) [*centrum*, the center; *fugere*, to fly]. A centrifugal machine.

Centripetal (*sen'-trip'-et-al*) [*centrum*, the center; *petere*, to seek]. Traveling toward the center from the periphery.

Centrolecithal (*sen'-tro-les'-ith-al*) [*κέντρον*, center; *λέκθος*, yolk]. In embryology, having the food-yolk located centrally.

Centrosome (*sen'-tro-söm*) [*κέντρον*, center; *σῶμα*, body]. A highly refractive body lying in the protoplasm of the ovum and other cells, and taking an active part in cell-division. It is also known as pole-capsule.

Centrum (*sen'-trum*) [L.]. The center or middle part; the body of a vertebra, exclusive of the bases of the neural arches. **C. ovale majus**, the large mass of white matter appearing when either of the cerebral hemispheres is cut down to the level of the corpus callosum. **C. ovale minus**, the white matter appearing when the upper part of a hemisphere of the brain is removed.

Cephaelis (*sef'-el-is*). See *Specuariaha*.

Cephalad (*sef'-al-ad*) [κεφαλή, head; ad, to]. Toward the head.

Cephalalgia (*sef'-al-ge-ah*) [κεφαλή, head; άλγος, pain]. Headache.

Cephalhematoma (*sef'-al-hem-at-o'-mah*) [κεφαλή, head; αἷμα, blood; ὄμα, tumor]. A collection of blood beneath the pericranium, forming a tumor-like swelling.

Cephalic (*sef'-al-ik*) [κεφαλή, head]. Pertaining to the head. **C. Index**. See *Index*.

C. Version. See *Version*.

Cephalodynia (*sef'-al-o-din'-e-ah*) [κεφαλή, head; ὁδύνη, pain]. Rheumatism affecting the occipitofrontalis muscle, the pain being chiefly experienced in the forehead or occiput, and at times involving the eyeballs.

Cephalomelus (*sef'-al-om'-el-us*) [κεφαλή, head; μέλος, a part]. A form of double monster in which there is a supernumerary limb attached to the head.

Cephalometer (*sef'-al-om'-el-er*) [κεφαλή, head; μέτρον, a measure]. An instrument for measuring the head.

Cephalopagus (*sef'-al-op'-ag-us*) [κεφαλή, head; παγίς, joined]. A double monstrosity having the heads united at the top.

Cephalorrhachidian (*sef'-al-o-rak-id'-e-an*). Same as *Cerebrospinal*.

Cephalotomy (*sef'-al-o'-o-mé*) [κεφαλή, head; τμήσις, section]. The opening or division of the head of the fetus to facilitate labor.

Cephalotribe (*sef'-al-o-trib*) [κεφαλή, the head; τρίβειν, to crush]. An instrument for crushing the fetal head.

Cephalotripsy (*sef'-al-o-trib-se*) [κεφαλή, head; τρίβειν, to crush]. The operation of crushing the fetal head when delivery is otherwise impossible.

Cera (*se'-rah*) [L.]. Wax. A mixture of cerotic acid, cerolein, and myricin, gathered by the honey bee from the pollen of flowers and the leaves of plants. **C. alba**, white wax, prepared by bleaching yellow wax. It is valuable as an ingredient of cerates and ointments. **C. flava**, yellow wax; it possesses an agreeable balsamic odor, and is soluble in ether, in hot alcohol, and in chloroform.

Cerasin (*ser'-as-in*) [*cerasus*, a cherry-tree]. 1. An ingredient of the gum of cherry, peach, and plum trees, apparently identical with bassorin. 2. A crude precipitate of choke-cherry.

Cerate (*se'-rât*) [*cera*, wax]. In pharmacy, an unctuous preparation consisting of wax mixed with oils, fatty substances, or resins, and of such a consistence that at ordinary temperatures it can be readily spread upon linen or muslin, and yet is so firm that it will not melt or run when applied to the skin. **C.**, Goulard's, ceratum plumbi subacetatis. See *Plumbum*.

Cerato- (*ser'-at-o-*). See *Kerato-*.

Cercomonas (*ser-kom'-o-nas*) [κέρκος, tail; μονάς, monad]. A genus of flagellate infusorians. **C. intestinalis**, a protozoon, occasionally found in the fecal discharges of patients suffering with typhoid fever, chronic diarrhea, and cholera. Its pathologic significance has not yet been ascertained.

Cereal (*se'-re-al*) [*Ceres*, the goddess of agriculture]. 1. Relating to edible grains. 2. Any edible grain.

Cerebellar (*ser-e-bel'-ar*) [dim. of *cerebrum*]. Relating to the cerebellum.

Cerebellum (*ser-e-bel'-um*) [dim. of *cerebrum*]. The inferior part of the brain lying below the cerebrum and above the pons and medulla. It consists of two lateral and a middle lobe.

Cerebral (*ser'-e-bral*) [*cerebrum*, the brain]. Relating to the cerebrum. **C. Index**. See *Index*. **C. Pneumonia**. See *Pneumonia*.

C. Vesicles. The embryonic vesicles from which the brain is developed.

Cerebrasthenia (*ser-e-bras-the'-ne-ah*) [*cerebrum*, the brain; ἀσθενής, without strength]. Cerebral asthenia; cerebral neurasthenia.

Cerebration (*ser-e-brat'-shun*) [*cerebrum*, the brain]. Mental activity. **C.**, Unconscious, mental activity of which the subject is not conscious.

Cerebrin (*ser'-e-brin*) [*cerebrum*, the brain], $C_{17}H_{33}NO_2$. A nitrogenous glucosid obtained from brain-tissue, nerves, and pus-corpuscles. It is a light, colorless, exceedingly hygroscopic powder. The term has also been applied to an organic extract of unknown constitution, derived from brain-tissue.

Cerebritis (*ser-e-bri'-tis*) [*cerebrum*, the brain; ιτις, inflammation]. Inflammation of the proper substance of the cerebrum.

Cerebromalacia (*ser-e-bro-mal'-a'-se-ah*) [*cerebrum*, the brain; μαλακία, softness]. Softening of the brain-tissue.

Cerebropathy (*ser-e-brop'-a-the*) [*cerebrum*, the brain; πάθος, illness]. 1. A train of symptoms following overwork, and approaching the character of insanity. 2. Cerebral disease in general.

Cerebroscopy (*ser-e-bros'-ko-pe*) [*cerebrum*, the brain; σκοπεῖν, to inspect]. Investigation of the condition of the brain by inspection, especially ophthalmoscopy, in the diagnosis of brain-disease.

Cerebrose (*ser'-e-brōs*) [*cerebrum*, the brain], $C_6H_{12}O_6$. A crystallized sugar isomeric with glucose, occurring in brain-tissue.

Cerebrosid (*ser'-e-bro-sid*) [*cerebrum*, the brain]. One of a class of substances occurring in brain-tissue, containing cerebrose, just as glucosids contain glucose.

Cerebrospinal (*ser'-e-bro-spi'-nal*) [*cerebrum*, the brain; *spina*, the spine]. Pertaining to the brain and spinal cord. **C. Axis**. See *Axis*. **C. Fluid**, the fluid between the arachnoid membrane and the pia-mater. **C. Meningitis**. See *Fever, Cerebrospinal*. **C. System**, the brain, spinal cord, and nerves.

Cerebrum (*ser'-e-brum*) [L.]. The chief portion of the brain, occupying the whole upper part of the cranium and consisting of the right and left hemispheres.

Cereus (*se'-re-us*) [L., "a wax candle"]. A genus of cactaceous plants. **C. grandiflorus**. See *Cactus grandiflorus*.

Cerevisia (*ser'-e-vis'-e-ah*) [L.]. Ale or beer. **C.**, Fermentum, beer-yeast. The ferment obtained in brewing beer, and produced by saccharomyces cerevisiae.

Cerin (*se'-rin*) [*cera*, wax]. An ether of cerotic acid; one of the substances found in wax.

Cerium (*se'-re-um*) [named after the planet *Ceres*]. $Ce = 140$; quantivalence II, IV. One of the rarer metals. See *Elements, Table of*. **C. oxalae**, $Ce_2(C_2O_4)_3 \cdot 9H_2O$, a white granular powder, insoluble in water or alcohol, but soluble in hydrochloric acid. It is useful in the vomiting of pregnancy. Dose gr. j-x, in pill (0.065-0.65). **C. valerianate** has been used in the same class of cases. Dose gr. iss (0.1).

Cerotic Acid (*se-ro'-tik*). See *Acid, Cerotic*.

Cerumen (*ser-u'-men*) [*cera*, wax]. The wax of the ear.

Ceruminous (*ser-u'-min-us*) [*cera*, wax]. Pertaining to cerumen.

Ceruse (*se'-rūs*) [*cerussa*, white lead]. 1. White lead: basic carbonate and hydrate of lead. 2. A white face-powder. **C. of Antimony**, white oxid of antimony; also antimoniac acid.

Cervical (*ser'-vik-al*) [*cervix*, the neck]. Pertaining to the neck, or to the cervix uteri. **C. Carcinoma**, carcinoma of the neck of the womb. **C. Endometritis**. See *Endometritis*. **C. Pregnancy**, a rare condition in which from atrophy of the decidual membranes the impregnated ovum is not properly held in place, and dropping, lodges in the cervical canal, where it develops until the uterus expels it.

Cervicitis (*ser-vis-i'-tis*) [*cervix*, the neck; *itis*, inflammation]. Inflammation of the cervix uteri.

Cervix (*ser'-viks*) [L.]. A constricted portion or neck. **C. uteri**, the neck of the womb. **C. vesicæ**, the neck of the bladder.

Cesarean, or Cæsarean, Section [from *Cæsar*, Cæsar, the Roman Consul, whose delivery is said to have been effected in this way]. Extraction of the fetus through an incision made in the abdomen. **C. S., Post-mortem**, extraction of the child after the mother's death.

Cestoid, or Cestode (*ses'-toid, or ses'-tōd*) [*κεστός*, a girdle; *ειδος*, likeness]. Shaped like a girdle, or ribbon; applied to worms of which *Tenia* is a type.

Cetacea (*se-ta'-se-ah*) [*cetus*, a whale]. An order of mammals living in the sea, as the whale, dolphin, etc.

Cetaceum (*se-ta'-se-um*) [*κῆτος*, a large fish]. **Spermaceti**. A fatty substance somewhat resembling paraffin in its physical properties. It is obtained from the head of the sperm-whale, *Phyzeter macrocephalus*. It is soluble in ether, in chloroform, and in boiling alcohol, and is employed as an emollient. **C.**, **Ceratum**, contains spermaceti 10, white wax 35, olive-oil 55 parts. **C., Ung.**, (B. P.), contains spermaceti, white wax, almond-oil, and benzoin.

Cetraria (*se-tra'-re-ah*) [*cetra*, a short Spanish shield]. 1. A genus of lichens. 2. Iceland moss,—a lichen, *C. islandica*, found in Iceland and other northern countries. It contains a form of starch, lichenin, that gelatinizes when boiled with water. It is a feebly tonic demulcent, sometimes used in pulmonary affections. **C., Decoct.** (B. P.), contains five per cent. of the lichen. Dose $\frac{1}{3}$ ij-iv (64.0-128.0).

Chagres Fever. See *Fever*.

Chalaza (*kal-a'-zah*) [*χαλαζα*, a hailstone]. One of the twisted cords binding the yolk-bag of an egg to the lining membrane of the shell; or that part of a seed where its coats unite with each other and the nucleus.

Chalazion (*kal-a'-se-on*) [*χαλῶζιον*, a small hailstone]. A tumor of the eyelid from retained secretion of the Meibomian glands; a Meibomian cyst.

Chalcosis (*kal-ik-o'-sis*) [*χαλῆξ*, gravel]. A disease of the lungs caused by the inhalation of dust or sand.

Chalk (*charok*) [*Calx*, limestone]. Carbonate of lime. See *Calcium*. **C. Stone**, gout-stone—a deposit beneath the skin in gouty patients.

Chalybeate (*ka-lib'-e-at*) [*χαλῦς*, iron]. Containing iron.

Chamber (*chām'-ber*) [*camera*, a chamber]. A cavity or space. **C., Anterior** (of the eye), the space between the cornea and the iris. **C., Posterior** (of the eye), the space between the iris and the lens.

The chambers of the eye contain no aqueous humor.

Chamomile (*anemone*) See *Anemone* and *Matricaria*.

Champagne (*anemone*) See *Anemone*. An effervescent wine made with ferment of grapes and containing.

Chancres (*anemone*) [Fr.] A term formerly used indiscriminately for any venereal venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*.

C., **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Chancroid (*anemone*) [Fr.] A term formerly used indiscriminately for any venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*. **C.**, **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Change (*anemone*) [Fr.] A term formerly used indiscriminately for any venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*. **C.**, **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Chantreuil's Method. In bacteriology, a method of ascertaining the distance between the tubercles of the *Salmonella* by estimating the size of the points between. The two tubercles are placed upon the water-soluble while an assistant measures the distance between them.

Chap (*anemone*) [Fr.] A term formerly used indiscriminately for any venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*. **C.**, **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Charcoal (*anemone*) [Fr.] A term formerly used indiscriminately for any venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*. **C.**, **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Charcoal (*anemone*) [Fr.] A term formerly used indiscriminately for any venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*. **C.**, **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Charcot's Disease. *See* *Charcot's Disease*. **Charcot's Joint.** or **Joint Disease.** *See* *Arthritis* and *Disease*. **Table of.** **C.**, **Leyden Crystals**, microscopic, colorless, octahedral or rhomboidal crystals found in semen in the shed blood of leukemic patients, and in the sputum of asthma and croupous bronchitis. Chemically, they are the phosphate of an organic base. **C.'s Method.** *See* *Hypertension*. **C.-Neumann Crystals.** **C.-Robin Crystals.** *See* *C.-Leyden Crystals*.

Charlatan (*shar-lat-an*) [Ital., *ciarlatano*, to

trick] A term formerly used indiscriminately for any venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*. **C.**, **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Chancres (*anemone*) [Fr.] A term formerly used indiscriminately for any venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*. **C.**, **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Chancroid (*anemone*) [Fr.] A term formerly used indiscriminately for any venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*. **C.**, **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Change (*anemone*) [Fr.] A term formerly used indiscriminately for any venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*. **C.**, **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Chantreuil's Method. In bacteriology, a method of ascertaining the distance between the tubercles of the *Salmonella* by estimating the size of the points between. The two tubercles are placed upon the water-soluble while an assistant measures the distance between them.

Chap (*anemone*) [Fr.] A term formerly used indiscriminately for any venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*. **C.**, **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Charcoal (*anemone*) [Fr.] A term formerly used indiscriminately for any venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*. **C.**, **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Charcoal (*anemone*) [Fr.] A term formerly used indiscriminately for any venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*. **C.**, **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Charcoal (*anemone*) [Fr.] A term formerly used indiscriminately for any venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*. **C.**, **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Charcot's Disease. *See* *Charcot's Disease*. **Charcot's Joint.** or **Joint Disease.** *See* *Arthritis* and *Disease*. **Table of.** **C.**, **Leyden Crystals**, microscopic, colorless, octahedral or rhomboidal crystals found in semen in the shed blood of leukemic patients, and in the sputum of asthma and croupous bronchitis. Chemically, they are the phosphate of an organic base. **C.'s Method.** *See* *Hypertension*. **C.-Neumann Crystals.** **C.-Robin Crystals.** *See* *C.-Leyden Crystals*.

Charlatan (*shar-lat-an*) [Ital., *ciarlatano*, to

trick] A term formerly used indiscriminately for any venereal ulcer but now generally limited to the initial lesion of syphilis see *Chancres*. **C.**, **Hard Hammer**, **Indurated**, **Infecting**, **Non-suppurating**, or **True** the ulcer of venereal origin that is followed by constitutional syphilis. **C.**, **Non-indurated**, **Non-infecting**, **Simple**, or **Soft** a benign tuberculous suppurating subcutaneous venereal ulcer, properly called *carbuncle*.

Chelidonin (*kel-id'-o-nin*) [*χελιδόνιον*, celandine], $C_{19}H_{17}N_3O_3 + H_2O$, or $C_{19}H_{17}NO_4$. A crystalline alkaloid of celandin (*Chelidonium majus*).

Chelidonium (*kel-id'-o-ne-um*) [*χελιδόνιον*, celandine]. Celandin. The leaves and stems of *C. majus*, with properties due to a number of alkaloids and acids. It is a drastic cathartic, and externally an irritant, and has been used in jaundice, whooping-cough, and catarrhal pneumonia. Dose of the plant gr. x-xxx (0.65-2.0); of the juice ℥v-xx (0.32-1.3). Unof.

Chelidoxanthin (*kel-id'-oks-an'-thin*) [*χελιδόνιον*, celandine]. One of the bitter, crystalline constituents of celandin.

Cheloid (*ke'-loid*) [*χηλῖδ*, a claw; *εἶδος*, form]. See *Keloid*.

Chelotomy (*ke-lat'-o-me*). See *Kelotomy*.

Chemic, Chemical (*kem'-ik*, *kem'-ik-al*) [*χημεία*, chemistry]. Of or pertaining to chemistry.

Chemiotaxis, Chimiotoxic (*kem'-e-o-taks-is*, *kim'-e-o-taks-is*). See *Chemotaxis*.

Chemistry (*kem'-is-tre*) [*χημεία*, chemistry]. The science of the molecular and atomic structure of bodies.

Chemosis (*ke-mo'-sis*) [*χημωσις*, a gaping]. Swelling of the conjunctiva.

Chemotactic (*kem-o-tak'-tik*) [*χημεία*, chemistry; *τάσσειν*, to arrange]. Pertaining or relating to chemotaxis.

Chemotaxis, Chimiotoxic (*kem'-o-taks-is*, *kem'-e-o-taks-is*) [*χημεία*, chemistry; *τάσσειν*, to order, arrange]. The property of cellular attraction and repulsion. It is displayed by the proteid constituents of the protoplasm of various species of bacteria, as well as by proteids from a great variety of sources. The qualifications positive and negative are added according as the phenomenon is one of attraction or repulsion.

Chemotic (*ke-mot'-ik*) [*χημωσις*, a gaping]. Pertaining to or marked by chemosis.

Chemotropism (*kem-at'-ro-pizm*) [*χημεία*, chemistry; *τροπή*, a turning]. The destruction of bacteria by phagocytes; the victory of the phagocytes over bacteria, or of bacteria over phagocytes. In biology, the attraction of leukocytes by certain chemic substances held in solution in the blood. Cf. *Chemotaxis*.

Chenopodium (*ken-o-po'-de-um*) [*χέν*, a goose; *πόδιον*, a little foot]. American Wormseed; the fruit of *C. ambrosioides*, or *anthelminticum*, a plant native to the United States, with properties due to a volatile oil, which is the only preparation used. It is an efficient anthelmintic against the round worm. Dose of the oil ℥v-xv (0.32-1.0).

Cherry (*cher'-e*) [*κέρρα*, cherry-tree]. The bark of the common cherry, *Prunus sero-*

tina, a mild bitter and tonic, containing tannin. Dose of fld. ext. f5ss-j (2.0-4.0). Prunin, a concentrated extract; dose gr. j-ij (0.065-0.2). See also *Choke-cherry* and *Prunus virginiana*. **C.-laurel**, the European evergreen cherry, *Prunus lauro-cerasus*. Water distilled from its leaves is used in the same way as dilute hydrocyanic acid. Dose ℥xxx-f5j (2.0-4.0).

Chest. See *Thorax*. **C., Barrel-**, a peculiar formation of the chest observed in cases of long-standing emphysema of the lungs; it is round like a barrel, and in respiration is lifted vertically instead of being expanded laterally.

Chestnut. See *Castanea*.

Cheyne-Stokes' Breathing or Respiration. See *Signs and Symptoms, Table of*. **C.-S.**

Nystagmus, a variety of nystagmus in which the oscillations of the eyeball have a rhythmic variation similar to the rhythm of Cheyne-Stokes' respiration.

Chiasm, Chiasma (*ki'-asm*, *ki-as'-mah*) [*χιάζω*, to make a cross, as an X]. The optic commissure.

Chickahominy Fever (*chik-a-hom'-in-e fe'-ver*). See *Fever*.

Chicken (*chik'-en*) [AS., *cicēn*, a chicken].

A domestic fowl. **C. Breast**, an abnormally prominent condition of the sternum and of the sternal region; pigeon-breast; it is seen in rachitic persons, etc. **C. Cholera**. See *Cholera*. **C.-fat Clot**, a clot of blood yellowish in color, consisting largely of fibrin, and containing but few red cells. **C.-pox**. See *Varicella*.

Chicory (*chik'-or-e*) [*κίχώρα*, chicory]. *Cichorium intybus*, a composite plant of Europe and Asia, naturalized and growing in the United States. Its ground root is used to adulterate coffee.

Chilblain (*chil'-blan*) [AS., *cele*, cold; *blēgen*, a boil]. Pernio; Erythema pernio. A congestion and swelling of the skin, due to cold, and attended with severe itching or burning; vesicles and bullæ may form, and these may lead to ulceration.

Child-bed. The popular term for the puerperal state. **C.-bed Fever**, puerperal fever. **C.-birth**, parturition. **C.-crowing**, the crowing sound of the respiration that characterizes laryngismus stridulus.

Chili Saltpeter. Sodium nitrate.

Chill (*chil*) [AS., *chle*, chilliness]. A sensation of cold accompanied by shivering, usually appearing shortly after exposure to cold or wet. It is frequently the initial symptom of acute disorders, as pneumonia, etc. It is a prominent symptom of various forms of malarial fever.

Chills and Fever. A popular term for intermittent fever.

Chiloplasty (*ki'-lo-plas-te*) [*χειλος*, lip; *πλάσ-*

seiv, to form]. Any plastic operation upon the lip.

Chimaphila (*ki-maf'-il-ah*) [*χειμα*, winter; *φιλος*, loving]. Pipsissewa, Prince's Pine; the leaves of *C. umbellata*, an evergreen found in the U. S., an astringent tonic and excellent diuretic. The bruised leaves are used as a rubefacient. It is valuable in dropsy, in renal disease, and in affections of the urinary passages. *C.*, Decoctum. Dose f3j-ij (32.0-96.0). Unof. *C.*, Ext., Fld. Dose f3ss-ij (2.0-8.0).

Chimiotaxis (*kim-e-o-taks'-is*). See *Chemotaxis*.

Chimney-sweep's Cancer. Epithelioma of the scrotum, occurring among chimney-sweepers.

Chin [AS., *cin*, *chin*]. The mentum; the lower part of the face, at or near the symphysis of the lower jaw. *C.*-jerk. See *Jaw-jerk*, and *Reflexes*, Table of. *C.* Reflex. See *Reflexes*, Table of.

China (*kin'-ah*, or *ke'-nah*) [Sp.]. Same as *Cinchona*.

Chinoidin, or **Chinoidinum** (*kin-oi'-din*, or *kin-oi-din'-um*) [Sp., *China*]. Quinoidin. A mixture of amorphous alkaloids obtained in the manufacture of quinin. It has the therapeutic properties of quinin. Dose gr. j-xx (0.065-1.3).

Chinolin (*kin'-o-lin*). See *Chinolina* and *Quinolin*.

Chinolina (*kin-o-li'-nah*) [*china*, quinin], C_9H_7N . Chinolin, Leukolin. An alkaloidal derivative of quinin and cinchonin, occurring also in coal-tar. It is an antiseptic and antipyretic. It is commonly used in the form of the tartrate because of the deliquescence of the other salts. Dose mjjj-x (0.2-0.65); of the tartrate gr. v-xx (0.32-1.0). The tartrate is used as a local application to the throat in strength of 1-10. Unof.

Chionablepsia (*ki-on-ab-lep'-se-ah*) [*χιών*, snow; *ἀβλεψία*, without sight]. Loss of sensibility of the retina resulting from the exposure of the eyes to reflection of the sunlight upon snow.

Chirata (*ke-ra'-tah*, or *che-ra'-tah*) [Hind., *chiraita*, a species of gentian]. Chiretta. The leaves and lighter stems of *C. ophelia*. It resembles gentian in its therapeutic properties, and is an excellent tonic. It does not contain any tannin. Dose of the powdered plant gr. xv-xxx (1.0-2.0). *C.*, Ext., Fld., in glycerol and alcohol. Dose mxxv-xxx (1.0-2.0). *C.*, Infusum (B. P.). Dose, a wineglassful (64.0). *C.*, Tinct., ten per cent. in strength. Dose f3ss-ij (2.0-8.0).

Chiropodist (*ki-rop'-o-dist*) [*χειρ*, hand; *ποδός*, foot]. A surgeon or person who professionally treats diseases of the hands and feet, especially corns, bunions, etc.

Chirurgion (*ki-rer'-jon*) [*χειρουργός*, a surgeon]. A surgeon.

Chirurgia (*ki-rer'-je-ah*) [*χειρ*, hand; *εργον*, work]. Surgery.

Chittim Bark (*chit'-im bark*). See *Cassara sagrada*.

Chloasma (*klo-az'-mah*) [*χλωάζειν*, to be pale green]. Melanoderma; melasma; discolorations. A deposit of pigment in the skin, occurring in patches of various sizes and shapes, and of a yellow, brown, or black color. *C.* hepaticum, liver-spots; a form following dyspepsia and popularly associated with hepatic disturbance. *C.* phthisicorum, the brown patches upon the skin of the forehead or upper portions of the cheeks in tuberculous patients. *C.* uterinum, chiefly located on the forehead, temples, cheeks, nipples, and median line of abdomen. They are marked during pregnancy, and often during menstruation.

Chloracetic Acid (*klo-ras'-el-tik*). See *Acid*.

Chloral (*klo'-ral*) [*chlorin*; *aldehyd*], C_2Cl_3HO . A pungent, colorless, mobile liquid. The name is often misapplied to chloral hydrate. *C.* butylicum, **Butyl-chloral Hydrate** (B. P.), croton-chloral, $C_4H_9Cl_3O$, H_2O , a solid occurring in crystalline scales, resembling chloral hydrate, but made with butyl, C_4H_9 , as a base, instead of ethyl, C_2H_5 . Its properties are parallel to those of chloral, but are much feebler. Dose gr. v-xx (0.32-1.3), in syrup. **Syrupus Butyl-chloral**, strength 1-30. Dose f3j iv (4.0-16.0). Unof. *C.* Hydrate, a colorless, crystalline solid having the composition $C_2H_5Cl_3(OH)_2$, the hydrate of chloral. It is a powerful hypnotic, antispasmodic, and depressant to the cerebral, medullary, and spinal centers, and, to a limited extent, is an anesthetic. It is serviceable in fevers accompanied by cerebral excitement, in chorea, convulsions, and in delirium tremens, but should be used with great caution. Dose gr. v-xx (0.32-1.3). *C.*, **Syrupus** (B. P.). Dose f3ss-ij (2.0-8). *C.*-urethane. See *Uralium*.

Chloralamid (*klo ral' am id*) [*chloral*; *amid*], $CCl_3CH < \begin{smallmatrix} OH \\ NHCHO \end{smallmatrix}$. Chloral formamide. In doses of 30 to 45 grains (2.0-3.0) it is an hypnotic.

Chloralimid (*klo ral' im id*) [*chloral*; *imid*], $CCl_3CH.NH$. An hypnotic allied to chloral, soluble in alcohol. The dose is the same as that of chloral hydrate. Unof.

Chloralism (*klo'-ral-izm*) [*chloral*]. 1. Chloral-poisoning, the morbid state caused by the injudicious use of chloral. 2. The habit of using chloral.

Chloralose (*klo'-ral-ose*) [*chloral*; *glucose*]. Anhydroglycochloral; a product of the

action of anhydrous chloral upon glucose. It is used as an hypnotic. Dose gr. iij-xiv (0.2-0.9).

Chloranemia (*klo-ran-e'-me-ah*). Synonym of *Chlorosis*.

Chloranodyne (*klor-an'-o-din*) [*chlorin*; *ân* priv.; *ôdinv*, pain]. A proprietary remedy introduced as an improvement on chlorodyne. It contains morphin hydrochlorate, tinct. cannab. ind., chloroform, dilute hydrocyanic acid, and aromatics. Dose for an adult $\mathfrak{m}\text{xv}$ (1.0). Unof.

Chlorate (*klo'-rat*) [*chlorin*]. A salt of chloric acid.

Chloric (*klo'-rik*) [*chlorin*]. Pertaining to or containing chlorin. **C. Acid.** See *Acid*. **C. Ether.** 1. See *Ethyl chlorid*. 2. See *Spiritus Chloroformi*.

Chlorid (*klo'-rid*) [*chlorin*]. A binary compound, one of the elements of which is chlorin. **C. of Methyl.** See *Anesthetic*.

Chlorin (*klo'-rin*) [*χλωρός*, green]. Chlorum $\text{Cl} = 35.5$; quantivalence 1. A greenish-yellow gas, prepared by decomposing sodium chlorid, NaCl . It is highly irritative to the skin and mucous membranes, producing spasmodic closure of the glottis. It is a valuable disinfectant. **Chlori, Aqua,** U. S. P.; **C., Liquor,** (B. P.), chlorin-water, contains four per cent. of the gas in solution. It is a good antiseptic wash. Dose, internally, $\mathfrak{m}\text{x-xxx}$ (0.65-2.0). **Calcis Chlorinatæ, Liq.,** (B. P.), contains 1 pound of the salt per gallon of water. **Calx Chlorata,** "chlorid of lime," a hypochlorite of calcium containing free chlorin; a valuable disinfectant. Dose, internally, gr. iij-vj (0.2-0.4). **C.-hunger,** the condition of the body when chlorin (usually in the form of common salt) is lacking. Among the immediate results of this deficiency are indigestion and albuminuria. **C.-vapor** (B. P.), used for inhalation. **Sodæ Chloratæ, Liq.,** Labarraque's Solution, sodium carbonate 10, calx chlorinata 8, water ad 100. Dose $\mathfrak{m}\text{x-f}\mathfrak{v}$ (0.65-4.0), in 20 parts of water.

Chlorobrom (*klo'-ro-brôm*) [*chlorin*; *brâmin*]. A solution each ounce of which contains 30 grains each of chloralamid and potassium bromid. It is hypnotic and useful in seasickness.

Chlorodyne (*klo'-ro-din*) [*χλωρός*, green; *ôdinv*, pain]. A proprietary remedy supposed to contain chloroform, ether, morphin, cannabis indica, hydrocyanic acid, and capsicum. It is anodyne and narcotic. Dose $\mathfrak{m}\text{x-xxx}$ (0.65-2.0). Unof.

Chloroform (*klo'-ro-form*) [*chloroformum*, chloroform]. Methyl terchlorid, CHCl_3 . A heavy, colorless liquid obtained by the action of chlorinated lime on methyl alcohol. The

commercial article, *C. venale*, contains 2 per cent. of impurities. Administered internally in large doses, chloroform produces narcosis and violent gastroenteritis. In small doses it is antispasmodic and carminative. Chloroform has an agreeable odor and a sweetish taste. It solidifies in the cold, boils at 61° , and has a specific gravity of 1.526. Externally, it is much employed as an ingredient of rubefacient and anodyne liniments. Mixed with a large per cent. of air, and inhaled, it is one of the most valuable of general anesthetics, but occasionally (1:3000) causes death by cardiac paralysis. See *Anesthetic*. Deep injections in the vicinity of the sciatic nerve are recommended in sciatica. **C., Alcoholized,** a mixture of chloroform and alcohol. **C., Ammoniated,** equal parts of ammonia in alcohol and chloroform; antipyretic and anodyne. Unof. **C., Aqua** (B. P.). Dose $\mathfrak{f}\mathfrak{ss-ij}$ (16.0-64.0). **C., Emulsum,** chloroform 4, expressed oil of almond 6, tragacanth 1.5, water sufficient to make 100 parts. **C.-liniment,** commercial chloroform 40, soap liniment 60 parts. **C., Liniment, Comp.,** chloroform 4, oil of turpentine 4, tincture of opium 2, tincture of aconite 1 part. Unof. **C., Spt.,** pure chloroform 10, alcohol 90 parts. Dose $\mathfrak{m}\text{x-f}\mathfrak{v}$ (0.65-4.0). **Tinct. Chloroformi et Morphinæ** (B. P.), a substitute for chlorodyne. Each dose of $\mathfrak{m}\text{x}$ contains chloroform $\mathfrak{m}\text{i}\mathfrak{v}$, ether $\mathfrak{m}\mathfrak{v}$, alcohol $\mathfrak{m}\text{i}\mathfrak{v}$, morphin hydrochlorate gr. \mathfrak{v} , dilute hydrocyanic acid $\mathfrak{m}\mathfrak{v}$, oil of peppermint $\mathfrak{m}\mathfrak{v}$, fluid extract of licorice $\mathfrak{m}\text{i}\mathfrak{v}$, treacle and syrup q. s. **Tinct. C. Comp.** (B. P.), chloroform 2, alcohol 8, comp. tinct. of cardamon 10. Dose $\mathfrak{m}\text{xx-f}\mathfrak{v}$ (1.3-4.0).

Chloroformism (*klo'-ro-form-izm*) [*chloroformum*, chloroform]. 1. The use of chloroform to excess for its narcotic effect. 2. The symptoms produced by this use of the drug.

Chloroma (*klo-ro'-mah*) [*χλωμός*, yellowish-green; *bma*, a tumor]. "Green cancer"; a rare variety of sarcoma, of a greenish tint, usually seated upon the periosteum of the bones of the head.

Chlorophane (*klo'-ro-fân*) [*χλωρός*, green; *phainômai*, to appear]. A yellowish-green chromophane. See *Chromophanes*.

Chlorophyl (*klo'-ro-fil*) [*χλωρός*, green; *φύλλον*, leaf]. The green coloring-matter of plants. It decomposes carbon dioxide, setting free oxygen, and forming new organic compounds. This decomposition takes place only or chiefly in the presence of sun-light. The chlorophyl is contained in certain parts of the protoplasm of the plant. It is the substance by the agency of which carbohydrates are formed in green plants.

Chlorosarcoma (*klo-ro-sar-ko'-mah*). See *Chloroma*.

Chlorosis (*klo-ro'-sis*) [*χλωρός*, green]. The "green sickness." A form of anemia, most common in young women, and characterized by a marked reduction of hemoglobin in the blood, with but a slight diminution of red corpuscles. In some cases there is a hyperplasia of the sexual organs and the heart and large blood-vessels. The symptoms are those of anemia, a greenish color of the skin, gastric and menstrual disturbances.

Chlorotic (*klo-ro'-ik*) [*χλωρός*, paleness]. 1. Relating to chlorosis. 2. A person affected by chlorosis.

Chlorphenol (*klor-fe'-nol*) [*χλωρός*, green; *φοίνις*, purple-red], $C_6H_4Cl.OH$. Monochlor-phenol, a substance possessed of antiseptic properties. It is recommended for inhalation in diseases of the respiratory passages.

Chlorum (*klo'-rum*) [*χλωρός*, green]. The official pharmacologic name of chlorine. See *Chlorin*.

Choana (*ko'-an-ah*) [*χόανη*, a funnel]. 1. A funnel-like opening. 2. A name applied to the posterior nasal orifices.

Chocolate (*chok'-o-lat*) [Mex., *chocolatl*, chocolate]. A dried paste prepared from the powder of cacao-seeds (see *Cacao*) with various mucilaginous and amylaceous ingredients. It is used to prepare a beverage, and also as a vehicle, especially for quinine.

Choked Disc (*chokd disk*). See *Papillitis*.

Cholagogue (*kol'-ag-og*) [*χολή*, bile; *ἀγωγός*, leading]. 1. Stimulating the flow or the secretion of bile. 2. Any agent that promotes the flow of bile.

Cholalic Acid (*kol-a'-lik*), $C_{18}H_{34}O_4$. A decomposition product of glycocholic or taurocholic acid, obtained when the latter is boiled with caustic potash, baryta water, or dilute mineral acids. It is dextrorotatory, and is colored blue by iodine.

Cholangitis (*kol-an'-ji'-tis*) [*χολή*, bile; *ἀγγεῖον*, a vessel; *τις*, inflammation]. Inflammation of a bile-duct.

Cholecystectomy (*kol-e-sis-tek'-to-me*) [*χολή*, bile; *κίστις*, bladder; *ἐκτομή*, a cutting off]. Excision of the gall-bladder.

Cholecystenterostomy (*kol-e-sis-ten-ter-os'-to-me*) [*χολή*, bile; *κίστις*, a bladder; *ἐντερον*, intestine; *στόμα*, a mouth]. The artificial establishment of a communication between the gall-bladder and the intestine.

Cholecystitis (*kol-e-sis-ti'-tis*) [*χολή*, bile; *κίστις*, bladder; *τις*, inflammation]. Inflammation of the gall-bladder.

Cholecystolithotripsy (*kol-e-sis-to-lith-ot'-rip-se*) [*χολή*, bile; *κίστις*, a bladder; *λίθος*, a stone; *τρίβειν*, to crush]. The crushing of gall-stones in the gall-bladder.

Cholecystorrhaphy (*kol-e-sist-or'-a-fe*) [*χολή*, bile; *κίστις*, bladder; *ράφή*, a seam]. Suture of the gall-bladder, especially suture to the abdominal wall.

Cholecystostomy (*kol-e-sist-os'-to-me*) [*χολή*, bile; *κίστις*, a bladder; *στόμα*, mouth]. The establishment of an opening into the gall-bladder.

Cholecystotomy (*kol-e-sist-ot'-o-me*) [*χολή*, bile; *κίστις*, bladder; *τέμνειν*, to cut]. Incision of the gall-bladder to remove gall-stones, etc.

Choledochus (*kol-ed'-o-kus*) [*χολή*, bile; *δέχεσθαι*, to receive]. Receiving or holding bile. C., **Ductus Communis**, the common excretory duct of the liver and gall-bladder.

Choleic (*kol-e'-ik*) [*χολή*, bile]. Pertaining to the bile.

Cholelithiasis (*kol-e-lith-i'-as-is*) [*χολή*, bile; *λίθος*, a stone]. The presence of, or a condition associated with, calculi in the gall-bladder or in a gall-duct.

Cholemia (*kol-e'-me-ah*) [*χολή*, bile; *αἷμα*, blood]. The presence of bile in the blood.

Cholera (*kol'-er-ah*) [*χολέρα*; from *χολή*, bile]. 1. A name given to a number of acute diseases characterized mainly by large discharges of fluid material from the bowels, vomiting, and collapse. 2. A synonym of *C. asiatica*. C. *asiatica*, an acute, specific, highly malignant disease, existing in India and the tropics of Asia during the entire year, and occasionally spreading as an epidemic over large areas. It is characterized by vomiting, alvine discharges resembling flocculent rice-water, severe cramps, and collapse. The rate of mortality varies from 10 to 66 per cent., the average being over 50 per cent. The cause is the comma bacillus of Koch, which is always found in the rice-water discharges. The germs commonly gain entrance into the system by means of the drinking-water. C., **Bilious**, a form of the disease attended with excessive discharge of bile. C., **Chicken**, a very fatal epidemic disease of fowls, marked by tumefaction of the lymphatic glands, with inflammation and ulceration of the digestive organs. C., **English**. See *C. nostras*. C., **Epidemic**. Synonym of *C. asiatica*. C., **Hog**, an infectious disease attacking swine and characterized by a patchy redness of the skin, with inflammation and ulceration of the bowels, enlargement of the abdominal glands, and congestion of the lungs. C. *infantum*, the "summer complaint" of infants and young children; an acute disease occurring in warm weather, and characterized by pain, vomiting, purgation, fever, and prostration. The cause of the disease is probably microorganismal, and is favored by the prolonged action of heat, to-

gether with errors in diet and hygiene. It is most common among the poor and in hand-fed babes. The disease is of short duration, death frequently ensuing in from three to five days. *C. morbus*, an acute catarrhal inflammation of the mucous membrane of the stomach and intestines, with pain, purging, vomiting, spasmodic contractions of the muscles, etc. It is a disease of the heated term and is very similar to Asiatic cholera in its symptomatology. *C. nostras*, same as *C. morbus*. *C. sicca*, a term applied to those cases of Asiatic cholera in which rice-water liquid is found in the intestines after death, though none had been voided during life.

Choleraic (*kol-er-ai'-ik*) [*χολέρα*, cholera]. Pertaining to or resembling cholera. **C.** **Diarrhea**, diarrhea characterized by a profuse, exhausting discharge of watery material. **Cholerine** (*kol'-er-en*) [dim. of *cholera*]. A mild form of Asiatic cholera, or the initial stage of a more severe form.

Cholesteatoma (*kol-es-te-at'-o'-mah*) [*χολή*, bile (cholesterin); *στέρμα*, fat; *δύμα*, tumor]. A teratoid tumor containing plates of cholesterol, epithelial cells, hair, and other dermal structures, and occurring most frequently in the brain.

Cholesteremia, **Cholesterinemia** (*kol-es-ter-e'-me-ah*, *kol-es-ter-in-e'-me-ah*) [*χολή*, bile; *στέρμα*, fat; *αἷμα*, blood]. The morbid state resulting from the retention of cholesterol in the blood. The condition is probably due to the retention of the bile-acids.

Cholesterol (*kol-es'-ter-in*) [*χολή*, bile; *στέρμα*, fat], $C_{26}H_{44}O$, or $C_{25}H_{42}O$. A monatomic alcohol, a constituent of bile, gallstones, nervous tissue, yolk, and blood, and sometimes found in foci of fatty degeneration. It is a glistening, white, crystalline substance, soapy to the touch, crystallizing in fine needles and rhombic plates. It is insoluble in water, soluble in hot alcohol, ether, or chloroform. It is held in solution in the bile by the bile-salts; it is levorotatory.

Choletelin (*kol-et'-el-in*) [*χολή*, bile; *τέλος*, completion], $C_{16}H_{18}N_2O_6$. An amorphous, soluble, yellow pigment derived from biliverdin. It is the final product of the oxidation of bile-pigments. It is readily soluble in alkalis, alcohol, and chloroform.

Choleuria (*kol-e-u'-re-ah*) [*χολή*, bile; *ούρον*, urine]. The presence of bile in the urine.

Cholic (*kol'-ik*) [*χολή*, bile]. Pertaining to the bile. **C. Acid**. See *Acid*.

Cholin (*kol'-in*) [*χολή*, bile], $C_5H_{13}NO_2$. A substance found among the products of the decomposition of hog's bile and ox-bile. It is also found in the extracts of the suprarenals, and is a product of the decomposition of lecithin.

Choloidinic Acid (*kol-oi-din'-ik*), $C_{24}H_{38}O_4$. A decomposition product of cholalic acid.

Chololith (*kol'-o-lith*) [*χολή*, bile; *λίθος*, stone]. A gall-stone.

Chondral (*kon'-dral*) [*χόνδρος*, cartilage]. Cartilaginous; relating to or composed of cartilage.

Chondrification (*kon-drif-ik-a'-shun*) [*χόνδρος*, cartilage; *facere*, to make]. The process of being converted into cartilage.

Chondrin (*kon'-drin*) [*χόνδρος*, cartilage]. A substance obtained from the matrix of hyaline cartilage by boiling. It resembles gelatin in general properties, but differs from it in not being precipitated by tannic acid.

Chondritis (*kon-dri'-tis*) [*χόνδρος*, cartilage; *εἶς*, inflammation]. Inflammation of a cartilage.

Chondroblast (*kon'-dro-blast*) [*χόνδρος*, cartilage; *βλαστός*, germ]. A cell of developing cartilage.

Chondrocostal (*kon-dro-kos'-tal*) [*χόνδρος*, cartilage; *costa*, a rib]. Relating to the ribs and their cartilages.

Chondrodendron (*kon-dro-dent'-dron*) [*χόνδρος*, cartilage; *δένδρον*, a tree]. A genus of S. American menispermaceae climbing plants. **C. glaberrimum** and **C. tomentosum** are among the plants that furnish Pareira brava.

Chondrogen (*kon'-dro-jen*) [*χόνδρος*, cartilage; *γενεῖν*, to produce]. A substance forming a part of the tissue of imperfectly developed cartilage.

Chondroma (*kon-dro'-mah*) [*χόνδρος*, cartilage; *δύμα*, tumor]. A cartilaginous tumor.

Chondromalacia (*kon-dro-mal-a'-se-ah*) [*χόνδρος*, cartilage; *μαλακία*, softening]. Softening of a cartilage. **C. auris**. Same as *Hematoma auris*.

Chondrosarcoma (*kon-dro-sar-ko'-mah*) [*χόνδρος*, cartilage; *σάρξ*, flesh; *δύμα*, tumor]. A tumor composed of cartilaginous and sarcomatous tissue.

Chondrotome (*kon'-dro-tôm*) [*χόνδρος*, cartilage; *τομή*, cutting]. An instrument for cutting cartilage.

Chondrotomy (*kon-drot'-o-me*) [*χόνδρος*, cartilage; *τομεῖν*, to cut]. The division of a cartilage.

Chondrus (*kon'-drus*) [*χόνδρος*, a grain; cartilage]. 1. Irish Moss. The substance of the alga *C. crispus* and *C. mammillatus*. These yield, on boiling with water, a soluble colloid consisting mainly of mucilage. This is demulcent and somewhat nutrient. Dose indefinite. Unof. 2. A cartilage; the ensiform cartilage.

Chopart's Amputation. An amputation of the foot consisting of a disarticulation through the tarsal bones, leaving only the os calcis and the astragalus.

Chorda (*kor'-dah*) [L.]. A cord, tendon, or a nerve-filament. *C. dorsalis*. See *Notochord*. *C. tendinea*, any one of the tendinous strings connecting the papillary muscles of the heart with the auriculoventricular valves. *C. tympani*. See *Nerves*, *Table of*.

Chordee (*kor-de'*) [*χορδή*, a cord]. A painful curved erection of the penis with concavity downward. The corpus spongiosum being infiltrated from urethral inflammation, does not fill with blood during erection, and so acts like a bow-string.

Chorea (*ko-re'-ah*) [*χορεία*, dancing]. St. Vitus's dance. A functional nervous disorder, usually occurring in youth, characterized by irregular and involuntary action of the muscles of the extremities, face, etc., with general muscular weakness. Frequently a mitral systolic murmur is heard, often hemic, but in a large proportion of cases due to endocarditis, and there seems to be a close relation between the two diseases. Rheumatism often coexists. Chorea may be caused by a number of conditions, among which are fright and reflex irritation. It affects girls about three times as frequently as boys. Occasionally a form of chorea is seen in the adult, and may become a serious complication of pregnancy, resulting in the death of both fetus and mother. When it occurs late in life it generally resists treatment. *C.*, *Button-maker's*, a form of chorea occurring in persons employed in making buttons. *C.*, *Chronic*. See *C.*, *Huntingdon's*. *C.*, *Electric*. See *Dubini's Disease*. *C.*, *Epidemic*, dancing mania. *C.*, *Habit*. See *Spasm*. *C.*, *Hereditary*. See *C.*, *Huntingdon's*. *C.*, *Hysteric*. See *C.*, *Major*. *C.*, *Huntingdon's*, an hereditary affection of adult or middle life, characterized by irregular movements, disturbance of speech, and gradual dementia. *C. insaniens*, maniacal chorea; a grave form of chorea usually seen in women, and associated with mania, and usually ending fatally. It may develop during pregnancy. *C.*, *Maniacal*. See *C. insaniens*. *C. major*, a form of hysteria in which there are continual regular oscillatory movements. *C.*, *Mimetic*, that which is caused by imitation. *C. minor*, simple chorea. *C.*, *Posthemiplegic*. *C.*, *Postparalytic*, a form of involuntary movement seen in patients after an attack of hemiplegia. *C.*, *School-made*, chorea resulting from overstimulation of children at school. *C.*, *Senile*, a choreiform affection coming on in old age. *C.*, *Sydenham's*, ordinary chorea.

Choreic (*ko-re'-ik*) [*χορεία*, dancing]. Relating to, of the nature of, or affected with, chorea.

Choreiform (*ko-re'-i-form*) [*χορεία*, dancing; *forma*, form]. Resembling chorea.

Choremania (*ko-re-ma'-ne-ah*) [*χορεία*, dancing; *μανία*, madness]. Synonym of *Choromania*.

Choriocapillaris (*ko-re-o-kap-il-a'-ris*) [*χόριον*, skin; *capillus*, a hair]. The network of capillaries over the inner portion of the choroid coat of the eye.

Chorioid (*ko'-re-oid*). See *Choroid*.

Chorioidal (*ko-re-oid'-al*). See *Choroid*.

Chorioideremia (*ko-re-oid-er-e'-me-ah*). See *Chorioideremia*.

Chorioiditis (*ko'-re-oid-i'-tis*). See *Chorioiditis*.

Chorion (*ko'-re-on*) [*χόριον*, fetal membrane]. The outermost of the fetal membranes, formed from the vitelline membrane, the false amnion, and the allantois. The chorion lies between the amnion and the deciduae (reflexa and vera). *C.*, *Cystic Degeneration of*, a myxoma of the chorion, producing the so-called "hydatid mole." It is characterized by rapid increase in the size of the uterus, hemorrhage, often profuse, beginning during the second month of pregnancy, and the discharge of small cysts, whitish in appearance, surrounded by bloody clots. These cysts vary in size from a pin's-head to a filbert. *C. leve*, the membranous portion of the chorion. *C.*, *Shaggy*, or *C. frondosum*, the part covered by villi. It helps to form the placenta.

Chorionic (*ko-re-on'-ik*) [*χόριον*, the chorion]. Relating to the chorion.

Chorionitis (*ko-re-on-i'-tis*). See *Scleroderma*.

Choroid (*ko'-roid*) [*χόριον*, the chorion; *eidos*, likeness]. The vascular tunic of the eye, continuous with the iris in front and lying between the sclerotic and the retina. *C. Plexus*, a vascular plexus in the ventricles of the brain.

Chorioideremia (*ko-roid-er-e'-me-ah*) [*χόριον*, the chorion; *eidos*, like; *ἐρημία*, desolation]. Absence of the choroid.

Choroiditis (*ko-roid-i'-tis*) [*χόριον*, the chorion; *ιτις*, inflammation]. Inflammation of the choroid coat of the eye. It may be anterior, the foci of exudation being at the periphery of the choroid; or central, the exudate being in the region of the macula lutea; diffuse or disseminated, characterized by numerous round or irregular spots scattered over the fundus; exudative or non-suppurative, when there are isolated foci of inflammation scattered over the choroid; metastatic, when due to embolism; and suppurative, when proceeding to suppuration.

C. serosa. Synonym of *Glaucoma*.

Choroidiritis (*ko-roid-o-i-ri'-tis*) [*χόριον*, the chorion; *eidos*, like; *ίρις*, the rainbow; *ιτις*, inflammation]. Inflammation of the choroid and the iris.

Choroidoretinitis (*ko-roid-o-ret-in-i'-tis*) [*χόριον*, the chorion; *ειδος*, like; *retina*, the retina; *ιτις*, inflammation]. Choroïditiſ associated with retinitis. **C.**, **Ametropic**, that caused by ametropia.

Choromania (*ko-ro-ma'-ne-ah*) [*χόρος*, a dance; *μανια*, madness]. A nervous disorder characterized by dancing or other rhythmic movements; epidemic chorea; dancing mania.

Christian Science (*kris'-te-an si'-ens*). An alleged system of therapy; a form of faith-cure.

Christison's Formula. A formula for estimating the amount of solids in the urine: Multiply the last two figures of the specific gravity expressed in four figures by 2.33 (or by 2, *Trapp*; or by 2.2, *Labisch*). This gives the amount of solids in every 1000 parts.

Chromate (*kro'-mat*) [*χρώμα*, color]. Any salt of chromic acid.

Chromatic (*kro-mat'-ik*) [*χρώμα*, color]. Relating to or possessing color. **C. Aberration**, a failure of a lens to bring to the focus some of the color-rays into which white light has been divided, due to the difference in refrangibility of rays of different color. **C. Audition**, luminous sensations aroused by sound.

Chromatin (*kro'-mat-in*) [*χρώμα*, color]. The portion of the protoplasm of a cell that takes the stain, forming a delicate reticular network or plexus of fibrils permeating the achromatin of a cell. It is also called karyomitome.

Chromatogenous (*kro-mat-ō'-en-us*) [*χρώμα*, color; *γενναι*, to beget]. Producing color.

Chromatopsia (*kro-mat-ōp'-se-ah*) [*χρώμα*, color; *ὄψις*, vision]. A disorder of vision in which color-impressions arise subjectively. It may be due to disturbance of the optic centers, or to drugs, especially *santonin*.

Chromic (*kro'-mik*) [*χρώμα*, color]. Pertaining to or made from chromium. **C. Acid.** See *Acid*.

Chromidrosis (*kro-mid-ro'-sis*) [*χρώμα*, color; *ιδρώς*, sweat]. A rare condition of the sweat in which it is variously colored, being bluish, blackish, reddish, greenish, or yellowish. *Black sweat* (*Seborrhœa nigricans*) occurs usually in hysteric women, the face being most often affected. It is associated with chronic constipation and is due to the presence of indican in the sweat. *Red sweat*, *hematidrosis*, may be due to an exudation of blood into the sweat-glands, or to the presence of a microorganism in the sweat.

Chromium (*kro'-me-um*) [*χρώμα*, color]. Cr = 52.2; valence III and IV. One of the elements of the iron group. The various salts of chromium, especially the derivatives of

C. trioxid or chromic acid, Cr₂O₃, are much used in the manufacture of pigments and as a caustic. All are poisonous. Potassium dichromate (bichromate of potash), K₂Cr₂O₇, is used as a caustic, antemetic, and alterative; as a preserving agent for animal tissues, as a reagent, and in battery-fluids.

Chromocytometer (*kro-mo-si-tom'-et-er*) [*χρώμα*, color; *κύτος*, cell; *μέτρον*, a measure]. An instrument for estimating the proportion of hemoglobin present in the blood.

Chromogen (*kro'-mo-jen*) [*χρώμα*, color; *γενναι*, production]. Any principle of the animal or vegetable economy which is susceptible, under suitable circumstances, of being changed into a coloring matter.

Chromophane (*kro'-mo-fan*) [*χρώμα*, color; *φαίνω*, to appear]. The pigment of the inner segments of the retinal cones of certain animals. There are at least three varieties, chlorophane, rhodophane, xanthophane.

Chromophile (*kro'-mo-fil*). Same as *Chromophilous*.

Chromophilous (*kro-moff'-il-us*) [*χρώμα*, color; *φίλος*, loving]. Readily stained; easily absorbing color.

Chromophose (*kro'-mo-fos*). See *Phase*.

Chromophytosis (*kro-mo-fō-to'-sis*) [*χρώμα*, color; *φυτόν*, a plant]. 1. Any microscopic plant-growth that produces a discoloration of the skin in which it grows; any pigmentary skin-disease caused by a vegetable parasite. 2. A synonym of *Tinea versicolor*, or *Eichstedt's Disease*.

Chronic (*kron'-ik*) [*χρόνος*, time]. Long-continued; of long duration; opposed to acute.

Chronicity (*kro-nis'-it-ē*) [*χρόνος*, time]. The state of being chronic or long-continued.

Chronograph (*kro'-no-graf*) [*χρόνος*, time; *γράφειν*, to write]. An instrument for graphically recording intervals of time, in physiologic and psychophysics experiments.

Chrysarobin, or **Chrysarobinum** (*kris-ar-ō'-bin*, or *kris-ar-o-bi'-num*) [*χρῆστος*, golden; *araroba* (nat. East Ind), bark of a leguminous tree], C₁₀H₈O₇. A reduction-product of chrysophanic acid; it occurs in goa-powder and araroba-powder. It is a yellow-colored powder, the product of the decay of *Andira araroba*, a Brazilian tree. It is a gastro-intestinal irritant; locally and internally it is useful in psoriasis, but stains the skin a dark yellowish-brown color. Dose, internally, gr. $\frac{1}{4}$ -1 (0.008-0.032). **C.**, **Ung.**, contains 10 per cent. of the drug with 90 per cent. benzoated lard.

Chrysophanic Acid (*kris-o-fan'-ik*). See *Acid*.

Chvostek's Sign. See *Tetany*; also *Signs and Symptoms*, Table of.

Chyle (*ki'*) [*χυλός*, juice]. The milk-white fluid absorbed by the lacteals during digestion. On standing, it separates into a thin, jelly-like clot and a substance identical with serum. **C.-corpuscle**, any floating cell of the chyle. These cells resemble, and are probably identical with, the colorless blood-corpuscles.

Chylification (*ki-le-fak'-shun*) [*chylus*, chyle; *facere*, to make]. The forming of chyle from food.

Chyliferous (*ki-lif'-er-us*) [*chylus*, chyle; *ferre*, to carry]. Containing or carrying chyle.

Chylification (*ki-le-fik'-a'-shun*) [*χυλός*, chyle; *facere*, to make]. The process by which chyle is formed, separated and absorbed by the villi of the small intestine.

Chylodochium (*kil-o-do'-ke-um*) [*χυλός*, chyle; *δοχείον*, receptacle]. The receptaculum chyli.

Chylopoiesis (*ki-lo-poi-e'-sis*) [*χυλός*, chyle; *ποιεῖν*, to make]. Chylification.

Chylopoietic (*ki-lo-poi-et'-ik*) [*χυλός*, chyle; *ποιεῖν*, to make]. Making or forming chyle.

Chylous (*ki-lus*) [*χυλός*, chyle]. Relating to or resembling chyle.

Chyluria (*ki-lu'-re-ah*) [*χυλός*, juice; *οὐρον*, urine]. The passage of chyle in the urine. It is thought to be caused by a disordered condition of the lacteals, and is also connected with the presence in the blood of *Filaria sanguinis hominis*, which blocks up the lymph-channels.

Chyme (*kim*) [*χυμός*, juice]. Food that has undergone gastric digestion and has not yet been acted upon by the biliary, pancreatic, and intestinal secretions.

Chymification (*ki-me-fik'-a'-shun*) [*χυμός*, chyme; *facere*, to make]. The change of food into chyme by the digestive process.

Cicatricial (*sik-at-rish'-al*) [*cicatrix*, a scar]. Pertaining to or of the nature of a cicatrix. **C. Deformities**, abnormal contractions caused by cicatrices. **C. Tissue**, a form of dense connective tissue, seen in cicatrices.

Cicatrix (*sik-a'-triks*) [L.]. A scar. The connective-tissue which replaces a localized loss of substance. Its color is usually whitish and glistening when old, red or purple when newly-developed.

Cicatritization (*sik-at-riz-a'-shun*) [*cicatrix*, a scar]. The process of healing.

Cilia (*sil'-e-ah*) [Pl. of *cilium*, the eyelid or lash]. 1. The eyelashes. 2. The locomotor and prehensile organs of certain microorganisms. 3. The hair-like appendages of certain epithelial cells, the function of which is to propel fluid or particles.

Ciliary (*sil'-e-a-re*) [*cilium*, an eyelash]. 1. Pertaining to the eyelid or eyelash. 2. Relating to ciliary movement. 3. Pertaining to

the **C. Apparatus**, or the structure related to the mechanism of accommodation. **C. Arteries**, anterior, posterior long, and posterior short, branches of the ophthalmic artery, supplying the recti muscles, the ciliary apparatus, and the posterior structures of the eye, with the exception of the retina. See *Arteries, Table of*. **C. Body**, the ciliary muscle and processes. **C. Canal**, the canal of Fontana. **C. Ganglion**, the ganglion at the apex of the orbit, supplying the ciliary muscle and iris. **C. Movement**, movement by means of vibratory cilia. **C. Muscle**, the muscle of accommodation, the contraction of which lessens the tension upon the suspensory ligament of the lens. **C. Nerves**, branches of the ophthalmic ganglion supplying the anterior structures of the eyeball and the accommodative apparatus. **C. Neuralgia**, neuralgic pain of the eye, brow, temple, etc. **C. Processes**, circularly arranged choroidal foldings continuous with the iris in front. **C. Region**, the pericorneal or "danger-zone," corresponding to the position of the ciliary body. **C. Zone**, the ciliary processes collectively.

Ciliated (*sil'-e-a-ted*) [*cilium*, an eyelash]. Having cilia.

Ciliospinal (*sil-e-o-spi'-nal*) [*cilium*, an eyelash; *spina*, the spine]. Relating to the ciliary zone and the spine. **C. Center**. See *Center*.

Cilium (*sil'-e-um*). [L.]. See *Cilia*.

Cimex (*si'-meks*) [L., a bug]. A genus of hemipterous insects. **C. lectularius**, the common bed-bug.

Cimicifuga (*sim-is-e-fu'-gah*) [*cimex*, a bug; *fugare*, to drive away]. Black snake-root, Black cohosh. The root of *C. racemosa*, ord. Ranunculaceæ. A stomatic, antispasmodic, aphrodisiac, expectorant, and diuretic. Its action on the heart is similar to that of digitalis. It has been used in cardiac diseases, functional impotence, chorea, and ovarian neuralgia. **C., Ext., Fld.** (alcoholic). Dose $\text{m}\text{v}-\text{xxx}$ (0.32-2.0). **C. Ext., Liquid.** (B.P.). Dose $\text{m}\text{ij}-\text{xxx}$ (0.2-2.0). **C., Tinct.**, 20 per cent. in strength. Dose $\text{m}\text{xv}-\text{f}\text{3j}$ (1.0-4.0). **Macrotin.** (unof.), a resinous extract. Dose $\text{gr } \frac{1}{2}-\text{ij}$ (0.032-0.13).

Cina (*si'-nah*) [L.]. The plant *Artemisia santonica*. See *Santonica*.

Cinchamidin (*sin-kam'-id-in*) [*cinchona*; *amidin*], $\text{C}_{20}\text{H}_{28}\text{N}_2\text{O}$. An alkaloid frequently present in commercial cinchonidin.

Cinchona (*sin-ko'-nah*) [from the Countess *Cinchon*]. Peruvian bark. The bark of several varieties of cinchona, a tree native to the eastern slopes of the Andes, the most valuable being *C. calisaya*. Other varieties are *C. succirubra*, red bark, *C. condaminea*, pale bark, *C. pitayensis*, pitaya bark, and *C. micran-*

tha. Cinchona bark contains 21 alkaloids, of which 4, quinin, cinchonin, quinidin, and cinchonidin, are the most important. Cinchona has the same physiologic action and therapeutic uses as its chief alkaloid, quinin. See *Quinin*. It is also an astringent, bitter and stomachic tonic, stimulating appetite and promoting digestion, beneficial in atonic dyspepsia and adynamia, but especially useful in malarial affections. **C.**, Decoctum (B. P.). Dose f 3 j-ij (32.0-64.0). **C.**, Ext., Fld. Dose m x-f 3 j (0.065-0.32). **C.**, Ext., Liquidum (B. P.). Dose m v-x (0.32-0.65). **C.**, Infusum, bark 6, arom. sulph. acid 1, water, 93 parts. Dose f 3 j-f 3 j (4.0-32.0). **C.**, Infus. acidum (B. P.). Dose f 3 j-ij (32.0-64.0). **C.**, Tinct., 20 per cent. of the bark. Dose f 3 ss-ij (2.0-8.0). **C.**, Tinct., Comp., Huxham's Tincture, bark 10, bitter-orange peel 8, serpentaria 1, alcohol 80 parts. Dose f 3 j-f 3 ss (4.0-16.0).

Cinchonamin (*sin-kon'-am-in*) [*cinchona; amin*], $C_{19}H_{24}N_2O$. An alkaloid of cuprea bark. It occurs in glistening, colorless crystals that are nearly insoluble in water, and but slightly soluble in ether.

Cinchonicin (*sin-kon'-is-in*) [*cinchona*], $C_{20}H_{24}N_2O$. An artificial alkaloid derived from cinchonin.

Cinchonidin (*sin-kon'-id-in*), $C_{20}H_{24}N_2O$. An alkaloid derived from cinchona. It is a crystalline substance resembling quinin in general properties. **C.** Salicylate (unof.) has decided antiperiodic properties. **C.** Sulphate, $(C_{20}H_{24}N_2O)_2H_2SO_4 \cdot 3H_2O$, less bitter than quinin and valuable as an antipyretic. Dose gr. j-xx (0.065-1.3) or more.

Cinchonin (*sin'-ko-nin*) [*cinchona*], $C_{19}H_{22}N_2O$. An official alkaloid derived from cinchona. It is a colorless, crystalline body, similar to quinin in therapeutic effects, but less active. **C.** Sulph., $(C_{19}H_{22}N_2O)_2H_2SO_4 \cdot 2H_2O$. It is soluble with difficulty in water, but soluble in acidulated water. Dose gr. v-xxx (0.32-2.0).

Cinchonism (*sin'-ko-nizm*) [*cinchona*]. The systemic effect of cinchona or its alkaloids when given in full doses. The symptoms produced are a ringing in the ears with deafness, headache, giddiness, dimness of sight, and a weakening of the heart's action.

Cinchonize (*sin'-ko-niz*) [*cinchona*]. To bring under the influence of cinchona or its alkaloids.

Cincture (*sing'-tūr*) [*cinctura*, a girdle]. A belt or girdle. **C.-feeling**, a sensation as if the waist were encircled by a tight girdle.

Cinerea (*sin'-e-re-ah*) [*cinereus*, ashen]. The gray substance of the brain, spinal cord, and ganglia. **C.**, Lamina, a thin layer of gray substance extending backward above the

optic commissure from the termination of the corpus callosum to the tuber cinereum.

Cinesia (*sin'-e-se-ah*). See *Kinesia*.

Cingulum (*sing'-gu-lum*) [*cingere*, to gird]. 1. A girdle or zone; the waist. 2. Herpes zoster, or shingles.

Cinnabar (*sin'-ab-ar*) [*κιννάβρις*, a pigment]. Mercuric sulphid, HgS .

Cinnamene (*sin-am'-en*). See *Styrol*.

Cinnamic (*sin-am'-ik*) [*κινναμιν*, cinnamon]. Pertaining to or derived from cinnamon. **C.** Acid. See *Acid*.

Cinnamomum, or **Cinnamon** (*sin-am-o'-mum*, or *sin'-am-on*) [*κιννάμωμον*, cinnamon].

The inner bark of the shoots of several species of *Cinnamomum*, native to Ceylon and China, the latter being known in commerce under the name of *cassia*. Its properties are due to a volatile oil. It is an agreeable carminative and aromatic stimulant, used for flatulence, colic, enteralgia, etc. **C.**, Aqua, 2 parts of oil in 1000 of water. Dose f 3 j-ij (32.0-64.0). **C.**, Ext. Aromat., Fld., contains aromatic powder 10, alcohol 8 parts. Dose m x-xxx (0.65-2.0). **C.**, Oleum, the volatile oil. Dose gtt. j-v (0.065-0.32). **C.**, Pulv., Comp. (B. P.), cinnamon bark, cardamom seeds, and ginger. Dose gr. iij-x (0.2-0.65). **C.**, Spt., 10 per cent. of the oil in spirit. Dose m v-xxx (0.32-2.0). **C.**, Tinct., 10 per cent. of the powdered bark in alcohol. Dose f 3 ss-ij (2.0-8.0). Pulvis aromaticus, aromatic powder, cinnamon, aa 35, cardamom, nutmeg, aa 15. Dose gr. x-xxx (0.65-2.0).

Circle (*si'-kl*) [*κύκλος*, a circle]. A ring; a line, every point of which is equidistant from a point called the center. **C.** of Diffusion. See *Diffusion*. **C.** of Haller. 1. The plexus of vessels formed by the short ciliary arteries upon the sclerotic, at the entrance of the optic nerve. 2. The circulus venosus mammae situated beneath the areola of the nipple. **C.** of Willis, the arterial anastomosis at the base of the brain, formed by the anterior communicating artery between the anterior cerebral arteries, the internal carotids and middle and posterior cerebral arteries, and the posterior communicating arteries.

Circular Insanity. See *Insanity*.

Circuit (*si'-kit*) [*circuitus*, a going round]. The course of an electric current.

Circulation (*si'-ku-la'-shun*) [*circulatio*, a circular course]. Passage in a circle, as the C. of the blood. **C.**, Collateral, that taking place through branches and secondary channels after stoppage of the principal route. **C.**, Fetal, that of the fetus, including the circulation through the placenta and umbilical cord. **C.**, First, or Primitive, that of the embryo, a closed system,

carrying nutriment and oxygen to the embryo. **C.**, **Placental**, the fetal circulation. **C.**, **Portal**, the passage of the blood from the gastrointestinal tract and spleen through the liver, and its exit by the hepatic vein. **C.**, **Pulmonary**, the circulation of blood through the lungs by means of the pulmonary artery and veins, for the purpose of oxygenation and purification. **C.**, **Second**, the fetal circulation, replacing the omphalomesenteric system. **C.**, **Systemic**, the general circulation, as distinct from the pulmonary circulation. **C.**, **Third**, that of the adult. **C.**, **Vitelline**, first, or primitive circulation.

Circum- (*sir'-kum-*) [L.]. A prefix meaning around, about.

Circumcision (*sir-kum-sish'-un*) [*circum*, around; *cadere*, to cut]. The removal of the foreskin.

Circumduction (*sir-kum duk'-shun*) [*circum*, around; *ducere*, to lead]. The movement of a limb in such a manner that its distal part describes a circle, the proximal end being fixed.

Circumflex (*sir'-kum-fleks*) [*circum*, around; *flectere*, to bend]. Winding around. The name given to a number of arteries, veins, and nerves, on account of their course.

Circumscribed (*sir'-kum-skribd*) [*circum*, around; *scribere*, to write]. Strictly limited or marked off; well defined; distinct from surrounding parts, as a circumscribed inflammation or tumor.

Circumvallate (*sir-kum-val'-at*) [*circum*, around; *vallum*, wall]. Surrounded by a wall or prominence. **C.** **Papillæ**, certain papillæ at the base of the tongue.

Cirrhosis (*sir'-o'-sis*) [*κίρρως*, reddish-yellow; from the color of the cirrhotic liver]. Chronic inflammation of an organ characterized by an overgrowth of the connective tissue. **C.**, **Atrophic**, a form of cirrhosis of the liver occurring in hard drinkers, characterized by great overgrowth of the interstitial substance with atrophy of the parenchyma. **C.**, **Biliary**, a form of cirrhosis of the liver due to chronic retention of bile from long-continued obstruction in the bile-ducts. **C.**, **Hypertrophic**, a form of cirrhosis in which the liver is permanently enlarged. The disease is probably infectious and is characterized by an overgrowth of the connective tissue which has no tendency to contract. **C.**, **Periportal**, atrophic cirrhosis, so-called because the hyperplasia of the connective tissue follows the portal vessels.

Cirrhotic (*sir'-o'-ik*) [*κίρρως*, yellow]. Affected with, or relating to, cirrhosis. **C.** **Kidney**, chronic interstitial nephritis.

Cirsocele (*sir'-so-sel*) [*κίρσος*, a varix; *κήλη*, tumor]. A varicose tumor, especially of the spermatic cord.

Cirroid (*sir'-soid*) [*κίρσος*, a varix; *εἶδος*, form]. Resembling a varix, or dilated vein. **Cirsocephalos** (*sir-som'-fal-os*) [*κίρσος*, varix; *ὀμφαλός*, navel]. A varicose condition of the navel.

Cistern (*sist'-tern*) [*cisterna*, a vessel; receptacle]. 1. A reservoir. 2. Any dilatation of the space between the pia and arachnoid. **C.** of **Pequet**, the receptaculum chyli.

Citrate (*sit'-rat*) [*citrus*, citron tree]. Any salt of citric acid.

Citric (*sit'-rik*) [*citrus*, citron-tree]. Pertaining to or derived from lemons or citrons. **C.** **Acid**. See *Acid*.

Citrine (*sit'-rin*) [*citrus*, a lemon]. Yellow; of a lemon-color. **C.** **Ointment**. See *Ily drargyrum*.

Citrus (*sit'-rus*) [L.]. A genus of aurantiaceous trees. See *Aurantium*, *Bergamot*, *Lime*, *Lemon*.

Cladosporium cancerogenes (*klad-o-spo'-um kan-ser-oh'-en-ēs*). A fungus said to be the cause of carcinoma; also termed canceromyces.

Cladothrix (*klad'-o-thriks*) [*κλάδος*, branch; *θρίξ*, a hair]. A genus of bacteria having long, apparently branching filaments.

Clamp (*klamp*) [Ger., *Klampe*]. An instrument for compressing the parts in surgical operations to prevent hemorrhage, etc.

Clap (*klap*) [OF., *clapoir*, a venereal sore]. Gonorrhea. **C.** **threads**, slimy threads consisting of mucus and pus-cells in the urine of gonorrheal patients.

Claret (*klar'-et*) [*clarus*, clear]. A light wine of a red color.

Clarify (*klar'-if-i*) [*clarus*, clear; *facere*, to make]. To free a liquid or solution from insoluble substances; to make clear.

Clarke's Intermediolateral Tract. See *Column of Clarke*. **C.** **Posterior Vesicular Column**. See *Column of Clarke*.

Clasmatocyte (*klas-mat'-o-sit*) [*κλάσμα*, fragment; *κύτος*, cell]. A form of very large connective-tissue corpuscles that tend to break up into granules or pieces.

Clasp (*klasp*) [ME., *clapen*, to grasp firmly]. **C.-knife Rigidity**, a spastic condition of a limb, as a result of which extension is completed with a "spring," as in a knife blade. It is met with in the cerebral palsies of children.

Clathrocystis (*klath-ro-sit'-tis*) [*κλίθρον*, a trellis; *κύστις*, pouch]. A genus of microorganisms with round or oval cells, forming zoogloæ in the form of circular layers.

Claudication (*claw-dic-ah'-shun*) [*claudicare*, to limp]. Lameness. **C.**, **Intermittent**, a spasmodic lameness due to imperfect circulation in the muscles of the lower limbs.

Claustrophobia (*klaw-tro-fo'-be-ah*) [*claudere*, to shut; *φοβία*, fear]. Morbid distress at being in a room or confined space.

Claustrum (*klaw'-trum*) [*claudere*, to shut].

A layer of gray matter in the cerebrum to the outer side of the lenticular nucleus.

Claviceps (*klav'is-eps*) [*clava*, club; *caput*, head]. A genus of fungi. **C. purpurea**, the fungus producing the ergot of rye.

Clavicle (*klav'ik-l*) [*clavicula*; *clavus*, a key]. The collar-bone.

Clavicular (*kla-vik'-u-lar*) [*clavus*, a key]. Relating to the clavicle.

Clavus (*kla'-vus*) [*clavus*, a nail, a wart, a corn]. Corn; a hyperplasia of the horny layer of the epidermis, in which there is an ingrowth as well as an outgrowth of horny substance, forming circumscribed epidermal thickenings, chiefly about the toes. Corns may be hard or soft, the latter being situated between the toes, where they are softened by maceration. Both forms are due to pressure and friction. **C. hystericus**, a local pain in the head, as if a nail were being driven in.

Claw (*klaw*) [*AS.*, *clawon*, a claw]. **C.-hand**, a condition of the hand characterized by over-extension of the first phalanges and extreme flexion of the others. The condition is a result of atrophy of the interosseous muscles, with contraction of the tendons of the common extensor and long flexor. (French, *main-en-griffe*.)

Clearing (*kler'-ing*) [*clarus*, clear]. **C. Agent**, a substance used in microscopy to render tissues transparent and suitable for mounting.

Cleavage (*kle'-vaj*) [*AS.*, *cleafan*, to split asunder]. 1. The linear clefts in the skin indicating the general direction of the fibers. They govern to a certain extent the arrangement of the lesions in skin-diseases. The lines of cleavage run, for the most part, obliquely to the axis of the trunk, sloping from the spine downward and forward; in the limbs they are mostly transverse to their longitudinal axis. 2. A mode of cell-division.

Cleft (*klet*) [*leel.*, *kluft*, a cleft]. Divided. A fissure. **C. Palate**, a congenital fissure of the palate. **C. Sternum**, congenital fissure of the sternum.

Cleft, Visceral, or Branchial. The four slit-like openings on each side in the cervical region in the fetus, sometimes called the Branchial openings. The slits close (in the human fetus), except the upper, from which are developed the auditory meatus, tympanic cavity and Eustachian tube.

Cleido- (*kli'-do-*) [*κλεις* the clavicle]. A prefix, meaning pertaining to the clavicle.

Cleptomania (*klep-to-ma'-ne-ah*). See *Kleptomania*.

Clergyman's Sore-Throat. A chronic hypertrophic form of pharyngitis, with more or less enlargement of the tonsils and lymph-follicles of the posterior wall, due to excessive or improper use of the voice.

Climacteric (*kli-mak-ter'-ik*, or *kli-mak'-ter-ik*) [*κλιμακτήριον*, the round of a ladder]. A period of the lifetime at which the system was believed to undergo marked changes. These periods were thought to occur every seven years. The word is now generally applied to the menopause. **C. Age**, puberty; also in women the time of cessation of the catamenia. **C. Epoch**. Same as *C. Age*. **C., Grand**, the 63d year.

Climatology (*kli-mat-ol'-o-je*) [*κλίμα*, climate, a clime; *λόγος*, science]. The science of climate.

Clinic (*klin'-ik*) [*κλινικός*, pertaining to a bed]. 1. Medical instruction given at the bedside, or in the presence of the patient whose symptoms are studied and whose treatment is considered. 2. A gathering of instructors, students, and patients, for the study and treatment of disease.

Clinical (*klin'-ik-al*) [*κλινικός*, pertaining to a bed]. Relating to bedside-treatment, or to a clinic.

Clinoid (*kli'-noid*) [*κλίση*, a bed; *ειδος*, likeness]. Resembling a bed; applied to sundry bony structures of the body, as the clinoid processes, plate, walls, etc.

Clisometer (*kliis-e-om'-e-ter*) [*κλίσις*, inclination; *μέτρον*, a measure]. An instrument for measuring the degree of inclination of the pelvic axis.

Clitoridectomy (*kliit-or-id-ek'-to-mē*) [*κλειτορίς*, clitoris; *ἐκτομή*, excision]. Excision of the clitoris.

Clitoris (*kliit'-or-is*) [*κλειτορίς*, clitoris]. The homologue in the female of the penis, attached to the ischiopubic rami by two crura or branches, which meet in front of the pubic joint to form the body, or corpus. It possesses erectility. **C. Crises**, paroxysms of sexual excitement in women suffering from tabes.

Clivus (*kli'-vus*) [*L.*, a slope]. A slope. **C. ossis**, or **C. of Blumenbach**, the slanting surface of the body of the sphenoid bone between the sella turcica and the basilar process of the occipital bone.

Cloaca (*klo-a'-kah*) [*cloaca*, a sewer]. 1. In early fetal life, the common orifice of the intestine and the allantois. 2. A fistulous tract in bone discharging pus from a sequestrum.

Clonic (*klon'-ik*) [*κλονος*, commotion]. Applied to convulsive and spasmodic conditions of muscles characterized by alternate contractions and relaxations.

Clonus (*klo'-nus*) [*κλονος*, commotion]. A series of movements characterized by alternate contractions and relaxations; a clonic spasm. Involuntary, reflex, irregular contractions of muscles when put suddenly upon the stretch. According to the part affected, the phenomenon is spoken of as *ankle*, *foot*,

rectus, or *wrist C.*, etc. See *Reflexes*, *Table of*.

Cloquet, Canal of. See *Canals*, *Table of*.

Clostridium (*klos-trid'-e-um*) [*κλωστήρ*, a spindle]. A genus of bacteria, differing from bacilli in the fact that their spores are formed in enlarged rods.

Closure (*klo'-zūr*) [*clausura*, a closing]. The act of completing or closing an electric circuit.

Clot (*klot*) [*AS.*, *clate*, a bur]. A peculiar solidification of the blood, such as takes place when it is shed. It is due to the formation of fibrin which entangles the blood corpuscles and, contracting, squeezes out the liquid portion of the blood.

Cloudy Swelling. Parenchymatous degeneration; a swelling up of the elements of a tissue, with the formation in them of fine granules due to the change of soluble albuminates into insoluble.

Clove (*klōv*). See *Caryophyllus*. **C.-Hitch Knot**, a form of double-knot in which two successive loops are made close to one another on the same piece of cord or bandage, a half-twist being given to the junction of each loop at the time of making it.

Clownism (*kloun'-izm*) [*Icel.*, *klunni*, a boorish fellow]. That stage of hysterio-epilepsy in which there is an emotional display and a remarkable series of contortions.

Club-foot. See *Talipes*. **C.-hand**, a deformity of the hand similar to that of club-foot.

Clubbed Fingers (*klubd fin'-gers*). Knobbed deformity of the finger-tips, with curvature of the nails over the finger-ends; seen in some cases of pulmonary and cardiac disease.

Clyster (*kliis'-ter*) [*κλύστήρ*, an injection]. An enema.

Coagulated (*ko-ag'-u-la-ted*) [*coagulare*, to curdle]. Clotted; curdled. **C. Proteids**, a class of proteids produced by heating solutions of egg-albumin or serum-albumin up to 70° C. or higher. At the body-temperature they are readily converted into peptones by the action of the gastric juice in an acid medium, or of pancreatic juice in an alkaline medium.

Coagulation (*ko-ag'-u-la'-shun*) [*coagulatio*, a clotting]. The formation of a coagulum or clot, as in blood or in milk.

Coagulative (*ko-ag'-u-la-tiv*) [*coagulare*, to curdle]. Causing or marked by coagulation.

C. Necrosis. See *Necrosis*.

Coagulum (*ko-ag'-u-lum*) [*coagulare*, to curdle]. A clot. The mass of fibrin, enclosing red and colorless corpuscles and serum, that forms from the blood after the latter has been drawn from the body. Also, the curd of milk, and the insoluble form of albumin.

Coal (*kōl*) [*ME.*, *cole*]. **C.-tar**, a by-pro-

duct in the manufacture of illuminating gas; it is a black, viscid fluid of a characteristic and disagreeable odor. The specific gravity ranges from 1.10 to 1.20. Its composition is extremely complex, and its principal constituents are separated, one from the other, by means of fractional distillation.

Coalescence (*ko-al-es'-ens*) [*coalescere*, to grow together]. The union of two or more parts or things.

Coaptation (*ko-ap'-ta'-shun*) [*con*, together; *aptare*, to fit]. The proper union or adjustment of the ends of a fractured bone, the lips of a wound, etc.

Coarctate (*ko-ark'-tat*) [*coarctare*, to press together]. Crowded together.

Coarctation (*ko-ark'-ta'-shun*) [*coarctare*, to put together]. A compression of the walls of a vessel or canal, narrowing or closing the lumen; reduction of the normal or previous volume, as of the pulse; shrivelling and consequent detachment, as of the retina. A stricture.

Coarse (*kōrs*) [*ME.*, *course*]. Not fine; gross. **C. Adjustment.** See *Adjustment*.

C. Features of Disease, macroscopic organic lesions, such as swelling, hemorrhage, etc.

Coat (*kōt*) [*cottus*, a tunic]. A cover, or membrane covering a part or substance. **C., Buffy**, the upper fibrinous layer of the clot of coagulated blood, characterized by its pale color, due to absence of red corpuscles.

Cobalt (*kō'-barult*) [*Kobold*, a German mythologic goblin]. A tough, heavy metal having some of the general properties of iron. Its oxides have been employed in medicine, but are now very little used. See *Elements*, *Table of*.

Coca (*kō'-kah*). See *Erythroxyton*.

Cocain (*kō'-kah-in*, or *kō'-kan*) [*S.A.*, *coca*], $C_{17}H_{21}NO_4$. The chief alkaloid of *Erythroxyton coca*. It is at first stimulant and afterward narcotic, and resembles caffeine in its action on the nerve-centers, and atropin in its effects on the respiratory and circulatory organs. Its long-continued use (cocain-habit) is followed by insomnia, decay of moral and intellectual power, emaciation, and death. It is a local anesthetic when applied to the surface of mucous membranes or given hypodermically. Applied to the conjunctiva of the eye it causes also dilatation of the pupil and paralysis of the function of accommodation. Dose gr. $\frac{1}{4}$ -ij (0.008-0.13). **C., Ceratum**, 1 in 30, for burns, etc. **C. Citras**, used to stop toothache. **C. Hydrochloras**, $C_{17}H_{21}NO_4Cl$, most commonly used for local anesthesia in 2-8 per cent. solution. Dose, internally, gr. $\frac{1}{4}$ -ij (0.008-0.13). **C. Hydrochlorat.**, *Liquor* (B. P.). Dose m℥ij-x (0.13-0.65). *Injectio C. Hydrochlorat.*

Hypoderm.; 1 in 20. **C.**, **Lamellæ** (B. P.), each containing gr. $\frac{1}{100}$ (0.00065) of cocain hydrochlorate. **C. Oleas**, a 10 per cent. solution in oleic acid, for external use. **C. Phenas** or **Carbolas**, a topic application in catarrhs and in rheumatism, used as a five to ten per cent. alcoholic solution; also internally. Dose, gr. $\frac{1}{12}$ - $\frac{1}{8}$ (0.005-0.016).

Cocainism (*ko-ka'-in-izm*) [S. Amer., *coca*]. The cocain-habit.

Cocainization (*ko-ka'-in-iz-a'-shun*) [S. Amer., *coca*]. The bringing of the system or organ under the influence of cocain.

Coccaceæ (*kok-kas'-e-e*) [*κόκκος*, a kernel]. A group of schizomycetous fungi, or bacteria, including as genera the *Micrococcus*, *Sarcina*, *Ascococcus*, and *Leuconostoc*.

Coccidiosis (*kok-sid-i'-o-sis*) [*κόκκος*, a berry; *vóσος*, disease]. The group of symptoms produced by the presence of coccidia in the body.

Coccidium (*kok-sid'-e-um*) [*κόκκος*, a berry]. A genus of protozoans, by some referred to as the so-called psorosperms (see *Psorosperm*). **C. oviforme** has been found in intestinal epithelium, and in the liver of man, and often in the liver of the rabbit. True coccidia are nonmotile cell-parasites. **C. sarkolytus**, the name given by Adamkiewicz to the so-called parasite of carcinoma.

Coccinella (*kok-sin-el'-ah*). See *Cochineal*.

Coccobacteria (*kok-o-bak-te'-re-ah*) [*κόκκος*, a kernel; *βακτηριον*, a little rod]. The rod-like or spheroidal bacteria found in putrefying liquids, and called **C. septica**. See *Bacteria*, *Table of*.

Cocculus indicus (*kok'-u-lus in'-dic-us*). The dried fruit of *Anamirta cocculus*. It is an active narcotic poison. It is employed as a destroyer of vermin. See *Picrotoxin*.

Coccus (*kok'-us*) [*κόκκος*, a berry]. 1. A genus of insects containing *C. cacti*, the cochineal insect. 2. A spheric bacterium, a micrococcus.

Coccydynia (*kok-se-din'-e-ah*). See *Coccygodynia*.

Coccygeus (*kok-sij'-e-us*) [*κόκκυξ*, cuckoo]. One of the pelvic muscles. See *Muscles*, *Table of*.

Coccygodynia (*kok-sij-o-din'-e-ah*) [*κόκκυξ*, cuckoo; *δύσπη*, pain]. Pain referred to the region of the coccyx; confined almost exclusively to women who have given birth to children.

Coccyx (*kok'-siks*) [*κόκκυξ*, the cuckoo (resembling the bill)]. The last bone of the spinal column, formed by the union of four rudimentary vertebrae.

Cochin Leg (*ko'-chin*). Synonym of *Elephantiasis arabum*.

Cochineal (*ko'-in-el*, or *koch-in-el'*) [ME., *cutchaneal*]. The dried insects of a species

of plant lice, *Coccus cacti*, parasitic upon a cactus of Mexico and Central America. It contains a rich red coloring-matter, carmin, used mainly as a dyeing agent. It is thought to be valuable in whooping-cough. Dose gr. $\frac{1}{2}$ (0.02).

Cochlea (*kok'-le-ah*) [*κόχλος*, a conch-shell]. A cavity of the internal ear resembling a snail-shell. It describes two and a half turns about a central pillar called the modiolus or columella, forming the spiral canal, about $1\frac{1}{2}$ inches in length. See also *Ear*.

Cochlear, or **Cochleare** (*kok'-le-ar*, or *kok-le-a'-re*) [L.]. A spoon; a spoonful. **C. magnum**, a tablespoon; **C. medium**, a dessert-spoon; **C. minimum**, a teaspoon.

Cockroach (*kok'-rach*). See *Blatta*.

Coco (*ko'-ko*). Synonym of *Yaws*. See *Frambesia*.

Coco, or **Cocoa** (*ko'-ko*) [Port., *cacao*]. See *Cacao* and *Theobroma*.

Cod (*kod*) [ME., *cod*]. The *Gadus morrhua*, a fish furnishing **C.-liver Oil**, an oil derived from the liver of the *Gadus morrhua*, and ranging in color according to the method of its preparation from pale-straw to dark-brown; its specific gravity is .923 to .924 or even .930 at 15° C. See *Morrhua*.

Codein (*ko'-de-in*) [*κώδεα*, the poppy-head], $C_{15}H_{21}NO_3$. A white, crystalline alkaloid of opium resembling morphin in action, but being weaker. It is used in cough and in diabetes mellitus. Dose gr. ss-ij (0.032-0.13). **C. Phosphate**, soluble in water. It is similar to morphin in action, but less toxic. Dose, hypodermically, gr. ss (0.032). **C. Sulphas**, dose gr. $\frac{1}{6}$ - $\frac{1}{4}$ (0.01-0.016). **C. Valerianate**, an antispasmodic and sedative. Dose gr. $\frac{1}{4}$ (0.016).

Cœliac (*se'-le-ah*). See *Celiac*.

Cœliotomy (*se-le-ah'-o-mie*). See *Celiotomy*.

Cœlom, **Cœloma** (*se'-lom*, *se-lo'-mah*). See *Celom*.

Cœnesthesis (*sen-es-the'-sis*). See *Cenesthesia*.

Cœnurus (*se-nûr'-us*) [*καόνος*, common; *οἰπά*, tail]. The larva of *Tœnia cœnurus* (Küchenmeister), producing the disease of sheep called staggers. **C. cerebralis**, an hydatid found mainly in the brain and spinal canal of the ox and sheep (mostly in young animals). Occasionally it has been discovered in the muscles of man. It is known to be the larva of the tapeworm, *tœnia cœnurus*.

Coffee (*kof'-e*). See *Caffea*. **C.-ground Vomit**, the material ejected by emesis in gastric carcinoma and other conditions that give rise to a slow hemorrhage into the stomach. It consists of blood changed by the action of the gastric juice, and mixed with other contents of the stomach.

Cogwheel Breathing or Respiration. A type of breathing characterized by a jerky, wavy inspiration.

Cohabitation (*ko-hab-it-a'-shun*) [*con*, together; *habitare*, to dwell]. The living together of a man and woman, with or without legal marriage; sexual connection.

Cohesion (*ko-he'-zhun*) [*coherere*, to stick together]. The force whereby molecules of matter adhere to each other; the attraction of aggregation.

Cohnheim's Areas. See *Area*. **C. Fields.** See *Cohnheim's Areas*. **C. Frog.** See *Salt frog*. **C. Theory**, a theory that all true tumors are due to faulty embryonal development. The embryonal cells do not undergo the normal changes, are displaced, or are superfluous. When the favorable conditions are presented later in life they take on growth, with the formation of tumors of various kinds.

Cohosh (*ko'-hosh*) [Am. Ind.]. A name given to several medicinal plants. **C., Black.** See *Cimicifuga*.

Coil (*koil*) [*colligere*, to gather together]. A spiral formed by winding. **C.-gland.** See *Sweat-gland*. **C., Induction**, rolls of wire used to produce an electric current by induction. **C., Leiter's.** See *Lleiter's Tubes*.

Coín-test (*koin'-test*). See *Tympany, Bell*.

Coition (*ko-ish'-un*) [*coire*, to come together]. Same as *Coitus*.

Coitophobia (*ko-it-o'-fo'-be-ah*) [*coire*, to come together; *φόβος*, fear]. Morbid dread of coitus from disgust or dyspareunia.

Coitus (*ko'-it-us*) [*coire*, to come together]. The act of sexual connection; copulation.

Coko Disease (*ko'-ko dis-ēz'*). A name applied in the Fiji Islands to a disease resembling yaws.

Cola (*ko'-lah*). See *Kola-nut*.

Colchicin (*kol'-chis-in*) [*κόλχικόν*, colchicum], $C_{17}H_{19}NO_6$. An alkaloid of colchicum; it is a pale, brownish-yellow, exceedingly bitter powder, freely soluble in water. It is a very active poison. Its dose is gr. $\frac{1}{16}$ (0.0032), hypodermatically.

Colchicum (*kol'-chik-um*) [*κόλχικόν*, colchicum]. Meadow-saffron. The corm and seed of *C. autumnale*, the properties of which are due to an alkaloid, colchicin. It is an emetic, diuretic, diaphoretic, and drastic cathartic. It is valuable in acute gout, and in some forms of rheumatism. Dose of the powdered root gr. ij-vij (0.13-0.52), of the powdered seeds gr. j-v (0.065-0.32). **C., Extract.** (B. P.). Dose gr. ss-ij (0.032-0.13). **C., Extract, Aceticum** (B. P.). Dose gr. ss-ij (0.032-0.13). **C. Rad., Ext.** Dose gr. ss-ij (0.032-0.13). **C. Rad., Ext.,**

Fld. Dose \mathfrak{m} ij-x (0.13-0.65). **C. Rad., Vinum**, 40 per cent. in strength. Dose \mathfrak{m} v-xv (0.32-1.0). **C. Sem., Ext., Fld.** Dose \mathfrak{m} j-v (0.065-0.32). **C. Sem., Vinum**, 15 per cent. in strength. Dose \mathfrak{m} x-xxx (0.65-2.0). **C., Tinct.**, prepared from the seed; strength 15 per cent. Dose \mathfrak{m} x-xxx (0.65-2.0). **C., Vinum** (B. P.). Dose \mathfrak{m} x-xxx (0.65-1.3).

Cold (*kold*) [AS., *ceald*]. 1. The comparative want of heat. 2. A term used popularly for coryza and catarrhal conditions of the respiratory tract. Cold is employed largely in various forms as a therapeutic agent, mainly for the purpose of lowering temperature and allaying irritation and inflammation. It may be used in the form of affusion, that is, the sudden application of a considerable volume of cold water to the body. Cold may be used as an anesthetic in baths (see *Bath*); in the form of compresses applied over the affected part; in the form of irrigation, especially in the treatment of bruised and injured members; as a lotion, for the purpose of relieving local heat, pain, and swelling; as an injection, in the form of ice-water, into the vagina or rectum, for various conditions; and as the cold pack, which is a valuable means of reducing the body-temperature in cases of hyperpyrexia. Cold may be applied in the dry form by means of the ice-cap or bladder, an india-rubber bag filled with ice, snow, or a freezing mixture. **C. Abscess.** See *Abscess*. **C. Cream.** See *Rosa*. **C. Pack.** See *Pack*. **C. Sore**, herpes labialis.

Colectomy (*ko-lekt'-to-me*) [*κόλον*, colon; *ἐκτομή*, cutting out]. Excision of a portion of the colon.

Colic (*kol'-ik*) [*κόλικός*, pertaining to the colon]. 1. Pertaining to the colon. 2. A severe gripping pain in the bowels, due to spasm of the intestinal walls; also any severe spasmodic pain in the abdomen. **C., Biliary**, that due to the passage of a gall-stone through the gall ducts. **C., Devonshire.** Synonym of *C., Lead*. **C., Hepatic.** See *C., Biliary*. **C., Lead**, or *C., Saturnine*, *Colica pictorum*, *painter's colic*. Intestinal colic due to lead-poisoning. It is characterized by excruciating abdominal pain, a hard and retracted condition of the abdomen, slow pulse, and increased arterial tension. **C., Menstrual**, the pain of menstruation. **C., Renal**, due to the presence of a calculus in the ureter. **C., Uterine**, colic-like pains experienced at the menstrual epochs, often coming on in paroxysms.

Colitis (*ko-lit'-tis*) [*κόλον*, the large intestine; *itis*, inflammation]. Inflammation of the colon.

Collagen (*kol'-aj-en*) [*κόλλα*, glue; *γεννάειν*,

to produce]. A substance existing in various tissues of the body, especially bone and cartilage; it is converted into gelatin by boiling.

Collapse (*kol-aps'*) [*collabere*, to fall together]. Extreme depression and prostration from failure of the circulation, as in cholera, shock, hemorrhage, etc. **C. of Lung**, return of a portion or the whole of a lung to its fetal or airless condition from some mechanic hindrance to the entrance of air. It is characterized by dyspnea, with more or less cyanosis, and is mainly encountered in bronchopneumonia.

Collapsing (*kol-aps'-ing*) [*collabere*, to fall together]. Suddenly breaking down. **C. Pulse**. See *Corrigan's Pulse*.

Collar-bone. The clavicle.

Collateral (*kol-at'-er-al'*) [*con*, together; *lateralis*, of the side]. Accessory or secondary; not direct or immediate.

Collecting Tubes of the Kidney. A name given to the ducts discharging into the calices of the kidneys.

Colles' Fascia. The deep layer of the perineal fascia. It is attached to the base of the triangular ligament, to the anterior lips of the rami of the pubes and ischia laterally, and anteriorly it is continuous with the dartos of the scrotum. **C. Fracture**. See *Diseases, Table of*. **C. Law**. See *Law*.

Collidin (*kol'-id-in*) [*κόλλα*, glue], $C_8H_{11}N$. A ptomain, isomeric but not identical with aldehyd-collidin. The ptomain was obtained from pancreas and gelatin allowed to putrefy together in water.

Collier's Lung. Synonym of *Anthraxis*.

Collinsonia (*kol-in-so'-ne-ah*) [after Peter Collinson, an English gentleman]. A genus of labiate herbs. **C. canadensis**. Stoneroot, heal-all, a coarse plant with a disagreeable smell; it has tonic, diuretic, and diaphoretic properties. Dose gr. xv-lx (1.0-4.0) in decoction; of the fluid extract $\mathfrak{m}\text{x}$ - $\mathfrak{f}\mathfrak{z}$ (0.65-4.0); of the tincture (1 in 10) $\mathfrak{f}\mathfrak{z}$ ss-ij (2.0-8.0).

Colliquation (*kol-ik-wa'-shun*) [*con*, together; *liqueare*, to melt]. The liquefaction or breaking down of a tissue or organ.

Colliquative (*kol-ik-wa-tiv*) [*con*, together; *liqueare*, to melt]. Profuse or excessive; marked by excessive fluid discharges. **C. Diarrhea**, a profuse watery diarrhea. **C. Necrosis**. See *Necrosis, Liquefaction*. **C. Sweat**, a profuse clammy sweat.

Colliquefaction (*kol-ik-we-fak'-shun*) [*con*, together; *liqueare*, to melt]. A melting, or fusing together.

Collodion (*kol-o'-de-on*) [*κολλώδης*, glue-like]. See *Pyroxilin*.

Colloid (*kol'-oid*) [*κόλλα*, glue]. 1. A nondializable organic substance. See *Dialysis*. 2. A substance formed by colloid degenera-

tion of epithelium. See *Degeneration*. 3. Having the nature of glue. 4. In chemistry, amorphous and non-crystalline. **C. Cancer**. See *Cancer*. **C. Degeneration**. See *Degeneration*. **C. Degeneration of the Skin**, a rare disease, occurring chiefly on the upper part of the face, in the form of small, glistening, translucent, flattish elevations, and yielding by pressure a small, jelly-like mass.

Colloma (*kol-o'-mah*) [*κόλλα*, glue; *δυμα*, a tumor]. A cystic tumor containing a gelatiniform substance.

Collum (*kol'-um*) [*L.*]. The neck; especially the anterior part of the neck. **C. distortum**. Synonym of *Torticollis*.

Collutory (*kol'-u-to-re*) [*collutorium*, *colluere*, to rinse]. A gargle or mouth-wash.

Collyrium (*kol ir'-e-um*) [*κόλλυριον*, an eye-salve]. A lotion for the eyes.

Coloboma (*kol-o-bo'-mah*) [*κολοβόειν*, to mutilate]. A congenital fissure of the iris, choroid, or eyelids.

Colocynthis (*kol-o-sin'-this*) [*κολοκυνθίς*, *colocynth*]. *Colocynth*. The fruit of *C. citrullus*, from which the seeds and rind have been removed. Its properties are due to a bitter glucosid, colocynthin, $C_{26}H_{34}O_{13}$, the dose of which is gr. $\frac{1}{2}$ - $\frac{1}{3}$ (0.003-0.013). It is a tonic and astringent purgative, and is used mainly as an ingredient of compound cathartic pills. **C., Ext.**, alcoholic. Dose gr. ss-ij (0.032-0.13). **C., Ext., Comp.**, contains colocynth extract 16, aloes 50, cardamom 6, resin of scammony 14, soap 14, alcohol 10 parts. Dose gr. v-xx (0.32-1.3). **C., Pilula, Comp.** (B. P.), contains colocynth, aloes, scammony, potassium sulphate and oil of cloves. Dose gr. v-x (0.32-0.65). **Pilulæ Catharticæ Comp.**, compound cathartic pills; contain each, comp. ext. of colocynth gr. 1.3, abstract jalap gr. j, calomel gr. j, gamboge gr. $\frac{1}{4}$. Dose 1-3 pills. **Pilula Colocynthis et Hyoscyam.** Dose gr. v-x (0.32-0.65).

Coloenteritis (*kol-o-en-ter-i'-tis*) [*κόλον*, colon; *εντερον*, an intestine; *ιτις*, inflammation]. Inflammation of the small and large intestine. See *Enterocolitis*.

Colomba, Colombo (*kol-am'-bah*, *kol-am'-bo*). See *Calumba*.

Colon (*kol'-lon*) [*κόλον*, the colon]. The part of the large intestine beginning at the cecum and terminating at the end of the sigmoid flexure. In the various parts of its course it is known as the ascending C., the transverse C., the descending C., and the sigmoid flexure.

Colony (*kol'-o-ne*) [*colonia*, a colony]. A collection or assemblage, as of microorganisms in a culture.

Colophony Resin (*kol'-o-fo-ne*) [*κολοφών*, a city of Ionia]. Rosin. The solid residue

left on distilling off the volatile oil from crude turpentine. See *Resin*.

Color-blindness. See *Blindness*.

Color-hearing. The excitation of the visual center for color through the auditory nerve.

Color-sensation. The perception of color; it depends on the number of vibrations of the ether.

Colorimeter (*kull-or-im'-et-er*) [*color*, *color*; *μέτρον*, measure]. An instrument for determining the quantity of coloring matter in a mixture, as in the blood.

Colostomy (*ko-lost'-to-me*) [*κόλον*, colon; *στόμα*, opening]. 1. The formation of an artificial anus by an opening into the colon. 2. Any surgical operation upon the colon that makes a permanent opening into it, whether internal or external.

Colostrum (*kol-os'-trum*) [L.]. The first milk from the mother's breasts after the birth of the child. It is laxative, and assists in the expulsion of the meconium. **C. Corpuscles**, small microscopic bodies contained in the colostrum. They are the epithelial cells of the mammary glands, full of oil-globules. After about the third day these cells burst and set free the fat-globules before they leave the gland, and in this way the true milk is formed.

Colotomy (*ko-lot'-o-me*) [*κόλον*, colon; *τέμνειν*, to cut]. Incision of the colon, either abdominal, lateral, lumbar, or iliac, according to the region of entrance.

Colpeurynter (*kol-pu-rin'-ter*) [*κόλπος*, vagina; *εὐρύνειν*, to widen]. An inflatable bag or sac used for dilating the vagina.

Colpitis (*kol-pi'-tis*) [*κόλπος*, vagina; *ιτις*, inflammation]. Inflammation of the vagina.

Colpocele (*kol'-po-sel*, or *-se'-le*) [*κόλπος*, vagina; *κλή*, tumor]. Hernia of the vagina.

Colpocleisis (*kol-po-kli'-sis*) [*κόλπος*, vagina; *κλείσις*, a closure]. The surgical closure of the vagina.

Colpohysterectomy (*kol-po-his-ter-ek'-to-me*) [*κόλπος*, vagina; *ὑστέρα*, womb; *ἐκτομή*, excision]. Removal of the uterus through the vagina.

Colpoperineorrhaphy (*kol-po-per-in-e-or'-af-e*) [*κόλπος*, vagina; *perineum*; *ράφή*, seam]. Repair of a perineal laceration by denuding and in part suturing the posterior wall of the vagina.

Colpoptosis (*kol-pop-to'-sis*) [*κόλπος*, vagina; *πτώσις*, a falling]. Prolapse of the vaginal walls.

Colporrhaphy (*kol-por'-a-fe*) [*κόλπος*, vagina; *ῥαφή*, seam]. Suture of the vagina.

Coltsfoot (*kolts'-foot*). The leaves of *Tussilago farfara*, a demulcent and tonic, sometimes prescribed for chronic cough. Dose of a decoction (℥j to Oj) a teacupful; of fld. ext. f℥j-ij (4.0-8.0). Unof.

Columbin (*kol-um'-bin*). An active principle of *Calumba*. It is a colorless crystalline substance of excessively bitter taste and neutral reaction.

Columbo (*kol-um'-bo*). See *Calumbo*.

Columella (*kol-u-mel'-ah*) [L., a little column]. The modiolus or central axis of the cochlea of the human ear. See *Cochlea* and *Modiolus*.

Column (*kol'-um*) [*columna*, a column]. A name given to several parts of the body that furnish support to surrounding parts, or that have the shape of pillars. **C.**, Anterior, the layer of white matter in either half of the spinal cord included between the anterior horn and nerve-roots, and the anterior median fissure. **C.**, Anterolateral Ascending. See *C. of Gowers*. **C. of Burdach**. See *C.*, *Postero-external*. **C. of Clarke**, a column of gray substance occupying the region to the outer and posterior side of the central canal of the spinal cord, at the inner part of the base of the posterior cornu. It contains fusiform cells, and is the trophic center for the direct cerebellar tract. **C.**, Direct Cerebellar, in the spinal cord, is situated outside of the lateral pyramidal tract. **C. of Goll**, the posteromedian column of the spinal cord. **C.**, Gowers', in the spinal cord, a group of fibers lying in front of the direct cerebellar tract and reaching around upon the periphery of the anterior pyramidal tract. **C.**, Lateral, the layer of white matter in either half of the spinal cord included between the posterior horn and nerve-roots, and the anterior horn and nerve-roots. **C. of Morgagni**, vertical folds of the rectal mucous membrane seen at the point of union of the latter with the skin of the anus. **C.**, Posterior, a collection of white matter situated in the spinal cord on either side between the posterior horns and posterior nerve-roots, and the posterior median fissure. **C.**, *Postero-external*, the outer wider division of the posterior column of the cord; the column of Burdach. **C.**, *Posteromedian*, the median division of the posterior column of the cord; the column of Goll. **C. of Spitzka-Lissauer**, in the spinal cord, a group of nerve-fibers lying just in front of and just behind the posterior horns. **C.**, *Postero-vesicular*. See *Column of Clark*. **C. of Türck**, the anterior or direct pyramidal tract. **C. of the External Ring**, the free borders of the aponeurosis of the external oblique muscle, forming the edges of the external abdominal ring.

Columna (*kol-um'-nah*) [L.]. A column or pillar. **C. Bertini**, that part of the cortical structure of the kidneys that separates the sides of any two pyramids. **Columnæ car-**

nae, the muscular columns projecting from the inner surface of the ventricles of the heart.

Coma (*ko'-mah*) [*kāpa*, deep sleep]. Unconsciousness from which the patient cannot be aroused by external stimulus. **C.**, **Alcoholic**, that due to poisoning by alcohol. **C.**, **Apoplectic**, that due to apoplexy. **C.**, **Uremic**, that due to uremia. **C.** **Vigil**, a comatose condition in which the patient lies with open eyes, but unconscious and delirious. This occurs occasionally in typhoid and typhus fevers and in delirium tremens.

Comatose (*ko'-mat-ās*) [*kāpa*, a deep sleep]. In a condition of coma.

Combustion (*kom-bus'-chun*) [*comburare*, to burn up]. The process of oxidation, attended with the liberation of heat and sometimes of light. It is loosely used as a synonym of inflammation.

Comedo (*kom'-e-dō*) [*comedo*, a glutton: *pl.*, *Comedones*]. Grubs; Grub-worms; Black-heads. A chronic disorder of the sebaceous glands characterized by yellowish or whitish pin-point and pin-head-sized elevations, containing in their center exposed blackish points. They are found usually on the face, back of the neck, chest, and back, and are often associated with acne. They occur as a rule in the young. Occasionally a parasite, the *Demodex folliculorum*, is found in each comedo.

Comes (*ko'-mēs*) [*L.*: *pl.*, *Comites*]. A companion, as a vein to an artery, or an artery to a nerve.

Comma-bacillus (*kom'-ah-bas-il'-us*). See *Bacteria*, *Table of*.

Commensal (*kom-en'-sal*) [*cum*, together; *mensa*, table]. In biology, the harmonious living together of two animals or plants. One of two such organisms.

Comminuted (*kom'-in-u-ted*) [*comminuere*, to break in pieces]. Broken into a number of pieces.

Commissure, Commissura (*kom'-iss-ar*, *kom-iss-u'-rah*) [*cum*, together; *mittere*, to send]. That which unites two parts. **C.**, **Anterior** (*of third ventricle*), a rounded cord of white fibers placed in front of the anterior crura of the fornix. **C.**, **Anterior White** (*of spinal cord*), a layer of fibers separating the posterior gray commissure from the bottom of the anterior median fissure. **C.**, **Gray** (*of spinal cord*), the transverse band of gray matter connecting the masses of gray matter of the two halves of the spinal cord. **C.** **magna**, the corpus callosum. **C.**, **Middle**, a band of soft gray matter connecting the optic thalami. **C.**, **Optic**, the union and crossing of the two optic nerves in front of the tuber cinereum. **C.**, **Posterior** (*of third ventricle*), a flattened

white band connecting the optic thalami posteriorly. **C.**, **Posterior White** (*of spinal cord*), a band of fibers separating the gray commissure from the bottom of the posterior median fissure.

Commissural (*kom-iss-u'-ral*) [*cum*, together; *mittere*, to send]. Having the properties of a commissure; uniting symmetric parts, as **C.** fibers of the brain.

Commotio (*kom-o'-she-o*) [*L.*]. A commotion or shock. **C.** **cerebri**, concussion of the brain.

Communicans (*kom-u'-nik-ans*) [*L.*]. Communicating. **C.** **noni**. See *Nerves*, *Table of*.

Commutator (*kom'-u-ta-tor*) [*commutare*, to exchange]. An instrument for automatically interrupting or reversing the flow of an electric current.

Compact Tissue. The external, hard part of bone.

Comparative Anatomy. See *Anatomy*.

Compatibility (*kom-pat-ib-il'-it-e*) [*compati*, to suffer with]. Of medicines, the relation of one substance to another, so that they may be mixed without chemic change or loss of therapeutic power.

Compensating (*kom'-pen-sa-tiŋ*) [*compensare*, to compensate]. Making good a deficiency. **C.** **Ocular**. See *Ocular*. **C.** **Operation**, in ophthalmology, tenotomy of the associated antagonist in cases of diplopia from paresis of one of the ocular muscles.

Compensation (*kom-pen-sa'-shun*) [*compensare*, to equalize]. The act of making good a deficiency; the state of counter-balancing a functional or structural defect.

Compensatory (*kom-pen'-sa-tō-re*) [*compensare*, to equalize]. Making good a deficiency. Restoring the balance after failure of one organ or part of an organ by means of some other organ or part of an organ.

Complemental, Complementary (*kom-ple-men'-tal*, *kom-ple-men'-tō-re*) [*cum*, together; *plere*, to fill]. Supplying a deficiency. **C.** **Air**. See *Air*. **C.** **Colors**, a term applied to any two colors which combined produce white light, as, *e. g.*, blue and yellow.

Complexus (*kom-pleks'-us*) [*complexus*, complex]. The totality of symptoms, phenomena, or signs of a morbid condition. **C.** **Muscle**. See *Muscles*, *Table of*.

Complication (*kom-plik-a'-shun*) [*complicare*, to fold together]. A disease occurring in the course of some other disease and more or less dependent upon it.

Compos Mentis (*kom'-pos men'-tis*) [*L.*]. Of sound mind.

Composite (*kom-pos'-it*) [*componere*, to put together]. 1. Composed of distinct portions. 2. In biology, belonging to the order *Compositae*.

Compound (*kom'-pound*, or *kom-pound'*)

[*cum*, together; *ponere*, to put]. 1. To mix, as drugs. 2. Composed of several parts. **C.**, **Cathartic Pills.** See *Colocynth*. **C. Fracture.** See *Fracture*.

Compress (*kom'-pres*) [*compressus*, pressed together]. Folded cloths, wet or dry, applied firmly to the part for relief of inflammation, or to prevent hemorrhage. **C. Graduated**, a compress composed of folds of a gradually increasing size.

Compression (*kom'-pres'-un*) [*compressus*, pressed together]. The state of being compressed. **C.-atrophy**, atrophy of a part from constant compression. **C.-myelitis**, myelitis due to compression of the spinal cord.

Compressor (*kom'-pres'-or*) [*comprimere*, to press together]. 1. An instrument for compressing an artery, vein, etc. 2. The term is also applied to muscles having a compressing function. **C. sacculi laryngis**, the inferior arytenoepiglottideus muscle. See *Muscles*, *Table of*. **C. urethræ.** See *Muscles*, *Table of*.

Conarium (*ko-na'-re-um*) [*κωνάριον*; dim. of *κωνος*, a cone]. The pineal gland.

Concave (*kon-kāv'*) [*cum*, together; *cavus*, hollow]. Hollow; incurved, as the inner surface of a hollow sphere.

Concavoconvex (*kon-kāh'-vo-kon'-veks*). Having one surface concave, the other convex, the convexity exceeding the concavity.

Conceive (*kon-sev'*) [*concipere*, to take in]. To become pregnant.

Concentration (*kon-sen-tra'-shun*) [*cum*, together; *centrum*, the center]. The act of making denser, as of a mixture, by evaporating a part of the liquid.

Concentric (*kon-sen'-trik*) [*cum*, together; *centrum*, the center]. Arranged in an equidistant manner about a center. **C. Hypertrophy of the Heart**, increase in the muscular texture of the heart, the capacity of the cavities remaining unchanged.

Conception (*kon-sep'-shun*) [*concipere*, to conceive]. 1. The fecundation of the ovum by the spermatozoid. 2. The abstract mental idea of anything; the power or act of mentally conceiving. **C., Imperative**, a false idea that a person dwells upon and cannot expel from his mind, even when he knows it to be absurd. It dominates his actions, and is a symptom of insanity.

Concha (*hong'-kah*) [*κόχχα*, a shell]. A shell. Applied to organs having some resemblance to a shell, as the patella, vulva, etc., and especially to **C. auris**, or the hollow part of the external ear.

Conchoscope (*hong'-ko-skōp*) [*κόχχα*, a shell; the *naris*; *σκοπεῖν*, to inspect]. A speculum and mirror for inspecting the nasal cavity.

Concomitant (*kon-kom'-it-ant*) [*concomitari*, to accompany]. Accompanying. **C. Strabismus.** See *Strabismus*. **C. Symptoms**, symptoms that are not in themselves essential to the course of a disease, but that may occur in association with the essential symptoms.

Concretion (*kon-kret'-shun*) [*concretio*; *concrecere*, to grow together]. The solidification or condensation of a fluid substance; a calculus; a union of parts normally separate, as the fingers.

Concussion (*kon-kush'-un*) [*concussio*, a violent shock]. Shock; the state of being shaken; a severe shaking or jarring of a part; also, the morbid state resulting from such a jarring. **C. of Brain**, a condition produced by a fall or blow on the head, and marked by unconsciousness, feeble pulse, cold skin, pallor, at times the involuntary discharge of feces and urine; this is followed by partial stupor, vomiting, and headache, and eventually recovery. In severe cases inflammation of the brain may follow, or a condition of feeble-mindedness. **C. of Spinal Cord**, a condition caused by severe shock of the spinal column, with or without appreciable lesion of the cord. It leads to functional disturbances analogous to railway-spine.

Condensed (*kon-densd'*) [*condensare*, to make thick]. Made compact; reduced to a denser form. **C. Milk**, milk that has had most of its watery elements evaporated. Condensed milk prepared with the addition of cane-sugar is a white or yellowish-white product of about the consistency of honey and ranging in specific gravity from 1.25 to 1.41. It should be completely soluble in from four to five times its bulk of water, without separation of any flocculent residue, and then possess the taste of fresh, sweetened milk. Condensed milk prepared without the addition of cane-sugar is not boiled down to the same degree, and therefore remains liquid.

Condenser (*kon-den'-ser*) [*condensare*, to make dense]. A lens or combination of lenses used in microscopy for gathering and concentrating rays of light.

Conduction (*kon-duk'-shun*) [*conductio*; *conducere*, to draw together]. The passage or transfer of force or material from one part to another.

Conductor (*kon-duk'-tor*) [*conducere*, to draw together]. 1. A body that transmits force-vibrations, such as those of heat or electricity. 2. A term applied to the electrodes and cords by which they are joined to the battery. 3. An instrument serving as a guide for the surgeon's knife.

Condurango Bark (*kon-du-rang'-go bark*) [native of S. Amer.]. Bark of *Gonolobus*

tetragonus, or of *Conglobus condurango*; a remedy much used in S. America as an alterative in syphilis. It was introduced into the U. S. as a remedy for carcinoma of the stomach, but yielded uncertain results. It is a stomachic tonic. Dose of the fluid extract f3 ss-j (2.0-4.0); of the tincture, f3j-ij (4.0-8.0). Unof.

Condy's Fluid. One part of sodium or potassium permanganate dissolved in 500 parts of water; it is a useful disinfectant.

Condylarthrosis (*kon-dil-ar-thro'-sis*) [*κόνδυλος*, a knuckle; *άρθρον*, a joint]. A form of diarthrosis, wherein a condyle is set in a shallow and elliptic cavity, and free and varied movement of the joint is possible; condylar articulation.

Condyle (*kon'-dil*) [*κόνδυλος*, a knuckle]. Any rounded eminence such as occurs in the joints of many of the bones, especially the femur, humerus, and lower jaw.

Condylloid (*kon'-dil-oid*) [*κόνδυλος*, a knuckle; *ειδος*, likeness]. Resembling or pertaining to a condyle.

Condyloma (*kon-dil-oid'-mah*) [*κονδυλωμα*, a swelling]. A wart-like growth or tumor usually near the anus or pudendum. **C. latum**, the syphilitic mucous patch.

Cone, Conus (*kón, kó'-nus*) [*κónος*, a cone]. A solid body having a circle for its base, and terminating in a point. **C. of Light**, the triangular reflection from the normal membrana tympani. **C., Retinal**, one of the rod-like bodies, which, with the associated rods, form one of the outer layers of the retina, the so-called rod-and-cone layer.

Confection (*kon-fek'-shun*) [*confectio*, a preparation]. In pharmacy, a mass of sugar and water, or of honey, used as an excipient with a prescribed medicinal substance.

Confinement (*kon-fín'-ment*) [*cum*, together; *finis*, boundary, limit]. The condition of women during childbirth.

Confluent (*kon'-flu-ent*) [*confluere*, to flow together]. Running together. The opposite of discrete. In anatomy, coalesced or blended; applied to two or more bones originally separate, but subsequently formed into one.

Confrontation (*kon-frun'-ta'-shun*) [*confrontari*, to be contiguous to]. The examination of a person by whom a diseased person may have been infected as a means of diagnosing the disease in the latter.

Confusion (*kon-fu'-shun*) [*cum*, together; *fundere*, to pour]. Mixing; confounding. **C. Colors**, a set of colors so chosen that they cannot be distinguished by one who is color-blind. **C. Letters**, test-type letters, such as C. G. O., or F. P. T., liable to be mistaken for each other.

Congelation (*kon-jel'-a'-shun*) [*congelatio*, a

freezing]. 1. Freezing; frost-bite; intense cold or its effect on the animal economy, or any organ or part. 2. The chilling or benumbing effect of any freezing-mixture or application; mainly employed for its local anesthetic effect. 3. Coagulation.

Congenital (*kon-jen'-it-al*) [*cum*, together; *gigni*, to be born]. Existing at birth.

Congestion (*kon-jes'-chun*) [*congerere*, to heap up]. An abnormal collection of blood in a part or organ. Congestion may be active or passive, atonic or inflammatory, functional or hypostatic. It is also named from the parts affected; the most important varieties of morbid congestion are the cerebral, spinal, pulmonary, hepatic, and renal.

Congestive (*kon-jes'-tiv*) [*congerere*, to heap up]. Marked by, due to, or of the nature of congestion.

Congius (*kon'-je-us*) [L.]. A Roman measure. A gallon.

Conglomerate (*kon-glom'-er-at*) [*conglomerare*, to heap up]. 1. Massed together; aggregated. 2. A mass of units without order.

C. Glands, acinous glands.

Conglutin (*kon-glu'-tin*) [*cum*, together; *gluten*, glue]. One of the proteids found in peas, beans, and other kinds of pulse.

Conglutination (*kon-glu-tin-a'-shun*) [*conglutinare*, to glue together]. The abnormal union of two contiguous surfaces or bodies, as of two fingers, or of the opposed surfaces of the pleural or pericardial sac.

Congo-red. A red coloring-matter which becomes blue in the presence of free HCl. It is used in chemic investigation of the gastric juice.

Conic, Conical (*kon'-ik, kon'-ik-al*) [*κωνικός*, pertaining to a cone]. Cone-shaped. **C. Cornea.** See *Keratoglobus*.

Coniin (*ko-ni'-in*). See *Conium* and *Conin*.

Conin (*kó'-nin*) [*κόνειον*, hemlock]. $C_8H_{10}N$. A liquid alkaloid which is the active principle of *Conium*.

Conium (*ko-ni'-um*) [*κόνιον*, hemlock]. Hemlock. The leaves and the fruit of the spotted hemlock, *C. maculatum*. It contains three alkaloids and a volatile oil. Its properties are mainly due to the alkaloids coniin, $C_8H_{13}N$, and methylconiin, $C_8H_{14}NCH_3$. It produces motor-paralysis, without loss of sensation or of consciousness. Toxic doses cause death by paralysis of the organs of respiration. It is valuable in acute mania, delirium tremens, tetanus, blepharospasm, asthma, and whooping-cough. **C., Abstract**, made from conium, 200, dilute hydrochloric acid 6, sugar of milk and alcohol q. s. to make 100 parts of abstract. Dose gr. ss-ij (0.032-0.2). **C., Cataplasma** (B. P.), made from the leaves; for external use. **C., Ext.** Dose gr. ss-ij (0.032-0.13). **C., Ext. Fld.**,

Dose m-jij (0.065-0.13). C., *Pilula*, *Comp.* (B. P.), contains extract of hemlock and ipecac. Dose gr. v-x (0.32-0.65). C., *Succus* (B. P.), made from the leaves. Dose mxxx-3ij (2.0-8.0). C., *Tinct.*, 15 per cent strength. Dose m-x-f3j (0.65-40). C. *Vapor* (B. P.), for inhalations. *Coniin*, volatile alkaloid of conium. Dose gr. $\frac{1}{16}$ - $\frac{1}{8}$ (0.001-0.006). Unof. *Coniin Hydrobrom.*, $C_8H_9(C_2H_5)$, recommended for spasmodic affections. Dose gr. $\frac{1}{16}$ - $\frac{1}{8}$ (0.002-0.004). Unof. All preparations are uncertain in strength.

Coni vasculosi. A series of conic masses of tubules that together form the globus major of the epididymis.

Conjugal (*kon'-ju-gal'*) [*cum*, together; *jugare*, to yoke]. Pertaining to marriage; connubial. C. *Diabetes*, diabetes affecting husband and wife together; this is said to be not infrequently observed.

Conjugate (*kon'-ju-gāt'*) [*cum*, together; *jugare*, to yoke]. Yoked or coupled. C.

Deviation. See *Deviation*. C. *Diameter* (of the pelvis). See *Diameter*. C., *True*, the minimum diameter of the pelvic inlet.

Conjunctiva (*kon-junk-ti'-vah*) [*conjunctivus*, connecting]. The mucous membrane covering the anterior portion of the globe of the eye, reflected upon the lids and extending to their free edges. Its parts are called palpebral and ocular, or bulbar.

Conjunctivitis (*kon-junk-tiv-i'-tis*) [*conjunctivus*, connecting; *itis*, inflammation]. Inflammation of the conjunctiva. C., *Acute Catarrhal*, the most common form, usually mild, resulting from cold or irritation. C., *Croupous*, a variety associated with the formation of a whitish-gray membrane that is easily removed. C., *Diphtheric*, a specific purulent inflammation of the conjunctiva, due to the Klebs-Löffler bacillus. C., *Egyptian*, See *Trachoma*. C., *Follicular*, a form characterized by numerous round, pinkish bodies found in the retrolarsal fold. C., *Gonorrheal*, a severe form of purulent C., caused by infection with gonococci. C., *Phlyctenular*, one characterized by the presence on the ocular conjunctiva of small vesicles surrounded by a reddened zone. C., *Purulent*, one characterized by a thick, creamy discharge.

Connective (*kon-ek'-tiv*) [*connectere*, to connect]. C. *Tissue*. See *Tissue*.

Conoid, or Conoidal (*ko'-noid*, or *ko-noi'-dal'*) [*κωνος*, cone; *ειδος*, shape]. Of a conic shape. C. *Ligament*, the lower and inner part of the coracoclavicular ligament. C. *Tubercle*, the eminence on the inferior surface of the clavicle to which the conoid ligament is attached.

Conquinamin (*kon-kwin'-a-min*). An alkaloid of *Cuprea* bark.

Consanguinity (*kon-san-gwin'-it-e*) [*cum*, together; *sanguinis*, of blood]. The relationship arising from common parentage; blood-relationship.

Consciousness (*kon'-shus-nes*) [*consci-us*, knowing]. The state of being aware of one's own existence, of one's own mental states, and of the impressions made upon one's senses; ability to take cognizance of sensations. C., *Double*, that morbid condition in which there are two separate and alternating states of mental consciousness, in either one of which the events that have occurred in the other state are not remembered by the patient.

Consensual (*kon-sen'-su-al'*) [*consensus*, agreement]. Excited reflexly by stimulation of another part, usually a fellow organ, as the C. reaction of one pupil when the iris of the other eye is stimulated.

Conservation (*kon-ser-va'-shun*) [*conservare*, to keep]. Preservation without loss. C., of *Energy*. See *Energy*.

Consolidation (*kon-sol-id-a'-shun*) [*consolidare*, to make firm]. The process of becoming firm or solid, as a lung in pneumonia.

Constant (*kon'-stant'*) [*constans*, steady]. Fixed. C. *Current*, one that goes continuously in one direction.

Constipation (*kon-stip-a'-shun*) [*constipare*, to crush tightly together]. A condition in which the bowels are evacuated at long intervals or with difficulty.

Constitution (*kon-stit-u'-shun*) [*constituere*, to dispose]. In chemistry, the atomic or molecular composition of a body. In pharmacy, the composition of a substance. In physiology, the general temperament and functional condition of the body.

Constitutional (*kon-stit-u'-shun-al'*) [*constituere*, to dispose]. Pertaining to the state of the constitution. C. *Diseases*, such diseases as are inherent, owing to an abnormal structure of the body. Also, a condition in which the disease pervades the whole system. General diseases, in contradistinction to local.

Constrictor (*kon-strik'-tor*) [*constringere*, to bind together]. Any muscle that contracts or tightens any part of the body. See *Muscles*, *Table of*.

Consumption (*kon-sump'-shun*) [*consumere*, to consume or wear away]. A wasting away, especially a wasting disease like tuberculosis, particularly pulmonary tuberculosis or tuberculosis of the bowels.

Contact (*kon'-takt*) [*contactus*, a touching]. A touching. C. *Breaker*, an instrument by means of which a galvanic circuit is broken.

Contagion (*kon-ta'-jun*) [*contagio*; *contingere*, to touch]. 1. The process by which a specific disease is communicated from one person to another, either by direct contact or by means of an intermediate agent. 2. Also

the specific germ or virus from which a communicable disease develops.

Contagious (*kon-ta'-jus*) [*contagium*, a touching]. Communicable or transmissible by contagion, or by a specific contagium.

Contagium (*kon-ta'-je-um*) [L.]. Any virus or morbid matter by means of which a communicable disease is transmitted from the sick to the well.

Contiguity (*kon-tig'-u'-it-e*) [*cum*, together; *tangere*, to touch]. Proximity. **C.**, **Amputation in the**, is performed at a joint, without section of a bone.

Continenence (*kon'-ti-nens*) [*continere*, to hold together]. Self-restraint, especially in regard to the sexual passion.

Continued (*kon'-tin'-u-d*) [*continuare*, to make continuous]. Persisted in. **C.**, **Fever**, a fever that is long-continued, without intermissions.

Continuity (*kon-tin'-u'-it-e*) [*cum*, together; *tenere*, to hold]. The state of being continuous or uninterrupted. **C.**, **Amputation in the**, amputation in which a bone is divided.

Contractility (*kon-trak-til'-it-e*) [*contrahere*, to draw together]. That property of certain tissues, especially muscle, of shortening upon the application of a stimulus.

Contraction (*kon-trak'-shun*) [*contrahere*, to draw together]. Approximation of the elements of a tissue or organ, thus diminishing its volume or contents. **C.**, **Anodal, Closing or Opening**, the contraction taking place at the anode on closing or opening the circuit. **C.**, **Carpopedal**, a variety of tetany occurring in infants, and generally associated with dentition or seat-worms. There is a flexing of the fingers, toes, elbows, and knees, and a general tendency to convulsions. **C.**, **Closing**, muscular contractions produced at the instant that the electric current is closed. **C.**, **Dupuytren's**, a contraction of the palmar fascia causing the fingers to fold into the palm. **C.**, **Front-tap**, a phenomenon often observed in cases with exaggerated knee-jerk. When the foot is placed at a right angle to the leg, and the muscles of the front of the leg are tapped, the foot is extended. See *Reflexes*, *Table of*. **C.**, **Hour-glass**, a contraction of an organ, as the stomach or uterus, at the middle. **C.**, **Opening**, the muscular contraction produced by opening or breaking the circuit. **C.**, **Paradoxic**, a phenomenon that consists in the contraction of a muscle, caused by the passive approximation of its extremities. **C.**, **remainder**, the stage of elastic after-vibration or residual contraction persisting in a muscle after withdrawal of the stimulus. **C.**, **ring**, the boundary line between the upper and lower segments of the parturient uterus.

Contracture (*kon-trak'-chur*) [*cum*, together; *trahere*, to draw]. Contraction; permanent shortening, as of a muscle; distortion or deformity due to the shortening of a muscle or of various muscles.

Contraindication (*kon-trah-in-dik-a'-shun*) [*contra*, against; *indicare*, to point out]. That modifying condition in which a remedy or a method of treatment is forbidden.

Contrecoup (*kön'-tr-koo*) [Fr.]. Counter-stroke. The transmission of a shock from the point struck to a point on the opposite side of the body or the part.

Control (*kon-tröl'*) [*contra*, against; *rotula*, a roll]. A standard by which to check observations and insure the validity of their results. Colloquially, the term is sometimes used as a noun for control animal, or control experiment. **C.**, **Animal**, one used in a control experiment. **C.**, **Experiment**, an experiment carried out under normal or common circumstances or conditions, to serve as a standard whereby to test the variation or value of another experiment carried out under peculiar or abnormal circumstances.

Contusion (*kon-tu'-shun*) [*contundere*, to bruise]. A bruise or injury inflicted without the integument being broken. **C.**, **pneumonia**, a form of pneumonia following traumatism.

Conus (*ko'-nus*) [*kónos*, a cone]. 1. A cone. 2. A crescentic patch of atrophic choroid tissue near the optic papilla in myopia. **C.**, **arteriosus**, the cone-shaped eminence of the right ventricle of the heart, whence arises the pulmonary artery. **C.**, **medullaris**, the cone-like termination of the spinal cord, continuous as the filum terminale. **Coni vasculosi**, a series of conic masses that together form the globus major of the epididymis.

Convalescence (*kon-val-es'-ens*) [*convalescere*, to become well]. A term applied to the restoration to health after disease.

Convallamarin (*kon-val-am'-ar-in*) [*convallis*, a valley; *amarus*, bitter], $C_{23}H_{44}O_{12}$. A glucosid derived from *Convallaria*. See *Convallaria majalis*.

Convallaria (*kon-val-a'-re-ah*) [*convallis*, a valley]. A genus of liliaceous plants. **C.**, **majalis**, lily of the valley. All parts of the plant are used in medicine. Its properties are due to convallarin, $C_{24}H_{40}O_{11}$, and convallamarin, $C_{23}H_{44}O_{12}$, glucosids. It is a cathartic, diuretic, and cardiac stimulant. **C.**, **Ext.**, soluble in water. Dose gr. ij-x (0.13-0.65). **C.**, **Ext.**, **Fld.**, alcoholic. Dose \mathfrak{m} ij-xj (0.13-0.7). **C.**, **Infusum**, prepared with three times its weight of water. Dose \mathfrak{f} $\frac{3}{4}$ ss-ij (16.0-64.0). **Convallamarin**, or **Convallamarinum**, is soluble in water. Dose gr. $\frac{1}{4}$ -ij (0.016-0.13). All unof.

Convallarin (*kon-val'-ar-in*) [*convallis*, a

valley], $C_{34}H_{62}O_{11}$. A crystalline purgative glucosid derived from *Convallaria*.

Convection (*kon-vek'-shun*) [*convehere*, to carry together]. A transmission or carrying, as of heat or electricity. **C.-current**, a current of a liquid or gas heated to a temperature above that of the surrounding medium; it rises to the surface because of its lesser density and thus the entire fluid or gas acquires the same temperature.

Convergence (*kon-ver'-jens*) [*cum*, together; *vergere*, to incline]. Inclination or direction toward a common point, center, or focus, as of the axes of vision upon the near-point. **C.-stimulus** Adduction, the power of adduction of the eyes provoked by fixation of the gaze upon an object placed at the near-point.

Convergent (*kon-ver'-jent*) [*cum*, together; *vergere*, to incline]. Tending to a common center. **C. Strabismus**. See *Strabismus*.

Convex (*kon-veks'*) [*convexus*, vaulted]. Having a surface approximating more or less a part of the surface of a sphere.

Convexity (*kon-veks'-it-e*) [*convexus*, convex]. A surface rounding outward; the quality of being convex.

Conexoconcave (*kon-veks'-o-kon-kav*). See *Lens*.

Convolution (*kon-vo-lu'-shun*) [*convolvere*, to roll together]. A fold, twist, or coil of any organ, especially any one of the prominent convex parts of the brain, separated from each other by depressions or sulci. **C., Angular**, the posterior part of a convolution situated between the intraparietal fissure in front and above, and the horizontal limb of the Sylvian fissure, and the hinder part of the first part of the first temporal fissure below. The anterior part is called the supramarginal convolution. **C., Annectant**, small convolutions which connect the occipital with the temporosphenoidal and parietal lobes. **C., Anterior Central**, **C., Ascending Frontal**, the convolution in front of the fissure of Rolando. **C., Ascending Parietal**, the convolution just behind the fissure of Rolando. **C., Broca's**, the inferior or third frontal convolution. **C., Fornicate**, a long convolution on the mesial surface of the brain above the corpus callosum. **C., Frontal**, the convolutions of the frontal lobe. **C., Hippocampal**, the part of the fornicate convolution that winds around the splenium of the corpus callosum. **C., Insular**, the small convolutions composing the island of Reil. **C., Marginal**, the mesial surface of the first frontal convolution. **C., Occipital**, the convolutions making up the occipital lobe. **C., Paracentral**, a convolution on the mesial surface of the brain, representing the junction of the upper ends of the ascending frontal and ascending parie-

tal convolutions. **C., Parietal**, the convolutions of the parietal lobe. **C., Posterior Central**. See *C., Ascending Parietal*. **C., Supramarginal**. See *C., Angular*. **C., Temporal**, the convolutions of the temporal lobe. **C., Uncinate**, the hook-like termination of the fornicate convolution.

Convolvulin (*kon-vol'-yu-lin*) [*convolvere*, to roll together], $C_{31}H_{50}O_{16}$. A glucosid derived from the roots of Jalap (*Convolvulus purga*). It is a gummy mass, with active purgative properties.

Convulsant (*kon-vull'-sant*) [*convellere*, to convulse]. A medicine that causes convulsions.

Convulsion (*kon-vull'-shun*) [*convellere*, to convulse]. An involuntary general paroxysm of muscular contraction. It is either tonic (without relaxation) or clonic (having alternate contractions of opposite groups of muscles). **C., Epileptiform**, one characterized by total loss of consciousness. **C., Hysteric**, one due to hysteria. Consciousness is only apparently lost. **C., Tetanic**, general tonic convulsions without loss of consciousness. **C., Uremic**, one that occurs in kidney disease due to retention in the blood of matters that should be eliminated by the kidney.

Coordination (*ko-or-din'-a'-shun*) [*cum*, together; *ordinare*, to regulate]. The harmonious activity and proper sequence of operation of those parts that cooperate in the performance of any function.

Copaiba (*ko-pa'-e-bah*) [Sp.]. Balsam of Copaiba. The oleoresin of *Copaifera officinalis*, *C. coriacea*, *C. guianensis*, *C. multi-juga*, *C. cordifolia*, *C. laxa*, *C. nitida*, *C. oblongifolia*, and *C. langsdorffii*, leguminous trees, native to South America. It is a stimulant, diuretic, diaphoretic, and an expectorant, and is much used in gonorrhea. **C., Massa**, copaiba 94, magnesia 6 parts. Dose gr. x-3j (0.65-4.0). **C., Mist.**, **Comp.**, Lafayette's mixture, unof., copaiba 3 vij, oil of cubebs 3j, glyceritum vitelli 3 vij, triturate and add syr. aq. menth. piperit. 3 liiss; then add, with constant stirring, liq. potass. 3 ss, tinct. cardamom comp. 3 ij, spt. æther. nitrosi, 3 ss, aq. menth. piperit. q. s. to make 3 viij. Dose f 3 j-f 3 ss (4.0-16.0). **C., Oleum**. Dose ℥ x-xv (0.65-1.0). **C., Resina**, mainly copaibic acid. Dose gr. j-v (0.065-0.30).

Copiopia (*kop-i-o'-pe-ah*) [*κόπος*, a straining; *ὤψ*, eye]. Eye-strain; weariness of the eyes. **C. hysterica**, a term applied to those symptoms that indicate hyperesthesia of the fifth and optic nerves.

Copper (*kop'-er*) [ME., *coper*, copper]. Cuprum. A reddish-brown metal existing in nature chiefly in the form of copper pyrites, which is a double salt of copper and iron

sulphid. Various salts are used in medicine. In toxic doses they are gastrointestinal irritants. In therapeutic doses they are used as astringents in inflammation of mucous membranes. They are also employed as emetics, and, externally, as caustics. See *Elements, Table of*. **C. Acetate**, $\text{Cu}(\text{C}_2\text{H}_3\text{O}_2)_2$, verdigris, used in pulmonary diseases, and as a lotion in skin-diseases. Dose gr. $\frac{1}{10}$ - $\frac{1}{4}$ (0.0065-0.016). **C. Acetoarsenite**, Paris Green, used as a pigment and an insecticide. **C. Ammoniate**, unof., ammonium carbonate 3, copper sulphate 4 parts, useful in chorea, hysteria, etc. Dose gr. $\frac{1}{6}$ - $\frac{1}{2}$ (0.01-0.065). **C. Arsenite**, a salt valuable in intestinal diseases. Dose gr. $\frac{1}{100}$ (0.00065). **C. Nitrate** (B. P.), $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$, is used for the same purposes as the sulphate. **C. sulphate**, $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, soluble in water, valuable as an emetic, tonic and astringent. Dose, as an emetic, gr. ij-v (0.13-0.32), as a tonic, gr. $\frac{1}{6}$ - $\frac{1}{2}$ (0.01-0.032). **C. Amalgam**, a metallic filling-material composed of copper and mercury. **C.-nose**. Synonym of *Aene rosacea*.

Copperas (*kop'-er-as*) [*cupri rosa*, rose of copper (?)]. A common name for ferrous sulphate.

Copremia (*kop'-re'-me-ah*) [*κόπρος*, dung; *αἷμα*, blood]. A form of general blood poisoning arising from chronic constipation. The symptoms are anemia, sallow complexion, anorexia, frontal headache, vertigo, nausea, flatulence, thirst, fetid breath, lassitude, hypochondriasis, and irritability of temper.

Coprolalia (*kop-ro-lá'-le-ah*) [*κόπρος*, filth; *λάλησις*, speech]. The use of filthy and offensive language as a manifestation of disease.

Coprolith (*kop'-ro-lith*) [*κόπρος*, dung; *λίθος*, stone]. A hard mass of fecal matter in the bowels.

Coprophagy (*kop-roff'-a-je*) [*κόπρος*, dung; *φαγεῖν*, to eat]. The eating of dung, a symptom seen in insane and hysteric patients.

Coprostasis (*kop-ras'-tas-is*) [*κόπρος*, dung; *στάσις*, a standing]. The accumulation of fecal matter in the bowel.

Coptis (*kop'-tis*) [*κόπτειν*, to cut]. Goldthread. The root of *C. trifolia*, a simple bitter tonic resembling quassia. It contains coptin, an alkaloid closely allied to berberin. Dose gr. x-xxx (0.65-2.0).

Copulation (*kop-u-lá'-shun*) [*copulare*, to couple]. The act of sexual intercourse.

Cot (*kor*) [L.: *gen.*, *cordis*]. The heart. See *Heart*. **C. adiposum**, a heart with a simple excess of the normal subpericardial fat. **C. bovinum**. See *Bovine Heart*. **C. villosum**, hairy heart; the peculiar shaggy appearance presented by the heart in acute plastic pericarditis, with the deposited fibrin existing in long shreds.

Coraco- (*kor'-ak-o-*) [*κόραξ*, a crow]. Pertaining to the coracoid process. **C.-Brachialis**. See *Muscles, Table of*.

Coracoid (*kor'-ak-oid*) [*κόραξ*, a crow; *ειδος*, likeness]. 1. Having the shape of a crow's beak. 2. The coracoid process. **C. Ligament**, a triangular ligament joining the coracoid process to the acromion. **C. Process**, a beak-shaped process of the scapula.

Cord [*chorda*, a string]. 1. Any string-like body. 2. Used as a synonym for the *Umbilical Cord*, the vascular, cord-like structure connecting the placenta and fetus. **C., False or Superior (vocal)**, a fold of mucous membrane on either side of the middle line of the larynx, inclosing the superior thyroarytenoid ligament. **C., Presentation of**, descent of the umbilical cord between the presenting part and the membranes, at the beginning of labor. **C., Prolapse of**, descent of the umbilical cord at the rupture of the bag of waters; incomplete, if remaining in the vagina, complete, if protruding therefrom. **C., Spinal**. See *Spinal Cord*. **C., True Vocal**; **C., Vocal**. See *Vocal Band*.

Cordial (*kor'-jal*) [*cor*, the heart]. 1. Pertaining to the heart; exhilarant; stimulant. 2. An aromatic, spirituous stimulant.

Core (*kor*) [*cor*, heart]. The central slough of a boil or carbuncle.

Corectopia (*kor-ek-tó'-pe-ah*) [*κόρη*, pupil; *ἐκτοπος*, misplaced]. An anomalous position of the pupil; displacement of the pupil.

Coredialysis (*ko-re-di-ál'-is-is*) [*κόρη*, pupil; *ἀνάλειψιν*, to liberate]. The production of an artificial pupil at the ciliary border of the iris.

Corelysis (*kor-el'-is-is*) [*κόρη*, pupil; *λίσις*, a loosening]. The detachment of iritic adhesions to the lens, or to the cornea.

Coremorphosis (*kor-e-mor-fó'-sis*) [*κόρη*, pupil; *μόρφωσις*, formation]. The operation for establishing an artificial pupil.

Corcometer (*kor-e-om'-et-er*) [*κόρη*, pupil; *μέτρον*, a measure]. An instrument for measuring the pupil of the eye.

Coriander, or **Coriandrum** (*ko-re-an'-der*, *ko-re-an'-drum*) [*κορίανδρον*, coriander]. Coriander. The fruit of *C. sativum*, an aromatic, carminative, and stimulant, used mainly to give flavor to other remedies and as a corrective to griping purgatives. Dose gr. x-xx (0.65-1.3). **C., Oleum**, the volatile oil. Dose ℥ ij-v (0.13-0.32).

Corium (*kor'-re-um*) [L., leather]. The deep layer of the skin.

Corm (*korm*) [*κόρυμβος*, the trunk of a tree]. The bulbous underground part of certain plants, as the crocus.

Corn [*cornu*, horn]. A local induration and thickening of the skin from friction or pressure. See *Clavus*. **C.-silk**. See *Zea Mays*.

C.-smut. See *Ustilago*. **C.-starch**, the commercial name of a starch derived from maize, and extensively used as an article of food, especially for invalids.

Cornea (*kor'-ne-ah*) [*corneus*, horny]. The transparent anterior portion of the eyeball, its area occupying about one-sixth the circumference of the globe. It is continuous with the sclerotic, and is nourished by lymph from the looped blood-vessels at its peripheral border. **C.**, **Conic.** See *Keratoglobus*. **C.**, **Transplantation of**, the operation of engrafting a section of transparent cornea from some animal into the space of an excised portion of human cornea.

Corneal (*kor'-ne-al*) [*corneus*, horny]. Relating to the cornea.

Corneitis (*kor-ne-i'-tis*). See *Keratitis*.

Corneous (*kor'-ne-us*) [*corneus*, horny]. Horny, or horn-like. **C. Tissue**, the substance of the nails.

Corniculum (*kor-nik'-u-lum*) [*cornicula*, a little horn]. A small cornu or horn-like process.

Cornification (*kor-nif-ik-a'-shun*) [*corneus*, horny; *facere*, to make]. The process of hardening or making horny.

Cornu (*kor'-nu*) [*L.*]. A horn. A name applied to any excrescence resembling a horn. **C. ammonis**, the hippocampus major of the brain. **C. cervi**, hartshorn or ammonium hydroxid. **C. cutaneum**, cornu humanum, a horn-like excrescence arising from the skin.

Cornual (*kor'-nu-al*) [*cornu*, a horn]. Relating to a cornu. **C. Myelitis**, myelitis affecting the anterior cornua of the spinal cord.

Cornus (*kor'-nus*) [*L.*]. Dogwood. The bark of the root of *C. florida*, the properties of which are due to a crystalline principle, cornin. It is a simple stomachic bitter and slightly antiperiodic. **C.**, **Ext. Fld.** Dose ℥x-fʒj (0.65-4.0).

Corona (*ko-ro'-nah*) [*corona*, a garland]. A crown. **C. glandis**, the ridge of the glans penis. **C. radiata**, a radiating mass of white nerve-fibers ascending from the internal capsule to the cortex cerebri. **C. veneris**, a circle of syphilitic blotches occurring on the forehead.

Coronal (*kor-o'-nal*) [*corona*, the crown]. Encircling like a crown; pertaining to the crown of the head. **C. Suture**, the suture joining the frontal with the two parietal bones.

Coronary (*kor'-o-na-re*) [*corona*, a crown]. A term applied to vessels, nerves, or attachments that encircle a part or organ.

Coroner (*kor'-o-ner*) [*corona*, a crown]. An officer who inquires by authority of the law into the causes of sudden or violent deaths. **C.**, **Inquest of**, the legal inquiry before a jury into the cause of a sudden or violent death.

Coronoid (*kor'-o-noid*) [*corona*, crown; *eidos*, likeness]. Crown-shaped; as the **C. process** of the ulna or of the jaw.

Corpora (*kor'-por-ah*) [*pl. of corpus*, a body]. A general term applied to certain parts of the body having a rounded or oval shape. **C. albicantia**, two white masses in the interpeduncular space at the base of the brain, the projections of the anterior pillars of the fornix. **C. geniculata**, two small eminences projecting from the optic thalami. **C. quadrigemina**, the optic lobes of the brain, the four rounded eminences situated under the corpus callosum. The anterior pair are called the nates, and the posterior, the testes.

Corpulency (*kor'-pu-len-se*) [*corpulentus*, corpulent]. Obesity; fatness of the body.

Corpus (*kor'-pus*) [*corpus*, a body: *pl.*, *Corpora*]. A body; the human body. **C. Arantii**, the fibrous tubercle in the center of each segment of the semilunar valves. **C. callosum**, the broad band of white matter uniting the hemispheres of the cerebrum. **C. dentatum**. 1. See *Olivary Body*. 2. The central folded gray nucleus of the cerebellum. **C. highmorianum**. See *Highmore, Body of*. **C. fimbriatum**, the lateral thin edge of the tenia hippocampi. **C. luteum**, the yellow body formed in the ovary in the site of a Graafian vesicle after the escape of the ovum. **C. luteum, False**, that resulting when pregnancy does not occur, called also the **C. L. of Menstruation**. **C. luteum, True**, that resulting when pregnancy takes place, called also the **C. L. of Pregnancy**. **C. spongiosum**, the spongy part of the penis encircling the urethra. **C. striatum**, a mass of gray matter extending into the lateral ventricles of the brain and composed of the caudate and lenticular nuclei.

Corpuscle (*kor'-pus-l*) [*dim. of corpus*]. A small body or particle. **C.**, **Bizzozero's**. See *Blood-platelets*. **C.**, **Red, of Blood**, biconcave, non-nucleated discs, circular in outline, and containing red coloring-matter, termed hemoglobin, to which the color of the blood is due. Red corpuscles have been divided, according to their size, into normocytes (normal in size), megalocytes (of excessive size), microcytes (abnormally small), and poikilocytes (of irregular shape and size). The red corpuscles in the blood of man are about $\frac{1}{2500}$ in. in diameter and $\frac{1}{1000}$ in. thick, and their number is about five millions to each cubic millimeter of blood. They consist of a colorless stroma infiltrated with the coloring-matter (hemoglobin). **C.**, **White or colorless**, flattened cells, about $\frac{1}{3500}$ in. in diameter, existing in the ratio of 1:500 compared with red corpuscles. Their protoplasm is granular, they have one or more nuclei and no cell-wall. They pos-

sess contractile power and alter their shape readily. The colorless corpuscles are variously designated as eosinophile, basophile, neutrophile, mononuclear, polynuclear, lymphocytes, transitional, large, small, etc. **C. of Donne.** See *Colostrum-corpuscles*. **C., Malpighian**, of the spleen, a name applied to the lymphoid nodules of the spleen. **C., Malpighian**, of the kidney, the tuft of blood-vessels surrounded by the expanded portion of the uriniferous tubule, the capsule of Bowman. **C., Norris's Invisible**, colorless, transparent, biconcave discs of the same size as the red corpuscles, invisible in the serum because their color and refractive index are the same as those of the liquor sanguinis. **C., Pacinian**, certain small corpuscles occurring in the subcutaneous cellular tissue of the fingers and toes. They consist of concentric lamellæ of connective tissue surrounding the termination of a sensory nerve. **C., Tactile**, of Wagner, the small, oval bodies found in the papillæ of the skin and enveloped by nerve-fibers.

Corpuscular (*kor-pus'-ku-lar*) [*corpusculum*, a little body]. Relating to or of the nature of a corpuscle.

Correctant, or Corrective (*kor-ek'-tant, kor-ek'-tiv*) [*corrigere*, to correct]. 1. Modifying favorably. 2. A substance used to modify or make more pleasant the action of a purgative or other remedy.

Corrigan's Disease. See *Diseases, Table of*. **C. Pulse**, water-hammer pulse. The jerking pulse of aortic regurgitation.

Corrigent (*kor'-ij-ent*). See *Corrective*.

Corrosion (*kor-d'-zhun*) [*cum*, together; *rodere*, to gnaw]. The process of corroding, or the state of being corroded. **C.-anatomy**, that branch of anatomy which demonstrates an anatomic specimen by means of a corrosive process that eats away those parts which it is not desired to preserve. In some cases a resisting-substance is injected, so as to preserve the vessels and ducts from corrosion. **C.-preparation**, one in which the vessels, ducts, or cavities of organs are filled by a fluid that will harden and preserve the shape of the vessel or cavity after the organ itself is corroded or digested or otherwise destroyed.

Corrosive (*kor-d'-ziv*) [*cum*, together; *rodere*, to gnaw]. 1. Eating away. 2. A substance that destroys organic tissue either by direct chemic means or by causing inflammation and suppuration. **C. Chlorid**, **C. Sublimite**, mercuric chlorid. See *Hydrargyrum*.

Corrugator (*kor'-u-ga-tor*) [*corrugere*, to wrinkle]. That which wrinkles. See *Muscles, Table of*.

Cortex (*kor'-tek-s*) [*L.*, bark]. 1. The bark of an exogenous plant. 2. The surface-layer of an organ. **C. aurantii**, orange-peel.

C. cerebri, the external layer of gray matter of the brain. **C. renalis**, the cortical substance of the kidney.

Corti's Arch; **C. Canal**; **C. Organ.** See *Ear*.

Cortical (*kor'-tik-al*) [*cortex*, bark]. Pertaining to the cortex or bark, or to the cortex of the brain, or of the kidney. **C. Epilepsy**, **C. Paralysis**, such as is due to a lesion of the cortical substance of the brain.

Coryza (*kor-i'-zah*) [*κόρυς, κόρυζα*, the head]. Catarrh of the mucous membrane of the nasal passages and adjacent sinuses, popularly called "cold in the head." See *Rhinitis*.

Cosmetic (*kos-met'-ik*) [*κοσμητικός, κοσμεῖν*, to adorn]. 1. Beautifying. 2. A remedy designed to hide defects of the skin or other external parts. **C. Operation**, a surgical operation to give a natural appearance to a defective or unsightly part.

Cosmolin (*koz'-mo-lin*). See *Petrolatum*.

Costa (*kos'-tah*) [*L.*]. A rib.

Costal (*kos'-tal*) [*costa*, a rib]. Pertaining to the ribs. **C. Arch**, the arch of the ribs.

C. Cartilages, the 12 cartilaginous extensions of the ribs.

Costive (*kos'-tiv*) [*constipare*, to be bound]. Constipated.

Costiveness (*kos'-tiv-ness*) [*constipare*, to be bound]. Constipation.

Costotome (*kos'-to-tom*) [*costa*, rib; *τομή*, cutting]. A strong knife or heavy shears with the under blade in the shape of a hook, for cutting the costal cartilages in dissection, etc.

Coto (*ko'-to*) [*Sp.*, a cubit]. Coto bark. The bark of a tree native to Bolivia. It contains a bitter principle, cotoin, $C_{22}H_{18}O_6$, irritant to the skin and mucous membranes. It is recommended for diarrhea and zymotic fevers, and for the night-sweats of pulmonary tuberculosis. Dose of the powder, gr. j-xv (0.065-1.0); of the fluid extract, gr. v-xv (0.32-1.0); of the tincture (1 in 10), gr. xxx (0.65-2.0); of cotoin, gr. ss-j (0.032-0.065).

Cotton (*kol'-n*) [*Arab. qutun*, cotton]. Gossypium, a white fibrous seed-hair that envelops the seeds of the cotton-plant. **C., Absorbent**, cotton so prepared that it readily absorbs water. **C.-root**. See *Gossypium*. **C.-seed Oil** or **C.-oil**, oleum gossypii seminis, an oil obtained by pressure from the bulled seeds of several species of *Gossypium*.

Cotunnus, Liqueur, or Water of. The perilymph. See *Aqua labyrinthi*. **C., Nerve of**, the nasopalatine nerve.

Cotyledon (*kol'-il'-el'-don*) [*κοτυληδών*, a socket]. Any one of the enlarged, vascular villi of the chorion which project into depressions of the decidua vera.

Cotyloid (*kol'-il'-oid*) [*κοτύλη*, a cup; *ειδος*,

form]. Cup-shape. **C. Fossa**, or Cavity, the acetabulum. **C. Ligament**, a ligament surrounding the acetabulum. **C. Notch**, a notch in the anterior and lower border of the acetabulum.

Couch-grass (*kowch'-gras*). See *Triticum*.

Couching (*kowch'-ing*) [Fr., *coucher*, to depress]. The operation, now fallen into disuse, of depressing a cataractous lens into the vitreous chamber, where it was left to be absorbed.

Cough (*kof*) [ME., *cough*, a cough]. A sudden, violent expulsion of air after deep inspiration and closure of the glottis. **C.**, Dry, that unattended by expectoration. **C.**, Ear-, cough excited reflexly from some morbid condition of the ear. **C.**, Moist, cough with free expectoration. **C.**, Reflex, cough produced by irritation of a remote organ, as **C.**, Ear, and **C.**, Stomach. **C.**, Stomach-. See **C.**, Reflex. **C.**, Winter-, a short troublesome cough of old people due to chronic bronchitis, and recurring every winter.

Coulomb (*koo-lom'*) [after *Coulomb*, a French physicist]. The unit of measurement of electric quantity; the quantity of electricity that passes during one second through a conductor having a resistance of one ohm, with one volt of electromotive force. The micro-coulomb is the millionth part of this amount.

Coumarin (*koo'-mar-in*), $C_9H_6O_2$. A vegetable proximate principle that occurs in *Asperula odorata*, in the Tonka bean, and in *Melilotus officinalis*. It conceals the odor of iodoform. Unof.

Counterextension (*koun'-ter-eks-ten-shun*). See *Extension*.

Counterindication (*koun'-ter-in-dik-a-shun*). See *Contraindication*.

Counterirritation (*koun'-ter-ir-it-a-shun*) [*contra*, against; *irritare*, to irritate]. Superficial inflammation produced artificially, in order to exercise a good effect upon some adjacent or deep-seated morbid process.

Counteropening (*koun'-ter-o-pen-ing*) [*contra*, against; AS., *open*, open]. An incision made in an abscess or cavity, opposite to another, generally for purposes of drainage.

Counterpoison (*koun'-ter-poi-zn*) [*contra*, against; *potio*, a drink]. A poison given as an antidote to another poison.

Counterstroke (*koun'-ter-strök*). See *Contre-coup*.

Coup de soleil (*koo-duh-so-lay'*) [Fr.]. Sun-stroke.

Court-plaster (*kört'-plas-ter*). See *Ichthyocolla*.

Cover (*kuv'-er*) [*cum*, together; *operire*, to shut]. **C.-glass**, in microscopy, the thin slip of glass covering the object mounted on the slide.

Cowage (*kow'-aj*) [Hind., *kawanch*, cow-

hage]. The external hairs of the pod of *Mucuna pruriens*, used in medicine as a mechanic vermifuge.

Cowhage, **Cowitch** (*kow'-aj*, *kow'-ich*). See *Mucuna* and *Cowage*.

Cowper's Glands. See *Gland*.

Cowperitis (*kow-per-i'-tis*) [*Cowper*, an English anatomist; *itis*, inflammation]. Inflammation of the glands of Cowper, usually gonorrheal in origin.

Cow-pock, a variety of pock seen in the cow, and thought to correspond with smallpox in man.

Coxa (*koks'-ah*) [*coxa*, hip]. The hip-joint or the hip.

Coxalgia (*koks-al'-je-ah*) [*coxa*, a hip; *âλγος*, pain]. Literally pain in the hip-joint, but generally used synonymously with hip-disease.

Coxalgic (*koks-al'-jik*) [*coxa*, the hip; *âλγος*, pain]. Relating to coxalgia.

Coxe's Hive-mixture. See *Scilla* and *Antimonium*.

Coxitis (*koks-i'-tis*) [*coxa*, the hip; *itis*, inflammation]. Inflammation of the hip-joint.

Coxofemoral (*koks-o-fem'-or-al*) [*coxa*, the hip; *femur*, the thigh-bone]. Relating to the hip and the femur, as the *C. joint*, the hip-joint.

Crab-louse (*krab'-lous*). See *Pediculus*.

Crab's-eyes (*krab'-iz*). Flat, calcareous concretions (*Lapides cancerorum*) derived from the stomach of the crab; they have been used as a means of removing foreign bodies from the eye. Also a name for the seeds of *Abrus precatorius*.

Cracked-pot Sound. A peculiar sound elicited by percussion over a pulmonary cavity communicating with a bronchus.

Cradle (*kra'-dl*) [AS., *cradol*, a cradle]. In surgery, a wire or wicker-frame so arranged as to keep the weight of the bed-clothing from an injured part of the body. It is employed in the treatment of fractures, wounds, etc. **C.**, Ice-, the suspension over a febrile patient, by means of iron frames, of a number of buckets, kept half-filled with ice, and enclosed in a light covering.

Cramp (*kramp*) [Teut., *krampf*]. A spasmodic tonic contraction of a muscle, attended with sharp pain. **C.**, Professional, spasm of certain groups of muscles, from their continuous use in different occupations, as Writer's **C.**, Hammerman's **C.**, Piano-player's **C.**, Dancer's **C.**, etc.

Crane's-bill Root. See *Geranium*.

Cranial (*kra'-ne-al*) [*κρᾶνιον*, the skull]. Relating to the cranium.

Craniectomy (*kra'-ne-ek'-to-me*) [*κρᾶνιον*, skull; *ἐκτομή*, a cutting out]. The surgical removal of strips or pieces of the cranial bones. It is performed in cases of microcephaly.

Cranioclast (*kra'-ne-o-klast*) [*κρᾱνίον*, skull; *κλάειν*, to break]. A heavy forceps for crushing the fetal head.

Craniology (*kra-ne-ol'-o-je*) [*κρᾱνίον*, cranium; *λόγος*, science]. A branch of anatomy comprising the study of skulls.

Craniometer (*kra-ne-om'-et-er*) [*κρᾱνίον*, skull; *μέτρον*, a measure]. An instrument for measuring the dimensions of the skull.

Craniometric, Craniometrical (*kra-ne-o-met'-rik*, *kra-ne-o-met'-rik-al*) [*κρᾱνίον*, skull; *μέτρον*, a measure]. Pertaining to craniometry. **C. Point**, any one of the points of measurement used in craniometry. A list of the craniometric points is given.

ACANTHION, a point in the median line of the skull at the base of the nasal spine. **ALVEOLAR POINT**, the point between the two middle incisors of the upper jaw. **ANTINION**, that point on the glabellum, and in the median line, that is farthest from theinion. **ASTERION**, the point behind the ear where the parietal, temporal and occipital bones meet.

AURICULAR POINT, the center of the orifice of the external auditory meatus. **BASION**, the middle point of the anterior margin of the foramen magnum. **BREGMA**, the point where the coronal and sagittal sutures meet.

DACRYON, or **DAKRYON**, the point beside the root of the nose where the frontal, lacrymal, and superior maxillary bones meet. **ENTOMION**, the point where the parietal notch of the temporal bone receives the anterior extension of the mastoid angle of the parietal bone. **GLABELLA**, or **GLABELLUM**, the point in the median line between the superciliary arches, marked by a swelling, sometimes by a depression. **GNATHION**, same as Mental Point. **GONION**, the point at the angle of the lower jaw. **HORMION**, the anterior point of the basilar portion of the united sphenoccipital bone, where it is crossed by the median line. **INION**, the external occipital protuberance. **JUGAL POINT**, the point situated at the angle that the posterior border of the frontal branch of the malar bone makes with the superior border of its zygomatic branch. **KORONION**, the apex of the coronoid process of the inferior maxilla. **LAMBDA**, the point of meeting of the sagittal and the lambdoid sutures. **MALAR POINT**, a point situated on the tubercle on the external surface of the malar bone, or at the intersection of a line drawn from the external extremity of the frontomalar suture to the tubercle at the inferior angle of the malar bone and a line drawn nearly horizontally from the inferior border of the orbit over the malar bone to the superior border of the zygomatic arch. **MAXIMUM OCCIPITAL POINT**, or **OCCIPITAL POINT**, the posterior extremity of the anteroposterior diameter of

the skull, measured from the glabella in front to the most distant point behind. **MENTAL POINT**, the middle point of the anterior lip of the lower border of the lower jaw. **METOPION**, or **METOPIC POINT**, a point in the middle line between the two frontal eminences. **NASION**, or **NASAL POINT**, the middle of the frontal suture at the root of the nose. **OBELION**, the part of the sagittal suture between the two parietal foramina. **OCCIPITAL POINT**. See *Maximum Occipital Point*. **OPHYRYON**, the middle of the supraorbital line, which, drawn across the narrowest part of the forehead, separates the face from the cranium. **OPISTHION**, the middle point of the posterior border of the foramen magnum. **PROSTHION**, the alveolar point. **PTERION**, the point where the frontal, parietal, temporal, and sphenoid bones come together. **RHINION**, the upper median point of the anterior nasal opening. **SPINAL POINT**. Same as Subnasal Point. **STEPHANION, INFERIOR**, the point where the ridge for the temporal muscle intersects the coronal suture. **STEPHANION, SUPERIOR**, the point where the coronal suture crosses the temporal ridge. **SUBNASAL POINT**, the middle of the inferior border of the anterior nares at the base of the nasal spine. **SUPRA-AURICULAR POINT**, the point vertically over the auricular point at the root of the zygomatic process. **SUPRANASAL POINT**. Same as Ophryon. **SUPRAORBITAL POINT**. Same as Ophryon. **SYMPHYSION**, the median point of the outer border of the alveolus of the lower jaw. **VERTEX**, the superior point of the skull. In obstetrics, that conic portion of the skull the apex of which is at the posterior fontanel and the base of which is formed by the biparietal and trachelobregmatic diameters.

Craniometry (*kra-ne-om'-et-re*) [*κρᾱνίον*, skull; *μέτρον*, a measure]. The ascertainment of the proportions and measurements of skulls.

Craniotabes (*kra-ne-o-la'-bez*) [*κρᾱνίον*, skull; *ταβέειν*, to waste away]. An atrophy of the cranial bones occurring in infancy, with the formation of small, shallow, conic pits in the bone-substance. Craniotabes results from rachitis, syphilis, or marasmus.

Craniotomy (*kra-ne-ol'-o-me*) [*κρᾱνίον*, skull; *τομή*, a cutting]. The operation of reducing the size of the fetal head by cutting or breaking it up, when delivery is otherwise impossible. **C., Linear**. See *Cranietomy*.

Craniotympanic (*kra-ne-o-tim-pan'-ik*) [*κρᾱνίον*, the skull; *tympanum*]. Pertaining to the skull and the tympanum.

Cranium (*kra'-ne-um*) [*κρᾱνίον*, the skull]. The skull. The cavity that contains the brain, its membranes and vessels.

Crassamentum (*kras-am-ent-tum*) [L., thickness]. A clot, as of blood.

Cravat (*kra-vat'*) [Fr., *cravate*]. A bandage of triangular shape, used as a temporary dressing for a wound or fracture. The middle is applied to the injured part, and the ends are brought around and tied.

Cream (*krem*) [*cremor*, thick juice or broth]. The rich, fat part of milk. **C.**, Cold. See *Rosa*. **C.** of Tartar. See *Potassium*.

Crease (*kres*) [*Celtic*]. A line made by folding. **C.**, Gluteofemoral. See *C.*, *Ileo-femoral*. **C.**, Ileo-femoral, is the crease that bounds the buttock below, corresponding nearly to the lower edge of the gluteus maximus muscle. It is of supposed significance in the diagnosis of hip-disease.

Creasol (*kre'-as-ol'*) [*κρεας*, flesh; *oleum*, oil], $C_8H_{10}O_2$. One of the principal phenols contained in creosote. It is formed from guaiacum-resin, and is found in beechwood-tar. It is a colorless, oily liquid of an agreeable odor and a burning taste, boiling at 226° C. It is very similar to guaiacol.

Creosote (*kre'-o-sot*). See *Creosote*.

Creatin (*kre-at-in*) [*κρεας*, flesh], $C_4H_7N_3O_2$. A neutral organic substance that occurs in the animal organism, especially in the juice of muscles.

Creatinin (*kre-at'-in-in*) [*κρεας*, flesh], $C_4H_7N_3O$. An alkaline substance, a normal constituent of urine. It crystallizes in rhombic prisms, and is a strong base. It is much more soluble than creatin.

Crede's Method (*kre-da*). A method of expelling the placenta by grasping the uterus firmly through the abdominal walls, kneading it to excite contraction, and then pressing downward toward the sacrum.

Cremaster (*kre-mas'-ter*) [*κρεμειν*, to support]. The muscle that draws up the testis. See *Muscles*, Table of.

Cremasteric (*kre-mas-ter'-ik*) [*κρεμειν*, to support]. Pertaining to the cremaster muscle. **C.** Reflex. See *Reflexes*, Table of.

Cremation (*kre-ma'-shun*) [*cremare*, to burn]. The destruction of the body by burning, as distinguished from interment.

Cremor (*kre'-mor*) [*cremor*, broth]. Cream. Any thick substance formed on the surface of a liquid. **C.** tartari, cream of tartar.

Crenation (*kre-na'-shun*) [*crena*, a notch]. A notched or mulberry-like appearance of the red corpuscles of the blood. It is seen when they are exposed to the air or strong saline solutions.

Crenothrix (*kren'-oth-riks*) [*κρίννυ*, a spring; *θρίξ*, hair]. A genus of Schizomycetes the filaments of which are enveloped in a gelatinous sheath.

Creolin (*kre'-o-lin*) [*κρεας*, flesh; *oleum*, oil]. A coal-tar product deprived of phenol.

It is an antiseptic used especially as a douche in obstetric practice. It has also been used in a solution of 5 : 1000 for irrigation of the bowel in dysentery and enterocolitis.

Creosote, or **Creosotum** (*kre'-os-ot, ot-tum*) [*κρεας*, flesh; *σῴζειν*, to preserve]. The product of the distillation of wood-tar, consisting of a mixture of phenol-compounds. It is an inflammable oily liquid, differing in this respect from phenol. It does not coagulate albumin or collodion. Most of the commercial creosote consists of phenol or contains a large percentage of it. It is valuable for its antiseptic, astringent, styptic, anesthetic, and escharotic properties. It is used extensively in pulmonary tuberculosis. Dose \mathfrak{m} j-ij (0.065-0.2). **Beechwood Creosote** is best for internal use. Dose \mathfrak{m} j (0.065). **C.**, **Aqua**, a one per cent. solution. Dose $\mathfrak{f}\mathfrak{v}$ -iv (4.0-16.0). **C.**, **Mistura** (B.P.). Dose $\mathfrak{f}\mathfrak{v}$ j-ij (4.0-8.0). **C.**, **Unguentum** (B.P.), for local application. **C.**, **Vapor** (B.P.), for inhalation.

Crepitant (*krept'-it-ant*) [*crepitare*, to crackle]. Possessing the character of crepitus. **C.** Râle. See *Râle* and *Breath-sounds*.

Crepitation, **Crepitus** (*krept'-it-ant-shun, krept'-it-us*) [*crepitare*, to crackle]. The grating of fractured bones. The crackling of the joints. The noise produced by pressure upon tissues containing an abnormal amount of air or gas, as in cellular emphysema. Also the sound heard at the end of inspiration in the first stage of croupous pneumonia. It closely resembles the sound produced by rubbing the hair between the fingers held close to the ear. **C.**, **Redux**, a crepitant râle heard in pneumonia during the stage of resolution; usually the first manifestation of the recession of the disease.

Crescent (*kres'-ent*) [*crescere*, to grow]. 1. Sickie-shaped, or shaped like the new moon. 2. A name given to one form of the malarial hematooon. **C.** of *Gianuzzi*, groups of deeply staining cells in the acinus of a gland, pushed to one side by the secreting cells, and probably representing exhausted cells. **C.** **Myopic**. See *Myopia*.

Crezol (*kre'-sol*) [*κρεας*, flesh; *oleum*, oil], $C_8H_8O_3$. Cresylic acid; a body obtained from the distillation of coal-tar. It is a colorless, caustic liquid, with properties similar to those of phenol, but is superior as an antiseptic. Unof.

Crest (*krest*) [*crista*, a crest]. A ridge or linear prominence, especially of bone. **C.**, **Frontal**, a ridge along the middle line of the internal surface of the frontal bone. **C.** of **Ilium**, the thickened and expanded upper border of the ilium. **C.**, **Lacrymal**, a vertical ridge dividing the external surface of the lacrymal bone into two parts. **C.**, **Nasal**,

a crest on the internal border of the nasal bone and forming part of the septum of the nose. **C.**, **Occipital**, a vertical ridge on the external surface of the occipital bone extending from the occipital protuberance to the foramen magnum. **C. of Pubes**, a crest extending from the spine to the inner extremity of the pubes. **C. of Tibia**, the prominent border or ridge on the front of the tibia; the shin. **C.**, **Turbinated**, a prominent horizontal ridge on the internal surface of the palate bone.

Creta (*kre'-tah*) [*L.*]. Chalk. Native calcium carbonate. **C.**, **Mistura**, consists of *C.*, pulv. cretæ. comp. 20, cinnamon-water 40, water 40. It is used in diarrhea. Dose [$\frac{3}{4}$ ss (16.0). **C. preparata**, chalk freed from impurities by washing. Dose gr. v-xx (0.32-1.3). **C.**, **Pulv.**, **Aromat.** (*B. P.*). Dose gr. x- $\frac{3}{4}$ j (0.65-4.0). **C.**, **Pulv.**, **Comp.**, compound chalk-powder, consists of *C.* prep. 30, acacia 20, sugar 50. Dose gr. v- $\frac{3}{4}$ j (0.32-4.0). **C.**, **Trochisci**, each contains prepared chalk gr. iv, acacia gr. j, sugar gr. vj, with a little nutmeg.

Cretin (*kre'-tin*) [*Fr.*, *crétin*, a simple-minded person]. A person affected with cretinism.

Cretinism (*kre'-tin-izm*) [*Fr.*, *crétin*, a simple-minded person]. A congenital disease, characterized by absence of the thyroid gland, diminutiveness of size, thickness of neck, shortness of arms and legs, prominence of abdomen, large size of face, thickness of lips, large and protruding tongue, and imbecility or idiocy. It occurs endemically in the goitrous districts of Switzerland, and sporadically in other parts of Europe and in America. Lack of the secretion of the thyroid gland seems to be the cause.

Cretinoid (*kre'-tin-oid*) [*crétin*, a simple-minded person]. 1. Resembling a cretin; resembling cretinism. 2. A person who resembles a cretin. **C. State**, the morbid state presented by a sufferer from cretinism; cretinism.

Cribriform (*kri'-re-form*) [*cribrum*, a sieve; *forma*, form]. Perforated like a sieve, as the cribriform plate of the ethmoid bone.

Crico- (*kri'-ko-*) [*κρίκος*, a ring]. A prefix denoting connection with the cricoid cartilage.

Cricoid (*kri'-koid*) [*κρίκος*, a ring; *εἶδος*, form]. Ring-shaped. **C. Cartilage**, the ring-shaped cartilage of the larynx.

Cricothyroid (*kri-co-thi'-roid*) [*κρίκος*, a ring; *θυρεοειδής*, shield-shaped]. Pertaining to the cricoid and thyroid cartilages. **C. Artery**, a small branch of the superior thyroid, crossing the cricothyroid membrane. **C. Membrane**, a ligamentous membrane that lies between the cricoid and thyroid cartilages.

C. Muscle. See *Muscles*, Table of.

Crisis (*kri'-sis*) [*κρίσις*, a decisive point]. A

turning-point, as that of a disease, fever, especially the sudden favorable termination of the acute symptoms of an infectious disease. **C.**, **Gastric**, attacks of intense, paroxysmal pain in the abdomen, often attended with vomiting. They occur in locomotor ataxia.

Crista (*kris'-tah*) [*L.*]. Crest. **C. acustica**, a yellow elevation projecting into the equator of the ampulla of the ear. **C. galli**, cock's crest, the superior triangular process of the ethmoid bone.

Crith (*kri-th*) [*κριθή*, barleycorn]. The assumed unit of mass for gases. It is the weight of one liter of hydrogen, which is .0896 of a gram or 1.37 grains.

Critical (*krit'-ik-al*) [*κρίσις*, a decisive point]. Pertaining to a crisis.

Crocus (*kro'-kus*) [*κρόκος*, crocus, saffron]. Saffron. The stigma of the flowers of *C. sativus*. It is an aromatic stimulant, emmenagogue, and antispasmodic. **C.**, **Tinct.**, 10 per cent. in strength. Dose [$\frac{5}{4}$ j-ij (4.0-8.0); of the drug, gr. v-xx (0.32-1.3), in infusion.

Cross-birth. Shoulder-presentation, or other presentation requiring version. **C.-eye**.

See *Strabismus*. **C.-legged Progression**, a method of walking in which one foot gets over or in front of the other.

Crossed (*kroid*) [*crux*, a cross]. Having the shape of a cross. Affecting alternate sides of the body. **C. Anesthesia**. See *Anesthesia*. **C. Hemiplegia**. See *C. Paralysis*. **C. Paralysis**. See *Paralysis*. **C. Reflexes**, reflex movements on one side of the body excited by stimulation of a part on the opposite side.

Crotchet (*krocht'-et*) [*ME.*, *crochet*, a little hook]. A hook used in extracting the fetus after craniotomy.

Croton (*kro'-ton*) [*κρότων*, a tick]. A great genus of euphorbiaceous plants. **C. eleuteria**, yields cascarrilla. **C. tiglium**. See *Tiglitum*. **C. Chloral**. See *Chloral butyricum*. **C. Oil**. See *Tiglitum*.

Croup (*kroop*) [*AS.*, *krōpan*, to cry aloud]. Membranous croup; pseudomembranous croup; true croup; a disease of the larynx and trachea of children, prominent symptoms of which are a harsh "croupy" cough, and difficulty in breathing; it is often accompanied by the development of a membranous deposit or exudate upon the parts. It is usually caused by the diphtheria-bacillus, sometimes by other microorganisms. **C.**, **Catarrhal**, a simple non-contagious inflammation of the larynx accompanied by the formation of membrane. **C.**, **False**, a spasm of the muscles of the larynx with a slight inflammation. **C.**, **Spasmodic**. See *C.*, *False*.

Crucial (*kru'-shal*) [*crux*, a cross]. Resembling or pertaining to a cross, as a crucial incision.

Crura (*kru'-rah*) [plural of *crus*, a leg]. See *Crus*.

Crural (*kru'-ral*) [*crus*, a leg]. Pertaining to the thigh. **C. Arch.** See *Ligament*. **C. Hernia**, femoral hernia. **C. Ring**, the femoral ring; the upper opening of the femoral canal, bounded in front by Poupart's ligament and the deep crural arch, behind by the pubis, internally by Gimbernat's ligament, externally by a fibrous band separating it from the femoral vein.

Crureus (*kru'-re-us*) [L.]. One of the muscles of the thigh.

Crus (*kru-s*) [L.]. A leg, limb, or support. **C. cerebelli**, any one of the cerebellar peduncles. **C. cerebri**, either of the two peduncles connecting the cerebrum with the pons. **C. of the Diaphragm**, either of the two fibromuscular bands arising in front of the vertebræ and inserted into the central tendon of the diaphragm. **C. of the Penis**, the corpus cavernosum.

Crusocreatinin (*kru-so-kre-at'-in-in*), $C_8H_8N_2O_4$. A leukomatin, isolated from muscle-tissue.

Crust (*krust*) [*crusta*]. A covering, especially a dried exudate on the skin.

Crusta (*kru-s'-tah*) [L. a crust]. The inferior portion of the crus cerebri. **C. petrosa**, a thin layer of bone covering the fang of a tooth. **C. phlogistica**, the yellowish layer of the upper stratum of a blood-clot coagulating slowly.

Crutch-paralysis. Paralysis of an upper extremity due to the pressure of the crutch-head upon the nerves of the axilla, especially the musculospiral nerve.

Cruveilhier's Disease. See *Diseases, Table of*.

Crypt (*kript*) [*κρυπτός*, hidden]. A small sac or follicle. **Crypts of Lieberkühn**, minute tubular depressions of the mucous membrane of the small intestine.

Cryptocephalus (*krip-to-sef'-al-us*) [*κρυπτός*, hidden; *κεφαλή*, head]. A fetal monster with an imperfectly formed and concealed head.

Cryptophthalmos (*krip-toff-thal'-mos*) [*κρυπτός*, hidden; *ὀφθαλμός*, the eye]. 1. Congenital union of the eyelids, usually over imperfect eyes. 2. A person who has congenital union of the eyelids.

Cryptopin (*krip'-to-pin*) [*κρυπτός*, hidden; *ὀπιον*, opium], $C_{11}H_{12}NO_8$. One of the alkaloids of opium, colorless and odorless. It is said to be anodyne and hypnotic, but it is less safe than morphin. Dose, $\frac{1}{8}$ gr. (0.008).

Cryptorchid, or **Cryptorchis** (*krip-tor'-kid*, *-kis*) [*κρυπτός*, hidden; *ὄρχις*, testicle]. A

person with retained testicles, *i. e.*, not descended into the scrotum.

Crystal (*kris'-tal*) [*κρύσταλλος*, clear ice]. In chemistry, a substance that assumes a definite geometric form. **C. Charcot-Leyden**, **C. Charcot's**, minute colorless crystals found in the sputa of asthma and bronchitis, and in other conditions. They consist of spermin.

Crystallin (*kris'-tal-in*) [*κρύσταλλος*, clear ice]. The globulin of the crystalline lens.

Crystalline (*kris'-tal-en* or *-in*) [*κρύσταλλος*, crystal]. Like a crystal. **C. Lens**. See *Lens*.

Crystallization (*kris-tal-is-a'-shun*) [*κρύσταλλος*, ice]. The process by which the molecules of a substance arrange themselves in geometric forms when passing from a gaseous or a liquid to a solid state. **C., Water of**, the water of salts that cannot be extracted without destruction of their crystalline nature.

Cubeb, or **Cubeba** (*ku'-beb*, or *ku-be'-bah*) [Pers., *ka-baba*, cubeb]. The unripe fruit of *C. officinalis*, cultivated in Java. Its properties are due to a volatile oil, $C_{15}H_{24}$, and an organic acid. It is an aromatic stimulant, diuretic in small doses, and is useful in affections of the bladder and urethra. It is also employed in catarrh of the air-passages, etc. Dose gr. x- $\frac{3}{4}$ ij (0.65-8.0). **C., Ext.**, **Fld.**, alcoholic. Dose \mathfrak{M} v-xxx (0.32-2.0). **C., Oleoresina**, ethereal. Dose \mathfrak{M} v-xxx (0.32-2.0). **C., Oleum**, the volatile oil. Dose \mathfrak{M} v-xx (0.32-1.3). **C., Tinct.**, 10 per cent. in strength. Dose \mathfrak{M} x-f $\frac{3}{4}$ iij (0.65-12.0). **C., Trochisci**, oleoresin gr. $\frac{1}{2}$, oil of sassafras gr. $\frac{1}{8}$, ext. of glycyrrhiza gr. iv, acacia gr. ij, syr. of tolu q.s., in each troche. Dose j-ij.

Cubebic Acid (*ku-beb'-ik*) [*cubeb*], $C_{15}H_{14}O_7$. A resinous acid body from cubebs; actively diuretic and cathartic. Unof.

Cubebin (*ku-beb'-in*) [*cubeb*], $C_{16}H_{16}O_8$. An odorless crystalline substance obtained from cubebs.

Cubitus (*ku'-bi-tus*) [*cubitus*, the elbow]. The forearm.

Cuboid (*ku'-boid*) [*κίβος*, cube; *εἶδος*, resemblance]. Resembling a cube. **C. Bone**, a bone of the foot situated at the outer anterior part of the tarsus.

Cuca (*koo'-kah*). See *Erythroxylon*.

Cucumber, Squirting. See *Eluterium*.

Cuirass (*kue-ras'*) [Fr.]. A close-fitting or immovable bandage for the front of the chest.

C. Cancer, a large, flat, carcinomatous growth upon the front of the chest.

Cul de sac, or preferably, **Culdesac** (*ku'-de-sak*) [Fr.]. A closed or "blind" pouch or sac. **Douglas's C.**, a pouch between the anterior wall of the rectum and the

posterior wall of the uterus, formed by the reflexion of the peritoneum.

Culex (*ku'-leks*) [L., a gnat]. A mosquito.

Culture (*kul'-chūr*) [*colere*, to till]. The growth of microorganisms on artificial media.

The act of cultivating microorganisms on artificial media. **C., Hanging-drop**, a culture in which the microorganism is inoculated into a drop of fluid on a cover-glass and the latter is inverted over a glass slide having a central concavity. **C. Media**, certain substances used for cultivating bacteria. They are either liquid or solid, bouillon and milk being the important liquid, and gelatin, agar-agar, blood-serum, and potato, the principal solid media. **C., Plate**, a culture of bacteria on a medium spread upon a flat plate or in a double dish. **C., Pure**, a culture of a single microorganism. **C., Stab**, one in which the medium is inoculated by means of a needle bearing the microorganisms, and which is inserted deep down into the medium.

Cumulative (*ku'-mu-la-tiv*) [*cumulare*, to heap up]. Increasing; adding to. **C. Action**, or **Effect**, the production of a marked and sudden result, after the administration of a considerable number of comparatively ineffective doses.

Cundurango (*kun-du-ran'-go*). See *Condurango*.

Cuneate (*ku'-ne-at*) [*cuneus*, a wedge]. Wedge-shaped.

Cuneiform (*ku'-ne'-if-orm*) [*cuneus*, a wedge; *forma*, shape]. Wedge-shaped, cuneate. **C. Bones**, three wedge-shaped bones at the anterior part of the tarsus. **C. Columns**. See *Columns of Burdach*.

Cuneus (*ku'-ne-us*) [L., a wedge]. A wedge-shaped convolution on the mesial aspect of the occipital lobe.

Cuniculus (*ku'-nik'-u-lus*) [L.]. The burrow of the itch-insect.

Cup (*kup*) [AS., *cuppe*, a cup]. 1. To bleed. 2. A cupping-glass. **C., Dry**, a cup for merely drawing the blood to the surface. **C., Favus**, a depression in a favus-scale surrounding a hair. **C., Glaucomatous**, a deep depression in the optic papilla seen in cases of glaucoma. **C., Physiologic**, the normal concavity of the optic papilla. **C., Wet**, a cup for abstracting blood through incisions in the skin.

Cupphosis (*ku'-fo'-sis*). See *Kyphosis*.

Cupped (*kup*) [AS., *cuppe*, a cup]. Having the upper surface depressed; applied to the coagulum of blood after phlebotomy. **C. Disc**, excavation of the optic papilla, normally present in slight degree, but pathologic if excessive.

Cupping (*kup'-ing*) [AS., *cuppe*, a cup]. A method of blood-derivation by means of the application of cupping-glasses to the surface

of the body. **C., Dry**, a form of counter-irritation in which the blood is drawn to the surface by means of a cup. This is used mainly in inflammatory affections of the lung. **C.-glass**, a small bell-shaped glass capable of holding three to four ounces, in which the air is rarefied either by heat or exhaustion, and the glass applied to the skin, either with or without scarification of the latter. **C., Wet**, with the abstraction of blood after scarification.

Cuprum (*ku'-prum*) [L.]. See *Copper*.

Curaçao (*ku'-ra-so'*) [Island of *Curaçao*, north of Venezuela]. A cordial or elixir prepared from brandy, and flavored, principally with orange-peel. It is used as a vehicle for certain medicines.

Curaçoa (*ku'-ras-o'-ah*). See *Curaçao*.

Curara, **Curare**, **Curari** (*koo-rah'-rah*, *koo-rah'-ret*, *koo-rah'-re*) [S. Am.]. Woorara.

A vegetable extract obtained from *Paullinia curare* and certain members of the *Strychnos* family. It is a powerful paralyzant of the motor nerves and of the voluntary muscles.

Its active principle is curarin, $C_{10}H_{15}N$; dose gr. $\frac{1}{100}$ – $\frac{1}{10}$, hypodermically. It is used in S. America and elsewhere as an arrow-poison. Toxic doses cause death by paralysis of the organs of respiration. It has been reported effectual in cases of hydrophobia and tetanus. Dose by hypodermic injection gr. $\frac{1}{3}$ – $\frac{1}{2}$ (0.003–0.01). For hypodermic injection a solution of 5 grains in 60 minims is employed. Dose m j–vj (0.065–0.4).

Curcuma (*ker'-ku-mah*) [L.]. Turmeric. The rhizome of *Curcuma longa* of India. Its action is similar to that of ginger. It is employed as a yellow dye; in chemistry, as a test for alkalies, which turn it brown; and in pharmacy, occasionally, to color ointments and other preparations.

Curd (*kerd*) [ME., *curd*, curds]. The coagulum of milk that separates on the addition of rennet or an acid to milk.

Cure (*kūr*) [*cura*, care]. The successful treatment of a disease; also, a system of treatment, as *Faith-cure*, *Mind-cure*, *Grape-cure*, *Water-cure* (see *Hydrotherapy*), *Hunger-cure*, *Rest-cure*, etc. **C., Potato**, a method of treating foreign bodies in the alimentary tract by the ingestion of mashed potatoes. The body becomes imbedded in the potato-mass that is formed.

Curet or Curette (*ku-ret'*) [Fr.]. An instrument shaped like a spoon or scoop, for scraping away exuberant or dead tissue.

Current-jelly Clot. A soft, red clot seen post-mortem in the heart and blood-vessels.

Current (*ku'-ent*) [*corere*, to run]. A term applied to the transference of electric force, which is likened to the flow of a liquid in a tube. **C., After**, a current produced in

nervous or muscular tissue when a constant current which has been flowing through the same has been stopped. **C., Alternating**, a term applied to a current which, by means of an interrupter, is alternately direct and reverse. **C., Ascending**, the current formed by placing the positive electrode upon the periphery of a nerve and the negative higher up on the trunk of the nerve, or on the surface over the nerve-center in the spinal cord. **C., Battery**, a galvanic current. **C., Centrifugal**, a descending current. **C., Centripetal**, an ascending current. **C., Constant**, See **C., Continuous**. **C., Continuous**, a constant, uninterrupted current in one direction. **C., Derived**, a current drawn off by a derivation-wire from the main current. **C., Descending**, one passing through a nerve centrifugally, the anode being placed proximally, the cathode distally. **C., Direct**, a current constant in direction, in contradistinction to an alternating current. **C., Faradic**, the current produced by an induction-coil, or by a magneto-electric machine. **C., Galvanic**, a current generated by the decomposition of acidulated water by means of metallic plates. **C., Induced**. See **C., Secondary**. **C., Interrupted**, a current that is alternately opened and closed. **C., Labile**, a current applied while moving one or both electrodes over the surface treated. **C., Reversed**, that produced by changing the poles. **C., Secondary**, momentary currents produced in a coil of insulated wire, surrounded within the field of another coil, when the current is made or broken in the second coil. **C., Stable**, a current applied with both electrodes at a fixed position. **C., Voltaic**, the continuous current.

Curschmann's Spirals. Spiral threads of mucus surrounded in the spinal cord by cells that are experienced during an epileptic paroxysm. They are supposed to be parts of the dendrites and axons of Purkinje-Leyden cells and ependymal cells.

Curvature. *curvatur* (L.) *curvus* (L.) *curvatus* (L.). A bending or curving. **C., Angular**. See **Synangium**. **C., From**, a angular curvature of the spine with a posterior prominence. **C. of Spine**, a bending of the vertebral column.

Concavity. *concavitas* (L.) *concavus* (L.) *concavatus* (L.). A concave surface.

Consp. eye [*consp. oculus*]. The prominence on the front of a tooth.

Consp. ear. See **Stapes**.

Cytaceous. *cytaceus* (L.) *cyta* (L.) *cytaceus* (L.). Pertaining to the skin. **C., Epidermal**. See **Epidermis**. **C., Peridermal**. See **Peridermis**. **C., Transdermal**. The transposition of gases through the skin.

Cytide. *cytides* (L.) *cytis* (L.) *cytis* (L.). The epidermis or cuticle.

Cutis (*ku'-tis*) [L.]. The derma, or true skin. **C. anserina**. See **Goose skin**. **C. testacea**, a variety of seborrhea in which the trunk and extensor surfaces of the extremities are covered with large, thick plates of greasy, inspissated sebum, usually greenish or blackish, from accumulation of dirt upon them. **C. unctuosa**. Synonym of **Seborrhea**.

Cutisector (*ku-te-tek'-tor*) [*cutis*, skin; *sector*, a cutter]. An instrument for taking small sections of skin from the living subject.

Cutol (*ku'-tol*). Aluminum boricotannicum. A powder, insoluble in water, used in dermatology.

Cyanic (*si-an'-ik*) [*ciavus*, blue]. Blue or bluish. **C. Acid**. See **Acid**, **Cyanic**.

Cyanid (*si-an'id*) [*ciavus*, blue]. Any compound of cyanogen with a metal, or a radicle. Most of the cyanids are actively poisonous.

Cyanogen (*si-an'-o-jen*) [*ciavus*, blue; *gennao*, to produce]. A radicle having the structure C.N., an acid compound of carbon and nitrogen, existing as a gas, called cyanide gas; it is exceedingly poisonous.

Cyanosis (*si-an'-o-sis*) [*ciavus*, blue]. A bluish coloration of the skin from deficient oxidation of the blood caused by loss or general circulatory disturbance. **C., Congenital**. See **Congenital cyanosis**. Due to a congenital lesion of the heart or the great vessels.

Cyclic (*si'-kik*) [*kyklos*, circular]. Having the character of a circle, or of a cycle, or of a series of cycles. **C. Albuminuria**. See **Cyclical albuminuria**. **C. Insanity**. See **Cyclical insanity**.

Cyclitis (*si'-kli-tis*) [*kyklos*, circular; *itis*, inflammation]. Inflammation of the iris, which leads to a loss of transparency of the internal eye, surrounded by the cornea. It may be serous, purulent, or suppurative. **Indolent** is the character of the disease, and it may vary in its intensity.

Cyclocephalus (*si'-klo-sef'-a-lus*) [*kyklos*, circular; *kephala*, head]. A genus of large aquatic mollusk, characterized by a large, rounded, circular, convex shell, which is regular in shape, with an internal siphon of expanded and retracted parts, and a large, muscular foot.

Cyclopora (*si'-klo-po'-ra*) [*kyklos*, circular; *pora*, pore]. A genus of small aquatic mollusk, characterized by a small, rounded, circular, convex shell, which is regular in shape, with an internal siphon of expanded and retracted parts, and a large, muscular foot.

Cyclops (*si'-klo-ps*) [*kyklos*, circular; *ops*, eye]. A genus of small aquatic mollusk, characterized by a small, rounded, circular, convex shell, which is regular in shape, with an internal siphon of expanded and retracted parts, and a large, muscular foot.

Cyclostoma (*si'-klo-sto-ma*) [*kyklos*, circular; *stoma*, mouth]. A genus of small aquatic mollusk, characterized by a small, rounded, circular, convex shell, which is regular in shape, with an internal siphon of expanded and retracted parts, and a large, muscular foot.

Cyclostomus (*si'-klo-sto-mus*) [*kyklos*, circular; *stoma*, mouth]. A genus of small aquatic mollusk, characterized by a small, rounded, circular, convex shell, which is regular in shape, with an internal siphon of expanded and retracted parts, and a large, muscular foot.

ployed mainly for the mucilage contained in the covering, which consists of a compound of gum and glucose. **C.**, **Mucilago**, quince-seeds 2, macerated in water 100 parts, is a bland demulcent, and is used as a hair-dressing.

Cyesis (*si-el'-sis*) [*κύσις*, pregnancy]. Pregnancy.

Cyetic (*si-el'-ik*) [*κυητικός*, pregnant]. Relating to pregnancy.

Cylinder (*sil'-in-der*) [*κύλινδρος*, a cylinder]. See *Lens*.

Cylindroid (*sil'-in-droid*) [*κύλινδρος*, cylinder; *ειδος*, likeness]. A name given to a mucous cast frequently found in the urine in cases of mild irritation of the kidney. Cylindroids are ribbon-like forms, usually of great length, and of about the same diameter as renal casts. They may assume various shapes. One extremity is usually pointed and may be drawn out into a long tail.

Cylindroma (*sil-in-dro'-mah*) [*κύλινδρος*, a cylinder]. A myxosarcoma in which the degeneration is confined to areas surrounding the blood-vessels.

Cynanche (*sin-ang'-ke*) [*κύνω*, a dog; *ἀγχειν*, to strangle]. An old name for any acute affection of the throat, as diphtheria, croup, tonsillitis, etc., in which the patient struggles for breath (as a panting dog). **C.** **maligna**, a fatal form of sore-throat. **C.** **suffocativa**. Synonym of *Croup*. **C.** **tonsillaris**. See *Quincy*.

Cynanthropia (*sin-an-thro'-pe-ah*) [*κύνω*, dog; *ἄνθρωπος*, a man]. A mania in which the patient believes himself a dog, and imitates the actions of one.

Cynic (*sin'-ik*) [*κυνικός*, dog-like]. Pertaining to a dog. **C.** **Spasm**, a contraction of the facial muscles upon one side, so as to expose the teeth in the manner of an angry dog.

Cynurenic Acid (*sin-u-ren'-ik*) [*κύνω*, a dog; *οὔρον*, urine], $C_{10}H_{13}N_2O_6 + 2H_2O$. A crystalline acid found in dog's urine. It is a decomposition-product of proteids. On heating it cynurin is evolved.

Cypripedium (*sip-re-pe'-de-um*) [*κύπρις*, Venus; *πέδιον*, a slipper]. Lady's slipper. The roots of *C. pubescens* and *C. parviflorum*, American valerian, the properties of which are due to a volatile oil and acid. It is an antispasmodic and stimulant tonic, used instead of valerian, which it resembles. **C.**, **Ext.**, **Fld.** Dose πλx-xxx (0.65-2.0). Unof.

Cyrtometer (*sir-tom'-et-er*) [*κυρτός*, curved; *μέτρον*, a measure]. An instrument adapted for measuring curves. One form is used to locate the fissures of the brain.

Cyst (*üst*) [*κύστις*, a pouch]. A cavity containing fluid and surrounded by a capsule. **C.**, **Blood**. See *Hematoma*. **C.**, **Cutaneous**.

See *C.*, *Dermoid*. **C.**, **Daughter**, any one of the small cysts developed by secondary growth from the walls of a large cyst. **C.**, **Dentigerous**, one containing teeth. **C.**, **Dermoid**, a congenital cyst containing bone, hair, teeth, etc. **C.**, **Echinococcus**, a cyst formed in various tissues and organs of man by the larva of the *Tænia echinococcus* of the dog, taken into the stomach. **C.**, **Extravasation**, a cyst formed by the encapsulation of a hemorrhage into the tissues. **C.**, **Follicular**, one due to the occlusion of the duct of a small follicle or gland. **C.**, **Hydatid**. See *C.*, *Echinococcus*. **C.**, **Meibomian**. See *Chalazion*. **C.**, **Mucous**, a retention-cyst containing mucus. **C.**, **Multilocular**, one composed of many separate compartments. **C.**, **Retention**, one that is due to the retention of the secretion of a gland, in consequence of closure of the duct, as in mucous or sebaceous cysts. **C.**, **Sebaceous**, a retention-cyst of a sebaceous gland. **C.**, **Softening**, one due to encapsulation of the fluid after liquefaction-necrosis. **C.**, **Sublingual**. See *Ranula*. **C.**, **Unilocular**, one having but a single cavity.

Cystadenoma (*sist-ad-en-o'-mah*) [*κύστις*, cyst; *ἀδην*, gland; *δύα*, a tumor]. An adenoma containing cysts.

Cystalgia (*sist-all'-je-ah*) [*κύστις*, bladder; *ἄλγος*, pain]. Pain in the bladder.

Cystic (*sist'-ik*) [*κύστις*, bladder]. 1. Pertaining to or resembling a cyst. 2. Pertaining to the urinary bladder or the gall-bladder.

Cysticercus (*sist-te-ser'-kus*) [*κύστις*, a bladder; *κέρκος*, a tail]. The embryo of a tapeworm when it has reached the encysted stage. An hydatid. **C.** **cellulose**, the larval parasite inhabiting the intermuscular connective tissue of the pig, producing the condition known as "measles." It is rarely found in the tissues of man. Its progenitor is the *tænia solium*.

Cystin (*sist'-in*) [*κύστις*, bladder], $C_2H_7N_3SO_2$. A substance found in the urine. It occurs in regular, colorless, six-sided tables, of very characteristic appearance.

Cystinuria (*sist-in-u'-re-ah*) [*κύστις*, bladder; *οὔρον*, urine]. The presence of cystin in the urine.

Cystitis (*sist-it'-tis*) [*κύστις*, bladder; *ιτις*, inflammation]. Inflammation of the bladder.

Cystitome (*sist-it'-om*). See *Cystotome*.

Cystocele (*sist-o-sel*) [*κύστις*, pouch; *κῆλη*, a tumor]. A hernia of the bladder.

Cystoma (*sist-o'-mah*) [*κύστις*, a cyst; *δύα*, a tumor]. A new growth made up of cysts; applied especially to ovarian cysts.

Cystopexy (*sist-o-peki-e*) [*κύστις*, bladder; *πέξις*, fixation]. Fixation of the bladder, an operation for the cure of cystocele.

Cystoscope (*sist-o-skop*) [*κύστις*, bladder;

σκοπεῖν, to examine]. An instrument for inspecting the interior of the bladder.

Cystotome (*sist'-o-tōm*) [*κυστίς*, bladder; *τέμνειν*, to cut]. A knife used in cystotomy; also a knife used in rupturing the capsule of the lens in cataract-operations.

Cystotomy (*sist'-ot'-o-me*) [*κυστίς*, bladder; *τομή*, a cutting]. Incision of the bladder.

Cytochrome (*si'-to-krōm*) [*κύτος*, cell; *χρῶμα*, color]. A term applied by Nissl to nerve-cells deficient in cell-protoplasm, the nucleus not being completely surrounded. The nucleus stains well and is about the size of the leukocyte-nucleus.

Cytode (*si'-tōd*) [*κύτος*, a cell; *εἶδος*, form]. The simplest, most primitive form of cell, without nucleus or nucleolus.

Cytogenesis (*si'-to-jen'-es-is*) [*κύτος*, a cell; *γενεσις*, production]. The formation or genesis of the cell.

Cytoglobin (*si-to-glo'-bin*) [*κύτος*, a cell; *globus*, a ball]. An albuminoid, obtainable in the form of a white, soluble powder. It forms about three per cent. of the pulp of the lymphatic glands.

Cytometer (*si-tom'-et-er*) [*κύτος*, cell; *μέτρον*, a measure]. A device for counting cells, especially blood-corpuscles. See also *Hemocytometer*.

Cytomitoma (*si-to-mi-to'-mah*) [*κύτος*, cell; *μίτος*, a fiber]. The fibrillar part of a cell-body. See *Mitoma*.

Cytoplasm, or **Cytoplasma** (*si'-to-plasm*, or *si-to-plas'-mah*) [*κύτος*, cell; *πλάσμα*, anything formed]. Protoplasm.

Cytozoon (*si-to-zō'-on*) [*κύτος*, cell; *ζῶον*, animal]. A protoplasmic cell-mass, probably parasitic in nature, with independent movement; found by Gaule in defibrinated blood and other structures.

D

D. An abbreviation for *dexter*, right; *dioptry*; and *detur*, let it be given.

Dacryadenitis, **Dacryoadenitis** (*dak'-re-ad-en-i'-tis*, *dak'-re-o-ad-en-i'-tis*) [*δάκρυον*, a tear; *ἀδην*, a gland; *ιτις*, inflammation]. Inflammation of the lacrimal gland.

Dacryocystitis (*dak'-re-o-sis-ti'-tis*) [*δάκρυον*, tear; *κυστίς*, a sac; *ιτις*, inflammation]. Inflammation of the lacrimal sac.

Dacryolith (*dak'-re-o-lith*) [*δάκρυον*, tear; *λίθος*, stone]. A calcareous concretion in the lacrimal passages.

Dacryops (*dak'-re-ops*) [*δάκρυον*, tear; *ὄψ*, eye]. 1. A watery eye. 2. A cyst of the duct of a lacrimal gland.

Dacryocystotome (*dak'-re-o-sis'-to-tōm*) [*δάκρυον*, a tear; *κυστίς*, a sac; *τομή*, a cutting]. An instrument for dividing strictures of the lacrimal passages.

Dacryocystoblennorrhœa (*dak'-re-o-sis-to-blēn-or-e'-ah*) [*δάκρυον*, a tear; *κυστίς*, a sac; *βλέννα*, mucus; *ροία*, a flow]. Chronic inflammation of the lacrimal sac with a mucopurulent discharge.

Dactylitis (*dak'-til-i'-tis*) [*δάκτυλος*, a finger; *ιτις*, inflammation]. Inflammation of a finger or toe.

Dæmonomania (*de-mo-no-ma'-ne-ah*). See *Demonomania*.

Dalton-Henry Law. See *Law*.

Daltonism (*dall'-ton-ism*) [*Dalton*, a physician]. Color-blindness.

Dam. See *Rubber-dam*.

Damiana (*dam-e-an'-ah*). The leaves of

Turnera aphrodisiaca, found in Mexico and Lower California; a stimulant tonic and aphrodisiac. It is the basis of a great number of quack remedies. **D.**, **Ext.** Dose gr. ij-x (0.13-0.65). **D.**, **Ext.**, **Fld.** Dose ηx-fʒj (0.65-4.0). Dose of the leaves ʒj (32.0) daily. All unof.

Dammar (*dām'-ar*) [*Hind.*, *dāmar*, resin]. A gum or resin produced by various species of *Dammara* and other trees. **D.**, **True**, is obtained from the *Dammara orientalis*, a coniferous tree, indigenous in the East Indies, and also from *Dammara australis*, in New Zealand.

Dance, **St. Vitus's**. See *Chorea*.

Dancing Mania. See *Choromania*.

Dandelion (*dān'-de-li-on*). See *Taraxacum*.

Dandruff (*dān'-druf*) [*Welsh*, *ton*, skin; *drug*, bad]. The scurf or scales formed upon the scalp in seborrhea.

Dandy (*dān'-de*) **Fever**. See *Dengue*.

Daphne (*daf'-ne*) [*δάφνη*, the laurel]. See *Mezerion*.

Darier's Disease. See *Diseases*, *Table of*.

Dartos (*dar'-tos*) [*δάρτος*, flayed]. The contractile musculobrous layer beneath the skin of the scrotum.

Dartrous (*dar'-trus*) [*Fr.*, *dartre*]. Of the nature of tetter or herpes; herpetic.

Darwinism (*dar'-win-ism*). The doctrine that higher organisms have been developed from lower forms by the influence of natural selection, a theory advocated by Charles Darwin.

Datura (*da-tu'-rah*) [Hind., *dhatāra*, a certain plant]. A genus of Solanaceæ, or night-shade family. **D. stramonium**. See *Stramonium*.

Daturin (*da-tu'-rin*) [Hind., *dhatāra*, a certain plant]. A poisonous alkaloid from the thorn-apple. See *Atropin*.

Daughter (*daŋ'-ter*). A female child or descendant. **D.-cell**. See *Cell*. **D.-cyst**, a cyst formed within a cyst. **D.-nuclei**. See *Karyokinesis*. **D.-star**, an amphiaster. See *Karyokinesis*.

Day-blindness. See *Nyctalopia*; also *Hemeralopia*.

Deaf (*def*) [A.S., *deaf*, deaf]. Lacking the sense of hearing; in a condition of impaired hearing. **D.-mutism**, the state of being both deaf and dumb; the deafness may be congenital or acquired, and prevent the individual from learning to speak. **D.-mutism**, **Hysteria**, a condition of deaf-mutism of sudden development, due to hysteria.

Deafness (*def'-nes*) [A.S., *deaf*, deaf]. The state of being deaf. Deafness may be due to disease of the external auditory canal, the middle ear, the internal ear, the auditory nerve, or the brain. **D.**, **Boilermakers'**, deafness resulting from working among machinery, and characterized by inability to hear ordinary conversation, while hearing-power is increased amidst loud noise. **D.**, **Cerebral**, that due to a brain-lesion. **D.**, **Cortical**, that due to disease of the cortical centers for hearing; it may be absolute, or consist in the inability to comprehend spoken language—a **psychic deafness**. **D.**, **Word**. Synonym of *D.*, *Psychic*.

Death (*deŋh*) [A.S., *deŋth*, death]. The cessation of life. **D.**, **Black**, an exceedingly fatal epidemic called the "Plague," which occurred in Europe during the 14th century, and during which, it is estimated, 20,000,000 persons died. **D.**, **Local**, death of a part. **D.**, **Molecular**, death of individual cells; ulceration. **D.**, **Somatic**, death of the organism as a whole.

Debove's Membrane. See *Membrane*.

Deca- (*dek'-a-*) [*deka*, ten]. Ten; prefixed to the units of weight, capacity, and length in the metric system, it signifies a measure ten times as large as the unit. See *Metric System*.

Decalcification (*de-kal'-sif-ik-a'-shun*) [*de* priv.; *calx*, lime; *facere*, to make]. The withdrawal of the lime-salts of bone.

Decalcify (*de-kal'-sif-i*) [*de* priv.; *calx*, lime; *facere*, to make]. To remove lime-salts from tissues.

Decantation (*de-kan-ta'-shun*) [*de*, down; *cantus*, a side]. The operation of removing the supernatant fluid from a sediment.

Decapitation (*de-kap-it-a'-shun*) [*de*, from; *caput*, head]. The act of beheading, espe-

cially as performed on the fetus when other means of delivery have failed.

Decay (*de-ka'*) [*de*, down; *cadere*, to fall]. 1. Putrefactive change. 2. The ultimate katabolic state; decline of life, of health, or of one or more functions.

Deci- (*des'-e-*) [*decem*, ten]. A prefix which, joined to the metric units of length, capacity, and weight, signifies a measure one-tenth as large as the unit. See *Metric System*.

Decidua (*de-sid'-u-ah*) [*deciduis*, a falling off]. The mucous membrane which lines the uterus and surrounds the ovum during pregnancy. **D. reflexa**, that part of the decidua growing about the ovum and enclosing it as a sac. **D. serotina**, that part of the decidua vera upon which the ovum lies, and from which the placenta is subsequently formed. **D. vera**, the thickened, vascular, spongy mucous membrane of the gravid uterus.

Deciduoma (*de-sid-u-o'-mah*) [*deciduis*, a falling off; *oma*, a tumor]. An intrauterine tumor containing decidual relics, and believed to arise from some hyperplasia of a retained portion of the decidua. By some it is considered a sarcoma.

Deciduous (*de-sid'-u-us*) [*de*, away, from; *cadere*, to fall]. Falling off. **D. Teeth**, the temporary teeth or milk-teeth.

Declination (*dek-lin-a'-shun*) [*declinare*, to decline]. The dip of the magnetic needle.

Decline (*de-klin'*) [*declinare*, to bend]. A gradual decrease, as of a fever; a wasting away of the bodily strength.

Decoction (*de-kok'-shun*) [*decoquere*, to boil down]. A liquid preparation obtained by boiling vegetable substances in water.

Decoloration (*de-kul-or-a'-shun*) [*decolorare*, to deprive of color]. Removal of color.

Decomposition (*de-kom-po-zish'-un*) [*decomponere*, to decompose]. 1. The separation of the component principles of a body. 2. Putrefactive fermentation.

Decortication (*de-kor-tik-a'-shun*) [*de*, from; *cortex*, the bark]. 1. The stripping of the bark or husk of a plant. 2. The stripping off of portions of the cortical substance of the brain from the summits of the gyri.

Decubitus (*de-kul'-bit-us*) [*decumbere*, to lie down]. 1. The position of a sick person while in bed. 2. A bed-sore, usually acute in onset and due to spinal lesions.

Decussate (*de-kus'-at*) [*decussatus*, crossed]. To intersect, to cross.

Decussation (*de-kus-a'-shun*) [*decussatio*, a crossing]. A chiasma or X-shaped crossing, especially of symmetric parts, as of nerve-fibers or nerve-tracts, or of nerve-filaments. The principal decussations are that of the optic nerve and that of the lateral pyramidal tracts in the medulla.

Deep Reflexes. See *Reflexes*, *Table of*. **D.** Water, water obtained from a porous layer beneath the first impervious stratum.

Defecation (*def-ek-a'-shun*) [*defecare*, to separate from the dregs]. The evacuation of the bowels.

Defensive Proteids. Those substances formed in the bodies of animals that render them immune against certain diseases.

Deferent (*def'-er-ent*) [*deferens*, carrying away]. Carrying away or down; efferent.

Defervescence (*de-fer-ves'-ens*) [*defervescere*, to cease boiling]. Disappearance of fever.

Defibrination (*de-fi-brin-a'-shun*) [*de*, from; *fibrā*, a fiber]. The removal of fibrin from blood or lymph.

Definition (*def-in-ish'-un*) [*definitio*; *definire*, to bound by limits]. In optics, the power of a magnifying lens to show clear outlines of the object examined, free from aberration or distortion.

Deflagration (*def-lag-ra'-shun*) [*deflagrare*, to be consumed]. A sudden, violent combustion, such as accompanies the oxidation of certain inorganic substances by mixing them with an easily decomposing salt, such as the alkaline chlorates and nitrates.

Defluvium capillorum (*de-flu'-ve-nim kap-il-or'-um*). Alopecia.

Defluxion (*de-fluk'-shun*) [*defluxio*; *de*, down; *fluere*, to flow]. A discharge.

Deformity, Anterior. See *Lordosis*.

Degeneration (*de-jen-er-a'-shun*) [*degenerare*, to become base]. 1. A morbid process consisting in the conversion of the elements of a tissue into some inert substance. 2. A term indicating imperfect or abnormal development of the psychic faculties. **D.**, **Albuminoid**. Synonym of **D.**, **Amyloid**. **D.**, **Amyloid**. See *Amyloid*. **D.**, **Ascending**, a trophic degeneration of nerve-fibers or tracts progressing from the site of the original lesion toward the cerebrum. **D.**, **Colloid**, the change of the protoplasm of epithelial cells into a substance that resembles mucus, but is not precipitated by alcohol or acetic acid. **D.**, **Descending**, a degeneration of nerve-fibers or tracts extending peripherally from the original lesion. **D.**, **Fatty**, a change of the proteids of the tissues into fat. **D.**, **Hyaline**, a degeneration affecting particularly the connective tissue of the walls of blood-vessels, and giving rise to a substance resembling amyloid material but lacking its reactions. See *Amyloid*. **D.**, **Mucoid**, the degeneration of tissue into a jelly-like, transparent substance containing mucin. **D.**, **Myxomatous**. Synonym of **D.**, **Mucoid**. **D.**, **Parenchymatous**. See *Cloudy Swelling*. **D.**, **Reactions of**. See *Reaction*. **D.**, **Signs of**, physical imperfections, such as asymmetry of corresponding parts, adherent lobules of the

ear, stammering, supernumerary or deficient digits, etc., observed in persons presenting psychic degeneration. **D.**, **Wallerian**, the degeneration of nerve-fibers and tracts after separation from their trophic centers.

Deglutitio impedita (*de-glu-tish'-e-o im-ped-it'-tah*). Synonym of *Dysphagia*.

Deglutition (*deg-lu-tish'-un*) [*deglutitio*, a swallowing]. The act of swallowing.

Debiscence (*de-his'-ens*) [*de*, off; *hiscere*, to gape or yawn]. The act of splitting open.

Dehydration (*de-hi-dra'-shun*) [*de*, away from; *idwā*, water]. The removal of water.

Deiters' Cells. 1. Certain cellular structures between the outer hair-cells of the organ of Corti. 2. The neuroglia-cells. **D. Process**, the process of a nerve-cell that goes to form an axis-cylinder.

Dejecta (*de-jek'-tah*) [*L.*]. Feces.

Dejection (*de-jek'-shun*) [*dejectio*; *de*, down; *jacere*, to throw]. The discharge of fecal matter; the matter so discharged.

Delamination (*de-lam-in-a'-shun*) [*de*, away; *lamina*, a plate]. The splitting into layers.

Delhi Boil (*del'-he boil*). See *Furunculus orientalis*.

Deligation (*del-ig-a'-shun*) [*deligatio*, a binding]. Ligation, as of an artery.

Deliquescent (*del-ik-wes'-ent*) [*deliquescere*, to melt away]. Dissolving. Applied especially to salts that absorb moisture from the air and liquefy.

Deliriant, Delirifacient (*de-le'-re-ant*, *de-le-re-fa'-she-ent*) [*delirium*, madness; *facere*, to make]. Producing delirium.

Delirious (*de-le'-re-us*) [*delirius*, mad, raving]. Affected with delirium.

Delirium (*de-le'-re-um*) [*delirium*, madness; *de*, out of; *lira*, the furrow]. A condition of mental excitement with confusion and usually hallucinations and illusions. **D.**, **Alcoholic**. See *D. tremens*. **D. cordis**, a violent, tumultuous beating of the heart. **D.**, **Febrile**, the delirium of fever. **D. of Grandeur**, a condition in which an individual has insanely exaggerated ideas of his own importance or his possessions. **D. of Persecution**, that in which the patient imagines himself the object of persecution. **D.**, **Toxic**, delirium caused by poisons. **D. tremens**, the delirium arising from alcoholic poisoning. It is characterized by constant tremor, insomnia, great exhaustion, distressing illusions, and hallucinations.

Delitescence (*del-it-es'-ens*) [*delitescere*, to lie hid]. The sudden disappearance of inflammation by resolution.

Deliver (*de-liv'-er*) [*de*, from; *liberare*, to free]. To free from something, especially to deliver a woman of a child, or of the after-birth. The word is also applied to the part

removed, as to deliver the placenta or a tumor.

Delivery (*de-liv'-er-e*) [Fr., *délivrer*; *deliberare*, to set free.] The act of delivering or freeing from something, especially the relieving of a woman from the contents of the uterus. Parturition; child-birth. **D.**, Post-mortem, the birth of a fetus after the death of the mother.

Delomorphous (*del-o-mor'-fus*) [*δηλος*, conspicuous; *μορφή*, form]. Having a conspicuous form. **D. Cells of Rollet**, large, well-defined cells, between the membrana propria and the chief cells of the fundus-glands of the gastric mucous membrane. They are supposed to secrete the hydrochloric acid.

Delphinin, Delphinium, Delphinoidin, Delphisin (*del'-fin-in*, *del-fin'-e-um*, *del-fin-oid'-in*, *del'-fis-in*). See *Staphisagria*.

Deltoid (*del'-toid*) [*delta*, the Greek letter Δ; *ειδος*, likeness]. Having the shape of the Greek letter *delta*; triangular, as the **D. Muscle**. See *Muscles, Table of*.

Delusion (*de-lu'-shun*) [*de*, from; *lusus*, play]. A false belief, the falsity of which is apparent, but out of which the person cannot be reasoned by indubitable evidence.

Delusional (*de-lu'-shun-al*) [*deludere*, to delude]. Of the nature of a delusion; characterized by delusions. **D. Stupor**. See *Insanity, Confusional*.

Demarcation (*de-mark-a'-shun*) [*demarcare*, to set the bounds of]. Separation. **D.**, Line of, a red line forming at the edge of a gangrenous area and marking the limit of the process.

Dement (*de'-ment*) [*demens*, insane]. A person suffering with dementia.

Dementia (*de-men'-she-ah*) [*de*, away from; *mens*, mind]. A form of insanity characterized by a deterioration or loss of the intellectual faculties, the reasoning power, the memory, and the will. **D.**, Paralytic, general paralysis of the insane. **D.**, Primary, that occurring independently of other forms of insanity. **D.**, Secondary, that following another form of insanity. **D.**, Senile, that due to the degenerations of old age. **D.**, Terminal, that coming on toward the end of other forms of insanity or certain nervous diseases.

Demi- [*dimidius*, half]. A prefix meaning half.

Demilune Cells (*dem'-e-lun*) [*demius*, half; *luna*, moon]. **D. C. of Heidenhain**, crescentic bodies lying between the cells and the membrana propria of an acinus of a salivary gland.

Demodex (*dem'-o-deks*) [*δημος*, fat; *δῆξ*, an insect]. A genus of parasitic insects. **D. folliculorum**, the pimple-mite, a minute parasite found in the sebaceous follicles, par-

ticularly of the face. It probably does not produce any symptoms.

Demography (*de-mog'-ra-fe*). [*δῆμος*, the people; *γράφειν*, to write]. The science of peoples collectively considered; social science, including that of vital statistics and the consideration of questions of state medicine.

Demonomania (*de-man-o-ma'-ne-a-ly*) [*δαίμων*, a devil; *μανία*, madness]. A form of madness in which a person imagines himself possessed of a devil.

Demulcent (*de-mul'-sent*) [*demulcere*, to soothe]. 1. Soothing; allaying irritation of surfaces, especially mucous membranes. 2. A soothing substance, particularly a slippery, mucilaginous liquid.

Denarcotized (*de-nar'-ko-tized*) [*de* priv.; *ναρκωτικός*, narcotic]. 1. Deprived of narcotizing qualities. 2. Of opium, deprived of narcotin.

Dendritic (*den-dri'-ik*) [*δένδρον*, a tree]. Branching like a tree.

Dengue (*deng'-ga*) [West Ind.]. Break-bone fever; dandy fever; an acute, epidemic, infectious disease, characterized by a febrile paroxysm, severe pains in the bones, joints, and muscles, and, at times, a cutaneous eruption. The period of incubation is from 3 to 5 days; the invasion is sudden, with high fever (106° F.), severe pains in the muscles, bones and joints, the last being swollen and reddened. After the fever has lasted 3 or 4 days it subsides, but at the end of from 2 to 4 days a second paroxysm accompanied with pain occurs. Convalescence is slow; complications are rare.

Density (*den'-sit-e*) [*densitas*, thickness]. Closeness; compactness, especially the degree of closeness of one body compared with an equal volume of another taken as a standard; specific gravity. In electricity, the amount of electricity accumulated on a unit of surface during a given time.

Dental (*den'-tal*) [*dens*, a tooth]. Pertaining to the teeth. **D. Engine**, a machine worked by a treadle and possessing a flexible cable and adjustable arm and hand-piece, which afford great facility of movement and adaptation. By means of attachments to the hand-piece, drills can be operated at various angles. **D. Tubuli**, the minute wavy tubes occurring in the dentin of teeth.

Dentate (*den'-tat*) [*dens*, a tooth]. Toothed; having a toothed or serrated edge. **D. Body**. See *Corpus dentatum*. **D. Convolution**, a convolution found in the hippocampal fissure. **D. Fascia**, the serrated free edge of the dentate convolution. **D. Fissure**, the hippocampal fissure.

Denticulate (*den-tik'-u-lat*) [*denticulus*, a small tooth]. Having minute dentations; furnished with small teeth or notches.

Dentifrice (*den'-tif-ris*) [*dens*, tooth; *fricere*, to rub]. A substance for cleansing the teeth.

Dentigerous (*den-tij'-er-us*) [*dens*, a tooth; *gerere*, to carry]. Bearing or containing teeth, as a dentigerous cyst.

Dentin (*den'-tin*) [*dens*, a tooth]. A modified osseous tissue forming the principal part of a tooth, and consisting, histologically, of dental tubuli and intertubular tissue, chemically, of the phosphates of calcium and magnesium, the carbonate and fluorid of lime, and organic matter, chiefly gelatin. The bony structure of the tooth, lying under the enamel of the crown and the cement substance of the root.

Dentinal (*den'-tin-al*) [*dens*, a tooth]. Pertaining to or composed of dentin.

Dentist (*den'-tist*) [*dens*, a tooth]. One who practises dentistry.

Dentistry (*den'-tis-tre*) [*dens*, a tooth]. Dental surgery, embracing everything pertaining to the treatment of diseases of the teeth.

Dentition (*den'-tish'-un*) [*dens*, a tooth]. Teething; the cutting of the teeth. **D.**, **Primary**, the cutting of the temporary or milk-teeth. **D.**, **Secondary**, the eruption of the permanent teeth.

Denture (*den'-chûr*) [*dens*, a tooth]. The entire set or group of teeth; the whole assemblage of teeth in both jaws; a set, or plate, of artificial teeth.

Denudation (*den-u-da'-shun*) [*denudare*, to denude]. A stripping or making bare.

Denutrition (*de-nu'-trish'-un*) [*de*, from; *nutriare*, to nourish]. An atrophy and degeneration of tissue arising from lack of nutrition.

Deobstruent (*de-ob'-struent*) [*de*, from; *obstruere*, to obstruct]. **1.** Removing obstruction. **2.** A medicine that removes obstruction; an aperient.

Deodorant (*de-o'-dor-ant*) [*de* priv.; *odorare*, to smell]. **1.** Removing or concealing offensive odors. **2.** A substance that removes or conceals offensive odors.

Deodorized (*de-o'-dor-izd*) [*de* priv.; *odorare*, to smell]. Deprived of odor.

Deorsum (*de-or'-sum*) [*L.*]. Downward.

Deoxidation (*de-oks-id-a'-shun*) [*de*, from; *oxygen*]. The removal of the oxygen from a chemie compound.

Depilatory (*de-pil'-at-or-e*) [*de*, from; *pilus*, the hair]. **1.** Having the power to remove the hair. **2.** A substance used to destroy the hair, usually a caustic alkali.

Depletion (*de-ple'-shun*) [*deplere*, to empty]. **1.** The act of diminishing the quantity of fluid in the body or in a part, especially by bleeding. **2.** The condition of the system produced by the excessive loss of blood or other fluids.

Depolarization (*de-po-lar-is-a'-shun*) [*de*, from; *polus*, pole]. The neutralization of polarity.

Depressant (*de-pres'-ant*) [*deprimere*, to press down]. **1.** Lowering. **2.** A medicine that diminishes functional activity.

Depression (*de-presh'-un*) [*depressio*; *deprimere*, to depress]. **1.** A hollow, or fossa. **2.** Inward displacement of a part, as of the skull. **3.** Lowering of vital functions under the action of some depressing agent.

Depressomotor (*de-pres-o-mo'-tor*) [*deprimere*, to depress; *movere*, to move]. An agent that diminishes the action of the motor apparatus.

Depressor (*de-pres'-or*) [*deprimere*, to depress]. **1.** A muscle, instrument, or apparatus that depresses. **2.** A nerve, stimulation of which lowers the functional activity of a part, as the depressor nerve of the heart.

Depurated (*dep'-u-ra-ted*) [*depurare*, to purify]. Purified, cleansed.

Depurative (*dep'-u-ra-tiv*) [*depurare*, to purify]. Purifying or cleansing.

DeR. A contraction and symbol of the term *Reaction of Degeneration*.

Deradelphus (*der-ad-el'-fus*) [*δέρην*, neck; *ἀδελφός*, brother]. A monocephalic dual monstrosity, with fusion of the bodies above the umbilicus, and with four lower extremities, and three or four upper.

Derbyshire Neck (*der'-be-shir nek*). See *Goiter*.

Derencephalus (*der-en-sef'-al-us*) [*δέρην*, neck; *ἐνέφαλον*, brain]. A variety of single autostic monsters of the species anencephalus, in which the bones of the cranial vault are rudimentary, the posterior portion of the occiput absent, and the upper cervical vertebrae bifid, the brain resting in them.

Derivation (*der-iv-a'-shun*) [*derivare*, to turn a stream from its banks]. The drawing away of blood or liquid exudates from a diseased part by creating an extra demand for them in some other part.

Derivative (*de-riv'-at-iv*). **1.** Producing derivation. **2.** An agent that produces derivation.

Derm. **Derma** (*der'-mah*) [*δέρμα*, the skin]. The true skin.

Dermal (*der'-mal*) [*δέρμα*, the skin]. Pertaining to the skin.

Dermalgia (*der-mal'-je-ah*). See *Dermatalgia*.

Dermatagra (*der-mat-a'-grah*) [*δέρμα*, skin; *ἀγγρα*, seizure]. See *Pellagra*.

Dermatalgia (*der-mat-al'-je-ah*) [*δέρμα*, skin; *ἀλγος*, pain]. Pain in the skin unaccompanied by any structural change, and caused by some nervous disease or reflex influence.

Dermatitis (*der-mat-it'-is*) [*δέρμα*, skin; *ίτις*, inflammation]. An inflammation of the skin.

D. **ambustionis**, **D.** **calorica**, the form due to burns and scalds. **D.** **congelationis**, same as

Frostbite. **D. contusiformis**. Erythema nodosum. **D. exfoliativa**, an acute or chronic inflammation of the skin, in which the epidermis is shed more or less freely in large or small scales. See *Pityriasis rubra*. **D. gangrænosa**, sphaceloderma; gangrenous inflammation of the skin. **D. herpetiformis**, an inflammatory skin-disease of an herpetic character, the various lesions showing a tendency to group. It is a protean disease, appearing as erythema, vesicles, blebs, and pustules, and is associated with fever, itching, and burning. **D. medicamentosa**, drug-eruptions; inflammatory eruptions upon the skin due to the action of certain drugs taken internally. **D. papillaris capillitii**, a chronic skin-disease, affecting the nape of the neck and adjacent parts, and characterized by minute red papules, which occasionally suppurate, and are usually traversed by a hair. They unite to form hard, white, or reddish keloid-looking elevations, from which a bundle of atrophied hairs protrudes. **D. traumatica**, that resulting from traumatism. **D. venenata**, that produced by the local action of irritant substances.

Dermato- [*déppa*, skin]. A prefix, signifying pertaining to the skin.

Dermatol (*der'-mat-ol*) [*déppa*, skin], $C_6H_2(OH)_3CO_2Bi(OH)_2$. The subgallate of bismuth, an astringent, antiseptic powder, of yellow color, used in affections of the skin and mucous membranes that are associated with excessive secretion; it is especially recommended for diarrhea in tuberculosis and typhoid fever. Dose internally, 3 ss-3 iss (2.0-6.0) daily.

Dermatology (*der-mat-ol'-o-jé*) [*déppa*, skin; *lógos*, science]. The science of diseases of the skin.

Dermatolysis (*der-mat-ol'-is-is*) [*déppa*, skin; *lósis*, a loosing]. An hypertrophy of the skin and subcutaneous tissue, with a tendency to the formation of folds.

Dermatome (*der'-ma-tôm*) [*déppa*, skin; *τέμνω*, to cut]. An instrument for incising the skin.

Dermatomycosis (*der-mat-o-mi-kô'-sis*) [*déppa*, skin; *μύκης*, fungus]. Any skin-disease caused by a vegetable parasite.

Dermatoplasty (*der'-mat-o-plas-te*) [*déppa*, skin; *πλάσσειν*, to form]. An operative replacement of destroyed skin by means of flaps or skin-grafts.

Dermatosclerosis (*der-mat-o-skle-ro'-sis*). See *Scleroderma*.

Dermatosis (*der-mat-ô'-sis*) [*déppa*, skin; *νόσος*, disease]. Any disease of the skin.

Dermic (*der'-mik*) [*déppa*, the skin]. Relating to the skin, or formed of skin.

Dermis (*der'-mis*). The corium or true skin.

Dermographia (*der-mo-graf'-ô-ah*) [*déppa*,

skin; *γράφειν*, to write]. A condition of the skin in which tracings made with the fingernail or a blunt instrument are followed by elevations at the points irritated. It is common in the condition termed vasomotor ataxia.

Dermoid (*der'-moid*) [*déppa*, skin; *εἶδος*, form]. Resembling skin. **D. Cyst**, a cyst containing elements of the skin, as hair, teeth, etc.

Dermoplasty (*der'-mo-plas-te*) [*déppa*, skin; *πλάσσειν*, to form]. See *Dermatoplasty*.

Dermoskeleton (*der-mo-skel'-et-un*) [*déppa*, skin; *σκελετόν*, a skeleton]. The exoskeleton.

Derodidymus (*der-o-did'-im-us*) [*déppa*, neck; *δίδυμος*, double]. A monstrosity with a single body, two necks and heads, two upper and lower extremities, with other rudimentary limbs occasionally present.

Descemet's Membrane. The elastic membrane lining the posterior surface of the cornea.

Descemetitis (*des-em-et-i'-tis*). Inflammation of Descemet's Membrane; serous iritis.

Descendens (*de-sen'-dens*) [*descendere*, to go down]. Downward. **D. noni**, a branch of the hypoglossal nerve. See *Nerves*, Table of.

Descending (*de-sen'-ding*) [*descendere*, to go down]. Passing downward. **D. Current**.

See *Current*. **D. Degeneration**. See *Degeneration*. **D. Tract**, a collection of nerve-fibers conducting impulses from the centers to the periphery.

Deshler's Salve. The compound resin cerate of the pharmacopeia.

Desiccant (*des-ik-ant*) [*desiccare*, to dry up]. Drying.

Desiccation (*des-ik-ant'-shun*) [*desiccare*, to dry up]. The process of drying.

Desmobacteria (*des-mo-bak-te'-re-ah*) [*δέσμος*, band; *βακτήριον*, a small staff]. A group of bacteria corresponding to the genus Bacilli.

Desmography (*des-mog'-ro-fe*) [*δέσμος*, a ligament; *γράφειν*, to write]. The description of the ligaments.

Desmoid (*des'-moid*) [*δέσμη*, a bundle; *εἶδος*, likeness]. Fibrous.

Desmoma (*des-mô'-mah*) [*δέσμος*, a band; *δύνα*, tumor]. A connective-tissue tumor.

Despumation (*des-fu-mat'-shun*) [*despumare*, to skim froth]. The purification of a liquid by removal of the scum or froth.

Desquamation (*des-kvam'-at'-shun*) [*desquamare*, to scale off]. The shedding of the superficial epithelium, as of the skin, mucous membranes, and renal tubules.

Desquamative (*des-kvam'-at-iv*) [*desquamare*, to scale off]. Characterized by desquamation.

Dessert Spoon. A domestic measure equal to about $\frac{3}{4}$ ij (8.0).

Detergent (*de-ter'-jent*) [*detergere*, to cleanse]. Purifying; cleansing.

Determination (*de-ter-min-a'-shun*) [*deter-minatio*, a directing]. Of the blood, a tendency to collect in a part, as determination of the blood to the head.

Deuter-, Deutero- (*du'-ter, du'-ter-o*) [*δευτερο*, second]. A Greek prefix indicating the second of two similar substances, or conditions, especially that one which contains more of the substance.

Deuteroalbumose (*du-ter-o-al'-bu-môs*). A form soluble in water and not precipitated by saturation with sodium chlorid or magnesium sulphate, but by ammonium sulphate.

Deuteropathy (*du-ter-op'-a-the*) [*δευτερος*, second; *πάθος*, a disease]. A disease that is secondary to another.

Deviation (*de-ve-a'-shun*) [*deviare*, to deviate]. A turning away from the regular course or standard. **D.**, Conjugate, the turning of eyes and head toward one side, observed in some lesions of the cerebrum.

Devitalize (*de-vi'-tal-iz*) [*de*, from; *vita*, life]. To destroy vitality.

Devonshire Colic (*dev'-on-shir kol'-ik*). Lead-colic.

Dew-point. The temperature at which the air is saturated with vapor; the temperature at which dew forms.

Dewees's Carminative. See *Asafetida*.

Dextiocardia (*deks-e-o-kar'-de-ah*) [*δεξιός*, on the right; *καρδια*, the heart]. Transposition of the heart to the right side of the thorax.

Dextrad (*deks'-trad*) [*dexter*, right; *ad*, to]. Toward the right side.

Dextran (*deks'-tran*) [*dexter*, right], $C_6H_{10}O_5$. A stringy, gummy substance formed in milk by the action of cocci, and also occurring in unripe beet-root.

Dextrin (*deks'-trin*) [*dexter*, right], $C_6H_{10}O_5$. A soluble carbohydrate into which starch is converted by diastase or dilute acids. It is a whitish substance, turning the plane of polarization to the right.

Dextrocardia (*deks-tro-kar'-de-ah*). See *Dextiocardia*.

Dextrogyrate (*deks-tro-jiv'-rät*). Same as *Dextrorotatory*.

Dextrorotatory (*deks-tro-ro'-lat-o-re*) [*dexter*, right; *rotare*, to whirl]. Turning the rays of light to the right.

Dextrose (*deks'-tros*) [*dexter*, right], $C_6H_{12}O_6$. Grape-sugar. A sugar belonging to the glucose-group, that rotates polarized light to the right.

Dextroversion (*deks-tro-ver'-zhun*) [*dexter*, right; *vertere*, to turn]. Version to the right side.

Diabetes (*di-ab-e'-tez*) [*διαβήτης*; *διά*, through; *βαίνειν*, to go]. A disease characterized by the habitual discharge of an excessive quantity of urine; used without qualification, the word indicates diabetes mellitus. **D.**, Bil-

iary. See *Biliary*. **D. decipiens**, diabetes mellitus in which there is no polyuria or polydipsia. **D.**, Gouty, a form of glycosuria occurring in gouty individuals. **D. insipidus**, a chronic disease characterized by the passage of a large quantity of normal urine of low specific gravity, associated with intense thirst. The etiology of the disease is obscure—injuries and diseases of the brain, emotional disturbances, syphilis, acute infectious diseases, and heredity have been assigned as causes. The urine at times contains inosit, whence the term diabetes inositus in contradistinction to diabetes mellitus. **D. mellitus**, a nutritional disease characterized by the passage of a large quantity of urine containing sugar; there is intense thirst, voracious appetite, progressive loss of flesh and strength, and a tendency to a fatal termination. The causes are not well understood—diseases of the nervous system, especially of the medulla, intense mental strain, gout, syphilis, malaria, diseases of the pancreas, have been etiologic factors in some cases. Complications are frequent in diabetes, the most important being pulmonary tuberculosis, furunculosis, gangrene, cataract, polyneuritis. Death, if not caused by one of the complications, is generally due to the so-called diabetic coma. The ultimate cause of the presence of sugar in the urine is believed to be the incomplete destruction or utilization of the sugar formed in the system. **D.**, Pancreatic, a variety of glycosuria associated with and probably dependent upon disease of the pancreas. **D.**, Phloridzin-, that form produced in animals by the administration of phloridzin. **D.**, Phosphatic, a condition characterized by polyuria, polydipsia, emaciation, and excessive excretion of phosphates in the urine.

Diabetic (*di-ab-et'-ik*) [*διά*, through; *βαίνειν*, to go]. 1. Pertaining to diabetes. 2. A person suffering from diabetes. **D. Cataract**, an opacity of the crystalline lens sometimes found in association with diabetes. **D. Coma**, the coma caused by diabetes mellitus. **D. Gangrene**, a moist gangrene sometimes occurring in persons suffering from diabetes. **D. Sugar**, $C_6H_{12}O_6$, the glucose present in the urine in diabetes mellitus. It is identical with grape-sugar. **D. Tabes**, a peripheral neuritis occurring in diabetic patients, and causing symptoms resembling tabes dorsalis. **D. Urine**, urine containing sugar.

Diacetic Acid (*di-as-e'-tik*). See *Acid*.

Diaceturia (*di-as-et-u'-re-ah*) [*di*, two; *acetum*, vinegar; *οὐρον*, urine]. The presence of diacetic acid in the urine.

Diachylon (*di-ak'-il-on*) [*διά*, through; *χυμός*, juice]. Lead-plaster. See *Plumbum*.

Diacid (*dī-as'-id*) [*dī*, two; *acidus*, acid]. Having two atoms of hydrogen replaceable by a base.

Diad (*dī'-ad*) [*dī*, two]. Having a quantivalence of two.

Diagnose (*dī-ag-nōs'*). See *Diagnosticate*.

Diagnosis (*dī-ag-nōs'-sis*) [*diá*, apart; *γνώσις*, knowledge]. The determination of the nature of a disease. **D.**, **Differential**, the distinguishing between two diseases of similar character, by comparing their symptoms. **D.** by **Exclusion**, the recognition of a disease by excluding all other known conditions. **D.**, **Pathologic**, the diagnosis of the structural lesions present in a disease. **D.**, **Physical**, the determination of disease by inspection, palpation, percussion, or auscultation.

Diagnostic (*dī-ag-nōs'-tik*) [*diá*, apart; *γνώσις*, knowledge]. Serving as evidence in diagnosis.

Diagnosticate, Diagnose (*dī-ag-nōs'-tik-at*, *dī'-ag-nōs*) [*diá*, apart; *γνώσις*, knowledge]. To make a diagnosis.

Diagnostician (*dī-ag-nōs'-tish'-an*) [*diá*, through; *γνώσις*, knowledge]. One skilled in making diagnoses.

Dialysis (*dī-al'-is-is*) [*diá*, through; *λίσσω*, to loose]. The separation of several substances from each other in solution by taking advantage of their differing diffusibility through porous membranes. Those that pass through readily are termed *crystalloids*, those that do not, *colloids*.

Dialyzed (*dī'-al'-īzd*) [*diá*, through; *λίσσω*, to loose]. Separated by dialysis.

Dialyzer (*dī'-al'-ī-zer*) [*diá*, through; *λίσσω*, to loose]. An apparatus for effecting dialysis; also the porous septum or diaphragm of such an apparatus.

Diamagnetic (*dī-ah-mag-net'-ik*) [*diá*, across; *μάγνης*, magnet]. Taking a position at right angles to the lines of magnetic force.

Diameter (*dī-am'-et-er*) [*diá*, through; *μέτρον*, a measure]. A straight line joining opposite points of a body or figure and passing through its center. **D.**, **Cranio-metric**, one of several lines connecting points on opposite surfaces of the cranium. *Biparietal*, that joining the parietal eminences; *Bitemporal*, that joining the extremities of the coronal suture; *Occipitofrontal*, that joining the root of the nose and the most prominent point of the occiput; *Occipitomen-tal*, that joining the external occipital protuberance and the chin; *Trachelobregmatic*, that joining the center of the anterior fontanel and the junction of the neck and floor of the mouth. **D.**, **Pelvic**, any one of the diameters of the pelvis. The most important are the following: *Anteroposterior* (of pelvic inlet), that which joins the sacrovertebral angle and the pubic sym-

physis; *Anteroposterior* (of pelvic outlet), that which joins the tip of the coccyx with the subpubic ligament; *Conjugate*, the *anteroposterior diameter* of the pelvic inlet; *Conjugate, diagonal*, that connecting the sacrovertebral angle and subpubic ligament; *Conjugate, external*, that connecting the depression above the spine of the first sacral vertebra and the middle of the upper border of the symphysis pubis; *Conjugate, true*, that connecting the sacrovertebral angle and the most prominent portion of the posterior aspect of the symphysis pubis; *Transverse* (of pelvic inlet), that connecting the two most widely separated points of the pelvic inlet; *Transverse* (of pelvic outlet), that connecting the ischial tuberosities.

Diamid (*dī'-am-id*) [*dī*, two; *amid*]. A double amid formed by replacing hydrogen in two ammonia molecules by an acid radicle.

Diamin (*dī'-am-in*) [*dī*, two; *amin*]. An amin formed by replacing hydrogen in two molecules of ammonia by a basic radicle.

Diapedesis (*dī-ah-ped'-ē-sis*) [*diá*, through; *πηδάω*, to leap]. The passage of the blood through the unruptured vessel walls.

Diaphanometer. See *Lactoscope*.

Diaphanoscope (*dī-af-an'-o-skōp*) [*διαφανής*, translucent; *σκοπεῖν*, to inspect]. An instrument for illuminating the interior of a body-cavity so as to render the boundaries of the cavity visible from the exterior.

Diaphoresis (*dī-ah-for'-ē-sis*) [*diá*, a part; *φορέω*, to carry]. Perspiration, especially perceptible perspiration.

Diaphoretic (*dī-ah-for'-ē-ik*) [*diá*, a part; *φορέω*, to carry]. 1. Causing an increase of perspiration. 2. A medicine that induces diaphoresis.

Diaphragm (*dī'-ah-frag*) [*diá*, across; *φράγμα*, wall]. 1. The musculomembranous partition that separates the thorax and abdomen. It is the chief muscle of respiration. 2. A thin septum, such as is used in dialysis. 3. In a microscope, a perforated plate placed between the mirror and object to regulate the amount of light that is to pass through the object.

Diaphragmatic (*dī-ah-frag-mat'-ik*) [*diá*, across; *φράγμα*, a wall]. Relating to the diaphragm. **D.** **Phenomenon**, Litten's sign; in a state of health there can be seen a shadow rising and falling from the vertebral column to the attachment of the diaphragm from the seventh rib to the convexity; this movement can be seen through the thoracic walls, and shows the intensity of respiration, and the limit of the diaphragm between its position on inspiration and that on expiration. The upper position corresponds to the liver-margin in the state of rest. Deviations of the extent of movement mark certain pathologic states.

Diaphtherin (*di-af'ther-in*) [*διαφθερίν*, to destroy]. Oxychinaseptol; a coal-tar derivative composed of two molecules of oxychinolin and one of aseptol. It is a yellow powder, with a phenol-like odor, and is used as an antiseptic in solutions varying in strength from 1 to 50 per cent.

Diaphysis (*di-af'-is-is*) [*διά*, between; *φύειν*, to grow]. The shaft of a long bone.

Diapophysis (*di-ap'-off'-is-is*) [*διά*, apart; *ἀποφύσις*, an outgrowth]. The superior or articular part of a transverse process of a vertebra.

Diarrhea (*di-ar-e'-ah*) [*διά*, through; *ῥέειν*, to flow]. A condition characterized by increased frequency and lessened consistency of the fecal evacuations. **D.**, Choleraic, severe, acute diarrhea with serous stools, and accompanied by vomiting and collapse. **D.**, Lienteric, a form of diarrhea characterized by the passage of fluid stools containing masses of undigested food. **D.**, Mucous, that characterized by the presence of mucus in the stools. **D.**, Summer, an acute form occurring particularly in children during the intense heat of summer.

Diarthrosis (*di-ar-thro'-sis*) [*διά*, through-out; *ἀρθρωσις*, articulation]. A form of articulation characterized by mobility. The various forms are: *Arthrodia*, in which the bones glide upon plane surfaces; *Enarthrosis*, ball-and-socket joint, with motion in all directions; *Ginglymus*, or hinge joint, with backward and forward motion; and *D. rotatoria*, with pivotal movement.

Diastase (*di'-as-tās*) [*διάστασις*, separation]. A nitrogenous vegetable ferment found in malt; it converts starch into glucose.

Diastasic, or **Diastatic** (*di-as-tas'-ik*, or *di-as-tat'-ik*) [*διάστασις*, separation]. 1. Pertaining to diastase. 2. Pertaining to diastasis.

Diastasis (*di-as'-tas-is*) [*διάστασις*, separation]. 1. The separation of an epiphysis from the body of a bone without true fracture. 2. A dislocation of an amphiarthrotic joint.

Diaster (*di'-as-ter*). See *Karyokinesis*.

Diastole (*di-as'-to-le*) [*διαστολή*, a drawing apart]. The period of dilatation of a chamber of the heart; used alone it signifies D. of the ventricles.

Diastolic (*di-as-tole'-ik*) [*διαστολή*, a drawing apart]. Pertaining to the diastole of the heart.

Diathermal, or **Diathermanous** (*di-ath-er'-mal*, *di-ath-er'-man-us*) [*διά*, through; *θέρμην*, heat]. Permeable by waves of radiant heat.

Diathesis (*di-ath'-es-is*) [*διά*, through; *ῥθί-ναι*, to arrange]. A state or condition of the body whereby it is especially liable to certain diseases, such as gout, calculus, dia-

betes, etc. It may be acquired or hereditary. **D.**, Hemorrhagic. See *Hemophilia*.

Diathetic (*di-ath-et'-ik*) [*διά*, through; *ῥθί-ναι*, to arrange]. Relating to a diathesis.

Diatomic (*di-at-om'-ik*) [*δίς*, two; *ἄτομος*, atom]. 1. Consisting of two atoms. 2. Having two hydrogen atoms replaceable by radicles.

Diazo- (*di-az'-o*) [*δίς*, two; *αζότιον*, nitrogen]. A prefix signifying that a compound contains phenyl, C_6H_5 , united with a radicle consisting of two nitrogen atoms. **D.-reaction** or **test**, a urinary test, valuable in the diagnosis of enteric fever. The solutions required are (a) a saturated solution of sulphuric acid in a 5 per cent. solution of hydrochloric acid; (b) a $\frac{1}{2}$ per cent. solution of sodium nitrite. Mix a and b in the proportion of 40 c.c. of a to 1 c.c. of b, and to a few cubic centimeters add an equal volume of urine, and, after shaking well, allow a few drops of ammonia to flow down the side of the tube. A garnet-red color at the point of contact denotes the reaction, or a rose-pink foam after shaking.

Dibasic (*di-ba'-sik*) [*δίς*, twice; *βάσις*, base]. Of a salt, containing two atoms of a monobasic element or radicle; of an acid, having two replaceable hydrogen atoms.

Dibromid (*di-brom'-id*). A compound consisting of an element or radicle and two atoms of bromine.

Dichloroacetic Acid (*di-klor-as-e'-tik*), $C_2H_3Cl_2O_2$. An acid produced by the replacement of two hydrogen atoms in acetic acid by chlorine. It is a caustic liquid.

Dichlorid (*di-klo'-rid*). A compound consisting of an element or radicle and two atoms of chlorine.

Dichroism (*di'-kro-izm*) [*δίς*, double; *χρῶμα*, color]. The state of presenting one color when seen by reflected light, and another when seen by transmitted light.

Dicinchonin (*di-sin'-ken-in*) [*δίς*, double; *cinchona*], $C_{20}H_{14}N_4O_2$. An alkaloid of Cinchona bark.

Dicrotic (*di-krot'-ik*) [*δίκροτος*, double beating]. Having a double beat. **D.** Pulse. See *Dicrotism*. **D.** Wave, the recoil-wave of the sphygmographic tracing, generated by closure of the aortic valves.

Dicrotism (*di'-kro-tizm*) [*δίκροτος*, a double-beat]. A condition of the pulse in which with every wave there is given to the finger of the examiner the sensation of two beats. It is present when the arterial tension is low.

Didactic (*di-dak'-tik*) [*διδασκτικός*, apt at teaching]. Teaching by description and theory.

Dielectric (*di-el-ek'-trik*) [*διά*, through; *ἤλεκτρον*, amber]. Transmitting electricity by induction and not by conduction.

Diet (*dī'-et*) [*diata*, a system or mode of living]. The food taken regularly by an individual; the food adapted to a certain state of the body, as fever-diet, convalescence-diet. **D., Diabetic**, a diet mostly of meats and green vegetables, starches and sugars being excluded. **D., Fever**, a nutritious, easily digestible liquid or semi-liquid diet, usually with milk and meat-broths as a basis. **D., Gouty**, a diet of simple nutritious food, avoiding wines, fats, pastries, and much meat. **D., Tuffnell's**, a highly nutritious diet including but a small amount of liquids, employed in the treatment of aneurysm.

Dietary (*dī'-et-a-re*) [*diata*, a system or mode of living]. A system of food-regulation intended to meet the requirements of the animal economy.

Dietetic (*dī'-et-ik*) [*diata*, a system or mode of living]. Pertaining to diet.

Diethylamin (*dī-eth-il'-am-in*) NC_2H_5 . A non-poisonous, liquid atomain obtained from putrefying fish.

Diethylene-diamin. See *Piperazin*.

Differential (*dif'-er-en'-shal*) [*differentia*, difference]. Pertaining to or creating a difference. **D. Diagnosis**. See *Diagnosis*. **D. Staining**, a method of staining tubercle-bacilli, syphilis-bacilli, etc., founded upon the fact that they retain the color in the presence of certain reagents that decolorize the surrounding tissues.

Differentiation (*dif'-er-en-she-a'-shun*) [*differentia*, difference]. 1. The act or process of distinguishing or making different. 2. Changing from general to special characters; specialization.

Diffraction (*dif'-rak'-shun*) [*dis*, apart; *fractus*, broken]. The deflection or the separation into its component parts that takes place in a ray of light when it passes through a narrow slit or aperture. **D. Grating**, a strip of glass closely ruled with fine lines; it is often used in the spectroscope in the place of the battery of prisms.

Diffuse (*dif'-ūs'*). Scattered; not limited to one tissue or spot; opposed to localized.

Diffusible (*dif'-ū-zibl*) [*dis*, away; *fundere*, to pour]. Spreading rapidly; capable of passing through a porous membrane; applied to certain quickly-acting stimulants, usually of transient effect.

Diffusion (*dif'-ū-shun*) [*diffundere*, to spread]. A spreading-out. **D.-circle**, the imperfect image formed by incomplete focalization, the position of the true focus not having been reached by some of the rays of light or else having been passed.

Digastric (*dī-gas'-trik*) [*dis*, double; *gastrop*, belly]. Having two bellies, as the D. Muscle.

Digestant (*dī-jest'-ant*) [*digerere*, to digest]. A substance that assists digestion of the food.

Digestion (*dī-jes'-chun*) [*digerere*, to digest].

Those processes whereby the food taken into an organism is made capable of being absorbed and assimilated by the body-tissues. **D., Artificial**, digestion carried on outside of the body. **D., Gastric**, digestion by the action of the gastric juice. **D., Intestinal**, digestion by the action of the intestinal juices, including the action of the bile and the pancreatic fluid. **D., Pancreatic**, digestion by the action of the pancreatic juice. **D., Peptic**. See *D., Gastric*. **D., Primary**, gastrointestinal digestion. **D., Salivary**, digestion by the saliva. **D., Secondary**, the assimilation by the body-cells of their appropriate pabulum.

Digestive (*dī-jes'-tiv*) [*digerere*, to digest].

1. Relating to or favoring digestion. 2. An agent that promotes digestion. **D. Tract**, the whole alimentary canal from the mouth to the anus.

Digit (*dij'-it*) [*digitus*, finger]. A finger or toe.

Digital (*dij'-it-al*) [*digitus*, a finger]. Pertaining to the fingers or toes. **D. Arteries**, the arteries of the hands and feet supplying the digits. See *Arteries*, *Table of*. **D. Compression**, the stoppage of a flow of blood by pressure with the finger. **D. Examination**, examination or exploration with the finger.

Digitalin (*dij'-it-a'-lin*) [*digitalis*, pertaining to the fingers]. 1. *Digitalinum* (U. S. P.); $\text{C}_8\text{H}_8\text{O}_4$ (?), the active principle of *Digitalis purpurea*. Dose gr. $\frac{1}{80}$ – $\frac{1}{10}$ (0.001–0.002). 2. A precipitate from a tincture of *Digitalis purpurea*. Unof.

Digitalis (*dij'-it-a'-lis*) [*digitalis*, pertaining to the fingers]. Foxglove. The leaves of *D. purpurea*. It contains an amorphous complex substance, digitalin, that does not, however, represent the full properties of the leaves. It is a powerful cardiac stimulant, strengthening the systole and lengthening the diastole of the heart. It also acts as a diuretic; in large doses it causes gastric disturbances. It is employed mainly in diseases of the heart when compensation is lost. **D., Extractum**, Dose gr. $\frac{1}{4}$ – $\frac{1}{2}$ (0.01–0.032). **D., Extractum, Fluidum**, Dose m i–iii (0.065–0.2). **D., Infusum**, $1\frac{1}{2}$ parts of the leaves in 100 parts. Dose f $\frac{3}{4}$ ss– f $\frac{3}{4}$ j (16.0–32.0). **D., Tinctura**, 15 parts of the leaves in 100 of diluted alcohol. Dose m x–xx (0.6–1.2).

Digitation (*dij'-it-a'-shun*) [*digitatus*, having digits]. A finger-like process, or a succession of such processes, especially of a muscle.

Digitus mortui (*dij'-it-i mor'-tu-i*) [*1.*].

Dead fingers; a cold and white state of the fingers.

Digitonin (*dij'-it-e'-nin*) [*digitalis*, pertaining to the fingers]. $\text{C}_{31}\text{H}_{52}\text{O}_{17}$. A white, amorphous mass obtained from digitalis.

Digitoxin (*dij'-it-aks'-in*) [*digitalis*, pertaining to the fingers; *toxikon*, poison]. $\text{C}_{31}\text{H}_{52}$

O_2 , or $C_{12}H_{22}O_{24}$. A highly poisonous glucosid obtained from *Digitalis purpurea*.

Dihydrocollidin (*di-hi-dro-kol'id-in*), $C_8H_{18}N$. A liquid substance isomeric with a ptomain obtained from putrid flesh and fish.

Dihydrocoridin (*di-hi-dro-kor'id-in*), $C_{10}H_{17}N$. A substance isomeric with a ptomain found in cultures of the *Bacillus alli*.

Dihydrolutidin (*di-hi-dro-lu'id-in*) [*dis*, double; *idrop*, water; *luteus*, yellow], $C_7H_{11}N$. One of the alkaloidal bodies found in cod-liver oil. It is slightly poisonous, in small doses diminishing general sensibility, in large doses causing tremor, paralysis of the hind limbs, and death.

Diiodid (*di-i'o-did*) [*di*, two; *iodum*, iodine]. A compound consisting of a basic element and two atoms of iodine.

Dilaceration (*di-las-er-a'-shun*) [*dilaceratio*, a tearing apart]. A tearing apart; division of a membranous cataract by a tearing operation.

Dilatation (*di-lat-a'-shun*) [*dilatare*, to spread out]. A spreading apart; the state, especially of a hollow part or organ, of being dilated or stretched. **D. of Heart**, an increase in the size of one or more of the cavities of the heart, arising from a relaxation or weakening of the heart-muscle. It is associated with evidences of failure of circulation, resulting in congestion of the lungs and other viscera.

Dilatator (*di-lat-a'-tor*). See *Dilator*.

Dilator (*di-la'-tor*) [*dilatare*, to spread out]. An instrument for stretching or enlarging a cavity or opening; also, a dilating muscle. See *Muscle*, *Table of*. **D., Barnes'**, an instrument for dilatation of the os and cervix uteri, consisting of a rubber bag, which, after insertion, is distended with water.

Dill (*dil*). See *Anethum*.

Diluent (*dil'-u-ent*) [*diluere*, to wash away]. 1. Diluting. 2. An agent that dilutes the secretions of an organ.

Dilution (*di-lu'-shun*) [*diluere*, to wash away]. 1. The process of adding a neutral fluid to some other fluid or substance, in order to diminish the qualities of the latter. 2. A diluted substance; the result of a diluting process.

Dimethylamin (*di-meth-il'-am-in*) [*dis*, two; *methyl*, *amin*], NC_2H_7 . A nontoxic ptomain found in putrefying gelatin, old decomposing yeast, etc.

Dimethylxanthin (*di-meth-yl'-san'-thin*), $C_8H_3(CH_3)_2N_4O_9$. Theobromin.

Dimorphous (*di-mor'-fus*) [*dis*, double; *μωρφή*, form]. Existing in two forms.

Dinitrate (*di'-ni-trat*). A compound resulting from the replacement of the hydrogen of two molecules of nitric acid by a base.

Dinitrocellulose (*di-ni-tro-sel'-u-lös*). See *Pyroxilin*.

Dioptr (*di-op'-ter*) [*diá*, through; *δψωθαι*, to see]. The unit of measurement of the refractive power of an optic lens. It is the refractive power of a lens that has a focal distance of one meter.

Dioptric (*di-op'-trik*) [*diá*, through; *δψωθαι*, to see]. 1. Pertaining to transmitted and refracted light. 2. A diopter.

Dioptry (*di-op'-tre*). See *Diopter*.

Dioxid (*di-ok's-id*) [*dis*, two; *δξιν*, sharp]. A molecule containing two atoms of oxygen.

Dip. The deviation from the horizontal position shown by a freely-suspended magnetic needle.

Diphtheria (*dif-thel'-re-ah*) [*διφθερα*, a skin or membrane]. An acute infectious disease caused by the Klebs-Löffler bacillus. It is characterized by the formation on a mucous membrane, most frequently that of the pharynx, of a false membrane, grayish or buff in color and quite firmly adherent. Any mucous membrane, as the laryngeal, nasal, conjunctival, and, more rarely, the gastrointestinal, vaginal, and that of the middle ear, may be the seat of the disease. The membrane may also be formed on wounds—*Surgical or Wound-D*. The constitutional symptoms result from the absorption of poisons elaborated by the bacteria, the bacilli themselves not being found in the blood as a rule. These poisons are a nuclealbumin and a nuclein, the former producing the acute, the latter the later symptoms of the disease. The general symptoms are, in brief, fever, profound prostration, weakness of the heart, and a rapidly-developing anemia. A frequent sequel is paralysis, which may be general, but is often confined to the palatal and ocular muscles. Diphtheria is eminently a disease of childhood and very fatal. Death results from systemic poisoning, from heart-failure, from obstruction to respiration by the presence of membrane in the air-passages, or from a complication, as pneumonia. **D., Brétonneau's**, true diphtheria of the pharynx, first described by P. Brétonneau (1826).

Diphtheric (*dif-thel'-ik*) [*διφθερα*, a skin or membrane]. Of or pertaining to diphtheria.

Diphtheritic (*dif-thel'-ik*). See *Diphtheric*.

Diphthongia (*dif-thon'-je-ah*) [*dis*, double; *φθόγγος*, a voice]. The production of a double tone of the voice due to incomplete unilateral paralysis of the recurrent laryngeal nerve, or to some lesion of the vocal bands that causes each to produce its own sound.

Diaplacusis (*dip-lak-u'-sis*) [*διπλωος*, double; *ἀκουσις*, hearing]. 1. The hearing of a tone as higher by one ear than by the other; called *D. binauralis*. 2. The hearing of two tones by one ear, when only one tone is produced; called *D. uniauralis*.

Diplegia (*di-ple'-je-ah*) [*dis*, double; *πληγή*,

stroke]. Paralysis of similar parts on the two sides of the body.

Diplo- [*διπλός*, double]. A prefix signifying double.

Diploblastic (*dip'-lo-blas'-tik*) [*διπλός*, double; *βλαστός*, sprout]. Having two germ-layers.

Diplococcus (*dip'-lo-kok'-us*) [*διπλός*, double; *κόκκος*, kernel]. A micrococcus that occurs in groups of two.

Diploe (*dip'-lo-e*) [*διπλοή*, a fold]. The cancellous bony tissue between the outer and inner tables of the skull.

Diploetic (*dip'-lo-et'-ik*) [*διπλοή*, a fold]. Relating to the diploe; diploic.

Diploic (*dip'-lo'-ik*) [*διπλοή*, a fold]. Pertaining to the diploe.

Diplomyelia (*dip'-lo-mi-e'-le-ah*) [*διπλός*, double; *μυελός*, marrow]. An apparent doubleness of the spinal cord, produced by a longitudinal fissure.

Diplopia (*dip'-lo-pe-ah*) [*διπλός*, double; *ὄψις*, sight]. Double vision, one object being seen by the eye or eyes as two. **D.**,

Binocular, the most frequent, is due to a derangement of the muscular balance, the images of the object being thereby thrown upon nonidentical points of the retina. **D.**,

Crossed, or **D.**, **Heteronymous**, the result of divergent strabismus, the image of the right eye appearing upon the left side, and that of the left eye upon the right side. **D.**, **Di-**

rect, or **D.**, **Homonymous**, the reverse of Crossed **D.**, found in convergent strabismus. **D.**, **Monocular**, diplopia with a single eye,

usually due to hysteria, or to double pupil, or beginning cataract.

Dipotassic (*di-po-tas'-ik*) [*di*, two; *potassium*]. Containing two atoms of potassium in a molecule.

Dippel's Animal Oil. *Oleum cornu cervi*. An oil obtained in distilling bone and deer's horn. It contains pyridin and lutidin. It is antispasmodic and stimulant. **Unof.**

Dipping Needle. A magnetic needle so hung that it can move freely in a vertical plane.

Dipsomania (*dip'-so-ma'-ne-ah*) [*δίψα*, thirst; *μανία*, madness]. The uncontrollable desire for spirituous liquors.

Dipterocarpus (*dip'-ter-o-kar'-pus*) [*διπτερος*, two-winged; *καρπός*, fruit]. A genus of trees chiefly found in Southern Asia, some of which furnish gurjun-balsam.

Direct (*di-rekt'*) [*directus*, straight]. In a right or straight line; without the interposition of some medium. **D.** **Current**, a galvanic current. **D.** **Image**. See *Image*. **D.**

Ophthalmoscopy. See *Ophthalmoscopy*. **D.** **Vision**, the perception of an object the image of which falls upon the macula.

Director (*di-rek'-tor*) [*dirigere*, to guide].

Anything that guides or directs. **D.**, **Grooved**, an instrument grooved to guide the knife in surgical operations.

Dis- [*dis*, twice]. 1. A prefix denoting two or double. 2. A prefix to denote apart from.

Disarticulation (*dis-ar-tik-u-la'-shun*) [*dis*, apart; *articulum*, a joint]. Separation at a joint; amputation at a joint.

Disassimilation (*dis-as-sim-il-a'-shun*) [*dis-assimilatio*]. The process of transformation of assimilated substances into waste products.

Disc (*disk*) [*discus*, *διακος*, a quoit or round plate]. A circular, plate-like organ or structure. **D.**, **Blood**. See *Blood-corporuscle*. **D.**,

Choked. See *Papillitis*. **D.**, **Germinal**, the small disc of the blastodermic membrane, in which the first traces of the embryo are seen. **D.**, **Optic**, the circular area in the retina that represents the termination of the optic nerve.

Discharge (*dis-charj'*) [*OF.*, *descharger*, to unload]. 1. A setting free; also that which is discharged. 2. A sudden setting free of a force.

Discharger (*dis-char'-jer*). An instrument for setting free electricity stored in a Leyden jar or other condenser.

Discharging (*dis-char'-jing*) [*OF.*, *descharger*, to unload]. Unloading; flowing out, as pus, etc. **D.** **Lesion**, a brain-lesion that causes sudden discharges of nervous motor impulses.

Dissection (*dis-ish'-un*) [*discissio*; *discindere*, to tear or cut apart]. An operation for soft cataract, in which the capsule is lacerated a number of times, to allow the lens-substance to be absorbed.

Discrete (*dis-kret'*) [*discretus*, separated]. Not running together; separate.

Discus (*dis'-kus*) [*διακος*, a disc]. A disc. **D.** **proligerus**, the mass of cells of the membrana granulosa of the Graafian vesicle that surround the ovum.

Discutient (*dis-ku'-shent*) [*discutere*, to shake apart]. 1. Capable of effecting resolution. 2. A medicine having the power of causing an exudation to disappear.

Disdiacblast (*dis-di'-ak last*) [*dis*, double; *διά*, through; *κλάνειν*, to break down]. One of the small doubly-refractive elements in the contractile discs of a muscle-fiber.

Disease (*dis-iz'*) [*dis*, negative; *ease*, a state of rest]. A disturbance of function or structure of any organ or part of the body. A *Table of Eponymic Diseases* is appended.

D., **Acute**, a disease marked by rapid onset and short course. **D.**, **Bleeders'**, synonym of *Hemophilia*. **D.**, **Chronic**, one that is slow in its course. **D.**, **Constitutional**, one in which a system of organs or the whole body is involved. **D.**, **Focal**, a localized disease. **D.**, **Functional**, a disease without

discoverable organic lesion. **D.**, Idiopathic, one that exists by itself without any connection with another disorder; one of which the cause is unknown. **D.**, Intercurrent, a disease occurring during the progress of another of which it is independent. **D.**, Organic, one due to structural changes. **D.**, Parasitic, one due to an animal or vegetal parasite. **D.**, Septic, one arising from the development of pyogenic or putrefactive organisms within the body. **D.**, Specific, one caused by the introduction of a specific virus or poison within the body; also a synonym of syphilis. **D.**, Venereal, one due to sexual intercourse. **D.**, Zymotic, a disease arising from the introduction and multiplication of some living germ within the body.

Disengagement (*dis-en-gāj'-ment*) [*Fr., des-engager, to disengage*]. Emergence from a confined state; especially the escape of the head of the fetus from the vaginal canal.

Disinfectant (*dis-in-fek'-tant*) [*dis, neg.; inficere, to corrupt*]. An agent that destroys the germs of disease, fermentation, and putrefaction.

Disinfection (*dis-in-fek'-shun*) [*dis, neg.; inficere, to corrupt*]. The destroying of disease-germs, especially by means of chemic substances.

Disintegrate (*dis-in'-te-grāt*) [*dis, apart; integer, the whole*]. To break up or decompose.

Dislocation (*dis-lo-ka'-shun*) [*dis, apart; locare, to place*]. The displacement of one or more bones of a joint, or of any organ from its natural position. **D.**, Complete, one in which the joint surfaces are entirely separated. **D.**, Compound, one in which the joint communicates with the external air through a wound. **D.**, Consecutive, one in which the displaced bone is not in the same position as when originally misplaced. **D.**, Old, one in which inflammatory changes have occurred. **D.**, Partial or Incomplete, one in which the articulating surfaces remain in partial contact; also called subluxation. **D.**, Pathologic, one the result of disease in the joint or of paralysis of the controlling muscles. **D.**, Primitive, one in which the bones remain as originally displaced. **D.**, Recent, one in which no inflammatory changes have ensued. **D.**, Simple, one in which there is no communication with the air through a wound. **D.**, Traumatic, that due to injury.

Disodic (*di-so'-dik*) [*di, two; sodium*]. Containing two atoms of sodium in the molecule.

Disparate (*dis'-par-at*) [*dispar, unequal*]. Not alike; unequal or unmated. **D.** Points, nonidentical points of the two retinæ. Diplopia is produced when the images of a single object fall upon such points.

Dispareunia (*dis-par-oo'-ne-ah*). See *Dyspareunia*.

Dispensary (*dis-pens'-ar-e*) [*dispensare, to distribute*]. A charitable institution where medical treatment is given to the poor.

Dispensatory (*dis-pens'-at-or-e*) [*dispensatorium, an apothecary's diary*]. A treatise on materia medica, and the composition, effects, and preparation of medicines.

Dispersion (*dis-per'-shun*) [*dispersio; dispersus, scattered*]. The act of scattering. In physics, the separation of a ray of light into its component parts by reflection or refraction; also, any scattering of light, as that which has passed through ground glass.

Dispirem (*dis-pi'-rem*) [*dis, two; spira, a spiral*]. The two skeins of a dividing nucleus formed from the nuclear loops and in development giving rise to the daughter-nuclei.

Displacement (*dis-plas'-ment*) [*Fr., displacer, to displace*]. A putting out of the normal position.

Dissect (*dis-ekt'*) [*dissecare to cut up*]. To cut tissues apart carefully and slowly, to allow study of the relations of a part.

Dissection (*dis-ekt'-shun*). The cutting apart of the tissues of the body for purposes of study. **D.-wound**, a septic wound acquired during dissection.

Disseminated (*dis-em'-in-a-ted*) [*disseminare, to scatter seed*]. Scattered; spread over a large area. **D.** Sclerosis, a disease of the central nervous system, in which the areas of sclerosis are irregularly scattered throughout the cord and brain. Synonyms—multiple or insular sclerosis.

Dissociation (*dis-so-se-a'-shun*) [*dis, apart; sociare, to associate*]. Separation, especially the separation of a complex compound into simpler molecules by the action of heat. **D.-symptom**, anesthesia to pain and to heat and cold, with preservation of tactile sensibility and of the muscular sense; it is observed in syringomyelia.

Dissolution (*dis-o-lu'-shun*) [*dissolutio; dissolvere, to set free*]. 1. The separation of a body or compound into its parts. 2. Death; decomposition.

Dissonance (*dis'-o-nans*) [*dissononare, to disagree in sound*]. The combination of such tones as are so different from each other as to produce beats.

Distal (*dis'-tal*) [*distare, to be at a distance*]. Extreme; at the greatest distance from a central point; peripheral.

Distichiasis (*dis-tik-i'-as-is*) [*δῖς, double; στίχος, a row*]. The condition in which there is a double row of eyelashes, the inner rubbing against the globe. See also *Trichiasis* and *Entropion*.

Distillate (*dis'-til-at*). The product obtained by distillation.

TABLE OF EPONYMIC DISEASES.

NAME.	PART AFFECTED.	SYMPTOMS.	PATHOLOGY.	SYNONYMS.
Addison's disease.	Suprarenal capsules, at times also abdominal sympathetic nerves and ganglia.	Brown pigmentation of the skin, progressive anemia, gastric disturbances, death from exhaustion.	Tuberculosis of the capsules.	Melasma suprarenale; dermatomelasma suprarenale; cutis aerea; bronzed-skin disease; copper-skin.
Addison's keloid.	Skin of breast, face, and neck.	Contractions of skin and fascia; scar-like growths, of rounded or irregular shape; pale-yellowish or brownish; fixation of joints from skin-induration.	Formation of scar-tissue.	Morphea; circumscribed scleroderma; scleroderma localis; true keloid.
Alibert's keloid.	Skin.	Fibrous outgrowths of the skin.	Formation of connective tissue.	Cheloid; kelis; fungoid mycosis.
Aran-Duchenne's disease.	Spinal cord.	Progressive muscular atrophy, spastic condition, fibrillation.	Degeneration of motor nuclei and pyramidal tracts of the cord; fatty and sclerotic changes in the muscles.	Cruveilhier's palsy; progressive muscular atrophy; Cruveilhier's atrophy; Duchenne's disease; amyotrophic lateral sclerosis.
Baelz's disease.	Labial mucous membrane.	Begins as small, painless papule which ulcerates; no glandular involvement.		
Balser's fatty necrosis.	Pancreas, omentum, mesentery.	Indefinite; those of acute pancreatitis.	Opaque, white, fatty areas in the organs.	
Barlow's disease.	Periosteum of bones.	Swelling and tenderness of extremities, chiefly lower; epiphyseal fractures; anemia.	Subperiosteal hemorrhages due to scurvy; rickets a predisposing cause.	Acute rachitis; rachitic scorbutus; periostitis hæmorrhagica; osteopathia hæmorrhagica infantum.
Barton's fracture.	Radius (lower extremity).	Silver-fork deformity.	A chipping off of the articular surface.	
Basedow's disease.	See <i>Graves' Disease</i> .			
Bazin's disease.	Mucous membrane of cheek.	Psoriatic eruption on the buccal mucous membrane.	That of psoriasis.	Buccal psoriasis.
Béclard's hernia.	Intestine.	Those of hernia.	Femoral hernia, through the saphenous opening.	
Bednar's aphthæ.	Mucous membrane of hard palate of the new-born.	Progressive marasmus and death.	Specific ulcerative stomatitis of the hard palate.	Ulcerative stomatitis of the new-born; apthæ of the palate.
Begbie's disease.	See <i>Graves' Disease</i> and	<i>Bergeron's Disease</i> .		
Belgel's disease.	See <i>Bergeron's Disease</i> .			
Bell's disease or mania.	Brain.	Acute delirium, wild hallucinations, insomnia, slight fever, dry tongue, rapid running pulse.	No lesions found.	Acute delirium; acute periencephalitis.

Bell's palsy.	Facial nerve.	Paralysis of entire half of face ; occasional neuralgia.	Lesion of the facial nerve ; a neuritis.	Peripheral facial paralysis.
Bérard's aneurysm.	Artery.	Varicose aneurysm with the sac in the tissues immediately around the veins.	That of aneurysm.	
Bergeron's disease.	Any group of muscles, as of abdomen, head, etc.	Localized rhythmic movements.	None. Hysteria.	Localized rhythmic chorea ; hysteretic chorea ; Begbie's disease.
Blainville's ear.	Ear.	Asymmetry of the ears.	Deformity of the ears ; congenital.	
Bostock's catarrh.	A synonym of <i>Hay-fever</i> .	<i>tis</i> .		
Bouillaud's disease.	A synonym of <i>Endocarditis</i> .	Painless and gradual enlargement of the bursa.		
Boyer's cyst.	Subhyoid bursa.	Edema, anemia, albuminuria, tubercasts.	Inflammation of the renal tissues.	Nephritis.
Bright's disease. See <i>Bright's Disease</i> .	Kidney.	Ataxia ; local anesthesia ; hysterical manifestations.	None.	Hysterical ataxia.
Briquet's ataxia.	Nervous system.	Loss of speech from incoordination of the muscles ; agaphia ; right-sided hemiplegia.	Lesion of third left frontal convolution of the brain.	Motor aphasia ; ataxic aphasia.
Broca's aphasia.	Brain.	Those of bone-abscess.	Ulceration of bone-tissue ; tuberculosis ; traumatism.	
Brodie's abscess.	Head of the tibia.	Those of synovitis.	Pulpy degeneration of the synovial membrane of a joint, especially of the knee.	
Brodie's disease or knee.	Knee and other joints.	Paralysis of a limb or side of the body, corresponding to the lesion in the cord ; anesthesia on the opposite side ; increased reflexes on side of lesion.	Lesion of one lateral half of the cord.	Brown-Séquard's paralysis.
Brown-Séquard's disease.	Spinal cord.	<i>low Atrophy of the liver. See Icterus</i>	<i>gravis</i> .	Icterus typhoides (of Lebert).
Budd's jaundice.	A synonym of <i>Acute Yellow</i>	Icterus, cyanosis, diarrhea, vomiting.	Acute fatty degeneration of all the organs ; septic infection.	Icterus of the new-born.
Buhl's disease.	All the organs of the new-born.	Those of pulmonary tuberculosis.	Racemose distribution of tuberculous infiltration about smaller bronchial tubes.	
Carswell's grapes.	Lungs.	Characteristic eruption on skin. See <i>Lupus</i> .	Inflammation of corium with cell-infiltration.	Lupus erythematodes ; lupus erythematosus.
Cazenave's lupus.	Skin of face and head ; occasionally of extremities, rarely on trunk.			

TABLE OF EPONYMIC DISEASES.—Continued.

NAME.	PART AFFECTED.	SYMPTOMS.	PATHOLOGY.	SYNONYMS.
Chabert's disease.	A synonym of <i>Symptomatic Anthrax</i> or <i>Black-leg</i> .			
Charcot's disease (<i>sclérose en plaques</i>).	Brain and spinal cord.	Spastic paraplegia with great increase in the reflexes; volitional tremor; scanning speech; nystagmus; mental debility; vertigo; attacks of coma.	Areas of grayish-red degeneration in the white matter of the cord and brain; excess of fibrous tissue.	Disseminated sclerosis; cerebro-spinal sclerosis; insular sclerosis; multilocular sclerosis; multiple sclerosis.
Charcot's disease.	Pyramidal tracts of medulla and spinal cord.	Progressive emaciation, and wasting of muscles; spastic condition of muscles; increased reflexes; tremor.	Atrophy of the anterior horns and sclerosis of the anterolateral columns.	Amyotrophic lateral sclerosis; primary lateral sclerosis; primary spastic paraplegia; lateral sclerosis; chronic anterior poliomyelitis; progressive muscular atrophy.
Charcot's joint.	Any joint.	Swelling; but slight local heat and pain; soreness on motion.	Chronic inflammation of the synovial membrane, later involving the bony substance. Occurs in the course of certain spinal diseases.	Spinal arthropathy.
Cloquet's hernia.	Intestines.	Those of hernia.	Femoral hernia resting on pectineus muscle, the tumor having turned within and behind femoral vessels.	Pectineal crural hernia.
Colles' fracture.	Radius (lower third).	Silver-fork deformity.	A splitting from $\frac{1}{8}$ inch to $1\frac{1}{2}$ inches above the articulation.	
Concato's disease.	Serous membranes, especially the pleura.	Vary with the membranes involved.	Progressive and malignant inflammation of various serous membranes.	Polyserositis; polyserositis.
Cooper's hernia.	Intestines.	Those of hernia.	Lodgment of the small intestine in the duodenojejunal fossa.	Mesenteric hernia; retroperitoneal hernia.
Cooper's irritable breast.	Synonym of <i>Mastodynia</i> .			
Cooper's irritable testicle.	Synonym of <i>Orchialgia</i> .			
Corrigan's disease.	A Synonym of <i>Aortic Insufficiency</i> .			
Craviellier's atrophy or palsy.	See <i>Avan-Duchenne's Disease</i> .			
Craviellier's disease.	Stomach.	Pain; hematemesis; vomiting; emaciation.	Ulceration of the gastric wall.	Simple ulcer of the stomach.

Curling's ulcer.	Duodenum.		Ulceration following burns of the sur- face.	Duodenal ulcer.
DeCosta's disease.	Gastrointestinal tract; heart, brain.	Vary with the part affected, and are preceded by a rapid disappearance of the local signs.	Vary with the localization of the dis- ease; gouty diathesis.	Suppressed goit; retrocedent gout; lithemia.
Darier's disease.	Skin of body.	Characteristic lesions of the skin.	Inflammation of the skin-layers, with cell-infiltration; probably parasitic.	Keratosis follicularis; ichthyosis sebacea cornea.
Darwin's ear.	Ear.	Absence of the upper part of the helix, and formation of cartilaginous nod- ules in the helix.	Malformation; congenital.	
Devergie's disease.	See <i>Gibert's pityriasis</i> .			
Donder's glaucoma.	Eye.	Those of glaucoma.	Atrophy of the optic papilla and hardening of the eyeball.	Simple atrophic glaucoma.
Dresler's disease.	Blood.	Recurring periodic attacks of hema- turia.	Presence of coloring-matter of the blood in the urine; malarial origin.	Paroxysmal hemoglobinuria; in- ferment hemoglobinuria; peri- odic hemoglobinuria; Harley's disease.
Dubini's disease.	Nervous system.	Progressive palsy and wasting, with sudden, shock-like movements.	Obscure.	Electric chorea.
Dubois' abscess.	Thymus gland.	Signs and symptoms of suppuration.	Abscess of gland in congenital syphi- lis.	
Duchenne's disease.	See <i>Aran-Duchenne's Dis-</i>	<i>ease</i> .		
Duchenne's disease.	A synonym of <i>Bulbar Par-</i>	<i>alysis</i> , and of <i>Locomotor Ataxia</i> .		
Duchenne's paralysis.	A synonym of <i>Pseudohy-</i>	<i>perrophic paralysis</i> .		
Duhring's disease, or impetigo.	Skin.	Protean eruption; itching and burn- ing; exhaustion.	Undetermined.	Dermatitis herpetiformis; pemphi- gus pruriginosus; hydroa herpeti- forme; herpes gestationis; herpes circinatus; bullous; pemphigus circinatus; pemphigus malignus; pemphigus hystericus.
Dupuytren's contrac- tion.	Palmar fascia and con- tiguous structures.	Contraction of one or more fingers.	Cicatrical contraction from injuries.	
Dupuytren's hydrocele.	Tunica vaginalis testis.	Those of hydrocele.	Same as hydrocele.	Bilocular hydrocele.
Eichhorst's neuritis.	Nerves.	Those of neuritis.	Obscure.	Neuritis fasciens.
Eichstedt's disease.	Skin.	Characteristic lesion. See <i>Tinea</i> <i>versicolor</i> .	Infiltration of cuticle with the para- site; microsporon furfur.	Pityriasis versicolor; chromophy- tosis; tinea versicolor.

TABLE OF EPONYMIC DISEASES.—Continued.

NAME.	PART AFFECTED.	SYMPTOMS.	PATHOLOGY.	SYNONYMS.
Erb's disease.	Spinal cord.	Progressive wasting of various groups of muscles in upper arm and shoulder, and gluteal and thigh muscles; later, hypertrophy.	Fatty and sclerotic degeneration of affected muscles; as a rule no degeneration of motor nuclei in cord.	Juvenile muscular atrophy; primary muscular dystrophy; idiopathic muscular atrophy.
Erb's palsy.	Pyramidal tracts of cord.	Loss of power; slight spasm of muscles of lower extremities; spastic gait; increased reflexes; bladder disturbance.	Degeneration of pyramidal tracts; meningeal changes; syphilis.	
Erichsen's disease.	Cerebrospinal axis.	Those of neurasthenia and hysteria.	Concussion of central nervous system from railroad-shock.	Railway-spine; railway-brain; traumatic hysteria.
Fauchard's disease.	Dental alveoli.	Pain; suppuration; swelling.	Slow and progressive destruction of alveoli; death of teeth.	Alveolodental periostitis; pyorrhea alveolaris; alveolar abscess; Rigg's disease.
Flajan's disease.	Synonym of <i>Exophthalmic goiter</i> .			
Fothergill's disease, or faceache.	Facial nerve.	Severe pain in face; tender points of Valleix; salivation; herpes; larymation.	Obscure.	Facial neuralgia; tic douloureux; trifacial neuralgia; prosopalgia; trigeminal neuralgia.
Fothergill's sore-throat.	Throat.	Pain; swelling; associated symptoms of scarlet fever.	Intense inflammation; membranous exudate.	Scarlatina anginosa; anginoid scarlet fever.
Friedreich's ataxia or disease.	Spinal cord.	Peculiar swaying, irregular gait; similar movements of head and arms; reflexes frequently lost; nystagmus; talipes equinus; lateral curvature of spine; scanning speech.	Sclerosis of posterior and lateral columns of cord; heredity.	Hereditary ataxia; family-ataxia; posterolateral sclerosis; hereditary ataxic paraplegia.
Friedreich's disease.	Synonym of <i>Paramyoclonus</i> .			
Gerlier's disease.	Nervous system.	Sudden paroxysms of ptoxis; rotary vertigo; muscular paresis; cervico-occipital pain.	Obscure.	Paralyzing vertigo.
Gibson's hydrocele.	Scrotum and bowel.	Those of hydrocele and hernia.	Hydrocele with hernia.	
Gilbert's pityriasis.	Skin.	Characteristic eruption; chronic course; itching; progressive emaciation; death.	A dermatitis.	Pityriasis rubra; dermatitis exfoliativa; pityriasis rubra aiguë; Devergie's disease; Hebra's pityriasis.
Giffard's disease.	Abdominal viscera.	Those of gastrointestinal disturbance with neurasthenia; prolapse of abdominal organs.	Looseness of mesenteric and peritoneal attachments; occurs in neurasthenia.	Enterptosis; splanchnoptosis.

Glissonian cirrhois.	Liver.	Those of cirrhois.	Liver much reduced in size from cirrhotic changes.	Inguinal intestinal hernia.
Gourand's (or Gourand's) disease.	The bowel.	Those of inguinal hernia.	That of hernia.	Exophthalmic goiter; Basedow's, Parry's, Basedow's, March's, Flajana, Parson's, Stokes's disease.
Graves's disease.	Thyroid gland, eyes, heart.	Palpitation; goiter; exophthalmos; Graefe's lid-sign; Stellwag's sign.	Increased vascularity and hypertrophy of thyroid gland.	Hydrocephaloid disease; spurious hydrocephalus; hydrencephaloid disease.
Gubler's tumor.	Wrist.	Prominence on back of wrist in wrist-drop.		Endemic syphilis.
Hall's (Marshall) disease.	Brain.	Occasional convulsions; semicomatose condition; irregular pulse; contracted pupils; flaccid, depressed fontanels; retracted head; Cheyne-Stokes' breathing.		Athetosis; posthemiplegic chorea.
Halstern's disease.	Entire body.	Those of syphilis.	That of syphilis.	Hypertrophic cirrhois of liver; biliary cirrhois.
Hammond's disease.	Brain and cord.	Constant rhythmic movements of fingers and toes.	Obscure; lesion of the brain and cord.	
Hanot's disease.	Liver.	Chronic jaundice; absence of ascites; enlargement of liver and spleen.	Increase of intralobular and interlobular connective tissue; liver large, smooth, firm.	
Harley's disease.	See Dressler's disease.			
Heberden's asthma.	Synonym of <i>Angina pectoris</i> .			
Heberden's disease or rheumatism.	Smaller articulations, as those of the fingers.	Those of chronic rheumatism, or rheumatoid arthritis with formation of nodosities.	Destruction of the cartilages; roughening of articular surfaces; thickening of the ligaments.	Osteoarthritis; rheumatic arthritis; digitum nodi.
Hebra's erythema.	Skin.	Characteristic eruption; slight constitutional manifestations.	Inflammatory effusion of serum and leukocytes.	Polymorphous erythema; erythema multiforme.
Hebra's pityriasis.	See <i>Gibert's pityriasis</i> .			Prurigo ferox; congenital prurigo.
Hebra's prurigo.	Skin.	Characteristic eruption; limbs feel like coarse brown paper; incurable; persists through life.	Unknown—probably a neurosis.	
Henoch's purpura.	Skin.	Purpuric eruption, associated with peculiar intestinal symptoms.		
Hesselbach's hernia.	Intestines.	Those of hernia. The hernia presents a lobulated appearance.	There is a diverticulum of the hernia through the cribriform fascia.	
Heubner's disease.	Cerebral arteries.	Cerebral disturbances, very variable.	Endarteritis from syphilis.	

TABLE OF EPONYMIC DISEASES.—Continued.

NAME.	PART AFFECTED.	SYMPTOMS.	PATHOLOGY.	SYNONYMS.
Hodgkin's disease.	Lymph-glands.	Enlargement of the lymphatic glands; progressive anemia.	Hyperplasia of lymphoid structures of the body; enlargement of spleen.	Pseudoleukemia; lymphadenoma; lymphadenosis; malignant lymphoma; aneemia lymphatica; adenoid disease; adenia; chronic recurrent fever (Ebstein).
Hodgson's aneurysm, or disease.	Aorta, first part.	Those of aneurysm, frequently with hypertrophy and dilatation of the heart.	Inflammatory changes.	Hodgson's dilatation.
Huguier's disease.	Uterus.	Those of fibroid tumor of the uterus.	Development of fibromyoma.	Fibromyoma of the uterus.
Huntingdon's chorea.	Brain.	Irregular movements; disturbance of speech; gradual dementia.	No characteristic lesions; heredity.	Hereditary chorea; chronic chorea.
Inman's myalgia.	Muscular system.	Muscular pain of any origin.	None.	Muscular rheumatism.
Jacksonian epilepsy.	Brain-cortex.	Spasm of a single group of muscles in the face, arm, or leg, or other region; no loss of consciousness; aura.	Lesion of the motor area of the brain.	Cortical epilepsy; symptomatic epilepsy; focal epilepsy; partial epilepsy.
Jacobi's ulcer.	Eyelid.	Slow ulceration at inner end of lower lid.	Cancer-formation; no lymphatic involvement.	Rodent ulcer of the eyelid; lupus of the eyelid; carcinoma of the eyelid.
Jaksch's disease.	See <i>Pseudoleukemia</i> , <i>Infantile</i>			Infantile pseudoleukemic anemia.
Kaposi's disease.	Skin.	Characteristic eruption, followed by atrophy of the skin and formation of carcinomata; emaciation; exhaustion; death.	Atrophic degeneration of the skin.	Xeroderma pigmentosum; parch-ment-skin; atrophoderma pigmen- tosum; angioma pigmentosum atrophicum.
Kopp's asthma.	See <i>Millar's Asthma</i> .			
Kriehaber's disease.	Nervous system.	Hyperesthesia; illusions of sense; vertigo; tachycardia.	Neurosis.	Cerebrocardiac neuropathy.
Krönlein's hernia.	Intestines.	Those of hernia.	A variety of inguinal hernia.	Inguinoproperitoneal hernia.
Kussmaul's coma.	A synonym of <i>Diabetic</i>			
Kussmaul's disease.	Spinal cord.	Sudden onset; rapid muscular wast- ing; reaction of degeneration; fever.	Acute hemorrhagic myelitis of the anterior horns, with degeneration and destruction of the large gar- gilon-cells.	Acute atrophic spinal paralysis; polio-myelitis anterior.
Lænnec's cirrhosis.	Liver.	Digestive derangement; diminution in size of liver; tendency to hemor- rhage; ascites.	Overgrowth of connective tissue be- tween acini.	Chronic interstitial hepatitis; atro- phic cirrhosis; portal cirrhosis.

L a n d o u z y - D é j é r i n e type of muscular atrophy.	Spinal cord.	Wasting of muscles of face and those of the scapulohumeral group.	Degeneration of the muscles involved; no degeneration of motor nuclei in spinal cord or of peripheral nerves.	Facioscapulohumeral type of idiopathic muscular atrophy.
Landry's paralysis.	Spinal cord.	Loss of motor power in the lower extremities gradually extending upward; exhaustion; loss of reflexes; sensation usually intact; generally fatal.	Obscure; toxic.	Landry's disease; acute ascending paralysis.
Laugier's hernia.	Intestines.	Those of hernia.	A femoral hernia perforating Gimbernat's ligament.	
Leber's disease.	Eye.	Those of optic neuritis.	Malformation of optic nerve with atrophy, and inflammatory changes.	Hereditary optic atrophy.
Legal's disease.	Meningotympanic region.	Headache.	Inflammatory changes.	
Leyden's neuritis.	Nerves.	Those of neuritis.	Destruction of the nerve-fibers; formation of fibrous connective tissue, with deposit of fat.	Lipomatous neuritis.
Leyden's type of muscular atrophy.	Spinal cord.	Wasting of muscles of lower extremities and back; hypertrophy of the calves.	Degeneration of the affected muscles; no degeneration of the motor nuclei of the spinal cord; difficult labor or premature delivery.	Hereditary muscular atrophy.
Little's disease.	Brain and cord.	Stiffness of the legs; talipes equinus with characteristic attitude; crossed-legged progression; lead-pipe contraction; increased reflexes; sensation normal; symptoms of cerebral defects—idiocy, imbecility, nystagmus.	Lesion of brain-cortex; meningeal hemorrhage; arrest of development of pyramidal tracts.	Congenital muscular rigidity; spastic tabes of children; spastic paraplegia of infants; paraplegia cerebrii spastica.
Littre's hernia.	Intestines.	Those of hernia.	Hernia of an intestinal diverticulum.	
Littre's hernia.	Synonym of <i>Richter's</i> .	<i>Hernia.</i>		
Lobstein's cancer.	Abdomen.	Large abdominal tumor, centrally situated; immovable.	Sarcomatous tissue.	Retropertitoneal sarcoma.
Louis's angina.	Synonym of <i>Ludwig's</i> .	<i>Angina.</i>		
Ludwig's angina.	Neck.	Swelling, especially in the submaxillary region; intense pain.	Suppurative inflammation of neck.	Louis's angina; cellulitis of the neck.
Malassez' disease.	Testicle.	Tumor of the gland.	Cyst-formation.	Cyst of the testicle.
March's disease.	See <i>Graves' Disease</i> .			
Marie's disease. See <i>Acromegaly</i>.	Bones of face and extremities.	Gradual hypertrophy of the hands, feet, and face; changes in the size of the thyroid gland; headache.	Hypertrophy of the bones; at times enlargement of the pituitary body.	Acromegaly; acromegalia; akromegaly.

TABLE OF EPONYMIC DISEASES.—Continued.

NAME.	PART AFFECTED.	SYMPTOMS.	PATHOLOGY.	SYNONYMS.
Marjolin's warty ulcer.	Skin.	Wart-like parallel growths; progress slow; glandular enlargement.	Malignant change of a cicatrix or old ulcer.	
McLeod's capsular rheumatism.	Joints.	Those of rheumatism.	Effusion into the synovial capsules and into the synovial sheaths and bursæ around the affected joints.	
Ménière's disease.	Semicircular canals.	Vertigo; nausea; vomiting; deafness; rotatory movements; temporary apopleptic symptoms; tinnitus aurium; ocular symptoms—nystagmus, diplopia.	Inflammatory processes in the canals, or perhaps a central lesion.	Aural vertigo; auditory vertigo; labyrinthine vertigo.
Millar's asthma.	Nervous system; reflexly the larynx.	Sudden paroxysms of suffocation, with short, sonorous respiration; high mortality in children.	Neurosis; rachitis.	Thymic asthma; spasm of the glottis; laryngismus stridulus; laryngospasm; laryngic suffocation; spasmodic croup; internal convulsions; spasmodic laryngitis; "passion-fits;" childhood; Kopp's asthma.
Möller-Barlow's disease.	Extremities.	<i>ease.</i>		
Morand's disease.	Foot.	Paresis of the extremities.	Disease of the nervous system.	
Morel's ear.	Ear.	The presence of eight toes on the foot.	Deformity of foot.	Flat ear.
Morton's foot or Morton's toe.	Metatarsophalangeal joint of the fourth toe.	Flattening of the folds of the ear; thinness of the rim; increase in the size of the ear.	Congenital abnormality of the helix, antihelix, scaphoid fossa, and crura furcata.	
Morvan's disease.	A limb.	Pain in the foot.	Local neuritis.	Metatarsalgia.
Nægele's pelvis.	Pelvis.	Loss of sensibility on one side; neuralgic pain and paresthesia of the affected limb; vasomotor and trophic disturbances, as shown by painless and destructive whitlows.	Disease of the posterior roots of the nerves of the affected limb. By some considered a form of leprosy or of syringomyelia.	Monoplegia anæsthetica.
Paget's disease.	Bone.	Deformity of pelvis.	Lack of development of one lateral mass of the sacrum.	Obliquely-contracted pelvis.
Paget's disease.	Nipple and areola.	Hypertrophy of the bone-tissue, with deformity.	Inflammatory changes.	Hypertrophic deforming osteitis; osteitis deformans.
		Eczema-like eruption; frequently terminates in carcinoma.	Inflammatory changes; proliferation of epithelial cells.	Paget's nipple.

Paget's recurrent fibroid.	Subcutaneous tissue; occasionally the breast.	Recurrent growth; painless; lymphatic glands not involved.	Spindle-cell sarcoma.	Spindle-cell sarcoma.
Park's aneurysm.	Arteries and veins.	Those of aneurysm.	Arteriovenous aneurysm; the artery communicates with two contiguous veins.	
Parkinson's disease. See <i>Paralysis agitans</i> .	Nervous system.	Muscular weakness; tremors; rigidity; festination; characteristic attitude.	No constant lesion; probably changes in the cerebral cortex.	Paralysis agitans; shaking-palsy.
Parrot's disease.	Nervous system.			Syphilitic pseudoparalysis.
Parry's disease.	See <i>Graves's Disease</i> .			
Parson's disease.	See <i>Graves's Disease</i> .			
Partridge's hernia.	Intestines.	Those of hernia.	Femoral hernia external to the femoral vessels.	
Pavy's disease.	Kidney.	Occasional appearance of a small amount of albumin in the urine.	Renal congestion.	Intermittent, paroxysmal, cyclic, physiologic, simple, functional, transparent albuminuria; albuminuria of adolescence.
Paxton's disease.	Hair of beard.	Nodular growth on the hairs; brittleness of the hairs.	Thickening and roughening of the hair-shaft.	Tinea nodosa.
Petit's hernia.	Intestines.	Those of hernia.	Hernia in the lumbar region.	
Physick's encysted rectum.	Rectum.	Burning pain in anus and adjacent parts; increased mucous secretion.	Hypertrophy and occasionally inflammation of the rectal sacs.	
Pott's aneurysm.	Arteries.	Pulsating tumor under the skin.	That of aneurysm.	Aneurysm by anastomosis; aneurysmal varix.
Pott's disease.	Spinal column.	Curvature of the spine in an antero-posterior direction; pain.	Inflammation and caries of the vertebral bodies and intervertebral discs.	Caries of the vertebrae; Pott's curvature.
Pott's fracture.	Fibula.	Those of fracture; dislocation outward of foot.	Fracture of lower end of bone; sometimes fracture of inner malleolus.	
Pott's gangrene.	Extremities.	Extreme pain; drying and discoloration of affected part.	Obstruction of blood-supply from atheroma, etc.; old age.	Senile gangrene.
Pott's paralysis or paraplegia.	Lower portion of body.	Inability to move the limbs; disturbances of sensation.	Compression of spinal cord.	
Quain's fatty heart.	Heart.	Those of fatty heart.	Fatty degeneration of heart-muscle.	
Quincke's disease.	Skin.	Sudden circumscribed swelling of the skin; a sense of tension.	Angioneurosis.	Urticaria oedematosa; giant urticaria; angioneurotic edema.

TABLE OF EPONYMIC DISEASES.—Continued.

NAME.	PART AFFECTED.	SYMPTOMS.	PATHOLOGY.	SYNONYMS.
Quinquaud's disease.	Scalp and hairy parts.	Formation of smooth, irregular patches of alopecia, dotted with red points.	Perifollicular inflammation.	Folliculitis decalvans; acne decalvans.
Ray's mania.	Brain.	A form of general moral insanity.	Changes in the cerebral cortex.	
Raynaud's disease. See <i>Raynaud's Disease</i> .	The extremities.	Local syncope; local asphyxia; local gangrene; hemoglobinuria.	Obscure; vasomotor changes.	Symmetric gangrene; sphaceloderma; dermatitis gangrenosa.
Raynaud's disease.	Throat-muscles.	Paralysis of affected muscles.	Secondary to parotitis.	
Reclus' disease.	Mammary gland.	Slight enlargement of both breasts; fluctuation; no pain.	Disseminated cystic dilatations of the ducts and acini.	Cystic disease of the breast.
Richter's hernia.	Intestines.	Those of hernia.	Enterocoele in which only a part of the circumference of the gut is constricted.	Parietal enterocoele; partial enterocoele.
Riga's disease.	Mouth.	Aphthous eruption beneath tongue; painless; grave constitutional symptoms; earthy pigmentation of skin; death.	Ulcer, gray in color, with irregular and poorly defined borders; enlargement and fatty degeneration of liver and spleen.	Cachectic aphthae.
Rigga' disease. See <i>Fauchard's Disease</i> .				
Ritter's disease.	Skin.	Characteristic eruption on skin; exhaustion.	Inflammation, with exfoliation.	Dermatitis exfoliativa infantum; dermatitis exfoliativa neonatorum.
Rivolta's disease.	Synonym of <i>Actinomyces</i> .			
Robert's pelvis.	Pelvis.	Deformity of pelvis.	Lack of development of both lateral masses of the sacrum.	Transversely-contracted pelvis; doubly-synostotic pelvis.
Rodrigues' aneurysm.	Arteries and veins.	Those of varicose aneurysm.	A varicose aneurysm with the sac immediately contiguous to the artery.	
Rokitansky's disease.	A synonym of <i>Acute Yel</i> .	<i>low Atrophy of the Liver</i> . See <i>Icterus</i> .	<i>gravis</i> .	
Rosbach's disease.	Stomach.	Cephalalgia; gastralgia; very acid vomiting, in paroxysms.	Neurosis.	Hyperchlorhydria; gastroxixia; gastrosynsial.
Sæmisch's ulcer.	Cornea.	Those of purulent keratitis.	An acute, serpiginous, or creeping ulceration; sharp margins; gray floor; sloughing of cornea.	Infectious ulcer of the cornea.
Saint Agatha's disease.	Synonym of <i>Mammitis</i> .			
Saint Aignan's disease.	Synonym of <i>Tinea favosa</i> , or <i>Favus</i> .			

Saint Anthony's dance.	Synonym of <i>Chorea</i> .			
Saint Anthony's fire.	Synonym of <i>Erysipelas</i> .			
Saint Anthony's fire.	Skin.	Gangrene of the affected parts ; pain.	Intense inflammation, with destruction of the affected parts ; chronic ergotism.	Epidemic gangrene.
Saint Apollonia's disease.	Teeth and jaws.	Toothache and pain in the jaws.	Neurosis.	Toothache.
Saint Avertin's disease.	Synonym of <i>Epilepsy</i> .			
Saint Avidus' disease.	Synonym of <i>Deafness</i> .			
Saint Blaise's disease.	Same as <i>Saint Blarius' Disease</i> .			
Saint Blasius' disease.	Synonym of <i>Quinsy</i> .			
Saint Clair's disease.	Synonym of <i>Ophthalmia</i> .	or <i>Sore Eyes</i> .		
Saint Dymphna's disease.	Synonym of <i>Insanity</i> .			
Saint Erasmus' disease.	Synonym of <i>Colic</i> .			
Saint Fiacre's disease.	Synonym of <i>Hemorrhoids</i> .	or of <i>Condylomata</i> .		
Saint Francis' fire.	Synonym of <i>Erysipelas</i> .			
Saint Gervasius' disease.	Synonym of <i>Rheumatism</i> .			
Saint Giles' disease.	Synonym of <i>Carcinoma</i> .	or <i>Leprosy</i> .		
Saint Guy's dance.	Synonym of <i>Chorea</i> .			
Saint Hubert's disease.	Synonym of <i>Hydrophobia</i> .			
Saint Job's disease.	Synonym of <i>Syphilis</i> .			
Saint John's dance.	Synonym of <i>Chorea</i> .			
Saint John's evil.	Synonym of <i>Epilepsy</i> .			
Saint Lazarus' disease.	Synonym of <i>Leprosy</i> .	<i>Timea</i> , and of <i>Measles of the Hog</i> .		
Saint Main's disease.	Synonym of <i>Scabies</i> .			
Saint Martin's evil.	Synonym of <i>Drunkennes</i> .			
Saint Mathurin's disease.	Synonym of <i>Insanity</i> , of	<i>Idiocy</i> , or of <i>Epilepsy</i> .		

TABLE OF EPONYMIC DISEASES.—Continued.

NAME.	PART AFFECTED.	SYMPTOMS.	PATHOLOGY.	SYNONYMS.
Saint Modestus' dance.	Synonym of <i>Chorea</i> .			
Saint Roch's disease.	Synonym of the <i>Bubonic Plague</i> .			
Saint Sement's disease.	Synonym of <i>Syphilis</i> .			
Saint Valentine's disease.	Synonym of <i>Epilepsy</i> .			
Saint Vitus' dance.	Synonym of <i>Chorea</i> .			
Saint With's dance.	Synonym of <i>Chorea</i> .			
Saint Zachary's disease.	Synonym of <i>Dumbness</i> .			
Schönlein's peliosis.	Joints and skin.	Multiple arthritis; purpuric eruption, with urticaria and erythema; sore throat; fever; edema; scanty and albuminous urine; relapses.	Vasomotor neurosis (?). Rheumatic poison.	Schönlein's disease; peliosis rheumatica; purpura rheumatica.
Stahl's ear.	Ear.	Helix flat and folded over, forming one mass with the anthelix at its bifurcation.	Deformity of ear.	
Stokes's disease.	See <i>Graves's Disease</i> .			
Storck's blennorrhea.	Respiratory tract.	Chronic mucous discharge.	Engorgement of the mucous membrane.	
Strumpell's disease.	Brain and spinal cord.	Those of inflammation of the cord and brain; ophthalmoplegia; progressive muscular atrophy.		Polioencephalitis; polioencephalomyelitis.
Sydenham's chorea.	Synonym of <i>Chorea</i> .			
Tay's choroiditis.	Eye.	Those of central choroiditis; chronic course; symmetric; numerous white, glistening dots on retina; contraction of field of vision; scotoma.	Changes resembling those in albuminuric retinitis; colloid degeneration, and calcareous formations in choroid.	Senile guttate choroiditis.
Thomsen's disease.	Muscles of arms and legs.	Tonic spasm or rigidity of the muscles; myotonic reaction; reflexes normal.	Hypertrophy of the muscles; increase in the number of the nuclei; congenital; hereditary.	Myotonia congenita.
Thornwaldt's disease.	Pharynx, Luschka's gland.	Hypersecretion of mucus; hawking; catarrh.	Nasolaryngeal stenosis; formation of a cyst-like cavity in the gland, containing pus or mucopus.	Thornwaldt's buritis.

Tourette's (Gilles de la) disease.	Nervous system.	Motor incoordination; echolalia; coprolalia; involuntary muscular movements; explosive utterances; echokinesis; fixed mental ideas.	Psychosis.	Tic convulsif.
Velpeau's hernia.	Intestines.	Those of femoral hernia.	Femoral hernia in front of the vessels.	
Volkman's deformity.	Foot.	Deformity.	Tibiotalar luxation.	
Votolini's disease.	Middle ear.	Acute inflammation; cerebral symptoms; permanent deafness.	Inflammation of the labyrinth.	Purulent labyrinthitis.
Wardrop's disease.	Matrix of nails.	Pain; swelling; redness; tenderness; exudation of a fetid, serous fluid; falling off of the affected nails; involvement of the bone.	Intense inflammation.	Malignant onychitis; onychia maligna.
Well's disease.	General disease.	Remitting fever; chill; muscular pains; jaundice; headache; course, 8-14 days; clay-colored stools; urine albuminous; occasionally hematuria; cerebral symptoms and coma.	Swelling of liver and spleen; frequently nephritis.	Acute infectious jaundice; acute febrile icterus.
Werthoff's disease.	Skin.	Headache; great debility; purpuric eruption; hemorrhages from mucous surfaces; slight fever; frequently fatal.	Obscure.	Purpura hemorrhagica; morbus maculosus Werthoffi; land-scurvy.
Wildermuth's ear.	Ear.	Prominence of the antihelix.	Deformity.	
Wilks' kidney.	The large white kidney of	<i>Nephritis, Chronic Parenchymatous.</i>		
Willard's lupus.	Skin.	Those of lupus.	A name formerly given to lupus of tuberculous, as opposed to that of nontuberculous, origin.	
Willis's disease.	A synonym of <i>Diabetes</i> .			
Wilson's (E.) disease.	Skin.	Those of <i>Pityriasis rubra</i> .	That of <i>Pityriasis rubra</i> .	Universal exfoliative dermatitis.
Winckel's disease.	All the organs of the newborn infant.	Acute hemoglobinuria; jaundice; cyanosis.	Fatty degeneration of the organs; infarcts of the kidney; microorganism, entering through umbilical cord.	
Woillez' disease.	Lungs.	Those of acute pulmonary congestion	Congestion of the lung-tissue.	

Distillation (*dis-til-a'-shun*) [*distillare*, to drop little by little]. The process of vaporizing and collecting the vapor by condensation. It is used mainly in purifying liquids by separating them from nonvolatile substances. **D.**, **Destructive**, the decomposition of a substance in a closed vessel in such a manner as to obtain liquid products. **D.**, **Fractional**, a method of separating substances from each other by distilling the mixture containing them at a gradually increased temperature, the different substances being vaporized and collected in the order of their volatility.

Distoma, or **Distomum** (*dis'-to-mah*, *dis'-to-mum*) [*dis*, double; *στόμα*, a mouth]. A genus of trematode worms which have an oral as well as a ventral sucker.

Distribution (*dis-trib-u'-shun*) [*distribuere*, to distribute]. The branching of a nerve or artery, and the arrangement of its branches within those parts that it supplies.

Disulphate (*di-sul'-fat*) [*dis*, two; *sulphur*]. A sulphate containing one atom of hydrogen that can be replaced by a base.

Disulphid (*di-sul'-fid*) [*dis*, two; *sulphur*]. A compound of an element or radicle with two atoms of sulphur.

Dita Bark (*di'-tah*) [*L.*]. The bark of *Alstonia scholaris*, native to the Philippine Islands. It is employed as a tonic and antiperiodic in intermittent fever. Dose of the tincture, fʒ j-ij (4.0-8.0); of the fluid extract, gr. ij-v (0.13-0.32); of the powder, gr. v (0.32). Unof.

Dithiosalicylate of Sodium (*di-theo-sa-lil'-sil-at*). See **Sodium**.

Dithymoldiiodid (*di-thi-mol-di-i'-o-did*). Aristol.

Ditrich's Plugs. Masses of granular matter, degenerated epithelium, and crystals of the fatty acids in the sputum of putrid bronchitis.

Diureid (*di-yu'-re-id*). A compound containing two molecules of urea.

Diuresis (*di-u-re'-sis*) [*diá*, through; *οὐρεῖν*, to make water]. Increase in the secretion of urine.

Diuretic (*di-u-ret'-ik*) [*diá*, through; *οὐρεῖν*, to make water]. 1. Increasing the flow of urine. 2. An agent that increases the secretion of urine. **D.**, **Alterative**, drugs eliminated by the kidney and used for their local action on the surfaces over which they pass. **D.**, **Hydragogue**, those that increase the flow of water from the kidneys. **D.**, **Refrigerant**, those that render the urine less irritating while not greatly increasing its flow.

Diuretin (*di-u'-re-tin*). $C_6H_7NaN_2O_7 \cdot C_6H_4 \cdot OHCOONa$. Theobromin sodiosalicylate. It has been found useful as a diuretic in pleuritic effusion and cardiac dropsy. Dose 90 grains (6.0) daily, in four doses.

Diurnule (*di-urn'-ul*) [*Fr.*]. A form of medicinal tablet or capsule that contains the maximum quantity of a toxic drug that may be administered in 24 hours.

Divergent (*di-ver'-jent*) [*divergere*, to diverge]. Moving in different directions from a common point. **D.** **Strabismus**. See **Strabismus**.

Divers' Paralysis. See **Caisson-disease**.

Diverticulum (*di-ver-tik'-u-lum*) [*divertere*, to turn]. A small pouch or sac springing from a main structure. **D.**, **Meckel's**, a congenital diverticulum of the ileum, the remains of the omphalomesenteric duct.

Divulsion (*di-vul'-shun*) [*divulsio*, a tearing apart]. A tearing asunder.

Dobell's Spray, or Solution. Liqueur sodii boratis compositus (N. F.); a solution of borax, sodium bicarbonate, and phenol in glycerol and water; it is used as a spray for nasal and throat troubles.

Dochmiasis (*dok-mi'-as-is*) [*δόχμος*, crumpled]. The diseased condition caused by the presence in the body of parasites belonging to the genus *Dochmius*. Cf. *Ankylostomiasis*.

Dochmius duodenalis (*dok'-me-us du-den'-al-is*). See *Ankylostomum duodenale*.

Dog's Bane. See **Apocynum**.

Dogwood (*dog'-wood*). See **Cornus**.

Dolichos. Same as *Mucuna*.

Dolichocephalic, or **Dolichocephalous** (*dol-ik-o-sef'-al'-ik*, or *-sef'-al-us*) [*δολιχός*, long; *κεφαλή*, head]. Long-headed; having a relatively long anteroposterior cephalic diameter.

Dolorosa (*do-lor-o'-sus*) [*L.*]. Full of pain.

Dondrus Ndugu [African]. Brother ulcer; a disease common on the east coast of Africa, due to some organism that infests stagnant water. It affects especially the leg, which becomes inflamed and swollen, and sloughs below the healthy tissue.

Donovan's Solution. See **Arsenic**.

Dorsad (*dor'-sad*) [*dorsum*, the back; *ad*, toward]. Toward the dorsal aspect.

Dorsal (*dor'-sal*) [*dorsum*, the back]. Pertaining to the back, or to the posterior part of an organ. **D.** **Artery**. See **Artery**. **D.** **Decubitus**, recumbency in the supine position. **D.** **Reflex**. See **Reflexes**, **Table of**.

Dorsalis pedis (*dor-sal'-lis pe'-dis*). See **Arteries**, **Table of**.

Dorsispinal (*dor-se-spi'-nal*) [*dorsum*, back; *spina*, spine]. Relating to the back and the spinal column.

Dorso- (*dor'-so-*) [*dorsum*, back]. A prefix used to signify pertaining to the back.

Dorsum (*dor'-sum*) [*L.*]. 1. The back. 2. Any part corresponding to the back; as the dorsum of the foot, hand, tongue, penis, etc.

Dosage (*do'-saj*) [*δοσις*, a portion given]. The determination of the proper amount of

a medicine or other agent for a given case or condition.

Dose (*dōs*) [*dōsis*, a portion]. The measured portion of medicine to be taken at one time.

D., **Divided**, a relatively small quantity of a drug taken at short intervals. **D.**, **Lethal**, a dose sufficient to kill. **D.**, **Maximum**, the largest dose consistent with safety. **D.**, **Minimum**, the smallest quantity of a medicine that will produce physiologic effects.

Dosimetric (*do-sim-et-rik*) [*dōsis*, a dose; *μέτρον*, a measure]. Relating to or characterized by dosimetry.

Dosimetry (*do-sim-et-re*) [*dōsis*, dose; *μέτρον*, a measure]. The accurate and systematic measurement of a dose.

Douche (*doosh*) [Fr.]. A stream of water directed against a part, or one used to flush a cavity of the body. **D.**, **Air**, a current of air directed against some organ for therapeutic purposes.

Douglas's Cul-de-sac or Pouch. See *Pouch*. **D.'s Semilunar Fold**, a thin curved margin that forms the lower part of the posterior wall of the sheath of the abdominal rectus muscle.

Dover's Powder. See *Opium*.

Doyère, Eminence, Hillock, Papilla, or Tuft of (*doy-yār*). The slight elevation in a muscular fiber corresponding to the entrance of a nerve-fiber.

Drachm (*dram*). See *Dram*.

Dracontium (*dra-kon'-she-um*) [*δράκων*, a dragon]. Skunk cabbage.

Dracunculus (*dra-kun'-ku-lus*) [dim. of *δράκων*, a dragon, serpent]. A genus of thread-worms, belonging to the family *Filariidae*.

Draft, Draught (*draft*) [AS., *dragon*, to draw]. A quantity of liquid drunk at one gulp. **D.**, **Black**, compound infusion of senna. **D.**, **Effervescing**, one containing sodium or potassium bicarbonate and a vegetable acid.

Dragon's Blood (*drag'-onz blud*). 1. The astringent resin of *Calamus rotang* and *C. draco*, E. Indian rattan-palms. 2. The resin of various species of *Dracena*. 3. The resin of *Pterocarpus draco*, a W. Indian tree. The various kinds of dragon's blood are astringent, but are no longer used internally.

Drain (*drän*) [ME., *drainen*, to drain]. A material that affords a channel of exit for the discharge from a wound or cavity.

Drainage (*drän'-äj*) [ME., *drainen*, to drain]. The method of effecting the exit of the discharges from a wound or cavities, by means of tubes or strands of fibers, or by a free incision. **D.-tube**, a rubber or glass tube with perforations for draining wounds or cavities.

Dram, or Drachm (*dram*) [*δραχμή*, a Greek weight]. The eighth part of the apothecaries' ounce, equal to 60 grains or 3.9 grams. Also the one-sixteenth part of the avoirdupois ounce, equal to 27.34 grains. **D.**, **Fluid**, the eighth part of a fluidounce, equal to 60 minims or 3.9 c.c.

Drastic (*dras'-tik*) [*δράειν*, to draw]. 1. Severe, harsh, powerful. 2. A powerful and irritating purgative.

Draught (*draft*). See *Draft*.

Drepanidium (*drop-an-id'-e-um*) [*δρεπάνη*, a sickle]. The sickle-shaped young of certain protozoans. **D. ranarum**, a (probably) parasitic cytozoon of frogs' blood.

Dressing (*dres'-ing*) [ME., *dressen*, to make straight]. 1. The application of various materials for protecting a wound and favoring its healing. 2. The material so applied.

Drop [AS., *dropa*, a drop]. A minute mass of liquid which in falling or in hanging from a surface, assumes the spheric form. **D.**, **Black**, acetum opii. See *Opium*. **D.-culture**, in bacteriology, a culture prepared by placing a little of the infected material in a drop of the culture-medium. **D.**, **Hanging**. See *Hanging Drop*.

Dropped Hand. A form of paralysis from lead-poisoning, consisting in the inability to contract the extensors of the wrist. **D. Foot**, extreme extension of the foot, especially observed in alcoholic neuritis, and dependent upon weakness of the flexors of the foot.

Dropsical (*drop'-sik-al*) [*ὑδρωψία*, dropsy]. Affected with or pertaining to dropsy.

Dropsy (*drop'-se*) [*ὑδρωψία*, dropsy]. An infiltration of the tissues with diluted lymph, or the collection of such lymph in the body cavities. ***D. of Belly**, ascites. **D.**, **Cardiac**, that due to failure of compensation in cardiac disease. **D. of Chest**, hydrothorax. **D. of Peritoneum**, ascites, hydroperitoneum. **D. of Pericardium**, hydropericardium.

Drosera (*dros'-er-ah*) [*δροσερός*, covered with dew]. Sundew; the *D. rotundifolia* and *D. longifolia* have been used in pulmonary tuberculosis. It is an antispasmodic, and is used in whooping-cough and other spasmodic coughs.

Drug [Fr., *drogue*, drug]. A substance used as a medicine. **D.**, **Antagonistic**, one that neutralizes the action of another.

Drum [ME., *drumme*]. The tympanum. See *Enr.* **D.-head**, the membrana tympani.

Drumstick-bacillus. See *Bacteria*, *Table of*.

Dry (*dri*) [AS., *dryge*, dry]. Free from moisture. **D. Cupping**, cupping without incising the skin. **D. Gangrene**. See *Gangrene*. **D. Labor**, one in which there is but a slight discharge of liquor amnii. **D.**

Pleurisy, pleurisy without effusion. **D.**
Wine, a wine containing little or no sugar.
Dubois' Abscess. See *Diseases, Table of*.
Dubini's Disease. See *Diseases, Table of*.
Duboisin (*du-bois'-in*) [after *Dubois*, a French botanist], $C_{17}H_{23}NO_3$. An alkaloid from *Duboisia myoporoides*, identical with hyoscyamin. It is mydriatic, sedative, and hypnotic. The dose of the sulphate, which is the salt generally used, is gr. $\frac{1}{10}$ – $\frac{1}{6}$ (0.00054–0.001).
Duchenne's Disease (*du-shenz'*). Glosso-labiolaryngeal paralysis.
Duct, or **Ductus** [*ducere*, to lead]. A tube or channel, especially one for conveying the secretions of a gland. **D. of Bartholin**, the larger and longer of the sublingual ducts, opening into the mouth near to, or in common with, Wharton's duct. **D.**, **Common Bile**, a duct formed by the union of the cystic and hepatic ducts. **D. of Cuvier**, two short transverse venous trunks in the fetus, one on either side, opening into the auricle of the heart. Each is formed by the union of a superior vein, the primitive jugular, and an inferior vein. The right one becomes the superior vena cava; the left one disappears. **D.**, **Cystic**, the excretory duct of the gall-bladder. **D.**, **Ejaculatory**, a duct formed by the union of the vas deferens and the duct of the seminal vesicle and carrying the semen into the urethra. **D.**, **Endolymphatic**, a tubular process of the membranous labyrinth of the ear, passing through the aqueduct of the vestibule into the cranial cavity, where it terminates below the dura mater in a blind enlargement, the sacculus endolymphaticus. **D.**, **Galactophorous**, one of the milk-ducts of the mammary gland. **D.**, **Hepatic**, a duct formed at the margin of the transverse fissure of the liver by the junction of the right and left hepatic ducts. **D.**, **Lacrymal**. See *D.*, *Nasal*. **D.**, **Lactiferous**. See *D.*, *Galactophorous*. **D.**, **Lymphatic, Right**, the vessel that receives the lymph from the lymphatics of the right arm, the right side of the head and neck, the chest, lung, and right side of the heart, and also from the upper surface of the liver. It terminates at the junction of the right subclavian and internal jugular veins. **D. of Müller**, a duct in the embryo, one on each side, parallel to the Wolffian duct and extending from the body-cavity to the cloaca. From them are derived the oviducts, uterus, and vagina. In the male they atrophy. **D.**, **Nasal**, the duct that conveys the tears from the lacrymal sac into the inferior meatus of the nose. **D.**, **Prostatic**, any one of the ducts conveying the secretion of the prostate into the urethra. **D. of Rivini**, one of the ducts of the sublingual salivary gland. **D.**,

Segmental, a tube, on either side of the body of the embryo, situated between the visceral and parietal layers of the mesoblast, opening anteriorly into the body-cavity, and posteriorly into the cloaca. **D.**, **Spermatic**, the vas deferens. **D. of Steno**. See *D. of Stenson*. **D. of Stenson**, the duct of the parotid gland. **D.**, **Thoracic**, a duct 18 to 20 inches long, beginning in the receptaculum chyli, passing upward, and emptying into the left subclavian vein at its junction with the left internal jugular vein. It receives all the lymph and chyle not received by the right lymphatic duct. **D. of Wharton**, the duct of the submaxillary salivary gland. **D. of Wirsung**, the main duct of the pancreas. It empties into the duodenum. **D.**, **Wolffian**, the duct of the Wolffian body of the embryo. It becomes the vas deferens.
Ductile (*duk'-til*) [*ducere*, to lead]. Capable of being drawn out thin, as a wire or thread.
Ductless Glands. The spleen, thyroid and thymus glands, suprarenal capsules, and pituitary body, which have no excretory duct.
Ductule (*duk'-tūl*) [*ductulus*, a small duct]. A small duct.
Ductus (*duk'-tus*) [*L.*]. A duct. **D. arteriosus**, a short vessel in the fetus connecting the pulmonary artery with the aorta. **D. auditorius** or **cochlearis**, the scala media of the cochlea. **D. cuvieri**. See *Duct of Cuvier*. **D. venosus**, a branch of the umbilical vein in the fetus which empties directly into the ascending vena cava.
Dulcamara (*dul-kam-a'-ra*) [*dulcis*, sweet; *amarus*, bitter]. Bittersweet. The young branches of *D. solanum*, containing an alkaloid, solanin. In overdoses it causes nausea, emesis, and convulsive muscular movements, and in toxic doses is a narcotic poison. It is employed in psoriasis and similar skin-diseases. **D.**, **Ext.**, **Fld.** Dose $\mathfrak{mxxx-f\bar{3}j}$ (2.0.4.0).
Dulcaminarin (*dul-kam-a'-rin*) [*dulcis*, sweet; *amarus*, bitter], $C_{20}H_{24}O_{10}$. A yellow, amorphous alkaloid found in dulcamara, with a sweetish-bitter taste, sparingly soluble in water, freely so in alcohol and acetic acid, insoluble in ether and in chloroform.
Dulness (*dul'-nes*) [*AS.*, *dol*, dull]. Lack of resonance on percussion.
Dumb (*dum*). Unable to utter articulate speech. **D. Ague**, a popular term for ague or malaria marked by obscure symptoms.
Duodenal (*du-o-dē'-nal*, or *du'-od'-en'-al*) [*duodeni*, twelve each]. Relating to the duodenum.
Duodenostomy (*du-o-den-ōs'-to-mē*) [*duodenum*; *stōma*, a mouth]. The operation of forming an opening into the duodenum through the abdominal walls.
Duodenum (*du-o-dē'-num*, or *du'-od'-en-um*)

[*duodeni*, twelve each; so called because it is about twelve finger-breadths long]. The first part of the small intestine beginning at the pylorus. It is from 8 to 10 inches long, is the most fixed part of the small intestine, consists of an ascending, descending, and transverse portion, and contains the openings of the pancreatic duct and the common bile-ducts.

Dupuytren's Contraction. See *Diseases*, *Table of*. **D. Splint**, a splint used in the treatment of Pott's fracture of the leg.

Dura, or **Dura Mater** (*du'-rah ma'-ter*) [*durus*, hard; *mater*, mother]. The fibrous membrane forming the outermost covering of the brain and spinal cord.

Dusting-powder. Any fine powder used to dust on the skin to absorb or diminish its secretions or allay irritation.

Dutch Liquid. See *Ethene Chlorid*.

Dyad (*di'-ad*) [*dyas*, two]. 1. Having a quantivalence of two. 2. An element or radicle having a quantivalence of two.

Dynamic (*di-nam'-ik*) [*dynamis*, power]. Pertaining to energy; characterized by energy or great force.

Dynamics (*di-nam'-iks*) [*dynamis*, power]. See *Mechanics*.

Dynamo (*di'-nam-o*) [*dynamis*, power]. A machine in which an electric current is generated by revolving coils of insulated wire through the field of a magnet intensified by the same current.

Dynamograph (*di-nam'-o-graf*) [*dynamis*, power; *graphein*, to write]. An instrument designed to measure and record graphically muscular strength.

Dynamometer (*di-nam-on'-et-er*) [*dynamis*, power; *metron*, a measure]. An instrument for the measurement of muscular strength, particularly of the hand.

Dyne (*din*) [*dynamis*, power]. A measure of force; it is the force that, when applied to a mass of one gram for one second, will give it a velocity of one centimeter a second.

Dys- [*dy-*], a prefix meaning bad, difficult, painful.

Dysacusma (*dis-ak-oo'-mah*) [*dy-*, painful; *akousia*, hearing]. A sensation of pain or discomfort caused by loud or even moderately loud noises.

Dysacusia, or **Dysacousis** (*dis-ak-oo'-se-ah* or *-sis*) [*dy-*, hard; *akousia*, hearing]. Difficulty of hearing.

Dysæsthesia (*dis-es-the'-se-ah*). See *Dysæsthesia*.

Dysalbumose (*dis-al'-bu-môz*). A variety of albumose, insoluble in hot or cold water or hydrochloric acid.

Dysarthria (*dis-ar'-thre-ah*) [*dy-*, difficult; *arthron*, articulation]. Impairment of articulation.

Dysarthrosis (*dis-ar-thro'-sis*) [*dy-*, bad; a joint]. A deformed joint.

Dysbasia (*dis-ba'-se-ah*) [*dy-*, difficult; *basia*, a step]. Impairment of the power of walking.

Dyschromatopsia (*dis-kro-mat-op'-se-ah*) [*dy-*, difficult; *chroma*, color; *opsis*, vision]. Partial color-blindness.

Dyscrasia (*dis-kra'-se-ah*) [*dy-*, bad; *krasis*, combination]. A depraved condition of the blood or system due to general disease.

Dysenteric (*dis-en-ter'-ik*) [*dy-*, bad; *enteron*, the bowels]. Of the nature of or affected with dysentery.

Dysentery (*dis'-en-ter-e*) [*dy-*, bad; *enteron*, the bowels]. An inflammation of the large intestine, probably infectious in origin, and characterized by pain, rectal tenesmus, and the frequent passage of small amounts of mucus and blood. Anatomically, three varieties may be distinguished: the catarrhal, the diphtheritic, and the gangrenous. The true cause of dysentery is not definitely known, but with many cases the *amœba coli* is intimately associated, particularly in the dysentery of the tropics (*amœbic dysentery*).

Dysæsthesia (*dis-es-the'-se-ah*) [*dy-*, difficult; *aischos*, sensation]. 1. Dulness of sensation. 2. Painfulness of any sensation not normally painful.

Dysgraphia (*dis-graf'-e-ah*) [*dy-*, difficult; *graphein*, to write]. Impairment of the power of writing as a result of a brain-lesion.

Dysidrosis (*dis-id-ro'-sis*) [*dy-*, bad; *idros*, sweating]. Synonym of pompholyx.

Dyslalia (*dis-la'-le-ah*) [*dy-*, difficult; *lalein*, to talk]. Impairment of the power of speaking, due to a defect of the organs of speech.

Dyslexia (*dis-leks'-e-ah*) [*dy-*, difficult; *lexis*, reading]. Impairment of the ability to read.

Dyslogia (*dis-lo'-je-ah*) [*dy-*, difficult; *logos*, speech]. Difficulty in the expression of ideas by speech.

Dysmenorrhea (*dis-men-or-e'-ah*) [*dy-*, difficult; *men*, month; *rhein*, to flow]. Difficult or painful menstruation. **D., Congestive**, a form of painful menstruation due to an intense congestion of the pelvic viscera. **D., Inflammatory**, that due to inflammation. **D., Mechanic**. See *D., Obstructive*.

D., Membranous, a very painful form characterized by the discharge of shreds of decidua. **D., Obstructive**, that due to mechanic obstruction to the free escape of the menstrual fluid. **D., Ovarian**, that form due to disease of the ovaries. **D., Spasmodic**, that form due to spasmodic uterine contraction.

Dysmimia (*dis-mim'-e-ah*) [*dy-*, difficult; *mimhai*, to mimic]. Impairment of the power to use signs and gestures.

Dysosmia (*dis-oo'-me-ah*) [*dy-*, ill; *osme*, odor]. Impairment of the sense of smell.

Dyspareunia (*dis-par-oo'-ne-ah*) [*δυσπαρευνος*, ill-mated]. Painful or difficult copulation.

Dyspepsia (*dis-pep'-se-ah*) [*δύς*, difficult; *πέπειν*, to digest]. Disturbed digestion. **D.**, **Atonic**, a form due to insufficient quantity or impaired quality of the gastric juice, or to deficient action of the gastric muscles. **D.**, **Catarrhal**, that form caused by inflammation of the stomach. **D.**, **Intestinal**, that due to imperfect digestive action of the intestinal juices, or to lack of tone in the muscular coat of the bowel. **D.**, **Nervous**, that form characterized by gastric pains, coming on often when the stomach is empty and relieved by eating, and by various reflex nervous phenomena, especially by palpitation.

Dyspeptic (*dis-pep'-tik*) [*δύς*, difficult; *πέπειν*, to digest]. 1. Relating to or affected with dyspepsia. 2. A person suffering from dyspepsia.

Dyspeptone (*dis-pep'-tōn*) [*δύς*, difficult; *πέπειν*, to digest]. An insoluble and unassimilable peptone.

Dysperistalsis (*dis-per-is-tal'-sis*) [*δύς*, painful; *περί*, around; *σπάσις*, compression]. Painful or perverted peristalsis.

Dysphagia (*dis-fa'-je-ah*) [*δύς*, ill; *φαγεῖν*, to eat]. Difficulty in swallowing, or inability to swallow.

Dysphasia (*dis-fa'-se-ah*) [*δύς*, ill; *φάσις*, speech]. Difficulty of speech depending on a central lesion.

Dysphonia (*dis-fo'-ne-ah*) [*δύς*, difficult; *φωνή*, voice]. An impairment of the voice.

Dysphrasia (*dis-fra'-se-ah*) [*δύς*, ill; *φράσις*, speech]. Imperfect speech, due to impairment of mental power.

Dyspnea (*disp-ne'-ah*) [*δύς*, difficult; *πνέειν*, to breathe]. Difficult or labored breathing.

D., **Cardiac**, that due to heart-disease. **D.**, **Renal**, that due to renal disease.

Dyspneic (*disp-ne'-ik*) [*δύς*, difficult; *πνέειν*, to breathe]. Affected with or caused by dyspnea.

Dyspermia (*dis-sper'-me-ah*) [*δύς*, difficult; *σπέρμα*, seed]. An abnormal condition of the semen.

Dystocia (*dis-to'-se-ah*) [*δύς*, difficult; *τόκος*, birth]. Difficult labor. **D.**, **Fetal**, difficult labor due to abnormalities of position or size and shape of the fetus. **D.**, **Maternal**, that dystocia the cause of which resides in the mother.

Dystrophic (*dis-tro'-fic*) [*δύς*, difficult; *τροφή*, nourishment]. Pertaining to dystrophy.

Dystrophy (*dis'-tro-fe*) [*δύς*, difficult; *τροφή*, nourishment]. Faulty nutrition.

Dysuria (*dis-u'-re-ah*) [*δύς*, difficult; *οὔρον*, urine]. Difficult or painful urination.

E

E. Abbreviation of Eye and Emmetropia.

Ear (*ēr*) [*ΜΕ.*, *ere*, ear]. The organ of hearing, consisting of the external ear, the middle ear or tympanum, and the internal ear or labyrinth. The outer ear is made up of an expanded portion, the pinna, and the external auditory canal. The middle consists of the tympanum, with the ear ossicles, the Eustachian tube, and the mastoid cells. The tympanum is lined by mucous membrane, and communicates with the pharynx by means of the Eustachian tube. It is divided into three parts, the atrium, the attic, and the antrum. Its outer end is closed by the tympanic membrane, from which sound is conducted along the ear-ossicles (the malleus, incus and stapes) to the fenestra ovalis, which communicates with the vestibule of the internal ear. By means of the fenestra rotunda, which is closed by the entotympanic membrane, it communicates with the cochlea of the internal ear. The mastoid cells, which are also part of the tympanum, are air-spaces

in the mastoid process of the temporal bone. The internal ear consists of the bony and membranous labyrinth, which are separated from each other by a space containing the perilymph. Each labyrinth consists of three parts: the vestibule, the semicircular canals, and the cochlea. The bony vestibule communicates with the tympanum by the fenestra ovalis, closed by the base of the stapes, and also with the other parts of the internal ear. The semicircular canals are three in number, the superior, the posterior, and the inferior. The cochlea, so named from its resemblance to a snail-shell, is a cylindric tube that winds around a central axis, the modiolus, which transmits the cochlear nerves and blood-vessels. The cochlear canal is divided by the spiral lamina into the scala vestibuli, communicating with the vestibule, and the scala tympani, communicating with the fenestra rotunda of the tympanum. The membranous labyrinth is made up of parts corresponding to the bony labyrinth. The

vestibule consists of two small sacs, the utricle and saccule, which communicate through the vestibular aqueduct, and are lined by an epithelium, the maculam acusticæ, which in places is largely specialized and receives the terminations of the vestibular nerve. The membranous cochlea, or cochlear duct, contains the acoustic organ of the cochlea, or organ of Corti, which consists of a series of epithelial arches formed by the interlocking of the ends of the pillars or rods of Corti. Upon the inner rods of Corti are the inner acoustic hair-cells; in relation with the outer rods are the outer hair-cells. The organ of Corti is covered by the membrana tectoria, or Corti's membrane.

Ear-wax. See *Cerumen*.

Earth (*erth*) [ME., *erthe*, earth]. A name given to various metallic oxides or silicates, not soluble in water and not affected by great heat. E., Alkaline, the oxides and hydrates of calcium, magnesium, strontium, barium, and other metals of the same group.

Earthy Phosphates. See *Phosphate*.

Eberth's Bacillus. See *Bacteria*, *Table of*.

Ebullition (*eb-ul-lish'-un*) [*ebullire*, to boil]. Boiling.

Eburnation (*e-bur-na'-shun*) [*ebur*, ivory]. An increase in the density of bone following inflammation.

Ecballium (*ek-ball'-e-un*). See *Elatarium*.

Ecbolic (*ek-bol'-ik*) [*ἐκβολή*, a throwing out].

1. Producing or accelerating labor. 2. Any agent producing this effect.

Eccentric (*ek-sen'-trik*) [*ἐκ*, out; *κέντρον*, center]. Situated away from the center.

E. Hypertrophy, hypertrophy of a hollow organ, as the heart, with dilatation.

Ecc hondroma (*ek-kon-dro'-mah*) [*ἐκ*, out; *χόνδρος*, cartilage; *ῥμα*, tumor]. A chondroma.

Ecchymoma (*ek-e-mo'-mah*) [*ἐκχύμαμα*, a bursting of a small blood-vessel]. A tumor-like swelling composed of extravasated blood.

Ecchymosis (*ek-e-mo'-sis*) [*ἐκχύμαμα*, a bursting of a small blood-vessel]. An extravasation of blood into the subcutaneous tissues. It is marked by a purple coloration of the skin, the color gradually changing to brown, green, and yellow.

Ecchymotic (*ek-e-mo'-ik*) [*ἐκχύμαμα*, a bursting of a small blood-vessel]. Relating to or resembling an ecchymosis.

Echinococcus (*e-ki-no-kok'-us*) [*ἰχίνος*, a hedgehog; *κόκκος*, a berry]. 1. The scolex or larval stage of the tænia echinococcus. 2. Hydatid.

Echo (*ek'-o*) [*ἠχώ*, a sound]. A reverberated sound. E., Amphoric, a vocal resonance in which the transmitted voice sounds

as if it were speaking into a narrow-necked bottle.

Echolalia (*ek-o-lal'-le-ah*) [*ἠχώ*, echo; *λαλέω*, talk]. A meaningless repetition by a person of words spoken to him by others.

Eclampsia (*ek-lamp'-se-ah*) [*ἐκλάμπειν*, to shine or burst forth]. An acute nervous affection, characterized by convulsions with loss of consciousness. E., Infantile, a reflex convulsion of childhood. E., Puerperal, a convulsion occurring toward the close of pregnancy, or after labor, believed to be caused by the irritation of the vasomotor centers by retained excrementitious substances.

Eclamptic (*ek-lamp'-tik*) [*ἐκλάμπειν*, to shine or burst forth]. Relating to, or of the nature of, eclampsia.

Eclectic (*ek-ek'-tik*) [*ἐκλεκτικός*, selected]. A name chosen by a school of physicians to denote their principle of selecting that which is good from all other schools.

Écraseur (*a-kras'-ser'*) [Fr.]. An instrument consisting of a chain or wire-loop which is placed about a projecting part, and by being tightened gradually cuts through the tissues.

Ecstasy (*eks'-ta-se*) [*ἔκστασις*, a trance]. A derangement of the nervous system characterized by an exalted visionary state, absence of volition, insensibility to surroundings, a radiant expression, and immobility in statuesque positions.

Ecstrophy (*ek'-stro-fe*) [*ἐκ*, out; *στρέφειν*, to turn]. Eversion; the turning inside-out of a part.

Ectasia, Ectasis (*ek'-ta'-se-ah*, *ek'-tis-is*) [*ἐκτασις*, extension]. Distention; dilatation.

Ectatic (*ek-tat'-ik*) [*ἐκτασις*, extension]. Distended or dilated.

Ecthyma (*ek-thi'-mah*) [*ἐκθύμα*, a pustule]. An inflammatory skin-disease attended with an eruption of large, flat, superficial pustules. They vary in size from a ten-cent piece to a silver quarter-dollar, and are surrounded by a distinct inflammatory areola. The eruption appears as a rule on the legs and thighs where the hairs are thick; it occurs in crops, and may persist for an indefinite period.

Ecto- (*ek'-to-*) [*ἐκτός*, without]. A prefix signifying without, upon the outer side.

Ectoderm (*ek'-to-derm*) [*ἐκτός*, outside; *δέρμα*, skin]. The outer of the two primitive layers of the embryo.

Ectodermal (*ek-to-derm'-al*) [*ἐκτός*, outside, *δέρμα*, skin]. Pertaining to or formed from the ectoderm.

-ectomy [*ἐκ*, out of; *τέμνειν*, to cut]. A suffix, meaning a cutting out.

Ectogenous (*ek-toj'-en-us*) [*ἐκτός*, out; *γενᾶν*, to produce]. Capable of growth outside of the body, applied especially to bacteria and other parasites.

Ectopagus (*ek-toj'-ag-us*) [*ἐκτός*, out; *πάγος*,

a fixture]. A twin monstrosity united laterally the full extent of the thorax.

Ectopia (*ek-to'pe-ah*) [*ἐκτοπος*, displaced]. Malposition.

Ectopic (*ek-to'pik*) [*ἐκτοπος*, displaced]. In an abnormal position. **E. Gestation**, extrauterine gestation.

Ectoplasm (*ek-to-plasm*) [*ἐκτός*, without; *πλάσσειν*, to form]. The outer, hyaline, more compact layer of protoplasm of a cell or unicellular organism.

Ectromelus (*ek-trom'-el-us*) [*ἐκτρωσις*, abortion; *μῆλος*, limb]. A single autositic monster characterized by the presence of imperfectly-formed limbs.

Ectropion (*ek-tro'-pe-on*) [*ἐκ*, out; *τρέπειν*, to turn]. Eversion of a part, especially of an eyelid.

Ectropionize (*ek-tro'-pe-on-ize*) [*ἐκ*, out; *τρέπειν*, to turn]. To produce by operation the condition of ectropion.

Eczema (*ek'-ze-mah*) [*ἐκζεῖν*, to boil over]. Tetter; an acute or chronic, noncontagious, inflammatory disease of the skin, characterized by multiformity of lesion, and the presence, in varying degrees, of itching, infiltration, and discharge. The skin is reddened, the redness shading off insensibly into the surrounding unaffected parts. **E. erythematousum**, the mildest form of eczema, in which the skin is reddened and slightly swollen. **E. fissum**, a form affecting the hands and skin over the articulations, and characterized by the formation of deep, painful cracks or fissures. **E. hypertrophicum**, a form characterized by permanent hypertrophy of the papillae of the skin, giving rise to general or limited warty outgrowths. **E. madidans**, **E. rubrum**; a form characterized by large, raw, weeping surfaces studded with red points. It follows **E. vesiculosum**. **E. papulosum**, a variety associated with the formation of minute papules of a deep-red color and firm consistence, and accompanied by intense itching. **E. pustulosum**, the stage of eczema characterized by the formation of pustules. **E. rubrum**. See *E. madidans*. **E. seborrhœicum**. Synonym of seborrhea. **E. solare**, that form due to irritation from the rays of the sun. **E. squamosum**, a variety characterized by the formation of adherent scales of shed epithelium. **E. vesiculosum**, an eczema characterized by the presence of vesicles.

Eczematous (*ek-ze-m'-at-us*) [*ἐκζεῖν*, to boil over]. Of the nature of or affected with eczema.

Edema (*e-de'-mah*) [*οἰδημα*; *οἰδέειν*, to swell]. An infiltration of serum in a part. **E.**, **Angioneurotic**. See *Angioneurotic*. **E.**, **Inflammatory**, that due to inflammation. **E.**, **Malignant**, an edematous inflammation that

occurs at times after serious injuries, and is characterized by its rapid spread, the speedy destruction of the tissue involved, and the formation of gas. It is due to the bacillus of malignant edema. **E.**, **Purulent**, a purulent infiltration in which there is a great deal of fluid.

Edematous (*e-dem'-at-us*) [*οἰδημα*, edema]. Characterized by edema.

Efferent (*ef'-er-ent*) [*efferens*, carrying from]. Carrying away, as **E. nerves**, nerves conveying impulses away from the central nervous system; of blood-vessels, conveying blood away from the tissues; of lymphatics, conveying lymph from the lymphatic glands.

Effervescing (*ef'-er-ves'-ing*) [*effervesce*, to boil up]. Giving off gas-bubbles; foaming.

E. Powder. See *Scitlitz Powder*.

Effleurage (*ef-flur-azh'*) [*Fr.*]. In massage, the stroking movement.

Efflorescence (*ef-flor-es'-enz*) [*efflorescere*, to bloom]. 1. The spontaneous conversion of a crystalline substance into powder by a loss of its water of crystallization. 2. The eruption of an exanthematous disease.

Effluvium (*ef-lu'-ve-um*) [*effluere*, to flow out]. An odor, especially one that is offensive.

Effusion (*ef-u'-shun*) [*effundere*, to pour out]. 1. A pouring-out, especially the pouring-out of blood or serum into the cellular tissues or the serous cavities. 2. The effused fluid.

Egesta (*e-jes'-tah*) [*egerere*, to cast out]. The discharges of the bowels or of other excretory organs.

Egg (*eg*) [*ME.*, *egge*, an egg]. See *Ovum*. **E.-albumin**, white of egg, constituting about 60 per cent. of the egg of the domestic fowl.

Egophony (*e-goff'-o-ne*) [*αἶψ*, a wild goat; *φωνή*, the voice]. A modification of bronchophony, in which the voice has a bleating character like that of a kid. It is heard in pleurisy with slight effusion.

Egyptian (*e-jip'-le-an*). Pertaining to Egypt.

E. Chlorosis. See *Anchylostomiasis*.

E. Ophthalmia. See *Ophthalmia, Purulent*.

Ehrlich's Anilin-water Solution, or **Ehrlich's Solution**. A solution of a basic anilin-dye in anilin-oil and water.

Eiloid (*i'-loid*) [*εἰλεῖν*, to coil; *εἶδος*, form]. Having a coiled structure, as an eiloid tumor.

Ejaculation (*e-jak-u-la'-shun*) [*ejaculatio*, a throwing out]. The ejection of the semen.

Ejaculatory (*e-jak'-u-la-tor-e*) [*ejaculatio*, a throwing out]. Throwing or casting out.

E. Duct. See *Duct*.

Elastic (*e-las'-tik*) [*ἐλαττειν*, to urge forward]. Returning to the original form after being stretched or compressed. **E. Bandage**, an India-rubber bandage exerting continuous compression of a part. **E. Stocking**,

an India-rubber stocking exerting continuous pressure. **E. Tissue**, a variety of connective tissue composed of yellow elastic fibers.

Elastin (*e-las'-tin*) [*ἐλαστίν*, to urge forward]. An albuminoid substance forming the basis of elastic tissue.

Elaterin, or **Elaterinum** (*el-at'-er-in*, *el-at-er-ī'-num*) [*ἐλατήριος*, driving away], $C_{20}H_{20}O_4$. A neutral principle obtained from *Ecballium elaterium*. It is a powerful hydragogue cathartic. Dose gr. $\frac{1}{16}$ (0.0032). **E.**, **Pulv.**, Comp. (B. P.), contains elaterin 1, sugar of milk 39 parts. Dose gr. ss-v (0.032-0.32). **E.**, **Trituratio**, elaterin 10, sugar of milk 90 parts, thoroughly mixed. Dose gr. ss- $\frac{1}{2}$ (0.032-0.04).

Elaterium (*el-at-e'-re-um*) [*ἐλατήριος*, driving away]. The dried sediment from the juice of the squirting cucumber, *Ecballium elaterium*. It is a powerful hydragogue cathartic. Dose $\frac{1}{8}$ gr. (0.008).

Elbow (*el'-bo*) [*ἄστρον*, *elboga*]. The region corresponding to the junction of the arm and forearm; the bend of the arm. **E.-jerk**. See *Reflexes*, Table of.

Elder (*el'-der*). See *Sambucus*.

Elecampane (*el-e-kam-pān'*). See *Inula*.

Electric (*e-lek'-trik*) [*ἤλεκτρον*, amber]. Having the nature of or produced by electricity. **E. Chorea**. See *Dubini's Disease*, in *Diseases*, Table of.

Electricity (*el-ek-tris'-it-e*) [*ἤλεκτρον*, amber]. One of the forces of nature developed or generated by chemism, magnetism, or friction. **E.**, **Chemic**. See *E.*, *Galvanic*. **E.**, **Faradic**, that produced by induction. **E.**, **Franklinic**, frictional or static electricity. **E.**, **Frictional**, that produced by friction. **E.**, **Galvanic**, that which is generated by chemic action in a galvanic cell. **E.**, **Inductive**, that produced in a body by proximity to an electrified body. **E.**, **Magnetic**, that developed by bringing a conductor near the poles of a magnet. **E.**, **Static**, frictional electricity. **E.**, **Voltaic**. Galvanic or chemic electricity.

Electrify (*el-ek'-trif-i*) [*ἤλεκτρον*, amber; *facere*, to make]. To make electric.

Electrization (*el-ek-tris-a'-shun*) [*ἤλεκτρον*, amber]. The application of electricity to the body.

Electro- (*el-ek'-tro-*) [*ἤλεκτρον*, amber]. A prefix denoting connection with or relation to electricity.

Electrobiology (*el-ek-tro-bi-ol'-o-je*) [*ἤλεκτρον*, amber; *βίος*, life; *λόγος*, science]. The science of the electric relations and laws of organic beings.

Electrobioscopy (*el-ek-tro-bi-os'-ko-pe*) [*ἤλεκτρον*, amber; *βίος*, life; *σκοπεῖν*, to view]. The test of the existence of life by means of electricity.

Electrocautery (*el-ek-tro-kaw'-ter-e*). See *Galvanocautery*.

Electrochemistry (*el-ek-tro-kem'-is-tre*) [*ἤλεκτρον*, amber; *χημεία*, chemistry]. The science treating of the chemic changes produced by electricity.

Electrode (*el-ek'-trōd*) [*ἤλεκτρον*, amber; *ὁδός*, a way]. The pieces of metal or other substance fastened to the conducting cords of a battery through which electricity is applied to the body.

Electrodiagnosis (*el-ek-tro-di-ag-nō'-sis*) [*ἤλεκτρον*, amber; *diagnosis*]. Diagnosis by examining the reaction of the excitable tissues of the body by means of electric currents.

Electrography (*el-ek-trog'-ra-fe*). See *Skio-graphy*.

Electrolysis (*el-ek-trol'-is-is*) [*ἤλεκτρον*, amber; *λύσις*, solution]. The dissolution of a chemic compound by an electric current.

Electrolyte (*el-ek-tro'-lit*) [*ἤλεκτρον*, amber; *λίσις*, solution]. A substance decomposed by an electric current.

Electrolytic (*el-ek-tro'-lit-ik*) [*ἤλεκτρον*, amber; *λύσις*, solution]. Relating to electrolysis.

Electromagnet (*el-ek-tro'-mag-net*) [*ἤλεκτρον*, amber; *magnet*]. A mass of soft iron surrounded by a coil of wire. A current passing through the wire will make the iron core magnetic.

Electromassage (*el-ek-tro-mas-azh'*) [*ἤλεκτρον*, amber; *massage*]. The transmission of a current of electricity through a kneading instrument.

Electrometer (*el-ek-trom'-et-er*) [*ἤλεκτρον*, electricity; *μέτρον*, a measure]. An instrument for measuring electric force.

Electromotive Force (*el-ek-tro-mō'-tiv*). The force which produces an electric current. Symbol, E. M. F.

Electronegative (*el-ek-tro-neg'-a-tiv*) [*ἤλεκτρον*, amber; *negare*, to deny]. Pertaining to the electric condition found at the negative pole of a battery.

Electropathology (*el-ek-tro-path-ol'-o-je*) [*ἤλεκτρον*, amber; *πάθος*, disease; *λόγος*, science]. The study of morbid conditions by the aid of electric irritation.

Electrophysiology (*el-ek-tro-fis-e-ol'-o-je*) [*ἤλεκτρον*, amber; *physiology*]. The study of electric reactions, properties, and relations of organs and organic tissues.

Electropositive (*el-ek-tro-pos'-it-iv*) [*ἤλεκτρον*, amber; *ponere*, to place]. Pertaining to the electric state which exists at the positive pole of a battery.

Electropuncturation, **Electropuncture** (*el-ek-tro-pung-tu-ra'-shun*, *-pung'-tūr*) [*ἤλεκτρον*, amber; *pungere*, to prick]. The use of needles as electrodes, which are thrust into an organ or tumor.

Electrostatics (*el-ek-tro-stat'-iks*) [*ἤλεκτρον*, amber; *στατικός*, causing to stand]. The science of static electricity, or that developed by friction.

Electroscope (*el-ek'-tro-skop*) [*ἤλεκτρον*, amber; *σκοπεῖν*, to view]. An instrument for detecting the presence of static electricity and determining whether it is positive or negative.

Electrotherapeutics (*el-ek-tro-ther-ap-u'-tiks*) [*ἤλεκτρον*, amber; *θεραπεία*, treatment]. The science and art of the application of electricity for therapeutic purposes.

Electrotonus (*el-ek-tro'-o-nus*) [*ἤλεκτρον*, amber; *τόνος*, tension]. The change of condition in a nerve during the passage of a cur-

rent of electricity. See *Anelectrotonus* and *Cutelectrotonus*.

Electuary (*e-lek'-tu-ar-e*) [*electuarium*, an electuary]. A soft or pasty mass, consisting of a medicinal substance, with sugar, honey, water, etc.

Eleidin (*e-le'-id-in*) [*ἐλαία*, olive-oil]. A material occurring in the form of granules in the stratum granulosum of the epidermis.

Element (*el'-em-ent*) [*elementum*, a first principle]. Any one of the ultimate parts of which anything is composed, as the cellular elements of a tissue. In chemistry, a body that cannot be decomposed into simpler substances. The recognized elements now number about 73. See *Table of Elements*.

TABLE OF CHEMIC ELEMENTS.

Arranged alphabetically from *Funk and Wagnall's Standard Dictionary* [copyrighted].
(Published by permission.)

NAME.	SYMBOL.	ATOMIC WEIGHT.	SPECIFIC GRAVITY.*	FUSING-POINT OR MELTING-POINT. DEG. C. AND F.	VALENCE.	WHERE AND HOW FOUND.
Aluminum,	Al	27.0	2.58	627° C.; 1160° F.	III	In many rocks. (The most abundant metal.)
Antimony (<i>stibium</i>),	Sb	120.0	6.70	432° C.; 808° F.	V	Chiefly as sulphid, and in various metallic ores.
Argentum (<i>silver</i>).	A	19.7	1.5†	—128.6° C.		Free in the atmosphere.
Argon,	As	75.0	5.71	Ab't 500° C.; 932° F.	V	Native, as sulphid, and in various metallic ores.
Aurum (<i>gold</i>).	Ba	137.0	3.75	Above redness.	II	In barite and witherite.
Barium,	Bi	208.9	9.80	268° C.; 517° F.	V	Native, as sulphid, and in rare minerals.
Beryllium (<i>glucinum</i>).	B	11.0	2.6	Very high.	III	In borax and various minerals.
Bismuth,	Br	79.95	3.19	—7.2° C.; —20° F.	I or VII	Mainly in sea-water and other natural brines.
Boron,	Cd	112.0	8.65	231° C.; 609° F.	II	In small amount in zinc ores.
Bromin,	Ca	40.0	1.6 to 1.8	Bright redness.	II	In limestone, and abundantly in other rocks.
Cadmium,	C	12.0	3.52‡	Infusible.	IV	In coal, limestone, and all organic matter.
Calcium,	Ce	140.2	6.7	Below silver.	III or IV	In cerite and other rare minerals.
Carbon,	Cs	132.9	1.88	26.5° C.; 80° F.	I	In lepidolite, pollucite, and mineral springs.
Cerium,	Cl	35.45	1.33†	—75.6° C.; —103° F.	I or VII	In common salt (NaCl), and other chlorids.
Cesium,	Cr	52.1	7.3	Above platinum.	II or VI	Mainly in chrome-iron ore.
Chlorin,	Co	59.0	8.96	1500° C.; 2732° F.	II or VIII	In many metallic ores.
Chromium,	Cb	94.0	Above 7	V	In columbite and other rare minerals.
Cobalt,	Cu	63.6	8.9	1054° C.; 1931° F.	I or II	Native, and in many ores.
Columbium (<i>niobium</i>),	Er	166.3	III	In rare minerals, as gadolinite, etc.
Copper (<i>cuprum</i>),	F	19.0	I or VII	In fluorite (CaF ₂) and other minerals.
Erbium,						
Ferrum (<i>iron</i>).						
Fluorin,						

TABLE OF CHEMIC ELEMENTS.—Continued.

NAME.	SYMBOL.	ATOMIC WEIGHT.	SPECIFIC GRAVITY.	FREEZING-POINT OR MELTING-POINT, DEG. C. AND F.	VALENCY.	WHERE AND HOW FOUND.
Gadolinium, . . .	Gd	156.1	III	In rare minerals, as gadolinite.
Gallium,	Ga	69.0	5.95	30.1° C.; 86° F.	III	In certain zinc-ores.
Germanium, . . .	Ge	72.3	5.47	900° C.; 1652° F.	IV	In argentic & rare minerals.
Glaucium (<i>beryl- lum</i>),	Gl	9.0	1.85	Above redness.	II	In beryl and several rare minerals.
Gold (<i>aurum</i>), . .	Au	197.3	19.3	1045° C.; 1913° F.	I or III	Generally free, rarely combined in various ores.
Hydrargyrum (<i>mercury</i>).						
Hydrogen,	H	1.008	0.085†	—200° C.; —320° F.	I	Mainly in water, H ₂ O.
Indium,	In	113.7	7.4	176° C.; 349° F.	III	In certain zinc-ores.
Iodine,	I	125.85	4.95	114° C.; 238° F.	I or VII	Mainly in ashes of seaweeds.
Iridium,	Ir	193.1	22.4	1950° C.; 3542° F.	II or IV	In iridosmin.
Iron (<i>ferrum</i>), . .	Fe	56.0	8.0	1600° C.; 2912° F.	II or IV	As oxide and sulphide, and in nearly all rocks.
Kalium (<i>potassium</i>).						
Lanthanum,	La	138.2	6.1	...	III	In cerite and other rare minerals.
Lead (<i>plumbum</i>), .	Pb	206.95	11.36	326° C.; 650° F.	II or IV	In galena, PbS, and other ores.
Lithium,	Li	7.02	0.585	180° C.; 350° F.	I	In lepidolite, spodumene, and some rare minerals.
Magnesium, . . .	Mg	24.3	1.75	Abt 1430° C.; 506° F.	II	In sea-water magnesite, and many rocks.
Manganese,	Mn	55.0	7.2	Above iron.	II or VII	In pyrolusite, and many other minerals.
Masrium,	Ms	238.	II	In bed of old river in upper Egypt, in masrite.
Mercury (<i>hydrargyrum</i>).	Hg	200.0	13.556	—38.8° C.; —38° F.	I or II	Native, and in cinnabar (HgS).
Molybdenum, . . .	Mo	96.0	8.6	Very high.	II or VI	Mainly as molybdenite (MoS ₂).
Sodium (<i>sodium</i>).						
Neodymium,	Nd	140.5	About 6.5	...	III or IV	In cerite and other rare minerals.
Nickel,	Ni	58.0	8.9	1450° C.; 2642° F.	II or VIII	In many metallic ores.
Niobium (<i>columbium</i>).						
Nitrogen,	N	14.03	0.38	...	V	In the atmosphere and organic matter.
Osmium,	Os	190.08	22.48	Nearly infusible.	II or VII	In iridosmin and native platinum.
Oxygen,	O	16.0	1.11	...	II or VI	Free in air. (Forms one-half the earth's crust, combined.)
Palladium,	Pd	106.6	12.1	1500° C.; 2732° F.	II or IV	Native, and with platinum and gold.
Phosphorus, . . .	P	31.0	1.84	44.2° C.; 112° F.	V	In bones, and in apatite and many minerals.
Platinum,	Pt	195.0	21.5	1775° C.; 3225° F.	II or IV	Mainly as native platinum in river gravels.
Plumbum (<i>lead</i>).						
Potassium (<i>kalium</i>).	K	39.11	0.86	62.5° C.; 144.5° F.	I	In wood-ashes and many rocks.
Praseodymium, . .	Pr	143.5	About 6.5	...	III or IV	In cerite and other rare minerals.
Rhodium,	Rh	103.0	12.1	2000° C.; 3632° F.	II or VIII	With platinum and iridosmin.
Rubidium,	Rb	85.5	1.52	38.5° C.; 101.5° F.	I	In lepidolite and some mineral springs.
Ruthenium, . . .	Ru	101.6	12.26	Nearly infusible.	—	With platinum and iridosmin.

TABLE OF CHEMIC ELEMENTS.—*Continued.*

NAME.	SYMBOL.	ATOMIC WEIGHT.	SPECIFIC GRAVITY.*	FUSING-POINT OR MELTING-POINT. DEG. C. AND F.	VALENCE.	WHERE AND HOW FOUND.
Samarium,	Sm	150.0	III	In samarskite, cerite, and other rare minerals.
Scandium,	Sc	44.0	III	In gadolinite and other rare minerals.
Selenium,	Se	79.0	4.5	217° C.; 425° F.	II or VI	Mainly in sulphur as an impurity.
Silicou,	Si	28.4	2.48	Above 800° C.; 1500° F.	IV	In quartz (SiO ₂). Most abundant element after oxygen.
Silver (<i>argentum</i>),	Ag	107.9	10.5	954° C.; 1750° F.	I	Native, and in many ores.
Sodium (<i>natrium</i>),	Na	23.05	0.97	95.6° C.; 204° F.	I	In common salt (NaCl) and many rocks.
Stannum (<i>tin</i>).						
Stibium (<i>antimony</i>).						
Strontium,	Sr	87.6	2.5	Red heat.	II	In celestite and strontianite.
Sulphur,	S	32.06	2.07	114.5° C.; 235° F.	II or VI	Native, and in many natural sulphids and sulphates.
Tantalum,	Ta	182.6	Above 10	V	In tantalite and other rare minerals.
Tellurium,	Te	125.0	6.23	455° C.; 851° F.	II or VI	In several rare minerals.
Terbium,	Tb	160.0	III	In rare minerals, as gadolinite, etc.
Thallium,	Tl	204.18	11.19	239.9° C.; 561° F.	I or III	In pyrites and in flue-dust of sulphuric-acid works.
Thorium,	Th	232.6	11.23	Almost infusible.	IV	In thorite and other rare minerals.
Thulium,	Tu	170.7	III	In rare minerals, as gadolinite, etc.
Tin (<i>stannum</i>), . .	Sn	119.0	7.25	233° C.; 551° F.	II or IV	Mainly in cassiterite (SnO ₂).
Titanium,	Ti	48.0	. . .	Not fusible.	V	Widely diffused in rocks and clays, in small amounts.
Tungsten (<i>wol-</i>	W	184.0	19.26	Very high.	IV or VI	Mainly in wolframite (MnFeWO ₄).
framium); . . . }						
Uranium,	U	239.6	18.69	Very high.	II or VI	In pitch-blende and other rare minerals.
Vanadium,	V	51.4	5.87	In oxyhydric flame.	V	In vanadinite and other rare minerals.
Wolframium (<i>tung-</i>						
sten).						
Ytterbium,	Yb	173.0	III	In rare minerals, as gadolinite, etc.
Yttrium,	Yt	89.1	III	In gadolinite and other rare minerals.
Zinc (<i>zincum</i>), . .	Zn	65.3	7.12	433° C.; 811.5° F.	II	In ores, as oxid, silicate, sulphid, and carbonate.
Zirconium,	Zr	90.06	4.15	Above sulphur.	IV	In zircon and other rare minerals.

* The factors in the columns of specific gravities and melting-points naturally vary with the form which the element takes (*e.g.*, in carbon the specific gravity varies as diamond, charcoal, or lampblack is taken), but as far as possible the factor of the most typical form is given.

† Of the liquid element. ‡ Diamond. § Of the liquid at 0° C. ¶ Of the liquid at -181° C.

Elementary (*el-e-men'-ta-re*) [*elementum*, a first principle]. Pertaining to or having the characters of an element.

Elemi (*el'-em-e*) [Arab.]. A resinous exudation probably derived from the *Canarium commune*, although its botanic source is still undetermined. It contains a crystalline re-

sin, elemin or amylin. Its action is similar to that of the turpentine. **E.**, Unguent. (B. P.), elemi and simple ointment; it is used as an application to indolent sores and boils.

Elephantiasis (*el-ef-an-ti'-as-is*) [*ἐλεφας*, an elephant]. A chronic affection of the cuta-

neous and subcutaneous tissues, due to obstruction of lymph-vessels, and characterized by enormous thickening of the affected parts. The disease occurs in successive attacks accompanied by fever, and by swelling of the affected parts, usually the lower extremities and genital organs; it is endemic in certain tropical countries, and seems to be connected, in many cases, with the presence in the blood of the *Filaria sanguinis hominis*. **E. arabum**. See *Elephantiasis*. **E. græcorum**. See *Leprosy*. **E. telangiectodes**, elephantiasis characterized by a great increase in the blood-vessels.

Elevator (*el'-ev-a-tor*) [*elevare*, to lift]. 1. The same as *Levator*. See *Muscles*, Table of. 2. An instrument for elevating or lifting a part.

Elimination (*e-lim-in-a'-shun*) [*e*, out; *limen*, threshold]. The process of expelling or casting out, especially waste-products.

Elixir (*el-iks'-ir*) [Arab., *el iksir*, the philosopher's stone]. A sweetened, aromatic, spirituous preparation, containing only a small amount of an active ingredient. **E. aromaticum** (U.S.). Compound spirit of orange, 1.2, syrup 37.5, deodorized alcohol, distilled water, each a sufficient quantity to make 100 c.c. It is used as a vehicle. **E. phosphori**, spirit of phosphorus 21, oil of anise 0.2, glycerol 55, aromatic elixir a sufficient quantity to make 100 cc.

Elkoplasty (*el'-ko-plas-ty*). See *Helcoplasty*. **Elm**. See *Ulmus*.

Elutriation (*e-lu-tre-a'-shun*) [*elutriare*, to wash out]. A process whereby the coarser particles of an insoluble substance are separated from the finer by decanting the fluid after the coarser particles have settled.

Elytritis (*el-it-ri'-tis*) [*ἐλντρον*, sheath; *ιτις*, inflammation]. Inflammation of the vagina.

Elytrophasty (*el'-it-ro-plas-te*) [*ἐλντρον*, sheath; *πλάσσειν*, to form]. A plastic operation upon the vagina.

Elytrophosis (*el'-it-rof-to'-sis*) [*ἐλντρον*, sheath; *πρῶσις*, a falling]. Prolapse of the vagina.

Elytrophaphy (*el'-it-rof-a-fe*) [*ἐλντρον*, sheath; *φαγή*, a seam]. Suture of the vaginal wall.

Emansio (*e-man'-se-o*) [L.]. A failing. **E. mensium**, delay in the first appearance of the menses.

Embedding (*em-bed'-ing*) [*in*, in; AS., *bed*, bed]. The fixation of a tissue-specimen in a firm medium, in order to keep it intact during the cutting of thin sections.

Embole (*em'-bo-le*). See *Emboly*.

Embolic (*em-bol'-ik*) [*ἐν*, in; *βάλλειν* to throw]. Relating to or caused by an embolus.

Emboliform (*em-bol'-if-orm*) [*ἐν*, in; *βάλ-*

λειν, to throw; *forma*, form]. Resembling an embolus.

Embolism (*em'-bo-lizm*) [*ἐν*, in; *βάλλειν*, to throw]. The obstruction of a blood vessel, especially an artery, by a fragment of matter brought from another point. **E.**, Air, obstruction of a vessel by a bubble of air. **E.**, Fat, obstruction of blood vessels by globules of fat. **E.**, Infective, embolism in which the emboli contain microorganisms and cause metastatic abscesses. **E.**, Miliary, a condition in which many small blood vessels are the seats of emboli.

Embololalia (*em-bo-lo-lal'-e-ah*) [*ἐν*, in; *βάλλειν*, to throw; *λαλέω*, talk]. The intercalation of meaningless words into the speech.

Embolus (*em'-bo-lus*) [*ἐν*, in; *βάλλειν*, to throw]. A particle of fibrin or other material brought by the blood current and forming an obstruction at its place of lodgment.

Emboly (*em'-bo-le*) [*ἐμβόλη*, insertion]. The process of invagination that gives rise to a gastrula from a blastosphere or vesicular morula.

Embrocation (*em-bro-ka'-shun*) [*ἐμβρῆσαι*, to soak in]. 1. The application, especially by rubbing, of a liquid to a part of the body. 2. The liquid so applied.

Embryo (*em'-bre-o*) [*ἐν*, within; *βρίσκειν*, to swell with]. 1. The product of conception up to the fourth month of pregnancy. 2. The fertilized germ of an animal.

Embryocardia (*em-bre-o-kar'-de-ah*) [*ἐν*, within; *βρίσκειν*, to swell with; *καρδία*, the heart]. A condition in which the heart sounds resemble those of the fetus, the first and second sounds being almost identical.

Embryogenetic (*em-bre-o-jen'-et-ik*) [*ἐν*, within; *βρίσκειν*, to swell with]. Giving rise to an embryo.

Embryology (*em-bre-ol'-o-jé*) [*ἐν*, within; *βρίσκειν*, to swell with; *λογία*, science]. The science dealing with the development of the embryo.

Embryonic (*em-bre-on'-ik*) [*ἐν*, within; *βρίσκειν*, to swell with]. Pertaining to the embryo.

E. Area, an opaque circular spot that forms on the blastoderm. **E. Tissue**, tissue in the undifferentiated state, consisting of small, round cells. **E. Spot**. See *E. Area*.

Embryotome (*em'-bre-o-tom*) [*ἐν*, within; *βρίσκειν*, to swell with; *τομή*, section]. An instrument for performing embryotomy.

Embryotomy (*em-bre-ot'-o-mé*) [*ἐν*, within; *βρίσκειν*, to swell with; *τομή*, section]. The cutting up of the fetus in utero for the purpose of reducing its size.

Emesis (*em'-es-is*) [*ἐμέειν*, to vomit]. Vomiting.

Emetic (*e-met'-ik*) [*ἰμετικός*, causing vomiting]. 1. Having the power to induce vomiting. 2. An agent causing emesis. **E.**,

granulations. This is also called hemorrhagic endometritis. **E., Hemorrhagic.** See *E., Fungous*. **E., Simple,** a catarrhal inflammation of the endometrium.

Endometrium (*en-do-mē'-tre-um*) [*ἐνδον*, within; *μήτρα*, uterus]. The mucous membrane lining the uterus.

Endomysium (*en-do-mīz'-e-um*) [*ἐνδον*, within; *μῦς*, muscle]. The connective tissue between the fibrils of a muscular bundle.

Endoneurium (*en-do-nū'-re-um*) [*ἐνδον*, within; *νεῖρον*, a nerve]. The delicate connective tissue holding together the fibrils of a bundle of nerves.

Endoparasite (*en-do-par'-as-īt*) [*ἐνδον*, within; *παράσιτος*, parasite]. A parasite living within its host.

Endoscope (*en'-do-skōp*) [*ἐνδον*, within; *σκοπεῖν*, to observe]. An instrument for the examination of a body-cavity through its natural outlet.

Endoskeleton (*en-do-skēl'-et-un*) [*ἐνδον*, within; *σκελετόν*, a dry body]. The internal supporting structure of an animal.

Endosmometer (*en-dos-mom'-et-er*) [*ἐνδον*, within; *ωμός*, a thrusting; *μέτρον*, a measure]. An instrument for measuring endosmosis.

Endosmosis (*en-dos-mō'-sis*) [*ἐνδον*, within; *ωμός*, impulsion]. The passage of a liquid through a porous septum from without inward.

Endosmotic (*en-dos-mōt'-ik*) [*ἐνδον*, within; *ωμός*, a thrusting]. Pertaining to endosmosis. **E. Equivalent,** the weight of distilled water that passes into the flask of the endosmometer in exchange for a known weight of the soluble substance.

Endospore (*en'-do-spōr*) [*ἐνδον*, within; *σπόρος*, seed]. A spore formed within the parent-cell.

Endosteum (*end-os'-te-um*) [*ἐνδον*, within; *ὀστέον*, bone]. The vascular membranous layer of connective tissue lining the medullary cavity of bones.

Endothelial (*en-do-the'-le-al*) [*ἐνδον*, within; *θηλή*, nipple]. Pertaining to endothelium.

Endothelioma (*en-do-the-le-o'-mah*) [*ἐνδον*, within; *θηλή*, a nipple; *ῥυμα*, a tumor]. A variety of sarcoma, formed by the multiplication of the endothelial cells of lymphatic spaces.

Endothelium (*en-do-the'-le-um*) [*ἐνδον*, within; *θηλή*, nipple]. See *Cell*, *Endothelial*.

Enema (*en'-em-ah*) [*ἐν*, in; *ἔναι*, to send]. A rectal injection for therapeutic or nutritive purposes.

Energy (*en'-er-ge*) [*ἐν*, in; *ἐργον*, work]. The capacity for doing work. All forms of energy are mutually convertible one into the other, without loss, a principle expressed in the term "conservation of energy." **E., Kinetic,** the power of a body in motion. **E.,**

Latent, E., Potential, the power possessed by a body at rest, by virtue of its position, as the potential *E.* of a suspended weight.

Engorged (*en-gorj'd*) [*Fr., engorgement*, a choking up]. Congested.

Engorgement (*en-gorj'-ment*) [*Fr., engorgement*, a choking up]. Over-distention of the vessels of a part with blood.

Enophthalmos (*en-off-thall'-mos*) [*ἐν*, in; *ὀφθαλμός*, the eye]. Recession of the eyeball into the orbit.

Enostosis (*en-os-to'-sis*) [*ἐν*, in; *ὀστέον*, bone]. A tumor or bony outgrowth within the medullary canal of a bone.

Ensiform (*en'-sif-orm*) [*ensis*, a sword; *forma*, form]. Shaped like a sword. **E. Appendix,** the cartilaginous process at the lower extremity of the sternum.

Ensomphalus (*en-som'-fal-us*) [*ἐν*, in; *σῶμα*, body; *ὀμφαλός*, navel]. A double monstrosity, with practically complete and functioning organs, but united with a more or less superficial bond.

Enteralgia (*en-ter-al'-je-ah*) [*ἐντερον*, intestine; *ἄλγος*, pain]. Pain in the bowels.

Enterectomy (*en-ter-ek'-to-me*) [*ἐντερον*, intestine; *ἐκτομή*, excision]. Excision of a part of the intestine.

Enteropilecele (*en-ter-ep'-ip'-lo-sēl*) [*ἐντερον*, intestine; *ἐπιπλοον*, caul; *κῆλη*, hernia]. Hernia in which both bowel and omentum are involved.

Enteric (*en-ter'-ik*) [*ἐντερον*, intestine]. Pertaining to the intestines. **E. Fever,** typhoid fever.

Enteritis (*en-ter-i'-tis*) [*ἐντερον*, bowel; *ιτις*, inflammation]. Inflammation of the intestine.

Entero- (*en'-ter-o-*) [*ἐντερον*, intestine]. A prefix denoting relation to the intestine.

Enterocoele (*en'-ter-o-sēl*) [*ἐντερον*, bowel; *κῆλη*, tumor]. A hernia containing a loop of intestine.

Entercholecystostomy (*en-ter-o-ko-le-sis-tōs'-to-me*). Same as *Cholecystenterostomy*.

Enteroclysis (*en-ter-ok'-lis-is*) [*ἐντερον*, bowel; *κλύσις*, a drenching]. Injection of a large quantity of fluid into the rectum to reach the small intestine.

Enterocolitis (*en-ter-o-ko-lī'-tis*) [*ἐντερον*, intestine; *κόλον*, colon; *ιτις*, inflammation]. Inflammation of the small intestine and of the colon.

Enteropilecele (*en-ter-o-e-pip'-lo-sēl*). See *Enteropilecele*.

Enterogastritis (*en-ter-o-gas-trī'-tis*) [*ἐντερον*, intestine; *γαστήρ*, stomach; *ιτις*, inflammation]. Inflammation of the stomach and intestine.

Enterolith (*en'-ter-o-līth*) [*ἐντερον*, bowel; *λίθος*, a stone]. A concretion formed in the intestines.

Enteropathy (*en-ter-op'-ath-e*) [*έντερον*, bowel; *πάθος*, disease]. Any disease of the intestines.

Enteroplasty (*en'-ter-o-plas-te*) [*έντερον*, bowel; *πλασσειν*, to form]. A plastic operation upon the intestine.

Enterorrhagia (*en-ter-ar-aj'-e-ah*) [*έντερον*, bowel; *ρήγνιναι*, to burst forth]. Intestinal hemorrhage.

Enterorrhaphy (*en-ter-ar'-a-fe*) [*έντερον*, bowel; *ράφη*, suture]. Suture of the intestine.

Enterostomy (*en-ter-os'-to-me*) [*έντερον*, bowel; *στόμα*, mouth]. The formation of an artificial opening into the intestine through the abdominal wall.

Enterotome (*en'-ter-o-tōm*) [*έντερον*, bowel; *τέμνειν*, to cut]. An instrument for cutting open the intestine.

Enterotomy (*en-ter-ot'-o-me*) [*έντερον*, bowel; *τέμνειν*, to cut]. Incision of the intestine.

Enterozoön (*en-ter-o-zō-on*) [*έντερον*, intestine; *ζῶον*, an animal]. An animal parasite of the intestine.

Enthetic (*en-thet'-ik*) [*ἐνθεῖναι*, to put in]. Introduced; coming from without; applied especially to syphilitic and other specific contagious diseases.

Entoblast (*en'-to-blast*) [*έντός*, within; *βλαστός*, a bud, germ]. 1. The nucleolus of a cell. 2. The endoderm.

Entophyte (*en'-to-fit*) [*έντός*, within; *φυτόν*, a plant]. A vegetable parasite living within the body of its host, as *e.g.*, a bacterium.

Entoplastic (*en-to-plas'-tik*) [*έντός*, within; *πλασσειν*, to form]. 1. Having an endoplast or nucleus. 2. Pertaining to the nucleus.

Entoptic (*ent-op'-tik*) [*έντός*, within; *ὀπτική*, pertaining to vision]. Pertaining to the internal parts of the eye. **E. Phenomena**, visual sensations generated within the eye.

Entotic (*ent-ot'-ik*) [*έντός*, within; *ὅς*, ear]. Pertaining to the internal parts of the ear.

Entozoön (*en-to-zō-on*) [*έντός*, within; *ζῶον*, an animal]. An animal parasite living within another animal.

Entropion (*en-tro'-pe-on*) [*έν*, in; *τρέπειν*, to turn]. Inversion of the eyelid, so that the lashes rub against the globe of the eye.

Enucleation (*e-nu-kle-a'-shun*) [*ε*, out of; *nucleus*, a kernel]. The shelling-out of a tumor or organ from its capsule. The excision of the eye-ball.

Enuresis (*en-u-re'-sis*) [*ἐνουρησις*, to be incontinent of urine]. The involuntary emptying of the bladder. **E., Nocturnal**, that occurring at night, during sleep.

Enzyme (*en'-zim*) [*έν*, in; *ζῆν*, to leaven]. 1. Any ferment formed within the living organism. 2. A chemie ferment, as distinguished from organized ferments, such as the yeasts.

Eosin (*e'-o-sin*) [*ῥῶς*, the dawn], $C_{20}H_2Br_2O_6$. Tetrabromfluorescein; an acid dye produced by the action of bromin on fluorescein suspended in glacial acetic acid. It occurs in red or yellowish crystals, and is used as a stain in histology.

Eosinophile (*e-o-sin'-o-fil*) [*eosin*, *φιλεῖν*, to love]. Showing a peculiar affinity for eosin-stain or for acid-stains in general.

Epencephalon (*ep-en-sef'-al-on*) [*ἐπι*, on; *ἐγκέφαλος*, brain]. The after-brain or hind-brain; the cerebellum and pons taken together.

Ependyma (*ep-en'-dim-ah*) [*ἐπένδυμα*, an upper garment]. The lining membrane of the cerebral ventricles and of the central canal of the spinal cord.

Ependymitis (*ep-en-dim-it'-tis*) [*ἐπένδυμα*, an upper garment; *τις*, inflammation]. Inflammation of the ependyma.

Ephedra (*ef'-e-drah*) [*ἐπί*, upon; *ἑδρα*, a seat]. A genus of plants of the Gnetaceæ. **E. anti-syphilitica**, has been used in gonorrhea. Dose of the fluid extract, f ʒj-ij (4.0-8.0). **E. vulgaris**, contains the alkaloid ephedrin, which is mydriatic.

Ephelis (*ef'-el-is*) [*ἐπί*, on; *ἥλος*, wart]. A freckle.

Ephemeral (*ef-em'-er-al*) [*ἐφήμερος*, living a day]. Temporary. Applied to fevers that pass away in a day.

Ephidrosis (*ef-id-ro'-sis*) [*ἐπί*, upon; *ἰδρωσις*, sweating]. Excessive perspiration. See *Hyperidrosis*.

Epiblast (*ep'-e-blast*) [*ἐπί*, upon; *βλαστός*, a sprout]. The external or upper layer of the blastoderm; called also the ectoderm, from which are developed the central nervous system and the epithelium of the sense-organs, the mucous membranes of the mouth and anus, the enamel of the teeth, the epidermis and its derivatives (hair, nails, glands, etc.).

Epiblastic (*ep-e-blast'-ik*) [*ἐπί*, upon; *βλαστός*, a sprout]. Pertaining to or derived from the epiblast.

Epibole, Epiboly (*ep-ib'-ol-e*) [*ἐπί*, upon; *βάλλειν*, to throw]. The enclosure of the large yolk-mass of an invertebrate ovum by the overgrowth of cleavage-cells.

Epicanthus (*ep-e-kan'-thus*) [*ἐπί*, on; *κανθός*, angle of the eye]. A fold of skin over the inner canthus of the eye.

Epicondyle (*ep-e-kon'-dl*) [*ἐπί*, upon; *κόνδυλος*, a knuckle]. An eminence upon a bone above its condyle.

Epicranium (*ep-e-kra'-ne-um*) [*ἐπί*, upon; *κρανιον*, cranium]. The structures covering the cranium.

Epicranius (*ep-e-kra'-ne-us*) [*ἐπί*, upon; *κρανιον*, cranium]. The occipito-frontalis muscle.

Epicystotomy (*ep-e-sis-tot'-o-me*) [*ἐπί*, upon; *κυστις*, a bladder; *τέμνειν*, to cut]. Suprapubic incision of the bladder.

granulations. This is also called hemorrhagic endometritis. **E., Hemorrhagic.** See *E., Fungous*. **E., Simple**, a catarrhal inflammation of the endometrium.

Endometrium (*en-do-mé'-tre-um*) [*éndov*, within; *μήτρα*, uterus]. The mucous membrane lining the uterus.

Endomysium (*en-do-miz'-e-um*) [*éndov*, within; *μῦς*, muscle]. The connective tissue between the fibrils of a muscular bundle.

Endoneurium (*en-do-nú'-re-um*) [*éndov*, within; *νεῖρον*, a nerve]. The delicate connective tissue holding together the fibrils of a bundle of nerves.

Endoparasite (*en-do-par'-as-ít*) [*éndov*, within; *παράσιτος*, parasite]. A parasite living within its host.

Endoscope (*en'-do-skóp*) [*éndov*, within; *σκοπεῖν*, to observe]. An instrument for the examination of a body-cavity through its natural outlet.

Endoskeleton (*en-do-skél'-et-un*) [*éndov*, within; *σκελετόν*, a dry body]. The internal supporting structure of an animal.

Endosmometer (*en-dos-mom'-et-er*) [*éndov*, within; *ώσμός*, a thrusting; *μέτρον*, a measure]. An instrument for measuring endosmosis.

Endosmosis (*en-dos-mó'-sis*) [*éndov*, within; *ώσμός*, impulsion]. The passage of a liquid through a porous septum from without inward.

Endosmotic (*en-dos-mó'-tik*) [*éndov*, within; *ώσμός*, a thrusting]. Pertaining to endosmosis. **E. Equivalent**, the weight of distilled water that passes into the flask of the endosmometer in exchange for a known weight of the soluble substance.

Endospore (*en'-do-spór*) [*éndov*, within; *σπόρος*, seed]. A spore formed within the parent-cell.

Endosteum (*en-dó'st'-e-um*) [*éndov*, within; *ὀστέον*, bone]. The vascular membranous layer of connective tissue lining the medullary cavity of bones.

Endothelial (*en-do-thé'-le-al*) [*éndov*, within; *θηλή*, nipple]. Pertaining to endothelium.

Endothelioma (*en-do-the-le-ó'-mah*) [*éndov*, within; *θηλή*, a nipple; *δύμα*, a tumor]. A variety of sarcoma, formed by the multiplication of the endothelial cells of lymphatic spaces.

Endothelium (*en-do-thé'-le-um*) [*éndov*, within; *θηλή*, nipple]. See *Cell*, *Endothelial*.

Enema (*en'-em-ah*) [*én*, in; *έναι*, to send]. A rectal injection for therapeutic or nutritive purposes.

Energy (*en'-er-ge*) [*én*, in; *έργον*, work]. The capacity for doing work. All forms of energy are mutually convertible one into the other, without loss, a principle expressed in the term "conservation of energy." **E., Kinetic**, the power of a body in motion. **E.,**

Latent, E., Potential, the power possessed by a body at rest, by virtue of its position, as the potential *E.* of a suspended weight.

Engorged (*en-gorj'd*) [*Fr.*, *engorgement*, a choking up]. Congested.

Engorgement (*en-gorj'-ment*) [*Fr.*, *engorgement*, a choking up]. Over-distention of the vessels of a part with blood.

Enophthalmos (*en-off-thal'-mos*) [*én*, in; *ὀφθαλμός*, the eye]. Recession of the eyeball into the orbit.

Enostosis (*en-os-to'-sis*) [*én*, in; *ὀστέον*, bone]. A tumor or bony outgrowth within the medullary canal of a bone.

Ensiform (*en'-sif-orm*) [*ensis*, a sword; *forma*, form]. Shaped like a sword. **E. Appendix**, the cartilaginous process at the lower extremity of the sternum.

Ensomphalus (*en-som'-fal-us*) [*én*, in; *σῶμα*, body; *ἀμφαλός*, navel]. A double monstrosity, with practically complete and functioning organs, but united with a more or less superficial bond.

Enteralgia (*en-ter-all'-je-ah*) [*έντερον*, intestine; *ἄλγος*, pain]. Pain in the bowels.

Enterectomy (*en-ter-ek'-to-me*) [*έντερον*, intestine; *ἐκτομή*, excision]. Excision of a part of the intestine.

Enteropilecele (*en-ter-ep-íp'-lo-sél*) [*έντερον*, intestine; *ἐπίπλοον*, caul; *κήλη*, hernia]. Hernia in which both bowel and omentum are involved.

Enteric (*en-ter'-ik*) [*έντερον*, intestine]. Pertaining to the intestines. **E. Fever**, typhoid fever.

Enteritis (*en-ter-ít'-tis*) [*έντερον*, bowel; *itis*, inflammation]. Inflammation of the intestine.

Entero- (*en'-ter-o-*) [*έντερον*, intestine]. A prefix denoting relation to the intestine.

Enterocoele (*en'-ter-o-sél*) [*έντερον*, bowel; *κήλη*, tumor]. A hernia containing a loop of intestine.

Enterocholecystostomy (*en-ter-o-ko-le-sis-tó'st'-to-me*). Same as *Cholecystenterostomy*.

Enteroclysis (*en-ter-ok'-lís-is*) [*έντερον*, bowel; *κλύσις*, a drenching]. Injection of a large quantity of fluid into the rectum to reach the small intestine.

Enterocolitis (*en-ter-o-ko-lí'-tis*) [*έντερον*, intestine; *κόλον*, colon; *itis*, inflammation]. Inflammation of the small intestine and of the colon.

Enteropilecele (*en-ter-o-e-íp'-lo-sél*). See *Enteropilecele*.

Enterogastitis (*en-ter-o-gas-trí'-tis*) [*έντερον*, intestine; *γαστήρ*, stomach; *itis*, inflammation]. Inflammation of the stomach and intestine.

Enterolith (*en'-ter-o-lith*) [*έντερον*, bowel; *λίθος*, a stone]. A concretion formed in the intestines.

Enteropathy (*en-ter-ōp' -ath-e*) [*έντερον*, bowel; *πάθος*, disease]. Any disease of the intestines.

Enteroplasty (*en' -ter-o- plas- te*) [*έντερον*, bowel; *πλάσσειν*, to form]. A plastic operation upon the intestine.

Enterorrhagia (*en-ter-or-aj' -e-ah*) [*έντερον*, bowel; *ρηγνύναι*, to burst forth]. Intestinal hemorrhage.

Enterorhaphy (*en-ter-or' -a-fe*) [*έντερον*, bowel; *ῥαφή*, suture]. Suture of the intestine.

Enterostomy (*en-ter-os' -to-me*) [*έντερον*, bowel; *στόμα*, mouth]. The formation of an artificial opening into the intestine through the abdominal wall.

Enterotome (*en' -ter-o-tōm*) [*έντερον*, bowel; *τέμνειν*, to cut]. An instrument for cutting open the intestine.

Enterotomy (*en-ter-ōt' -o-me*) [*έντερον*, bowel; *τέμνειν*, to cut]. Incision of the intestine.

Enterozoon (*en-ter-o-zō-on*) [*έντερον*, intestine; *ζῶον*, an animal]. An animal parasite of the intestine.

Enthetic (*en-thet' -ik*) [*ένθῆναι*, to put in]. Introduced; coming from without; applied especially to syphilitic and other specific contagious diseases.

Entoblast (*en' -to-blast*) [*έντός*, within; *βλαστός*, a bud, germ]. 1. The nucleolus of a cell. 2. The endoderm.

Entophyte (*en' -to-fit*) [*έντός*, within; *φυτόν*, a plant]. A vegetable parasite living within the body of its host, as *e.g.*, a bacterium.

Entoplastic (*en-to-plas' -tik*) [*έντός*, within; *πλάσσειν*, to form]. 1. Having an endoplast or nucleus. 2. Pertaining to the nucleus.

Entoptic (*ent-ōp' -tik*) [*έντός*, within; *ὀπτικός*, pertaining to vision]. Pertaining to the internal parts of the eye. **E. Phenomena**, visual sensations generated within the eye.

Entotic (*ent-ōt' -ik*) [*έντός*, within; *ὄψ*, ear]. Pertaining to the internal parts of the ear.

Entozoon (*en-to-zō-on*) [*έντός*, within; *ζῶον*, an animal]. An animal parasite living within another animal.

Entropion (*en-tro' -pe-on*) [*έν*, in; *τρέπειν*, to turn]. Inversion of the eyelid, so that the lashes rub against the globe of the eye.

Enucleation (*e-nu-kle-ō' -shun*) [*έν*, out of; *nucleus*, a kernel]. The shelling-out of a tumor or organ from its capsule. The excision of the eye-ball.

Enuresis (*en-u-re' -sis*) [*ένουρῆσαι*, to be incontinent of urine]. The involuntary emptying of the bladder. **E., Nocturnal**, that occurring at night, during sleep.

Enzyme (*en' -zim*) [*έν*, in; *ζυμῶν*, leaven]. 1. Any ferment formed within the living organism. 2. A chemie ferment, as distinguished from organized ferments, such as the yeasts.

Eosin (*e' -o-sin*) [*ήώς*, the dawn], $C_{20}H_8Br_2O_3$. Tetrabromfluorescein; an acid dye produced by the action of bromin on fluorescein suspended in glacial acetic acid. It occurs in red or yellowish crystals, and is used as a stain in histology.

Eosinophile (*e-o-sin' -o-fil*) [*eosin*, *φιλεῖν*, to love]. Showing a peculiar affinity for eosin-stain or for acid-stains in general.

Epencephalon (*ep-en-sef' -al-on*) [*ἐπι*, on; *ἐγκέφαλος*, brain]. The after-brain or hind-brain; the cerebellum and pons taken together.

Ependyma (*ep-en' -dim-ah*) [*ἐπένδυμα*, an upper garment]. The lining membrane of the cerebral ventricles and of the central canal of the spinal cord.

Ependymitis (*ep-en-dim-i' -tis*) [*ἐπένδυμα*, an upper garment; *ίτις*, inflammation]. Inflammation of the ependyma.

Ephedra (*ef' -e-drah*) [*ἐπί*, upon; *ἐδρα*, a seat]. A genus of plants of the Gnetaceae. **E. anti-syphilitica**, has been used in gonorrhoea. Dose of the fluid extract, f ʒj-ij (4.0-8.0). **E. vulgaris**, contains the alkaloid ephedrin, which is mydriatic.

Ephelis (*ef' -el-is*) [*ἐπί*, on; *ἥλος*, wart]. A freckle.

Ephemeral (*ef-eni' -er-al*) [*ἐφήμερος*, living a day]. Temporary. Applied to fevers that pass away in a day.

Ephidrosis (*ef-id-ro' -sis*) [*ἐπί*, upon; *ἰδρωσις*, sweating]. Excessive perspiration. See *Hyp-eridrosis*.

Epiblast (*ep' -e-blast*) [*ἐπί*, upon; *βλαστός*, a sprout]. The external or upper layer of the blastoderm; called also the ectoderm, from which are developed the central nervous system and the epithelium of the sense-organs, the mucous membranes of the mouth and anus, the enamel of the teeth, the epidermis and its derivatives (hair, nails, glands, etc.).

Epiblastic (*ep-e-blast' -ik*) [*ἐπί*, upon; *βλαστός*, a sprout]. Pertaining to or derived from the epiblast.

Epibole, **Epiboly** (*ep-ib' -ol-e*) [*ἐπί*, upon; *βάλλειν*, to throw]. The enclosure of the large yolk-mass of an invertebrate ovum by the overgrowth of cleavage-cells.

Epicanthus (*ep-e-kan' -thus*) [*ἐπί*, on; *καίνθος*, angle of the eye]. A fold of skin over the inner canthus of the eye.

Epicondyle (*ep-e-kon' -dl*) [*ἐπί*, upon; *κόνδυλος*, a knuckle]. An eminence upon a bone above its condyle.

Epicranium (*ep-e-kra' -ne-um*) [*ἐπί*, upon; *κράνιον*, cranium]. The structures covering the cranium.

Epicranius (*ep-e-kra' -ne-us*) [*ἐπί*, upon; *κράνιον*, cranium]. The occipito-frontalis muscle.

Epicystotomy (*ep-e-sis-tōt' -o-me*) [*ἐπί*, upon; *κύστις*, a bladder; *τέμνειν*, to cut]. Suprapubic incision of the bladder.

granulations. This is also called hemorrhagic endometritis. **E., Hemorrhagic.** See *E., Fungous*. **E., Simple**, a catarrhal inflammation of the endometrium.

Endometrium (*en-do-mé'-tre-um*) [*éndov*, within; *μήτρα*, uterus]. The mucous membrane lining the uterus.

Endomysium (*en-do-mis'-e-um*) [*éndov*, within; *μῦς*, muscle]. The connective tissue between the fibrils of a muscular bundle.

Endoneurium (*en-do-nu'-re-um*) [*éndov*, within; *νεῖρον*, a nerve]. The delicate connective tissue holding together the fibrils of a bundle of nerves.

Endoparasite (*en-do-par'-as-ít*) [*éndov*, within; *παράσιτος*, parasite]. A parasite living within its host.

Endoscope (*en'-do-skóp*) [*éndov*, within; *σκοπεῖν*, to observe]. An instrument for the examination of a body-cavity through its natural outlet.

Endoskeleton (*en-do-skél'-et-um*) [*éndov*, within; *σκελετόν*, a dry body]. The internal supporting structure of an animal.

Endosmometer (*en-dos-mom'-et-er*) [*éndov*, within; *ώσμός*, a thrusting; *μέτρον*, a measure]. An instrument for measuring endosmosis.

Endosmosis (*en-dos-mo'-sis*) [*éndov*, within; *ώσμός*, impulsion]. The passage of a liquid through a porous septum from without inward.

Endosmotic (*en-dos-mot'-ik*) [*éndov*, within; *ώσμός*, a thrusting]. Pertaining to endosmosis. **E. Equivalent**, the weight of distilled water that passes into the flask of the endosmometer in exchange for a known weight of the soluble substance.

Endospore (*en'-do-spór*) [*éndov*, within; *σπόρος*, seed]. A spore formed within the parent-cell.

Endosteum (*en-dos'-te-um*) [*éndov*, within; *όστέον*, bone]. The vascular membranous layer of connective tissue lining the medullary cavity of bones.

Endothelial (*en-do-the'-le-al*) [*éndov*, within; *θηλή*, nipple]. Pertaining to endothelium.

Endothelioma (*en-do-the-le-ó'-mah*) [*éndov*, within; *θηλή*, a nipple; *όμα*, a tumor]. A variety of sarcoma, formed by the multiplication of the endothelial cells of lymphatic spaces.

Endothelium (*en-do-the'-le-um*) [*éndov*, within; *θηλή*, nipple]. See *Cell*, *Endothelial*.

Enema (*en'-em-ah*) [*év*, in; *έναι*, to send]. A rectal injection for therapeutic or nutritive purposes.

Energy (*en'-er-ge*) [*év*, in; *έργον*, work]. The capacity for doing work. All forms of energy are mutually convertible one into the other, without loss, a principle expressed in the term "conservation of energy." **E., Kinetic**, the power of a body in motion. **E.,**

Latent, **E., Potential**, the power possessed by a body at rest, by virtue of its position, as the potential **E.** of a suspended weight.

Engorged (*en-gorj'd*) [*Fr.*, *engorgement*, a choking up]. Congested.

Engorgement (*en-gorj'-ment*) [*Fr.*, *engorgement*, a choking up]. Over-distention of the vessels of a part with blood.

Enophthalmos (*en-óff-thal'-mos*) [*év*, in; *όφθαλμός*, the eye]. Recession of the eyeball into the orbit.

Enostosis (*en-os-to'-sis*) [*év*, in; *όστέον*, bone]. A tumor or bony outgrowth within the medullary canal of a bone.

Ensiform (*en'-sif-orm*) [*ensis*, a sword; *forma*, form]. Shaped like a sword. **E. Appendix**, the cartilaginous process at the lower extremity of the sternum.

Ensomphalus (*en-som'-fal-us*) [*év*, in; *σώμα*, body; *όμφαλός*, navel]. A double monstrosity, with practically complete and functioning organs, but united with a more or less superficial bond.

Enteralgia (*en-ter-al'-je-ah*) [*έντερον*, intestine; *άλγος*, pain]. Pain in the bowels.

Enterectomy (*en-ter-ek'-to-me*) [*έντερον*, intestine; *εκτομή*, excision]. Excision of a part of the intestine.

Enteropileoce (*en-ter-ep-íp'-lo-sel*) [*έντερον*, intestine; *επιπλοον*, caul; *κήλη*, hernia]. Hernia in which both bowel and omentum are involved.

Enteric (*en-ter'-ik*) [*έντερον*, intestine]. Pertaining to the intestines. **E. Fever**, typhoid fever.

Enteritis (*en-ter-i'-tis*) [*έντερον*, bowel; *ιτις*, inflammation]. Inflammation of the intestine.

Entero- (*en'-ter-o-*) [*έντερον*, intestine]. A prefix denoting relation to the intestine.

Enterocoele (*en'-ter-o-sel*) [*έντερον*, bowel; *κήλη*, tumor]. A hernia containing a loop of intestine.

Entercholecystostomy (*en-ter-o-ko-le-sis-tos'-to-me*). Same as *Cholecystenterostomy*.

Enteroclysis (*en-ter-ók'-lis-is*) [*έντερον*, bowel; *κλίσαις*, a drenching]. Injection of a large quantity of fluid into the rectum to reach the small intestine.

Enterocolitis (*en-ter-o-ko-li'-tis*) [*έντερον*, intestine; *κόλον*, colon; *ιτις*, inflammation]. Inflammation of the small intestine and of the colon.

Enteropileoce (*en-ter-o-e-íp'-lo-sel*). See *Enteropileoce*.

Enterogastitis (*en-ter-o-gas-tri'-tis*) [*έντερον*, intestine; *γαστήρ*, stomach; *ιτις*, inflammation]. Inflammation of the stomach and intestine.

Enterolith (*en'-ter-o-lith*) [*έντερον*, bowel; *λίθος*, a stone]. A concretion formed in the intestines.

Enteropathy (*en-ter-op'-ath-e*) [*έντερον*, bowel; *πάθος*, disease]. Any disease of the intestines.

Enteroplasty (*en'-ter-o-plas-te*) [*έντερον*, bowel; *πλασσειν*, to form]. A plastic operation upon the intestine.

Enterorrhagia (*en-ter-ar-aj'-e-ah*) [*έντερον*, bowel; *ρήγνιναι*, to burst forth]. Intestinal hemorrhage.

Enterorrhaphy (*en-ter-ar'-a-fe*) [*έντερον*, bowel; *ράφω*, suture]. Suture of the intestine.

Enterostomy (*en-ter-os'-to-me*) [*έντερον*, bowel; *στόμα*, mouth]. The formation of an artificial opening into the intestine through the abdominal wall.

Enterotome (*en'-ter-o-tom*) [*έντερον*, bowel; *τέμνειν*, to cut]. An instrument for cutting open the intestine.

Enterotomy (*en-ter-at'-o-me*) [*έντερον*, bowel; *τέμνειν*, to cut]. Incision of the intestine.

Enterozoon (*en-ter-o-zo'-on*) [*έντερον*, intestine; *ζών*, an animal]. An animal parasite of the intestine.

Enthetic (*en-thet'-ik*) [*ένθελναι*, to put in]. Introduced; coming from without; applied especially to syphilis and other specific contagious diseases.

Entoblast (*en'-to-blast*) [*έντός*, within; *βλαστός*, a bud, germ]. 1. The nucleolus of a cell. 2. The endoderm.

Entophyte (*en'-to-fit*) [*έντός*, within; *φυτόν*, a plant]. A vegetable parasite living within the body of its host, as *e.g.*, a bacterium.

Entoplastic (*en-to-plas'-tik*) [*έντός*, within; *πλασσειν*, to form]. 1. Having an endoplast or nucleus. 2. Pertaining to the nucleus.

Entoptic (*ent-op'-tik*) [*έντός*, within; *οπτικός*, pertaining to vision]. Pertaining to the internal parts of the eye. **E. Phenomena**, visual sensations generated within the eye.

Entotic (*ent-at'-ik*) [*έντός*, within; *οὖς*, ear]. Pertaining to the internal parts of the ear.

Entozoon (*en-to-zo'-on*) [*έντός*, within; *ζών*, an animal]. An animal parasite living within another animal.

Entropion (*en-tro'-pe-on*) [*έν*, in; *τρέπειν*, to turn]. Inversion of the eyelid, so that the lashes rub against the globe of the eye.

Enucleation (*e-nu-kle-at'-shun*) [*e*, out of; *nucleus*, a kernel]. The shelling-out of a tumor or organ from its capsule. The excision of the eye-ball.

Enuresis (*en-u-re'-sis*) [*ένουρπειν*, to be incontinent of urine]. The involuntary emptying of the bladder. **E., Nocturnal**, that occurring at night, during sleep.

Enzyme (*en'-zim*) [*έν*, in; *ζύμν*, leaven]. 1. Any ferment formed within the living organism. 2. A chemie ferment, as distinguished from organized ferments, such as the yeasts.

Eosin (*e'-o-sin*) [*ήώς*, the dawn], $C_{20}H_8Br_2O_5$. Tetrabromfluorescein; an acid dye produced by the action of bromin on fluorescein suspended in glacial acetic acid. It occurs in red or yellowish crystals, and is used as a stain in histology.

Eosinophile (*e-o-sin'-o-fil*) [*eosin*, *φιλειν*, to love]. Showing a peculiar affinity for eosin-stain or for acid-stains in general.

Epencephalon (*ep-en-sef'-al-on*) [*έπι*, on; *εγκέφαλος*, brain]. The after-brain or hind-brain; the cerebellum and pons taken together.

Ependyma (*ep-en'-dim-ah*) [*έπένδυμα*, an upper garment]. The lining membrane of the cerebral ventricles and of the central canal of the spinal cord.

Ependymitis (*ep-en-dim-it'-tis*) [*έπένδυμα*, an upper garment; *τις*, inflammation]. Inflammation of the ependyma.

Ephedra (*ef'-e-drah*) [*έπι*, upon; *έδρα*, a seat]. A genus of plants of the Gnetaceæ. **E. anti-syphilitica**, has been used in gonorrhea. Dose of the fluid extract, fʒj-ij (4.0-8.0). **E. vulgaris**, contains the alkaloid ephedrin, which is mydriatic.

Ephelis (*ef'-el-is*) [*έπι*, on; *ήλος*, wart]. A freckle.

Ephemeral (*ef-em'-er-al*) [*εφήμερος*, living a day]. Temporary. Applied to fevers that pass away in a day.

Ephidrosis (*ef-id-ro'-sis*) [*έπι*, upon; *ιδρωσις*, sweating]. Excessive perspiration. See *Hyperidrosis*.

Epiblast (*ep'-e-blast*) [*έπι*, upon; *βλαστός*, a sprout]. The external or upper layer of the blastoderm; called also the ectoderm, from which are developed the central nervous system and the epithelium of the sense-organs, the mucous membranes of the mouth and anus, the enamel of the teeth, the epidermis and its derivatives (hair, nails, glands, etc.).

Epiblastic (*ep-e-blast'-ik*) [*έπι*, upon; *βλαστός*, a sprout]. Pertaining to or derived from the epiblast.

Epibole, Epiboly (*ep-ib'-ol-e*) [*έπι*, upon; *βάλλειν*, to throw]. The enclosure of the large yolk-mass of an invertebrate ovum by the overgrowth of cleavage-cells.

Epicanthus (*ep-e-kan'-thus*) [*έπι*, on; *καιθός*, angle of the eye]. A fold of skin over the inner canthus of the eye.

Epicondyle (*ep-e-kon'-dl*) [*έπι*, upon; *κόνδυλος*, a knuckle]. An eminence upon a bone above its condyle.

Epicranium (*ep-e-kra'-ne-um*) [*έπι*, upon; *κρανιον*, cranium]. The structures covering the cranium.

Epicranius (*ep-e-kra'-ne-us*) [*έπι*, upon; *κρανιον*, cranium]. The occipito-frontalis muscle.

Epicystotomy (*ep-e-sis-tot'-o-me*) [*έπι*, upon; *κυστις*, a bladder; *τέμνειν*, to cut]. Suprapubic incision of the bladder.

granulations. This is also called hemorrhagic endometritis. **E., Hemorrhagic.** See *E., Fungous*. **E., Simple**, a catarrhal inflammation of the endometrium.

Endometrium (*en-do-mē'tre-um*) [*ēvdov*, within; *μήτρα*, uterus]. The mucous membrane lining the uterus.

Endomysium (*en-do-miē'-e-um*) [*ēvdov*, within; *μῦς*, muscle]. The connective tissue between the fibrils of a muscular bundle.

Endoneurium (*en-do-nū're-um*) [*ēvdov*, within; *νεῖρον*, a nerve]. The delicate connective tissue holding together the fibrils of a bundle of nerves.

Endoparasite (*en-do-par'-as-it*) [*ēvdov*, within; *παράσιτος*, parasite]. A parasite living within its host.

Endoscope (*en'-do-skōp*) [*ēvdov*, within; *σκοπεῖν*, to observe]. An instrument for the examination of a body-cavity through its natural outlet.

Endoskeleton (*en-do-skēl'-et-un*) [*ēvdov*, within; *σκελετόν*, a dry body]. The internal supporting structure of an animal.

Endosmometer (*en-dos-mōm'-et-er*) [*ēvdov*, within; *ώσμός*, a thrusting; *μέτρον*, a measure]. An instrument for measuring endosmosis.

Endosmosis (*en-dos-mō'-sis*) [*ēvdov*, within; *ώσμός*, impulsion]. The passage of a liquid through a porous septum from without inward.

Endosmotic (*en-dos-mōt'-ik*) [*ēvdov*, within; *ώσμός*, a thrusting]. Pertaining to endosmosis. **E. Equivalent**, the weight of distilled water that passes into the flask of the endosmometer in exchange for a known weight of the soluble substance.

Endospore (*en'-do-spōr*) [*ēvdov*, within; *σπόρος*, seed]. A spore formed within the parent-cell.

Endosteum (*en-dōs'-te-um*) [*ēvdov*, within; *ὀστέον*, bone]. The vascular membranous layer of connective tissue lining the medullary cavity of bones.

Endothelial (*en-do-thē'-le-al*) [*ēvdov*, within; *θηλή*, nipple]. Pertaining to endothelium.

Endothelioma (*en-do-the-le-ō'-mah*) [*ēvdov*, within; *θηλή*, a nipple; *δῆμα*, a tumor]. A variety of sarcoma, formed by the multiplication of the endothelial cells of lymphatic spaces.

Endothelium (*en-do-the'-le-um*) [*ēvdov*, within; *θηλή*, nipple]. See *Cell*, *Endothelial*.

Enema (*en'-em-ah*) [*iv*, in; *έναι*, to send]. A rectal injection for therapeutic or nutritive purposes.

Energy (*en'-er-ge*) [*iv*, in; *έργον*, work]. The capacity for doing work. All forms of energy are mutually convertible one into the other, without loss, a principle expressed in the term "conservation of energy." **E., Kinetic**, the power of a body in motion. **E.,**

Latent, **E., Potential**, the power possessed by a body at rest, by virtue of its position, as the potential **E.** of a suspended weight.

Engorged (*en-gorj'd*) [*Fr.*, *engorgement*, a choking up]. Congested.

Engorgement (*en-gorj'-ment*) [*Fr.*, *engorgement*, a choking up]. Over-distention of the vessels of a part with blood.

Enophthalmos (*en-off-thal'-mos*) [*iv*, in; *ὀφθαλμός*, the eye]. Recession of the eyeball into the orbit.

Enostosis (*en-os-to'-sis*) [*iv*, in; *ὀστέον*, bone]. A tumor or bony outgrowth within the medullary canal of a bone.

Ensiform (*en'-sif-orm*) [*ensis*, a sword; *forma*, form]. Shaped like a sword. **E. Appendix**, the cartilaginous process at the lower extremity of the sternum.

Ensomphalus (*en-som'-fal-us*) [*iv*, in; *σῶμα*, body; *ὀμφαλός*, navel]. A double monstrosity, with practically complete and functioning organs, but united with a more or less superficial bond.

Enteralgia (*en-ter-all'-je-ah*) [*ēvterov*, intestine; *ἄλγος*, pain]. Pain in the bowels.

Enterectomy (*en-ter-ek'-to-me*) [*ēvterov*, intestine; *ἐκτομή*, excision]. Excision of a part of the intestine.

Enteropilecele (*en-ter-ep-ip'-lo-sēl*) [*ēvterov*, intestine; *ἐπιπλοον*, caul; *κῆλη*, hernia]. Hernia in which both bowel and omentum are involved.

Enteric (*en-ter'-ik*) [*ēvterov*, intestine]. Pertaining to the intestines. **E. Fever**, typhoid fever.

Enteritis (*en-ter-i'-tis*) [*ēvterov*, bowel; *τις*, inflammation]. Inflammation of the intestine.

Entero- (*en'-ter-o-*) [*ēvterov*, intestine]. A prefix denoting relation to the intestine.

Enterocoele (*en'-ter-o-sēl*) [*ēvterov*, bowel; *κῆλη*, tumor]. A hernia containing a loop of intestine.

Enterocholecystostomy (*en-ter-o-ko-le-sis-tōs'-to-me*). Same as *Cholecystenterostomy*.

Enteroclysis (*en-ter-ōk'-līs-is*) [*ēvterov*, bowel; *κλίσις*, a drenching]. Injection of a large quantity of fluid into the rectum to reach the small intestine.

Enterocolitis (*en-ter-o-ko-lī'-tis*) [*ēvterov*, intestine; *κόλον*, colon; *τις*, inflammation]. Inflammation of the small intestine and of the colon.

Enteropilecele (*en-ter-o-e-pip'-lo-sēl*). See *Enteropilecele*.

Enterogastitis (*en-ter-o-gas-trī'-tis*) [*ēvterov*, intestine; *γαστήρ*, stomach; *τις*, inflammation]. Inflammation of the stomach and intestine.

Enterolith (*en'-ter-o-lith*) [*ēvterov*, bowel; *λίθος*, a stone]. A concretion formed in the intestines.

Enteropathy (*en-ter-op'-ath-e*) [*έντερον*, bowel; *πάθος*, disease]. Any disease of the intestines.

Enteroplasty (*en'-ter-o-plas-te*) [*έντερον*, bowel; *πλασσειν*, to form]. A plastic operation upon the intestine.

Enterorrhagia (*en-ter-or-aj'-e-ah*) [*έντερον*, bowel; *ρήγνυναι*, to burst forth]. Intestinal hemorrhage.

Enterorrhaphy (*en-ter-or'-a-fe*) [*έντερον*, bowel; *ράφω*, suture]. Suture of the intestine.

Enterostomy (*en-ter-os'-to-me*) [*έντερον*, bowel; *στόμα*, mouth]. The formation of an artificial opening into the intestine through the abdominal wall.

Enterotome (*en'-ter-o-tōm*) [*έντερον*, bowel; *τέμνειν*, to cut]. An instrument for cutting open the intestine.

Enterotomy (*en-ter-ot'-o-me*) [*έντερον*, bowel; *τέμνειν*, to cut]. Incision of the intestine.

Enterozoon (*en-ter-o-zō'-on*) [*έντερον*, intestine; *ζῷον*, an animal]. An animal parasite of the intestine.

Ethnetic (*en-thet'-ik*) [*ένθιθίναί*, to put in]. Introduced; coming from without; applied especially to syphilitic and other specific contagious diseases.

Entoblast (*en'-to-blast*) [*έντός*, within; *βλαστός*, a bud, germ]. 1. The nucleolus of a cell. 2. The endoderm.

Entophyte (*en'-to-fit*) [*έντός*, within; *φυτόν*, a plant]. A vegetable parasite living within the body of its host, as *e.g.*, a bacterium.

Entoplastic (*en-to-plas'-tik*) [*έντός*, within; *πλασσειν*, to form]. 1. Having an endoplast or nucleus. 2. Pertaining to the nucleus.

Entoptic (*ent-op'-tik*) [*έντός*, within; *ὀπτικός*, pertaining to vision]. Pertaining to the internal parts of the eye. **E. Phenomena**, visual sensations generated within the eye.

Entotic (*ent-ot'-ik*) [*έντός*, within; *οὖς*, ear]. Pertaining to the internal parts of the ear.

Entozoon (*en-to-zō'-on*) [*έντός*, within; *ζῷον*, an animal]. An animal parasite living within another animal.

Entropion (*en-tro'-pe-on*) [*έν*, in; *τρέπειν*, to turn]. Inversion of the eyelid, so that the lashes rub against the globe of the eye.

Enucleation (*e-nu-kle-a'-shun*) [*ε*, out of; *nucleus*, a kernel]. The shelling-out of a tumor or organ from its capsule. The excision of the eye-ball.

Enuresis (*en-u-re'-sis*) [*ένονρεύν*, to be incontinent of urine]. The involuntary emptying of the bladder. **E., Nocturnal**, that occurring at night, during sleep.

Enzyme (*en'-zim*) [*έν*, in; *ζύμη*, leaven]. 1. Any ferment formed within the living organism. 2. A chemic ferment, as distinguished from organized ferments, such as the yeasts.

Eosin (*e'-o-sin*) [*ήώς*, the dawn], $C_{20}H_2Br_4O_5$. Tetrabromfluorescein; an acid dye produced by the action of bromin on fluorescein suspended in glacial acetic acid. It occurs in red or yellowish crystals, and is used as a stain in histology.

Eosinophile (*e-o-sin'-o-fil*) [*eosin*, φιλεῖν, to love]. Showing a peculiar affinity for eosin-stain or for acid-stains in general.

Epencephalon (*ep-en-sef'-al-on*) [*ἐπί*, on; *ἐγκέφαλος*, brain]. The after-brain or hind-brain; the cerebellum and pons taken together.

Ependyma (*ep-en'-dim-ah*) [*ἐπέδνυμα*, an upper garment]. The lining membrane of the cerebral ventricles and of the central canal of the spinal cord.

Ependymitis (*ep-en-dim-it'-tis*) [*ἐπέδνυμα*, an upper garment; *ιτις*, inflammation]. Inflammation of the ependyma.

Ephedra (*eff'-e-drah*) [*ἐπί*, upon; *ἔδρα*, a seat]. A genus of plants of the Gnetaceæ. **E. anti-syphilitica**, has been used in gonorrhea. Dose of the fluid extract, fʒj-ij (4.0-8.0). **E. vulgaris**, contains the alkaloid ephedrin, which is mydriatic.

Ephelis (*eff'-el-is*) [*ἐπί*, on; *ἦλος*, wart]. A freckle.

Ephemeral (*ef-em'-er-al*) [*ἐφήμερος*, living a day]. Temporary. Applied to fevers that pass away in a day.

Ephidrosis (*ef-id-ro'-sis*) [*ἐπί*, upon; *ἰδρωσις*, sweating]. Excessive perspiration. See *Hyperidrosis*.

Epiblast (*ep'-e-blast*) [*ἐπί*, upon; *βλαστός*, a sprout]. The external or upper layer of the blastoderm; called also the ectoderm, from which are developed the central nervous system and the epithelium of the sense-organs, the mucous membranes of the mouth and anus, the enamel of the teeth, the epidermis and its derivatives (hair, nails, glands, etc.).

Epiblastic (*ep-e-blast'-ik*) [*ἐπί*, upon; *βλαστός*, a sprout]. Pertaining to or derived from the epiblast.

Epibole, **Epiboly** (*ep-ib'-ol-e*) [*ἐπί*, upon; *βάλλειν*, to throw]. The enclosure of the large yolk-mass of an invertebrate ovum by the overgrowth of cleavage-cells.

Epicanthus (*ep-e-kant'-thus*) [*ἐπί*, on; *κανθός*, angle of the eye]. A fold of skin over the inner canthus of the eye.

Epicondyle (*ep-e-kon'-dil*) [*ἐπί*, upon; *κόνδυλος*, a knuckle]. An eminence upon a bone above its condyle.

Epicranium (*ep-e-kra'-ne-um*) [*ἐπί*, upon; *κρανίον*, cranium]. The structures covering the cranium.

Epicranius (*ep-e-kra'-ne-us*) [*ἐπί*, upon; *κρανίον*, cranium]. The occipito-frontalis muscle.

Epicystotomy (*ep-e-sis-tot'-o-me*) [*ἐπί*, upon; *κυστις*, a bladder; *τέμνειν*, to cut]. Suprapubic incision of the bladder.

Epicyte (*ep'-e-sīt*) [*ἐπί*, upon; *κύτος*, cell]. The cell-wall.

Epidemic (*ep-e-dem'-ik*) [*ἐπί*, upon; *ἄμος*, people]. Of a disease, affecting large numbers, or spreading over a wide area.

Epidemiography (*ep-e-dem-e-og'-ra-fe*) [*ἐπί*, upon; *ἄμος*, people; *γράφειν*, to write]. A description of epidemic diseases.

Epidemiology (*ep-e-dem-e-ol'-o-je*) [*ἐπί*, upon; *ἄμος*, people; *λόγος*, science]. The science of epidemic diseases.

Epidermatic, Epidermic (*ep-e-der-mat'-ik, -der'-mik*) [*ἐπί*, upon; *δέρμα*, skin]. Relating to the epidermis. **E. Method**, a method of administering medicinal substances by applying them to the skin.

Epidermis (*ep-e-der'-mis*) [*ἐπί*, upon; *δέρμα*, the skin]. The outer layer of the skin. The scarf-skin, consisting of a layer of horny cells, that protects the true skin.

Epididymis (*ep-e-did'-im-is*) [*ἐπί*, upon; *διδυμοί*, the testes]. The small body lying above the testis. The superior end is the globus major, the inferior, the globus minor.

Epididymitis (*ep-e-did-im-i'-tis*) [*ἐπιιδιδυμίς*, epididymis; *ιτις*, inflammation]. Inflammation of the epididymis.

Epidural (*ep-e-dur'-ral*) [*ἐπί*, upon; *durus*, hard]. Situated upon or over the dura.

Epigastric (*ep-e-gas'-trik*) [*ἐπί*, upon; *γαστήρ*, belly]. Relating to the epigastrium. **E. Reflex**. See *Reflexes*, *Table of*.

Epigastrium (*ep-e-gas'-tre-um*) [*ἐπί*, upon; *γαστήρ*, stomach]. The upper and middle part of the abdominal surface corresponding to the position of the stomach; the epigastric region. See *Abdomen*.

Epigea, or Epigæa (*ep-e-je'-ah*) [*ἐπί*, upon; *γῆα*, earth]. A genus of trailing ericaceous plants. **E. repens**, trailing arbutus of N. America; it has diuretic properties. See *Trailing Arbutus*.

Epiglottic (*ep-e-glōt'-ik*) [*ἐπί*, upon; *γλωττίς*, glottis]. Relating to the epiglottis.

Epiglottis (*ep-e-glōt'-is*) [*ἐπί*, upon; *γλωττίς*, glottis]. A fibrocartilaginous structure that aids in preventing food and drink from passing into the larynx.

Epignathus (*ep-i-g'-na-thus*) [*ἐπί*, upon; *γνάθος*, jaw]. A monstrosity in which the rudimentary organs of a twin are united to the superior maxillary bone.

Epihyal Bone (*ep-e-hi'-al*) [*ἐπί*, upon; *ὑαλῆ*]. The stylohyoid ligament when it is ossified.

Epilation (*ep-il-a'-shun*) [*ε*, out of; *pilus*, a hair]. The extraction of hair.

Epilating Forceps. Forceps for plucking out hairs.

Epilepsy (*ep'-il-ep-se*) [*ἐπιληψία*, a laying hold of]. A chronic nervous affection characterized by sudden loss of consciousness with general

tonic and clonic convulsions, the paroxysms lasting but a short time. An epileptic seizure is often preceded by a peculiar sensation, or aura, and as the patient falls he sometimes makes an outcry, the epileptic cry. **E., Cortical, E., Focal, or E., Jacksonian**, spasmodic contractions in certain groups of muscles, with retention of consciousness, due to local disease of the cortex. **E., Masked, E. larvata**. In this, involuntary actions, often violent, replace the convulsion. **E., Nocturnal**, epilepsy in which the attack occurs during sleep. **E., Procrusive**, a form in which the patient runs rapidly forward before falling. **E., Spinal**, paroxysms of clonic spasm in the lower extremities sometimes observed in the course of spastic paraplegia.

Epileptic (*ep-il-ep'-tik*) [*ἐπιληψικός*, a laying hold of]. 1. Pertaining to or like epilepsy. 2. One affected with epilepsy. **E. Aura, E. Cry**. See *Epilepsy*. **E. Dementia**, the dementia which is frequently the terminal stage of epilepsy. **E. Equivalents**, transient psychic disturbances replacing the typical convulsions. **E. Mania**, mania following or taking the place of the fit.

Epileptiform (*ep-il-ep'-tif-orm*) [*ἐπιληψικός*, a laying hold of; *forma*, form]. Resembling an epileptic attack.

Epileptogenous (*ep-il-ep-toj'-en-oi*) [*ἐπιληψικός*, epilepsy; *γενῆναι*, to produce]. Producing epilepsy.

Epileptoid (*ep-il-ep'-toid*) [*ἐπιληψικός*, a laying hold of; *εἶδος*, likeness]. Resembling epilepsy.

Epimysium (*ep-e-mi'-ze-um*) [*ἐπί*, upon; *μῦς*, a muscle]. The sheath of areolar tissue surrounding a muscle.

Epineurium (*ep-e-nu'-re-um*) [*ἐπί*, upon; *νεῖρον*, a nerve]. The connective-tissue sheath of a nerve-trunk.

Epipastic (*ep-e-pas'-tik*) [*ἐπιπάσσειν*, to sprinkle]. Having the qualities of a dusting powder.

Epiphenomenon (*ep-e-fe-nom'-en-on*) [*ἐπί*, on; *φαινόμενον*, phenomenon]. An exceptional sequence or unusual complication arising in the course of a disease.

Epiphora (*ep-i-f'-or-ah*) [*ἐπί*, upon; *φέρειν*, to bear]. A persistent overflow of tears, due to excessive secretion or to impeded outflow through the lacrymal channels.

Epiphyseal, or Epiphysial (*ep-e-fis'-e-al*) [*ἐπί*, upon; *φύειν*, to grow]. Relating to or of the nature of an epiphysis.

Epiphyseitis (*ep-e-fis-e-i'-tis*) [*ἐπιφύσις*, an epiphysis; *ιτις*, inflammation]. Inflammation of an epiphysis.

Epiphysis (*ep-i-f'-is-is*) [*ἐπί*, upon; *φύειν*, to grow]. A process of bone attached for a time to another bone by cartilage, but in most

cases soon becoming consolidated with the principal bone. **E. cerebri**, the pineal gland.

Epiphysitis (*ep-if-is-it-tis*). See *Epiphyseitis*.

Epiphyte (*ep'-e-fit*) [*épi*, upon; *φύτον*, a plant]. A vegetable parasite growing on the exterior of the body.

Epipial (*ep-e-pe'-al*) [*épi*, upon; *pia mater*]. Upon the pia mater.

Epiplocele (*ep-ip'-lo-sel*) [*ἐπιπλοον*, caul; *κηλη*, hernia]. A hernia containing omentum.

Epiploic (*ep-ip'-lo'-ik*) [*ἐπιπλοον*, caul]. Relating or belonging to the omentum. **E.**

Appendages, small pouches of peritoneum filled with fat, found on the colon.

Epiploon (*ep-ip'-lo-on*) [*ἐπιπλεῖν*, to float upon]. The omentum.

Episcleral (*ep-e-skle'-ral*) [*ἐπι*, upon; *σκληρός*, hard]. Situated on the outside of the sclerotic coat.

Episcleritis (*ep-e-skle-rít-is*) [*ἐπι*, upon; *σκληρός*, hard; *ιτις*, inflammation]. An inflammation of the subconjunctival tissues, or of the sclera itself.

Episiorrhaphy (*ep-is-e-or'-a-fe*) [*ἐπισείων*, pubes; *ραφή*, seam]. An operation for the repair of tears about the vulva.

Episiostenosis (*ep-is-e-o-stem-o'-sis*) [*ἐπιστεῖων*, pubes; *στενός*, narrow]. Contraction or narrowing of the vulva.

Episiotomy (*ep-is-e-ol'-o-me*) [*ἐπισείων*, vulva; *τομή*, section]. Incision through the vulva in childbirth, to prevent rupture of the perineum and to facilitate labor.

Epispadias (*ep-e-spád'-e-as*) [*ἐπί*, above; *σπάειν*, to pierce]. A condition in which the urethra opens on the upper part of the penis, either on the dorsum or on the glans.

Epispastic (*ep-e-spás'-tik*) [*ἐπι*, upon; *σπάσις*, a drawing]. 1. Blistering. 2. A substance producing a blister.

Epistaxis (*ep-is-taks'-is*) [*ἐπιστάζειν*, to cause to drop]. Hemorrhage from the nose.

Episternal (*ep-e-stur'-nal*) [*ἐπί*, upon; *στέρνον*, the sternum]. Above the sternum.

Epithelial (*ep-e-the'-le-al*) [*ἐπί*, upon; *θηλή*, a nipple]. Pertaining to or made up of epithelium.

Epithelioid (*ep-e-the'-le-oid*) [*ἐπί*, upon; *θηλή*, nipple; *εἶδος*, likeness]. Resembling epithelium.

Epithelioma (*ep-e-the-le-o'-mah*) [*ἐπί*, upon; *θηλή*, nipple; *μα*, a tumor]. Properly, any tumor in which epithelium forms the prominent element; by usage the word is restricted to carcinoma of the skin and mucous membranes.

Epithelium (*ep-e-the'-le-um*) [*ἐπί*, upon; *θηλή*, nipple]. A term applied to the group of cells that forms the epidermis, that lines all canals having communication with the external air, and that are specialized for secretion

in certain glands, as the liver, kidneys, etc. Epithelium is divided according to the shape and arrangement of the cells into columnar, cuboidal, flat, pavement, squamous, stratified, tessellated, and transitional epithelium; according to function into protective and glandular or secreting. **E.**, **Ciliated**, a form in which the cells bear vibratile filaments or cilia on their free extremities. **E.**, **Transitional**, that intermediate between simple and stratified.

Epitrochlea (*ep-e-trok'-le-ah*) [*ἐπι*, upon; *τροχαλία*, a pulley]. The internal condyle of the humerus.

Epizoon (*ep-e-zo'-on*) [*ἐπι*, upon; *ζῶον*, an animal]. An animal parasite living upon the exterior of the body.

Epizootic (*ep-e-zo-ol'-ik*) [*ἐπι*, upon; *ζῶον*, animal]. An epidemic disease of the lower animals.

Epoophoron (*ep-o-off'-or-on*) [*ἐπι*, upon; *ᾠόν*, egg; *φέρειν*, to bear]. The parovarium.

Epsom Salt (*ep'-sum*). See *Magnesium*.

Epulis (*ep-ul'-lis*) [*ἐπι*, upon; *ὄλσ*, the gums]. A tumor of the alveolar processes of the jaws. **E.**, **Malignant**, a giant-cell sarcoma of the jaw.

Equation (*e-kwa'-shun*) [*æquare*, to make equal]. In chemistry, a collection of symbols so arranged as to indicate the reaction that will take place if the bodies represented by the symbols be brought together. **E.**, **Personal**, an allowance for individual peculiarity or error in an observer's work.

Equator (*e-kwa'-tor*) [*æquare*, to make equal]. An imaginary circle surrounding a sphere so as to divide it into equal halves.

E. of a Cell, the boundary of the plane through which division takes place. **E. of the Eye**, the æquator oculi; a line joining the four extremities of the transverse and vertical axes of the eye.

Equilibrating Operation (*e-kwil'-ib-ra-ting*). An operation on the ocular muscles to equalize their action in cases of squint.

Equilibration (*e-kwil-ib-ra'-shun*) [*æquilibrare*, to balance equally]. The maintenance of equilibrium.

Equilibrium (*e-kwil'-ib'-re-um*) [*æquus*, equal; *libra*, balance]. A state of balance.

Equinia (*e-kwin'-e-ah*) [*equus*, a horse]. Glanders; farcy.

Equivalence, Equivalency (*e-kwiv'-al-ens, e-kwiv'-al-en-se*) [*æquus*, equal; *valere*, to be worth]. The property possessed by an element or radicle of replacing another element or radicle in a compound body in definite proportions.

Equivalent (*e-kwiv'-al-ent*) [*æquus*, equal; *valere*, to be worth]. Of equal valency; having the same value.

Erasion (*e-ra'-shun*) [*e*, out; *radere*, to scrape]. The act of scraping.

Erb's Palsy. A paralysis involving the deltoid, biceps, brachialis anticus, and supinator longus; often also the supinator brevis, and at times the infraspinatus; rarely the subscapularis. It is traumatic in origin; it may occur during birth.

Erectile (*e-rek'-til*) [*erigere*, to set up]. Having the quality of becoming erect. **E. Tissue**, a tissue consisting of a network of expansile capillaries that under stimulus become engorged with blood and cause erection of the part.

Erection (*e-rek'-shun*) [*erigere*, to set up]. The state of being erect, as erection of the penis or clitoris.

Erector (*e-rek'-tor*) [*erigere*, to erect]. A muscle that produces erection of a part. See *Muscles*, *Table of*. **E. pili**, the unstriped muscular fibers causing the erection of the hair and the phenomenon called goose-flesh or goose-skin.

Eremacausis (*er-e-mak'-aw'-sis*) [*ἔρεμα*, slowly; *καίω*, burning]. Slow oxidation or gradual decay, without combustion.

Erethism, Erethismus (*er-e-thizm, er-e-thiz'-mus*) [*ἐρεθισμός*, irritation]. An abnormal increase of nervous irritability.

Erethismic or Erethistic (*er-e-thiz'-mik, er-e-thiz'-tik*) [*ἐρεθιστικός*, irritation]. Relating to, or affected with, erethism.

Erg [*ἐργον*, work]. A unit of work, representing the work done in moving a body against the force of one dyne through a space of one centimeter.

Ergot, or Ergota (*ur'-got, or ur-got'-tah*) [*Fr., ergot*, a spur]. The sclerotium of the *Claviceps purpurea*, a fungus growing on rye. It is a vasomotor stimulant and causes contraction of the involuntary muscles. It is used to control hemorrhage, and to cause uterine contraction; it is also employed in cerebral and spinal congestion, in diabetes insipidus, and in night-sweats. Dose gr. x- $\frac{3}{4}$ (0.65-4.0). **E., Ext., Fld.** Dose $\frac{1}{3}$ ss-f $\frac{3}{4}$ ss (2.0-16.0). **E., Ext.,—Ergotin.** Dose gr. ij-xx (0.13-1.3), hypodermically gr. $\frac{1}{4}$ -v (0.016-0.32). **E., Infus.** (B. P.). Dose f $\frac{3}{4}$ j-ij (32.0-64.0). **Injectio ergotinæ hypodermica** (B. P.), ergotin 1, camphor-water 2. Dose subcutaneously m ij-x (0.2-0.65). **E., Tinctura** (B. P.). Dose m x-f $\frac{3}{4}$ j (0.65-4.0).

Ergotin (*ur'-go-tin*). See *Ergot*.

Ergotinine (*ur'-got'-in-in*) [*Fr., ergot*, a spur], $C_{25}H_{40}N_2O_6$. An alkaloid from ergot of rye.

Ergotism (*ur'-got-izm*) [*Fr., ergot*, a spur]. The constitutional effects following the prolonged use of ergot, or of grain containing the fungus, *claviceps purpurea*. The symptoms are of two types, either a spasmodic form with contractions and cramps of the muscles, or a form characterized by dry gangrene.

Erigeron (*er-ij'-er-on*) [*ἔριγινον*, groundsel]. Fleabane. The plant *E. canadense*, having physiologic actions like those of oil of turpentine, but less irritant. It contains oil of erigeron, and is used as a hemostatic. **E. bellidifolium** and **E. philadelphicum** afford similar oils, and have the same properties. **Erigerontis, Oleum.** Dose m x-f $\frac{3}{4}$ ss (0.65-2.0). **Unof.** **Eriodictyon** (*er-e-o-dik'-te-on*) [*ἔριον*, wool; *δικτυον*, a net]. Verba Santa or mountain-balm. The leaves of *E. glutinosum*, a shrub of California, an expectorant, and an excipient for quinin, the taste of which it largely conceals. **E., Ext., Fld.** Dose m xv-f $\frac{3}{4}$ j (1.0-4.0) **E., Ext.** Dose gr. ij-x (0.13-0.65).

Erosion (*e-ro'-shun*) [*erodere*, to eat out]. The eating away of tissue.

Erotic (*er-ot'-ik*) [*ἔρως*, love]. Pertaining to the sexual passion.

Erotomania (*er-of-o-ma'-ne-ah*) [*ἔρως*, love; *μανία*, madness]. Morbid exaggeration of the affections, usually toward the opposite sex.

Errhine (*er'-in*) [*iv, in*; *ῥίς*, the nose]. 1. Causing discharges from the nose. 2. A medicine that increases nasal secretions; a sternutatory.

Eruclation (*e-ruk'-ta'-shun*) [*eructare*, to belch]. Belching.

Eruption (*e-rup'-shun*) [*erumpere*, to burst out]. A bursting forth, especially applied to the skin-lesions of the exanthematous diseases.

Eruptive (*e-rup'-tiv*) [*erumpere*, to burst out]. Attended by an eruption, as an eruptive fever.

Erysipelas (*er-is-ip'-el-as*) [*ἐρυθρός*, red; *πέλλα*, skin]. An acute infectious disease, due to the streptococcus-erysipelas (which is probably identical with the streptococcus pyogenes), and characterized by an inflammation of the skin and subcutaneous tissues. **E., ambulans.** See *E., Wandering*. **E., Facial**, erysipelas of the face, the most common form. After an initial chill the temperature rises very high; there may be vomiting and delirium, and the disease may rapidly spread over a great part of the body. The affected area is swollen, has a deep-red color, an elevated margin, and itches. **E., Idiopathic**, erysipelas occurring without any visible wound. **E. migrans.** See *E., Wandering*. **E., Phlegmonous**, a form of erysipelas in which there is pus-formation. **E., Surgical or Traumatic**, erysipelas occurring in the site of a wound. **E., Wandering**, a form in which the erysipelas process successively disappears from one part of the body to appear subsequently at another part.

Erysipelatous (*er-is-ip-el'-at-us*) [*ἐρυθρός*, red; *πέλλα*, skin]. Of the nature of or affected with erysipelas.

Erythema (*er-ith-e'-mah*) [*ἐρυθμαίνω*, to make red]. A redness of the skin occurring in patches of variable size and shape. **E. annulare**, a form of *E. multiforme*, in which the lesions shrink and desquamate at the center, but continue to extend at the periphery by a raised margin. **E. intertrigo**, intertrigo; a hyperemia of the skin occurring where the folds of the integument come in contact. The epidermis may be abraded. **E. multiforme**, an acute inflammatory skin-disease, characterized by reddish macules, papules, or tubercles, usually appearing on the legs and forearms. It is often ushered in by gastric distress and rheumatic pains. **E. nodosum**, dermatitis contusiformis, an inflammatory disease characterized by the formation, especially on the tibial surfaces, of rounded, elevated, erythematous nodules. **E., Symptomatic**, a hyperemia of the skin either diffuse or in nonelevated patches. It is either idiopathic, as when arising from the action of the sun, **E. solare**, or due to various poisons, **E. venenatum**, or it is symptomatic of systemic disease or gastrointestinal disorder.

Erythematous (*er-ith-em'-at-us*) [*ἐρυθμαίνω*, to make red]. Of the nature of erythema.

Erythrea (*er-ith-red'-ah*) [*ἐρυθρός*, red]. A genus of plants of the *Gentianaceae*. *E. centaureum*, is the European centaur.

Erythremelalgia (*er-ith-rem-el-al'-je-ah*). See *Erythromelalgia*.

Erythroblast (*er-ith'-ro-blast*) [*ἐρυθρός*, red; *βλαστός*, germ]. A rudimentary red blood-corpuscle.

Erythrocyte (*er-ith'-ro-sit*) [*ἐρυθρός*, red; *κύτος*, cell]. A red blood-corpuscle.

Erythrodextrin (*er-ith-ro-deks'-trin*) [*ἐρυθρός*, red; *dexter*, right]. A dextrin formed by the action of saliva on starch. It yields a red color with iodine.

Erythrogranulose (*er-ith-ro-gran'-u-las*) [*ἐρυθρός*, red; *granulum*, a little grain]. A granular substance, found in starch-grains, coloring red with iodine.

Erythromelalgia (*er-ith-ro-mel-al'-je-ah*) [*ἐρυθρός*, red; *μέλος*, limb; *ἄλγος*, pain]. An affection of the distal parts of the extremities, particularly the feet, characterized by redness and neuralgic pain. The disease is very obstinate; its pathology is not well understood. It may be a vasomotor neurosis, a neuritis of the peripheral nerves, or it may be due to changes in the spinal cord.

Erythrophlein (*er-ith-rof'-le-in*) [*ἐρυθρός*, red; *φλοιός*, bark]. A poisonous alkaloid from *Casca bark*.

Erythrophleum (*er-ith-rof'-le-um*). *Casca Bark*.

Erythroptasia (*er-ith-rof'-se-ah*) [*ἐρυθρός*, red; *ὄψις*, vision]. An abnormality of vision in which all objects appear red; red vision.

Erythroxylin (*er-ith-roks'-il-in*) [*ἐρυθρός*, red; *ξύλον*, wood]. Synonym of cocaine.

Erythroxyton (*er-ith-roks'-il-on*) [*ἐρυθρός*, red; *ξύλον*, wood; *γεν.*, *Erythroxyli*]. *Coca*, *Cuca*. The leaves of *E. coca*, a shrub indigenous to the Andes. It contains an alkaloid, cocaine, $C_{17}H_{21}NO_4$, to which its properties are mainly due. It is an aromatic tonic and cerebral stimulant. **E. coca** (B. P.). Dose gr. ij-xv (0.13-1.0). **Cocæ**, **Ext.**, **Fld.** Dose ℥xx-℥j (1.3-4.0). **Extractum cocæ liquid.** (B. P.). Dose ℥xx-℥j (1.3-4.0). See *Coca*.

Eschar (*es'-kar*) [*ἑσchara*, a scab]. A slough, especially that produced by the thermocautery. **E., Neuropathic**. A bedsore.

Escharotic (*es-kar-et'-ik*) [*ἑσcharωτικός*]. 1. Caustic; producing a slough. 2. A substance that produces an eschar; a caustic.

Eserin (*es'-er-in*) [*Esere*, native name of the plant or bean]. See *Physostigma*.

Esmarch's Bandage, or **Apparatus**. An elastic rubber bandage used upon a limb to be amputated, in order to drive the blood out of it by the pressure of progressive turns about the limb toward the trunk. **E.'s Tubes**, tubes on the sides of which agar or gelatin has been solidified in a thin layer, by rapid turning of the tube on ice or under ice-water.

Esodic (*es-sod'-ik*) [*ἑς*, into; *ὁδός*, way]. Affluent.

Esophageal (*e-sof-aj'-e-al*) [*οἰσophageος*, esophagus]. Pertaining or belonging to the esophagus.

Esophagismus (*e-sof-aj'-is'-mus*) [*οἰσophageος*, esophagus]. Spasmodic contraction of the esophagus.

Esophagitis (*e-sof-aj'-i'-tis*) [*οἰσophageος*, esophagus; *ῖτις*, inflammation]. Inflammation of the esophagus.

Esophagocèle (*e-sof'-ag-o-sel*) [*οἰσophageος*, esophagus; *κῆλη*, hernia]. An abnormal distention of a portion of the esophagus.

Esophagomycosis (*e-sof-ag-o-mi'-ko'-sis*) [*οἰσophageος*, esophagus; *μύκης*, a fungus]. Disease of the esophagus caused by fungi.

Esophagoscope (*e-sof-ag'-o-skōp*) [*οἰσophageος*, esophagus; *σκοπεῖν*, to view]. An instrument for examining the interior of the esophagus by artificial light.

Esophagospasm (*e-sof'-ag-o-spazm*). See *Esophagismus*.

Esophagostenosis (*e-sof-ag-o-sten'-o'-sis*) [*οἰσophageος*, esophagus; *στενωσις*, constriction]. Constriction of the esophagus.

Esophagostomy (*e-sof-ag-o'-stō-me*) [*οἰσophageος*, esophagus; *στόμα*, mouth]. The formation of an artificial opening in the esophagus.

Esophagotomy (*e-sof-ag-o'-o-me*) [*οἰσophageος*, esophagus; *τομή*, a cutting]. Opening of the esophagus by an incision.

Esophagus (*e-sof'-ag-us*) [*φάρυγξ, οἶστρον*, to carry; *φαγεῖν*, to eat]. The gullet, a musculomembranous canal, about nine inches in length, extending from the pharynx to the stomach.

Esophoria (*es-o-fo'-re-ah*) [*ἴσθω*, inward; *φορεῖν*, to bear]. See *Heterophoria*.

Esotropia (*e-so-tro'-pe-ah*) [*ἴσθω*, inward; *τροπεῖν*, to turn]. Convergent strabismus.

Essence (*es'-ens*) [*essentia*, essence]. 1. That which gives to anything its character or peculiar quality. 2. The peculiar qualities of a drug extracted and reduced to a small compass. 3. A solution of an essential oil in alcohol.

Essential (*es-en'-shal*) [*essentia*, essence].

1. Pertaining to the essence of a substance. 2. Of diseases, occurring without a known cause. **E. Oils**, the volatile oils, obtained from aromatic plants by distillation or fermentation.

Esthesiometer (*es-the-se-om'-e-ter*) [*αἰσθησις*, sensation; *μέτρον*, a measure]. An instrument for measuring tactile sensibility.

Esthiomene (*es-the-om'-en-e*) [*ἰσθητικὴν*, eating]. *Lupus exedens*.

Estlander's Operation. An excision of portions of one or more ribs for the relief of empyema.

Etat Mamelonné (*et-ah-mah-mel-on-ā*) [Fr.]. A condition of the stomach in chronic gastritis in which there is a projection of small elevations consisting of hyperplastic mucous membrane.

Ethene (*eth'-en*). Same as *Ethylene*. **E. Chlorid**, $C_2H_2Cl_2$. Dutch liquid. An anesthetic resembling chloroform, but less dangerous.

Ether (*e'-ther*) [*æther*; *αιθήρ*, the upper air].

1. The subtle fluid filling space and penetrating all bodies, the medium of transmission of light, heat, electricity, and magnetism. 2. A compound formed hypothetically from H_2O by the substitution of two alcohol-radicles for the H. 3. Diethylidic acid (C_2H_5O), a thin, colorless, volatile, and highly inflammable liquid, known also as ethylic ether or sulphuric ether, as sulphuric acid is used in its manufacture. Its chief use is as an anesthetic, it being less dangerous than chloroform. It is also employed as a cardiac stimulant in sudden heart-failure, and as a carminative. Dose by the mouth \mathfrak{m}_{xxx} - $\mathfrak{f}\mathfrak{ss}$ (2.0-16.0) in ice-water. **E.**, Acetic, has properties like those of ethylic ether. Dose \mathfrak{m}_x - $\mathfrak{f}\mathfrak{z}$ (0.65-4.0). **E. fortior**, contains 94 per cent. of ethylic acid. Dose \mathfrak{m}_x - $\mathfrak{f}\mathfrak{z}$ (0.65-4.0). **E.**, Hydriodic, unof. Dose for inhalation \mathfrak{m}_{xv} (1.0). **E.**, Hydrobromic, unof. Dose \mathfrak{m}_x - \mathfrak{z} (0.65-4.0). **Spiritus ætheris compositus**. See *Hoffmann's Anodyne*.

Ethereal (*e-the'-re-al*) [*æther*, the upper air].

1. Pertaining to the ether. 2. Made of ether, as *E. tinctures*. 3. Volatile.

Etherization (*e'-ther-i-z'-a'-shun*) [*æther*, ether]. The administration of ether to produce anesthesia. This is effected by inhalation of the vapor.

Etherize (*e'-ther-ize*) [*æther*, ether]. To administer ether.

Ethidene (*eth'-id-en*) [*æther*, ether], C_2H_2 . Ethylidene; a bivalent radicle. **E. Chlorid**, or *Dichlorid*, a colorless fluid, tasting and smelling like chloroform. It has been used as a general anesthetic. See *Anesthetic*.

Ethmocephalus (*eth-mo-sef'-al-us*) [*ἔθμος*, ethmoid; *κεφαλή*, head]. A variety of single autositic monsters in which there is a rudimentary nose in the shape of a proboscis terminating anteriorly in two imperfect nostrils or in a single opening.

Ethmoid (*eth'-moid*) [*ἔθμος*, a sieve; *εἶδος*, likeness]. The sieve-like bone of the nose, perforated for the transmission of the olfactory nerve; it forms a part of the base of the skull.

Ethmoid or Ethmoidal (*eth'-moid*, *eth-moi'-dal*) [*ἔθμος*, sieve; *εἶδος*, likeness]. Relating to the ethmoid bone.

Ethnology (*eth-nol'-o-jē*) [*ἔθνος*, a nation; *λόγος*, science]. The comparative study of the races of mankind.

Ethoxycaffein (*eth-oks-e-kaf'-e-in*), $C_{10}H_{14}N_4O_3$. A remedy recommended in herpes zoster and migraine. Dose gr. iv (0.26). Unof.

Ethyl (*eth'-il*) [*αιθήρ*, ether; *ἔλγν*, material]. The alcohol-radicle, C_2H_5 . **E. Alcohol**, ordinary alcohol of the pharmacopeia. See *Alcohol*. **E. Bromid**, C_2H_5Br , Bromethyl; monobromethane. A rapid and transient anesthetic. See *Anesthetic*. **E. Bromid**, *Liq.*, 1 in 200, used in angina pectoris. Dose $\mathfrak{f}\mathfrak{ss}$ - \mathfrak{ij} (16.0-64.0). Unof. **E. Carbamate**, Urethane. **E. Chlorid**, C_2H_5Cl , an anesthetic resembling chloroform in action. **E. Iodid**, C_2H_5I , hydriodic ether, used to relieve the dyspnea of bronchitic asthma and edematous laryngitis. Dose to be inhaled, \mathfrak{m}_v (0.32), three or four times daily. Unof. **E. Oxid**. See *Ether*.

Ethylate (*eth'-il-at*). A compound of ethylic alcohol in which the H of the hydroxyl is replaced by a base.

Ethyl-chloral-urethane. See *Somnal*.

Ethylene (*eth'-il-en*) [*αιθήρ*, ether; *ἔλγν*, matter]. Olefiant gas, C_2H_4 . A colorless, poisonous gas, which burns with a bright, luminous flame, and when mixed with air explodes violently. It is one of the constituents of illuminating gas. **E. Bromid**, a light, brownish-colored liquid with the formula, $C_2H_4Br_2$. It has been used in epilepsy. Dose

$m\frac{3}{4}-2$ (0.05-0.13). Unof. **E. Chlorid**, **E. Bichlorid**. See *Ethene Chlorid*. **E.-diamin**, a non-poisonous base isomeric with ethylidene-diamin.

Ethylene-imid (*eth'-il-en-im'-id*), C_2H_5N . A nonpoisonous base found in cholera-cultures.

Ethylidene (*eth-il'-id-en*). See *Ethidene*.

Ethylidene-diamin (*eth-il'-id-en di'-a-min*) $C_2H_4(NH_2)_2$. A poisonous ptomain obtained from decomposing haddock. Injections into mice and guinea-pigs produce hypersecretion from mouth, nose, and eyes, mydriasis, exophthalmos, great dyspnea, and death.

Ethyl-phenyl-carbamate, **Ethyl-phenyl-urethane**. See *Euphorin*.

Etiologic (*e-te-ol-og'-ik*) [*aitia*, a cause; *lógos*, science]. Pertaining to etiology.

Etiology (*e-te-ol'-o-je*) [*aitia*, a cause; *lógos*, science]. The causation of disease.

Eucalyptol (*u-kal-ipt'-tol*) [*ev*, well; *καλύπτειν*, to cover], $C_{10}H_{18}O$. A neutral principle obtained from the volatile oil of *Eucalyptus globulus*, and of some other species of *Eucalyptus*. It is used in bronchitis and malaria, and also in ear-diseases and in urethritis, and externally in various liniments and washes. Dose $m\frac{v}{x}$ (0.32-0.65), in capsules, three times daily.

Eucalyptus (*u-kal-ipt'-tus*) [*ev*, well; *καλύπτειν*, to cover]. The leaves of *E. globulus*, native to Australia, but now cultivated in California. It contains a volatile oil, from which eucalyptol is obtained. The properties largely depend on the volatile oil. *Eucalyptus* has been used as an antiseptic, as a stimulant to mucous membranes, as an antispasmodic in asthma, in migraine, and, with doubtful success, in malaria. **E.**, **Oleum**, the volatile oil. Dose $m\frac{v}{x}$ (0.32) in capsules or emulsion. **E.**, **Unguentum** (B. P.), contains 20 per cent. of *E. Oi*.

Euchlorhydria (*u-klor-hy'-dre-a*) [*ev*, well; *χλωρός*, green; *ἵδωρ*, water]. The presence of a normal amount of hydrochloric acid in the gastric juice.

Eugenia (*u-je'-ne-ah*) [after Prince *Eugene*, of Savoy]. A genus of trees and shrubs, mostly tropical, among which are *E. caryophyllata*, which yields caryophyllus, and *E. pimenta*, which produces pimenta.

Eugenic Acid (*u-jeu'-ik*). See *Eugenol*.

Eugenol (*u'-jen-ol*) [See *Eugenia*]. $C_{10}H_{12}O_2$. Eugenic Acid; a phenol-like compound that occurs in clove-oil and in allspice, and is convertible into vanillin. It is used as an antiseptic, and as a local anesthetic in dentistry. Dose gr. xv (1.0) well diluted. Unof.

Eunuch (*u'-nuk*) [*ευνούχος*, guardian of the couch]. A male whose genital organs have been removed or mutilated so as to render him impotent.

Euonymin (*u-on'-im-in*). A precipitate from the tincture of wahoo; it is tonic, laxative, and expectorant. Dose $\frac{1}{2}$ to 3 grains (0.032-0.2). Unof.

Euonymus (*u-on'-im-us*) [*εὐόνομος*, having a good name]. Wahoo, the bark of *E. atropurpureus*, a mild purgative and cholagogue. **E.**, **Ext.** Dose gr. j-v (0.065-0.32).

Eupatorium (*u-pat'-o'-re-um*) [*εὐπατόριον*, agrimony]. 1. A genus of composite-flowered plants. The leaves and flowering-tops of *E. perfoliatum*, thoroughwort, or boneset. It is a bitter tonic, diaphoretic, and feeble emetic. Dose of the powder gr. xx-xxx (1.3-2.0). **E.**, **Ext.**, **Fld.** Dose $m\frac{x}{x}$ -f3j (0.65-4.0).

Euphorbia (*u-for'-be-ah*). A genus of trees, shrubs, and herbs, yielding a milky juice. **E. resinifera**, of Africa, affords euphorbium. **E. corollata**, **E. ipecacuanha**, American species, have been employed in medicine on account of their emetic, diaphoretic, and expectorant properties. **E. pilulifera**, of S. America and Australia, is used in asthma and bronchitis. Dose of the extract gr. j (0.065); of the fluid extract f3 ss-j (2.0-4.0); of the tincture f3 ss-j (2.0-4.0).

Euphorbium (*u-for'-be-um*). An acrid gum-resin obtained from *Euphorbia resinifera*. It is strongly purgative and vesicant, and is now mainly employed in veterinary medicine. **Euphoria** (*u-for'-e-ah*) [*εὐφορος*, easily carried]. The sense of well-being; health.

Euphorin (*u'-for-in*) [*εὐφορος*, easily carried], $C_{10}H_{14}$. Phenyl-urethan, a white crystalline powder derived from anilin. It is recommended as an analgesic and antipyretic in neuralgia and rheumatism. Dose gr. v-xxx (0.32-2.0) daily.

Euplastic (*u-plas'-tik*) [*ev*, well; *πλασσειν*, to form]. Capable of being transformed into healthy tissue.

Eupnea (*ūp-ne'-ah*) [*ev*, well; *πνεειν*, to breathe]. Normal or easy respiration.

Euophen (*u'-ro-fen*). Diisobutylorthocresol iodid. An amorphous yellow powder recommended as a substitute for iodoform. It is used hypodermatically in doses of gr. $\frac{1}{4}$ -iss in syphilis.

Eustachian (*u-stal'-ke-an*) [*Eustachio*, an Italian anatomist]. **E. Catheter**, a catheter for introduction into the Eustachian tube; it is introduced along the floor of the nose. **E. Tube**, the osseocartilaginous canal extending from the tympanum to the pharynx. **E. Valve**, a prolongation of the lining membrane of the inferior vena cava into the right auricle of the fetal heart.

Euthanasia (*u-than-a'-se-ah*) [*ev*, well; *θάνατος*, death]. An easy or calm death.

Eutocia (*u-to'-ke-ah*) [*ev*, well; *τόκος*, childbirth]. Natural or easy childbirth.

Evacuant (*e-vak'-u-ant*) [*evacuare*, to empty]. 1. Emptying. 2. A medicine that causes the emptying of an organ, especially the bowels; a purgative.

Evacuation (*e-vak-u-a'-shun*) [*evacuare*, to empty]. 1. The act of emptying, especially of the bowels. 2. That which is evacuated.

Evacuator (*e-vak'-u-a-tor*) [*evacuare*, to empty]. An agent to produce emptying, especially an instrument for removing from the bladder fragments of stone after litholapaxy.

Evaporation (*e-vap-or-a'-shun*) [*e* priv.; *vapor*, vapor]. The conversion of a liquid into vapor.

Eventration (*e-ven-tra'-shun*) [*e*, out of; *venter*, the belly]. Protrusion of the abdominal viscera through the abdominal walls.

Eversion (*e-ver'-shun*) [*eversio*, a turning out]. A turning outward.

Evisceration (*e-vis-er-a'-shun*) [*e*, out; *viscera*, the bowels]. The removal of the viscera. **E**, of the Eye, removal of the entire contents of the globe of the eye, leaving the sclerotic intact. **E**, Obstetric, the removal

of the abdominal or thoracic viscera of the fetus to facilitate delivery.

Evolution (*ev-o-lu'-shun*) [*evolvere*, to unroll]. The process of unfolding or developing from a simple to a complex, specialized, perfect form. **E.**, Spontaneous, a series of changes whereby a shoulder-presentation is transformed within the pelvis into a combined breech-and-shoulder-presentation and delivery effected without artificial aid.

Evulsion (*e-vul'-shun*) [*evellere*, to pluck out]. The forcible tearing or plucking away of a part.

Exacerbation (*eks-as-er-ba'-shun*) [*exacerbare*, to be violent]. An increase in the symptoms of a disease.

Exalgin (*eks-al'-jin*) [*ēξ*, out; *ἀλγος*, pain], $C_9H_{11}NO$. Methylacetanilid; a benzene derivative allied to phenacetin. It is an analgesic and antipyretic in doses of from one-half to four grains (0.032-0.26). Unof.

Exanthema, or **Exanthem** (*eks-an'-them-ah*, *eks-an'-them*) [*ἐξάνθημα*, eruption]. 1. An eruption upon the skin. 2. Any exanthematous or eruptive fever.

TABLE OF EXANTHEMATA.

NAME.	PERIOD OF INCUBATION.	STAGE OF INVASION.	TIME OF APPEARANCE OF ERUPTION.	CHARACTER OF ERUPTION.	DURATION OF ERUPTION.	LOCATION.	DESQUAMATION.	DURATION OF DISEASE.	CONVALESCENCE.
Cerebro-spinal Meningitis.	Unknown.	Sudden.	2d to 4th day.	Herpes labialis, purpuric spots, dusky erythema.		Herpes on lips; purpuric spots over entire body.		Variable; many sequelae.	Lysis.
Erysipelas.	Few hours to 3 or 4 days.	1 to 3 days.	Within 24 hours.	Bright-red, shining patches, with well-defined raised margin.	4 to 8 days.	Begins usually on face; may spread to trunk and arms. In the traumatic form begins at the wound.	Branny or in large flakes.	1 to 3 weeks.	Crisis.
Measles.	10 to 12 days.	4 days.	4th day.	Small, dark-red maculae arranged in crescentic form. Complete in 24 hours.	4 to 5 days.	Face; then downward over body.	Branny. 8-11 days.	2 weeks.	Crisis.
Rötheln.	8 to 17 days.	24 to 48 hours.	Within 48 hours.	Rose-colored, rounded, discrete maculae.	3 days.	Face and scalp; then downward over body.	Slightly branny.	4 to 7 days.	Crisis.
Scarlet Fever.	1 to 21 days.	1 to 2 days.	Within 24 hours.	Diffuse, scarlet, punctate.	7 to 10 days.	Neck, chest, face; then over body.	Scales or large flakes. About one week.	3 to 3 weeks.	Lysis.

TABLE OF EXANTHEMATATA.—Continued.

NAME.	PERIOD OF INCUBATION.	STAGE OF INVASION.	TIME OF APPEARANCE OF ERUPTION.	CHARACTER OF ERUPTION.	DURATION OF ERUPTION.	LOCATION.	DESQAMATION.	DURATION OF DISEASE.	CONVALESCENCE.
Typhoid Fever.	5 to 35 days.	6 to 8 days.	7th day.	Rose-colored, lenticular spots, coming on in successive crops.	Each crop 3 to 5 days. Lasts 10 to 20 days or throughout the whole course of the fever.	Abdomen, chest, and back.	Slightly branny or none.	3 to 4 weeks.	Lysis.
Typhus Fever.	4 to 12 days.	5 days.	Usually 5th day; may be on 3d or not till 7th day.	Dusky spots or papules, or petechiæ.	Few days, or may last throughout the course of the disease.	Sides of chest and abdomen, arms, back.	Slightly branny.	2 to 4 weeks.	Crisis.
Varicella.	4 to 14 days.	1 to 2 days.	Within 12 to 24 hours.	Vesicles appearing in crops.	5 to 8 days.	Back, chest, arms.	Crusts. 5-8 days.	2 weeks.	Lysis.
Variola.	8 to 14 days.	3 days.	4th day.	First, shot-like papules, then vesicles, then umbilicated pustules.	21 to 25 days.	Face, and over body.	Crusts. 12-22 days.	4 to 5 weeks.	Lysis.

Exanthematous (*eks-an-them'-at-us*) [*ἐξάνθημα*, eruption]. Of the nature of or characterized by exanthema, or eruption; of the nature of an eruptive fever.

Exarticulation (*eks-ar-tik-u-la'-shun*) [*ex*, out; *articulus*, joint]. 1. Dislocation of a joint. 2. Amputation at a joint.

Excavation (*ex-kav-a'-shun*) [*excavare*, to hollow out]. A hollow or cavity. E. of the Optic Nerve, a hollowing or "cupping" of the optic disc, that may be physiologic, congenital, or pathologic, the result of glaucoma, optic atrophy, etc.

Excentric (*eks-sen'-trik*) [*ex*, out; *centrum*, center]. See *Eccentric*.

Excipient (*ek-sip'-e-ent*) [*excipere*, to take up]. Any substance combined with an active drug to give the latter an agreeable or convenient form.

Excision (*ek-sish'-un*) [*excisio*, a cutting out]. The cutting out of a part.

Excitability (*ek-si-ta-bil'-it-e*) [*excitare*, to rouse]. The property of reacting to a stimulus.

Excitant (*ek-si'-tant*) [*excitare*, to rouse]. 1. Stimulating. 2. A remedy that stimulates the activity of an organ.

Excitation (*ek-si-ta'-shun*) [*excitare*, to rouse]. The act of stimulating or irritating. E., Direct, the stimulation of a muscle by placing an electrode on the muscle itself. E.,

Indirect, the stimulation of a muscle through its nerve.

Exciting (*ek-si'-ting*) [*excitare*, to rouse]. Calling forth directly, as an exciting cause.

Excitomotor (*ek-si-to-mo'-tor*). Exciting or arousing motor function; also, a drug or agent that increases the activity of the motor nerve centers.

Exclusion (*eks-klu'-shun*) [*excludere*, to shut out]. A shutting out. E., **Diagnosis by**, the reaching of a diagnosis by excluding one hypothesis after another till only one remains.

Excoriation (*eks-ko-re-a'-shun*) [*ex*, from; *corium*, the skin]. Abrasion of a portion of the skin.

Excrement (*eks'-kre-ment*) [*excernere*, to separate]. An excreted substance; the feces.

Excrementitious (*eks-kre-men-tish'-us*) [*excernere*, to separate; to excrete]. Pertaining to excrement.

Excrecence (*eks-kres'-ens*) [*excrecere*, to grow out]. An abnormal outgrowth upon the body.

Excreta (*eks-kre'-tah*) [*excernere*, to separate]. The natural discharges of the body, particularly those of the bowels.

Excrete (*eks-kre't*) [*excernere*, to separate]. To remove from the body useless substances.

Excretin (*eks'-kre-tin*), $C_{20}H_{26}O$. A crystalline substance found in feces.

Excretion (*eks-kre'-shun*) [*excernere*, to excrete]. The discharge of waste-products.

Excretory (*eks'-kre-to-re*) [*excernere*, to excrete]. Pertaining to excretion.

Excursion (*eks-kur'-shun*) [*ex*, out of; *currere*, to run]. A wandering from the usual course.

Exencephalus (*eks-en-sef'-al-us*) [*ēξ*, out; *ἐνκεφαλος*, brain]. A species of monsters characterized by a malformed brain, situated without the cranial cavity.

Exenteration (*eks-en-ter-a'-shun*) [*ēξ*, out; *ἐντερον*, intestine]. Removal of the intestines of the fetus, to allow delivery.

Exfoliation (*eks-fo-le-a'-shun*) [*exfoliare*, to shed leaves]. The separation of bone or other tissue in thin layers.

Exhalation (*ex-hal-a'-shun*) [*exhalare*, to breathe out]. The giving off of matters in the form of vapor.

Exhaustion (*eg-sawst'-yun*) [*exhaurire*, to pour out]. Loss of vital and nervous power from fatigue or protracted disease.

Exhibit (*ek-zib'-it*) [*exhibere*, to give]. To administer, as a medicine.

Exhumation (*ex-hu-ma'-shun*) [*ex*, out of; *humus*, the ground]. The removal of a corpse from the ground.

Exocardiac, or Exocardial (*eks-o-kar'-de-ak*, or *eks-o-kar'-de-al*) [*ēξω*, out; *καρδια*, heart]. Originating or situated outside of the heart.

Exodic (*eks-od'-ik*) [*ēξω*, out; *ὁδός*, a way]. Transmitting; efferent.

Exogenetic (*eks-o-jen-et'-ik*) [*ēξω*, outward; *γενᾶν*, to produce]. Due to an external cause; not arising within the organism.

Exogenous (*eks-oj'-en-us*) [*ēξω*, out; *γενᾶν*, to produce]. Growing by accretions to the outer surface.

Exomphalos (*eks-om'-fal-as*) [*ēξ*, out; *ὀμφαλός*, navel]. Undue prominence of the navel; also, umbilical hernia.

Exophoria (*eks-o-fo'-re-ah*). See *Heterophoria*.

Exophthalmic (*eks-off-thal'-mik*) [*ēξ*, out; *ὀφθαλμός*, eye]. Pertaining to exophthalmos. **E. Goiter.** See *Goiter*.

Exophthalmos, Exophthalmus (*eks-off-thal'-mos, -mus*) [*ēξ*, out; *ὀφθαλμός*, eye]. Abnormal prominence of the eyeballs.

Exoskeleton (*eks-o-skel'-et-un*) [*ēξω*, outside; *σκελετόν*, a dried body]. The rigid outer envelop of many of the lower forms of life for the protection and attachment of organs.

Exosmosis (*eks-os-mo'-sis*) [*ēξ*, out; *ὠμός*, thrust]. Outward osmosis. See *Osmosis*.

Exostosis (*eks-os-to'-sis*) [*ēξ*, out; *ὀστέον*, bone]. A bony outgrowth from the surface of a bone.

Exotropia (*eks-o-tro'-pe-ah*). See *Strabismus*.

Expectant (*eks-pek'-tant*) [*expectare*, to look out for]. Awaiting or expecting. **E. Treat-**

ment, watching the progress of a disease, and not interfering unless warranted by special symptoms.

Expectation of Life. The average number of years that persons of a given age live.

Expectorant (*eks-pek'-to-rant*) [*ex*, out; *pectus*, breast]. 1. Promoting expectoration. 2. A remedy that promotes or modifies expectoration.

Expectoration (*eks-pek-tor-a'-shun*) [*ex*, out; *pectus*, breast]. 1. The ejection from the mouth of material brought into it from the air-passages. 2. The fluid or semi-fluid matters from the lungs and air-passages expelled by coughing and spitting. **E., Prune-juice**, a sputum containing altered blood expectorated in gangrene and cancer of the lung, and in grave pneumonias in the aged. **E., Rusty.** See *Sputum*.

Expiration (*eks-pi-ra'-shun*) [*expirare*, to breathe out]. The act of breathing forth, or expelling air from the lungs.

Expiratory (*eks-pi'-ra-to-re*) [*expirare*, to breathe out]. Relating to expiration.

Exploration (*eks-plo-ra'-shun*) [*explorare*, to search out]. The act of exploring; investigation of a part hidden from sight by means of touch, by artificial light, etc.

Exploratory (*eks-plor'-at-or-e*) [*explorare*, to search out]. Pertaining to exploration. **E. Puncture**, the puncture of a cavity or tumor and extraction therefrom of some of the contents to learn their nature.

Exploring Needle. A needle with a grooved side to allow the passage of fluid along it after it is plunged into a part where fluid is suspected.

Expression (*eks-fresh'-un*) [*expressus*; *exprimere*, to press out]. A pressing out. **E. of Fetus or Placenta**, assisting the expulsion of fetus or placenta by pressure upon the uterus through the abdominal walls.

Expulsive (*eks-pul'-siv*) [*expellere*, to drive out]. Forcing out.

Exsanguination (*ek-sang-gwin-a'-shun*) [*ex*, out; *sanguis*, blood]. The act of making bloodless.

Exsection (*ek-sek'-shun*) [*ex*, out of; *secare*, to cut]. The act of cutting a part out from its surroundings.

Exsiccation (*ek-sik'-a'-shun*) [*ex*, out; *siccus*, dry]. The act of drying; especially the depriving of a crystalline substance of its water of crystallization.

Exstrophy of the Bladder (*ek'-strof'-e*) [*ēξ*, out of; *στροφήν*, to turn]. A congenital condition in which the lower part of the abdominal wall, the anterior wall of the bladder, and usually the symphysis pubis are wanting, and the posterior wall of the bladder is pressed through the opening.

Extension (*eks-ten'-shun*) [*extendere*, to

stretch out]. A straightening out, especially the muscular movement by which a flexed limb is made straight. **Counterextension**, traction made on a part in a direction opposite to that in which traction is made by another force.

Extensor (*eks-ten'-sor*) [*extendere*, to extend]. That which stretches out or extends, as E. muscles. See *Muscles*, Table of.

External (*ex-tur'-nal*) [*externus*, outward]. On the exterior, or on the side removed from the center or middle line of the body.

Extirpation (*eks-ter-pa'-shun*) [*extirpare*, to root out]. Complete removal of a part.

Extra- (*eks'-trah-*) [L.]. A prefix meaning outside, without. **E. Current**, the induced electric current.

Extraarticular (*eks-trah-ar-tik'-u-lar*). Outside of the proper structures of a joint.

Extracapsular (*eks-trah-cap'-su-lar*). Outside of the capsular ligament of a joint.

Extract, Extractum (*ek'-trakt, -trakt'-um*) [*extrahere*, to extract]. In pharmacy, a solid or semi-solid preparation, made by extracting the soluble principles of a drug with water or alcohol and evaporating the solution. **E., Alcoholic**, that in which alcohol is the solvent. **E., Aqueous**, that prepared by using water as the solvent. **E. ferri pomatum**, N. F.; this is made from iron, in the form of fine, bright wire, 1 part; ripe sour apples 50 parts; water a sufficient quantity. **E., Fluid**, a solution of the solid principles of a vegetable drug, of such strength that 1 gram of the drug is fully represented by one cubic centimeter of the fluid extract.

Extraction (*ex-trak'-shun*) [*extractio*; *ex*, out; *trahere*, to draw]. 1. The act of drawing out. 2. The process of making an extract. **E. of Cataract**, removal of a cataractous lens by surgical operation.

Extractive (*eks-trak'-tiv*) [*ex*, out; *trahere*, to draw]. Any organic substance that may be extracted in small amount from animal tissues.

Extractor (*eks-trak'-tor*) [*ex*, out; *trahere*, to draw]. An instrument for extracting bullets, sequestræ, etc.

Extradural (*eks-trah-du'-ral*) [*extra*, outside; *durus*, hard]. Situated outside of the dura mater.

Extramedullary (*eks-trah-me-dull'-ar-i*) [*extra*, without; *medulla*, marrow]. Situated or occurring outside of the medulla.

Extraneous (*eks-tra'-ne-us*) [*extraneus*, external]. Existing or belonging outside the organism.

Extrapolar (*eks-trah-po'-lar*) [*extra*, without; *polus*, a pole]. Not lying in the space between the electrodes of a battery.

Extrauterine (*eks-trah-yu'-ter-in*) [*extra*, outside; *uterus*]. Outside of the uterus. **E. Pregnancy**. See *Pregnancy*.

Extravasation (*eks-trav-as-a'-shun*) [*extra*, outside; *vas*, a vessel]. 1. The passing of fluid outside of the cavity or space normally containing it. 2. The fluid that has passed out.

Extravascular (*eks-trah-vas'-ku-lar*) [*extra*, outside; *vas*, a vessel]. Outside of the vessels.

Extrinsic (*eks-trin'-sik*) [*extrinsicus*, from without]. External; not directly belonging to a part. **E. Muscles**, those situated on the exterior of an organ.

Extroversion (*eks-tro-ver'-shun*). See *Ex-strophy*.

Exudate (*eks'-u-dat*) [*exudare*, to sweat]. The material that has passed through the walls of vessels into the adjacent tissues.

Exudation (*eks'-u-da'-shun*) [*exudare*, to sweat]. The passing out of serum or pus; the material that has passed out.

Exudative (*eks'-u-da-tiv*) [*exudare*, to exude]. Of the nature of or characterized by exudation.

Eye (*i*) [AS., *cāge*]. The organ of vision. It occupies the anterior part of the orbit, is nearly spheric in outline, and is composed of three concentric coats: the sclerotic and cornea, the choroid and iris, and the retina. The sclerotic is an opaque, dense, white, fibrous membrane, into the anterior part of which the transparent cornea is fitted. The choroid is the vascular tissue, and is continuous with the iris in front. The latter is a circular membrane with a central perforation, the pupil. Within the choroid is the retina, a delicate transparent membrane containing the terminations of the optic nerve. The greater part of the eyeball is filled with a mucoid substance, the vitreous humor, against the anterior surface of which rests the crystalline lens. The space between the lens and the cornea is divided by the iris into two compartments, communicating through the pupillary opening, the anterior and posterior chambers, which contain the aqueous humor. Anteriorly the eye is covered by conjunctiva, posteriorly by a fibrous capsule (capsule of Tenon). The eyeball is moved by a series of muscles attached on the outer surface. Changes in the curvature of the lens are brought about by the ciliary muscle, while the size of the pupil is modified by the action of dilator and constrictor fibers in the iris. **E., Diagrammatic, of Listing**, a scheme for simplifying optic problems by representing the two nodal points and the two principal points of the eye by a mean nodal point and a mean principal point. **E.-brow**, the hair-covered skin on the upper margin of the orbit. **E.-lashes**, the hairs of the eyelid. **E.-lid**, the protective covering of the eyeball, composed of

skin, glands, connective and muscular tissue, the tarsus and conjunctiva, with the cilia at the free edge. **E.-piece**. Synonym of *Ocular*. **E., Reduced, of Donders**, a representation of the eye in such a way that all the

distances needed in the calculations are represented by whole numbers. **E.-speculum**, an instrument for retracting the eyelids. **E.-teeth**, the canine teeth of the upper jaw.

F

F. 1. Abbreviation of Fahrenheit. 2. Symbol for fluorin.

Ft. Abbreviation of *fiat* or *fiant*, let there be made.

Face (*fas*) [*facies*, the face]. A name applied to the lower and anterior part of the head, including the eyes, nose, mouth, cheeks, lips, etc.

Facet (*fas'-et*) [*Fr.*, *facette*, a little face]. A small plane surface, especially on a bone or a hard body, like a calculus.

Facial (*fa'-shal*) [*facies*, the face]. Pertaining to the face. **F. Angle**, an angle measured in different ways by different authorities. That of Virchow and Holder is formed by the union of a line joining the frontonasal suture and the most prominent point of the lower edge of the superior alveolar process, and a line joining the superior border of the external auditory meatus and the lower portion of the orbit. That of Camper is formed by the union of Camper's line (a line touching the most prominent points of the upper and lower face) and a line joining the acantion and the auricular point.

Facies (*fa'-she-tes*) [*L.*, face]. The appearance of the face. **F. hippocratica**, an appearance of the face indicative of the rapid approach of dissolution: the nose is pinched, the temples hollow, the eyes sunken, the ears leaden and cold, the lips relaxed, the skin livid. **F. leontina**. See *Leontiasis*.

Factitious (*fak'-tish'-us*) [*facere*, to make]. Artificial.

Facultative (*fak'-ul-ta-tiv*) [*facultas*, capability]. Voluntary; optional. **F. aerobia**. See *Aerobic*. **F. anaerobia**. See *Anaerobia*.

F. Manifest Hyperopia, that part of the manifest hyperopia that can be concealed by the accommodation. **F. Parasite**, an organism that while usually parasitic, can also live outside of its host.

Fæcal, Fæces (*fe'-kal, fe'-sts*). See *Fecal, feces*.

Fænumgræcum. See *Fenugreek*.

Fahrenheit's Thermometer. See *Thermometer*.

Falciform (*fal'-si-form*) [*falx*, a sickle; *forma*, form]. Having the shape of a sickle.

F. Ligament. See *Ligament*. **F. Process**, a process of the dura mater that separates the hemispheres of the brain; the falx.

Falling (*faul'-ing*) [*AS.*, *feallan*, to fall]. Dropping down. **F.-sickness**, epilepsy.

F. of Womb, a descent of the uterus into the vagina.

Fallopian (*fal'-d'-pe-an*) [*Fallopia*, an Italian anatomist]. Described by Fallopio or Fallopius. **F. Tubes**. See *Oviducts*.

Fallopian, Aqueduct of. See *Aqueduct*.

F., Hiatus of, an opening in the petrous portion of the temporal bone, for the transmission of the petrosal branch of the Vidian nerve.

False (*fauls*) [*falsus*, deceptive]. Not genuine; not real; imitating. **F. Aneurysm**. See *Aneurysm*. **F. Ankylosis**, ankylosis due to rigidity of the soft tissues. **F. Image**, in diplopia, the image of the deviating eye. **F. Membrane**, a fibrinous exudate upon a surface. **F. Pains**, pains that precede true labor-pains. **F. Passage**, a passage formed by the laceration of any canal. **F. Pelvis**, the portion of the pelvic cavity situated above the linea ileopectinea. **F. Ribs**. See *Ribs*.

Falx (*falks*) [*L.*]. A sickle; a sickle-shaped structure. **F. cerebelli**, a sickle-like process of dura mater between the lobes of the cerebellum. **F. cerebri**, the process of the dura, separating the hemispheres of the cerebrum.

Famine Fever. Same as *Relapsing Fever*.

Fang [*AS.*, *fangan*, to seize]. The root of a tooth.

Farad (*far'-ad*) [after Michael Faraday, a physicist]. The unit of electric capacity; a capacity sufficient to hold one coulomb of current having a potential of one volt.

Faradaic, Faradic (*far-ad-a'-ik, far-ad'-ik*) Pertaining to induced electric currents. **F. Current**, the induced electric current.

Faradism (*far'-ad-izm*) [after Michael Faraday, a physicist]. 1. The electricity produced in an induced or faradic current. 2. Faradization.

Faradization (*far-ad-is-a'-shun*). The application of the induced current to a part.

Farcy (*far'-se*) [*farrire*, to stuff]. The form of glanders that attacks the skin and lymphatic glands.

Farfara (*far'-far-ah*). See *Coltsfoot*.

Farina (*far'-e'-nah*) [*farina*, meal]. The ground or powdered starchy part of seeds, especially that of corn, barley, rye, and wheat.

Farinaceous (*far-in-a'-se-us*) [*farina*, meal]. Having the nature of or yielding flour.

Far Point. The most distant point at which an eye can see distinctly when accommodation is completely relaxed.

Far-sightedness. Hypermetropia.

Fascia (*fa'-she-ah*) [*fascia*, a band]. The areolar tissue forming layers beneath the skin (superficial F.), or between muscles (deep F.). F., Anal. See F., *Ischio-rectal*. F., Cervical, Deep, that which invests the muscles of the neck and encloses the vessels and nerves. F., Colles', the deep layer of the superficial fascia of the perineum. F., Cremasteric, a thin covering of the spermatic cord, formed by the stretched fibers of the cremaster muscle. F., Cribriform, the sieve-like covering of the saphenous opening. F. dentata, a serrated band of gray matter of the hippocampal gyrus of the cerebrum. F., Infundibuliform, the process of the transversalis fascia extended over the spermatic cord. F., Intercolumnar, a fascia attached to the margins of the external abdominal ring and forming a sheath for the cord and testis. F., Ischio-rectal, covers the perineal aspect of the levator ani muscle. F. lata, the dense fascia surrounding the muscles of the thigh. F., Transversalis, that lying between the transversalis muscle and the peritoneum.

Fascial (*fa'-she-al*) [*fascialis*, of a fascia]. Pertaining to or of the nature of a fascia.

Fasciculus (*fus-ik'-u-lus*) [dim. of *fascis*, a bundle]. A little bundle, particularly of muscle-fibers. F., Cuneate, the continuation of Burdach's column, or the posteromedian column of the spinal cord. F., Gracile, the posterior pyramid of the medulla. F., Olivary. See *Fillet, Olivary*. F., Postero-longitudinal, fibers connecting the corpora quadrigemina and the nuclei of the fourth and sixth nerves with the parts below. F., Solitary, fibers connecting the internal capsule and the lenticular nucleus with parts below. F. subcallosus (of Muratoff), a tract of long association-fibers lying under the corpus callosum, and connecting the frontal, parietal, and occipital lobes. F., Uncinate, fibers connecting the frontal and the temporo-sphenoidal lobes.

Fastigium (*fas-tij'-e-um*) [L., summit]. The acme of a disease.

Fat [ME., *fat*, fat]. A greasy substance, a compound of oleic, palmitic, or stearic acid with glycerol. F.-cell, a connective-tissue cell containing oil-globules. F.-necrosis, a peculiar form of necrosis of a fatty tissue occurring in pin-point sized areas of a dead-white color.

Fatigue (*fa-tig'*) [*fatigo*, weariness]. Weariness. F. Fever, fever following excessive exertions, and supposed to be caused by the absorption of waste-products.

Fatty (*fat'-e*) [ME., *fat*, fat]. Containing fat, or derived from fat. F. Acids, a series of acids with the formula $C_nH_{2n-1}O(OH)$ some of the members of which combine with glycerol to form fats. F. Degeneration. See *Degeneration*.

Fauces (*faw'-sez*) [L., the upper part of the throat]. The space surrounded by the palate, tonsils, and uvula. F., Isthmus of, the space at the back of the mouth enclosed by the margin of the palate, the back of the tongue and the pillars of the fauces. F., Pillars of, the folds formed by the palatoglossus muscle in front of the tonsils and by the palatopharyngeus behind them.

Faucial (*faw'-se-al*) [*fauces*, the upper part of the throat]. Pertaining to the fauces.

Favus (*fa'-vus*) [L., a honeycomb]. Tinea favosa; a parasitic skin-disease due to the presence of a vegetable parasite, the *Achorion schönleini*. It is characterized by the presence of round, sulphur-yellow, cup-shaped crusts, having a peculiar musty odor, and which are found on microscopic examination to be composed almost entirely of the elements of the fungus. The disease affects most frequently the scalp, but may occur anywhere. F.-cup, any one of the cup-shaped crusts that characterize favus.

Febricula (*fe-brik'-u-lah*) [*febris*, fever]. A slight fever of short duration, most frequently encountered among children.

Febrifuge (*feb'-rif-uj*) [*febris*, fever; *fugare*, to dispel]. 1. Dispelling fever. 2. An agent that lessens fever.

Febrile (*feb'-ril*) [*febrilis*; *febris*, a fever]. Pertaining to or characterized by fever.

Febris (*feb'-ris*) [L.]. See *Fever*.

Fecal (*fe'-kal*) [*fecex*, sediment]. Pertaining to, consisting of, or discharging, feces.

Feces (*fe'-sez*) [*fecex*, sediment]. The excretions of the bowels. Feces consists of excretions and secretions from the intestines and of undigested food, the latter being made up of digestible substances that escaped digestion and of undigestible matters, such as nuclein, cellulose, chlorophyll, and mineral salts.

Fechner's Law. See *Law*.

Fecula (*fek'-u-lah*) [dim. of *fecex*, sediment]. The starchy part of a seed. Also the sediment subsiding from an infusion.

Fecundation (*feh-un-du'-shun*) [*fecundatio*; *fecundus*, fruitful]. The act of fertilizing.

F., Artificial, fecundation brought about by the injection of semen into the vagina or uterus through a syringe or other instrument.

Fecundity (*fe-kun'-dit-e*) [*fecunditas*, fruitfulness]. The ability to produce offspring.

Fehling's Solution. A solution used as a test for glucose, especially in the urine. The solution is prepared as follows: (a) Copper sulphate, 34.652 grams, is dissolved in about 200 c.c. of water. (b) Sodid potassium tartrate (Rochelle salt), 173 grams, is dissolved in 480 c.c. of sodium hydrate solution of specific gravity 1.14. The two solutions are mixed and sufficient water added to make 1000 c.c. One c.c. of this solution is reduced by 0.005 of glucose. To apply the test, a small quantity of the solution is placed in a test-tube, diluted with water, boiled, and the urine added drop by drop. The presence of sugar is indicated by a reddish or yellow precipitate, due to the reduction of the copper sulphate to the copper oxid or hydrate.

Fel [*L.*]. Bile. **F. bovis purificatum**, purified ox-gall; it is said to be tonic and laxative. Dose gr. iij-v (0.2-0.32).

Felon (*fel'-on*). See *Paronychia*.

Female (*fe'-mal'*) [*femina*, woman]. 1. Pertaining to the sex that conceives and bears young; pertaining to a woman. 2. Denoting that part of a double-limbed instrument that receives the complementary part. **F. Catheter**, a catheter for emptying the female bladder.

Femoral (*sem'-or-al*) [*femur*, the thigh bone]. Pertaining to the femur. **F. Arch**, Poupart's ligament. **F. Artery**. See *Arteries*, *Table of*. **F. Canal**, a canal in the sheath of the femoral vessels, to the inner side of the femoral vein, through which at times a hernia descends. **F. Hernia**. See *Hernia*. **F. Ligament of Hey**. See *Ligament*. **F. Ring**, the abdominal end of the femoral canal, normally closed by the crural septum and the peritoneum. **F. Sheath**, a continuation downward of the fasciæ that line the abdomen. It contains the femoral vessels.

Femorotibial (*sem-or-o-tib'-e-al*). Relating to the femur and the tibia.

Femur (*fe'-mur*) [*L.*]. The thigh-bone.

Fenestra (*fe-nest'-trah*) [*L.*, a window]. A window-like opening, as the open space in the blade of an obstetric forceps, or an opening in a bandage or dressing for drainage, etc. **F. ovalis**, **F. rotunda**. See *Ear*.

Fenestrated (*fe-nest'-tra-ted*) [*fenestra*, a window]. Perforated. **F. Membrane of Henle**, the layer of elastic tissue in the intima of large arteries.

Fennel (*fen'-el*) [*ME.*, *fenel*]. See *Faniculum*.

Fenugreek (*fen'-u-grêk*). The *Trigonella fenumgræcum*, a leguminous plant cultivated in France and Germany, the seeds of which contain two alkaloids, cholin and trigonellin. The seeds are employed for the preparation of emollient poultices, enemata, ointments, and plasters. They are not used internally.

Ferment (*fer'-ment*) [*fermentum*, leaven, yeast]. Any substance which in contact with another substance is capable of setting up changes (fermentation) in the latter, without itself undergoing much change. Ferments are classified into unorganized, or soluble, and organized, or living, ferments. According to the character of the fermentation, the unorganized ferments are divided into amylolytic, proteolytic, fat-decomposing, milk-curdling, and coagulating ferments. *Amylolytic ferments* (sugar-producing or diastatic ferments) convert starch into sugar. The most important is ptyalin of the saliva; but similar ferments are found in the pancreatic and intestinal juice, bile, blood, milk, urine, etc. *Proteolytic ferments* convert proteids into peptones and albumoses. They are found in the stomach (*pepsin*), in the pancreatic juice (*trypsin*), and elsewhere. A *fat-decomposing ferment* is found in the stomach and in the pancreatic juice. *Milk-curdling ferment* is found in the stomach and the intestinal juice. The best example of a coagulating ferment is the fibrin-ferment. The organized or living ferments are the yeasts and bacteria. See *Fermentation*.

Fermentation (*fer-men-tat'-shun*) [*fermentum*, leaven]. The decomposition of complex molecules under the influence of ferments. **F., Acetic**, the fermentation whereby weak alcoholic solutions are converted into vinegar; caused by the *Bacillus aceti*, etc. **F., Alcoholic**, that occurring in carbohydrates under the action of various saccharomycetes. **F., Ammoniacal**, that giving rise to ammoniacal gas and carbon dioxide, which combine to form ammonium carbonate. The agent of the ammoniacal fermentation of urine is the *Micrococcus ureæ*. **F., Butyric**, the conversion of sugars, starches, milk, etc., into butyric acid, due to various microorganisms, especially the *Bacillus butyricus*. **F., Lactic**, the "souring" of milk, caused by the *Bacillus lacticus*. **F., Propionic**, the *Bacillus cavidia*, decomposes saccharine solutions, producing propionic acid. **F., Test**, a test for glucose in the urine. The glucose is decomposed by yeast into alcohol and CO₂, the escape of the latter causing a reduction in the specific gravity of the urine. From the loss in weight the amount of glucose is calculated. **F., Viscous**, a fermentation characterized by the production of a gummy substance.

Fern. Any cryptogamous plant of the order *Filices*. See *Aspidium*.

Ferralia (*fer-a'-le-ah*) [*ferrum*, iron]. Preparations containing iron.

Ferrein, Pyramids of. See *Pyramid*.

Ferric (*fer'-ik*) [*ferrum*, iron]. Containing iron as a quadrivalent element.

Ferricyanid (*fer-i-si'-an-id*). A compound of ferricyanogen $\text{Fe}_3(\text{CN})_6$ with an element or radicle.

Ferro- (*fer'-o-*) [*ferrum*, iron]. A prefix used with the names of ferrous compounds.

Ferrocyanid (*fer-o-si'-an-id*). A compound of ferrocyanogen, $\text{Fe}(\text{CN})_2$, with an element or radicle.

Ferrous (*fer'-us*) [*ferrum*, iron]. Containing iron as a bivalent element.

Ferrum (*fer'-um*) [L.]. Iron, $\text{Fe} = 56$. Quantivalence 11, IV. The most familiar and most useful of all metals; it is found in many minerals, in nearly all soils, in many mineral waters, and also occurs pure, especially in the form of meteoric iron. Pure iron is rare, nearly all commercial irons containing carbon in various proportions. In pharmacy, iron is used in the form of fine, bright, non-elastic wire, as reduced iron, a metallic iron with a variable amount of iron oxid, and in the form of salts. The therapeutic properties of iron depend on its power to build up the blood, it being a normal constituent of the red corpuscles. Hence it is useful in all forms of anemia and in the diseases depending upon the latter. Externally many of the soluble salts of iron are used as styptic and astringent lotions. The following preparations of iron are chiefly employed: *F. acetatis*, Liq. Dose $\mathfrak{m}\text{ij}-\text{x}$ (0.13-0.65). *F. albuminas*; it contains 5 per cent. ferric oxid, or 10 per cent. ferric chlorid. Dose gr. xx-xxx (1.3-2.0). Unof. *F. arsenias*, (B. P.), is chiefly valuable for the arsenic it contains. Dose gr. $\frac{1}{10}-\frac{1}{2}$ (0.006-0.008). *F. bromidum*, used in solution, in doses of 20 drops, three times daily, as an alterative tonic in chorea and scrofula. Unof. *F. carbonas saccharatus*, U. S. P. Dose gr. v-xxx (0.32-2.0). *F. carbonatis*, Massa, Vallet's Mass. Dose gr. iij-v (0.2-0.32). *F. carbonatis*, Pilulæ, Bland's Pill; it consists of ferrous sulphate, potassium carbonate, and sugar. *F.*, Mistura, Composita, Griffith's Mixture. Dose $\mathfrak{f}\text{ss}-\mathfrak{f}\text{ss}$ (16.0-32.0). *F. chloridum* is used in the preparation of *F. chloridi*, Liquor, dose $\mathfrak{m}\text{ij}-\text{x}$ (0.13-0.65), and *F. chloridi*, Tinct., dose $\mathfrak{m}\text{x}-\text{xxx}$ (0.65-2.0). *F. citras*. Dose gr. iij-v (0.2-0.32). *F. citratis*, Liquor. Dose $\mathfrak{m}\text{x}$ (0.65). *F. et ammonii citras*. Dose gr. v (0.32). *F. et quininæ citras* (B. P.). Dose gr. v-xv (0.32-1.0). *F. et quininæ citras solubilis*. Dose gr. v (0.32). *F. et strychninæ citras*. Dose gr. j-iij (0.065-0.2). *F.*,

Vinum amarum, is made from the soluble citrate of iron and quinin. Dose $\mathfrak{f}\text{ss}$ j-iij (4.0-8.0). *F. citratis*, Vinum (U. S. P.). Dose $\mathfrak{f}\text{ss}$ j-iij (4.0-8.0). *F. et magnesi citras*. Dose gr. v-x (0.32-0.65). Unof. *F. et bismuthi citras*. Unof. *F. dialysatum*. Dose of the solution 10-20 drops (0.32-0.65). *F. dialysatus*, Liq. (B. P.). Dose $\mathfrak{m}\text{x}-\text{xxx}$ (0.65-2.0). *F. hypophosphis* is used to prepare *F. hypophosphitis*, Syrupus, N. F. Dose $\mathfrak{f}\text{ss}$ ij-vj (8.0-24.0). *F. iodidum*. Dose gr. j-vij (0.065-0.52). Unof. *F. iodidum saccharatum*. Dose gr. ij-v (0.13-0.32). *F. iodidi*, Pil., Blancard's pills; each contain one grain of ferrous iodid. *F. iodidi*, Syrupus. Dose $\mathfrak{m}\text{xv}-\text{xxx}$ (1.0-2.0). *F. lactas*. Dose gr. v (0.32). Syrupus hypophosphitum cum Ferro contains *F. lactas*. Dose $\mathfrak{f}\text{ss}$ ss-j (2.0-4.0). *F. oxalas*. Dose gr. ij-iij (0.13-0.20). Unof. *F. oxidum hydratum* is an antidote to arsenic. *F. oxidum hydratum cum magnesio* is an antidote to arsenic. *F. oxidum magnetium*. Dose gr. v-xx (0.32-1.3). Unof. *F. phosphas* (B. P.). Dose gr. v-x (0.32-0.65). *F. phosphas solubilis*. Dose gr. v-x (0.32-0.65). *F. pyrophosphas solubilis*. Dose gr. ij-v (0.13-0.32). Syrupus ferri, quininæ et strychninæ phosphatum. Dose $\mathfrak{f}\text{ss}$ j (4.0). *F. reductum*. Dose gr. iij-vj (0.2-0.4). *F. succinas*. Dose gr. v (0.32). Unof. *F. sulphas*. Dose gr. v (0.32). *F. sulphas exsiccatus*. Dose gr. iij (0.2). *F. sulphas granulatus*. Dose gr. iij (0.2). *F. tannas*. Dose gr. viij-xxx (0.52-2.0) in 24 hours. *F. valerianas*. Dose gr. ij-v (0.13-0.32).

Fertile (*fer'-til*) [*fertilis*, fruitful]. Prolific, fruitful.

Fertilization (*fer-til-is-a'-shun*) [*fertilis*, fruitful]. The art of making fertile; impregnation.

Ferula (*fer'-u-lah*) [L.]. A genus of the order Umbellifere. See *Asa fetida* and *Galbanum*.

Festination (*fes-tin-a'-shun*) [*festinare*, to hasten]. A gait that increases in rapidity; it is seen in paralysis agitans.

Fetal (*fe'-tal*) [*fetus*, offspring]. Pertaining to the fetus.

Feticide (*fe'-tis-id*) [*fetus*, offspring; *cedere*, to kill]. The killing of the fetus in the womb.

Fetus (*fe'-tus*) [*fetus*, offspring]. The unborn offspring of viviparous animals in the later stages of development.

Fever (*fe'-ver*) [*febris*, a fever]. 1. An elevation of the body-temperature above the normal. 2. A disease the distinctive characteristic of which is elevation of temperature, accompanied also by quickened pulse and respirations, increased tissue-waste, and dis-

ordered secretions. **F.**, **Asthenic**, one in which there is a weak circulation, a clammy skin, and a low state of the nervous system. **F.-blister**. See *Herpes labialis*. **F.**, **Brain**, fever associated with inflammation of the cerebral meninges; meningitis. **F.**, **Catarrhal**, influenza. **F.**, **Catheter**. See *F.*, *Urethral*. **F.**, **Cerebrospinal**, an acute infectious disease characterized by inflammation of the meninges of the brain and cord with involvement of the superficial layers of nerve-substance. See *Exanthemata*, *Table of*. **F.**, **Chagres**, a malignant form of malaria. **F.**, **Continued**, one the course of which is free from remissions or intermissions. **F.**, **Dandy**. Synonym of *Dengue*. **F.**, **Enteric**. See *F.*, *Typhoid*. **F.**, **Eruptive**, one that is accompanied by an eruption on the skin. **F.**, **Famine**. See *F.*, *Relapsing*. **F.**, **Gastric**, a term used indefinitely to indicate any febrile ailment associated with abdominal symptoms. **F.**, **Hay**. See *Hay-fever*. **F.**, **Hectic**, a diurnally intermittent fever with the highest temperature in the evening and accompanied by sweats and chills. It is found in tuberculosis and other diseases associated with the absorption of septic products. **F.**, **Low**, fever of an asthenic type. **F.**, **Lung**, croupous pneumonia. **F.**, **Milk**, a slight form of puerperal septicemia, formerly thought to be due to the formation of milk in the mother's breast. **F.**, **Puerperal**. See *Puerperal*. **F.**, **Relapsing**. See *Relapsing Fever*. **F.**, **Remittent**, a paroxysmal fever with exacerbations and remissions, but not intermissions; usually applied specifically to remittent malarial fever. **F.**, **Rheumatic**. See *Rheumatism*. **F.**, **Scarlet**. See *Scarlet Fever*. **F.**, **Septic**, one due to the entrance of septic matter into the system. **F.**, **Simple** **Continued**, a continued, noncontagious fever, varying in duration from one to twelve days, and usually ending in recovery. **F.**, **Spirillum**. Synonym of *F.*, *Relapsing*. **F.**, **Splenic**. Synonym of *Anthrax*. **F.**, **Sthenic**, a fever characterized by rapid, full pulse, heat and dryness of the skin, high temperature, scanty urine, and delirium. **F.**, **Traumatic**, that following traumatism. **F.**, **Typhoid**. See *Typhoid Fever*. **F.**, **Typhus**. See *Typhus Fever*. **F.**, **Urethral**, the febrile disturbance that follows the use of the catheter or bougie. **F.**, **Yellow**. See *Yellow Fever*. **Fiber** (*fī'-ber*) [*fibra*, a thread]. A filamentary or thread-like structure. **F.** of Gerdy, a transverse band of fibers supporting the hand crosswise and forming the web of the fingers. **F.**, **Muscle**. See *Muscle*. **F.**, **Nerve**. See *Nerve*. **F.** of Remak, the nonmedullated nerve-fibers. **F.** of Sharpey, delicate processes or rods that pass through and rivet together the concentric laminae of

bone. **F.** of Tome, elongated and branched processes of the odontoblasts of the pulp, filling the dentinal tubules of teeth.

Fibril (*fī'-bril*) [*fibra*, a fiber]. A little fiber, particularly one of the ultimate fibers of muscle or of fibrous connective tissue.

Fibrillar (*fī'-bril-ar*) [*fibrilla*, a small fiber]. Pertaining to fibrils. **F.** **Contractions**, spontaneous contractions successively taking place in different bundles of muscular fibers; they are seen in progressive muscular atrophy and other diseases.

Fibrillary (*fī'-bril-a-re*). Same as *Fibrillar*.

Fibrin (*fī'-brin*) [*fibra*, a fiber]. A proteid formed in shed blood, lymph, in other body-fluids, and in tissues when these coagulate. It exists in the shape of fibrils, granules, plates, or as a homogeneous material. Fibrin forms about 0.2 per cent. of the blood.

F.-factors, the substances necessary for and concerned in the formation of fibrin. They are fibrinogen, fibrin-ferment, and certain salts. **F.-ferment**, a ferment obtained from blood-serum after clotting has occurred. It is one of the fibrin-factors, and is probably derived from the leukocytes.

Fibrinogen (*fī-brin'-o-jen*) [*fibra*, a fiber; *γεννᾶν*, to produce]. A proteid of the globulin class, obtained from blood-plasma and serous transudations. It is one of the chief elements in the formation of fibrin.

Fibrinogenous (*fī-brin-ōj'-en-us*) [*fibra*, a fiber; *γεννᾶν*, to produce]. Forming or producing fibrin.

Fibrinoplastic (*fī-brin-o-plas'-tik*) [*fibra*, a fiber; *πλάσσειν*, to form]. Of the nature of fibrinoplastin.

Fibrinoplastin (*fī-brin-o-plas'-tin*). See *Paraglobulin*.

Fibrinous (*fī'-brin-us*) [*fibra*, a fiber]. Of the nature of or containing fibrin.

Fibroareolar (*fī-bro-ar-e'-o-lar*) [*fibra*, fiber; *areola*, dim. of *area*, an open space]. Containing fibrous tissue with an areolar arrangement.

Fibrocartilage (*fī-bro-kar'-sil-aj*) [*fibra*, a fiber; *cartilago*, gristle]. Cartilage with an intermixture of fibrous elements.

Fibrocellular (*fī-bro-sel'-u-lar*) [*fibra*, fiber; *cella*, cell]. Both fibrous and cellular; fibroareolar.

Fibrocyst (*fī'-bro-sist*) [*fibra*, fiber; *κυστίς*, cyst]. A fibroma that has undergone cystic degeneration.

Fibroglioma (*fī-bro-gli-o'-mah*) [*fibra*, fiber; *γλῖα*, glue; *δῶμα*, tumor]. A tumor having the elements of a fibroma and a glioma.

Fibroid (*fī'-broid*) [*fibra*, fiber; *ειδός*, likeness]. Resembling fibers or composed of fibers. Also, a fibroid tumor. **F.** **Heart**, a chronic form of myocarditis in which there is a development of fibrous connective tissue

in the cardiac muscle. **F. Induration.** See *Induration*. **F. Phthisis**, a chronic form of phthisis, in which there is a formation of fibrous tissue.

Fibroma (*fi-bro'-mah*) [*fibra*, a fiber; *δμα*, a tumor]. A benign tumor composed of fibrous tissue. **F. Hard**, one containing few cells, being chiefly composed of fibers. **F. lipomatodes**. Same as *Xanthoma*. **F. molluscum**. Synonym of *Molluscum fibrosum*. **F. Soft**, one rich in cells.

Fibromyoma (*fi-bro-mi-o'-mah*) [*fibra*, fiber; *μῦς*, muscle; *δμα*, a tumor]. A tumor composed of fibrous and muscular tissue.

Fibromyxoma (*fi-bro-miks-o'-mah*) [*fibra*, fiber; *μύξα*, mucus; *δμα*, a tumor]. A tumor composed of fibrous and myxomatous tissue.

Fibroneuroma (*fi-bro-nu-ro'-mah*) [*fibra*, fiber; *νεῖρον*, nerve; *δμα*, tumor]. A tumor composed of fibrous tissue and nerve-fibers.

Fibroplastic (*fi-bro-plas'-tik*) [*fibra*, fiber; *πλασσειν*, to form]. Tending to form fibers.

Fibrosarcoma (*fi-bro-sar-ko'-mah*) [*fibra*, fiber; *σάρξ*, flesh; *δμα*, tumor]. A sarcoma containing fibrous tissue.

Fibrosis (*fi-bro'-sis*) [*fibra*, fiber]. The development of fibrous tissue. **F., Arterio-capillary**. See *Endarteritis*.

Fibrous (*fi'-brus*) [*fibra*, a fiber]. Containing fibers; of the character of fibrous tissue.

Fibula (*fi'b'-u-lah*) [*L.*, a buckle]. The slender bone at the outer part of the leg articulating above with the tibia and below with the astragalus and tibia.

Ficus (*fi'-kus*) [*L.*, a fig-tree]. The fig. The fleshy receptacle of *F. carica*, native of Asia Minor, and cultivated throughout Europe and tropic America. It is laxative and nutritious, and is a constituent of Confectio sennæ.

Field (*feld*) [*ME., feeld*]. 1. An open space or area. 2. The field of vision. **F. of Fixation.** See *Fixation*. **F. Magnetic**, the portion of space about a magnet in which its action is felt. **F. of a Microscope**, the area that can be seen through a microscope at one time. **F. of Vision**, the space in which the patient can see when the eye is steadily fixed on the object held in the direct line of vision.

Fifth Nerve. See *Trifacial*. **F. Ventricle.** See *Ventricle*.

Fig. See *Ficus*. **F.-wart**, a moist condyloma. **F.-wort**, the herb *Scrophularia nodosa*, an alterative, diuretic, and anodyne. It is sometimes used in the form of an ointment for piles. Dose of the fld. ext. f ʒ ss-j (2.0-4.0). Unof.

Filament (*fil'-am-ent*) [*filum*, a thread]. A small, thread-like structure. **F., Spermatic**, the caudal filament of a spermatozoon.

Filaria (*fi-la'-re-ah*) [*filum*, a thread]. A genus of nematode or thread-worms of the

family Filariidæ. **F. medinensis**, Guinea Worm, an animal parasite, the female of which works its way from the intestinal tract to the subcutaneous tissue, where, after developing its embryos, it is sooner or later set free by abscess-formation and discharge. **F. sanguinis hominis**, the larva or embryo of the nematode worm, *Filaria bancrofti*. The embryos are about 0.35 mm. long, and inhabit the lymph-channels of the lower extremities and the scrotum. They lead to dilatation of the lymphatics, to hyperplasia of the tissues, to chyluria, hematuria, abscesses, etc. They are found in the blood at night. Elephantiasis arabum and lymph-scrotum are said to be due to the filaria.

Filariasis (*fi-lar-i'-as is*) [*filum*, a thread]. A diseased state due to the presence in the body of *Filaria sanguinis hominis*, or allied species.

Filicic Acid, $C_{14}H_{16}O_5$. An acid extracted from *Filix mas*.

Filiform (*fil'-if-orm*) [*filum*, a thread; *forma*, form]. Thread-like. **F. Bougie.** See *Bougie*. **F. Papillæ**, the smallest and most numerous of the papillæ of the tongue, occurring over its whole surface.

Filix (*fi'-liks*) [*L.*]. A fern. **F. mas**, male fern. See *Aspidium*.

Fillet (*fil'-et*) [*Fr., filet*, a thread]. 1. A loop for the purpose of making traction on the fetus. 2. The lemniscus, a band of nerve-fibers connected below with the nucleus gracilis and nucleus caudatus of the medulla and running upward through the pons and crus cerebri to the cerebrum, a portion of the fibers (lateral fillet) entering the posterior corpora quadrigemina, another (mesal fillet) passing to the anterior corpora quadrigemina and the optic thalamus. A part of the mesal fillet is continued into the cortex. **F., Olivary**, a fasciculus of nerve-fibers enclosing the olivary body of the medulla.

Filter (*fil'-ter*) [*filtrum*, felt]. An apparatus for straining water or other liquids to remove any undissolved matters. **F., Pasteur-Chamberland**, a hollow column of unglazed porcelain through which solutions are filtered by means of a vacuum-exhaust or by pressure. **F.-paper**, an unglazed paper used for filtration.

Filtrate (*fil'-trat*) [*filtrum*, felt]. The liquid that has passed through a filter.

Filtration (*fil'-tra'-shun*) [*filtrum*, felt]. The operation of straining through a filter.

Filum terminale (*fil'-lum ter-min-al'-e*). A long, slender thread of pia mater, the termination of the spinal cord.

Fimbria (*fin'-bre-ah*) [*fimbria*, a fringe]. A fringe. **F. of Fallopian Tube**, the fringe-like processes of the outer extremity of the oviduct.

Fimbriated (*fim'-bre-a-ted*) [*fimbria*, fringe]. Fringed.

Finger (*ing'-ger*) [ME., *finger*]. A digit of the hand. **F.**, **Clubbed**, a finger the terminal phalanx of which is short and broad with overhanging nail. It is seen in cases of pulmonary tuberculosis, congenital heart-disease, etc.

Fir (*fur*) [ME., *fir*, the fir-tree]. See *Abies*. **F.-balsam**. See *Abies balsamea*.

First Intention. See *Healing*.

Fish-berry. See *Cocculus indicus*.

Fish-skin Disease. See *Ichthyosis*.

Fissiparous (*sis-ip'-ar-us*) [*findere*, to split; *parere*, to produce]. Propagating by fission.

Fissure (*fish'-ur*) [*fissura*; *findere*, to cleave]. A groove or cleft. A term applied to the clefts or grooves in various organs, as the skull, the brain, the liver, the spinal cord. Also to cracks in the skin or linear ulcers in mucous membranes. **F. of the Anus**, a linear ulcer at the mucocutaneous junction of the anus, giving rise to intense suffering on defecation. **F. of Bichat**, the transverse fissure of the brain, between the fornix and the upper surface of the cerebellum. Through it the pia enters the ventricles. **F. of Broca**, a fissure surrounding Broca's lobe. **F.**, **Calcarine**, a fissure on the mesal aspect of the cerebrum, between the lingual lobule and the cuneate lobe. **F.**, **Callosomarginal**, one on the mesal surface of the cerebral hemisphere, dividing the area between the corpus callosum and the margin into nearly equal parts. **F.**, **Longitudinal**, the deep fissure that divides the cerebrum into two hemispheres. **F.**, **Occipital**, a deep fissure, situated between the parietal and occipital lobes of the brain. **F.**, **Palpebral**, the space between the eyelids extending from the outer to the inner canthus. **F.**, **Portal**. See *F.*, *Transverse (of liver)*. **F.**, **Precentral**, a fissure in front of the *F.* of Rolando and parallel to it. **F. of Rolando**, a fissure on the lateral aspect of the cerebrum, extending downward from near the longitudinal fissure at about its middle point. It separates the frontal from the parietal lobe. **F. of Sylvius**, a deep fissure of the brain beginning on the outer side of the anterior perforated space, and extending outward to the lateral surface of the hemisphere. It has two branches, a short vertical and a long horizontal, the latter separating the parietal from the temporo-sphenoidal lobe. Between the branches lies the Island of Reil. **F.**, **Transverse (of liver)**, a fissure crossing transversely the lower surface of the right lobe of the liver. It transmits the portal vein, hepatic artery and nerves, and hepatic duct.

Fistula (*sis'-tu-lah*) [L., a pipe]. A narrow canal or tube left by the incomplete healing

of abscesses or wounds, and usually transmitting some fluid, either pus or the secretions or contents of some organ or body-cavity.

F., **Anal**, a fistula in the neighborhood of the anus which may or may not communicate with the bowel. **F.**, **Blind**, a fistula open at one end only. **F.**, **Blind, External**, one the only opening of which is on the exterior of the body. **F.**, **Blind, Internal**, one which opens only upon an internal surface. **F.**, **Branchial**, an opening that extends from the surface of the neck to the pharynx; it is an unclosed branchial cleft. **F.**, **Complete**, one having two openings, an internal and external. **F.**, **Fecal**, a fistula communicating with the intestine.

Fit [ÄS., *fit*, a struggle]. A name applied to any sudden paroxysm of a disease, but especially to an epileptic convulsion.

Fixation (*fix'-a'-shun*) [*fixus*, fixed]. The act of fixing or making firm. **F.**, **Field of**, in optics, the region bounded by the utmost limits of distinct or central vision, and which the eye has under its direct control through its excursions, without movements of the head. **F.-forceps**, those used for fixing or holding a part in position during a surgical operation.

Fixing (*fix'-ing*) [*fixus*, fixed]. The preparation of tissue for microscopic study by means of some agent that hardens it and preserves the form and arrangement of the cells.

Flagellate (*flaj'-el-ät*) [*flagellum*, a whip]. Furnished with slender, whip-like processes.

Flagellum (*flaj'-el'-um*) [L., a whip]. A whip-like mobile process; the organ of locomotion of certain bacteria and infusoria.

Flag, Sweet. See *Calamus*.

Flank (*flank*) [ME., *flank*, from L. *flaccus*, soft]. The part of the body between the ribs and the upper border of the ilium.

Flap [ME., *flap*, a loose, flexible part]. A loose and partly detached portion of the skin or other soft tissue. **F.-amputation**, one in which flaps of soft tissues are left to cover over the end of the bone. **F.-extraction**, a method of extracting the crystalline lens so as to make a flap of the cornea.

Flat [ME., *flat*]. 1. Lying on one plane; having an even surface. 2. Of a percussion-sound, having no resonance. **F.-foot**, depression of the plantar arch.

Flatness (*flat'-nes*) [ME., *flat*]. The sound obtained by percussing over an airless organ or a large effusion.

Flatulence (*flat'-u-lens*) [*flatus*, breath]. A condition marked by the presence of gas in the stomach and intestinal canal.

Flatulent (*flat'-u-lent*) [*flatus*, breath]. Characterized by flatulence.

Flatus (*fla'-tus*) [L.]. Gas, especially gas in the gastrointestinal canal.

Flax [AS., *flax*]. See *Linum*. **F.-dresser's Phthisis**, a fibroid pneumonia resulting from the inhalation of particles in flax-dressing. **F.-seed**. See *Linum*.

Flea (*fle*). See *Pulex*. **F.-bane**. See *Erigeron*.

Fleam (*flem*) [*φλεβοτόμον*, a lancet, from *φλέψ*, a vein; *τέμνειν*, to cut]. A phlebotome; a lancet for venesection.

Fleece of Stilling. A group of white fibers surrounding the corpus dentatum of the cerebellum.

Fleming's Tincture. An alcoholic preparation of aconite, stronger than the official tincture. Dose, two minims (o.13).

Flemming's Solution. A mixture used as a fixing agent for tissue in histologic study. It consists of 15 parts of 1 per cent. solution of chromic acid, 4 parts of a 2 per cent. solution of osmic acid, 1 part of glacial acetic acid.

Fleah [AS., *flase*, flesh]. The soft tissues of the body, especially the muscles. **F., Proud**, the soft and exuberant granulations of a wound or ulcer.

Flexibilitas (*fleks-i-b-il'-it-as*) [L.]. Flexibility. **F. cerea**, a condition of the limbs in catalepsy in which they seem as if made of wax.

Flexible (*fleks'-ib-l*) [*flexus*, bent]. That which may be bent, as a F. catheter, F. colloid.

Flexion (*flek'-shun*) [*flexus*, bent]. The act of bending; the condition of being bent.

Flexor (*fleks'-or*) [*flexus*, bent]. A muscle that bends or flexes a limb or part. See *Muscles*, *Table of*.

Flexure (*fleks'-ūr*) [*flexura*, a bending]. A bending. **F., Caudal**, the bend at the lower portion of the embryo. **F., Cephalic**, the arching over of the cephalic end of the embryo. **F., Sigmoid**. See *Sigmoid Flexure*.

Floating (*flō'-ting*) [AS., *floatan*, to float]. Swimming; free to move around. **F. Kidney**, one that is detached from its normal position and abnormally movable. **F. Liver**, one with abnormal mobility; movable liver. **F. Ribs**. See *Ribs*.

Floccitation (*flok-sit'-a'-shun*). Same as *Carphology*.

Flocculent (*flok'-u-lent*) [*floccus*, a flock of wool]. Flaky, downy, or woolly; coalescing in floccy masses.

Flocculus (*flok'-u-lus*) [dim. of *floccus*, a tuft of wool]. A prominent lobe of the cerebellum situated behind and below the middle cerebellar peduncle on either side of the median fissure.

Flood's Ligament. See *Ligament*.

Flooding (*flud'-ing*) [AS., *flod*, a flood]. A copious bleeding from the womb.

Flores (*flō'-rēs*) [L.]. Plural of *flor*, a flower. 1. The flowers or blossoms of a plant.

2. A flocculent or pulverulent form assumed by certain substances after sublimation, as *flores sulphuris*, flowers of sulphur.

Florida Allspice, the leaves of *Calycanthus floridus*, having the properties of an aromatic stimulant. Unof.

Flour (*flour*) [*flor*, a flower]. The finer part of the ground grain, especially of wheat.

Flow (*flo*) [AS., *flowan*, to flow]. The free discharge of a liquid, as the blood; the menses.

Flower (*flou'-er*). See *Flores*.

Fluctuation (*fluk-tu-a'-shun*) [*fluctuare*, to float or roll]. The wave-like motion produced when a body containing fluid is tapped between the fingers or hands.

Fluor albus (*flu'-or al'-bus*) [L., a white flow]. Leukorrhea.

Fluorescein (*flu-or-es'-e-in*), $C_{20}H_{12}O_6 + H_2O$. An anhydrid of resorcin, prepared by heating phthalic anhydrid with resorcin to 200° C. It has the property of coloring abrasions of the cornea greenish, and has been used on this account for diagnostic purposes.

Fluorescence (*flu-or-es'-ens*) [*fluere*, to flow]. A property possessed by certain substances of converting obscure actinic rays (such as the ultraviolet) into luminous rays.

Fluorin (*flu'-or-in*) [*fluor*, a flow]. $F = 19$; quantivalence 1. An element belonging to the chlorin group. Its intense chemic affinity has so far rendered its isolation impossible. The salts formed with the alkaline metals, fluorides, have been used in goiter and in rheumatism. See *Elements*, *Table of*. Hydrofluoric acid is a powerful caustic.

Flux (*fluks*) [*fluxus*, flowing]. 1. An excessive flow of any of the excretions of the body, especially the feces. 2. Dysentery.

Focal (*fo'-kal*) [*focus*, a fire-place]. Pertaining to or occupying a focus. **F. Depth**, the power of a lens to give clear images of objects at different distances from it. **F. Distance**, the distance from the focus to a reflecting or refracting surface, or in the case of a lens to the principal point of the lens. **F. Disease**, **F. Lesion**, one that is limited to a small area. **F. Epilepsy**, epilepsy due to a focal lesion of the brain; Jacksonian epilepsy.

Focus (*fo'-kus*) [*focus*, a fire-place]. 1. The principal seat of a disease. 2. The point (called principal focus or real focus), at which rays of light converge that pass through a convex lens or are reflected from a concave mirror. **F., Negative** or **Virtual**, the point at which divergent rays would meet if prolonged in a backward direction.

Fœniculum (*fen-ik'-u-lum*) [L.]. Fennel. The fruit of *F. vulgare*, the properties of which are due to a volatile oil. It is a mild

stimulant and aromatic carminative. **F.**, **Aqua**, 2 parts of the oil in 1000 of water. Dose $\frac{f\text{ss}}{ss}$ – $\frac{f\text{ss}}{ss}$ (2.0–32.0). **F.**, **Oleum**, Dose $\frac{m\text{ij}}{ij}$ – $\frac{v}{v}$ (0.13–0.32).

Fœnum Græcum (*fœ'-num gree'-kum*).

See *Penugreek*.

Fœtal (*fœ'-tal*). See *Fetal*.

Fœtus (*fœ'-tus*). See *Fetus*.

Folia (*fœ'-le-ah*) [*folium*, leaf]. Leaves.

Follicle (*fol'-ik-l*) [*folliculus*, dim. of *follix*, bellows]. 1. A small lymphatic gland, the tissue of which is arranged in the form of a little sac; also a small tubular gland. 2. A simple tubular gland. **F.**, **Graafian**, one of the small vesicular bodies in the ovary, each of which contains an ovum. **F.** of **Liebkühn**. See *Crypts* of Liebkühn. **F.**, **Sebaceous**, the sebaceous glands of the skin. **F.**, **Solitary**, small discrete lymph-follicles found in the mucous membrane of the intestines.

Follicular (*fol'-ik'-u-lar*) [*folliculus*, a little follicle]. Pertaining to a follicle.

Folliculitis (*fol'-ik-u li'-tis*) [*folliculus*, a follicle; *itis*, inflammation]. Inflammation of a group of follicles. **F.** *barbæ*. See *Sycosis*.

Fomentation (*fo-men-tat'-shun*) [*fomentare*, to foment]. The application of heat and moisture to a part to relieve pain or reduce inflammation. It may be by cloths soaked in hot water or medicated solution, or by a poultice. Also the substance applied.

Fontana's Spaces. Spaces between the processes of the ligamentum pectinatum of the iris.

Fontanel, or **Fontanelle** (*fon-tan-el'*) [dim. of *fontana*, a fountain]. A membranous space between the cranial bones in fetal life and infancy. **F.**, **Anterior**, that at the point of union of the frontal, sagittal, and coronal sutures. **F.**, **Posterior**, that at the point of junction of the lambdoid and the sagittal sutures.

Food [*AS.*, *fôda*, food]. Anything which when taken into the body is capable of building up tissue, or by oxidation, of supplying heat.

Foot [*ME.*, *foot*]. 1. The terminal extremity of the leg. It consists of the tarsus, metatarsus, and phalanges, or toes. 2. The base of a microscope. 3. A measure of length equal to twelve inches, or 30.479 cm. **F.-clonus**. See *Ankle-clonus*. **F.-drop**, a falling of the foot due to a paralysis of the flexors of the ankle. **F.**, **Fungus**-. See *Madura-foot*. **F.-and-mouth Disease**, a febrile affection of sheep, cows, pigs, and horses, rarely of man, manifesting itself by the appearance of vesicles and bullæ in the mouth and on the feet. It is probably due to a special microorganism. **F.-pound**, the work equal to that of raising a pound to the height of one foot.

Foramen (*for-a'-men*) [*forare*, to pierce]. A perforation or opening, especially in a bone.

TABLE OF FORAMINA.

NAME.	LOCATION.	TRANSMITS.
Auditory, external.	The external meatus of the auditory canal.	Sound-waves to tympanic membrane.
Auditory, internal.	The petrous portion of temporal bone.	The auditory and facial nerves.
Of Bichat.	Connects subarachnoid space and third ventricle (?).	Cerebrospinal fluid.
Of Botallo.	Between the two auricles in the fetal heart.	
Cecal (of frontal bone).	Formed by the frontal bone and the crista galli of the ethmoid; a small foramen.	A vein, occasionally.
Cecal (of medulla oblongata).	A depression at the termination of the anterior median fissure.	
Cecal (of pharynx).	Median line of posterior wall of the pharynx; a depression in the mucous membrane.	
Cecal (of tongue).	Posterior termination of the median raphe of tongue.	A number of small glands open into it.
Condylod, anterior (sometimes double).	Anterior to and to the outer side of each occipital condyle, passing downward, outward, and forward through the basilar process.	Hypoglossal nerve; occasionally a meningeal branch of ascending pharyngeal artery.

TABLE OF FORAMINA.—*Continued.*

NAME.	LOCATION.	TRANSMITS.
Condylold, posterior.	The fossa behind the occipital condyles.	A vein to the lateral sinus.
Cotylold.	A notch in the acetabulum converted into a canal by a ligament.	Vessels and nerves.
Dental, inferior.	The external aperture of the inferior dental canal, in the ramus of the inferior maxilla.	Inferior dental vessels and nerves.
Esophageal.	The diaphragm.	The esophagus.
Ethmoid, anterior.	A canal formed between the ethmoid and frontal bones.	Nasal branch of the ophthalmic nerve; anterior ethmoidal vessels.
Frontal.	The supraorbital notch of the frontal bone when it is converted into a canal by a bony process.	Supraorbital vessels and nerves.
Incisor.	Aperture of the incisor canal in the alveolar margin.	Nerves and vessels to the incisor teeth.
Infraorbital.	In the superior maxilla, the external aperture of the infraorbital canal.	The infraorbital nerve and artery.
Intervertebral, anterior.	The aperture formed by the notches opposite to each other in the laminae of adjacent vertebrae.	The spinal nerves and vessels.
Intervertebral, posterior.	The space between the articular processes of adjacent vertebrae, except the first cervical.	
Jugular.	See <i>F., Lacerated, posterior.</i>	
Lacerated, anterior.	See <i>F., Lacerated, middle.</i>	
Lacerated, middle.	An irregular aperture between the apex of the petrous portion of the temporal bone and the body and great wing of the sphenoid, and the basilar process of the occipital bone.	The carotid artery and the large superficial petrosal nerve.
Lacerated, posterior.	The space formed by the jugular notches of the occipital and temporal bones, divided into two portions.	The <i>posterior portion</i> , the internal jugular vein; the <i>anterior portion</i> , the 9th, 10th, and 11th cranial nerves, and the inferior petrosal sinus.
Of Magendie.	An orifice in the pia mater of the roof of the fourth ventricle.	
Magnum.	A large oval aperture, centrally placed in the lower and anterior part of the occipital bone.	The spinal cord and its membranes; the spinal accessory nerves; the vertebral arteries.
Mastoid.	A small foramen behind the mastoid process.	A small artery from the dura; a vein opening into the lateral sinus.
Medullary.	See <i>Nutrient.</i>	
Mental.	A foramen in the inferior maxilla, external to the incisive fossa.	Mental nerve and vessels.
Of Monro.	An opening behind the anterior pillars of the fornix through which the lateral ventricle of the brain communicates with the third ventricle.	The choroid plexus.
Nutrient.	The canal connecting the medullary cavity of the bone with its external surface.	The nutrient vessels.
Obturator.	The large ovoid opening between the ischium and the pubis, internal and inferior to the acetabulum; it is partly closed in by a fibrous membrane.	Obturator vessels and nerves.

TABLE OF FORAMINA.—*Continued.*

NAME.	LOCATION.	TRANSMITS.
Olfactory.	Numerous foramina in the cribriform plate of the ethmoid.	Olfactory nerves.
Omental, lesser or small.	See <i>Foramen of Winslow</i> .	
Optic.	The canal at the apex of the orbit, the anterior termination of the optic groove, just beneath the lesser wing of the sphenoid bone.	Optic nerve; ophthalmic artery.
Ovale (of the heart).	In the fetus, the opening between the two auricles of the heart, situated at the lower posterior portion of the septum.	
Ovale (of the sphenoid).	An ovoid aperture near the posterior margin of the great wing of the sphenoid.	The inferior maxillary division of the trigeminal nerve; the small meningeal artery; occasionally, the small petrosal nerve.
Palatine, anterior.	The orifice of the incisor canal in the anterior part of the roof of the mouth.	The nasopalatine nerve and a branch of the posterior palatine artery.
Palatine, great posterior.	The orifice of the posterior palatine canal upon the posterior part of the hard palate.	The descending palatine artery.
Parietal.	Near the posterior superior angle of the parietal bone; inconstant.	An emissary vein of the superior longitudinal sinus; occasionally a small branch of the occipital artery.
Pterygopalatine.	The external aperture of the pterygopalatine canal.	Pterygopalatine vessels; pharyngeal nerve.
Quadrata.	The diaphragm.	The inferior vena cava.
Rotundum.	A round opening in the great wing of the sphenoid bone.	The superior division of the fifth nerve.
Sacral, anterior (<i>four on either side</i>).	On the anterior surface of the sacrum, connecting with the sacral canal.	The anterior branches of sacral nerves.
Sacral, posterior (<i>four on either side</i>).	The posterior surface of the sacrum, external to the articular processes.	Posterior branches of sacral nerves.
Sacrosciatic, great.	The oval space between the lesser sacrosciatic ligament and the innominate bone.	The pyriformis muscle, the gluteal, sciatic, and pudic vessels and nerves.
Sacrosciatic, small.	The space included between the greater and lesser sacrosciatic ligaments and the portion of the innominate bone between the spine and tuberosity of the ischium	Internal obturator muscle; the internal pudic vessels and nerves.
Of Scarpa.	Bony canals opening into the incisor canal.	The nasopalatine nerves.
Of Sæmmering.	The <i>Fovea centralis</i> .	
Sphenopalatine.	The space between the sphenoidal and orbital processes of the palate bone; it opens into the nasal cavity.	Branches from Meckel's ganglion; the nasal branch of the internal maxillary artery.
Spinosum.	The great wing of the sphenoid bone, near its posterior angle.	The middle meningeal artery.
Of Stensen.	The lateral pair of the four orifices of the anterior palatine foramen.	Terminal branches of the posterior palatine arteries.

TABLE OF FORAMINA.—*Continued.*

NAME.	LOCATION.	TRANSMITS.
Supraorbital.	A notch in the superior orbital margin at the junction of the middle with the inner third, sometimes converted into a foramen by a bony process or a ligamentous band.	Supraorbital artery, veins, and nerve.
Of Thebesius.	Small depressions on the inner surface of the heart, especially in the right auricle.	Some of these are cul-de-sacs; others are the mouths of the venæ minimæ cordis.
Thyroid.	See <i>Obturator</i> .	
Vertebral.	The space included between the body and arch of a vertebra.	The spinal cord and its appendages.
Vertebralarterial.	The foramina in the transverse processes of the cervical vertebræ.	Vertebral artery and vein.
Of Winslow.	An aperture situated between the liver and stomach, bounded in front by the portal vein, hepatic artery and duct, behind, by the inferior vena cava, below, by the hepatic artery, and above, by the liver. It is formed by folds of the peritoneum, and establishes communication between the greater and lesser cavities of the peritoneum.	

Force (*fors*) [*fortis*, strong]. That which produces or arrests motion.

Forced (*forst*) [*fortis*, strong]. Accomplished by an exertion of force. **F. Feeding.** 1. Systematic overfeeding as a therapeutic measure. 2. Feeding performed against the will of the patient.

Forceps (*for'-seps*) [*forceps*, a pair of tongs]. An instrument with two blades and handles used for purposes of drawing on or compressing an object. **F., Bone-**, a cutting forceps for use in operations upon bones. **F., Hemostatic**, a forceps for controlling hemorrhage.

Forcippresure (*for'-se-presh-ür*) [*forceps*, a pair of tongs; *pressura*, a pressing]. The catching the end of the divided vessel with a pair of spring-forceps, which is left on for some time for the purpose of preventing hemorrhage.

Fore (*ör*) [*AS.*, *fore*, before]. In front, coming first. **F.-arm**, that part of the arm between the wrist and the elbow. **F.-brain**, the anterior of the encephalic vesicles into which the primary nervous axis of the embryo divides at an early stage; the prosencephalon. **F.-finger**, the index finger. **F.-gut**, the embryonic tube corresponding to the pharynx, esophagus, stomach, and duodenum. **F.-head**, that part of the face above the orbits of the eyes. **F.-skin**, the prepuce.

Foreign Body. A substance not normally present in the place where it is found.

Forensic (*for-en'-sik*) [*forensis*, belonging to the forum]. Pertaining to a court of law. In medicine, that part of the science connected with judicial inquiry.

Formal (*for'-mal*). See *Methylal*.

Formaldehyd (*form-al'-de-hid*). See *Aldehyd*, *Formic*.

Formalin (*for'-mal-in*) [*formica*, an ant]. A proprietary substance composed of a 40 per cent. solution of formaldehyd. It is used as an antiseptic and as a fixing-agent in histologic studies.

Formate (*for'-mät*). A salt of formic acid.

Formatio (*for-ma'-she-o*) [*L.*]. A formation. **F. reticularis**, the intercrossing of the fibers of the anterior columns in the medulla.

Formative (*for'-ma-tiv*) [*formare*, to form]. Concerned in the formation of tissue.

Formic Acid (*for'-mik*). See *Acid*.

Formication (*for-mik-a'-shun*) [*formica*, an ant]. A sensation like that produced by ants or other insects crawling upon the skin.

Formol (*for'-mol*). See *Formalin*.

Formula (*form'-u-lah*) [dim. of *forma*, a form]. 1. A prescribed method. 2. The representation of a chemie compound by symbols. **F., Constitutional**, or **Rational**, one that indicates by means of symbols the exact proportion by weight in which the elements enter into a compound. **F., Empiric**, one that indicates only the constituents and their proportions in a molecule, as HNO_3 , nitric acid. **F., Graphic**. See *F., Structural*. **F., Rational**. Same as *F., Constitutional*. **F., Structural**, one in which the symbols are united by the bonds of affinity according to their quantivalence, as H-O-H .

Formulary (*for'-mu-la-re*) [*formula*; dim. of *forma*, form]. A collection of formulas

or recipes. **F.**, National, a collection of widely-used and well-known preparations, omitted from the U. S. Pharmacopeia, but collected and published by the American Pharmaceutical Association.

Formyl (*for' mil*) [*formica*, the ant; *ῥη*, matter], **CHO**. The radicle of formic acid. **F.**, Bromid, bromoform. **F.**, Chlorid, chloroform. **F.**, Iodid, iodoform.

Fornicate (*for'-nik-at*) [*fornix*, an arch]. Arched. **F. Gyrus**. See *Convolution*.

Fornix (*for'-niks*) [*fornix*, an arch]. A triangular body of white matter beneath the corpus callosum. From the apex, situated anteriorly, the anterior pillars arise and descend to form the corpora mamillaria. From the extremities of the base the posterior pillars descend into the lateral ventricles. The fornix serves as an anteroposterior commissure between the optic thalamus and the hippocampus major and the uncinate gyrus.

Fortification-spectrum (*for-tif-ik-a'-shun-spek'-trum*). Teichopsia; the appearance of a peculiar subjective visual sensation, sometimes followed by migraine. The luminous shape assumes a zigzag form, with angles resembling those of a fortification.

Fossa (*fos'-ah*) [*fossa*, a ditch]. A depression or pit. **F.**, Canine, a depression on the external surface of the superior maxilla, above and to the outer side of the socket of the canine tooth. **F.**, Cranial, any of the three depressions in the base of the skull for the reception of the lobes of the brain. **F.**, Digastric, a deep groove on the inner aspect of the mastoid process. **F.**, Digital, a depression at the base of the inner surface of the great trochanter of the femur. **F.**, Glenoid, the fossa in the temporal bone that receives the condyle of the lower jaw. **F.**, Hyaloid, a depression in the anterior surface of the vitreous body for the crystalline lens. **F.**, Iliac, the smooth internal surface of the ilium. **F.**, Ischio-rectal, the depression on either side of the anus, bounded on the outer side by the tuberosity of the ischium. **F.**, Lacrymal, the depression in the orbital plate of the frontal bone for the reception of the lacrymal gland. **F. navicularis**, the dilated portion of the urethra in the glans penis. In the vulva, the depression between the posterior commissure and the fourchet. **F. ovalis**, an oval depression in the right auricle of the heart. **F. patellaris**. See *F. Hyaloid*. **F.**, Pituitary, a depression in the sphenoid bone lodging the pituitary body. **F.**, Sublingual, a depression on the internal surface of the inferior maxillary bone, for containing the sublingual gland. **F.**, Submaxillary, the oblong depression on the internal surface of the inferior maxillary bone, containing the submaxillary gland.

Fourchet, or **Fourchette** (*four-shet'*) [*Fr.*, a fork]. 1. A fold of mucous membrane just inside of the posterior commissure of the vulva. 2. A forked instrument used in division of the frenum linguae.

Fourth Cranial Nerve. The pathetic nerve, supplying the superior oblique muscle.

Fovea (*fo-ve'-ah*) [*l.*, a small pit]. A small depression or pit. Applied to many depressions in the body, but more particularly to the fovea centralis retinae, a little pit in the macula lutea, opposite the visual axis, the spot of most distinct vision. **F. hemielliptica**, a small depression on the inner wall of the labyrinth. It is perforated for the passage of filaments of the auditory nerve. **F. hemisphaerica**, a depression in the roof of the labyrinth.

Fowler's Solution. See *Arsenic*.

Fox-glove (*foks'-gluv*). See *Digitalis*.

Fractional (*frak'-shun-al*) [*fractio*, a breaking]. Divided. **F. Cultivation**, the isolation of microorganisms from each other by diluting the mixture containing them to such a degree that a given quantity contains but few organisms. **F. Distillation**. See *Distillation*. **F. Sterilization**, intermittent sterilization.

Fracture (*frak'-tur*) [*frangere*, to break]. A breaking, especially of a bone. **F.**, Barton's. See *Diseases, Table of*. **F.-bed**, a bed designed for patients having fractures. It usually has a hole in the center to transmit the discharges. **F.-box**, a long box without ends or cover, used in the immobilization of fractured legs. **F.**, Comminuted, one in which the bone is splintered. **F.**, Colles'. See *Diseases, Table of*. **F.**, Complete, one in which the bone is entirely broken through. **F.**, Complicated, one associated with injury of adjacent parts. **F.**, Compound, one in which the point of fracture is in communication with the external air through a wound of the overlying parts. **F. by Contre-Coup**, a fracture of the skull caused by transmitted violence, and occurring at a distance from the point struck, usually opposite. **F.**, Depressed, one in which the fractured part is depressed below the normal level, as in fracture of the skull. **F.**, Double, the existence of two fractures in the same bone. **F.**, Greenstick, one in which one side of the bone is broken, the other bent. **F.**, Impacted, one in which one fragment is driven into the other so as to be held fast. **F.**, Pott's, fracture of the fibula about three inches above the ankle-joint, usually with a splitting-off of the tip of the inner malleolus, and at times rupture of the internal lateral ligament. **F.**, Simple, one in which the overlying integument is intact. **F.**, Smith's, fracture of the lower end of the radius, from a quarter of an inch to an inch from the articular surface.

F., Spontaneous, one due to a slight force, as when there is disease of the bone. **F., Ununited**, one in which bony union has failed to occur.

Frænulum (*fren'-u-lum*). See *Frenulum*.

Frænium (*fren'-ium*). See *Frenum*.

Fragilitas (*frag-il'-it-as*) [L.]. Brittleness.

F. crinium, an atrophic condition of the hair in which the individual hairs split into numerous fibrillæ. **F. ossium**, abnormal brittleness of the bones.

Fragmentation (*frag-men-ta'-shun*) [*fragmentum*, a piece]. The breaking up into fragments.

Frambesia (*fram-be'-se-ah*) [*frambesia*, raspberry]. Yaws; Pian. A tropical contagious disease of the skin, of long duration, characterized by dirty or bright-red raspberry-like tubercles, appearing usually on the face, toes, and genital organs. It is most frequent in young negroes.

Frangula (*frang'-gu-lah*) [L.]. The bark of *Rhamnus frangula*, or alder-buckthorn. The young bark is very irritant; bark at least a year old is laxative, resembling rhubarb in action. **F., Ext., Fld.** Dose $\mathfrak{M}\text{x-xx}$ (0.65-1.3). **Ext.** rhamni frangulæ liq. (B. P.). Dose $\mathfrak{M}\text{x-xx}$ (0.65-1.3).

Frankincense (*frangk'-in-sens*) [*francum incensum*, pure incense]. See *Olibanum*. **F., Common**, a concrete turpentine obtained from *Pinus palustris* and other species of *Pinus*.

Franklin Spectacles. [Named after *B. Franklin*, their inventor]. Spectacles with each eye-glass divided horizontally into an upper lens, suited for far vision, and a lower for close work.

Franklinic Electricity. Static or frictional electricity.

Frasera (*fra'-ze-rah*). The root of American Columbo; it is a mild tonic and simple bitter. Dose in powder $\mathfrak{z}\text{j}$ (4.0). Unof.

Fraunhofer's Lines. See *Absorption*.

Fraxinus (*fraks-in'-us*). See *Manna*.

Freckles (*frek'-ls*). See *Lentigo*.

Freezing Microtome. A microtome provided with a contrivance for freezing artificially the tissue to be cut.

Freezing-mixture, a mixture of salt and snow or ice, which absorbs a great deal of heat in undergoing solution.

Freezing-point, the temperature at which a liquid freezes.

Fremitus (*frem'-it-us*) [L., a murmur]. A vibration. **F., Friction**, the vibrations produced by the rubbing together of two dry surfaces, and felt by the hand. **F., Rhonchal**, vibrations produced by the passage of air through a large bronchial tube containing mucus. **F., Tactile**, the vibratory sensation conveyed to the hand applied to the chest of

a person speaking. **F., Vocal**, the sounds of the voice transmitted to the ear applied to the chest of a person speaking.

Frenulum (*fren'-u-lum*) [dim. of *frenum*, a bridle]. A small frenum; a slight ridge on the upper part of the valve of Vieussens. **F. pudendi**, the fourchet.

Frenum (*fren'-ium*) [L., a bridle]. A fold of integument, or mucous membrane, that checks or limits the movements of any organ.

F. linguæ, the vertical fold of mucous membrane under the tongue. **F. of Penis**, the fold on the lower surface of the glans penis connecting it with the prepuce.

Frenzy (*fren'-ze*) [*φρην*, mind]. Violent mania.

Friable (*fri'-ab-l*) [*friare*, to break into pieces]. Easily broken or crumbled.

Friction (*frik'-shun*) [*fricare*, to rub]. The act of rubbing. **F.-fremitus**. See *Fremitus*.

F.-sound, the sound observed in auscultation as a result of the rubbing together of adjacent parts, *e. g.*, of the pleural folds, the pericardium, or the peritoneum, when the layers are dry or roughened.

Friedreich's Ataxia or Disease. See *Diseases, Table of*. **F.'s Sign**. See *Signs and Symptoms, Table of*.

Frigorific (*frig'-or-if'-ik*) [*frigus*, cold; *facere*, to make]. Producing extreme cold.

Frog's Belly. The flaccid abdomen seen in children suffering from rickets or from atony of the abdominal cells, the result of dyspepsia with flatulent distention.

Frommann's Lines. Transverse markings on the axis-cylinder of medullated nerve-fibers, seen after treatment with silver nitrate.

Fron's Reagent. The double iodid of potassium and bismuth, used as a test for alkaloids.

Front-tap Contraction of Gowers. See *Reflexes, Table of*.

Frontal (*frunt'-al*) [*frons*, forehead]. 1. Pertaining to the anterior part or aspect of an organ or body. 2. Belonging to the forehead. **F. Bone**, the anterior bone of the skull and superior bone of the face. **F. Eminence**. See *Eminence*. **F. Lobe**. See *Lobe*. **F. Section**. See *Section*. **F. Sinuses**, the hollow air-spaces in the frontal bone.

Fronto- (*frunt'-to-*) [*frons*, the forehead]. A prefix denoting anterior position, or expressing a relation with the forehead.

Frost-bite, the morbid condition of a part, the result of extreme cold.

Frost-itch, a name for *Prurigo himialis*.

Frosted Feet (*fros'-ted fet*). See *Chilblain*.

Fructose (*fruk'-tös*) [*fructus*, fruit], $\text{C}_6\text{H}_{12}\text{O}_6$. Fruit-sugar, formerly called levulose; it is found in nearly all sweet fruits.

Fruit (*frute*) [*fructus*, fruit]. 1. The developed ovary of a plant, and especially the suc-

culent, fleshy parts gathered about the same.
2. The offspring of animals. **F.**-sugar. See *Fructose*.

Frumentum (*fru-men'-tum*) [L.]. Wheat or other grain. *Spiritus frumenti*, whisky.

Ft. Abbreviation for *fiat* or *fiant*, L. for Let there be made.

Fuchsin (*fook'-sin*) [after Leonhard *Fuchs*], $C_{20}H_{19}N_3, C_2H_4O_2$. The hydrochlorid or acetate of rosanilin, a lustrous, green, crystalline salt imparting an intense red color to solutions. It is employed as a staining agent in microscopy, and has been used internally in albuminuria. Dose gr. $\frac{1}{10}$ - $\frac{1}{4}$ (0.006-0.01).

Fucus (*fu'-kus*) [*φῦκος*, seaweed]. A genus of marine algae, the rock-weeds. **F.** *vesiculosus*, bladder-wrack, sea-wrack. It is employed in goiter and glandular enlargements, and in obesity, under the name of antiflat. Dose of the solid extract, gr. x (0.65); of the fluid extract, f3j-ij (4.0-8.0). Unof.

Fulgorant (*ful'-gu-rant*) [*fulgur*, lightning]. Lightning-like. **F.** Pains, pains that are excruciating and come on with lightning-like suddenness, and disappear as quickly.

Fulminant, or Fulminating (*ful'-min-ant*, or *ful'-min-a-ting*) [*fulminare*, to lighten]. Sudden, severe, and rapid in course; as fulminant glaucoma.

Fumigation (*fu-mig-a'-shun*) [*fumigare*, to smoke]. Disinfection by exposure to the fumes of a vaporized disinfectant.

Fuming (*fu'-ming*) [*fumus*, smoke]. Emitting smoke or vapor.

Function (*fungk'-shun*) [*fungi*, to perform]. The normal or special action of a part.

Functional (*fungk'-shun-al*) [*fungi*, to perform]. Pertaining to the special action of an organ. **F.** Disease, a derangement of the normal action of an organ without structural alteration.

Fundament (*fun'-dam-ent*) [*fundus*, bottom]. 1. The foundation or base of a thing. 2. The rudiment. 3. The breech.

Fundamental (*fun-dam-ent'-al*) [*fundus*, bottom]. Pertaining to the foundation.

Fundus (*fun'-dus*) [*fundus*, the bottom]. The base of an organ; the part farthest removed from the opening of the organ. **F.** oculi, the posterior portion of the interior of the eye seen by the ophthalmoscope. **F.** uteri, the part of the womb remotest from the cervix. **F.** vesicæ, the floor, or bas fond of the urinary bladder.

Fungiform (*fun'-ji-form*) [*fungus*, a mushroom; *forma*, form]. Having the form of a mushroom, as the *F.* papillæ of the tongue.

Fungoid (*fung'-oid*) [*fungus*, a toadstool; *eidōs*, likeness]. Resembling a fungus.

Fungus (*fung'-gus*) [L.]. 1. One of the lowest orders of plants, without stems, leaves,

or roots, and destitute of chlorophyl. The chief classes of fungi are the molds, or hyphomycetes, the yeasts, or saccharomycetes, and the bacteria, or schizomycetes. 2. A spongy, morbid excrescence, as proud flesh. **F.** of Brain, hernia cerebri. **F.** of Dura Mater, a tumor of the skull, of malignant nature, originating in the layers of osteal cells. **F.** Foot. See *Madura-foot*. **F.** hæmatodes, a bleeding tumor, generally a soft carcinoma. **F.**, Ray-, the actinomyces.

Funic (*fu'-nik*) [*funis*, a rope]. Pertaining to the funis.

Funicular (*fu-nik'-u-lar*) [*funis*, a rope]. Relating to the umbilical or spermatic cord.

F. Process, the portion of the tunica vaginalis that surrounds the spermatic cord.

Funiculus (*fu-nik'-u-lus*) [dim. of *funis*, a rope]. 1. A cord-like structure, as the spermatic cord, the umbilical cord. 2. A bundle of nerve-fibers in a sheath of perineurium. 3. A name for the different columns of the spinal cord and medulla oblongata; as the *F.* cuneatus, *F.* gracilis, *F.* of Rolando, *F.* teres. **F.** cuneatus. Column of Burdach. **F.** gracilis. Column of Goll.

Funis (*fu'-nis*) [L.]. A cord; the umbilical cord.

Funnel (*fun'-el*) [Etymology doubtful; Welsh, *ffynel*, an air-hole]. 1. A wide-mouthed, conic vessel ending in an open tube, used to transfer liquids from one vessel to another, and as a support for paper-filters.

Furfuraceous (*fur-fur-a'-se-us*) [*furfur*, bran]. Resembling the scales of bran, as *F.* desquamation.

Furrow (*fur'-o*) [AS., *furh*, a groove]. A groove. **F.**, Digital, one of the transverse lines or furrows on the palmar surface of the fingers. **F.**, Genital, a groove appearing on the genital tubercle of the fetus at the end of the second month.

Furuncle (*fu-rung'-kl*). See *Furunculus*.

Furunculosis (*fu-rung-ku-le'-sis*) [*furunculus*, a boil]. A condition associated with the formation of furuncles.

Furunculus (*fu-rung'-ku-lus*) [L.]. A boil. A local inflammatory affection, commonly involving a skin-gland or hair-follicle, and ending in suppuration. It begins with a painful induration, followed by swelling, suppuration of the corium and subcutaneous connective tissue, and the discharge of a central slough or core. **F.** orientalis, Oriental boil, Aleppo boil, Delhi boil, Biscara button, Pendergore sore, Natal sore. A disease, marked by the successive formation of papule, tubercle, scab, and sharply circumscribed ulcer on the face, especially the cheeks and angle of the mouth. It is common along the shores of the Mediterranean Sea.

Fusel Oil (*fu'-zel*), $C_5H_{11}HO$. Amylic alco-

hol. An oily liquid of strong odor. It is an ingredient of crude alcohol obtained by distilling grain and potatoes.

Fusible (*fu'-sibl'*) [*fusus*, melted]. Capable of being melted. **F.** Calculus, a calculus that can be liquefied by heat.

Fusiform (*fu'-sif-orm'*) [*fusus*, a spindle; *forma*, shape]. Spindle-shaped. **F.** Papillæ, the slender papillæ covering the anterior two-thirds of the tongue. **F.** Lobule, the

convolution on the median aspect of the brain below the collateral fissure.

Fusion (*fu'-shun*) [*fundere*, to pour out]. The process of melting; the act of uniting, or cohering.

Fustigation (*fus-tig-a'-shun*) [*fustigare*, to beat]. Flagellation. **F.**, Electric, an application of electricity in which the surface of the body is rapidly tapped with the electrodes of an induced current.

G

G. Abbreviation of *gram*.

Gadinin (*gad'-in-in*), $C_7H_{17}NO_2$. A ptomain obtained from decomposing haddock, and also from cultures of the bacteria of human feces.

Gaduin (*gad'-u-in*) [*gadus*, the codfish], $C_{55}H_{98}O_8$. A fatty principle occurring in cod-liver oil.

Gadus (*ga'-dus*) [*γάδος*, the whiting]. A genus of fish. **G.** morrhua, the cod; a fish from the livers of which cod-liver oil is obtained.

Gaertner, Duct of. See *Duct*.

Gag [Welsh, *cegio*, to choke]. An instrument placed between the teeth to prevent closure of the jaws.

Gait (*gât*) [Icel. *gata*, a way]. The manner of walking. **G.**, Ataxic, a gait in which the foot is raised high, thrown forward, and brought down suddenly, the whole sole striking the ground at once. **G.**, Cerebellar, a gait associated with a staggering movement. **G.**, Spastic, a gait in which the legs are held close together and move in a stiff manner, and the toes tend to drag and catch. **G.**, Steppage, a gait observed in certain neurotic conditions, in which the foot is thrown forcibly forward, the toe lifted high in the air, the heel being first brought down and then the entire foot.

Galactagogue (*gal-ak'-tag-og*) [*γάλα*, milk; *ἀγωγός*, leading]. 1. Inducing the secretion of milk. 2. An agent that increases the secretion of milk.

Galactidrosis (*gal-ak'-tid-ro'-sis*) [*γάλα*, milk; *ιδρώς*, sweat]. The sweating of a milk-like fluid.

Galactin (*gal-ak'-tin*) [*γάλα*, milk], $C_{54}H_{102}N_2O_{26}$. An amorphous alkaloid found in milk.

Galactoblast (*gal-ak'-to-blast*) [*γάλα*, milk; *βλαστός*, a germ]. A peculiar fat-containing globule found in the acini of the mammary gland.

Galactocoele (*gal-ak'-to-sel*) [*γάλα*, milk; *κύλη*, tumor]. 1. A cystic tumor of the female breast due to a collection of milk resulting from closure of a milk-duct. 2. Hydrocele with milky contents.

Galactometer (*gal-ak'-tom'-et-er*). See *Lactometer*.

Galactophlysis (*gal-ak'-toff'-lis-is*) [*γάλα*, milk; *φλύσις*, eruption]. A vesicular eruption containing a milk-like fluid.

Galactophorous (*gal-ak'-toff'-or-us*) [*γάλα*, milk; *φέρειν*, to bear]. Milk-bearing. **G.** Ducts, the excretory ducts of the mammary.

Galactophygous (*gal-ak'-toff'-ig-us*) [*γάλα*, milk; *φυγή*, flight]. Arresting the secretion of milk.

Galactoplasia (*gal-ak'-to-pla'-ne-ah*) [*γάλα*, milk; *πλάσις*, a wandering]. The metastasis of milk; a disease due to the suppression of lactation and the metastasis of the milk.

Galactopoietic (*gal-ak'-to-poi-et'-ik*) [*γάλα*, milk; *ποιεῖν*, to make]. Galactagogue.

Galactorrhœa (*gal-ak'-tor-el'-ah*) [*γάλα*, milk; *ροία*, a flow]. An excessive flow of milk.

Galactoscope (*gal-ak'-to-sköp*). See *Lactoscope*.

Galactose (*gal-ak'-täs*) [*γάλα*, milk], $C_6H_{12}O_6$. A sugar formed by boiling milk-sugar with dilute acids. It readily reduces alkaline copper-solutions and is fermentable with yeast.

Galactostasis (*gal-ak'-tos'-tas-is*) [*γάλα*, milk; *στάσις*, stoppage]. A stoppage of the milk.

Galactotherapy (*gal-ak'-to-ther'-ap-e*) [*γάλα*, milk; *θεραπεῖν*, treatment]. 1. The treatment of disease in suckling infants by the administration of the remedies to the mother or wet-nurse. 2. Milk-cure.

Galanga, or **Galangal** (*gal-ang'-gah*, or *gal'-an-gal*) [Chin.]. The rhizome of *Alpinia officinarum*, and of *Maranta galganga* (great galangal). The active principles are a

volatile oil and a resin; the actions those of a stimulant aromatic. Dose gr. xv-xxx (1.0-2.0).

Galbanum (*gal'-ban-um*) [Heb., *khelbenah*, white milk]. 1. A gum-resin of *Ferula galbaniflua*, and *F. rubricaulis*. It is expectorant, stimulant, and antispasmodic, and is useful in chronic bronchitis, amenorrhea, and chronic rheumatism. Locally it is employed in the form of a plaster for indolent swellings. Dose gr. x-xx (0.65-1.3). **G.**, **Pil.**, **Comp.**, now termed **Pil. asafoetida** comp.; each pill contains galbanum gr. jss, myrrh gr. jss, asafoetida gr. ss, and syrup. Dose ij-ijj. **G.**, **Emplastrum**, galbanum 16, turpentine 2, Burgundy pitch 6, lead-plaster 76 parts.

Galea (*ga'-le-ah*) [*galea*, helmet]. The aponeurotic portion of the occipito-frontalis muscle.

Gall (*gael*) [AS., *gealla*, bile]. The bile. **G.-bladder**, the pear-shaped sac on the under surface of the right lobe of the liver, constituting the reservoir for the bile. **G.-cyst**, the gall-bladder. **G.-stones**, the concretions occasionally formed in the gall-bladder and bile-ducts.

Galla (*gal'-ah*) [L.]. Nut-gall. An excrescence on the leaves of dyer's oak, *Quercus lusitanica*, caused by the deposited ova of an insect. It contains tannic acid from 10-75 per cent., gallic acid 5 per cent. Dose gr. v-xv (0.32-1.0). **G.**, **Tinct.**, 20 per cent. Dose fʒ ss-ijj (2.0-12.0). **G.**, **Unguentum**, 10 per cent. See *Acid, Tannic*. **G.**, **Unguentum**, cum opio (B.P.), an astringent and sedative ointment.

Gallic Acid (*gal'-ik*). See *Acid, Gallic*.

Gallipot (*gal'-e-pot*) [OD., *gley-pot*, a gallipot]. An apothecary's pot for holding ointments or confections.

Gallon (*gal'-on*) [ME., *galon*; L., *congius*]. A standard unit of volumetric measurement, having in the United States a capacity of 231 cubic inches.

Galvanic (*gal-van'-ik*) [*Galvani*, an Italian

scientist]. Pertaining to galvanism. **G. Battery**. See *Battery*.

Galvanism (*gal'-van-izm*) [*Galvani*, an Italian scientist]. Primary electricity produced by chemic action.

Galvano- (*gal-van'-o-*) [*Galvani*, an Italian scientist]. A prefix denoting a galvanic or primary current of electricity. **G.-cautery**, a form of thermal cautery in which the heat is produced by a galvanic current. **G.-contractility**, the property of being contractile under stimulation by the galvanic current. **G.-faradization**, the simultaneous excitation of a nerve or muscle by both a galvanic and a faradic current. **G.-puncture**, the introduction of fine needles that complete an electric circuit. **G.-therapeutics**, treatment by means of the galvanic current.

Galvanometer (*gal-van-am'-et-er*) [*Galvani*, an Italian scientist; *μετρον*, a measure]. An instrument used for the qualitative determination of the presence of an electric current.

Galvanoscope (*gal-van'-o-skop*) [*Galvani*, an Italian scientist; *σκοπεῖν*, to view]. An instrument for detecting the presence and direction of a galvanic current.

Gamboge (*gam-bōzh'*). See *Cambogia*.

Gammacism (*gam'-as-izm*) [*gammacismus*; *γάμμα*, the letter G]. Difficulty in pronouncing the letters g and k.

Gamogenesis (*gam-o-jen'-es-is*) [*γάμος*, marriage; *γενεσις*, origin]. Sexual reproduction.

Gangliform (*gang'-gle-form*) [*γάγγλιον*, ganglion; *forma*, form]. Having the shape of a ganglion.

Ganglion (*gang'-gle-on*) [*γάγγλιον*, a knot]. 1. A well-defined collection of nerve cells and fibers forming a subsidiary nerve-center. 2. An enlarged bursa in connection with a tendon. **G.-cell**, the large nerve-cell characteristic of the ganglia; similar cells are found in other parts of the nervous system.

Ganglia, Basal, the ganglia at the base of the brain, comprising the corpora striata (caudate and lenticular nuclei), and optic thalami.

TABLE OF GANGLIA.

NAMR.	LOCATION.	ROOTS.	DISTRIBUTION.
Andersch's.	See <i>Petræus</i> .		
Arnold's.	See <i>Otic</i> .		
Auricular.	See <i>Otic</i> .		
Basal.	At base of cerebrum; they include corpora striata and optic thalami.		
Bidder's (2).	In the auricular septum of the frog's heart.	Termination of the cardiac branches of the vagus.	Heart.

TABLE OF GANGLIA.—*Continued.*

NAME.	LOCATION.	ROOTS.	DISTRIBUTION.
Bochdalek's.	Above the root of one of the upper incisor teeth, at the point of communication between the posterior nasal branch of Meckel's ganglion and the anterior dental nerve.	Dental nerves.	Dental nerves.
Cardiac, superior (Wrisberg).	Beneath the arch of the aorta.	Superficial cardiac plexus.	Cardiac plexus.
Carotid.	In the lower part of the cavernous sinus, beneath the carotid artery.	Filaments from the carotid plexus.	Carotid plexus.
Carotid, inferior.	Lower portion of the carotid canal.	Carotid plexus.	Filaments to the carotid artery.
Carotid, superior.	Upper portion of the carotid canal.	Carotid plexus.	Filaments to the carotid artery.
Casserian.	See <i>Gasserian</i> .		
Cephalic.	Sympathetic ganglia of the head. They include the ciliary, sphenopalatine, otic, and submaxillary.		
Cervical, inferior.	Between the neck of the first rib, and the transverse process of the last cervical vertebra.	Three lower cervical, first dorsal.	Cardiac nerves and plexus, etc.
Cervical, middle, or thyroid.	Opposite the fifth cervical vertebra, near the inferior thyroid artery.	Fifth and sixth cervical nerves, spinal nerves and ganglia.	Thyroid gland, cardiac nerve, cavernous plexus.
Cervical, superior.	Opposite the second and third cervical vertebrae.	Four upper cervical, petrosal, vagus, glossopharyngeal, and hypoglossal nerves.	Superior, inferior, external, internal branches; carotid and cavernous plexuses.
Cervical (of uterus).	Near the cervix uteri.	Filaments from the hypogastric plexus, sacral ganglia, and sacral nerves.	Uterine nerves.
Ciliary.	See <i>Ophthalmic</i> .		
Of Cloquet.	Incisor foramen.	Nasopalatine nerves.	Nasopalatine nerves.
Coccygeal.	See <i>Impar</i> .		
Gasser's.	See <i>Gasserian</i> .		
Gasserian, or semilunar.	Fossa on the anterior part of the petrous portion of the temporal bone, near the apex.	Fifth cranial nerve, carotid plexus.	Ophthalmic, superior and inferior maxillary nerves.
Geniculate.	Aqueduct of Fallopius.	Large and small superficial petrosal.	Facial.
Hepatic.	Around the hepatic artery.	Hepatic branches of the semilunar ganglion.	Liver.
Impar, or coccygeal.	Anterior surface of the tip of the coccyx, uniting the two sympathetic nerves.	Sympathetic.	Sympathetic.
Inferior (of vagus).	Near jugular foramen.	Hypoglossal and cervical nerves and various plexuses.	Vagus.
Inframaxillary, anterior.	Near the incisor teeth.	Inferior maxillary nerve.	Filaments to the teeth.

TABLE OF GANGLIA.—*Continued.*

NAME.	LOCATION.	ROOTS.	DISTRIBUTION.
Inframaxillary, posterior.	Near the last molar tooth.	Inferior maxillary nerve.	Filaments to the teeth.
Jugular (Ehrenritter's).	Upper part of the jugular foramen.	Glossopharyngeal.	Continuation of the glossopharyngeal.
Jugular (of vagus).	Jugular foramen.	Vagus.	Vagus.
Ludwig's.	Right auricle of the heart.	Cardiac plexus.	Cardiac plexus.
Lumbar (4 or 5).	On each side and behind the abdominal aorta.	Sympathetic.	Sympathetic.
Meckel's.	See <i>Sphenopalatine</i> .		
Mesenteric, inferior.	The inferior mesenteric artery.	Inferior mesenteric plexus.	Mesentery and intestine.
Mesenteric, lateral.	In connection with superior mesenteric plexus, on left side.	Superior mesenteric plexus.	Mesentery and bowel.
Mesenteric, superior.	Near the origin of the superior mesenteric artery.	Superior mesenteric plexus.	Sympathetic.
Nasal.	See <i>Meckel's</i> .		
Ophthalmic.	Posterior part of the orbit.	Nasal branch of the ophthalmic, third nerve, cavernous plexus, and Meckel's ganglion.	Short ciliary.
Orbital.	See <i>Ophthalmic</i> .		
Otic (Arnold's).	Below the foramen ovale.	Inferior maxillary, auriculotemporal, glossopharyngeal, facial, sympathetic, and internal pterygoid.	Tensor tympani, tensor palati, chorda tympani.
Petrous (Andersch's).	Petrous portion of the temporal bone, lower border.	Glossopharyngeal.	Tympanic, sympathetic, and vagus.
Pharyngeal.	Near ascending pharyngeal artery.	Carotid plexus.	Carotid plexus.
Phrenic.	Under the diaphragm at the junction of the right phrenic nerve and phrenic plexus.	Sympathetic.	To the diaphragm, inferior vena cava, suprarenal capsule, hepatic plexus.
Prostatic (of Müller).	On the prostate.	Prostatic plexus.	Filaments to seminal vesicles and cavernous tissue of penis.
Pterygopalatine.	See <i>Meckel's</i> .		
Remak's.	Tissue of the heart, near the superior vena cava.	Sympathetic.	Sympathetic.
Renal.	Around the renal artery.	Renal plexus.	Renal artery.
Of Ribes.	Anterior communicating artery of the brain. (It is the upper termination of the sympathetic.)	Branches from the two superior ganglia.	Sympathetic.
Sacral (4 or 5).	Ventral surface of the sacrum.	Sympathetic.	Sympathetic.
Scarpa's.	Near the internal auditory meatus, at the point of junction of the facial nerve and the vestibular branch of the auditory nerve.	Facial and auditory nerves.	Internal ear.
Schacher's.	See <i>Ophthalmic</i> .		

TABLE OF GANGLIA.—*Continued.*

NAME.	LOCATION.	ROOTS.	DISTRIBUTION.
Semilunar (2, right and left).	Near the suprarenal bodies, in front of the crura of the diaphragm.	Solar plexus and great splanchnic nerves.	Solar plexus.
Solar.	See <i>Semilunar</i> .		
Sphenopalatine (Meckel's).	Sphenomaxillary fossa, near the sphenopalatine foramen.	Superior maxillary, facial, sympathetic.	Anterior, posterior, and external palatine, nasopalatine, superior nasal, Vidian, pharyngeal.
Splanchnic.	See <i>Semilunar</i> .		
Submaxillary.	Above the submaxillary gland.	Gustatory, chorda tympani, submaxillary, sympathetic.	Mouth, submaxillary gland, and Wharton's duct.
Superior (of vagus).	Jugular foramen.	Superior cervical and petrous ganglia, and spinal accessory.	Vagus.
Suprarenal.	Junction of the great splanchnic nerves.	Solar plexus.	Suprarenal plexus.
Thoracic (12).	Between the transverse processes of the vertebræ and the heads of the ribs.	Sympathetic.	Splanchnic nerves and branches to spinal nerves and plexuses.
Thyroid, inferior.	See <i>Cervical, middle</i> .		
Thyroid, superior.	See <i>Cervical, superior</i> .		
Tympanic.	Canal between lower surface of the petrous portion of the temporal bone and the tympanum.	Tympanic branch of the glossopharyngeal.	Tympanum.
Vagus: (1) of the root, see <i>Jugular</i> ; (2) of the trunk (<i>Vagus</i>).	Below the jugular foramen.	Vagus.	Vagus.
Valentin's.	Above the root of the second bicuspid, at the junction of the middle and posterior dental nerves.	Posterior and middle dental nerves.	Filaments to the teeth.
Ventricular.	See <i>Bidder's</i> .		
Vestibular.	Aqueduct of Fallopius.	Geniculate ganglion.	Vestibular nerve.
Walter's.	See <i>Impar</i> .		
Of Wrisberg.	See <i>Cardiac</i> .		

Ganglionic (*gang-gle-on'-ik*) [γάγγλιον, a ganglion]. Pertaining to or of the nature of a ganglion.

Gangrene (*gang'-gren*) [γάγγραινα, a sore, from γάγρειν, to gnaw]. The putrefactive fermentation of dead tissue. **G.**, Diabetic, a moist gangrene sometimes occurring in diabetic persons. **G.**, Dry, shriveling and desiccation from insufficiency of arterial blood. **G.**, Embolic, that caused by an embolus that cuts off the supply of blood. **G.**, Hospital-, a contagious, rapidly fatal form arising under crowded conditions, par-

ticularly in military hospitals. **G.**, Moist, a form with abundance of serous exudation and rapid decomposition. **G.**, Nosocomial, hospital-gangrene. **G.**, Primary, that without preceding inflammation of the part. **G.**, Pulp. See *G.*, *Hospital*-. **G.**, Secondary, a form with preceding inflammation. **G.**, Senile, that attacking the extremities of the aged. **G.**, Symmetric, that attacking corresponding parts of opposite sides. It is also called Raynaud's disease.

Gangrenous (*gang'-gren-us*) [γάγγραινα,

gangrene]. Pertaining to or of the nature of gangrene. **G. Emphysema.** See *Edema, Malignant*.

Gargle (*gar'-gl'*) [OF., *gargouiller*, to gargle]. A solution used for rinsing the pharynx and nasopharynx.

Garlic (*gar'-lik*). See *Allium*.

Garrot (*gar'-ot*) [Fr., *garotter*, to bind]. An instrument for compression of an artery by twisting a circular bandage about the part.

Gas (*gas*) [a word coined by the Belgian chemist, Van Helmont]. An air-like fluid. The word is especially applied to those fluids that under normal conditions are aeriform; while those that can be readily condensed to liquids are termed vapors. **G., Laughing**, nitrous oxid. **G., Marsh**, methane. **G., Olefiant**, ethylene. **G., Permanent**, a term formerly applied to those gases, as oxygen, nitrogen, hydrogen, that were thought nonliquefiable.

Gaseous (*gas'-e-us*) [see *Gas*]. Of the nature of a gas.

Gasserian Ganglion. See *Ganglia, Table of*.

Gasser's Ganglion. See *Ganglia, Table of*.

Gastro- (*gas'-ter-o-*) [*gastēr*, belly]. See *Gastro-*.

Gastralgia (*gas-tral'-je-ah*) [*gastēr*, stomach; *ἀλγος*, pain]. Paroxysmal pain in the stomach.

Gastrectomy (*gas-trek'-to-me*) [*gastēr*, stomach; *ἐκτομή*, a cutting out]. Excision of the whole or a part of the stomach.

Gastric (*gas'-trik*) [*gastēr*, stomach]. Pertaining to the stomach. **G. Artery.** See *Arteries, Table of*. **G. Crisis**, a severe, paroxysmal attack of pain in the stomach, accompanied by obstinate vomiting, occurring in the course of locomotor ataxia. **G. Juice**, the secretion of the glands of the stomach. It is a clear, colorless liquid, having an acid reaction and a specific gravity of 1002.5, and containing five per cent. of solid matter. A small amount of hydrochloric acid, 0.2 to 0.4 per cent., and a ferment called pepsin, are the essential elements. It digests proteids and precipitates the casein of milk.

Gastritis (*gas-tri'-tis*) [*gastēr*, stomach; *τις*, inflammation]. Inflammation of the stomach. It may be acute or chronic; catarrhal, suppurative or phlegmonous, or diphtheric. **G., Atrophic**, a chronic form with atrophy of the mucous membrane. **G., Hypertrophic**, the early stage of chronic gastritis in which there is a hyperplasia of the mucous membrane. **G. polyposa**, a form of chronic gastritis characterized by a great overgrowth of the connective tissue of the organ, giving rise to polypoid projections of fibrous tissue covered by epithelium.

Gastro- (*gas'-tro-*) [*gastēr*, stomach, belly]. A prefix denoting relation with the stomach.

Gastrocele (*gas'-tro-sel'*) [*gastēr*, stomach; *κῆλη*, hernia]. A hernia of the stomach.

Gastrocnemius (*gas-trok-ne'-me-us*) [*gastēr*, belly; *κνήμη*, leg]. A double-headed muscle forming the greater part of the calf of the leg.

Gastrocolic (*gas-tro-kol'-ik*) [*gastēr*, stomach; *κόλον*, the large intestine]. Pertaining to the stomach and the colon.

Gastrocolitis (*gas-tro-ko-lit'-tis*) [*gastēr*, stomach; *κόλον*, colon; *τις*, inflammation]. Inflammation of the stomach and colon.

Gastrocolpotomy (*gas-tro-kol'-pot'-o-me*) [*gastēr*, belly; *κόπος*, vagina; *τέμνειν*, to cut]. The operation of Cesarean section in which the opening is made through the linea alba into the upper part of the vagina.

Gastrodiaphane (*gas-tro-di'-af-an*) [*gastēr*, stomach; *διά*, through; *φαίνειν*, to show]. See *Gastrodiaphany*.

Gastrodiaphany (*gas-tro-di-af'-an-e*) [*gastēr*, stomach; *διά*, through; *φαίνειν*, to show]. A method of exploration of the stomach by means of an electric lamp known as the gastrodiaphane.

Gastroduodenal (*gas-tro-du-o-de'-nal*) [*gastēr*, stomach; *duodeni*, twelve each]. Pertaining to the stomach and the duodenum.

Gastroduodenitis (*gas-tro-du-o-d-en-i'-tis*) [*gastēr*, stomach; *duodeni*, twelve each; *τις*, inflammation]. Inflammation of the stomach and duodenum.

Gastrodynia (*gas-tro-din'-e-ah*) [*gastēr*, stomach; *ὄδυν*, pain]. Pain in the stomach.

Gastroenteritis (*gas-tro-en-ter-i'-tis*) [*gastēr*, stomach; *έντερον*, bowel; *τις*, inflammation]. Inflammation of stomach and bowels.

Gastroenterostomy (*gas-tro-en-ter-os'-to-me*) [*gastēr*, stomach; *έντερον*, bowel; *στόμα*, mouth]. The formation of a communication between the stomach and the small intestine.

Gastroenterotomy (*gas-tro-en-ter-ot'-o-me*) [*gastēr*, belly; *έντερον*, bowel; *τέμνειν*, to cut]. Incision of the intestines through the abdominal wall.

Gastroepiploic (*gas-tro-ep-ip-lo'-ik*) [*gastēr*, stomach; *ἐπιπλόον*, caul]. Pertaining to the stomach and omentum.

Gastrohepatic (*gas-tro-he-pat'-ik*) [*gastēr*, stomach; *ήπαρ*, the liver]. Relating to the stomach and liver.

Gastrohysterectomy (*gas-tro-his-ter-ek'-to-me*) [*gastēr*, the belly; *ήστέρα*, the womb; *ἐκτομή*, a cutting out]. Removal of the uterus through the abdominal wall.

Gastrohysterotomy (*gas-tro-his-ter-ot'-o-me*) [*gastēr*, belly; *ήστέρα*, womb; *τέμνειν*, to

cut]. Incision of the uterus through the abdominal wall, usually for the purpose of removing a fetus.

Gastrointestinal (*gas-tro-in-test-in-al*) [*γαστήρ*, stomach; *intestine*]. Pertaining to the stomach and intestine.

Gastrolith (*gas'-tro-lith*) [*γαστήρ*, stomach; *λίθος*, a stone]. A calcareous formation in the stomach.

Gastromalacia (*gas-tro-mal-a'-se-ah*) [*γαστήρ*, stomach; *μαλακία*, softening]. An abnormal softening of the walls of the stomach.

Gastromelus (*gas-trom'-el-us*) [*γαστήρ*, belly; *μέλος*, limb]. A monster with accessory limbs attached to the abdomen.

Gastropathy (*gas-trop'-ath-e*) [*γαστήρ*, stomach; *πάθος*, disease]. Any disease or disorder of the stomach.

Gastrophrenic (*gas-tro-fren'-ik*) [*γαστήρ*, stomach; *φρήν*, diaphragm]. Relating to the stomach and diaphragm.

Gastrorrhagia (*gas-tro-raj'-e-ah*) [*γαστήρ*, stomach; *ρηνύναι*, to break forth]. Hemorrhage from the stomach.

Gastrophaphy (*gas-trof'-a-fe*) [*γαστήρ*, stomach; *ῥαφή*, suture]. Suture of a wound of the stomach or abdominal wall.

Gastrorrhea (*gas-tro-r'-e-ah*) [*γαστήρ*, stomach; *ῥοία*, a flow]. Excessive secretion of mucus or gastric juice in the stomach.

Gastroschisis (*gas-tros'-kis-is*) [*γαστήρ*, belly; *σχίσσις*, cleft]. A congenital malformation in which the abdomen remains open.

Gastroscope (*gas'-tro-skôp*) [*γαστήρ*, stomach; *σκοπεῖν*, to see]. An instrument for examining the interior of the stomach.

GastroscoPy (*gas-tros'-ko-pe*) [*γαστήρ*, stomach; *σκοπεῖν*, to see]. The inspection of the interior of the stomach by means of the gastroscope.

Gastrosplenic (*gas-tro-splen'-ik*) [*γαστήρ*, stomach; *σπλήν*, spleen]. Relating to the stomach and the spleen.

Gastrostenosis (*gas-tro-ste-no'-sis*) [*γαστήρ*, stomach; *στενός*, narrow]. A narrowing or stricture of the stomach.

Gastrostomy (*gas-tros'-to-mé*) [*γαστήρ*, stomach; *στόμα*, mouth]. The establishing of a fistulous opening into the stomach.

Gastrotomy (*gas-trof'-o-me*) [*γαστήρ*, stomach; *τέμνειν*, to cut]. Incision of the abdomen or the stomach.

Gastroxia, Gastroxynsis (*gas-troks'-e-ah*, *gas-troks'-in'-sis*) [*γαστήρ*, stomach; *ὀξύς*, acid]. Excessive secretion of hydrochloric acid by the stomach, a condition that characterizes a form of dyspepsia.

Gastrula (*gas'-tru-lah*) [*γαστήρ*, belly]. The embryo at that stage of its development when it consists of two cellular layers formed by the invagination of the blastula.

Gastrulation (*gas-tru-la'-shun*) [*γαστήρ*,

belly]. The process of formation of the gastrula by the invagination of the blastula.

Gathering [*AS.*, *gædrian*, to gather]. A collection of pus beneath the surface.

Gaucher's Disease (*gô-shâs*). So-called primary epithelioma of the spleen.

Gaultheria (*gawt-thel'-re-ah*). Wintergreen, teaberry. The plant, *G. procumbens*, the leaves of which yield a volatile oil, oleum gaultheriæ, which contains 90 per cent. of methyl salicylate, and is used in acute rheumatism, and as a local antiseptic. Dose of oleum gaultheriæ ℥ij-xx (0.2-0.65). *Spiritus gaultheriæ* consists of oil of gaultheria 5, alcohol 95 parts. It is used chiefly as a flavoring agent. Dose ℥x-xx (0.65-1.3).

Gauze (*gawz*) [so called because first imported from *Gaza* in Palestine]. A thin, open-meshed cloth used for surgical dressings. When impregnated with antiseptic substances it is called antiseptic gauze, or, according to the substance used, it is spoken of as iodoform-gauze, sublimate-gauze, etc.

Gavage (*gav-ahzh'*) [*Fr.*]. The administration of liquid nourishment through the stomach-tube.

Gay-Lussac's Laws. See *Law*.

Gelatin (*jel'-at-in*) [*gelare*, to congeal]. An albuminoid substance of jelly-like consistence, obtained by boiling connective tissue in water.

G., Bone, the gelatin extracted from osseous tissue. **G. Capsules**, capsules of gelatin designed for containing medicines of nauseating taste. **G. Culture-medium**, a culture medium for bacteria containing from 8 to 15 per cent. of gelatin, in order to give it a solid consistence. **G., Medicated**, gelatin-discs, or Lamellæ (B.P.), containing traces of alkaloids, for introduction into the conjunctival sac. They are dissolved by the tears, the effects of the alkaloids being thus obtained.

Gelatiniferous (*jel-at-in-if'-er-us*) [*gelatina*, gelatin; *ferre*, to bear]. Producing gelatin.

Gelatinize (*jel'-at-in-iz*) [*gelatina*, gelatin]. To convert into a jelly-like mass.

Gelatinous (*jel-at'-in-us*) [*gelatina*, gelatin]. Resembling or having the nature of gelatin.

Gelose (*jel'-ôz*) [*gelare*, to freeze]. The gelatinizing principle of agar-agar.

Gelsemium (*jel-sem'-e-um*) [*gelseminum*, jasmine]. Yellow jasmine. The root of *G. sempervirens*, the properties of which are mainly due to a bitter alkaloid, gelsemin, $C_{11}H_{19}NO_7$, a powerful motor depressant, antispasmodic, and diaphoretic. In toxic doses it produces diplopia, extreme muscular weakness, and anesthesia, death occurring from asphyxia. Gelsemium is used in neuralgia, especially in migraine, in dysmenorrhea, hysteria, chorea, delirium tremens, and in malarial and typhoid fevers. **G., Ext.,**

Alcoholic (B. P.). Dose gr. $\frac{1}{4}$ – $\frac{1}{2}$ (0.016–0.032). **G.**, **Ext.**, **Fld.** Dose mjj–xx (0.13–1.3). **G.**, **Tinct.** Dose mxx (0.65).

Gelsemin. Dose gr. $\frac{1}{80}$ – $\frac{1}{40}$ (0.001–0.003).

Gemellus (*jem-el'-us*) [dim. of *geminus*, twin]. Applied to one of two muscles, G. superior and G. inferior; also to the gastrocnemius muscle on account of its two heads of origin.

Gemmation (*jem-a'-shun*) [*gemmare*, to put forth buds]. Budding; a mode of reproduction seen in low forms of animal and vegetal life, and characterized by the formation of a small projection from the parent-cell, which becomes constricted off and forms an independent individual.

Gemmule (*jem'-ul'*) [*gemma*, dim. of *gemma*, a bud]. A small bud.

Genera (*jen'-er-ah*) [L.]. Plural of *Genus*.

General (*jen'-er-al*) [*genus*, race]. Common to a class; distributed through many parts; diffuse. **G. Anatomy**, anatomy of the tissues in general, as distinguished from special anatomy, that dealing with special organs. **G. Paralysis**, or **Paresis**. See *Paralysis*. **General**. **G. Pathology**. See *Pathology*.

Generation (*jen'-er-a'-shun*) [*generare*, to beget]. 1. The act of begetting offspring. 2. A period extending from the birth of an individual to the birth of his offspring, usually estimated at a third of a century. **G.**, **Alternate**, the alternation of asexual with sexual generation in the same species of animals or plants, the offspring of one process differing from that of the other. **G.**, **Asexual**, reproduction without previous union of two sexual elements; reproduction by fission or gemmation. **G.**, **Organs of**, those that are functional in reproduction. **G.**, **Sexual**, reproduction by the union of a male and female element. **G.**, **Spontaneous**, the generation of living from nonliving matter.

Genesial, Genesic (*jen'-e'-se-al, jen'-e'-sic*) [*γενεσις*, origin]. Pertaining to generation.

Genesis (*jen'-es-is*) [*γενεσις*, production]. Begetting; development; origin; formation; generation.

Genetic (*jen'-e'-ik*) [*γενεσις*, generation]. Pertaining to generation; producing.

Genial (*je'-ne-al*) [*γενειον*, chin]. Pertaining to the chin. **G. Tubercles**, four prominent tubercles on the internal surface of the lower jaw.

Geniculate, Geniculated (*jen-ik'-u-lat, -ed*) [*genu*, the knee]. Abruptly bent. **G. Bodies**, the corpora geniculata; two oblong, flattened bodies, the external and internal G. bodies, on the posterior inferior part of the optic thalamus. **G. Ganglion**. See *Ganglia*, *Table of*.

Genio- (*je'-ne-o-*) [*γενειον*, the chin]. A prefix denoting connection with the chin. **G.-**

hyoglossus Muscle. See *Muscles*, *Table of*.

G.-hyoid Muscle. See *Muscles*, *Table of*.

Genital (*jen'-it-al*) [*genitalis*, pertaining to generation, from *gignere*, to beget]. Pertaining to the organs of generation or to reproduction. **G. Eminence**, or **Tubercle**, an elevation appearing about the sixth week of embryonic life, in front of the cloaca, and from which the penis or clitoris is developed.

G. Furrow, a furrow extending from the genital eminence of the embryo to the cloaca.

Genitalia (*jen-it-a'-le-ah*) [*genitalis*, pertaining to generation; from *gignere*, to beget]. The organs of generation. In the male these consist of two testicles or seminal glands, with their excretory ducts, the prostate, the penis, and the urethra. The female genitals include the vulva, the vagina, the ovaries, the Fallopian tubes, and the uterus.

Genito- (*jen'-it-o-*) [*gignere*, to beget]. A prefix denoting connection with or relation to the genital organs. **G.-crural**. See *Nerves*, *Table of*. **G.-urinary**, relating to the genitalia and the urinary organs.

Gentian (*jen'-she-an*) [*gentiana*, gentian]. The root of *G. lutea*, containing a neutral bitter principle, gentiopicroin, $C_{20}H_{30}O_{12}$, and gentianin, or gentisin, $C_{44}H_{70}O_{25}$. There are several other species of gentian (*G. purpurea*, *G. catesbaei*), very similar in action to *G. lutea*. Gentian is an excellent tonic, simple bitter. **G.**, **Ext.** Dose gr. j–v (0.065–0.32); dose of the British extract gr. ij–x (0.13–0.65). **G.**, **Ext.**, **Fld.** Dose f3 ss–j (2.0–4.0). **G.**, **Infusum**, **Comp.**, unof. Dose f3 j–f3 j (4.0–32.0). **G.**, **Tinct.**, **Comp.** Dose f3 ss–ij (2.0–8.0). **G.-violet**, a basic anilin dye, staining tissues violet.

Genu (*je'-nu*) [*genu*, the knee]. 1. The knee. 2. Any structure bent like a knee, as the genu of the corpus callosum, or of the optic tract. **G. extrorsum**, out-knee; outward bowing of the knee; bow-leg. **G.-pectoral**, relating to the knee and the chest. **G. recurvatum**, the backward curvature of the knee-joint. **G. valgum**, inward curving of the knee; knock-knee; in-knee. **G. varum**. Same as *G. extrorsum*.

Geophagism (*je'-off'-aj-izm*) [*γη*, earth; *φαγειν*, to eat]. The practice of eating earth or clay.

Geranium (*je'-ra'-ne-um*) [*γερανιον*, geranium]. Cranesbill-root. The root of *G. maculatum*, the properties of which are due to tannic and gallic acids. It is an astringent, useful in diarrhea, etc. **G.**, **Ext.**, **Fld.** Dose f3 ss–j (2.0–4.0).

Gerdy, Fibers of. A transverse band of fibers supporting the web of the fingers.

Gerlier's Disease. See *Diseases*, *Table of*.

Germ (*jern*) [L., *germen*, sprig, offshoot].

1. A portion of matter capable of developing into a living organism; a spore, seed, or embryo. 2. A bacterium. **G.-area**, the spot on an ovum where the development of the embryo begins. **G.-epithelium**, a thickening on the ventromesial aspect of the Wolfian body, giving rise to the male and female sexual elements. **G.-layer**, any one of the layers of a developing embryo. **G.-plasm**, the reproductive or hereditary substance of living organisms, which is passed on from the germ-cell in which an organism originates in direct continuity to the germ-cells of succeeding generations. **G.-theory**, the doctrine of the origin of every organism from a germ or germ-plasm; also the theory that certain diseases are due to the development of micro-organisms in the body.

German (*jer'-man*) [*Germanus*, German]. Pertaining to Germany. **G. Measles**. See *Rubeola*.

Germicidal (*jer'-mis-i-dal*) [*germen*, a germ; *cadere*, to kill]. Destroying germs.

Germicide (*jer'-mis-id*) [*germen*, germ; *cadere*, to kill]. An agent that destroys germs.

Germinal (*jer'-min-al*) [*germen*, a germ]. Pertaining to a germ or to the development of a tissue or organ. **G. Area**, a round spot upon one side of the vitelline membrane in which the development of the embryo begins.

G. Disc. See *Disc*. **G. Membrane**, the blastoderm. **G. Spot**, the nucleolus of the ovule. **G. Vesicle**, the blastodermic vesicle.

Germination (*jer-min-a'-shun*) [*germinatio*, sprouting, budding]. Sprouting of a seed or spore.

Gerontoxon (*jer-on-tok'-on*) [*γέρων*, an old man; *τόξον*, a bow]. The arcus senilis.

Gestation (*jes-ta'-shun*) [*gestare*, to bear]. Pregnancy. **G.**, **Abdominal**, the form of extrauterine gestation in which the product of conception is developed in the abdominal cavity. **G.**, **Double**. 1. Twin pregnancy. 2. The coexistence of uterine and extrauterine pregnancy. **G.**, **Ectopic**. Same as *G.*, *Extrauterine*. **G.**, **Extrauterine**, pregnancy in which the product of conception is not contained in the uterine cavity.

Giant (*ji'-ant*) [*γίγας*, giant]. A being or organism of great size. **G.-cell**. See *Cell*.

Gibbosity (*gib-as'-it-i*) [*gibbus*, a hump]. The condition of being hump-backed.

Gibbous (*gib'-us*) [*gibbus*, a hump]. Hump-backed.

Gibson's Bandage. A bandage for fracture of the lower jaw.

Giddiness (*gid'-e-nes*) [ME., *gidi*, dizzy]. A sensation of whirling or unsteadiness of the body; vertigo.

Gigantoblast (*ji-gan'-to-blast*) [*γίγας*,

giant; *βλαστός*, a germ]. A large nucleated red corpuscle, found in the blood in pernicious anemia.

Gimbernat's Ligament. See *Ligament*.

Gin (*jin*) [OF., *gencvre*, juniper]. Common grain-spirit distilled and flavored with juniper-berries. It is a stimulant and diuretic. *Spiritus juniperi compositus* is its official substitute in U. S. P. **G.-drinker's Liver**. The liver of atrophic cirrhosis.

Ginger (*jin'-jer*). See *Zingiber*.

Gingiva (*jin-jiv'-vah*) [L.]. The gum; the vascular tissue surrounding the necks of the teeth and covering the alveoli.

Gingival (*jin'-jiv-al*) [*gingiva*, the gum]. Pertaining to the gums. **G. Line**, a line along the gums, seen in chronic metallic poisoning, as the blue line of lead.

Gingivitis (*jin-jiv-i'-tis*) [*gingiva*, the gum; *itis*, inflammation]. Inflammation of the gums.

Ginglymus (*ging'-glim-us*). See *Diarthrosis*.

Ginseng (*jin'-seng*) [Chinese, *jintian*, ginseng]. The root of several species of *Panax* or *Aralia*. It has no other medicinal virtues than those of a demulcent, but it has a wonderful reputation in China, to which country most of it is exported.

Giraldensian Organ or **Organ of Giralde's**. Irregular tubules lying in the convolutions of the epididymis. They are the atrophic remains of the tubes of the Wolfian body.

Girdle (*gir'-dl*) [AS., *gyrdel*, a waistband]. A band designed to go around the body; a structure resembling a circular belt or band.

G.-pain, a sensation as if a girdle were drawn tightly around the body. **G.**, **Pelvic**, the bones (the two ossa innominata) forming the support for the lower limbs. **G.-sensation**. Same as *G.-pain*. **G.**, **Shoulder-**, the system of bones supporting the upper limbs or arms.

Gizzard (*gis'-ard*) [L., *gigeria*, the cooked entrails of poultry]. The strong muscular stomach of birds used for triturating the food. A proprietary substance, ingluvin, prepared from it, has been used in dyspepsia.

Glabella (*gla-bel'-ah*) [dim. of *glaber*, smooth]. The smooth triangular space between the eyebrows, just above the root of the nose.

Glacial (*gla'-she-al*) [*glacies*, ice]. Icy; resembling ice in appearance, as glacial acetic or phosphoric acid.

Gladiolus (*glad-e-o'-lus*) [dim. of *gladius*, a sword]. The middle or second piece of the sternum.

Glairin (*glar'-in*) [OF., *glaire*, the white of egg, from *clarus*, clear]. A peculiar organic, gelatinous substance found on the surface of some thermal waters. It is also called baregin.

Glairy (*glâr'-e*) [OF., *glaire*, the white of egg, from *clarus*, clear]. Slimy; albuminous.

Gland [*glans*, an acorn]. 1. An organ which secretes something essential to the system or excretes waste materials the retention of which would be deleterious to the body. The word is also applied to structures which were formerly thought to secrete, as the suprarenal capsules, the pineal gland, etc., but which have no visible secretion. In structure glands may be tubular or racemose, simple or compound. 2. The bulbous end of the penis and clitoris. **G.**, **Absorbent**. See *G.*, *Lymphatic*. **G.**, **Accessory Thyroid**, a small mass of gland-tissue connected with the thyroid gland. **G.**, **Acinous**. See *G.*, *Racemose*. **G.**, **Agminated**. See *G.*, *Peyer's*. **G.**, **Axillary**, the lymph-glands situated in the axilla. **G.** of **Bartholin**, a small gland on either side of the vagina, opening through a duct, on the inner side of the nymphæ; the vulvovaginal gland. **G.**, **Blood**. See *G.*, *Hematopoietic*. **G.**, **Bronchial**, the lymph-glands of the root of the bronchi. **G.**, **Brunner's**, the racemose glands found in the wall of the duodenum. **G.**, **Ceruminous**, the glands secreting the cerumen of the ear. **G.**, **Cervical**, the lymph-glands of the neck. **G.**, **Coccygeal** (or *Luschka's G.*), a small vascular body at the tip of the coccyx. **G.**, **Compound**, one composed of a number of small pouches; a gland the duct of which is branched. **G.**, **Conglobate**. See *G.*, *Lymphatic*. **G.**, **Ductless**, a gland without a duct, as the suprarenal capsule. In reality ductless glands are not true glands. **G.**, **Duodenal**. See *G.*, *Brunner's*. **G.**, **Duverney's**. See *G.* of *Bartholin*. **G.**, **Hematopoietic**, the so-called glands that are supposed to take part in the formation of the blood, as the spleen, thymus, suprarenal capsules, etc. **G.**, **Intestinal**, **Solitary**, the isolated lymph-glands distributed through the intestinal mucous membrane. **G.**, **Lacrymal**, a compound racemose gland in the upper and outer portion of the orbit, the function of which is to secrete the tears. **G.** of *Lieberkühn*. See *Crypts of Lieberkühn*. **G.** of *Littre* (or of *Morgagni*), the small racemose muciparous glands in the mucous membrane of the urethra. **G.**, *Luschka's*. See *G.*, *Coccygeal*. **G.**, **Lymphatic**, small oval masses of lymphatic tissue in the course of lymphatic vessels. Their functions are to act as filters to the blood, retaining foreign particles, and also to form white corpuscles. **G.**, **Mammary**, the glands that secrete milk. **G.**, **Meibomian**, the minute sebaceous follicles between the cartilage and conjunctiva of the eyelids. **G.**, **Montgomery's**, the sebaceous glands of the areola of

the breast. **G.**, **Mucous**, the glands in mucous membranes, secreting mucus. **G.**, **Parotid**, a large salivary gland situated in front of the ear. **G.**, **Peyer's**, a well-defined collection of solitary lymphatic glands, varying in length from 1 to 4 inches, and situated chiefly in the ileum. **G.**, **Pituitary**, a term for the hypophysis of the brain. **G.**, **Prostate**. See *Prostate Gland*. **G.**, **Pyloric**, the glands of the stomach situated near the pylorus and secreting pepsin. **G.**, **Racemose**, a gland composed of a number of acini communicating with several excretory ducts, which usually join to form a common duct. **G.**, *Rivini's*. See *G.*, *Sublingual*. **G.**, **Salivary**, a gland that secretes saliva. **G.**, **Sebaceous**, the glands in the corium of the skin, secreting sebum. **G.**, **Seminal**, the testicle. **G.**, **Serous**, a secreting gland, the cells of which are granular and spheric in form, with central nuclei, and which secrete a thin watery fluid. **G.**, **Sublingual**, the smallest of the salivary glands, situated one on either side beneath the tongue. **G.**, **Submaxillary**, a salivary gland situated below the angle of the jaw. **G.**, **Sudoriparous**, the convoluted glands in the skin that secrete the sweat. **G.**, **Thymus**. See *Thymus*. **G.**, **Thyroid**. See *Thyroid*. **G.**, **Tubular**, a gland having a tube-like structure. **G.**, **Tubular, Compound**, one composed of a number of small tubules with a single duct. **G.**, **Urethral**. See *G.* of *Littre*. **G.**, **Vaginal**, one of the glands in the vaginal mucous membrane. **G.**, **Vulvovaginal**. See *G.* of *Bartholin*.

Glanderous (*glân'-der-us*) [*glans*, acorn]. Affected with glanders.

Glanders (*glân'-ders*) [*glans*, acorn]. A contagious disease of horses and asses, but communicable to man, and due to the bacillus of glanders or bacillus mallei. It appears in two forms—as glanders proper, when affecting the mucous membranes, and as farcy, when limited to the skin and lymphatic glands. On mucous membranes, especially the nasal, it manifests itself as isolated nodules which coalesce and break down into deep ulcers that involve the cartilages and bones. It is apt to extend down to the lungs and give rise to suppuration and pneumonic processes. In man the disease usually runs an acute, febrile course, typhoid in type, and terminates fatally. Farcy is characterized by nodules (farcy-buds) in the skin and lymphatic glands, which break down into irregular chronic ulcers.

Glandula (*glând'-u-lah*) [L.]. A little gland.

Glandular (*glând'-u-lar*) [*glandula*, a little gland]. Relating to, or of the nature of, a gland.

Glans (*glans*) [*glans*, an acorn]. An acorn-shaped body. **G. clitoridis**, the rounded end of the clitoris analogous to the glans penis of the male. **G. penis**, the conical-shaped body forming the head of the penis.

Glass (*glas*) [A.S., *glæs*, glass]. 1. A brittle, hard, transparent substance, consisting usually of the fused amorphous silicates of potassium and calcium, or sodium and calcium, with an excess of silica. 2. Any article made from glass. **G., Crown**, a very hard glass, is a silicate of sodium and calcium. **G., Flint**, that composed of lead and potassium silicates. **G., Soluble**, potassium or sodium silicate, used as a substitute for plaster of Paris.

Glasses (*glas'es*) [A.S., *glæs*, glass]. A synonym of spectacles or eye-glasses. **G., Bifocal**, those that have a different refracting power in the upper part from that in the lower. **G., Prismatic**, those formed of prisms; used in insufficiency and paralysis of the ocular muscles.

Glauber's Salt (*glaw'-berz*) [from *Glauber*, a German chemist]. Sodium sulphate.

Glaucoma (*glaw'-ko'-mah*) [*γλαυκός*, sea-green; *ōma*, tumor]. A disease of the eye characterized by heightened intraocular tension, resulting in hardness of the globe, excavation of the papilla or optic disc, a restriction of the field of vision, corneal anesthesia, colored halo about lights, and lessening of visual power that may proceed to blindness. The etiology is obscure. **G. absolutum**, or **consummatum**, the completed glaucomatous process when the eyeball is exceedingly hard and totally blind. **G. hæmorrhagicum**, or **apoplecticum**, that associated with retinal hemorrhage. **G., Secondary**, that consequent upon other ocular diseases. **G. simplex**, that form without inflammatory symptoms.

Glaucomatous (*glaw'-kom'-at-us*) [*γλαυκός*, sea-green]. Affected with or pertaining to glaucoma.

Gleet (*glet*) [A.S., *glidan*, to glide]. The chronic stage of urethritis characterized by a slight mucopurulent discharge.

Gleety (*gle'-te*) [A.S., *glidan*, to glide]. Resembling the discharge of gleet.

Glénard's Disease. See *Diseases*, *Table of*.

Glenoid (*gle'-noid*) [*γλήνη*, a cavity; *ειδός*, form]. Having a shallow cavity; resembling a shallow cavity or socket. **G. Cavity**, the depression in the scapula for the reception of the head of the humerus. **G. Fossa**, a depression in the temporal bone for articulation with the condyle of the lower jaw.

Glia (*gli'-ah*) [*γλία*, glue]. The neuroglia. **Gliadin** (*gli'-ad-in*) [*γλία*, glue]. A proteid found in wheat-gluten.

Glioma (*gli'-o'-mah*) [*γλία*, glue; *ōma*, a tu-

mor]. A tumor composed of neuroglia-cells and occurring in the brain, spinal cord, retina, nerves, and suprarenal capsules. In the brain it closely resembles the brain-substance, but is usually more gelatinous and darker. In the retina it is often combined with sarcoma (gliosarcoma). It may also be combined with fibroma, myxoma, and neuroma. The last combination is known as neuroglioma ganglionare.

Gliomatosis (*gli'-o'-mat'-o'-sis*) [*γλία*, glue; *ōma*, a tumor]. The development of exuberant masses of glioma-like tissue in the nerve-centers. It is seen in the spinal cord in some cases of syringomyelia.

Gliomatous (*gli'-o'-mat-us*) [*γλία*, glue; *ōma*, a tumor]. Of the nature of, or affected with, glioma.

Glisson's Capsule. See *Capsule*.

Globin (*glo'-bin*) [*globus*, a globe]. A proteid derived from hemoglobin.

Globule (*glob'-ul*) [dim. of *globus*, a ball]. A small spheric particle, as a blood-corpuscle or lymph-corpuscle. Also a small pill or pellet.

Globulicidal (*glob-u-lis-i'-dal*) [*globulus*, a little ball; *cadere*, to kill]. Destructive to the blood-corpuscles.

Globulin (*glob'-u-lin*) [*globulus*, a little ball].

1. A general name for various proteids comprising globulin, vitellin, paraglobulin, or serum-globulin, fibrinogen, myosin, and globin, which differ from the albumins in not being soluble in water, but soluble in dilute neutral saline solutions. These solutions are coagulated by heat, and precipitated by a large amount of water. 2. Specifically, a proteid found in the crystalline lens.

Globulinuria (*glob-u-lin-ū'-re-ah*) [*globulus*, a little ball; *ουρον*, urine]. The presence of globulin in the urine.

Globus (*glo'-bus*) [L.]. A ball or globe.

G. hystericus, the "lump" or choking sensation occurring in hysteria, caused probably by spasmodic contraction of the esophageal and pharyngeal muscles. **G. major**, the larger end or head of the epididymis. **G. minor**, the lower end of the epididymis.

Glomerate (*glom'-er'-at*) [*glomerare*, to wind around]. Rolled together like a ball of thread.

Glomerulitis (*glom-er-u-lit'-tis*) [*glomerulus*, a small ball; *ιτις*, inflammation]. Inflammation of the glomeruli of the kidney.

Glomerulonephritis (*glom-er-u-lo-nef-rit'-tis*) [*glomerulus*, a little ball; *νεφρός*, the kidney; *ιτις*, inflammation]. Inflammation of the Malpighian bodies of the kidney.

Glomerulus, or **Glomerule** (*glom-er'-u-lus*, or *glom'-er-ul*) [dim. of *glomus*, a ball]. 1. A small rounded mass. 2. A coil of blood-vessels projecting into the expanded end

(Bowman's capsule) of each uriniferous tubule and with it composing the Malpighian body.
Glonin (*glo'-no-in*) [from Gl = glyceryl; O = oxygen; N = nitrogen, in the formula $\text{GlO}_2(\text{NO})_2$, in which Gl stands for glyceryl]. Nitroglycerin.

Glossal (*glos'-al*) [*γλωσσα*, tongue]. Pertaining to the tongue.

Glossalgia (*glos-al'-je-ah*) [*γλωσσα*, tongue; *ἀλγος*, pain]. Pain in the tongue.

Glossitis (*glos-i'-tis*) [*γλωσσα*, tongue; *τις*, inflammation]. Inflammation of the tongue.

Glossopiglottidean (*glos-o-ep-e-glot-id'-e-an*) [*γλωσσα*, tongue; *ἐπι*, upon; *γλωττις*, glottis]. Pertaining to both tongue and epiglottis.

Glossograph (*glos'-o-graf*) [*γλωσσα*, tongue; *γραφειν*, to write]. An instrument for registering the movements of the tongue in speech.

Glossohyal (*glos-o-hi'-al*) [*γλωσσα*, tongue; *ὑοειδής*, the hyoid bone]. Pertaining to the tongue and the hyoid bone.

Glossolabiolaryngeal Paralysis (*glos-o-lab-o-lar-in'-je-al*). See *Palsy, Bulbar*.

Glossology (*glos-o'-lo-je*) [*γλωσσα*, tongue; *λογία*, a treatise]. A treatise concerning the tongue.

Glossopathy (*glos-op'-ath-e*) [*γλωσσα*, tongue; *πάθος*, disease]. Any disease of the tongue.

Glossopharyngeal (*glos-o-far-in'-je-al*) [*γλωσσα*, tongue; *φάρυγξ*, pharynx]. 1. Pertaining to the tongue and the pharynx.

2. Pertaining to the glossopharyngeal nerve.

Glossophytia (*glos-o-f'-te-ah*) [*γλωσσα*, tongue; *φυτόν*, a plant]. A dark discoloration of the tongue, due to the accumulation of spores and dead epithelium; black tongue.

Glossoplegia (*glos-o-ple'-je-ah*) [*γλωσσα*, tongue; *πληγή*, stroke]. Paralysis of the tongue.

Glossospasm (*glos'-o-s-pazm*) [*γλωσσα*, tongue; *σπασμός*, spasm]. Spasm of the tongue.

Glossy Skin. A peculiar shining condition of the skin, due to trophic changes following injury or disease of the cutaneous nerves.

Glottis (*glot'-is*) [*γλωττις*, glottis]. The space between the vocal bands.

Glove-area. The area of anesthesia of the fingers, hand, and forearm, in multiple neuritis. It corresponds to the region of skin covered by gloves of various lengths.

Glovers' Stitch. The continuous suture used especially in repairing wounds of the intestines.

Glucose (*glu'-kōs*) [*γλυκίς*, sweet], $\text{C}_6\text{H}_{12}\text{O}_6$. Grape-sugar; dextrose. 1. A form of sugar found in many fruits, in blood and lymph, and in the urine in diabetes. It is crystalline; its solution turns the plane of polarized light to the right; it is less soluble and less sweet than cane-sugar, and ferments readily.

It can be obtained from starch by the action of diastatic ferments, or by boiling with dilute mineral acids, and crystallizes in nodular masses, melting at 86°F . 2. A generic name for a class of carbohydrates having the composition $\text{C}_6\text{H}_{12}\text{O}_6$, and of which ordinary glucose is the type.

Glucosid (*glu'-ko-sid*) [*γλυκίς*, sweet]. Any member of a series of compounds that may be resolved by an acid into glucose and another principle. The more important ones are amygdalin, arbutin, myronic acid, and salicin.

Glucosin (*glu'-ko-sin*) [*γλυκίς*, sweet]. Any one of a series of bases obtained by the action of ammonia on glucose.

Glue (*glu*) [Low L., *glutem*, accus. of *glus*, glue]. An impure gelatin prepared by boiling the skin, hoofs, and horns of animals. It is a very adhesive substance, and when cold, holds firmly together the surfaces between which it is placed.

Gluge's Corpuscles (*glu'-ges*) [*Gluge*, a professor of the University of Brussels]. Compound granular corpuscles; compound granule-cells occurring in tissues that are the seat of fatty degeneration.

Glutamic Acid (*glu-tam'-ik*), $\text{C}_5\text{H}_9\text{NO}_4$. A crystalline acid found in gluten and other proteids.

Glutaric Acid (*glu-tar'-ik*), $\text{C}_5\text{H}_8\text{O}_4$. A crystalline acid found in decomposed pus.

Gluteal (*glu-te'-al*) [*γλουτός*, the buttock].

Pertaining to the buttocks. **G. Artery**. See *Arteries, Table of*. **G. Muscles**. See *Muscles, Table of*. **G. Nerve**. See *Nerves, Table of*. **G. Reflex**. See *Reflexes, Table of*.

Gluten (*glu'-ten*) [*gluten*, glue]. A nitrogenous substance found in the seed of cereals. It consists mainly of gluten-fibrin, gluten-casein, gliadin, and mucedin. **G.-bread**, bread made from wheat-flour from which all the starch has been removed; it is used as a substitute for ordinary bread in diabetes.

Gluteofemoral (*glu-te-o-fem'-or-al*) [*γλουτός*, the buttock; *femur*, the femur]. Relating to the buttock and the thigh.

Gluteus (*glu-te'-us*) [*γλουτός*, the buttock]. One of the large muscles of the buttock. See *Muscles, Table of*.

Glutin (*glu'-tin*). 1. See *Gelatin*. 2. Synonym of gluten-casein. See *Gluten*.

Glycerid (*glis'-er-id*) [*γλυκερός*, sweet]. A compound of glycerol and an acid; the neutral fats are glycerids.

Glycerin, Glycerinum (*glis'-er-in, glis-er-i-num*) [*γλυκίς*, sweet]. 1. See *Glycerol*. 2. In the B. P., a solution of a medicinal substance in glycerol; a glycerite. **G., Suppositoria**, U. S. P., each contains six grams of glycerol; they are used in constipation.

Glycerite, or Glyceritum (*glis'-er-it, or glis-er-i'-tum*) [*γλυκίς*, sweet]. A mixture of

medicinal substances with glycerin. The following glycerites are official: *G. acidi carbolici*; *G. acidi tannici*; *G. amyli*; *G. boro-glycerini*; *G. hydrastis*; *G. vitelli*.

Glycerol (*glis'er-ol*) [*γλυκός*, sweet], $C_3H_5(OH)_3$. Glycerin. A colorless substance, of syrupy consistence, sweetish to the taste, obtained from fats and fixed oils. Chemically it is a triatomic alcohol, and may be looked upon as propenyl alcohol. It is soluble in water and in alcohol; and has a specific gravity of 1.25 at 15° C. It is used as a vehicle in pharmaceutical preparations, as an emollient application to chaps of the skin, as a laxative administered by the mouth or in suppository, for tampons in pelvic congestion, as a substitute for sugar in diabetes, as a mounting-medium in microscopy, and as an addition to bacteriologic culture-media.

Glyceryl (*glis'er-il*) [*γλυκός*, sweet]. The trivalent radicle, C_3H_5 , of glycerol, combining with the fatty acids to form the neutral fats.

Glycin (*gli'sin*). Synonym of *Glycocol*.

Glycocholic Acid (*gli-ko-kol'-ik*) [*γλυκός*, sweet; *χολή*, bile]. An acid found in the bile. See *Acid*.

Glycocol (*gli'-ko-kol*) [*γλυκός*, sweet; *κόλλα*, glue], $C_5H_9NO_3$. Also termed glycin, gelatin-sugar, or amidoacetic acid. It is obtained when glycocholic acid is boiled with caustic potash, baryta-water, or with dilute mineral acids; also by boiling gelatin with dilute acids. It is capable of acting as a base and as an acid.

Glycogen (*gli'-ko-jen*) [*γλυκός*, sweet; *γεννάν*, to produce], $6(C_6H_{10}O_5) + H_2O$. A carbohydrate found in the form of amorphous granules in the liver-cells, in all tissues of the embryo, in the testicle, muscles, leukocytes, fresh pus-cells, cartilage, and other tissues. It is formed from carbohydrates and probably also from proteids, and is stored in the liver, where it is converted, as the system requires, into sugar (glucose). Glycogen is soluble in water, dextrorotatory, and is colored red by iodine.

Glycogeny (*gli-koj'-en-e*) [*γλυκός*, sweet; *γένεσις*, production]. The normal production of glycogen.

Glycol (*gli'-kol*) [*γλυκός*, sweet]. A diatomic alcohol; a compound intermediate in its properties and chemic relations between monohydric alcohol and trihydric glycerol.

Glycolytic (*gli-kol-il'-ik*) [*γλυκός*, sweet; *λυσίς*, dissolution]. Splitting up glucose.

Glyconin (*gli'-ko-nin*) [*γλυκός*, sweet]. The Glyceritum vitelli of the U. S. P. It is a mixture of yolk of egg 45, and glycerol 55 parts.

Glycosuria (*gli-ko-su'-re-ah*) [*γλυκός*, sweet; *ούρον*, urine]. The presence of grape-sugar in the urine.

Glycuronic Acid (*gli-ku-ran'-ik*). See *Acid*.

Glycyrrhiza (*glis-ir-i'-zah*) [*γλυκός*, sweet; *ρίζα*, root]. Liquorice-root. The root of *G. glabra*, a demulcent and mild laxative, of sweet taste. It is used in catarrhal affections, and as an ingredient of pills. *G.*, *Ext.*, *Liquid* (B. P.). Dose f 3j-ij (4.0-8.0). *G.*, *Ext.*, *Fld.*, a vehicle for administering quinin. *G.*, *Ext.*, *Purum*. This is used for making pills. *G.*, *Mist.*, *Comp.*, Brown mixture. Dose f 3 ss-f 3j (16.0-32.0). *G. et opii*, *Trochisci*. Dose 1-2. *G.*, *Pulv.*, *Comp.*, compound liquorice-powder. Dose 3j (4.0).

Glycyrrhizinum (*glis-ir-is-i'-num*) [*γλυκός*, sweet; *ρίζα*, root]. The active principle of liquorice-root. It is in reality an acid, glycyrrhizic acid, $C_{42}H_{68}NO_{16}$. *G. ammoniatum* (U. S. P.), ammoniated glycyrrhizin, a sweet preparation used as a substitute for liquorice. Dose gr. v-xv (0.32-1.0).

Gmelin's Test. Same as *Gmelin-Heintz' Reaction*.

Gmelin-Heintz' Reaction. A test for bile-pigments, consisting in the development of a series of spectral colors, green, blue, and violet, when strong nitric acid is added to a solution containing bile-pigments.

Gnathic (*na'-thik*) [*γνάθος*, jaw]. Pertaining to the jaw. *G. Index*. See *Index*.

Goa-powder. See *Araroba*.

Goblet-cells. Beaker-shaped cells found on mucous membranes.

Goggles (*gog'-lz*) [Irish and Gael., *gog*, a nod]. Spectacles with colored lenses and wire or cloth sides, to protect the eyes from excessive light or dust.

Goiter (*goi'-tr*) [*guttur*, throat]. Enlargement of the thyroid gland. It is also called bronchocoele, tracheocoele, and Derbyshire neck. *G.*, *Exophthalmic*, Basedow's disease, Graves' disease.

Gold [AS., *gold*, gold]. See *Aurum*. *G.-beater's Skin*, a thin membrane prepared from the cecum of the ox.

Golden Seal. See *Hydrastis*.

Goll, Columns of. See *Column*.

Gomphosis (*gom-fol'-sis*) [*γόμφος*, nail]. See *Synarthrosis*.

Gonagra (*gon-a'-grah*) [*γόναυ*, knee; *ἀγρα*, seizure]. Gout of the knee-joint.

Gonarthrit (*gon-ar-thri'-tis*) [*γόναυ*, knee; *ἀρθρον*, a joint; *ιτις*, inflammation]. Inflammation of the knee-joint.

Gonocyst (*gon'-e-sit*) [*γονή*, semen; *κύστις*, cyst]. A seminal vesicle.

Gonocystitis (*gon-e-sis-ti'-tis*) [*γονή*, semen; *κύστις*, cyst; *ιτις*, inflammation]. Inflammation of the seminal vesicles.

Gonepoietic (*gon-e-poi-et'-ik*) [*γονή*, semen; *ποιεῖν*, to make]. Pertaining to the secretion of semen.

Gonion (*go'-ne-on*) [*γωνία*, an angle]. The angle of the lower jaw.

Gonococcus (*gon-o-kok'-us*) [*γωνή*, semen; *κόκκος*, kernel]. The organism causing gonorrhea. See *Micrococcus gonorrhea*, *Bacteria*, *Table of*.

Gonorrhea (*gon-or-el'-ah*) [*γωνή*, semen; *ῥοία*, a flow]. A specific infectious inflammation of the mucous membrane of the urethra and adjacent cavities, due to the gonococcus of Neisser. The disease is characterized by pain, burning urination, a profuse mucopurulent discharge, and a protracted course. It is apt to become chronic, and is frequently accompanied by complications—prostatitis, periurethral abscess, epididymitis, cystitis, purulent conjunctivitis. It may also cause arthritis (gonorrheal rheumatism), endocarditis; and in women, salpingitis. **G.**, **Dry**, a form unassociated with discharge.

Gonorrheal (*gon-or-el'-al*) [*γωνή*, semen; *ῥοία*, a flow]. Relating to gonorrhea, as **G. ophthalmia**. **G. Rheumatism**, inflammation of one or more joints as a sequel of gonorrhea.

Goose [*AS.*, *gos*, goose]. A bird of the family of *Anserines*. **G.-flesh**, **G.-skin**, *cutis ansera*; a condition of the skin marked by prominence about the hair-follicles.

Gorget (*gor'-jet*) [*gorges*, a chasm]. A channeled instrument similar to a grooved director used in lithotomy.

Gossypium (*gos-ip'-e-um*) [*L.*]. The Cotton-tree, *Gossypium herbaceum*, and other species of *Gossypium*, of the order *Malvaceae*. **G.** *purificatum*, cotton-wool, the hairs of the seed of *Gossypium herbaceum*, and of other species of *Gossypium*, used as a dressing and as a substitute for sponges in surgery; in pharmacy, as a filtering medium. See *Cotton*. **Gossypii radices cortex**; this is used as an emmenagogue, especially in the form of **G. radices**, *Ext.*, *Fld.* Dose [*ss*—*f 3 j*] (2.0—4.0). **G. seminis, Oleum**, cotton-seed oil. It is used in ointments.

Goulard's Cerate. See *Plumbum*. **G.'s Extract**. See *Plumbum*.

Gout (*gout*) [*gutta*, a drop]. A disease characterized by a paroxysmal painful inflammation of the small joints, particularly the great toe, accompanied by the deposit of sodium urate. The attack usually comes on at night, is attended by a dusky, glazed swelling of the joint and agonizing pain, and disappears with a sweat in the morning, to recur again at night. In some cases gout presents an atypical form, appearing as dyspepsia, bronchitis, or intestinal catarrh; at times it produces pneumonia and inflammation of the serous membranes. The cause of gout is not definitely known, but is connected with an excess of uric acid or urates

in the blood. The disease is most common in high livers. **G.**, **Latent** or **Masked**, lithemia, a condition ascribed to a gouty diathesis, but not presenting the typical symptoms of gout. **G.**, **Retrocedent**, that form that presents severe internal manifestations, without the customary arthritic symptoms. **G.**, **Rheumatic**. See *Arthritis*, *Rheumatoid*.

Gouty (*gow'-te*) [*gutta*, a drop]. Of the nature of gout; affected with gout. **G. Kidney**, chronic interstitial nephritis due to gout.

Graafian Follicles, or **Vesicles** [Ragner de Graaf, a Dutch anatomist]. Vesicular bodies existing in the ovary, and each containing an ovum.

Gracilis (*gras'-il-is*) [*L.*, slender]. See *Muscles*, *Table of*.

Graduate (*grad'-u-at*) [*gradus*, a step]. A vessel upon which the divisions of liquid measure have been marked.

Graduated (*grad'-u-a-ted*) [*gradus*, a step]. Arranged in degrees or steps. **G. Compress**, a compress made of pieces decreasing progressively in size, the apex or smallest piece being applied to the focus of pressure.

Graefe's Knife [*gra-fe*]. A narrow knife for the performance of the operation for cataract. **G.'s Signs or Symptoms**. See *Signs and Symptoms*, *Table of*.

Graft [*ME.*, *graffe*, graft, from *γρᾶναι*, to write]. A small portion of skin, bone, periosteum, nerve, etc., used to replace a defect in a corresponding structure. **G.**, **Sponge**, the insertion of a piece of sponge into the tissues to act as a framework for granulations.

Grain (*gran*) [*granum*, grain]. 1. Seed, as that of the cereals. 2. A body resembling a seed, as a starch-grain. 3. The unit of weight of the Troy and avoirdupois system of weights. See *Weights and Measures*. **G. of Paradise**, the unripe fruit of *Amomum melegueta*, and of *A. granum paradisi*, brought from W. Africa. It is an aromatic stimulant and diuretic, useful in some cases of neuralgia. *Unof.*

Gram, **Gramme** (*gram*) [*γράμμα*, inscription]. The gravimetric unit of the metric system of weights and measures, equivalent to the weight of a cubic centimeter of distilled water at its maximum density.

Gram's Method. A method for staining bacteria. The bacteria on the cover-glass or in the section are stained first with Ehrlich's solution, and then are treated with Gram's solution (iodin 1, potassium iodid 2, water 300), and then with alcohol. Some bacteria give up the color when washed with alcohol.

Granatum (*gran-a'-tum*). The bark of the stem and root of *Punica granatum*. The bark contains punicotannic acid and mannite,

but the active principle is the alkaloid pelletierin, $C_8H_{13}NO$. The chief use of pomegranate and its preparations is as a tennicide. **G.**, Decoctum; this consists of $\frac{3}{4}$ ij of bark in Oij of water, boiled down to a pint. Pelletierin tannate, dose gr. $\frac{1}{4}$ – $\frac{3}{4}$ (0.032–0.048) (from gr. v–viii (0.32–0.52), according to some authorities).

Grand Mal [Fr., great evil]. Epilepsy. See, also, *Petit Mal*.

Granular (*gran'-u-lar*) [*granum*, a grain]. Made up of, or containing, granules. **G.** Lids, trachoma. **G.** Pharyngitis, pharyngitis characterized by the presence of prominent follicles.

Granulation (*gran-u-lat'-shun*) [*granula*, dim. of *granum*, a grain]. 1. A capillary loop of blood-vessels surrounded by a group of connective-tissue cells. 2. Also the process by which these are formed. 3. The formation of new or cicatricial tissue in the repair of wounds or ulcers, the surface of which has a granular appearance; also, any one of the elevated points of such a surface or formation. **G.-tissue**, the material consisting of granulations by which the repair of loss of substance or the healing together of surfaces is brought about.

Granule (*gran'-ul*) [*granula*, a little grain]. A small grain, body, or particle, as the granules of a cell; also a small pill. **G.-layer**. See *Retina*.

Granuloma (*gran-u-lø'-mah*) [*granulum*, a small grain; *δμα*, a tumor]. A tumor or tumor-like nodule made up of granulation-tissue. **G.**, Infectious, that due to a specific microorganism, as tubercle, gumma, etc.

Granulose (*gran'-u-løs*) [*granula*, a little grain]. The material that forms the inner portion of starch-granules.

Granum (*gra'-num*). See *Grain*.

Grape-cure (*grāp'-kūr*). A treatment of pulmonary tuberculosis, consisting in the ingestion of large quantities of grapes. **G.-sugar**. See *Glucose* and *Dextrose*.

Graphite (*graf'-it*) [*γράφειν*, to write]. Plumbago, or black lead, an impure allotropic form of carbon. It has been applied externally in skin diseases.

Graphospasm (*graf'-o-spazm*) [*γράφω*, writing; *σπασμός*, spasm]. Writers' cramp.

Grating (*gra'-ting*) [*grata*, a grating]. 1. A frame or screen composed of bars. 2. A sound produced by the friction of very rough surfaces against each other. 3. A glass ruled with exceedingly fine parallel lines to produce chromatic dispersion in the rays of light reflected from it.

Grattage (*grah-ta(h)zh'*) [Fr.]. A method of removing morbid growths, as polypi or trachomatous granulations, by rubbing with a harsh sponge or brush.

Gravel (*grav'-el*) [Fr., *gravelle*, from Bret., *grouan*, gravel]. A granular, sand-like material forming the substance of urinary calculi, and often passed in the form of detritus with the urine.

Graves's Disease. See *Diseases, Table of*.

Gravid (*grav'-id*) [*gravare*, to load]. Pregnant. **G. Uterus**, the womb during pregnancy.

Gravida (*grav'-id-ah*) [*gravidus*, pregnant]. A pregnant woman.

Gravimetric (*grav-e-met'-rik*) [*gravis*, heavy; *μέτρον*, measure]. Pertaining to measurement by weight. **G. Analysis**, an analysis in which the component substances are weighed.

Gravity (*grav'-it-e*) [*gravis*, heavy]. Weight. **G.**, Specific, the measured weight of a substance compared with that of an equal volume of another taken as a standard. For gaseous fluids, hydrogen is taken as the standard; for liquids and solids, distilled water at its maximum density.

Gray (*gra*) [AS., *græg*, gray]. Of a color between white and black. **G. Hepatization**. See *Hepatization*. **G. Matter**, that forming the outer part of the brain and the inner part of the cord, containing the specialized cells of these parts. **G. Powder**, hydrargyrum cum creta. See *Mercury*.

Green (*grēn*) [ME., *grene*, green]. Of the color of grass. **G. Blindness**, a variety of color-blindness in which green is not distinguished. **G.**, Paris-, the acetoarsenite of copper. **G.**, Scheele's, cupric arsenite. **G.**, Schweinfurth's, synonym of Paris-green. **G. Sickness**, chlorosis.

Gregarina (*greg-ar-i'-nah*) [*greg*, a herd]. A genus of protozoa.

Grindelia (*grin-dē'-le-nā*) [after H. Grindel, a German botanist]. The leaves and flowering tops of *G. robusta*, wild sunflower, or gum-plant, and *G. squarrosa*. **G.**, Ext., Fld. Dose [$\frac{3}{4}$ ss–f $\frac{3}{4}$] (2.0–4.0). It is used in asthma, bronchitis, and whooping-cough, and locally in rhus-poisoning.

Grinder (*grin'-der*) [AS., *grindan*, to grind]. A molar tooth.

Grinder's Asthma. A fibroid pneumonia, a chronic affection of the lungs resulting from the inspiration of metallic or siliceous dust.

Grip or Grippe (*grip*). See *Influenza*.

Gristle (*gris'-l*) [AS., *gristel*, cartilage]. Cartilage.

Groin [Icel., *grein*, a branch or arm]. The depression between the abdomen and thigh.

Groove (*groov*) [D., *groef*, a channel]. A furrow or channel. **G.**, Bicipital, the deep groove on the anterior surface of the humerus separating the greater and lesser tuberosities and containing the long tendon of the biceps.

G., Cavernous, a broad groove on the superior surface of the sphenoid bone lodging the internal carotid artery and the cavernous sinus. **G., Medullary**, a long shallow furrow that appears along the dorsal line of the neural tube of the embryo.

Gross (*grōs*) [*Fr., gros, great*]. Coarse; large. **G. Anatomy**, anatomy as studied with the naked eye.

Grove Cell. See *Cell*.

Growing-pains (*gro'-ing*) [*AS., grown, to grow*]. A term applied to pains in the limbs occurring during youth, and perhaps of rheumatic origin.

Gruel (*gru'-el*) [*O. Low G., grut, groats*]. A decoction of corn-meal or oat-meal boiled in water to a thick paste.

Grumous (*gru'-mus*) [*grumus, a little heap*]. Clotted; consisting of lumps.

Grutum (*gru'-tum*) [*grutum, grit*]. See *Milium*.

Gtt. Abbreviation of *gutta*, or *guttæ*, drop, or drops.

Guachamacá (*gwaw-shaw-maw'-kah*). The bark of an apocynaceous tree. **G. toxifera**, or *Malouetia nitida*, furnishes a virulent arrow-poison, somewhat resembling curare; it has been employed in tetanus and hydrophobia. Unof.

Guaco (*gwaw'-ko*). The *Mikania guaco* and other species of *Mikania* and *Aristolochia*, used in South America for snake-bites; it has been employed in rheumatism, gout, and in various skin-diseases. Dose of a watery extract gr. iij (0.19).

Guaiacol (*gwai'-ak-ol*) [*S. Amer.*], $C_8H_8O_2$. Methyl-pyrocatechin, a substance obtained from beechwood creosote and also, synthetically, from pyrocatechin and methyl-sulphuric acid. It is used as a substitute for creosote in tuberculosis in doses of mjj-v (0.19-0.32); externally it has been employed as an antipyretic, 20-40 drops being painted on the skin.

Guaiacum (*gwai'-ak-um*). A genus of trees belonging to the order Zygophyllæ. **G., Lignum**, guaiac-wood, is the heart-wood of *G. officinale* and *G. sanctum*, and yields guaiac-resin—*Resina guaiaci*. The last contains guaiacic acid, $C_{13}H_{16}O_6$, guaiac-yellow, guaiacene, C_8H_8O , guaiacol, and pyroguaiacin. Guaiac is alterative, expectorant, and diaphoretic. It is used in syphilis, chronic rheumatism, and gout.

Guanin (*gwaw'-nin*) [*Peruv., huannu, dung*], $C_4H_5N_5O$. A leukomatin found in the pancreas, liver, and muscle-extract, as a decomposition-product of nuclein. It also occurs in guano, and is non-poisonous.

Guano (*gwaw'-no*) [*Peruv., huannu, dung*]. The excrement of sea-fowl found on certain islands in the Pacific Ocean. It contains guanin and alkaline urates and phosphates,

and is used externally in certain skin-diseases.

Guarana (*gwaw-rah'-nah*) [*Braz.*]. A dried paste prepared from the seeds of *Paullinia sorbilis*, found in Brazil. It contains an alkaloid, guaranin, $C_8H_{10}N_4O_2 \cdot H_2O$, identical with caffein. It is employed in nervous sick-headaches. Dose of the fluid extract, m̄v-xxx (0.32-2.0); of guaranin gr. j-ij (0.065-0.19).

Gubernaculum testis (*gu-ber-nak'-u-lum tes'-tis*) [*L.*]. The conical-shaped cord attached above the lower end of the epididymis, below to the bottom of the scrotum, and governing the descent of the testes.

Gubler's Line. A line connecting the superficial origin of the trifacial nerves. A lesion of the pons below this line causes crossed paralysis. **G.'s Tumor**, a prominence over the dorsum of the wrist in chronic wrist-drop.

Gullet (*gul'-et*) [*gula, throat*]. See *Esophagus*.

Gum [*gummi, gum*]. A concrete vegetable juice exuded from many plants. When treated with nitric acid it yields mucic acid. Gums are either entirely soluble in water, or swell up in it into a viscid mass. Various names are given to gums, usually indicating the place whence exported. **G. Arabic**. See *Acacia*. **G. Benjamin**. See *Benzoïn*. **G., Blue**. See *Eucalyptus*. **G. Tragacanth**. See *Tragacanth*.

Gumma (*gum'-ah*) [*gummi, gum*]. The gummy tumor characterizing the tertiary stage of syphilis. It consists of granulation-tissue, with giant-cells, and is the seat of a peculiar degeneration which causes the gummy appearance.

Gummatous (*gum'-at-us*) [*gummi, gum*]. Of the nature of or affected with gummata.

Gurgling [*gurges, a whirlpool*]. The peculiar sound caused by the passage of gas through a liquid. It is observed upon palpation of the abdomen in enteric fever and other conditions in which the bowel is distended with gas and contains liquid. **G. Râle**, a sound heard over the chest when the bronchi or pulmonary cavities contain fluid.

Gurjun Balsam (*ger'-jun*). Balsamum dipterocarpi, wood-oil. An oleoresin obtained from several species of *Dipterocarpus*, trees native to Southern Asia. It is similar to copaiba, but more decided in therapeutic effects, and is less unpleasant. It is used as an expectorant, and in leprosy and gonorrhea. Dose m̄xv-xl (1.0-2.6).

Gustatory (*gus'-ta-to-re*) [*gustare, to taste*]. Pertaining to taste. **G. Bud**, a taste-bud.

G. Nerve. See *Nerve*.

Gut [*ME., gut, intestine*]. The intestine.

Gutta (*gut'-ah*) [*L.*]. A drop. **G. rosacea**, rosacea. **G. serena**, amaurosis. **G.-percha**,

the concrete juice of *Dichopsis Gutta* and other species of the natural order Tapotaceæ. It is used to make splints, as a dressing for wounds, and as a vehicle for caustic substances.

Guttatim (*gut-a'-tim*) [*gutta*, a drop]. Drop by drop.

Guttur (*gut'-er*) [L.]. The throat.

Guttural (*gut'-u-ral*) [*guttur*, the throat]. Pertaining to the throat.

Gutturotetany (*gut-u-ro-tet'-an-e*) [*guttur*, throat; *tetanus*, tetanus]. A form of stuttering in which the pronunciation of such sounds as *g*, *k*, *q*, is difficult.

Gymnastics (*jim-nas'-tiks*) [*γυμνός*, naked]. Physical exercise, especially systematic exercise for the purpose of restoring or maintaining the bodily health. *G.*, Ocular, regular muscular exercise of the eye by the use of prisms or other means to overcome mus-

cular insufficiency. *G.*, Swedish, a system of exercises to restore strength to paretic muscles, consisting in movements made by the patient against the resistance of an attendant.

Gynecian (*jin-e'-se-an*) [*γυνή*, a woman]. Pertaining to women.

Gynecology (*jin-e'-kol'-o-je*) [*γυνή*, a woman; *λόγος*, science]. The science of the diseases of women, especially of those affecting the sexual organs.

Gynephobia (*jin-e-fo'-be-ah*) [*γυνή*, woman; *φόβος*, fear]. Morbid aversion to the society of women.

Gypsum (*jip'-sum*) [*γύψος*, chalk], $\text{CuSO}_4 + 2\text{H}_2\text{O}$. Native calcium sulphate. Deprived of its water of crystallization it constitutes plaster of Paris.

Gyrus (*ji'-rus*) [*γῦρος*, a circle]. A convolution of the brain. See *Convolution*.

H

H. 1. Symbol for Hydrogen. 2. Abbreviation for Hypermetropia, and for *Haustus*, a draught.

Habenula (*hab-en'-u-lah*) [*habena*, a rein]. 1. A ribbon-like structure; a name applied to different portions of the basilar membrane of the internal ear. 2. A peduncle of the pineal gland. Ganglion habenulæ, a small club-shaped body on the mesial surface of the optic thalamus, in which the corresponding peduncle of the pineal gland terminates.

Habit (*hab'-it*) [*habere*, to have]. 1. The general condition or appearance of an individual, as a full habit, a condition of plethora indicated by congestion of the superficial vessels and obesity. 2. The tendency to repeat an action or condition. *H.-spasm*, or habit-chorea, a spasmodic, constantly recurring movement of certain voluntary muscles, usually seen in children.

Habitat (*hab'-it-at*) [*habitare*, to dwell]. The natural home of an animal or vegetal species.

Habitus (*hab'-it-us*) [*habere*, to have]. General appearance or expression.

Hachement (*haksh-mon(g')*) [Fr., hacking, or chopping]. A form of massage consisting of a succession of strokes.

Hacking (*hak'-ing*). See *Hachement*.

Hadernkrankheit (*hak'-dern-kronk-hit*) [G.]. A disease of rag-pickers, by some supposed to be anthrax. Others look upon it as malignant edema.

Hæma- (*hem'-ah-*) [*αἷμα*, blood]. A prefix signifying relationship with the blood. See *Hema*.

Haidinger's Brushes. A brush-like image seen on directing the eye toward a source of polarized light, due to the doubly refractive character of the elements of the macula.

Hair [AS., *hær*]. A delicate filament growing from the skin of mammals; collectively, all the filaments forming the covering of the skin. Hair is a modified epidermal structure, and consists of a shaft and a root, the latter expanded at its end into the hair-bulb, which is concave and caps the hair-papilla. *H.-cell*, an epithelial cell with delicate hair-like processes, as, *e.g.*, the hair-cells of the organ of Corti. *H.-follicle*, the depression in the corium and subcutaneous connective tissue containing the root of the hair.

Haliteresis (*hal-is-ter-e'-sis*) [*ἅλς*, salt; *στέρησις*, privation]. The loss of lime-salts of bone.

Halitus (*hal'-it-us*) [L., a vapor]. A vapor, as that expired from the lung.

Haller [a German physiologist]. *H.*, Circle of, arterial and venous circles in the eye.

Hall's (Marshall) Disease. See *Diseases, Table of*.

Hallucination (*hal-lu-sin-a'-shun*) [*hallucinari*, to wander in mind]. A false sense-perception; it is the perception of an object or phenomenon which has no external existence, as *H.* of sight, sound, smell, taste, or touch.

Hallux (*hal'-uks*) [L.]. The great toe. **H. valgus**, displacement of the great toe toward the other toes. **H. varus**, displacement of the great toe away from the other toes.

Halo (*hal'-lo*) [*ἅλως*, a round threshing-floor]. 1. The areola of the nipple. 2. The luminous circles seen about a light.

Halogen (*hal'-o-jen*) [*ἅλς*, salt; *γεννᾶν*, to produce]. A univalent element that forms a compound of a saline nature by its direct union with a metal. The halogens are chlorine, iodine, bromine, and fluorine. **H. Acid**, an acid formed by the combination of a halogen with hydrogen.

Haloid (*hal'-oid*) [*ἅλς*, salt; *εἶδος*, likeness]. Resembling sea-salt. **H. Salts**, any one of those compounds that consist of a metal directly united to chlorine, bromine, iodine, or fluorine.

Ham [AS., *hamm*, the ham]. The back part of the knee; the popliteal space.

Hamamelis (*ham-am-e'-lis*) [*ἅμα*, together with; *μήλον*, apple-tree]. The leaves of *H. virginica*; witch-hazel. It is styptic, and sedative, and is used in uterine and other hemorrhages; as an application to hemorrhoids and to contused parts. **Ext. hamamelidis fld.** Dose ℥x-xx (0.65-1.3).

Hammer, Thermal. A hammer-shaped cautery-iron.

Hammer-toe. A term applied to a condition of the second toe in which the proximal phalanx is extremely extended while the two distal phalanges are flexed.

Hamstring. The tendons bounding the ham above on the outer and inner side. **H., Inner**, the tendons of the semimembranosus, sartorius, gracilis, and semitendinosus muscles. **H., Outer**, the tendons of the biceps flexor cruris.

Hamular (*ham'-u-lar*) [*hamus*, a hook]. Pertaining to or shaped like a hook.

Hamulus (*ham'-u-lus*) [dim. of *hamus*, a hook]. A hook-shaped process, as of a bone. **H. of the Cochlea**, the hook-like process of the osseous lamina at the cupola.

Hand [ME., *hand*, hand]. The organ of prehension in bipeds and quadrupeds, composed of the carpus, the metacarpus, and the phalanges. **H., Ape, H., Claw, H., Monkey.** See *Main en griffe*.

Handkerchief (*hang'-ker-chif*) [ME., *hand*, hand; *kerchief*, a kerchief]. A square piece of cloth for wiping the face or nose. **H. Dressing**, a form of temporary dressing for wounds and fractures, made of handkerchiefs.

Hangnail (*hang'-nail*). A partly detached piece of epidermis at the root of the nail, the friction against which has caused inflammation of the abraded surface.

Haphalgnesia (*haf-al-je'-se-ah*) [*ἄφῃ*, touch; *ἄλγος*, pain]. A feeling of pain produced by merely touching an object.

Hardening (*hard'-en-ing*) [AS., *hard*, hard]. A stage in the preparation of tissues for microscopic examination, in which they are rendered firm, so that they may, after embedding, be readily cut.

Harelip-suture. A figure-of-8 suture about a pin thrust through the lips of the freshened edges of the cleft.

Harmony (*har'-mo-ne*) [*ἁρμονία*, harmony]. A form of articulation between two bones that are closely and immovably apposed.

Hartshorn (*harts'-horn*). 1. Cornu cervi, the horn of the stag, formerly a source of ammonia, or spirits of hartshorn. 2. A name popularly given to ammonium hydroxid. See *Ammonium*.

Hashish (*hash'-esh*). See *Cannabis*.

Hassall, Concentric Corpuscles of. Peculiar bodies found in the medulla of the thymus gland. They consist of concentric layers of flattened epithelial cells.

Haunch (*hau'-ench*) [Fr., *hanche*, haunch]. The part of the body including the hips and the buttocks. **H.-bone**, the innominate bone.

Hautus (*haws'-tus*) [*haurire*, to draw]. A draught.

Havers, Canals of. See *Canal*. **H., Spaces of**, large and irregular spaces found in growing bone, and resulting from the absorption of the primary osseous formation. In them the Haversian systems are deposited.

Haversian (*hav-er'-zhe-un*) [*Havers*, an English anatomist]. Described by Havers.

H. Canal. See *Canal*. **H. System**, an Haversian canal with canaliculi radiating from it and the surrounding concentric lamellæ.

Hawking (*haw'-king*) [ME., *hawk*, to hawk]. Clearing the throat by a forcible expiration.

Hay-asthma, Hay-cold, Hay-fever. An acute affection of the conjunctiva and upper air-passages, coming on periodically at certain seasons of the year, especially in summer and autumn, in persons predisposed to the disease. The exciting factor is in some cases the pollen of grasses; in others the disease seems to be caused reflexly by polypi and other diseased conditions of the nose. The chief symptoms are coryza, sneezing, headache, cough, and asthmatic attacks.

Hay's Method. A method of removing dropsical effusions by producing frequent serous evacuations by means of saline cathartics combined with a dry diet.

Head (*hed*) [ME., *hed*, the head]. 1. The uppermost part of the body; that part of the body containing the brain, the organ of sight of smell, taste, and hearing, and part of the

organs of speech. 2. The top, beginning, or most prominent part of anything, as the head of the femur, the head of a muscle etc. **H.-drop**, a peculiar disease seen in Japan during the spring and early summer, supposed to be miasmatic in origin. It is attended with inability to hold the head erect, paralytic symptoms in the limbs, and optic disorders. One attack predisposes to others. **H.-gut**. See *Fore-gut*. **H.-kidney**, pronephros. **Head-louse**, the pediculus capitis.

Headache (*hed'-ak*). Pain in the head; cephalalgia. The following varieties of headache are described: local, general; organic, functional; toxemic—alcoholic, caffeinic, diabetic, lithemic, malarial, rheumatic, uremic; reflex—gastric, ocular, nasal, uterine; cardiac, pulmonic, anemic, congestive; hysterical, neurasthenic. **H.**, Biliary, migraine. **Healing** (*he'-ling*) [*AS.*, *hēlan*, to heal]. The process or act of getting well, or of making whole; especially the getting well of an ulcer or wound. **H.** by First Intention or Primary Healing, the union of two accurately apposed surfaces without any visible granulating process. **H.** by Second Intention, healing through the medium of granulations, which fill up the gap of the wound. **H.** by Third Intention, the two granulating surfaces are approximated so as to unite and heal readily.

Health (*hel'h*) [*AS.*, *hēlan*, to heal]. That condition of the body in which all the functions are performed normally.

Hearing (*he'-ring*) [*AS.*, *hýran*, to hear]. The special sense by which the sonorous vibrations of the air are communicated to the mind. The organ of hearing is the ear, whence the vibrations are carried by the auditory nerve to the center of hearing, situated in the temporosphenoidal lobe of the cerebrum.

Heart (*hart*) [*AS.*, *heorte*, heart]. A hollow, muscular organ, the function of which is to pump the blood through the vessels. It is enveloped by a serous sac called the pericardium, and consists of two symmetric halves, a right auricle and ventricle and a left auricle and ventricle. The right auriculo-ventricular orifice is guarded by the tricuspid valve; the left by a valve with two leaflets, the mitral. These valves are broad and thin, consisting of two layers of the lining membrane of the heart, the endocardium, separated by a slight amount of connective tissue. To support them, thin chordæ tendinæ join their free margins to the muscles in the wall of the ventricle, the columnæ carneæ. The outlet of the right ventricle into the pulmonic artery, and the left into the aorta, are guarded by stout, short, tricuspid valves, the semi-

lunar valves. **H.-burn**, a burning feeling at the epigastrium and lower part of the chest, caused by fermentation of the gastric contents. **H.**, Fatty, a name given to two distinct pathologic conditions of the heart-tissue. In the first there is a true fatty degeneration of the muscular fibers of the heart; and in the second there is an increase in the quantity of subpericardial fat—a fatty infiltration.

Heart-berg (*hart'-berg*). The vulgar name of the thoracic portion of the thymus gland of animals. See *Neck-berg*.

Heat (*het*) [*AS.*, *hatu*, heat]. A form of kinetic energy communicable from one body to another; it is that form of molecular motion which is appreciated by a special thermal sense. **H.**, Atomic. See *Atomic*. **H.**, Capacity for, the number of heat-units required to raise the temperature of a body 1° C. **H.-centers**, centers in the brain for stimulating heat-production or heat-elimination, and for regulating the relation of these. **H.**, Latent, the quantity of heat necessary to convert a body into another state without changing its temperature. **H.**, Molecular, the product of the molecular weight of a compound multiplied by its specific heat. **H.**, Prickly, or **H.-rash**, miliaria. **H.**, Specific, this is the ratio of the capacity for heat of a body to that of an equal weight of water. **H.-stroke**, the symptoms produced by exposure to great heat, either that of the sun or that of heated rooms. **H.-unit**, the amount of heat required to raise the temperature of one kilogram of water from 0° to 1° C.; it is technically called a calorie.

Hebephrenia (*he-be-fre'-ne-ah*) [*ἡβη, puberty*; *φρήν, mind*]. A form of mental derangement occurring in young persons at or soon after the age of puberty, and characterized by mental deterioration, and a gradually increasing egoism. It may end in a permanent dementia.

Heberden's Nodes. See *Nodes*.

Hebetude (*heb'-e-tūd*) [*hebetudo*, bluntness]. Dulness of the special senses and intellect; a condition present in grave fevers.

Hectic Fever (*hek'-tik*) [*ἡκτικός, habitual*]. A fever caused by absorption of toxic substances formed in the process of suppuration, and characterized by daily intermissions and frequent drenching sweats. It occurs in pulmonary tuberculosis, in pyemia, etc. **H. Flush**, the flushed cheek seen in hectic fever.

Hectogram (*hek'-to-gram*) [*ἑκατόν, one hundred*; *γράμμα, an inscription*]. One hundred grams, or 1543.2349 grains.

Hectoliter (*hek'-to-le-ter*) [*ἑκατόν, hundred*; *λίτρα, a pound*]. One hundred liters, equal to 22.009 imperial gallons, or 26.4 U. S. gallons.

Hectometer (*hek'-to-me-ter*) [*ἑκατόν*, hundred; *μέτρον*, a measure]. One hundred meters, or 328 feet, 1 inch.

Hedeoma (*he-de-o'-mah*) [*ἡδύς*, sweet; *ὀσμή*, smell]. American pennyroyal. The leaves and tops of *H. pulegioides*, the properties of which are due to a volatile oil. It is stimulant, carminative, and emmenagogue. It is used in suppression of the menses, the flatulent colic of children, and, on account of its pungent odor, to drive off fleas and mosquitoes. **H.**, **Oleum**, the volatile oil. Dose πj -x (0.13-0.65).

Hedge-hog Crystals. Crystals of ammonium urate found as a urinary deposit in the form of globular crystals, with spiny projections.

Hedrocele (*hed'-ro-sel'*) [*ἕδρα*, breech; *κύλη*, tumor]. A hernia through the notch of the ischium. Also, prolapsus of the anus.

Heel (*hel'*) [*AS.*, *hela*, heel]. The hinder part of the foot. **H.-bone**, the calcaneum.

Heidenhain, Demilune Cells of. See *Demilune* and *Crescent of Giannuzzi*.

Helcoid (*hel'-koid*) [*ἕλκος*, ulcer; *εἶδος*, likeness]. Resembling an ulcer.

Helcoma (*hel-ko'-mah*) [*ἕλκος*, an ulcer]. An ulcer.

Helicine (*hel'-is-in*) [*ἕλιξ*, a spiral]. 1. Spiral in structure. 2. Pertaining to the helix.

H. Arteries, spirally-winding arteries supplying the erectile tissue of the penis.

Helicoid (*hel'-ik-oid*) [*ἕλιξ*, a spiral; *εἶδος*, form]. Spiral; coiled like a snail-shell.

Helicotrema (*hel-ik-o-tre'-mah*) [*ἕλιξ*, helix, coil; *τρήμα*, hole]. The opening connecting the scala tympani and vestibuli of the spiral canal of the cochlea.

Heliophobia (*he-le-o-fo'-be-ah*) [*ἥλιος*, sun; *φόβος*, fear]. Morbid fear of exposure to the sun's rays.

Heliotherapy (*he-le-o-ther'-ap-e*) [*ἥλιος*, sun; *θεραπεία*, treatment]. The treatment of disease by exposure of the body to sunlight. Sun-bathing.

Helix (*he'-iks*) [*ἕλιξ*, a coil]. 1. The rounded, convex margin of the pinna of the ear. 2. A coil of wire, as that of a faradic battery or of an electromagnet.

Hellebore, Black. See *Helleborus*. **H.**, **White.** See *Veratrum album*.

Helleborus (*hel-eb'-or-us*) [*ἑλλέβορος*, hellebore]. The root of *H. niger*, black hellebore, the properties of which are due to two glucosids, helleborin, $C_{26}H_{42}O_{16}$, and helleborein, $C_{26}H_{44}O_{15}$. It is a drastic hydragogue cathartic, and an emmenagogue, and has been used as a drastic purge in insanity, dropsy, and amenorrhea. Dose of the powdered root, as a purge, gr. x-xx (0.65-1.3).

Heller's Test. 1. A test for albumin in the urine. A little nitric acid is placed in a test-

tube and the urine allowed carefully to flow down the side of the tube, so as to form a layer on the acid without mixing. The development of an opaque white ring indicates albumin. 2. A test for the presence of blood-coloring matter in the urine. The urine is boiled with half its volume of caustic potash, —whereby the phosphates are precipitated. The precipitate is colored red if blood is present.

Helminth (*hel'-minth*) [*ἔλμινξ*, worm]. A worm.

Helminthagogue (*hel-minth'-ag-ag*). See *Anthelmintic*.

Helminthiasis (*hel-min-thi'-as-is*) [*ἔλμινξ*, a worm]. The diseased condition produced by the presence of worms in the body.

Helminthology (*hel-min-thol'-o-jé*) [*ἔλμινξ*, a worm; *λόγος*, science]. A treatise on worms, especially those parasitic within the body.

Hemabrometer (*hem-ab-ar-om'-et-er*) [*αἷμα*, blood; *βαρὺς*, heavy; *μέτρον*, measure]. An instrument for the determination of the specific gravity of the blood.

Hem-, Hemato- (*hem-, hem'-at-o-*) [*αἷμα*, blood]. A prefix signifying of or pertaining to the blood.

Hemacyanin (*hem-as-i'-an-in*). See *Hemocyanin*.

Hemad (*hem'-ad*) [*αἷμα*, blood]. Toward the hemal aspect; opposed to neurad.

Hemadrometer (*hem-ad-ro-mom'-et-er*) [*αἷμα*, blood; *δρόμος*, course; *μέτρον*, measure]. An instrument for measuring the velocity of the blood-current.

Hemadynamometer (*hem-ad-i-nam-om'-et-er*) [*αἷμα*, blood; *δύναμις*, strength; *μέτρον*, a measure]. An instrument for measuring the tension or pressure of blood within the arteries.

Hemagogue (*hem'-ag-ag*) [*αἷμα*, blood; *ἀγωγός*, leading]. Promoting the menstrual or hemorrhoidal discharge of blood.

Hemal (*hem'-al*) [*αἷμα*, blood]. 1. Pertaining to the blood or vascular system. 2. Pertaining to the ventral aspect of the body, that part containing the heart and blood-vessels.

Hemameba (*hem-am-e'-bah*) [*αἷμα*, blood; *ἀμβιβή*, a change]. A white blood-cell.

Hemangioma (*hem-an-je-o'-mah*) [*αἷμα*, blood; *ἄγγειον*, vessel; *όμα*, a tumor]. An angioma made up of blood-vessels.

Hemaphein (*hem-af-e'-in*) [*αἷμα*, blood; *φαῖδς*, dusky]. A brown coloring-matter from blood, regarded as a decomposition product of hematin.

Hemapoiesis (*hem-ap-oi-e'-sis*). See *Hematopoiesis*.

Hemapoietic (*hem-ap-oi-et'-ik*). See *Hematopoietic*.

Hemarthrosis (*hem-ar-thro'-sis*) [*haima*, blood; *arthrosis*, a joint]. Effusion of blood into a joint.

Hematachometer (*hem-at-ak-om'-et-er*). See *Hemotachometer*.

Hematemesis (*hem-at-em'-es-is*) [*haima*, blood; *emein*, to vomit]. The vomiting of blood.

Hematic (*hem-at'-ik*) [*haima*, blood]. Bloody; pertaining to, full of, or having the color of, blood.

Hematidrosis (*hem-at-id-ro'-sis*) [*haima*, blood; *idrosis*, sweat]. A sweating of blood.

Hematimeter (*hem-at-im'-et-er*) [*haima*, blood; *metron*, measure]. An instrument for counting the corpuscles in a given volume of blood.

Hematin (*hem'-at-in*) [*haima*, blood], $C_{78}H_{72}N_4FeO_4$. A decomposition-product of hemoglobin. It is bluish-black, amorphous, contains iron, and is soluble in dilute alkalies and acids, insoluble in water, in alcohol, and in ether.

Hematinic (*hem-at-in'-ik*) [*haima*, blood]. 1. Relating to hematin. 2. An agent which tends to increase the proportion of hematin or coloring-matter in the blood.

Hematinometer (*hem-at-in-om'-et-er*). Same as *Hemoglobinometer*.

Hematuria (*hem-at-in-u'-re-ah*) [*haima*, blood; *ouron*, urine]. Same as *Hemoglobinuria*.

Hemato- (*hem'-at-o-*) [*haima*, blood]. A prefix signifying pertaining to the blood.

Hematobium (*hem-at-o'-be-um*) [*haima*, blood; *bios*, life]. 1. A blood-corpuscle. 2. A blood-parasite.

Hematoblast (*hem'-at-o-blast*) [*haima*, blood; *blastos*, germ]. A blood-plate; an immature red blood-corpuscle.

Hematocoele (*hem'-at-o-sel*) [*haima*, blood; *khele*, a tumor]. A tumor formed by the extravasation and collection of blood in a part, especially in the tunica vaginalis testis, or in the pelvic cavity (Pelvic H.).

Hematocephalus (*hem-at-o-sef'-al-us*) [*haima*, blood; *kephale*, head]. 1. An effusion of blood into the brain. 2. A monstrosity characterized by an effusion of blood into the cerebral hemispheres.

Hematochyluria (*hem-at-o-ki-lu'-re-ah*) [*haima*, blood; *chylus*, chyle; *ouron*, urine]. The presence of blood and chylous material in the urine.

Hematocolpos (*hem-at-o-kol'-pos*) [*haima*, blood; *kolos*, vagina]. A collection of blood within the vagina.

Hematocrit (*hem'-at-o-krit*) [*haima*, blood; *kriuein*, to separate, to judge]. An instrument for making volumetric estimation of the blood-corpuscles by separating, by cen-

trifugal action, the corpuscles from the plasma.

Hematocyst (*hem'-at-o-sist*) [*haima*, blood; *kystis*, bladder]. 1. A cyst containing blood. 2. An effusion of blood into the bladder.

Hematocyte (*hem'-at-o-sit*) [*haima*, blood; *kytos*, cell]. A blood-corpuscle.

Hematocytometer (*hem-at-o-sit-om'-et-er*). See *Hemacytometer*.

Hematodes (*hem-at-o'-des*) [*haimatodes*, bloody]. Bloody.

Hematogenesis (*hem-at-o-jen'-es-is*) [*haima*, blood; *genesis*, production]. The development of blood or blood corpuscles.

Hematogenic (*hem-at-o-jen'-ik*) [*haima*, blood; *gennan*, to produce]. Pertaining to the formation of blood.

Hematogenous (*hem-at-o-jen'-us*) [*haima*, blood; *gennan*, to produce]. Derived from, or having origin in, the blood.

Hematoglobin (*hem-at-o-glo'-bin*). See *Hemoglobin*.

Hematoid (*hem'-at-oid*) [*haima*, blood; *eidos*, form]. Resembling blood.

Hematoidin (*hem-at-oid-in*) [*haima*, blood; *eidos*, resemblance], $C_{16}H_{16}N_4O_4$. An iron-free derivative of hemoglobin, occurring in old blood-clots as yellowish-brown, rhombohedral crystals.

Hematokrit (*hem'-at-o-krit*). See *Hematocrit*.

Hematology (*hem-at-ol'-o-je*) [*haima*, blood; *logos*, science]. The science of the blood, its nature, functions, and diseases.

Hematolysis (*hem-at-ol'-is-is*) [*haima*, blood; *lysis*, a solution]. Destruction or disorganization of the blood or of the corpuscles.

Hematoma (*he-mat-o'-mah*) [*haima*, blood; *ma*, tumor]. A tumor or swelling containing blood. **H. auris**, insane ear; an effusion of blood or serum between the cartilage of the ear and its covering, occurring in various forms of insanity as the result of injuries or trophic changes. **H. of the Dura mater**, an effusion of blood beneath the dura mater, forming membranous layers. **H., Pelvic**, an effusion of blood into the cellular tissue of the pelvis.

Hematomediastinum (*hem-at-o-me-de-as-ti'-num*) [*haima*, blood; *mediastinum*, the mediastinum]. An effusion of blood into the mediastinal spaces.

Hematometra (*hem-at-o-me'-trah*) [*haima*, blood; *metra*, womb]. An accumulation of blood within the uterine cavity.

Hematomonas (*hem-at-om'-on-as*) [*haima*, blood; *monas*, a unit]. A genus of protozoan parasites inhabiting the blood.

Hematomphalocele (*hem-at-om-fal'-o-sel*) [*haima*, blood; *omphalos*, navel; *khele*, tumor]. A hernia at the navel, distended with blood.

Hematomyelia (*hem-at-o-mi-el'-le-ah*) [*aiua*, blood; *μυελός*, marrow]. Hemorrhage into the spinal cord; an accumulation of blood in the central canal of the spinal cord.

Hematomyelitis (*hem-at-o-mi-el'-tis*) [*aiua*, blood; *μυελός*, marrow; *τις*, inflammation]. An acute myelitis attended with an effusion of blood into the spinal cord.

Hematopericardium (*hem-at-o-per-ik-ar'-de-um*) [*aiua*, blood; *περί*, about; *καρδία*, heart]. An effusion of blood into the pericardium.

Hematophagous (*hem-at-off'-ag-us*) [*aiua*, blood; *φαγείν*, to eat]. Feeding on blood; blood-sucking.

Hematophyte (*hem'-at-o-fit*) [*aiua*, blood; *φύτον*, a plant]. A vegetable organism, such as a bacterium, living in the blood.

Hematoplastic (*hem-at-o-plas'-tik*) [*aiua*, blood; *πλαστικός*, plastic]. Blood-forming.

Hematopoiesis (*hem-at-o-poi-el'-sis*) [*aiua*, blood; *ποιεῖν*, to make]. The formation of blood.

Hematopoietic (*hem-at-o-poi-el'-ik*) [*aiua*, blood; *ποιεῖν*, to make]. Relating to the processes of blood-making. **H. Organs**, the blood-making organs.

Hematoporphyrin (*hem-at-o-por'-fe-rin*) [*aiua*, blood; *πύρρονος*, purple], $C_{88}H_{74}N_8O_{12}$. Iron-free hematin, a decomposition-product of hemoglobin occurring in the urine in conditions associated with destruction of red corpuscles. It is produced by dissolving hematin in concentrated sulphuric acid.

Hematoporphyrinuria (*hem-at-o-por'-fe-rin-ul'-re-ah*) [*aiua*, blood; *πύρρονος*, purple; *οὐρὸν*, urine]. The presence of hematoporphyrin in the urine.

Hematoposia (*hem-at-o-po'-ze-ah*) [*aiua*, blood; *ποσις*, a drinking]. The drinking of blood.

Hematorrhachis (*hem-at-or'-ak-is*) [*aiua*, blood; *ράχη*, spine]. Hemorrhage within the vertebral canal.

Hematorrhea (*hem-at-or'-el'-ah*) [*aiua*, blood; *ῥοα*, a flow]. A copious flow or discharge of blood.

Hematosalpinx (*hem-at-o-sal'-pinkz*) [*aiua*, blood; *σαλπίς*, a trumpet]. A collection of blood in a Fallopian tube.

Hematoscope (*hem'-at-o-skōp*) [*aiua*, blood; *σκοπεῖν*, to inspect]. An instrument used in the spectroscopic examination of the blood, by means of which the thickness of the layer of blood can be regulated.

Hematoscopy (*hem-at-o-skō-pe*) [*aiua*, blood; *σκοπεῖν*, to inspect]. Visual examination of the blood; examination of the blood by means of the hematoscope.

Hematosepsis (*hem-at-o-sep'-sis*) [*aiua*, blood; *σήψις*, putrefaction]. Septicemia.

Hematosis (*hem-at-o'-sis*) [*aiua*, blood].

The process of the formation of blood and the development of blood-corpuscles.

Hematospectroscope (*hem-at-o-spek'-tro-skōp*) [*aiua*, blood; *spectrum*, an image; *σκοπεῖν*, to view]. A spectroscope adapted to the study of the blood.

Hematospermia (*hem-at-o-sper'-me-ah*) [*aiua*, blood; *σπέρμα*, seed]. The discharge of bloody semen.

Hematothorax (*hem-at-o-thō'-raks*). See *Hemothorax*.

Hematoxylin (*hem-at-oks'-il-in*) [*aiua*, blood; *ξύλον*, wood], $C_{16}H_{14}O_6$. The coloring-matter of logwood. It is a crystalline substance and is used as a stain in microscopy.

Hematoxylon (*hem-at-oks'-il-on*) [*aiua*, blood; *ξύλον*, wood]. Logwood; the heartwood of *H. campechianum*; it contains tannic acid and a coloring-principle, hematoxylin, $C_{16}H_{14}O_6$, and is a mild astringent. **H.**, **Ext.** Dose gr. v-xx (0.32-1.3). **H.**, **Decoctum**, strength 1 to 17. Dose f3j-ij (32.0-64.0). **Unof.**

Hematozoon (*hem-at-o-zō'-on*) [*aiua*, blood; *ζῶον*, animal]. Any animal parasite in the blood.

Hematuria (*hem-at-u-re'-sis*) [*aiua*, blood; *οὐρὸν*, urine]. The passage of bloody urine.

Hematuria (*hem-at-u'-re-ah*) [*aiua*, blood; *οὐρὸν*, urine]. The discharge of urine containing blood. When only the coloring-matter of the blood is found in the urine, it is termed hemoglobinuria or hematuria.

Hemautography (*hem-aw-tog'-ra-fe*) [*aiua*, blood; *αὐτός*, self; *γράφειν*, to write]. The tracing produced by a jet of blood from a divided artery caught upon paper drawn in front of it.

Hemeralopia (*hem-er-at-o'-pe-ah*) [*ἡμέρα*, day; *ὤψ*, eye]. Night-blindness; a condition in which the individual sees relatively much better in the day than at night. **H.** is a symptom of retinitis pigmentosa.

Hemichromatopsia (*hem-e-ah-kro-mat-op'-se-ah*) [*ἡμι*, half; *ἀ* priv.; *χρῶμα*, color; *ὄψις*, vision]. Color-blindness in one-half of the field of vision.

Hemialbumin (*hem-e-al'-bu-min*). See *Antialbumin*.

Hemialbuminose (*hem-e-al'-bu'-min-ōs*). The same as *Hemialbumose*.

Hemialbumose (*hem-e-al'-bu-mōs*) [*ἡμι*, half; *albumen*, the white of egg]. The most characteristic and most frequently obtained by-product of proteid-digestion. It is the fore-runner of hemipeptone.

Hemianesthesia (*hem-e-an-es-thē'-ze-ah*) [*ἡμι*, half; *ἀναισθησία*, want of feeling]. Anesthesia of one lateral half of the body.

Hemianopia, **Hemianopsia** (*hem-e-an-o'-p-ah*, *hem-e-an-op'-se-ah*) [*ἡμι*, half; *ὀψ* priv.].

ὄψις, sight]. Blindness in one-half of the visual field. It may be bilateral (binocular) or unilateral (uniocular). **H.**, **Binasal**, blindness on the nasal side of the visual field, usually due to disease of the outer sides of the optic commissure. **H.**, **Bitemporal**, blindness on the temporal side of the visual field, due to disease of the central parts of the commissure. **H.**, **Crossed** or **Heteronymous**, a general term for either binasal or bitemporal hemianopsia. **H.**, **Homonymous**, the form affecting the inner half of one field and the outer half of the other.

Hemiataxia (*hem-e-at-aks'-e-ah*) [*ἥμι*, half; *ἄ* priv.; *τάξις*, order]. Ataxia limited to one side of the body.

Hemiathetosis (*hem-e-ath-et-ō'-sis*) [*ἥμι*, half; *ἄθετος*, without fixed position]. Athetosis of one side of the body.

Hemiatrophy (*hem-e-at'-ro-fe*) [*ἥμι*, half; *ἀτροφία*, lack of nourishment]. Atrophy confined to one side of the body.

Hemic (*hem'-ik*) [*αἷμα*, blood]. Pertaining to or developed by the blood.

Hemichorea (*hem-e-ko-re'-ah*) [*ἥμι*, half; *χορεία*, chorea, dance]. A form of chorea in which the convulsive movements are confined to one side of the body.

Hemicrania (*hem-e-kra'-ne-ah*) [*ἥμι*, half; *κράνιον*, head]. 1. Neuralgia of one-half of the head; migraine. 2. Imperfect development or absence of the anterior or posterior part of the skull.

Hemidiaphoresis (*hem-e-di-af-or-e'-sis*) [*ἥμι*, half; *διαφύρεσις*, sweating]. Sweating of one lateral half of the body.

Hemidrosis (*hem-id-ro'-sis*). See *Hemididrosis*.

Hemidysesthesia (*hem-e-dis-es-the'-ze-ah*) [*ἥμι*, half; *δυσ*, difficult; *αἴσθησις*, sensation]. Dysesthesia of a lateral half of the body.

Hemiglossitis (*hem-e-glos-sit'-is*) [*ἥμι*, half; *γλῶσση*, tongue; *ῖτις*, inflammation]. Inflammation of one-half of the tongue.

Hemihyperesthesia (*hem-e-hi-per-es-the'-ze-ah*) [*ἥμι*, half; *ὑπέρ*, over; *αἴσθησις*, sensation]. Hyperesthesia confined to one lateral half of the body.

Hemimelus (*hem-im'-el-us*) [*ἥμι*, half; *μέλος*, limb]. A monster with incomplete or stunted extremities.

Hemin [*αἷμα*, blood], $C_{32}H_{30}N_4FeO_8$. Hematin's crystals, or chlorid of hematin. It is a refractive crystals derived from a drop of blood containing a trace of lactic acid.

Pertaining to hemianopsia. **H.** **Papillary Reaction**, Wernicke's reaction.

Hemiparesis (*hem-e-par'-es-is*) [*ἥμι*, half; *παρασις*, impairment of strength]. Paresis of one side of the body.

Hemipeptone (*hem-e-pep'-ton*). See *Peptones*.

Hemiplegia (*hem-e-ple'-je-ah*) [*ἥμι*, half; *πληγή*, stroke]. Paralysis of one side of the body, due usually to a lesion in the internal capsule or corpus striatum, but at times caused by an extensive lesion of the cortex, or a lesion of the crus, pons, or medulla, or of the upper part of the spinal cord. If in the brain, the lesion is on the side opposite to the paralysis. **H.**, **Alternate**, paralysis of the facial muscles upon one side, with paralysis of the trunk and extremities upon the opposite side of the body. **H.**, **Bilateral Spastic**. See *Paraplegia*, *Infantile Spasmodic*. **H.**, **Cerebral**, the ordinary form first described. **H.**, **Crossed**. Same as *H.*, *Alternate*. **H.**, **Facial**, motor paralysis of one side of the face. **H.**, **Spastic**, a form occurring in infants, in which the affected extremities are spastically contracted. **H.**, **Spinal**. See *Brown-Sequard's Paralysis*.

Hemispasm (*hem'-e-spazm*) [*ἥμι*, half; *σπασμός*, a spasm]. A spasm affecting only one side of the body.

Hemisphere (*hem'-is-fer*) [*ἥμι*, half; *σφαῖρα*, a ball]. Half a sphere. **H.**, **Cerebellar**, either lateral half of the cerebellum. **H.**, **Cerebral**, either lateral half of the cerebrum.

Hemisystole (*hem-e-sis'-to-le*) [*ἥμι*, half; *συστολή*, a contraction]. A peculiar kind of irregular action of the heart-muscle, in which, with every two beats of the heart, only one beat of the pulse is felt.

Hemiterata (*hem-it'-er-at-ah*) [*ἥμι*, half; *τίρας*, monster]. A class of malformations, not grave enough to be called monstrous.

Hemiteratic (*hem-it'-er-at'-ik*) [*ἥμι*, half; *τίρας*, a monster]. Pertaining to hemiterata.

Hemlock (*hem'-lok*). See *Conium*.

Hemocyanin (*hem-o-si'-an-in*) [*αἷμα*, blood; *κυανός*, blue]. A coloring matter found in the blood of certain invertebrates. It contains copper, and gives to the blood a blue color.

Hemocyte (*hem'-o-sit*) [*αἷμα*, blood; *κύτος*, a cell]. A red blood corpuscle.

Hemocytometer (*hem-o-si'-tom'-et-er*) [*αἷμα*, blood; *κύτος*, a cell; *μέτρον*, a measure]. An instrument for estimating the number of corpuscles in the blood.

Hemodromometer (*hem-o-dro-mom'-et-er*). See *Hemadromometer*.

Hemodynamometer (*hem-o-di-nam-om'-et-er*). See *Hemadynamometer*.

Hemogenesis (*hem-o-jen'-es-is*). See *Hematogenesis*.

Hemogenic (*hem-o-jen'-ik*). See *Hematogenic*.

Hemoglobin (*hem-o-glo'-bin*) [*aiua*, blood; *globus*, a round body]. The coloring-matter of the red corpuscles. It is an exceedingly complex body, containing iron; it crystallizes in rhombic plates or prisms, and is composed of hematin and a proteid substance, called globulin. It has a strong affinity for oxygen, and the greater part of the oxygen in the blood is in combination with it as oxyhemoglobin. When it gives up the oxygen to the tissues it becomes reduced hemoglobin.

Hemoglobinemia (*hem-o-glo-bin-e'-me-ah*) [*aiua*, blood; *globus*, a round body; *aiua*, blood]. A condition in which the hemoglobin is dissolved out of the red corpuscles, probably as the result of the destruction of the latter, and is held in solution in the serum. The blood is "lake-colored." It occurs in some infectious diseases, and after injecting certain substances into the blood.

Hemoglobinometer (*hem-o-glo-bin-om'-et-er*) [*aiua*, blood; *globus*, a round body; *μετρον*, a measure]. An instrument for the quantitative estimation of hemoglobin in the blood. **H. of Fleischl**; in this a certain dilution of the blood is made, and its color then compared with that of different thicknesses of a sliding wedge of red glass. **H. of Gowers**; in this the calculation is made by measuring the amount of dilution necessary to make the blood of the same shade as a standard solution of carmin in gelatin.

Hemoglobinuria (*hem-o-glo-bin-ur'-e-ah*) [*aiua*, blood; *globus*, a round body; *ουρον*, urine]. The presence of hemoglobin in the urine, due either to its solution out of the red corpuscles, or to disintegration of the red corpuscles. **H., Epidemic**, hemoglobinuria of the new-born associated with jaundice, cyanosis, and nervous symptoms. **H., Intermittent** or **H., Paroxysmal**, a form characterized by recurring periodic attacks. It is related to cold, and is also closely associated with Raynaud's disease. **H., Toxic**, that form occurring in consequence of poisoning by various substances.

Hemology (*hem-ol'-o-je*) [*aiua*, blood; *λογος*, science]. The science treating of the blood.

Hemolysis (*hem-ol'-is-is*). See *Hematolysis*.

Hemometer (*hem-om'-et-er*). See *Hemoglobinometer*.

Hemophilia (*hem-o-fi'l'-e-ah*) [*aiua*, blood; *φιλειν*, to love]. Bleeder's disease, an abnormal tendency to hemorrhage. It is usually hereditary, and though it is most common in males, the hereditary influence is transmitted through the mother.

Hemophthalmia (*hem-off-thal'-me-ah*) [*aiua*, blood; *ὀφθαλμός*, eye]. A hemorrhage into the interior of the eye.

Hempneumothorax (*hem-o-nu-mo-tho'-raks*) [*aiua*, blood; *πνευμα*, wind; *θώραξ*, the chest]. A collection of air and blood within the pleural cavity.

Hemoptysis (*hem-op'tis-is*) [*aiua*, blood; *πτειν*, to spit]. The spitting of blood from the larynx, trachea, bronchi, or lungs.

Hemorrhage (*hem'-or-aj*) [*aiua*, blood; *ῥηγνίναν*, to burst forth]. An escape of blood from the vessels, either by diapedesis through intact walls, or by rhexis through ruptured walls. **H., Accidental**, hemorrhage during pregnancy from premature detachment of the placenta when normally situated. **H., Capillary**, oozing of blood from the capillaries. **H., Concealed**, a variety of accidental hemorrhage in which the bleeding takes place between the ovum and the uterine walls, without escape from the genital tract.

H., Consecutive, one ensuing some time after injury. **H., Critical**, occurring at the turning-point of a disease. **H., Petechial**, hemorrhage under the surface in the form of minute points. **H., Postpartum**, hemorrhage occurring shortly after labor. **H., Primary**, that immediately following any traumatism. **H., Secondary**, that occurring some time after the traumatism. **H., Unavoidable**, hemorrhage from detachment of a placenta previa. **H., Vicarious**, a discharge of blood from a part owing to the suppression of a flow in another part, as vicarious menstruation.

Hemorrhagic (*hem-er-a'-jik*) [*aiua*, blood; *ῥηγνίναν*, to burst forth]. Relating to or accompanied by hemorrhage. **H. Diathesis**. See *Hemophilia*. **H., Infarct**. See *Infarct*.

Hemorrhea (*hem-or-e'-ah*) [*aiua*, blood; *ῥοια*, a flow]. A hemorrhage.

Hemorrhoid (*hem'-or-oid*) [*αιμόρροος*, flowing with blood]. A pile. An enlarged and varicose condition of the veins of the lower portion of the rectum and the tissues about the anus. **H., External**, those situated without the sphincter ani. **H., Internal**, those within the anal orifice.

Hemorrhoidal (*hem-or-oi'-dal*) [*αιμόρροος*, flowing with blood]. Pertaining to or affected with hemorrhoids. **H. Veins**, the three veins which form a plexus about the lower end of the rectum. Through them the general venous system and the portal system communicate.

Hemosiderin (*hem-o-sid'-er-in*) [*aiua*, blood; *σίδηρος*, iron]. A granular pigment, a product of the decomposition of hemoglobin. It is found where blood is extravasated in contact with active cells, and contains iron.

Hemospasia (*hem-o-spa'-se-ah*) [*aiua*, blood; *σπάειν*, to draw]. The drawing of blood from a part by dry or wet cupping.

Hemostatic (*hem-o-stat'-ik*) [*αἷμα*, blood; *στατικός*, stationary]. 1. Arresting hemorrhage. 2. An agent or remedy that arrests hemorrhage.

Hemotachometer (*hem-o-tak-om'-et-er*) [*αἷμα*, blood; *τάχος*, swiftness; *μέτρον*, a measure]. An instrument for measuring the rate of flow of blood.

Hemothorax (*hem-o-tho'-raks*) [*αἷμα*, blood; *θώραξ*, the chest]. An accumulation of blood in a pleural cavity.

Hemp [ME., *hemp*, hemp]. *Cannabis sativa*, the bast-fiber of which is used for textile purposes. **H., Indian.** See *Cannabis indica*.

Henbane. See *Hyoscyamus*.

Henle's Fenestrated Membrane [*Henle*, a German anatomist]. The layer of longitudinal elastic fibers of the inner coat of the arteries. **H.'s Layer**, a layer of flat, horny, polygonal, nonnucleated cells situated between the cuticula and the external root-sheath of a hair-bulb. **H.'s Loop**, a loop in the uriniferous tubules. **H.'s Sheath**, a connective-tissue sheath surrounding terminal nerve-fibers.

Henry's Law. See *Law*.

Hepar (*he'-par*) [*ήπαρ*, the liver]. 1. The liver. 2. A substance having the color of liver, as *H. sulphuris*.

Hepatalgia (*hep-at-āl'-je-ah*) [*ήπαρ*, liver; *ἄλγος*, pain]. Neuralgic pain in the liver.

Hepatectomy (*hep-at-ek'-to-me*) [*ήπαρ*, liver; *ἐκτομή*, a cutting out]. Excision of the liver, wholly or in part.

Hepatic (*hep-at'-ik*) [*ήπαρ*, liver]. Pertaining to the liver.

Hepatitis (*hep-at-i'-tis*) [*ήπαρ*, liver; *τις*, inflammation]. Inflammation of the liver. **H., Chronic Interstitial, cirrhosis** of the liver. **H., Acute Parenchymatous**, acute yellow atrophy of the liver. **H., Suppurative**, abscess of the liver.

Hepatization (*hep-at-iz'-a'-shun*) [*ήπαρ*, liver]. A change of a tissue into a condition in which it resembles the liver, as *H.* of the lung. **H. Gray**, that in which the hepatized lung-tissue is gray. **H. Red**, that in which it is red from an excess of blood.

Hepatocele (*hep'-at-o-sel*) [*ήπαρ*, liver; *κήλη*, a tumor]. Hernia of the liver.

Hepatocirrhosis (*hep-at-o-sir'-o'-sis*) [*ήπαρ*, liver; *κίρρως*, yellow]. Cirrhosis of the liver.

Hepatogenic, Hepatogenous (*hep-at-o-jen'-ik, hep-at-ō'-en-us*) [*ήπαρ*, liver; *γεννάω*, to beget]. Produced by or in the liver. **H. Icterus**, jaundice caused by the absorption of bile from the liver.

Hepatolithiasis (*hep-at-o-lith-i'-as-is*) [*ήπαρ*, liver; *λίθος*, a stone]. A diseased condition characterized by the formation of gall-stones in the liver.

Hepatomalacia (*hep-at-o-mal'-a'-se-ah*) [*ήπαρ*, liver; *μαλακία*, softness]. Softening of the liver.

Hepatopathy (*hep-at-ōp'-ath-e*) [*ήπαρ*, liver; *πάθος*, disease]. Any disease of the liver.

Hepatorrhaphy (*hep-at-or'-a-fe*) [*ήπαρ*, liver; *ρᾶφή*, suture]. Suture of the liver.

Hepatotomy (*hep-at-ōt'-o-me*) [*ήπαρ*, liver; *τέμνειν*, to cut]. Incision of the liver.

Hephestic (*hef-es'-tik*) [*Hephestus*, Vulcan]. Prevailing or occurring among hammermen, as *H. cramp*.

Heptad (*hep'-tad*) [*ἑπτὰ*, seven]. An element having a quantivalence of seven.

Heptane (*hep'-tān*) [*ἑπτὰ*, seven], *C₇H₁₆*. A liquid hydrocarbon of the paraffin-group, contained in petroleum and also obtained from the resin of *Pinus sabiniana* by dry distillation. It is also called abietene.

Herb (*erb* or *herb*) [*herba*, an herb]. A plant the stem of which contains but little wood and dies down to the ground at the end of the season.

Herbivorous (*her-biv'-or-us*) [*herba*, grass; *vorare*, to devour]. Living on vegetable food.

Hereditary (*he-red'-it-a-re*) [*heres*, an heir]. Transmitted from parent to offspring, as *H. disease*. **H. Ataxia.** See *Friedreich's Disease, Diseases, Table of*. **H. Syphilis.** See *Syphilis*.

Heredity (*he-red'-it-e*) [*hereditas*, heredity]. The transmission of physical or mental qualities or tendencies from ancestor to offspring; the principle or force by reason of which the offspring resembles the parent.

Hermaphroditism, or Hermaphroditism (*her-maf'-ro-dism, her-maf'-ro-dit-tism*) [*Ἑρμῆς*, Mercury; *Ἀφροδίτη*, Venus]. The coexistence in a single individual of ovaries and testicles. **H., Complex**, a condition in which there are present both the internal and external organs of either sex.

H., Lateral, a form in which male organs (especially a testicle) are more or less developed on one side, and female organs (especially an ovary) on the opposite side.

H., Spurious, a condition in which the individual is of one sex, but presents the outward signs of the other. **H., Transverse**, an instance in which the external organs indicate the one sex, and the internal the opposite. **H., Unilateral**, on one side an ovary and a testicle, on the other an ovary or a testicle.

Hermaphrodite (*her-maf'-ro-dit*) [*Ἑρμῆς*, Mercury; *Ἀφροδίτη*, Venus]. An individual affected with hermaphroditism; usually the condition is due to some congenital malformation of the genital organs, such as epispadias, hypospadias, cleft of the scrotum, etc., that makes the determination of sex somewhat doubtful.

Hermetic (*her-met'-ik*) [*Ἑρμῆς*, the god Mercury]. Protected from exposure to air; air-tight, as the H. sealing of a wound.

Hernia (*her'-ne-ah*) [*hernia*, from *ἑρῆναι*, sprout]. A protrusion of a viscus through an abnormal opening in the wall of the containing cavity; used without qualification, the word refers to hernia of the intestines. **H.**, **Abdominal**, a protrusion of a portion of the abdominal viscera through some portion of the parietes. **H. of the Bladder**, the protrusion of a part of the bladder through one of the openings of the abdominal cavity. **H. of the Brain**, or **H. cerebri**, *Fungus cerebri*, a protrusion of the brain through the skull. **H.**, **Complete**, a hernia in which the sac and its contents have passed the hernial orifice. **H.**, **Congenital**, a form of indirect inguinal hernia in which, the vaginal process of the peritoneum having remained patulous, the bowel descends at once into the scrotum, in direct contact with the testicle. **H.**, **Crural**. See *H.*, *Femoral*. **H.**, **Cystic**. Same as *Cystocele*. **H.**, **Diaphragmatic**, a protrusion of a portion of some of the abdominal viscera into the thorax, through a congenital defect in the diaphragm, or through a dilatation or laceration of one of the natural openings. **H.**, **Encysted**; in this the pouch forming the tunica vaginalis is closed at its upper end but open below. The hernia in descending along the inguinal canal, enters the scrotum behind the tunica vaginalis, and is more or less completely surrounded by its posterior layer. **H.**, **Femoral**, a hernia through the femoral canal, the tumor appearing on the upper inner aspect of the thigh, below Poupart's ligament. **H.**, **Incarcerated**, a hernia which has become occluded by the accumulation of gas, feces, or undigested food, thus causing obstruction of the bowels. **H.**, **Incomplete**, one that has not entirely passed through the hernial orifice. **H.**, **Infantile**. See *H.*, *Encysted*. **H.**, **Inguinal**, a hernia occupying the inguinal canal. **H. of Iris**, a protrusion of a portion of the iris after iridectomy, trauma, etc. **H.**, **Irreducible**, one in which the protruded viscus cannot be returned by manipulation. **H.**, **Ischiatic**, a protrusion of the bowel through the great sacrosacral foramen. **H.-knife**, a probe pointed knife for incising the constriction of a hernial sac. **H.**, **Labial**, a protrusion of the bowel between the vagina and the ramus of the ischium into a labium majus. **H. of Littre**, a hernia in which only a diverticulum of the bowel is affected. The term is improperly applied to Richter's H. **H.**, **Obstructed**. See *H.*, *Incarcerated*. **H.**, **Obturator**, a protrusion of bowel through the obturator foramen. **H.**, **Omental**, a

hernia containing omentum; *epiplocele*. **H.**, **Perineal**, a protrusion of the abdominal contents between the fibers of the levator ani muscle in front of or to one side of the anus. **H.**, **Pudendal**. Same as *H.*, *Labial*. **H.**, **Reducible**, one that may be returned by manipulation. **H.**, **Retroperitoneal**, a hernia in which the intestine lodges in the fossa duodenojejunalis. **H.**, **Richter's**, one in which only a portion of the caliber of the gut enters the orifice. **H.**, **Scrotal**, that form of inguinal hernia in which the protrusion has entered the scrotum. **H.**, **Strangulated**, a hernia which is so tightly constricted at its neck as to interfere with its return, with the circulation of blood, and the passage of feces. **H.**, **Umbilical**, a protrusion of the abdominal contents through the umbilicus. **H.**, **Ventral**, the name applied to protrusions of the abdominal contents through the abdominal walls in situations not usually subject to herniæ.

Hernial (*her'-ne-al*) [*hernia*, a rupture]. Pertaining to hernia. **H. Sac**, the diverticulum of the peritoneum which the hernia pushes before it or into which it descends.

Herniopuncture (*her'-ne-o-punk'-chur*) [*hernia*, hernia; *punctura*, a pricking]. The puncture of a hernia.

Herniotome (*her'-ne-o-tōm*) [*hernia*, hernia; *τομή*, cutting]. A hernia-knife.

Herniotomy (*her'-ne-o'-o-me*) [*hernia*, hernia; *τμήναι*, to cut]. Operation for the relief of hernia by section of the constriction.

Heroic (*he-ro'-ik*) [*ἥρωας*, a hero]. Bold or daring; rash or unusually severe, as heroic treatment or measures.

Herophilus, **Torcular of**. See *Torcular*.

Herpes (*her'-pez*) [*ἑρπης*; *ἑρπεω*, to creep]. An acute inflammatory affection of the skin or mucous membrane, characterized by the development of groups of vesicles on an inflammatory base. **H. circinatus**. See *Tinea circinata*. **H. facialis**, herpes febrilis; an acute, noncontagious, inflammatory disorder of the skin that appears in the form of one or more groups of vesicles. It is commonly called fever-blisters, a form of herpes appearing especially on the lips. It is frequent in "cold," malaria, croupous pneumonia, and cerebrospinal meningitis. **H. febrilis**. See *H. facialis*. **H. Iris**, a form of erythema with vesicles growing in a ring. It is usually seen on the backs of the hands and feet. **H. labialis**. See *H. facialis*. **H. preputialis**, **H. pro-genitalis**, a form of herpes in which vesicles occur upon the glans and prepuce, from the size of a pin's head to that of a small pea. **H. tonsurans**. See *Tinea tonsurans*. **H. zoster**, *zoster*, *zona*, shingles, *ignis sacer*; herpes in which the lesions are distributed in relation

to the course of a cutaneous nerve, and, as a rule, unilateral. They are usually seen in the line of the intercostal nerves, but may follow the course of any nerve. The outbreak of the eruption is generally preceded by severe neuralgic pain.

Herpetic (*her-pet'-ik*) [*ἑρπης*, herpes]. Pertaining to herpes. **H. Sore-throat.** See *Tonsillitis, Herpetic*.

Herpetiform (*her-pet'-if-orm*) [*ἑρπης*, herpes; *forma*, form]. Resembling herpes.

Hesselbach's Triangle. See *Triangles, Table of*.

Heteradelphus (*het-er-ad-el'-fus*) [*ἑτερος*, other; *ἀδελφός*, brother]. A joined twin-monster, consisting of an autositic monster with an attached parasite, the head of the latter being absent.

Heteradenia (*het-er-ad-el'-ne-ah*) [*ἑτερος*, other; *ἀδὴν*, gland]. 1. Normal glandular structure occurring in a part normally not provided with glands. 2. Glandular structure departing from the normal type.

Heteradenic (*het-er-ad-el'-nik*) [*ἑτερος*, other; *ἀδὴν*, gland]. Pertaining to or consisting of tissue that is unlike normal glandular tissue, or to glandular tissue occurring in an abnormal place.

Heteralius (*het-er-al'-le-us*) [*ἑτερος*, other; *ἄλως*, a disc]. A double monster in which the parasite is very incomplete, and with no direct connection with the umbilical cord of its host.

Hetero- (*het'-er-o-*) [*ἑτερος*, other]. A prefix denoting unlikeness.

Heteroalbumose (*het-er-o-al'-bu-mōs*) [*ἑτερος*, other; *albumose*]. A variety of albumose soluble in salt-solutions, insoluble in water, and precipitated by saturation with sodium chlorid or magnesium sulphate.

Heteroautoplasty (*het-er-o-au'-to-plas-te*) [*ἑτερος*, other; *αὐτός*, self; *πλάσσειν*, to form]. Grafting of skin from one person upon the body of another.

Heteroblastic (*het-er-o-blas'-tik*) [*ἑτερος*, other, different; *βλαστός*, germ, bud]. Arising from tissue of a different kind.

Heterodymus (*het-er-od'-im-us*) [*ἑτερος*, other; *δίδυμος*, twin]. A double monster, the accessory part being but an imperfect head, with a neck and thorax by which it is implanted in the anterior abdominal wall of its host.

Heterogeneous (*het-er-o-je'-ne-us*) [*ἑτερος*, other; *γένος*, kind]. Differing in kind or nature; composed of different substances; not homogeneous.

Heterogenesis (*het-er-o-je'-es-is*) [*ἑτερος*, different; *γένεσις*, generation]. A mode of reproduction in which the living parent gives rise to offspring that pass through totally different series of states from those exhibited by

the parents, and do not return into the cycle of the parents.

Heteroinfection (*het-er-o-in-fek'-shun*) [*ἑτερος*, other; *inficere*, to taint]. 1. Infection transmitted by a person who is himself not affected. 2. Infection of any organism by a poison not produced within itself; opposed to autoinfection.

Heteroinoculation (*het-er-o-in-ok'-u-la'-shun*) [*ἑτερος*, other; *inoculatio*, an ingrafting]. Inoculation of one person by another.

Heterologous (*het-er-ol'-o-gus*) [*ἑτερος*, other; *λόγος*, relation]. Differing in structure or form from the normal. **H. tumors**, tumors constituted of a different tissue from that of the part in or on which they are situated.

Heteromorphous (*het-er-o-mor'-fus*) [*ἑτερος*, other; *μορφή*, form]. Differing from the normal in form.

Heteronymous (*het-er-on'-im-us*) [*ἑτερος*, other; *ὄνομα*, name]. On opposite sides.

Heteropagus (*het-er-op'-ag-us*) [*ἑτερος*, other; *πηγνύναι*, to unite]. A double monster in which the parasite, having a head and extremities, is attached to the anterior abdominal wall of its host.

Heteropathy (*het-er-op'-ath-e*) [*ἑτερος*, other; *πάθος*, affection]. 1. The treatment of a disease by inducing a different morbid condition to neutralize it. 2. Abnormal reaction to stimulus or irritation.

Heterophoralgia (*het-er-o-for-al'-je-ah*) [*ἑτερος*, other; *φορός*, tending; *ἄλγος*, pain]. Eye-strain or ocular pain caused by heterophoria.

Heterophoria (*het-er-o-for-re-ah*) [*ἑτερος*, different; *φορός*, tending]. A relation of the visual lines of the two eyes other than that of parallelism. It may be **Esophoria**, a tending of the lines inward; or **Exophoria**, outward; or **Hyperphoria**, a tending of the right or left visual line in a direction above its fellow; **Hyperesophoria**, a tending of the visual lines upward and inward; **Hyperexophoria**, upward and outward.

Heteroplasia (*het-er-o-pla'-se-ah*) [*ἑτερος*, other; *πλάσις*, shaping]. The presence in a part of a tissue that does not belong there normally.

Heteroplasty (*het-er-o-plas-te*) [*ἑτερος*, other; *πλάσσειν*, to form]. 1. Heteroplasia. 2. The operation of grafting parts taken from another species.

Heterotaxis (*het-er-o-taks'-is*) [*ἑτερος*, other; *τάξις*, order]. The anomalous disposition or transposition of organs.

Heterotopia (*het-er-o-to'-pe-ah*) [*ἑτερος*, other; *τόπος*, place]. A misplacement of normal tissue, especially a congenital malformation of the brain, in which masses of gray matter are found transplanted into the white.

Heterotrophia, or **Heterotrophy** (*het-er-o-tro'-fe-ah*, or *het-er-at'-ro-fe*) [*ἑτερος*, other; *τροφή*, sustenance]. Any perversion or disorder of nutrition.

Heterotropia (*het-er-o-trôp'-e-ah*). See *Strabismus*.

Heterotypus (*het-er-o-tîl'-pus*) [*ἑτερος*, different; *τύπος*, a type]. A double monster having the parasitic fetus hanging from the ventral wall of the principal subject.

Heteroxanthin (*het-er-o-zan'-thin*) [*ἑτερος*, other; *ξανθός*, yellow], $C_{40}H_{56}N_4O_2$. A leukomatin that can be isolated in crystalline form from urine.

Heubner's Disease. Syphilitic disease of the cerebral arteries. See *Diseases, Table of*.

Hexad (*heks'-ad*) [*ἑξάς*, six]. An element the atom of which has a quantivalence of six.

Hexane (*heks'-ân*) [*ἑξάς*, six], C_6H_{14} . The sixth member of the paraffin series of hydrocarbons. It is a liquid, boiling at about $71^{\circ}C.$, found in various natural oils.

Hexatomic (*heks-at-om'-ik*) [*ἑξάς*, six; *ἄτομος*, an atom]. Consisting of six atoms; also applied to atoms that are hexavalent, and to alcohols or other compounds having six replaceable hydrogen atoms.

Hexavalent (*heks-av'-al-ent*) [*ἑξάς*, six; *valens*, having power]. Having the same combining power as six hydrogen atoms.

Hey's Ligament. See *Ligament*.

Hiatus (*hi-a'-tus*) [*hiare*, to gape]. A space or opening. Also, the vulva. **H. of Fallopius**, a shallow groove on the petrous portion of the temporal bone for the passage of the petrosal branch of the Vidian nerve.

Hiccup, or **Hiccough** (*hik'-up*) [*hic*, a mimic word; *cough*, a cough]. A spasmodic contraction of the diaphragm causing inspiration, followed by a sudden closure of the glottis. Synonym *Singultus*.

Hide-bound Disease. See *Scleroderma*.

Hidropoiesis (*hid-ro-poi-e'-sis*) [*ιδρώς*, sweat; *ποιέω*, to make]. The formation of sweat.

Hidropoietic (*hid-ro-poi-et'-ik*) [*ιδρώς*, sweat; *ποιέω*, to make]. Relating to hidropoiesis.

Hidrosadenitis (*hid-rôs-ad-en-î'-tis*) [*ιδρώς*, sweat; *αδην*, gland; *τις*, inflammation]. Inflammation of the sweat glands.

Hidrosis (*hid-ro'-sis*) [*ιδρώς*, sweat]. 1. The formation and excretion of sweat. 2. Abnormally profuse sweating. 3. Any skin-disease marked by disorder of the sweat-glands.

Hiera picra (*hi'-er-ah pî'-kraîk*) [L., sacred bitters]. Pulvis aloes et canellæ. See *Aloes*.

Highmore, Antrum of (*hi'-môr*) [W. Highmore, an English physician]. A cavity in the superior maxillary bone, communicating with the middle meatus of the nose. **H., Body of**, a thickening of the tunica albu-

ginea at the posterior part of the testis, from which connective-tissue septa diverge.

Hilum (*hi'-lum*) [*hilum*, a little thing]. A pit, recess, or opening in an organ, usually for the entrance and exit of vessels or ducts.

Hind (*hind*) [AS., *hindan*, at the back of]. Pertaining to the rear or posterior extremity. **H.-brain**, a division of the brain in the embryo, that becomes the cerebellum and the medulla oblongata. **H.-gut**, that part of the embryonic intestine from which the cecum, vermiform appendix, colon, and rectum are developed.

Hinge-joint (*hinj'-joint*). See *Diarthrosis*.

Hip [AS., *hype*, hip]. 1. The upper part of the thigh at its junction with the buttocks. 2. The hip-joint. **H.-bath**. See *Bath*. **H.-bone**, the ischium. **H.-girdle**, the pelvic arch. **H.-joint**, the articulation of the femur with the haunch-bone or innominate bone. **H.-joint Disease**, coxitis, an inflammation of the hip-joint, usually tuberculous, and occurring most commonly in the young, and, according as it begins in the head of the femur, the acetabulum, or in the synovial membrane and proper structures of the joint, divided into femoral, acetabular, and arthritic. Symptoms: Shuffling gait, pain often referred to the inner side of the knee, pain in the hip elicited by jarring the heel, deformity, abduction and eversion of the thigh, slight flexion of the knee, and arching of the lumbar spine; later, adduction and inversion of the thigh, with flexion of the knee and shortening of the limb. Suppuration with formation of fistulae occurs in the advanced stages.

Hippocampal (*hip-o-kam'-pal*) [*ἵππος*, horse; *κάμπος*, a sea-monster]. Relating to the hippocampus.

Hippocampus (*hip-o-kam'-pus*) [*ἵππος*, horse; *κάμπος*, a sea-monster]. A name applied to two elevations, H. major and H. minor, the former situated in the middle, and the latter in the posterior horn of the ventricles of the brain.

Hippocratic (*hip-o-krat'-ik*) [*Hippocrates*, a Greek physician of the fifth century B. C.]. Described by Hippocrates. **H. Expression**, an anxious, pinched expression of the countenance, described as characteristic of peritonitis. **H. Sound**. See *Succession-sound*.

Hippuric Acid (*hip-u'-rik*). See *Acid*.

Hippus (*hip'-us*) [*ἵππος*, horse, from analogy to the movements of this animal]. Spasmodic pupillary movement, independent of the action of light.

Hirsute (*her-sut'*) [*hirsutus*, shaggy]. Shaggy, hairy.

Hirsuties (*her-su'-te-tes*) [*hirsutus*, shaggy]. Excessive growth of hair.

Histioid (*his'-te-oid*). See *Histoid*.

Histochemistry (*his-to-kem'-is-tre*) [*ιστός*, tissue; *χημεία*, chemistry]. The chemistry of the histologic elements of the body.

Histodialysis (*his-to-di-al'-is-is*) [*ιστός*, web; *διάλυσις*, a resolution]. The dissolution of organic tissue.

Histogenesis (*his-to-jen'-es-is*) [*ιστός*, web; *γένεσις*, generation]. The formation of tissues.

Histogenetic (*his-to-jen-et'-ik*) [*ιστός*, web; *γεννάν*, to produce]. Relating to histogenesis.

Histohematin (*his-to-hem'-at-in*) [*ιστός*, web; *αἷμα*, blood]. A pigment found in muscles, suprarenal capsules, and other organs, and believed to have a respiratory function.

Histoid (*his'-toid*) [*ιστός*, web; *εἶδος*, likeness]. Composed of only one kind of tissue.

Histologic (*his-to-log'-ik*) [*ιστός*, tissue; *λόγος*, science]. Relating to histology.

Histology (*his-tol'-o-je*) [*ιστός*, tissue; *λόγος*, science]. The minute anatomy of tissues.

Histolysis (*his-tol'-is-is*) [*ιστός*, tissue; *λύσις*, dissolution]. Disintegration and dissolution of organic tissue.

Histonomy (*his-ton'-o-me*) [*ιστός*, tissue; *νόμος*, a law]. The laws of the development and arrangement of organic tissue.

Histopathology (*his-to-path-ol'-o-je*) [*ιστός*, tissue; *πάθος*, disease; *λόγος*, science]. The study of minute pathologic changes in tissues.

Histophysiology (*his-to-fiz-e-ol'-o-je*) [*ιστός*, web, tissue; *φύσις*, nature; *λόγος*, science]. The science of the functions of the various tissues.

Histotome (*his'-to-tom*) [*ιστός*, tissue; *τέμνειν*, to cut]. An apparatus for cutting tissue for the study of its minute structure; a microtome.

Histotomy (*his-tot'-o-me*) [*ιστός*, tissue; *τέμνειν*, to cut]. The dissection of tissues.

Histrionic (*his-tre-on'-ik*) [*histrion*, an actor]. Dramatic. **H. Mania**, insanity with affectation and lofty manner. **H. Muscles**, the muscles of expression of the face. **H. Spasm**, spasm of the muscles of expression.

Hives (*hivz*) [origin uncertain]. 1. Urticaria. 2. In Great Britain the term is also applied to croup, laryngitis, and to chicken-pox.

Hob-nail Liver. The liver of advanced atrophic cirrhosis, so called on account of the small projections on the surface. It is called also gin drinker's liver.

Hodgkin's Disease. See *Diseases, Table of*.

Hoffmann's Anodyne (*hof'-mans an'-o-din*). Spiritus atheris compositus (U. S. P.), compound spirit of ether. It is anodyne, stimulant, and antispasmodic, and is used in nervous irritation, angina pectoris, and asthma. Dose ℥xxx-fʒij (2.0-8.0).

Hollyhock (*hol'-e-hok*). See *Althra*.

Holmgren's Test. A test for color-blindness. The patient is requested to match worsted skeins of different colors, and if color-blind, always selects characteristic shades.

Holoblastic (*hol-o-blas'-tik*) [*ὅλος*, whole; *βλαστός*, germ]. Applied to ova in which the entire yolk is included in the process of segmentation; one in which there is no separate food-yolk.

Homatropin (*ho-mat'-ro-pin*), $C_{10}H_{21}NO_3$. An alkaloid derived from atropin. It causes dilatation of the pupil and paralysis of accommodation like atropin, but its effects pass off more quickly, usually in two or three days. The hydrobromate is the salt generally employed.

Homeopathy (*ho-me-op'-ath-e*) [*ὁμοιος*, like; *πάθος*, ailment or disease]. A system of treatment of disease by the use of agents that, administered in health, would produce symptoms similar to those for the relief of which they are given.

Homocentric (*ho-mo-sen'-trik*) [*ὁμός*, same; *κέντρον*, center]. Concentric; having the same center. **H. Rays**, light rays that have a common focus or are parallel.

Homocerebrin (*ho-mo-ser'-e-brin*) [*ὁμός*, like; *cerebrum*, cerebrum]. A substance derived from brain-tissue, closely resembling cerebrin, but more soluble in alcohol.

Homæo- (*ho'-me-o-*). For words thus beginning, see *Homæo*.

Homogeneous (*ho-mo-je'-ne-us*) [*ὁμός*, like; *γένος*, kind]. Having the same nature or qualities; having a uniform character in all parts.

Homogenesis (*ho-mo-jen'-es-is*) [*ὁμός*, like; *γεννάν*, to beget]. Reproduction in which a parent gives rise to offspring that pass through the same cycle of changes as itself.

Homogentisic Acid (*hom-o-jen-tiz'-ik*). See *Acid*.

Homologous (*ho-mol'-o-gus*) [*ὁμός*, the same; *λόγος*, relation]. Corresponding in structure, either directly or as referred to a fundamental type. In chemistry, being of the same type or series; differing by a multiple or an arithmetic ratio in certain constituents. **H. Tissues**, those identical in type of structure. **H. Tumor**, a tumor consisting of tissue identical with that of the organ whence it springs.

Homologue (*hom'-o-log*) [*ὁμός*, same; *λόγος*, proportion]. An organ which has the same relative structure, position, or development as another. The same organ in different organisms under every variety of form and function.

Homology (*ho-mol'-o-je*) [*ὁμός*, same; *λόγος*, proportion]. The quality of being homologous; also, the morphologic identity of parts or organs in different animals.

Homonymous (*ho-mon'-im-us*) [*ὁμός*, same; *ὄνομα*, a name]. 1. Having the same sound or name; having the same relative position.

H. Diplopia, a form of diplopia in which the image seen by the right eye is on the right side and that seen by the left eye is on the left side. **H.** Hemianopia. See *Hemianopia*.

Homothermic (*ho-mo-ther'-mik*) [*ὁμός*, same; *θερμη*, heat]. Having a uniform temperature.

Homotonic (*ho-mo-ton'-ik*) [*ὁμός*, same; *τόνος*, tone, tension]. Having a uniform or even course.

Homotype (*ho'-mo-tip*) [*ὁμός*, same; *τύπος*, a pattern]. A part corresponding and similar to another part, as the humerus to the femur.

Honduras Bark (*hon-du'-ras bark*). *Cas-cara amara*.

Honey (*hun'-e*) [ME., *hony*, honey]. See *Mel*.

Hook [AS., *hōc*, a hook]. A curved instrument. **H.**, Blunt, an instrument for exercising traction upon the fetus in an arrested breech-presentation. **H.**, Malgaigne's, two pair of hooks connected by a screw for approximating the fragments of a fractured patella. **H.**, Tyrrel's, a blunt, slender hook for operations upon the eye.

Hop. See *Humulus*.

Hope's Camphor-mixture. A mixture containing nitric acid, camphor-water, and tincture of opium. It is used in the treatment of serous or choleraic diarrheas.

Hordeolum (*hor-de'-o-lum*) [*hordeum*, a grain of barley]. A sty; a furuncular inflammation of the connective tissue of the lids, near a hair-follicle.

Hordeum (*hor'-de-um*) [L.]. Barley.

Horehound (*hor'-hoy-und*). See *Marrubium*.

Horizontal (*hor-iz-on'-tal*) [*ὁριζων*, the horizon]. Parallel to the horizon.

Horn. See *Cornu*.

Horner's Muscle. See *Muscles*, Table of.

Horny (*horn'-e*) [ME., *horn*, horn]. Composed of, or resembling, horn.

Horo-pter (*hor'-op'-ter*) [*ὅρος*, boundary; *ὁπτήρ*, an observer]. The sum of all the points seen singly by the two retinæ while the fixation-point remains stationary.

Horripilation (*hor-ip-il'-at-shun*) [*horre-re*, to stand on end; *pilus*, the hair]. Erection of the hairs of the skin produced by the contraction of arrectores pili muscles.

Horse-radish (*hors'-rad-ish*). *Cochlearia armoracia*, a plant of the order Crucifere. The root (*A Armoracia radix*, U. S. P.) contains a volatile oil, and is a gastric stimulant and diuretic. It is chiefly used as a condiment, but has been employed in medicine in dropsy, chronic rheumatism, and scurvy. Dose of the root ʒ ss (2.0) or more.

Horseshoe-kidney. A kidney having somewhat the shape of a horseshoe, due to a fusion of the two kidneys at one of their ends, usually the lower.

Horseshoe-magnet. "A magnet bent in the shape of a horseshoe.

Horseweed. 1. The *Collinsonia canadensis*, an indigenous plant, the root of which (in decoction) is used in cystitis, leukorrhea, dropsy, gravel, etc. 2. The *Erigeron canadense*.

Horsley's Method. A method of determining the position of the fissure of Rolando by means of an instrument called a cyrtometer, encircling the head, and having an arm fixed at an angle of 67°, which indicates the position of the fissure. **H.** Test, a test for glucose. The urine rendered alkaline is boiled with potassium dichromate; if sugar is present, a green color is developed.

Hospital (*hos'-pit-al*) [*hospes*, a guest]. A building for the care and treatment of sick or infirm persons. **H.** Gangrene, a contagious, phagedenic gangrene occasionally attacking wounds or open sores. It is confined mainly to military hospitals, and is of microbic origin.

Hospitalism (*hos'-pit'-al-izm*) [*hospes*, a guest]. The morbid conditions arising from the gathering of diseased persons in a hospital.

Host (*hōst*) [*hostis*, a stranger, a landlord]. The organic body upon which parasites live.

Hot [ME., *hot*, hot]. Having or yielding the sensation of heat; stimulating; biting.

H.-air Bath. See *Bath*. **H. Bath**. See *Bath*.

Hottentotism (*hōt'-n-tot-izm*). An extreme form of congenital stammering.

Hour-glass Contraction. A contraction of a hollow organ, as the uterus or stomach, near the middle, producing a condition resembling an hour-glass.

Housemaid's Knee. A chronic inflammation of the bursa in front of the patella with an accumulation of serous fluid.

Howship's Lacunæ, or Pits [John Howship, an English surgeon]. Minute depressions or pits in bone undergoing absorption, produced by the action of osteoclasts.

Huguier's Canal. See *Canal*.

Humeral (*hu'-mer-al*) [*humerus*, the shoulder]. Pertaining to the humerus.

Humerus (*hu'-mer-us*) [L.]. 1. The bone of the upper arm. 2. The shoulder.

Humidity (*hu-mid'-it-ē*) [*humor*, moisture]. The state or quality of being moist; moisture; dampness. **H.**, Absolute, the actual amount of water present in the air at any moment. **H.**, Relative, the relative amount of water present in air as compared to what the air would contain at the existing temperature were its condition that of saturation.

Humor (*hu'-mor*) [L., moisture]. 1. Any fluid or semi-fluid part of the body. **H.**, **Aqueous**, the transparent fluid of the anterior chamber of the eye. **H.**, **Crystalline**. See *Lens*, *Crystalline*. **H.**, **Vitreous**. The transparent gelatin-like substance filling the greater part of the globe of the eye. 2. Disposition, temperament, as the four humors of Galen, the choleric, melancholy, phlegmatic, and sanguine.

Humoral (*hu'-mor-al*) [*humor*, moisture]. Pertaining to the natural fluids of the body. **H. Pathology**, that system of pathology according to which all diseases result from a disordered or abnormal condition of the fluids or humors of the body. **H. Theory**, that theory which ascribes the production of immunity to the antitoxic or bactericidal action of the fluids of the body.

Humpback (*hump'-bak*). See *Kyphosis*.

Humulus (*hu'-mu-lus*) [L.]. Hop. The fruit-cones or strobiles of *H. lupulus*, which yield a powder, lupulin, a volatile oil, and tannin. Hops are tonic and slightly narcotic, and are used internally in dyspepsia, delirium tremens, and insomnia; locally, as emollient poultices. **H.**, **Tinct.** Dose ʒj-ij (4.0-8.0). **Lupulinum**, the glandular powder. Dose gr. v-xv (0.32-1.0). **Lupuli, Ext.** (B. P.). Dose gr. v-x (0.32-0.65). **Lupuli, Infusum** (B. P.). Dose f ʒj-ij (32.0-64.0).

Hunchback (*hunch'-bak*). See *Kyphosis*.

Hunger (*hung'-ger*) [AS., *hungor*, hunger]. A condition marked by a sensation of emptiness of the stomach, with a longing for food.

Hunter's Canal [John *Hunter*, an English scientist and surgeon]. See *Canal*. **H.'s Method**, a method of treating aneurysm by ligating the artery on the proximal side of the sac.

Hunarian Chancre. See *Chancre*.

Huntingdon's Chorea. See *Chorea*.

Hunyadi Janos Water (*hun-yah'-de yah'-nos*). An aperient mineral water from Hungary, containing the sulphates of potassium, sodium, and magnesium, sodium chlorid and carbonate, iron oxid, and alumina.

Hutchinson's Teeth [Jonathan *Hutchinson*, an English physician]. A notched or furrowed condition of the free edges of the permanent teeth, especially the central incisors of the upper jaw, due to inherited syphilis.

Huxley's Layer [Thomas *Huxley*, an English biologist]. A layer of nucleated, elongated, polygonal cells lying within Henle's layer of the inner root-sheath of hairs.

Huygenian Ocular (*hi'-je'-ne-an*). See *Ocular*.

Hyalin (*hi'-al-in*) [*ὑαλος*, glass]. A translucent substance forming the walls of hydatid cysts.

Hyaline (*hi'-al-in*) [*ὑαλος*, glass]. Resembling glass. **H. Cartilage**. See *Cartilage*. **H. Cast**, or **Cylinder**, a clear, nearly transparent urinary tube-cast. **H. Degeneration**. See *Degeneration*.

Hyalitis (*hi'-al-i'-tis*) [*ὑαλος*, glass; *ιτις*, inflammation]. Inflammation of the hyaloid membrane of the vitreous humor.

Hyaloid (*hi'-al-oid*) [*ὑαλος*, glass; *ειδος*, form]. Transparent; glass-like. **H. Artery**, a branch of the arteria centralis retinae in the embryo, traversing the vitreous humor to the posterior capsule of the lens. **H. Membrane**, a delicate, transparent membrane surrounding the vitreous humor, except in front, where it becomes fibrous and strong and forms a leaflet of the zonula of Zinn.

Hyaloplasm (*hi'-al-o-plasm*) [*ὑαλος*, glass; *πλάσμα*, plasm]. The fluid portion of the protoplasm of a living cell.

Hybrid (*hi'-brid*) [*ὑβρις*, insult]. The offspring of two individuals of distinct but closely related species.

Hydantoin (*hi-dan-to-in*), $C_6H_8N_2O_2$. Glycolyl urea. A crystalline substance derived from allantoin and related to urea.

Hydatid (*hi'-dat-id*) [*ὑδαρίς*, vesicle]. A cyst-like body with clear contents, especially that formed by the larva of the *Tænia echinococcus*. **H. Disease**, a disease characterized by the presence in various portions of the body of cysts containing the embryo of the *Tænia echinococcus*. **H. Fremitus**, **H. Thrill**, a fremitus occasionally obtained on palpating an hydatid cyst. **H. Mole**. See *Chorion*, *Cystic Degeneration of*, and *Mole*. **H. of Morgagni**, a small cyst connected with the fimbriated extremity of the Fallopian tube. In the testicle it is found between the testicle proper and the epididymis. It represents the remains of the Müllerian duct.

Hydatidiform (*hi-dat-id'-if-orm*) [*ὑδαρίς*, hydatid; *forma*, form]. Having the form of an hydatid. **H. Mole**. See *Chorion*, *Cystic Degeneration of*.

Hydracetic (*hi-dras'-et-in*), $C_6H_5N_2H_2(C_2H_5O)$. A coal-tar derivative. It is antipruritic and antirheumatic. Dose gr. ½ to ij (0.032-0.13).

Hydracid (*hi-dras'-id*) [*ὑδωρ*, water; *ατιδ-um*, acid]. An acid formed by a combination of hydrogen and an acid element or radicle other than oxygen.

Hydræmia (*hi-dre'-me-ah*). See *Ilydremia*.

Hydragogue (*hi'-drag-og*) [*ὑδωρ*, water; *αγωγος*, leading]. 1. Expelling water. 2. A purgative that causes copious liquid discharges.

Hydramnios (*hi-itrant'-ne-os*) [*ὑδωρ*, water; *αμνιον*, fetal membrane]. An abnormal amount of amniotic fluid.

Hydrangea (*hi-dran'-je-ah*) [*ὑδωρ*, water;

ἀγγειον, vessel]. A genus of saxifragaceous shrubs. The root of *H. arborescens* contains hydrangin, and is employed in lithiasis.

Hydrargism, Hydrargyria, Hydrargyriasis, Hydrargyris (*hi-drar'-gism, hi-drar-jir'-e-ah, hi-drar-jir'-i'-as-is, hi-drar'-jir-ism*). Chronic mercurial poisoning. See *Mercurialism*.

Hydrargyrum (*hi-drar'-jir-um*) [*ὑδράργυρος*, mercury]. See *Mercury*.

Hydrarthrosis (*hi-drar'-thro'-sis*) [*ὑδάρθρωσις*, water; *ἀρθρῶν*, joint]. An accumulation of fluid in a joint.

Hydrastis (*hi-dras'-tis*). Golden seal, a plant of the order Ranunculaceae. The rhizome and roots of *H. canadensis*, which contain the alkaloids hydrastin and berberin. It is tonic, antiperiodic, cholagogue, and diuretic, and has been employed in leukorrhea, cystitis, constipation, menorrhagia, gonorrhea, dyspepsia, etc. Dose of the hydrastin of commerce, an impure body, gr. v-x (0.32-0.65); of the alkaloid, gr. 1-3 (0.008-0.022). *H.*, *Ext.*, *Fld.*, Dose 13 ss-j (2.0-4.0). *H.*, *Glyceritum*. Dose 13 ss-j (2.0-4.0).

Hydrate (*hi'-drat'*) [*ὑδρῶς*, water]. 1. A compound which may be considered to be formed by the replacement of half the hydrogen of one or more molecules of water by an electropositive element or radicle. 2. A salt containing water of crystallization.

Hydrated (*hi'-dra-ted'*) [*ὑδρῶς*, water]. Combined with water.

Hydration (*hi-dra'-shun*) [*ὑδρῶς*, water]. The process of combining with water, or of converting into a hydrate.

Hydrazin (*hi-dras'-in*) [*ὑδρῶς*, water; *azotum*, nitrogen], H_4N_2 . 1. Diamid; a colorless, stable, gas, soluble in water, having a peculiar odor, and a strong alkaline reaction. 2. One of a class of bodies derived from hydrazin by replacing one or more of its hydrogen atoms by a radicle.

Hydremia (*hi-dre'-me-ah*) [*ὑδρῶς*, water; *αἷμα*, blood]. A watery condition of the blood.

Hydrencephalocoele (*hi-dren-sef'-al-o-sel*) [*ὑδρῶς*, water; *ἐνδοφθαλμῶς*, brain; *κῆλη*, hernia]. Hernia of the brain, in which the tumor is in part composed of a watery fluid.

Hydrencephalus (*hi-dren-sef'-al-us*). See *Hydrocephalus*.

Hydric (*hi'-drik*) [*ὑδρῶς*, water]. Containing water.

Hydrid (*hi'-drid'*) [*ὑδρῶς*, water]. A chemic compound containing hydrogen united to an element or radicle.

Hydriodic (*hi-dre-od'-ik*) [*ὑδρῶς*, water; *ἰώδης*, like a violet]. Containing hydrogen and iodine. *H. Acid*, III, a heavy, colorless gas, with a suffocating odor and an acid reaction. The syrup of hydriodic acid (Syrupus

acidi hydriodici, U.S.P.) contains 10 per cent. of absolute H.I. Dose ℥xx-xl (1.3-2.6). It is used as an alterative in scrofula, rickets, etc.

Hydro- (*hi'-dro-*) [*ὑδρῶς*, water]. A prefix signifying combined with water or with hydrogen.

Hydroa (*hi-dro'-ah*) [*ὑδρῶς*, water; *ὄν*, egg]. A chronic inflammatory disease of the skin characterized by erythema, papules, pustules, vesicles, bullae, or combinations of these, and by intense itching. It is also known as dermatitis herpetiformis or pemphigus pruriginosus.

Hydrobilirubin (*hi-dro-bil-e-rui'-bin*) [*ὑδρῶς*, water; *bilis*, bile; *ruber*, red], $C_{23}H_{40}N_4O_7$. A brown-red pigment formed by treating a solution of bilirubin with sodium-amalgam. It is probably identical with stercobilin, the coloring-matter of the feces, and urobilin, the pigment of the urine.

Hydrobromate (*hi-dro-bro'-mat'*) [*ὑδρῶς*, water; *βρωμῶς*, a stench]. A salt of hydrobromic acid.

Hydrobromic (*hi-dro-bro'-mik*) [*ὑδρῶς*, water; *βρωμῶς*, stench]. Composed of hydrogen and bromine. *H. Acid*. See *Acid, Hydrobromic*. *H. Ether*, C_2H_5Br . See *Ethyl Bromid*.

Hydrobromid (*hi-dro-bro'-mid*). Same as *Hydrobromate*.

Hydrocarbon (*hi-dro-kar'-bon*) [*ὑδρῶς*, water; *carbo*, charcoal]. Any compound composed of hydrogen and carbon. All other organic compounds may be considered as derivatives of the hydrocarbons. The chief derivatives are (1) those containing hydrogen; (2) those containing oxygen; (3) those containing sulphur; and (4) those containing nitrogen. The principal hydrocarbons are: (1) the saturated hydrocarbons of the paraffin or marsh-gas series (C_nH_{2n+2}); (2) the unsaturated hydrocarbons of the olefin series (C_nH_{2n}); and (3) the benzene series or aromatic hydrocarbons (C_nH_{2n-6}).

Hydrocele (*hi'-dro-sel*) [*ὑδρῶς*, water; *κῆλη*, tumor]. A collection of serous fluid about the testicle or spermatic cord.

Hydrocephalic (*hi-dro-sef'-al'-ik*) [*ὑδρῶς*, water; *κεφαλή*, head]. Pertaining to or affected with hydrocephalus. *H. Cry*, the loud cry of a child, indicating pain in the head.

Hydrocephalocoele (*hi-dro-sef'-al-o-sel*) [*ὑδρῶς*, water; *κεφαλή*, head; *κῆλη*, a tumor]. See *Hydrencephalocoele*.

Hydrocephaloid (*hi-dro-sef'-al-oid*) [*ὑδρῶς*, water; *κεφαλή*, head; *ειδός*, form]. Pertaining to or resembling hydrocephalus. *H. Disease*. See *Hall's Disease, Disenses, Table of*.

Hydrocephalus (*hi-dro-sef'-al-us*) [*ὑδρῶς*, water; *κεφαλή*, head]. A collection of fluid

in the cerebral ventricles (internal H.) or outside the brain-substance (external H.). The symptoms are: Progressive enlargement of the head, bulging of the fontanels, prominent forehead, thinness of hair and scalp, distention of the superficial veins, mental impairment, muscular weakness, convulsions. Acute external H. is due to inflammation of the meninges, usually tuberculous; acute internal H. is caused by ependymitis. Chronic H. is either congenital or acquired.

Hydrochinone (*hi-dro-ki'-non*) [*idwup*, water; *quinone*]. See *Hydroquinone*.

Hydrochlorate (*hi-dro-klo'-rat*) [*idwup*, water; *χλωρος*, green]. Any salt of hydrochloric acid.

Hydrochloric Acid (*hi-dro-klo'-rik*). See *Acid*, *Hydrochloric*.

Hydrochlorid (*hi-dro-klo'-rid*) [*idwup*, water; *χλωρος*, green]. Same as *Hydrochlorate*.

Hydrocirsocele (*hi-dro-sir'-so-sel*) [*idwup*, water; *κίρσος*, venous enlargement; *κύλη*, tumor]. Hydrocele accompanied with varicose veins of the spermatic cord.

Hydrocollidin (*hi-dro-kol'-id-in*) [*idwup*, water; *κόλλα*, glue], $C_8H_{13}N$. A highly poisonous ptomain obtained from putrefying mackerel, horse-flesh, and ox-flesh, and said to be identical with one obtained from nicotine.

Hydrocyanic (*hi-dro-se-an'-ik*). See *Cyanogen*, and *Acid*, *Hydrocyanic*.

Hydroelectric (*hi-dro-e-lek'-trik*) [*idwup*, water; *ηλεκτρον*, electricity]. Pertaining to electricity developed in connection with water. **H. Bath**, a bath in which the metallic lining of the tub is connected with one pole of a battery, the other pole being in contact with the person of the patient.

Hydrofluoric Acid (*hi-dro-flu-or'-ik*). See *Acid* and *Fluorin*.

Hydrogen (*hi'-dro-jen*) [*idwup*, water; *γεννᾶν*, to produce]. Symbol H; atomic weight 1; quantivalence 1. A combustible, gaseous element, feebly basic, and occurring in nature in greatest abundance combined with oxygen in the form of water, H_2O . It is present in nearly all organic compounds, and is a constant constituent of acids. **H. Monoxid**, H_2O , water. **H. Dioxid**, or **H. Peroxid**, H_2O_2 , an unstable liquid which readily yields up an atom of oxygen, and hence is a powerful oxidizer. It is strongly antiseptic. **Aqua hydrogenii dioxid** (U. S. P.), contains 3 per cent. of pure H_2O_2 . It is useful as an antiseptic application to inflamed mucous membranes in diphtheria, scarlet fever, gonorrhea, etc., and as a cleansing agent of suppurating cavities.

Hydrolein (*hi-dro'-e-in*) [*idwup*, water; *oleum*, oil]. A proprietary remedy, an emulsion of cod-liver oil with pancreatin and borax.

Hydrology (*hi-drol'-o-je*) [*idwup*, water; *λόγος*, science]. A treatise on the nature and uses of water.

Hydrolytic (*hi-dro-lit'-ik*) [*idwup*, water; *λύνειν*, to dissolve]. Pertaining to the decomposition of water, or the liberation of water during a chemic reaction. **H. Ferments**, those causing a combination with the elements of water in the substances they decompose.

Hydromel (*hi'-dro-mel*) [*idwup*, water; *μέλι*, honey]. A mixture of honey and water with or without a medicinal substance.

Hydromeningitis (*hi-dro-men-in-jit'-tis*) [*idwup*, water; *μηνιγξ*, a membrane; *τις*, inflammation]. 1. Inflammation of the membranes of the brain or cord, accompanied by effusion of serous fluid. 2. Inflammation of the membrane of Descemet.

Hydromeningocele (*hi-dro-men-in'-go-sel*) [*idwup*, water; *μηνιγξ*, membrane; *κύλη*, a tumor]. 1. A cystic tumor of the meninges, protruding through the skull. 2. A form of spina bifida, in which the sac contains cerebrospinal fluid.

Hydrometer (*hi-drom'-et-er*) [*idwup*, water; *μέτρον*, a measure]. An instrument for determining the specific gravity of liquids.

Hydrometra (*hi-dro-me'-truh*) [*idwup*, water; *μήτρα*, the womb]. A collection of watery fluid in the uterus.

Hydromphalus (*hi-drom'-fal-us*) [*idwup*, water; *δμφαλός*, navel]. A tumor at the navel distended with water.

Hydromyelia (*hi-dro-mi-e'-le-ah*) or **Hydromyelus** (*hi-dro-mi'-el-us*) [*idwup*, water; *μυελός*, marrow]. Dilatation of the central canal of the spinal cord with an accumulation of fluid.

Hydronaphtol (*hi-dro-naf'-tol*) [*idwup*, water; *νάφθα*, asphalt]. A substance probably identical with beta-naphtol, and like it used as an intestinal antiseptic. Dose gr. iiij-iv (0.20-0.26).

Hydronephrosis (*hi-dro-nef-ro'-sis*) [*idwup*, water; *νεφρός*, kidney]. A collection of urine in the pelvis of the kidney from obstructed outflow. The pressure of the fluid causes in time atrophy of the kidney-structure, and the whole organ is converted into a large cyst.

Hydronephrotic (*hi-dro-nef-rot'-ik*) [*idwup*, water; *νεφρός*, kidney]. Relating to, affected with, or of the nature of, hydronephrosis.

Hydropathy (*hi-drop'-ath-e*) [*idwup*, water; *πάθος*, disease]. The treatment of diseases by the use of water, externally and internally.

Hydropericardium (*hi-dro-per-ik-ar'-de-um*) [*idwup*, water; *περί*, around; *καρδια*, heart]. A collection of serum within the pericardial cavity.

Hydroperitoneum (*hi-dro-per-it-on-e'-um*) [*idwop*, water; *peritoneum*]. Ascites.

Hydrophobia (*hi-dro-fo'-be-ah*) [*idwop*, water; *phobos*, dread]. An acute infectious disease communicated to man by the bites of animals suffering from rabies. The period of incubation is variable, the average being from three to six months. The disease begins with malaise, slight spasmodic movements, and enlargement of the lymphatic glands in the neighborhood of the original wound. Soon the true hydrophobic state sets in—it is characterized by tonic spasm of the muscles of deglutition, excited by an effort at swallowing water or even at the sight of water. The pulse is rapid, there is a slight fever, and usually profuse salivation. Toward the end paralysis may develop. The disease is nearly always fatal. Pasteur has proposed a method of prophylaxis, consisting in the injection of material from the spinal cord of animals dead of rabies.

Hydrophobic (*hi-dro-fo'-bik*) [*idwop*, water; *phobos*, dread]. Pertaining to, or of the nature of, hydrophobia. **H. Tetanus.** See *Tetanus*.

Hydrophobophobia (*hy-dro-fo-bo-fo'-be-ah*) [*idwop*, water; *phobos*, fear]. An intense dread of hydrophobia; a condition producing a state simulating true hydrophobia.

Hydrophthalmia (*hi-dro-ftal'-me-ah*) [*idwop*, water; *ophthalmos*, eye]. An increase in the fluid contents of the eye causing the organ to become distended, resulting in glaucoma, keratoglobus, staphyloma, etc.

Hydropic (*hi-droft'-ik*) [*idwop*, dropsy]. Pertaining to dropsy, dropsical.

Hydropneumatoxis (*hi-dro-nu-mat'-o'-sis*) [*idwop*, water; *pneumatosis*, inflation]. A collection of fluid and air or other gas within the tissues.

Hydropneumopericardium (*hi-dro-nu-mo-per-ik-ar'-de-um*) [*idwop*, water; *pneuma*, air; *perikardion*, pericardium]. A collection of serum and air or other gas within the pericardium.

Hydropneumothorax (*hi-dro-nu-mo-tho'-raks*) [*idwop*, water; *pneuma*, air; *thorax*, thorax]. The presence of serous fluid and air or gas in the pleural cavity.

Hydrops (*hi'-drops*) [*idwop*, dropsy]. Dropsy. **H. articuli**, a watery effusion into the synovial cavity of a joint.

Hydroquinin (*hi-dro-kwin'-in*) [*idwop*, water; *quina*, Peruvian bark], $C_{20}H_{14}N_2O_2$. An alkaloid obtained from cinchona, and frequently contaminating quinin.

Hydroquinone (*hi-dro-kwin'-on*), $C_6H_6O_2$. An isomer of resorcin and pyrocatechin, found in arbutin (see *Uva ursi*), and also obtained from quinin and quinone, etc. It is antipyretic and antiseptic. Dose gr. xv-xx (1.0-1.3). Unof.

Hydorrhea (*hi-dror-e'-ah*) [*idwop*, water; *rhoia*, flow]. A flow of watery liquid. **H., gravidarum**, a discharge from the pregnant uterus of thin mucus that accumulates as a result of excessive secretion of the uterine glands.

Hydrosalpinx (*hi-dro-sal'-pink*) [*idwop*, water; *salpinx*, trumpet]. A distention of the Fallopian tube with fluid.

Hydrosarcocele (*hi-dro-sar'-ko-sel*) [*idwop*, water; *sarx*, flesh; *kēlē*, hernia]. Sarcocele with hydrocele.

Hydrostat (*hi-dro'-stat*) [*idwop*, water; *istānai*, to stand]. An apparatus for preventing the spilling of the fluid of electric batteries during transportation.

Hydrostatic or Hydrostatics (*hi-dro-stat'-ik* or *iks*) [*idwop*, water; *istānai*, to stand]. The science treating of the properties of liquids in a state of equilibrium.

Hydrosulphuric Acid (*hi-dro-sul'-fu'-rik*). See *Acid*.

Hydrotherapeutics (*hi-dro-ther-ap-u'-tik*) [*idwop*, water; *therapeuein*, to heal]. The treatment of disease by means of water, or the use of water in the treatment of disease.

Hydrotherapy (*hi-dro-ther'-ap-e*). See *Hydrotherapeutics*.

Hydrothionammonemia (*hi-dro-thi-on-am-o-ne'-me-ah*) [*idwop*, water; *theion*, sulphur; *ammonia*, ammonia; *aima*, blood]. The condition produced by the presence of ammonium sulphid in the blood.

Hydrothionemia (*hi-dro-thi-on-e'-me-ah*) [*idwop*, water; *theion*, sulphur; *aima*, blood]. The condition produced by the presence of hydrogen sulphid in the blood.

Hydrothionuria (*hi-dro-thi-on-u'-re-ah*) [*idwop*, water; *theion*, sulphur; *urion*, urine]. The presence of hydrogen sulphid in the urine.

Hydrothorax (*hi-dro-tho'-raks*) [*idwop*, water; *thorax*, chest]. The presence of serous fluid in the pleural cavity, due to a passive effusion, as in cardiac, renal, and other diseases.

Hydrotomy (*hi-drot'-o-mē*) [*idwop*, water; *trōmein*, to cut]. A method of dissecting tissues by the forcible injection of water into the arteries and capillaries, whereby the structures are separated.

Hydroxid (*hi-droks'-id*) [*idwop*, water; *deixis*, sharp]. A metallic or basic radicle combined with one or more hydroxyl groups. Hydroxids may be regarded as formed from water (HOH) by the substitution for one of its hydrogen atoms of a metal or basic radicle. Synonym of *Hydrate*.

Hydroxyl (*hi-droks'-il*) [*idwop*, water; *deixis*, sharp]. The univalent radicle, OH, the combination of which with basic elements or radicles forms the hydroxids.

Hydroxylamin (*hi-droks-il'-am-in*) [*idwop*,

water; *δξύς*, sharp; *amin*]. NH_4OH . A basic substance, known only in solution in water or in combination with acids. Its hydrochlorid has been used as a substitute for chrysarobin in skin-diseases.

Hydruria (*hi-dru'-re-ah*) [*ὕδωρ*, water; *οὐρον*, urine]. The discharge of a large quantity of urine of low specific gravity.

Hygiene (*hi-je-èn*) [*ὑγιεινός*, good for the health]. The science that treats of the laws of health and the methods of their observance.

Hygienic (*hi-je-en'-ik*) [*ὑγιεινός*, good for the health]. Pertaining to hygiene, as H. treatment, that which simply guards against infraction of the laws of health.

Hygrin (*hi'-grin*) [*ὕγρος*, moist; from its liquid form], $\text{C}_{12}\text{H}_{13}\text{N}$. A liquid alkaloid, derived from coca.

Hygroma (*hi-gro'-mah*) [*ὕγρος*, moist; *δμα*, tumor]. A bursa, or newly-formed sac, distended with fluid.

Hygrometer (*hi-grom'-et-er*) [*ὕγρος*, moist; *μέτρον*, measure]. An instrument for determining quantitatively the amount of moisture in the air.

Hygrometric (*hi-gro-met'-rik*) [*ὕγρος*, moist; *μέτρον*, measure]. 1. Pertaining to hygrometry. 2. Readily absorbing water; hygroscopic.

Hygrometry (*hi-grom'-et-re*) [*ὕγρος*, moist; *μέτρον*, measure]. The measurement of the moisture of the air.

Hygroscopic (*hi-gro-skop'-ik*) [*ὕγρος*, moist; *σκοπεῖν*, to see]. Having the property of absorbing moisture from the air.

Hymen (*hi'-men*) [*ὑμῆν*, membrane]. The fold of mucous membrane that partially occludes the vaginal orifice.

Hymenal (*hi'-me-nal*) [*ὑμῆν*, membrane]. Pertaining to the hymen.

Hymenitis (*hi-men-i'-tis*) [*ὑμῆν*, membrane; *τις*, inflammation]. Inflammation of the hymen, or of any membranous structure.

Hymenology (*hi-men-ol'-o-je*) [*ὑμῆν*, membrane; *λόγος*, science]. The science of the nature, structure, functions, and diseases of membranes.

Hyoepiglottic (*hi-o-ep-e-glot'-ik*) [*ὕοειδής* hyoid; *ἐπιγλωττις*, epiglottis]. Relating to the hyoid bone and the epiglottis.

Hyoglossal (*hi-o-glos'-al*) [*ὕοειδής*, hyoid; *γλῶσσα*, tongue]. Extending from the hyoid bone to the tongue.

Hyoglossus (*hi-o-glos'-us*). See *Muscles*, Table of.

Hyoid (*hi'-oid*) [*Υ*, the Greek letter upsilon; *εἶδος*, resemblance]. Having the form of the Greek letter upsilon. **H. Bone**, a bone situated between the root of the tongue and the larynx, supporting the tongue and giving attachment to its muscles.

Hyoscin (*hi'-o-sin*) [*ὕς*, hog; *κνᾶμος*, bean], $\text{C}_{17}\text{H}_{23}\text{NO}_5$. A liquid alkaloid found in hyoscyamus. It is a powerful depressant of the cerebrum and the motor centers of the cord, and is employed in insomnia, mania, and excessive sexual excitement. The hydrobromate is most commonly administered. Dose gr. $\frac{1}{15}$ – $\frac{1}{8}$ (0.0005–0.00075).

Hyoscyamin (*hi-o-si'-am-in*) [*ὕς*, hog; *κνᾶμος*, bean], $\text{C}_{17}\text{H}_{23}\text{NO}_5$. An alkaloid occurring in hyoscyamus. It is isomeric with atropin, is a mydriatic, narcotic, and sedative. Dose gr. $\frac{1}{15}$ – $\frac{1}{8}$ (0.0005).

Hyoscyamus (*hi-os-si'-am-us*) [*ὕς*, hog; *κνᾶμος*, bean]. Henbane, a plant of the order Solanaceæ. The leaves and flowering tops of *H. niger* yield the alkaloids hyoscyamin and hyoscin. It is sedative to the nervous system, and has been employed in hysteria, cough, and colic, and to relieve pain in rheumatism, headache, and malignant tumors. **H.**, **Ext.** Dose gr. j–ij (0.065–0.13). **H.**, **Ext.**, **Fld.** Dose $\frac{m}{v}$ –x (0.32–0.65). **H.**, **Succus** (**B. P.**). Dose f3 ss–j (2.0–4.0). **H.**, **Tinct.** Dose f3 ss–j (2.0–4.0).

Hypacusis, **Hypacusia**, **Hypacusis**, or **Hypakusis** (*hip-ak-u'-sis*, *hip-ak-u'-ze-ah*) [*ὕπῶς*, under; *ἀκουσις*, hearing]. Impairment of hearing.

Hypalbuminosis (*hip-al-bu-min-o'-sis*) [*ὕπῶς*, under; *albumen*]. Diminution in the proportion of albumin in the blood.

Hyper- (*hi'-per-*) [*ὑπέρ*, above]. A Greek prefix signifying above, beyond, or excessive.

Hyperacidity (*hi-per-as-id'-it-e*) [*ὑπέρ*, over; *acidum*, acid]. Excess of acidity.

Hyperacusis, **Hyperacusia**, **Hyperacusis**, or **Hyperakusis** (*hi-per-ak-u'-sis*, *hi-per-ak-u'-ze-ah*) [*ὑπέρ*, over; *ἀκουσις*, hearing]. Morbid acuteness of the sense of hearing; auditory hyperesthesia.

Hyperactivity (*hi-per-ak-tiv'-it-e*) [*ὑπέρ*, over; *agere*, to do]. Excessive or abnormal activity.

Hyperæmia (*hi-per-e'-me-ah*). See *Hyperæmia*.

Hyperæsthesia (*hi-per-es-the'-ze-ah*). See *Hyperesthesia*.

Hyperakusis (*hi-per-ak-u'-sis*). See *Hyperacusis*.

Hyperalbuminosis (*hi-per-al-bu-min-o'-sis*) [*ὑπέρ*, over; *albumen*, albumin]. An increase in the amount of albumin in the blood.

Hyperalgia (*hi-per-al-je'-ze-ah*) [*ὑπέρ*, over; *ἀλγισις*, pain]. Excessive sensibility to pain.

Hyperbulia (*hi-per-bu'-le-ah*) [*ὑπέρ*, over; *βουλῆ*, will]. Exaggerated wilfulness; abnormal development of will-power.

Hypercatharsis (*hi-per-kath-ar'-sis*) [*ὑπέρ*, over; *κάθαρσις*, cleansing]. Excessive purging.

Hypercholia (*hi-per-ko'-le-ah*) [*ὑπέρ*, over; *χολή*, bile]. An excessive secretion of bile.

Hypercyesis (*hi-per-si-e'-sis*) [*ὑπέρ*, over; *κίησις*, conception]. Superfetation.

Hyperdicrotic (*hi-per-di-krot'-ik*) [*ὑπέρ*, over; *δικροτός*, a double beat]. Affected with marked or delayed diastole; a condition in which the aortic notch is below the base-line.

Hyperdistention (*hi-per-dis-ten'-shun*) [*ὑπέρ*, over; *distendere*, to stretch]. Forcible or extreme distention.

Hyperdynamia (*hi-per-di-nam'-e-ah*) [*ὑπέρ*, over; *δύναμις*, energy]. Excessive strength or exaggeration of nervous or muscular function.

Hyperemesis (*hi-per-em'-es-is*) [*ὑπέρ*, over; *ἔμεσις*, vomiting]. Excessive vomiting.

Hyperemia (*hi-per-e'-me-ah*) [*ὑπέρ*, over; *αἷμα*, blood]. Excessive blood in a part. **H.**, Active, that caused by an excessive supply of blood going to a part. **H.**, Passive, that caused by an impediment to the removal of the blood.

Hyperencephalus (*hi-per-en-sef'-a-lus*) [*ὑπέρ*, over; *ἐγκέφαλος*, brain]. A variety of single autostic monsters in which the upper portion of the skull is entirely lacking.

Hypererythrocythemia (*hi-per-er-ith-ro-si-the'-me-ah*) [*ὑπέρ*, over; *ἐρυθρός*, red; *κύτος*, cell; *αἷμα*, blood]. Excess of red corpuscles in the blood.

Hyperesthesia (*hi-per-es-the'-se-ah*) [*ὑπέρ*, over; *αἰσθησις*, sensational]. Excessive sensibility of the skin.

Hypergeusia (*hi-per-gu'-se-ah*) [*ὑπέρ*, over; *γεῖσις*, taste]. Abnormal acuteness of the sense of taste.

Hyperglobulia (*hi-per-glo-bu'-le-ah*) [*ὑπέρ*, over; *globus*, a ball]. An increase in the number of the red blood-corpuscles.

Hyperhidrosis, or **Hyperidrosis** (*hi-per-hid-ro'-sis*, or *hi-per-id-ro'-sis*) [*ὑπέρ*, over; *ἰδρῶς*, sweat]. Excessive sweating.

Hyperinosis (*hi-per-in-o'-sis*) [*ὑπέρ*, over; *ῖς*, fiber]. An excessive increase in the fibrin-factors in the blood.

Hyperinvolution (*hi-per-in-vo-lu'-shun*) [*ὑπέρ*, over; *involvere*, to roll around]. Excessive involution of an organ after enlargement, as of the uterus after pregnancy, resulting in a reduction below the normal size.

Hyperkeratosis (*hi-per-ker-at-o'-sis*) [*ὑπέρ*, over; *κέρας*, horn, cornea]. 1. Hypertrophy of the cornea. 2. Hypertrophy of the horny layer of the skin.

Hyperkinesia (*hi-per-kin-e'-se-ah*) [*ὑπέρ*, over; *κίνησις*, energy]. Excessive movement, as that associated with muscular spasm.

Hyperleukocythemia (*hi-per-lu-ko-si-the'-me-ah*). See *Hyperleukocytosis*.

Hyperleukocytosis (*hi-per-lu-ko-si-to'-sis*)

[*ὑπέρ*, over; *λευκός*, white; *κύτος*, cell]. An increase in the number of leukocytes in the blood.

Hypermastia (*hi-per-mas'-te-ah*) [*ὑπέρ*, over; *μαστός*, breast]. Excessive development of the mammary gland.

Hypermetropia (*hi-per-me-tro'-pe-ah*). Same as *Hyperopia*.

Hypermetropic (*hi-per-me-trop'-ik*) [*ὑπέρ*, over; *μέτρον*, measure; *ὤψ*, eye]. Affected with, or pertaining to, hypermetropia.

Hyperonychchia (*hi-per-o-nik'-e-ah*) [*ὑπέρ*, over; *ὄνυξ*, nail]. Hypertrophy of the nails.

Hyperope (*hi'-per-op*) [*ὑπέρ*, over; *ὤψ*, sight]. One who is affected with hyperopia.

Hyperopia (*hi-per-o'-pe-ah*) [*ὑπέρ*, over; *ὤψ*, sight]. That condition of the refractive media of the eye in which, with suspended accommodation, the focus of parallel rays of light is behind the retina. It is due to an abnormally short anteroposterior diameter of the eye, or to a subnormal refractive power of its media. **H.**, Absolute, that which cannot be corrected completely by accommodation, so that there is indistinct vision even for distance. **H.**, Axial, that due to abnormal shortness of the anteroposterior diameter of the eye, the refractive power being normal.

H., Facultative, that which may be corrected by the accommodation, so that there is distinct vision at a distance. **H.**, Latent, that part of the total H. that cannot be overcome by the accommodation, or the difference between the manifest and the total hyperopia.

H., Manifest, that which the accommodation can overcome, or that corrigible by a convex glass aided by the accommodation.

H., Relative, a high hyperopia in which distinct vision is possible only when excessive convergence is made. **H.**, Total, the entire hyperopia, both latent and manifest.

Hyperosmia (*hi-per-oe'-me-ah*) [*ὑπέρ*, over; *ὁσμή*, smell]. An abnormally acute sense of smell.

Hyperostosis (*hi-per-os-to'-sis*) [*ὑπέρ*, over; *ὀστέον*, bone]. A bony outgrowth.

Hyperplasia (*hi-per-pla'-ze-ah*) [*ὑπέρ*, over; *πλασις*, molding]. Excessive formation of tissue; an increase in the size of a tissue or organ owing to an increase in the number of cells; also termed numerical hypertrophy.

Hyperplastic (*hi-per-plas'-tik*) [*ὑπέρ*, over; *πλαστικός*, fit for molding]. Pertaining to hyperplasia.

Hyperpnea (*hi-per-pne'-ah*) [*ὑπέρ*, over; *πνέω*, breath]. Panting or exaggerated respiration.

Hyperpraxia (*hi-per-praks'-e-ah*) [*ὑπέρ*, over; *πράξις*, exercise]. The restlessness of movement characterizing certain forms of mania.

Hyperpselaphesia (*hi-per-sel-af-e'-ze-ah*)

[*ὑπέρ*, over; *ψηλάφησις*, touch]. Abnormal increase of tactile sensibility.

Hyperpyrexia (*hi-per-pi-rek's-e-ah*) [*ὑπέρ*, over; *πῦρ*, fire; *ἔχειν*, to have]. Excessively high body-temperature. By some the term is used only when the temperature is above 106° F.

Hyperresonance (*hi-per-res'-o-nans*) [*ὑπέρ*, over; *resonance*]. Increased resonance on percussion.

Hypersecretion (*hi-per-se-kre'-shun*) [*ὑπέρ*, over; *secerere*, to separate]. Excessive secretion.

Hyperthermia (*hi-per-ther'-me-ah*). Elevation of temperature above the normal.

Hypertrichiasis, Hypertrichosis (*hi-per-trik-i'-as-is, hi-per-trik-o'-sis*). Excessive growth of hair of a part or the whole of the body.

Hypertrophy (*hi-per'-tro-fe*) [*ὑπέρ*, over; *τροφή*, nourishment]. An increase in the size of a tissue or organ independent of the general growth of the body. **H., Compensatory**, that resulting from the increased activity of an organ to make up some deficiency in a paired organ, or in itself. **H., Concentric**, of the heart, increase in the thickness of the walls, without increase in the size of the organ, but with diminution in the capacity of its chambers. **H., Eccentric**, of the heart, hypertrophy with dilatation. **H., False**, an increase in some one constituent tissue of an organ, usually the connective tissue. **H., Numeric**, H. due to an increase in the number of cells. **H., Simple**. 1. That in which there is increase in the size of the individual cells. 2. Of the heart, increased thickness of the walls, the size of the cavities remaining unchanged. **H., True**, an increase of all the component tissues of an organ giving increased power.

Hypesthesia (*hip-es-the'-ze-ah*) [*ὑπό*, under; *αἰσθησις*, sensation]. Impairment of sensation; lessened tactile sensibility.

Hyphemia (*hi-fe'-me-ah*) [*ὑπό*, under; *αἷμα*, blood]. 1. Oligemia; deficiency of blood. 2. Hemorrhage in the anterior chamber of the eye.

Hyphomycetes (*hi-fo-mi-se'-tez*) [*ὑφώ*, web; *μύκης*, fungus]. The molds.

Hypinosis (*hip-in-o'-sis*) [*ὑπό*, under; *ἵς*, ἵνός, fiber]. A deficiency of fibrin-factors in the blood.

Hypnal (*hip'-nal*) [*ὑπνος*, sleep]. A drug composed of antipyrin and chloral hydrate, and used as an hypnotic. Dose gr. xv (1.0). Unof.

Hypnic (*hip'-nik*) [*ὑπνικός*, producing sleep]. 1. Pertaining to or inducing sleep. 2. An agent that induces sleep.

Hypnogenetic, Hypnogenic, Hypnogenous (*hip-no-jen-et'-ik, hip-no-jen'-ik, hip-*

noj'-en-us) [*ὑπνος*, sleep; *γεννάν*, to produce]. 1. Producing or inducing sleep. 2. Inducing hypnotism.

Hypnolepsy (*hip'-no-lep-se*) [*ὑπνος*, sleep; *λήψις*, seizure]. Excessive sleepiness.

Hypnology (*hip-nol'-o-je*) [*ὑπνος*, sleep; *λόγος*, science]. The science dealing with sleep or with hypnotism.

Hypnone (*hip'-non*). See *Acetophenone*.

Hypnosis (*hip-no'-sis*) [*ὑπνος*, sleep]. 1. The condition produced by hypnotizing. 2. The production of sleep; also, the gradual approach of sleep.

Hypnotic (*hip-not'-ik*) [*ὑπνος*, sleep]. 1. Inducing sleep. 2. Pertaining to hypnotism. 3. A remedy that causes sleep.

Hypnotism (*hip'-not-izm*) [*ὑπνος*, sleep]. A state of artificial somnambulism or trance, induced in certain persons by concentrating the gaze on a small object, or on a revolving mirror, or by complete subjection of their will to that of another, at whose command the hypnotic state develops. Three stages are described,—the cataleptic, the lethargic, and the somnambulist.

Hypnotization (*hip-not-iz-a'-shun*) [*ὑπνος*, sleep; *facere*, to make]. The induction of hypnotism.

Hypo- (*hi'-po-*) [*ὑπό*, under]. A prefix denoting: 1. Deficiency or lack. 2. Below or beneath, opposed to *epi-*, upon. 3. Of acids and salts, denoting those having a less number of atoms of oxygen than other compounds of the same elements.

Hypoblast (*hi'-po-blast*) [*ὑπό*, under; *βλαστός*, sprout]. The internal layer of the blastoderm, also called the endoderm, endoblast, or entoderm. From it is developed the intestinal epithelium (except that of the mouth and anus), and that of the glands opening into the intestines, and the epithelium of the air-passages.

Hypobromite (*hy-po-bro'-mit*) [*ὑπό*, under; *βρώμιος*, stench]. A salt of hypobromous acid. **H. Method**, a method of estimating the quantity of urea in urine, based upon the fact that when urea is acted upon by sodium hypobromite it is decomposed into nitrogen, carbon dioxide, and water. From the volume of nitrogen evolved the quantity of urea can be determined.

Hypobulia (*hi-po-bu'-le-ah*) [*ὑπό*, under; *βούλη*, will]. Deficiency of will-power.

Hypochlorhydria (*hi-po-klor'-hi'-dre-ah*) [*ὑπό*, under; *χλωρός*, green; *ὕδωρ*, water]. A condition in which there is a diminished amount of hydrochloric acid in the gastric juice.

Hypochlorite (*hi-po-klo'-rit*) [*ὑπό*, under; *χλωρός*, green]. Any salt of hypochlorous acid, HClO. The most important are those of calcium and sodium.

Hypochondriac (*hi-po-kon'-dre-ak*) [*ὑπό, under; χόνδρος, cartilage*]. 1. Pertaining to the hypochondrium. 2. A person who is affected with hypochondriasis.

Hypochondriasis (*hi-po-kon-dri'-as-is*) [*ὑπο-χονδριασμός, affected in the hypochondria*]. A condition in which the patient believes himself suffering from grave bodily diseases.

Hypochondrium (*hi-po-kon'-dre-um*) [*ὑπό, under; χόνδρος, cartilage*]. The upper lateral region of the abdomen beneath the lower ribs.

Hypodermatic (*hi-po-der-mat'-ik*) [*ὑπό, under; δέρμα, skin*]. Placed or introduced beneath the skin, as H. injection.

Hypodermatoclysis or **Hypodermoclysis** (*hi-po-der-mat-ok'-lis-is, hi-po-der-mok'-lis-is*) [*ὑπό, under; δέρμα, skin; κλύσις, injection*]. The introduction into the subcutaneous tissues of large quantities of fluids, especially of normal saline solution.

Hypodermic (*hi-po-der'-mik*). See *Hypodermatic*.

Hypogastric (*hi-po-gas'-trik*) [*ὑπό, under; γαστήρ, the belly*]. Pertaining to the hypogastrium.

Hypogastrium (*hi-po-gas'-tre-um*) [*ὑπό, under; γαστήρ, stomach*]. The lower median anterior region of the abdomen.

Hypogeusia (*hi-po-gu'-se-ah*) [*ὑπό, under; γεύσις, taste*]. Diminution in the sense of taste.

Hypoglossal (*hi-po-glos'-al*) [*ὑπό, under; γλῶσσα, tongue*]. Situated under the tongue.

H. Nerve. See *Nerves, Table of*.

Hypoglossus (*hi-po-glos'-us*) [*ὑπό, under; γλῶσσα, tongue*]. The hypoglossal nerve.

Hypoglossitis (*hi-po-glos'-is*) [*ὑπό, under; γλῶσσα, tongue*]. 1. The under part of the tongue. 2. A swelling at the under part of the tongue, as a ranula.

Hypognathous (*hi-pog'-na-thus*) [*ὑπό, under; γνάθος, jaw*]. Having the lower mandible longer than the upper.

Hypognathus (*hi-pog'-na-thus*) [*ὑπό, under; γνάθος, jaw*]. A double monstrosity in which the parasite is attached to the inferior maxillary bone.

Hypohidrosis (*hi-po-hid-ro'-sis*) [*ὑπό, under; ἰδρώσις, sweating*]. Scanty perspiration.

Hypokinesia, or **Hypokinesis** (*hi-po-kin-e'-se-ah, or hi-po-kin-e'-sis*) [*ὑπό, under; κίνησις, motion*]. Deficiency in motor reaction under stimulation.

Hypoleukocytosis (*hi-po-lu-ko-si'-to'-sis*) [*ὑπό, under; λευκός, white; κύτος, cell*]. A diminution of the number of leukocytes in the blood.

Hypomania (*hi-po-ma'-ne-ah*) [*ὑπό, under; μανία, madness*]. A moderate degree of maniacal exaltation.

Hypomelancholia (*hi-po-me-lan-ko'-le-ah*)

[*ὑπό, under; μελαγχολία, melancholia*]. Moderate melancholia; melancholia without delusions.

Hypometropia (*hi-po-me-tro'-pe-ah*) [*ὑπό, under; μέτρον, measure; ὥψ, vision*]. Myopia.

Hypopepsia (*hi-po-pep'-se-ah*) [*ὑπό, under; πέψις, digestion*]. Subnormal digestive power.

Hypophosphite (*hi-po-fos'-fit*) [*ὑπό, under; phosphorus*]. A salt of hypophosphorous acid. Those of calcium, iron, potassium, and sodium are official. Syrupus hypophosphitum contains the hypophosphites of calcium, potassium, and sodium. Dose f3 j-ij (4.0-8.0). Syr. hypophosphitum cum ferro contains syrupus hypophosphitum, ferrous lactate, and potassium citrate. Dose f3 j-ij (4.0-8.0). Both are used in wasting diseases, in scrofula, rickets, etc.

Hypophosphorous Acid (*hi-po-fos'-fo'-rus*) H_3PO_2 . A monobasic acid, used chiefly in the form of its salts. See *Hypophosphite*. Acidum hypophosphorosum dilutum (U.S.P.) is tonic. Dose ℥x-xxx (0.65-2.0).

Hypophysis (*hi-poff'-is-is*) [*ὑπό, under; οἰσιν, to grow*]. An outgrowth. **H. cerebri**, the pituitary body.

Hypoplasia (*hi-po-pla'-ze-ah*) [*ὑπό, under; πλάσσειν, to mold*]. Defective development of any organ or tissue.

Hypopselaphesia (*hi-pop-sel'-af'-ze-ah*) [*ὑπό, under; ψηλάφησις, touch*]. Diminution of sensitiveness to tactile impressions.

Hypopyon (*hi-po'-pe-on*) [*ὑπό, under; πύον, pus*]. A collection of pus in the anterior chamber of the eye.

Hyposmia (*hi-pas'-me-ah*) [*ὑπό, under; ὀσμή, smell*]. Diminution of the sense of smell.

Hypospadia, or **Hypospadias** (*hi-po-spa'-de-ah, hi-po-spa'-de-as*) [*ὑπό, under; σπᾶν, to draw*]. A condition in which the urethra opens upon the under surface of the penis.

Hypostasis (*hi-pas'-tas-is*) [*ὑπό, under; stasis*]. 1. A settling; also, the sediment. 2. The settling of blood in the dependent parts of the body.

Hypostatic (*hi-po-stat'-ik*) [*ὑπό, under; stasis*]. Due to, or of the nature of, hypostasis. **H. Congestion**. See *Hypostasis*.

H. Pneumonia. See *Pneumonia*.

Hyposthenia (*hi-po-sthe-ne'-ah*) [*ὑπό, under; ἀθίως, strength*]. Weakness.

Hypothenar (*hi-poth'-en-ar*) [*ὑπό, under; θύρον, palm*]. The fleshy eminence on the palm of the hand over the metacarpal bone of the little finger. Also, the prominences on the palm at the base of the fingers.

Hypothermal (*hi-po-ther'-mal*) [*ὑπό, under; θερμ, heat*]. Slightly hot; tepid.

Hypothermia (*hi-po-ther'-me-ah*) [*ὑπό, under; θερμ, heat*]. Subnormal temperature.

Hypotonia, or **Hypotonus** (*hi-po-to'-ne-ah*, or *hi-po'-o-nus*) [*ὑπό*, under; *τόνος*, tension]. Decrease of normal tonicity or tension; especially diminution of intraocular pressure.

Hypoxanthin (*hi-po-san'-thin*) [*ὑπό*, under; *ξανθός*, yellow], $C_8H_4N_4O$. A nonpoisonous leukomatin, also known as sarcin or sarkin. It occurs, accompanying adenin and guanin, in nearly all the animal tissues and organs rich in nucleated cells. In minute quantities it is a normal constituent of urine. It has also been found in plants, seeds, ferments, and wines. It is a crystalline body, soluble in cold and boiling water, insoluble in cold alcohol or ether. Hypoxanthin appears to be one of the products formed by the decomposition and successive oxidation of proteid matter previous to the formation of uric acid and urea. It is produced from adenin by the action of nitrous acid.

Hyssop (*his'-op*) [*ῥύσσωπος*, an aromatic plant]. Hedge-hyssop. The leaves and tops of *Hyssopus officinalis*, an aromatic stimulant, carminative, and tonic, employed in chronic catarrh of the respiratory tract. Dose of the fld. ext. f3j-ij (4.0-8.0). Unof.

Hysteralgia (*his-ter-al'-je-ah*) [*ὑστέρα*, womb; *ἄλγος*, pain]. Neuralgic pain in the uterus.

Hysterectomy (*his-ter-ek'-to-me*) [*ὑστέρα*, womb; *ἐκτομή*, a cutting out]. Excision of the uterus through the abdomen (abdominal H.), or the vagina (vaginal H.).

Hysteria (*his-te'-re-ah*) [*ὑστέρα*, womb, from the ancient belief that the condition depended upon uterine disease]. A diseased state of the mind manifesting itself in countless disturbances of the psychic, sensory, motor, and vasomotor functions. The etiology is not definitely known; heredity and mental shock play an important part. All ages and both sexes are subject to the disease, but it is most common in young women. The psychic disturbances consist in increased irritability, tendency to exaggeration, a heightened imagination, hallucinations, and somnambulistic and hypnotic states. Among sensory symptoms are various neuralgias, as clavius, hemicrania, and coccygodynia; anesthetics; hyperesthesias; diminution of the visual field; diplopia; deafness; loss of the sense of taste, etc. The motor symptoms comprise paralyses and contractions of the limbs, tremor, convulsions, catalepsy, aphonia, etc. The chief vasomotor phenomena are cyanosis, cutaneous hemorrhages, and edema. In addition to

these symptoms many others are at times noted, as anorexia, vomiting, salivation, polyuria, anuria, etc.

Hysteria (*his-ter'-ik*) [*ὑστέρα*, womb]. Pertaining to hysteria.

Hysterics (*his-ter'-iks*) [*ὑστέρα*, womb]. A popular term for the hysteric attack.

Hysteritis (*his-ter-i'-tis*). See *Metritis*.

Hystero- (*his'-ter-o-*) [*ὑστέρα*, uterus]. A prefix signifying relation to the uterus or to hysteria.

Hysteroceleisis (*his-ter-o-kli'-sis*) [*ὑστέρα*, womb; *κλείσις*, closure]. The closure of the uterus by suturing the edges of the os.

Hysteroepilepsy (*his-ter-o-ep'-e-lep-se*) [*ὑστέρα*, womb; *ἐπιληψία*, a laying hold of]. A form of hysteria accompanied by convulsions resembling those of epilepsy.

Hysterogenic, Hysterogenous (*his-ter-o-jen'-ik*, *his-ter-og'-en-us*) [*ὑστέρα*, womb; *γεννᾶν*, to beget]. Causing or producing an hysteric attack, as H. zones, certain regions pressure upon which excites an hysteric paroxysm.

Hysteroid (*his'-ter-oid*) [*ὑστέρα*, womb; *εἶδος*, form]. 1. Resembling hysteria. 2. Pertaining to hysteroepilepsy.

Hysterometry (*his-ter-oni'-et-re*) [*ὑστέρα*, womb; *μέτρον*, measure]. The measurement of the size of the uterus.

Hysteroneurosis (*his-ter-o-nu-ro'-sis*) [*ὑστέρα*, womb; *νεῦρον*, nerve]. A reflex neurosis resulting from irritation of the uterus.

Hysteropathy (*his-ter-op'-ath-e*) [*ὑστέρα*, womb; *πάθος*, disease]. Any disease or disorder of the uterus.

Hysteropexy (*his'-ter-o-peks-e*) [*ὑστέρα*, womb; *πήξις*, a fastening]. Fixation of the uterus by a surgical operation to correct displacement.

Hysteroptosis (*his-ter-op-to'-sis*) [*ὑστέρα*, womb; *πτῶσις*, a falling]. Falling or inversion of the uterus.

Hysterorrhaphy (*his-ter-or'-ra-fe*) [*ὑστέρα*, womb; *ραφή*, suture]. 1. The closure of a uterine incision or rent by suture. 2. Hysteropexy.

Hysterotomy (*his-ter-ot'-o-me*) [*ὑστέρα*, womb; *τομή*, a cutting]. Incision of the uterus.

Hysterotrachelorrhaphy (*his-ter-o-tra-kel-or'-ra-fe*) [*ὑστέρα*, womb; *τράχηλος*, neck; *ραφή*, suture]. A plastic operation for the restoration of a lacerated cervix uteri.

Hystericiasis (*his-tris-i'-as-is*) [*ὑστρίξ*, a hedgehog]. A disease of the hair in which it stands out stiffly like the hair of the hedgehog.

I. The symbol of Iodin.

Iatraliptic (*i-at-rat-ip'-tik*) [*iarpós*, physician; *ἀλειπτής*, an anointer]. Curing by using ointments and frictions.

Iatrochemic (*i-at-ro-kem'-ik*) [*iarpós*, physician; *χημεία*, chemistry]. Treating disease on the principles of chemistry.

Iatrochemistry (*i-at-ro-kem'-ist-re*) [*iarpós*, physician; *χημεία*, chemistry]. 1. The application of chemistry to therapeutics; the treatment of disease by chemic means. 2. The theory that disease and its treatment are explicable on a chemic basis.

Iatrophysics (*i-at-ro-fis'-iks*) [*iarpós*, physician; *φυσικός*, pertaining to nature]. 1. The treatment of disease by physical measures. 2. The theory that disease and its treatment are explicable on a materialistic or physical basis. The materialistic explanation of disease; applied especially to an obsolete theory of the 17th century, that sought to explain physiologic and therapeutic facts by means of the principles of physics (dynamics and statics).

Iatrotechnics (*i-at-ro-tek'-niks*) [*iarpós*, physician; *τέχνη*, art]. The art of healing.

Ice (*is*) [*AS., is, ice*]. Water in its solid state, which it assumes at a temperature of 0° Centigrade, or 32° Fahrenheit. It is used in medicine in the forms of **Ice-bag**, **Ice-cap**, **Ice-compress**, to reduce temperature, to lessen inflammatory action, to check hemorrhage, and to relieve pain; internally as a refrigerant and to combat nausea.

Iceland (*is'-land*) [*AS., is, ice; land, land*]. An island of the Arctic Ocean. **I. Moss**. See *Cetraria*. **I. Spar**, a crystalline form of calcium carbonate, having doubly refracting properties, and used in instruments for studying polarized light.

Ichor (*i'-kor*) [*ichōr*, serum, or pus]. An acrid, thin, puriform discharge.

Ichorous (*i'-kor-us*) [*ichōr*, serum, or pus; Resembling or relating to ichor.

Ichorrhemia (*i-kor.-e'-me-ah*) [*ichōr*, ichor; *αἷμα*, blood]. The presence of septic matter in the blood.

Ichthyocolla (*ik-the-o-kol'-ah*) [*ichthys*, fish; *κόλλα*, glue]. Isinglass. The air-bladder of the sturgeon, *Acipenser huso*, occurring in horny, translucent, white sheets, that form a jelly with hot water. It is a form of gelatin, and is used as a food, for clarifying liquids, and as a test for tannic acid. **I., Emplastrum**, court-plaster.

Ichthyoid (*ik'-the-oid*) [*ichthys*, fish; *ειδος*, form]. Fish-like.

Ichthyol (*ik'-the-ol*) [*ichthys*, fish; *oleum*, oil].

The ammonium or sodium salt of a tarry substance obtained in the distillation of a bituminous mineral containing fossil fish. The chemic formula is $C_{20}H_{36}S_3O_6(NH_4)_2$, or $C_{20}H_{36}S_3O_6Na_2$. It contains about 15 per cent. of sulphur, and is used as an alterative and antiphlogistic, especially in eczema, acne, lupus, and other dermal diseases. Internally it has been employed in rheumatism, syphilis, leprosy, tuberculosis, etc. Dose gr. x-xxx (0.65-2.0) in 24 hours.

Ichthyophagous (*ik-the-off'-ag-us*) [*ichthys*, fish; *φαγεῖν*, to eat]. Fish-eating.

Ichthyosis (*ik-the-o'-sis*) [*ichthys*, fish]. A chronic skin-disease characterized by the development of epidermal plates somewhat resembling the scales of a fish. **I. hystrix**, a form characterized by warty growths, consisting of elongated and hypertrophied papillae, covered by greatly thickened epidermis. **I. simplex**, the common form of ichthyosis, in which the surface has a tessellated appearance, from being covered with large, finely corrugated, papery scales.

Ichthyotoxicon (*ik-the-o-toks'-ik-on*) [*ichthys*, fish; *τοξικόν*, poison]. A poisonous principle obtained from the flesh of certain fishes.

Icteric (*ik-ter'-ik*) [*ikteros*, jaundice]. Pertaining to or characterized by jaundice.

Icteritious (*ik-ter-ish'-us*) [*ikteros*, jaundice]. Affected with or resembling icterus.

Ictericoid (*ik'-ter-oid*) [*ikteros*, jaundice; *ειδος*, form]. Resembling the color of, or having the nature of, jaundice.

Icterus (*ik'-ter-us*) [*ikteros*, jaundice]. Jaundice. **I. gravis**, acute yellow atrophy of the liver, an acute disease characterized by jaundice, marked nervous symptoms, diminution in size of the liver, and a rapidly fatal termination. The urine contains bile and crystals of leucin and tyrosin. Microscopically, the liver shows intense fatty degeneration of the cells, together with areas of congestion. The disease is most common in women, is in many cases associated with pregnancy, and is probably infectious in origin. A similar condition of the liver may occur in phosphorus-poisoning. **I. neonatorum**, that which is sometimes observed in infants during the first few days after birth. The causes are obscure, particularly in the mild form; it may be due to the absorption of biliary pigment from the meconium and its entrance into the circulation through an open ductus venosus; a severe form is due to absence of the large bile-ducts, or to septic infection.

Ictus (*ik't-us*) [L., a stroke]. A sudden attack. **I. epilepticus**, an epileptic fit. **I. paralyticus**, a paralytic stroke. **I. solis**, sunstroke.

-id (*id'*). A suffix used in chemistry to denote a combination of two elementary substances.

Ideation (*i-de-a'-shun*) [*idéa*, form or semblance]. The formation of a mental conception; the cerebral action by which, or in accord with which, an idea is formed.

Identical (*i-den'tik-al*) [*idem*, the same]. Being the same, corresponding exactly. **I. Points**, corresponding points of the two retinae, upon which the rays from an object must be focused in order that it may be seen as one.

Ideomotor (*i-de-o-mo'tor*) [*idéiv*, to see; *movere*, to move]. Pertaining conjointly to ideation and movement. **I. Center**, that part of the cortex which, influenced by ideation, excites muscular movement. **I. Movements**, unconscious movements, due to impulses of the mind when the attention is otherwise absorbed.

Ideophrenia (*i-de-o-fre'-ne-ah*) [*idéiv*, to see; *phrén*, mind]. Insanity with marked perversion of ideas.

Idio- (*id'-e-o-*) [*idios*, one's own]. A prefix signifying pertaining to one's self; peculiar to the individual.

Idiocy (*id'-e-o-se*) [*idiótēs*, a private person]. A congenital condition of mental deficiency, usually accompanied by physical defects, and characterized by an almost total absence of intelligence.

Idiomuscular (*id'-e-o-mus'-ku-lar*) [*idios*, one's own; *musculus*, muscle]. Peculiar to muscular tissue; not involving any nerve-stimulus or any function of the organism, except those of the muscle itself. **I. Contraction**, the contraction of a fatigued or weakened muscle under certain conditions of extraneous stimulus.

Idioneurosis (*id'-e-o-nu-ro'-sis*) [*idios*, one's own; *neuron*, a nerve]. An affection due to some disturbed or abnormal condition of the nerves supplying the affected part; a simple and uncomplicated neurosis.

Idiopathic (*id'-e-o-path'-ik*) [*idios*, one's own; *pathos*, disease]. Not dependent upon another disease, or upon a known or recognized cause.

Idioplasm (*id'-e-o-plazm*) [*idios*, one's own; *plasma*, a thing formed]. A reproductive substance not contained in the body of the cell, but in the chromosomes of the nucleus, controlling and determining the actual characters of the particular cell, and also those of all of its descendants.

Idiosyncrasy (*id'-e-o-sin'-kra-se*) [*idios*, one's own; *σύν*, together; *κράσις*, a mingling].

1. Any special or peculiar characteristic or temperament by which a person differs from other persons. 2. A peculiarity of constitution that makes an individual react differently to drugs or other influences from most persons.

Idiot (*id'-e-ot*) [*idiótēs*, a private person]. A person congenitally almost destitute of intelligence.

Idrosis (*id-ro'-sis*). See *Hidrosis*.

Ignatia (*ig-na'-she-ah*) [from *Ignatius* Loyola, the founder of the Jesuits]. St. Ignatius' bean. The seed of *Strychnos ignatii*, containing the alkaloids strychnin and brucin. Its therapeutic effects are similar to those of nux vomica. **I., Abstractum**. Dose gr. ss-j (0.032-0.065). Unof. **I., Tinct.** Dose ℥ij-x (0.13-0.65). Unof.

Ignipuncture (*ig-ne-punk'-tūr*) [*ignis*, fire; *punctura*, puncture]. Puncture with platinum needles heated to whiteness by the electric current.

Ignis (*ig'-nis*) [L.]. Fire. **I. sacer**, erysipelas. **I. sancti Antonii**, Saint Anthony's fire, an old name for erysipelas; also for anthrax.

Ignition (*ig-nish'-un*) [*ignis*, fire]. The act of setting on fire.

Ileo- (*il'-e-o-*) [*εἰλεiv*, to roll]. A prefix signifying relation to the ileum.

Ileocecal (*il-e-o-se'-kal*) [*εἰλεiv*, to roll; *cecus*, blind]. Pertaining to both ileum and cecum. **I. Fossa**, a depression in the lower part of the small intestine at the base of the vermiform process. **I. Valve**, a valve consisting of two folds of mucosa that guards the passage between the ileum and cecum.

Ileocolic (*il-e-o-kol'-ik*) [*εἰλεiv*, to roll; *κόλον*, colon]. Pertaining conjointly to the ileum and the colon.

Ileocolitis (*il-e-o-ko-lit'-is*) [*εἰλεiv*, to roll; *κόλον*, colon; *ιτις*, inflammation]. Inflammation of the ileum and the colon.

Ileocolostomy (*il-e-o-ko-lost'-to-me*) [*εἰλεiv*, to roll; *κόλον*, colon; *στόμα*, mouth]. The establishment of an artificial communication between the ileum and the colon.

Ileoilcostomy (*il-e-o-il-e-ost'-to-me*) [*εἰλεiv*, to roll; *στόμα*, mouth]. The operation of establishing an artificial communication between two different parts of the ileum.

Ileum (*il'-e-um*) [*εἰλεiv*, to roll]. The lower portion of the small intestine, terminating in the cecum.

Ileus (*il'-e-us*) [*εἰλεiv*, to roll]. Volvulus.

Iliac (*il'-e-ak*) [*ilium*, the flank]. Pertaining to the ilium or to the flanks. **I. Artery**. See *Arteries*, Table of. **I. Crest**, the upper free margin of the ilium to which the abdominal muscles are attached. **I. Fascia**, the fascia lining the posterior part of the abdominal cavity and covering the psoas and iliacus

muscles. **I. Fossa.** See *Fossa*. **I. Muscle.** See *Muscles*, Table of. **I. Region,** one of the regions into which the abdomen is divided. See *Abdomen*.

Iliadelphus (il'-e-ad-el'-fus) [*ilia*, flanks; ἀδελφός, brother]. A monstrosity double from the pelvis upward.

Ilio- (il'-e-o-) [*ilium*, the flank]. A prefix denoting relation to the ilium.

Iliofemoral (il'-e-o-fem'-or-al) [*ilium*, the flank; *femur*]. Pertaining conjointly to the ilium and the femur. **I. Ligament.** See *Ligament*.

Iliohypogastric (il'-e-o-hi-po-gas'-trik) [*ilium*, the flank; ὑπό, beneath; γαστήρ, stomach]. Pertaining conjointly to the ilium and the hypogastrium.

Ilioinguinal (il'-e-o-in'-gue-nal) [*ilium*, flank; *inguen*, groin]. Pertaining to the ilium and the groin.

Iliopectineal (il'-e-o-pek-tin'-e-al) [*ilium*, flank; *pecten*, comb]. Pertaining conjointly to the ilium and the pubes. **I. Line.** See *Line*.

Iliopsoas (il'-e-o-so'-as) [*ilium*, flank; ψῶα, loin]. Pertaining conjointly to the ilium and the loins. **I. Muscle,** the psoas and iliacus muscles considered as a single muscle.

Iliotibial (il'-e-o-tib'-e-al) [*ilium*, flank; *tibia*, tibia]. Pertaining to or connecting the ilium and the tibia. **I. Band,** a thickened portion of the fascia lata extending from the outer tuberosity of the tibia to the iliac crest.

Ilium (il'-e-um) [L.]. 1. The flank. 2. The superior broad portion of the os innominatum, properly the os ilii.

Illaqueation (il'-ak-soe-a'-shun) [*illaqueare*, to ensnare]. A method of changing the direction of misplaced cilia, by withdrawing them by means of a loop through an opening in the tissue of the lid.

Illicium (il'-ik'-e-um) [*illicere*, to entice]. Star-anise. The fruit of *Illicium verum* (U. S. P.), or *Illicium anisatum* (B. P.), of the natural order Magnoliaceæ. It is the source of star-anise. **I. religiosum** and **I. parviflorum** are poisonous.

Illumination (il'-u-min-a'-shun) [*illuminare*, to make light]. 1. The act of illuminating or lighting up. 2. The quantity of light thrown on an object. **I., critical,** in microscopy, an illumination in which the lamp-flame is focused on the object. **I., Direct,** illumination of an object by light thrown upon it from in front. **I., Oblique,** illumination of an object by throwing light upon it obliquely, usually by means of a lens.

Illuminator, Abbé's. A system of lenses placed beneath the stage of the microscope for the purpose of condensing the light thrown upon the object; it is also known as Abbé's condenser.

Illusion (il'-lu'-shun) [*illusio*, a mocking]. A false interpretation by the mind of a real sensation.

Illusional (il'-lu'-shun-al) [*illusio*, a mocking]. Of the nature of an illusion.

Image (im'-āj) [*imago*, a likeness]. 1. A more or less accurate representation of an object. 2. The picture of an object formed by rays of light reflected, refracted, or passed through a small aperture. **I., After-** See *After-image*. **I., Direct,** or **I., Erect,** a picture obtained from rays that have not yet come to a focus. **I., Inverted,** one turned upside down. Nearly all real images are inverted. **I., Real,** that formed at the place where the rays meet. **I., Virtual,** an apparent image formed in the direction in which the rays enter the eye, the rays not actually converging at the point where the image is seen. The images formed by plane or convex mirrors and by concave lenses, when the object is placed within the principal focus, are virtual.

Imbecility (im'-bes-il'-i-e) [*imbecillitas*, imbecility]. Mental weakness, similar to but less great than that of idiocy.

Imbed (im'-bed'). In histology, to treat a tissue with some substance, as paraffin or celloidin, which shall give it support during the process of section-cutting.

Imbibition (im'-be-bish'-un) [*in*, in; *bibere*, to drink]. The act of sucking up moisture.

Imbricated (im'-brak-a-ted) [*imbrex*, a roof-tile]. Overlapping, like shingles on a roof.

Immature (im-at-ūr') [*in*, not; *maturus*, ripe]. Unripe; not yet of an adult age or growth.

Immediate (im-e'-de-at) [*in*, not; *medius*, middle]. Direct; without the intervention of anything. **I. Contagion,** that from a direct source. **I. Union,** union by first intention.

Immersion (im'-er'-shun) [*in*, in; *mergere*, to dip]. The plunging of a body into a liquid. **I.-bath,** a plunge-bath. **I.-lens,** a lens, usually of high power, the lower end of which is immersed in a drop of some liquid, such as water or oil, that has nearly the same refractive index as glass, and is placed on the cover-glass of the object under examination.

Immobilization (im'-ob-il-iz-a'-shun) [*im*, not; *movere*, to move]. The act of making firm, or of rendering motionless, as **I. of a joint**.

Immune (im-an') [*in*, not; *munis*, serving]. Safe from attack; protected against a disease by a natural or an acquired peculiarity.

Immunity (im-u'-nit-e) [*in*, not; *munis*, serving]. Exemption from disease; the condition of the body, wherein it resists the development of morbid processes. **I., Ac-** tive, that possessed by an individual after

recovering from certain infectious diseases. **I.**, Congenital, or Natural, that with which the individual is born. **I.**, Passive, that conferred by the introduction of antitoxins or vaccines.

Immunization (*im-u-nis-a'-shun*) [*in*, not; *munis*, serving]. The act of rendering immune.

Immunize (*im'u-niz*) [*in*, not; *munis*, serving]. To give immunity.

Immunizing Unit. See *Unit*.

Impact (*im'-pakt*) [*impingere*, to drive into or against]. A forcible striking against.

Impacted (*im-pak'-ted*) [*impingere*, to drive into or against]. Driven against and retained, as a wedge. **I.** Fracture. See *Fracture*.

Impaction (*im-pak'-shun*) [*impingere*, to drive into]. The state of being impacted or fixed in a part, as **I.** of the feces, or **I.** of a fragment of bone into another fragment.

Impalpable (*im-pal'-pa-bl*) [*in*, not; *palpare*, to feel]. Not capable of being felt; unappreciable by touch. **I.** Powder, a powder so fine that its separate particles cannot be felt.

Impaludism (*im-pal'-u-dizm*) [*in*, in; *palus*, a marsh]. Chronic malarial poisoning.

Impar (*im'-par*) [*in*, not; *par*, equal]. Odd or unequal, or without a fellow. **I.**, Ganglion, a small ganglion on the coccyx.

Imperative (*im-per'-at-iv*) [*imperare*, to command]. Peremptory, absolute, compulsory, binding. **I.** Conception, a conception or thought that dominates the actions of an individual, although the falsity of the conception may be recognized.

Imperforate (*im-per'-for-at*) [*in*, not; *perforare*, to pierce]. Without opening; not open or pervious, as **I.** anus.

Impermeable (*im-per'-me-a-bl*) [*in*, not; *per*, through; *meare*, to go]. Not permitting passage; not capable of being traversed.

Impervious (*im-per'-ve-us*) [*in*, not; *per-vius*, capable of passage]. Not permitting passage, especially passage of fluids.

Impetiginous (*im-pet-ij'-in-us*) [*impetigo*; *impetere*, to attack]. Affected with or resembling impetigo.

Impetigo (*im-pe-ti'-go*) [*impetigo*; *impetere*, to attack]. An acute inflammatory disease of the skin characterized by discrete, rounded pustules, unattended, as a rule, by itching or other subjective symptoms. **I.** contagiosa, an acute inflammatory contagious disease, characterized by the appearance of vesicles or blebs, that dry into flat, straw colored crusts. **I.** herpetiformis, a rare disease of the skin, characterized by the formation of superficial liliary pustules, that may be discrete, but tend to form circular groups. It is most common in pregnant women. **I.**, Syphilitic,

a syphilitic eruption having the characters of small flat pustules.

Implantation (*im-plan-ta'-shun*) [*in*, in; *plantare*, to set]. The act of setting in, as the transplantation of a tooth from the jaw of one person to that of another; the engrafting of epidermis from the skin of one person upon the body of another; the repair of a wounded intestine by uniting the divided ends. **I.**, Hypodermic, the introduction of a medicine under the skin. **I.**, Teratologic, a monstrosity consisting of an imperfect, joined to a perfect fetus.

Imponderable (*im-pon'-der-a-bl*) [*in*, not; *pondus*, weight]. Incapable of being weighed; without weight. **I.** Fluids, an obsolete term, formerly applied to light, heat, and electricity.

Impotence (*im'-pu-tens*) [*in*, not; *posse*, to be able]. Lack of power, especially lack of sexual power in the man.

Impregnate (*im-preg'-nat*) [*impregnare*]. 1. To render pregnant. 2. To saturate or charge with.

Impregnation (*im-preg-na'-shun*) [*impregnare*, to impregnate]. 1. The act of rendering pregnant; fecundation. 2. The process of saturating with or charging with.

Impressio (*im-presh'-e-o*) [*L.*]. An impression. **I.** colica, an impression on the under surface of the right lobe of the liver for the hepatic flexure of the colon. **I.** renalis, an impression on the under surface of the liver for the right kidney and suprarenal capsule.

Impression (*im-presh'-un*) [*imprimere*, to press upon]. 1. A hollow or depression. 2. The effect produced upon the mind, the body, or a disease, by external influence.

I., Maternal, the effects produced upon the fetus in utero by mental impressions received by the mother during pregnancy.

I.-preparation, Klatschpräparat. A cover-glass upon which an entire bacterial colony has been fixed by pressing the glass lightly upon the colony.

Impulse (*im'-puls*) [*impellere*, to drive against]. 1. A push or communicated force. 2. A sudden mental feeling that urges onward to an action. **I.**, Cardiac, the beat of the heart felt in the fifth intercostal space to the left of the sternum. **I.**, Morbid, a sudden, almost uncontrollable desire to do some unlawful or improper act.

In- [*L.*]. 1. A prefix signifying in or within. 2. A prefix signifying negation.

Inadequacy (*in-ad'-e-qua-se*) [*in*, not; *adequare*, to make equal]. Insufficiency. **I.**, Renal, that state of the kidney in which it is unable to remove from the blood a sufficient proportion of the effete matters that are normally excreted by it.

Inanimate (*in an'-im-at*) [*in*, not; *animus*, life]. Not animate; dead; without life.

Inanition (*in-an-ish'-un*) [*inanire*, to make empty]. Emptiness; want of food; wasting of the body from starvation.

Inappetence (*in-ap'-et-ens*) [*in*, not; *appetere*, to desire]. Loss of appetite.

Inarticulate (*in-ar-tik'-u-lar*) [*in*, not; *articulus*, a joint]. Not jointed or articulated; especially applied to vocal sounds not arranged into syllables.

In articulo mortis (*in ar-tik'-u-to mor'-tis*) [L.]. In the agony of death; at the point of dying.

Inassimilable (*in-as-im'-il-a-bl*) [*in*, not; *ad*, to; *simulare*, to make like]. Incapable of assimilation.

Incandescent (*in-kan-des'-ent*) [*incandescere*, to become white-hot]. Glowing; emitting luminous heat-rays; heated to the degree of emitting light. **I. Light**, one in which light is produced by the passage of an electric current through a strip of carbon or platinum suspended in a vacuum.

Incarcerated (*in-kar'-ser-a-ted*) [*incarcerare*, to imprison]. Imprisoned; held fast, as **I. hernia**.

Incidence (*in'-sid-ens*) [*incidere*, to fall upon]. A falling upon. **I., Angle of**, in optics, the angle at which a ray of light strikes a reflecting or refracting surface. **I., Point of**, the point upon which a ray or projectile strikes a reflecting or refracting surface.

Incident (*in'-sid-ent*) [*incidere*, to fall upon]. Falling upon.

Incineration (*in-sin-er-a'-shun*) [*in*, in; *cinere*, ashes]. The process of heating organic substances until all organic matter is driven off, and only the ash remains.

Incipient (*in-sip'-e-ent*) [*incipiens*, beginning]. Beginning to exist.

Incised (*in-sid'-*) [*in*, into; *cadere*, to cut]. Cut or notched. **I. Wound**, one made by a sharp-edged instrument.

Incision (*in-sizh'-un*) [*in*, into; *cadere*, to cut]. **1.** The act of cutting into any thing. **2.** A wound made with a cutting instrument.

Incisive (*in-sil'-iv*) [*in*, into; *cadere*, to cut]. **1.** Cutting. **2.** Pertaining to the incisor teeth. **I. Bone**, that part of the superior maxilla between the two clefts in double hare-lip; called also the intermaxillary bone.

Incisor (*in-sil'-zor*) [*in*, into; *cadere*, to cut]. **1.** Anything that cuts, especially an **I. tooth**. See **Tooth**. **2.** That which supplies the incisor teeth, as the **I. nerve**.

Incisura (*in-si-sul'-rah*) [*in*, into; *cadere*, to cut]. A notch; an incision. **I. cerebelli anterior**, **I. cerebelli posterior**, the notches

separating the hemispheres of the cerebellum, in front and behind.

Incisure (*in-sil'-zhur*) [*in*, into; *cadere*, to cut into]. A slit or notch. **I.'s of Schmidt and Lantermann**, oblique lines running across the white substance of the internodal segments of medullated nerve-fibers.

Inclusio fœtalis (*in-klu'-ze-o fe-tal'-tis*) [L.]. A form of fetal parasitism in which the parasite is more or less included and overgrown by the developed fetus.

Inclusion (*in-klu'-shun*) [*in*, in; *claudere*, to shut]. **1.** The state of being shut in. **2.** The act of shutting in. **3.** That which is shut in. **I., Fetal**, a monstrosity in which one fetus is included in and overgrown by the tissues of the other fetus.

Incoherent (*in-ko-he'-rent*) [*in*, not; *coherere*, to stick together]. Not connected; without proper sequence.

Incompatibility (*in-kom-pat-ib-il'-it-e*) [*in*, not; *cum*, together; *pati*, to suffer]. The state of being incompatible. It may be chemic or physiologic.

Incompatible (*in-kom-pat'-ib-l*) [*in*, not; *cum*, together; *pati*, to suffer]. Of two substances, not miscible without chemic change that destroys the usefulness of either or both; nor capable of being administered together on account of antagonistic properties.

Incompetence, Incompetency (*in-kom'-pet-ens, in-kom'-pe-ten-se*) [*in*, not; *cum*, together; *petere*, to seek]. Incapacity; inadequacy; inability to perform the natural functions. **I. of the Cardiac Valves**, an imperfect state of the valves of the heart in which they permit the return of blood into the cavity from which it came.

Incontinence (*in-kon'-tin-ens*) [*in*, not; *continere*, to contain]. **1.** Inability to control the escape of anything, as of the feces or the urine; involuntary evacuation. **2.** Venereal indulgence; lewdness.

Incoordination (*in-ko-or-din-a'-shun*) [*in*, not; *cum*, together; *ordinare*, to order]. Inability to produce voluntary muscular movements in proper order or sequence.

Incorporation (*in-kor-por-a'-shun*) [*in*, in; *corpus*, a body]. The process of intimately mixing the particles of different bodies into a practically homogeneous mass.

Increment (*in'-kre-ment*) [*in*, in; *crecere*, to grow]. Increase or growth.

Incrustation (*in-krus-ta'-shun*) [*in*, upon; *crusta*, crust]. The formation of a crust, especially a crust-like deposit of mineral salts.

Incubation (*in-ku-ba'-shun*) [*in*, on; *cumbere*, to lie]. **1.** The process of sitting upon eggs to favor hatching. **2.** The period of a disease between the implanting of the contagium and the development of the symptoms.

Incubator (*in'-ku-ba-tor*) [*in*, on; *cumbere*, to lie]. A device for the artificial hatching of eggs, or for the cultivation of bacteria; a contrivance for rearing prematurely-born children.

Incubus (*in'-ku-bus*) [*in*, upon; *cumbere*, to lie]. 1. Nightmare. 2. Anciently, a male demon supposed to have sexual connection with women in their sleep.

Incudal (*ing'-ku-dal*) [*incus*, anvil, *incus*]. Relating to the incus.

Incudomalleal (*ing'-ku-do-mal'-e-al*) [*incus*, *incus*; *malleus*, hammer]. Relating to the incus and the malleus.

Incudostapedial (*ing'-ku-do-sta-pe'-de-al*) [*incus*, *incus*; *stapes*, *stapes*]. Relating to the incus and the stapes.

Incurable (*in-kü'-ra-bl*) [*in*, not; *curare*, to care for]. Not curable.

Incurvation (*in-kur-va'-shun*) [*incurvare*, to bend]. The state of being bent or curved in.

Incus (*ing'-kus*) [L., an anvil]. The middle one of the chain of bones in the middle ear, so termed from its resemblance to an anvil.

Indentation (*in-den-ta'-shun*) [*in*, in; *dens*, a tooth]. A condition of being notched or serrated. 1. of Tongue, the notching of the borders of the tongue made by the teeth.

Index (*in'-deks*) [L.]. 1. The first finger. 2. The relation or ratio of one part to another taken as a standard. 1., **Alveolar**, the degree of prominence of the jaws, measured by the basialveolar length multiplied by 100 and divided by the basinasal length. When the alveolar index is less than 98, the skull is orthognathic, when more than 103, prognathic, when intermediate, mesognathic. 1., **Cephalic**, the breadth of a skull multiplied by 100 and divided by its length. When this is below 75, the skull is called dolichocephalic, when above 80, it is called brachycephalic, between these limits, mesaticephalic. 1., **Cerebral**, the ratio of the greatest transverse to the greatest anteroposterior diameter of the cranial cavity, multiplied by 100. 1., **Refractive**, the coefficient of refraction. 1., **Thoracic**, the ratio of the anteroposterior diameter to the transverse, expressed in percentage. 1., **Vertical**, the ratio of the vertical diameter of the skull to the maximum anteroposterior diameter, multiplied by 100.

Indian (*in'-de-an*) [*India*]. 1. Pertaining to India. 2. Pertaining to the aboriginal Americans. 1. **Hemp**. See *Cannabis*. 1. **Tobacco**. See *Lobelia*.

India Rubber. See *Caoutchouc*.

Indican (*in'-dik-an*) [*induxōn*, indigo], $C_{16}H_{11}NO_3$. 1. A glucosid occurring in indigo-plants, and by the decomposition of which

indigo is produced. 2. Potassium indoxylsulphate, $C_{16}H_9NSO_3K$, a substance occurring in the urine and sweat, and formed from indol.

Indication (*in'-dik-a'-shun*) [*indicare*, to point out]. That which points out; a guide, especially that which points out the course of treatment.

Indicator (*in'-dik-a-tor*) [*indicare*, to point out]. 1. The index-finger. 2. The extensor indicis muscle. 3. In chemistry, a substance used to show by a color-change when a change of reaction has taken place or a chemic affinity has been satisfied.

Indifferent (*in-dif'-er-ent*) [*in*, not; *differens*, different]. 1. Not differentiated; not tending to build up tissue, as I. cells. 2. Not readily acted upon by agents.

Indigenous (*in-dij'-en-us*) [*indu*, within; *gignere*, to beget]. Native; originating or belonging to a certain locality or country.

Indigestion (*in-di-jes'-chun*) [*in*, not; *digerere*, to digest]. Imperfect digestion.

Indigitation (*in-dij'-it-a'-shun*) [*in*, in; *digitus*, a finger]. Intussusception.

Indigo (*in'-dig-o*) [*induxōn*, indigo], $C_{16}H_{10}N_2O_2$. A blue pigment formed by the decomposition of the indican contained in various species of Indigofera (I. tinctoria, I. anil, I. argentea), or in the urine and sweat.

Indigocarmin (*in-dig-o-car'-min*) [*induxōn*, indigo; *carmin*]. Potassium sulphindigotate, used as a stain in microscopy and as a test for sugar.

Indirect (*in'-di-rekt*) [*in*, not; *directus*, straight]. Not direct; not in a direct line; acting through an intervening medium. 1. **Cell-division**. See *Karyokinesis*. 1. **Vision**, vision by some other part of the retina than the macula.

Indisposition (*in-dis-po-zish'-un*) [*in*, not; *dis*, apart; *ponere*, to place]. A slight illness not confining the patient to bed.

Indol (*in'-dol*) [*induxōn*, indigo], C_8H_7N . A substance produced in pancreatic digestion, in intestinal putrefaction, and in certain bacterial cultures. It occurs in the feces, giving to them in part their odor, and is eliminated in the urine in the form of indican, being especially increased in intestinal obstruction.

Indolent (*in'-do-lent*) [*in*, not; *dolere*, to feel pain]. Sluggish.

Indoxylsulphuric Acid (*in-dok-sil-sul-fu'-rik as'-id*) [*induxōn*, indigo; *ὀξύς*, sharp], $C_{16}H_9NSO_4$. A combination of indoxyl ($C_8H_7(OH)N$) with the radicle of sulphuric acid (HSO_3), the potassium salt of which occurs in the urine. See *Indican*.

Induced (*in'-dūsd'*) [*inducere*, to lead into]. 1. Produced by induction, as I. electricity. 2. Produced artificially, as I. labor.

Induction (*in-duk'-shun*) [*inducere*, to lead in]. 1. The act of bringing on. 2. The process of drawing general conclusions from special facts. 3. The production of electricity or magnetism in a body by proximity to another body, which is electrified or magnetized, but not in direct contact with it. **I.-balance**, an instrument used for detecting the presence of metallic bodies by the electric disturbance which they cause. **I.-coil**, a wire wound around a bobbin, used for conducting a galvanic current, by means of which electricity is induced in a second coil.

Indurated (*in'-du-ra-ted*) [*in*, into; *durus*, hard]. Hardened, as **I. chancre**.

Induration (*in-du-ra'-shun*) [*in*, into; *durus*, hard]. Hardening; the state of being or becoming hard; a hardened mass or lump. **I., Brown**, a form of interstitial pneumonia in which there is, in addition to the new growth of fibrous tissue, a deposit of altered blood-pigment. **I., Gray**, a similar condition without the deposit of pigment. **I., Red**, an interstitial pneumonia in which the lung is red from congestion.

Inebriant (*in-e'-bri-ant*) [*inebriare*, to make drunk]. 1. Intoxicant; causing inebriation. 2. An agent that causes inebriation.

Inebriation (*in-e-bre-a'-shun*) [*inebriare*, to make drunk]. The condition of drunkenness.

Inebriety (*in-e-bri'-et-ē*) [*inebriare*, to make drunk]. Habitual drunkenness.

Inertia (*in-er'-she-ah*) [*iners*, inactive]. Sluggishness; inability to move except by means of an external force. In physics, that property of matter by virtue of which it is incapable of changing its condition of rest or motion. **I., Uterine**, sluggishness of uterine contractions during labor.

In extremis (*in eks-tre'-mis*) [*in*, in; *extremus*, last]. At the end; at the last; at the point of death.

Infant (*in'-fant*) [*infans*, a little child]. 1. A babe. 2. According to English law, one not having attained the age of twenty-one.

Infanticide (*in-fant'-is-īd*) [*infans*, a little child; *cadere*, to kill]. The murder of an infant.

Infantile (*in'-fan-tīl*) [*infans*, a little child]. Pertaining to infancy. **I. Paralysis**. See *Paralysis*.

Infarct (*in'-farkt*) [*infarcire*, to stuff in]. A wedge-shaped area, either of hemorrhage into an organ (hemorrhagic **I.**), or of necrosis in an organ (anemic **I.**), produced by the obstruction of a terminal vessel. **I., Uric-acid**, the deposition of crystals of uric acid in the renal tubules of the newborn.

Infarction (*in-fark'-shun*) [*infarcire*, to stuff in]. The production of an infarct; also the infarct itself.

Infecting (*in-fek'-ting*) [*in*, into; *facere*, to make]. Causing infection, as an **I. embolus**.

Infection (*in-fek'-shun*) [*infectio*, from *in*, into; *facere*, to make]. 1. The communication of disease from one body to another, or from one part to another part of the same individual (autoinfection). 2. The material conveying the disease; the disease-producing agent. **I.-atrium**, the point of entrance of an infection. **I., Consecutive**, septic infection implanted upon an already established morbid process.

Infectious (*in-fek'-shus*) [*in*, into; *facere*, to make]. 1. Communicating disease. 2. Caused by an infection.

Infecundity (*in-fe-kun'-dit-ē*) [*in*, not; *fecundus*, fruitful]. Sterility, barrenness.

Inferior (*in-fe'-re-or*) [comp. of *inferus*, low]. Lower.

Infiltrate (*in'-fil-trat*) [*in*, in; *filtrare*, to strain]. 1. To ooze into the spaces of a tissue. 2. The substance that has oozed out.

Infiltration (*in-fil-tra'-shun*) [*in*, into; *filtrare*, to strain]. 1. The entrance into the tissue-spaces or into the tissue-elements of some abnormal substance or of a normal substance in excess. 2. The material thus deposited. **I., Calcareous**, the deposit of lime and magnesium salts in the tissues. **I., Cellular**, an infiltration of the tissues with round cells. **I., Fatty**, the deposit of fat in the tissues; the presence of oil or fat-globules in the interior of a cell. **I., Glycogenic**, the deposit of glycogen-granules in the cells. **I., Pigmentary**, the deposit of pigment in the tissues, derived either from without or from within. **I., Serous**, an infiltration of the tissues with diluted lymph. **I., Tuberculous**, a confluence of tuberculous nodules.

Infinite (*in'-fin-īd*) [*in*, not; *finis*, boundary]. Immeasurable or innumerable; unlimited, when compared with any known or conceivable quantity. **I. Distance**, a term in optics indicating a distance of 20 feet (6 meters) or more. Rays from an object at that distance and entering the eye are practically parallel, as they would be completely if coming from a point at a really infinite distance.

Infirm (*in-firm'*) [*in*, not; *firmus*, firm]. Weak or feeble.

Infirmity (*in-fer'-ma-re*) [*infirmarium*, an infirmary]. A hospital; an institution where ill and infirm persons are maintained during the period of treatment.

Infirmity (*in-fer'-mit-ē*) [*infirmitas*, weakness]. 1. Weakness, feebleness. 2. A disease producing feebleness.

Inflammation (*in-flam-a'-shun*) [*inflamare*, to set on fire, to inflame]. A term applied to those tissue-changes by which irritants are eliminated, and which include overfilling of the blood-vessels, alteration in the blood-vessel walls, outwandering of leukocytes,

exudation of plasma, and multiplication of the cells of the surrounding connective-tissue.

I., Acute, that in which the processes are active; usually this form is characterized by the cardinal symptoms of inflammation—heat, redness, swelling, and pain. **I., Catarrhal**, one occurring on a mucous surface and causing the shedding of its epithelium. **I., Chronic**, that in which there is a building-up of new connective tissue. **I., Interstitial**, one affecting chiefly the connective tissue of an organ. **I., Parenchymatous**, one affecting chiefly the parenchyma of an organ. **I., Reactive**, an inflammation set up around a focus of degeneration to limit the spread of the degenerative process; also the inflammation around a foreign body. **I., Specific**, one due to a special microorganism, and characterized by the formation of a tumor-like nodule that tends to degenerate. **I., Suppurative**, that attended by the formation of pus.

Inflation (*in-fla'-shun*) [*inflare*, to puff up]. The act of distending with air.

Influenza (*in-flu-en'-zah*) [Ital., an influence]. An epidemic affection characterized by catarrhal inflammation of the mucous membrane of the respiratory tract, accompanied by a mucopurulent discharge, fever, pain in the muscles, and prostration. At times symptoms referable to the gastrointestinal system predominate; at others the symptoms are mainly referred to the nervous system. The cause of the disease is the bacillus of influenza, discovered by Pfeiffer and Canon. Complications are common, pneumonia being the most frequent; pleurisy, otitis media, and neuritis also occur. An occasional sequel is insanity. Synonyms: la grippe, grip.

Infra- (*in'-frah-*) [*infra*, below]. A prefix meaning below or beneath.

Infraaxillary (*in'-frah-aks'-il-a-re*) [*infra*, below; *axilla*, the arm-pit]. Below the arm-pit.

Infraclavicular (*in'-frah-klav-ik'-u-lar*) [*infra*, below; *clavicula*, the collar-bone]. Below the collar-bone.

Infracortical (*in'-frah-kor'-tik-al*) [*infra*, below; *cortex*, a bark]. Lying beneath the cortical substance of the brain.

Infracostal (*in'-frah-kos'-tal*) [*infra*, below; *costa*, a rib]. Below the ribs.

Inframammary (*in'-frah-mam'-ar-e*) [*infra*, below; *mamma*, the breast]. Situated beneath the mamma.

Inframaxillary (*in'-frah-mak'-il-a-re*) [*infra*, below; *maxilla*, the jaw]. Below or under the jaw.

Infraorbital (*in'-frah-or'-bit-al*) [*infra*, below; *orbita*, orbit]. Beneath or below the floor of the orbit. **I. Canal**, the canal in the superior maxillary bone that transmits

the infraorbital vessels and nerve. **I. Foramen**. See *Foramina*, Table of.

Infrascapular (*in'-frah-skap'-u-lar*) [*infra*, below; *scapula*, shoulder-blade]. Below the shoulder-blade.

Infraspinous (*in'-frah-spi'-nus*) [*infra*, below; *spina*, a spine]. Beneath a spine, as of the scapula or a vertebra. **I. Fascia**, the dense membranous fascia covering the infraspinous muscle. **I. Fossa**, the shallow depression on the dorsal surface of the scapula, below the spine, and lodging the infraspinatus muscles. **I. Muscle**. See *Muscles*, Table of.

Infrasternal (*in'-frah-ster'-nal*) [*infra*, below; *sternum*, breast-bone]. Below the sternum.

Infundibuliform (*in-fun-dib-ul'-le-form*) [*infundibulum*, a funnel; *forma*, a form]. Funnel-shaped. **I. Fascia**, the funnel-shaped membranous layer that invests the spermatic cord.

Infundibulum (*in-fun-dib'-u-lum*) [*infundere*, to pour into]. A funnel-shaped passage or part. **I. of Brain**, a tubular mass of gray matter attached to the pituitary body. **I. of the Ethmoid Bone**, a canal connecting the anterior ethmoidal cells with the middle meatus of the nose. **I. of Heart**, the arterial cone from which the pulmonary artery arises. **I. of the Kidney**, one of the primary divisions of the pelvis of the kidney. **I. of the Lung**, one of the air-spaces into which a terminal bronchiole divides, and which is composed of an aggregation of air-vesicles.

Infusion (*in-fu'-zhun*) [*in*, into; *fundere*, to pour]. 1. The process of extracting the active principles of a substance by means of water, but without boiling. 2. The product of such a process, known in pharmacy as infusum.

Infusum (*in-fu'-sum*) [*in*, into; *fundere*, to pour]. An infusion. The following are official in U. S. P.: **I. cinchonæ**. Dose f ʒ ij (64.0). **I. digitalis**. Dose f ʒ ss (16.0). **I. pruni virginianæ**. Dose f ʒ ij-ijj (64.0-96.0). **I. sennæ compositum**, black draught. Dose f ʒ iv (128.0).

Ingesta (*in-jes'-tah*) [*in*, into; *gerere*, to carry]. Substances introduced into the body, especially foods.

Ingestion (*in-jes'-chun*) [*in*, into; *gerere*, to carry]. The act of taking substances, especially food, into the body.

Ingluvies (*in-glu'-ve-ēs*) [L.]. 1. The crop or craw of birds. 2. The paunch or rumen of ruminating mammals.

Ingluvin (*in'-glu-vin*) [*ingluvies*, crop or craw of a bird]. A preparation obtained from the gizzard of the fowl, *Pallus gallinaceus*, used as a substitute for pepsin and pan-

creatin, and also in the vomiting of pregnancy. Dose gr. x-xx (0.65-1.3).

Ingrassias, Processes, or Wings of [*in-grassias*, an Italian physician of the sixteenth century]. The lesser wings of the sphenoid bone.

Ingravescent (*in-grav-es'-ent*) [*ingravesce*, to become heavy]. Increasing in severity, as I. apoplexy.

Ingrowing Nail. See *Nail*.

Inguen (*in'-gwen*) [L.]. The groin.

Inguinal (*in'-gwin-al*) [*inguen*, the groin]. Pertaining to the groin. **I. Canal**, the canal transmitting the spermatic cord in the male, and the round ligament in the female. It is situated parallel to and just above Poupart's ligament. **I. Glands**, the superficial and the deep glands of the groin. **I. Hernia.** See *Hernia*.

Inhalation (*in-ha-la'-shun*) [*inhalare*, to draw in]. The breathing in of air or other vapor. **I.-diseases**, those due to the inspiration of air containing dust or any finely divided matter.

Inhaler (*in-ha'-ler*) [*inhalare*, to breathe in]. An instrument for inhaling a gas or vapor.

Inherent (*in-he'-rent*) [*in*, to; *herere*, to cleave]. Innate; natural to the organism.

Inhibition (*in-hib-ish'-un*) [*inhibere*, to check]. The act of checking or restraining; a restraint.

Inhibitory (*in-hib'-it-o-re*) [*inhibere*, to check]. Checking; restraining.

Iniac, or Inial (*in'-e-ac, in'-e-al*) [*iniov*, occiput]. Pertaining to the inion.

Iniencephalus (*in-e-en-sef'-al-us*) [*iniov*, occiput; *ἐγκεφαλος*, brain]. A fetal monstrosity in which there is a posterior fissure of the skull, with protrusion of the brain-substance, combined with spinal fissure.

Inion (*in'-e-on*) [*iniov*, occiput]. The external protuberance of the occipital bone. See *Cranimetric Points*.

Initial (*in-ish'-al*) [*in*, into; *ire*, to go]. Beginning; early; primary, as the I. lesion of syphilis,—the chancre. **I. Cells**, germ-cells. **I. Sclerosis**, the hard chancre.

Inject (*in-jekt'*) [*in*, into; *jacere*, to throw]. To throw or force in, as to inject fluids into the tissues; also, to fill the vessels of an organ.

Injection (*in-jekt'-shun*) [*in*, into; *jacere*, to throw]. 1. The act of injecting or throwing in. 2. The substance injected. According to the organ into which the injection is made, different terms are employed, as *e. g.*, urethral I., intramuscular I., uterine I., vaginal I., etc. In the B. P. **Injections** are solutions of active substances used for hypodermic injection. **I. Hypodermic**, an injection made under the skin.

Injector (*in-jekt'-tor*) [*injicere*, to throw in]. An apparatus used in injecting.

Injury (*in-ju'-re*) [*in*, not; *jus*, a right]. A harm or hurt to the body.

Innervation (*in-ner-va'-shun*) [*in*, in; *nervus*, nerve]. Nerve-supply.

Innominate (*in-nom'-in-at*) [*in*, without; *nomen*, a name]. Unnamed; unnamable. **I.**

Artery. See *Arteries, Table of*. **I. Bone**, the irregular bone forming the sides and anterior wall of the pelvic cavity, and composed of the ilium, ischium, and pubis.

Innominatum (*in-nom-in-a'-tum*) [L., nameless (*os*, bone, understood)]. The innominate bone.

Innutrition (*in-nu-trish'-un*) [*in*, not; *nutrire*, to nourish]. Want of nutrition or nourishment.

Inoblast (*in'-o-blast*) [*ic*, fiber; *βλαστος*, germ]. Any one of the cells from which connective tissue is derived.

Inoculability (*in-ok-u-la-bil'-it-e*) [*in*, into; *oculus*, a bud]. The quality of being inoculable.

Inoculable (*in-ok-u-la-bl*) [*in*, into; *oculus*, a bud]. Capable of being inoculated; communicable by inoculation.

Inoculation (*in-ok-u-la'-shun*) [*in*, into; *oculus*, a bud]. 1. The act of introducing the virus of a disease into the body. 2. Specifically, the intentional introduction of a virus for the purpose of producing a mild form of a disease which is severe when spontaneously introduced, as the I. of small-pox virus. This is known as preventive I.

Inogen (*in'-o-jen*) [*ic*, fiber; *γενῶν*, to produce]. A hypothetic substance believed to occur in muscular tissue, and to be decomposed, during contraction, into carbon dioxide, sarcosolactic acid, and myosin.

Inorganic (*in-or-gan'-ik*) [*in*, not; *ὄργανον*, an implement]. Not organic; not produced by animal or vegetal organisms, as an I. compound. **I. Chemistry**, chemistry dealing with inorganic compounds.

Inosculate (*in-os'-cu-lāt*) [*in*, in, on; *oculum*, dim. of *os*, mouth]. To unite by small openings.

Inosculation (*in-os-ku-la'-tion*) [*in*, in; *oculum*, a small mouth]. The joining of blood-vessels by direct communication.

Inosite (*in'-o-sit*) [*ic*, fiber]. $C_6H_{12}O_6 + 2H_2O$. Muscle-sugar; a saccharine substance occurring in muscles, rarely in urine.

Inosituria (*in-o-sit-u'-re-ah*) [*ic*, fiber; *urina*, urine]. The presence of inosite in the urine.

Inquest (*in'-kwest*) [*in*, into; *querere*, to ask]. A judicial inquiry, especially one for the purpose of determining the cause of death of one who has died by violence or in some unknown way.

Inquisition (*in-kwis-ish'-un*) [*in*, into; *querere*, to ask]. An inquiry, especially one into the sanity or lunacy of a person.

Insalivation (*in-sal-iv-a'-shun*) [*in*, in; *sal-iva*, the spittle]. The mixture of the food with saliva during mastication.

Insane (*in-sān'*) [*in*, not; *sanus*, sound]. Deranged or diseased in mind. **I. Ear.** See *Hematoma auris*.

Insanitary (*in-san'-it-a-re*) [*in*, not; *sanitas*, health]. Not sanitary; not in a proper condition as respects the preservation of health.

Insanity (*in-san'-it-e*) [*in*, not; *sanus*, sound]. A derangement of the mental faculties, with or without loss of volition and of consciousness. Insanity may be due to defective development, to acquired disease, or to natural decay. It is characterized, according to its form, by a variety of symptoms, the most common of which are change of character and habits, moroseness, confusion, elation, melancholy, mania, delusions, and hallucinations. Melancholia, mania, delusional insanity, and dementia are the four principal types of the affection. **I., Acquired**, that arising after a long period of life of mental integrity. **I., Alcoholic**, that induced by alcoholic excess, usually a result of hereditary tendencies. **I., Alternating.** See *I., Circular*. **I., Circular**, cyclothymia; alternating insanity; a form of insanity recurring in cycles varying in length from a few days to many months. The arrangement of the cycle varies in different individuals, but is constant in a given case. Thus melancholia may be followed by mania, and this by a lucid interval, the passage from one mental condition to the other being abrupt or gradual. **I., Climacteric**, insanity occurring at or near the menopause. **I., Communicated**, that transmitted by association with an insane person. **I., Confusional**, an acute insanity produced by nervous shock or exhausting disease, without distinct constant emotional depression or exaltation, with marked failure of mental power or complete imbecility, often accompanied by hallucinations and loss of physical power. Recovery is usually complete. **I., Cyclic.** Same as *I., Circular*. **I., Doubting**, a form closely allied to delusional insanity, consisting in an uncontrollable doubt and indecision regarding the occupations, duties, or events of the day, of religion, etc. **I., Emotional**, insanity characterized by derangement of the emotions, either depressing or exalting in character. **I., Epidemic**, a form occasionally manifested among a number of persons in common association, as in convents or schools. **I., Hereditary**, that transmitted from parent to child, and not induced by other apparent cause. **I., Impulsive**, a form in which the patient possesses an uncontrollable desire to commit acts of vio-

lence. **I., Menstrual.** See *I., Periodic*. **I., Moral**, a form marked by perversion and depravity of the moral sense, apparently without impairment of the reasoning and intellectual faculties. **I., Periodic**, a condition dependent upon original or acquired psychopathy, in which attacks of insanity occur at regular or irregular intervals. If occurring in women at the menstrual epoch, it is called menstrual insanity. **I. of Pregnancy**, a form occurring during pregnancy, characterized by melancholia, suicidal intent, and abhorrence of friends and relatives. **I., Primary**, a form, often congenital, that arises with the development of the body. It may also proceed from injury or disease of the brain in early life. **I. of Puberty.** See *Hebephrenia*. **I., Puerperal**, a term sometimes applied to the delirium of childbirth, but more properly to the insanity occurring after delivery. **I., Recurrent**, that marked by recurrent attacks of mental aberration with intervening lucid intervals. **I., Religious**, that associated with religious subjects. **I., Stuporous**, a primary acute form of dementia, characterized by a tendency to stupor; a disease chiefly met with in youth and early maturity. **I., Surgical**, that coming on after surgical operations. **I., Toxic**, an acute form due to systemic poisoning by certain drugs.

Inscriptiones tendineæ (*in-skrip-te-o'-ne ten-din'-e-e*). The lineæ transversæ of the rectus abdominis muscle.

Insect (*in'-sekt*) [*in*, into; *secare*, to cut]. Any member of the class of animals called Insecta. **I. Powder**, a powder employed to destroy or ward off insects, and consisting usually of the powdered flowers of species of *Pyrethrum*.

Insecticide (*in-sek'-tis-id*) [*insectum*, insect; *cædere*, to kill]. A substance that is destructive to insects.

Insemination (*in-sem-in-a'-shun*) [*inseminare*, to plant seed]. 1. The planting of seed. 2. The introduction of semen.

Insensible (*in-sen'-sib-l*) [*in*, not; *sentire*, to feel]. 1. Incapable of being perceived or recognized by the senses. 2. Unconscious.

Insertion (*in-ser'-shun*) [*inserere*, to set in]. 1. The act of setting or placing in. 2. That which is set in. 3. The point at which anything, as a muscle, is attached; the place or the mode of attachment of an organ to its support.

Insidious (*in-sid'-e-us*) [*insidiæ*, ambush]. Coming on stealthily or imperceptibly. **I.**

Disease, one, the onset of which is gradual or inappreciable.

In situ (*in si'-tu*) [*in*, in; *situs*, position]. In a given or natural position.

Insolation (*in-so-la'-shun*) [*in*, in; *sol*, sun].

1. Exposure to the rays of the sun. 2. Sun-stroke or heatstroke.

Insoluble (*in-sol'-u-bl*) [*in*, not; *soluere*, to solve]. Incapable of dissolving.

Insomnia (*in-som'-ne-ah*) [*in*, not; *somnus*, sleep]. Want of sleep; inability to sleep.

Inspection (*in-spek'-shun*) [*inspicere*, to look]. In medicine, the examination of the body or any part of it by the eye.

Inspiration (*in-spir'-a'-shun*) [*in*, in; *spirare*, to breathe]. The drawing in of the breath.

Inspiratory (*in-spi'-ra-to-re*) [*in*, in; *spirare*, to breathe]. Pertaining to the act of inspiration.

Inspissate (*in'-spis'-at*) [*inspissare*, to thicken]. To make thick by evaporation or by absorption of fluid.

Instep (*in'-step*) [*instop*, the bend of the foot]. The arch on the upper surface of the foot.

Instillation (*in-stil'-a'-shun*) [*instillare*, to put in little by little]. The pouring of a liquid into a cavity drop by drop.

Instinct (*in'-stinkt*) [*instinguere*, to impel]. A natural impulse, which, though unassociated with reason, prompts a useful act.

Instinctive (*in-stinkt'-tiv*) [*instinguere*, to impel]. Prompted or determined by instinct; of the nature of instinct.

Institutes of Medicine. The philosophy of the science of medicine, of physiology, pathology, therapeutics, and hygiene, or the general and elementary principles of the same. The term is used sometimes as a synonym of physiology.

Instrument (*in'-stru-ment*) [*in*, in; *struere*, to build]. Any mechanical tool or device used to assist in the performance of a certain act.

Instrumental (*in'-stru-men'-tal*) [*in*, in; *struere*, to build]. Pertaining to or performed with instruments, as I. labor.

Instrumentation (*in-stru-men-ta'-shun*) [*in*, in; *struere*, to build]. The care or employment of instruments.

Insufficiency (*in-suf-fish'-en-se*) [*insufficiencia*; *in*, not; *sub*, under; *facere*, to make]. The state of being inadequate; incapacity to perform a normal function. I. of the Cardiac Valves, imperfect closure of the valves, permitting regurgitation. Depending upon the valve affected, the I. may be aortic, mitral, tricuspid, or pulmonary. I. of a Muscle, inability on a part of a muscle to contract sufficiently to produce the normal effect. The term is applied especially to the eye-muscles. I. of the Externi, a condition in which the contraction of the externi muscles of the eye is weak and is overbalanced by that of the interni, producing esophoria. I. of the Interni, defective power on the part of the interni muscles, producing exophoria.

Insufflation (*in-suf-fla'-shun*) [*in*, in; *sufflare*, to puff]. The act of blowing into, as the I. of a powder into a cavity; also, the blowing of air into a cavity, as I. of the middle ear. I., Mouth-to-mouth, the blowing of air into the mouth of a person, usually a new-born infant, to distend the lungs and counteract asphyxia.

Insufflator (*in'-suf-la-tor*) [*in*, in; *sufflare*, to blow]. An instrument for blowing air or powders into a cavity.

Insula (*in'-su-lah*) [L.]. The island of Reil.

Insular (*in'-su-lar*) [*insula*, an island]. I. Pertaining to the island of Reil. 2. Isolated; occurring in patches. I. Sclerosis. See Sclerosis.

Insulate (*in'-su-lat*) [*insula*, an island]. To isolate or separate from surroundings. In electricity, to surround a conductor with a nonconducting substance.

Integument (*in-teg'-u-ment*) [*in*, upon; *tegere*, to cover]. A covering, especially the skin. I., Fetal, the fetal membranes.

Intellect (*in'-tel-ekt*) [*intellectus*; *inter*, between; *legere*, to choose]. The mind or the reasoning power.

Intemperance (*in-tem'-per-ans*) [*in*, not; *temperare*, to moderate]. Want of moderation; immoderate indulgence, especially in alcoholic beverages.

Intensification (*in-ten-sif-ik'-a'-shun*) [*in*, upon; *tendere*, to stretch]. The act of making intense, or of increasing the strength of anything.

Intensity (*in-ten'-sit-e*) [*in*, upon; *tendere*, to stretch]. 1. The state of being intense or high-strung. 2. The degree to which a force is capable of rising. 3. A high degree of energy or power.

Intensive (*in-ten'-siv*) [*in*, upon; *tendere*, to stretch]. Gradually increased in force or intensity, as the I. method of inoculation.

Intention (*in-ten'-shun*) [*in*, upon; *tendere*, to stretch]. The end or purpose. See Healing. I.-tremor, a tremor coming on when attempts at voluntary motion are made.

Inter- (*in'-ter-*) [*inter*, between]. A prefix signifying between.

Interarticular (*in-ter-ar-tik'-u-lar*) [*inter*, between; *artculus*, a joint]. Situated between joints. I. Fibrocartilage, the flattened cartilaginous plates between the articular cartilages of certain joints.

Interauricular (*in-ter-aw-rik'-u-lar*) [*inter*, between; *auricula*, auricle]. Situated between the auricles.

Interbrain (*in'-ter-bran*). See *Thalamencephalon*.

Intercadence (*in-ter-kat'-dens*) [*inter*, between; *cadere*, to fall]. An irregular beating of the pulse, in which an additional beat is interposed between two pulsations.

Intercalary, Intercalated (*in-ter'-kal-a-re, in-ter'-kal-a-ted*) [*inter*, between; *calare*, to insert]. Placed or inserted between.

Intercarotid (*in-ter-kar-ot'-id*) [*inter*, between; *καρτεν*, to produce sleep]. Situated between the external and internal carotid arteries, as the I. ganglion.

Intercellular (*in-ter-sel'-u-lar*) [*inter*, between; *cellula*, a small cell]. Between cells, as I. substance of tissue.

Intercentral (*in-ter-sen'-tral*) [*inter*, between; *centrum*, a center]. Between centers.

Interchondral (*in-ter-kon'-dral*) [*inter*, between; *χόνδρος*, cartilage]. Between cartilages.

Interclavicular (*in-ter-klav-ik'-u-lar*) [*inter*, between; *clavicula*, the collar-bone]. Between the clavicles.

Intercolumnar (*in-ter-kol-um'-nar*) [*inter*, between; *columna*, column]. Between pillars, as the I. fascia, between the pillars of the external abdominal ring.

Intercondyloid (*in-ter-kon'-dil-oid*) [*inter*, between; *κόνδυλος*, a knob; *ειδος*, likeness]. Between condyles. I. Fossa, the notch between the condyles of the femur.

Intercostal (*in-ter-kos'-tal*) [*inter*, between; *costa*, a rib]. Between the ribs. I. Arteries, the arteries of the intercostal spaces. See *Arteries, Table of*. I. Muscles. See *Muscles, Table of*. I. Nerves, the anterior divisions of the dorsospinal nerves.

Intercostohumeral (*in-ter-kos-to-hu'-mer-al*) [*inter*, between; *costa*, rib; *humerus*, the bone of the upper arm]. Pertaining to the arm and the space between the ribs, as the I. nerve.

Intercurrent (*in-ter-kur'-ent*) [*inter*, between; *currere*, to run]. Occurring or taking place between. I. Disease, a disease arising or progressing during the existence of another disease in the same person.

Interdigital (*in-ter-dij'-it-al*) [*inter*, between; *digitus*, a finger]. Between the fingers.

Interdigitation (*in-ter-dij-it-a'-shun*) [*inter*, between; *digitus*, a finger]. The locking or dovetailing of similar parts, as the fingers of one hand with those of the other; or of the ends of the obliquus externus muscle with those of the serratus magnus.

Interfascicular (*in-ter-fas-ik'-u-lar*) [*inter*, between; *fasciculus*, a bundle]. Situated between fasciculi.

Interference (*in-ter-fe'-rens*) [*inter*, between; *ferire*, to strike]. The act of interfering or preventing. I. of Light, the mutual neutralization of waves of light, when the crest of one wave falls upon the trough of another. I. of Sound, the neutralization of two sound waves, one by the other.

Interfibrillar (*in-ter-fi'-bril-ar*) [*inter*, between; *fibra*, a fiber]. Situated between the fibrillæ of tissues.

Interganglionic (*in-ter-gang-le-on'-ik*) [*inter*, between; *γάγγλιον*, a ganglion]. Connecting one ganglion with another; lying between ganglia.

Interlamellar (*in-ter-lam-el'-ar*) [*inter*, between; *lamella*, a layer]. Between the lamellæ.

Interlobar (*in-ter-lo'-bar*) [*inter*, between; *lobus*, a lobe]. Situated between lobes, as I. pleurisy.

Interlobular (*in-ter-lob'-u-lar*) [*inter*, between; *lobulus*, a lobule]. Between lobules.

Intermarriage (*in-ter-mar'-āj*) [*inter*, between; *maritare*, to marry]. Marriage between persons related by consanguinity, or between persons of different races.

Intermaxillary (*in-ter-maks'-il-a-re*) [*inter*, between; *maxilla*, jaw-bone]. Between the maxillary bones. I. Bone, the small bone that receives the incisors, situated between the superior maxillary bones of the fetus.

Intermediate (*in-ter-me'-de-at*) [*inter*, between; *medius*, middle]. Situated between.

Intermediolateral (*in-ter-me-de-o-lat'-er-al*) [*inter*, between; *medius*, middle; *latus*, side]. Both lateral and intermediate, as the I. tract of the spinal cord.

Intermeningeal (*in-ter-men-in'-je-al*) [*inter*, between; *μηνιγξ*, membrane]. Between the dura and the arachnoid, or between the latter and the pia. I. Hemorrhage, a hemorrhage between the meninges of the brain or spinal cord.

Intermenstrual (*in-ter-men'-stru-al*) [*inter*, between; *mensis*, month]. Between the menstrual periods.

Intermetacarpal (*in-ter-met-a-kar'-pal*) [*inter*, between; *μετά*, beyond; *καρπός*, the wrist]. Between the metacarpal bones.

Intermetatarsal (*in-ter-met-a-tar'-sal*) [*inter*, between; *μετά*, beyond; *ταρσός*, tarsus]. Between the metatarsal bones.

Intermission (*in-ter-mish'-un*) [*inter*, between; *mittere*, to send]. An interval, as between the paroxysms of a fever, or between the beats of the pulse.

Intermittent (*in-ter-mit'-ent*) [*inter*, between; *mittere*, to send or occur]. Occurring at intervals; characterized by intermissions or intervals, as I. fever, I. insanity, I. pulse, I. sterilization.

Intermuscular (*in-ter-mus'-ku-lar*) [*inter*, between; *musculus*, a muscle]. Situated between muscles.

Intern (*in'-tern*) [Fr., *interne*]. An in-door or resident physician in a hospital.

Internal (*in-ter'-nal*) [*internus*, inward]. Situated within or on the inside. I. Capsule, the band of white nerve-matter between the

optic thalamus and caudate nucleus on the inner, and the lenticular nucleus on the outer side. It is the continuation of the crus cerebri, and consists of an anterior and a posterior limb joined at an angle, termed the knee. It is composed of fibers coming from and going to the cortex cerebri. **I. Medicine**, that branch of medicine which treats of disease affecting the internal organs. **I. Rectus**. See *Muscles*, Table of. **I. Resistance**. See *Resistance*.

Internodal (*in-ter-no'-dal*). See *Internode*.

Internode (*in'-ter-nod'*) [*inter*, between; *nodus*, a knot]. The space between two nodes of a nerve-fiber, as the I. between the nodes of Ranvier, also termed internodal segment.

Internuncial (*in-ter-nun'-she-al*) [*inter*, between; *nuncius*, a messenger]. Serving as a connecting or announcing medium, as I. fibers, nerve-fibers connecting nerve-cells.

Internus (*in-ter'-nus*). **I.** See *Internal*. **2.** The internal rectus muscle of the eye.

Interorbital (*in-ter-or'-bit-al*) [*inter*, between; *orbita*, the orbit]. Situated between the orbits.

Interosseous (*in-ter-os'-e-us*) [*inter*, between; *os*, a bone]. Between bones, as I. arteries, membrane, muscles, or nerves.

Interparietal (*in-ter-par-i'-e-tal*) [*inter*, between; *paries*, walls]. Between walls; between the parietal bones, as I. suture; between parts of the parietal lobe, as I. fissure.

Interpeduncular (*in-ter-pe-dung'-ku-lar*) [*inter*, between; *pedunculus*, a little foot]. Situated between the cerebral or cerebellar peduncles. **I. Space**, the pons Varolii, or posterior perforated space that forms the posterior portion of the floor of the third ventricle.

Interphalangeal (*in-ter-fa-lan'-je-al*) [*inter*, between; *phalanx*, a finger]. Between the fingers or the toes.

Interpubic (*in-ter-pu'-bi-*) [*inter*, between; *pubis*, pubis]. Situated between the pubic bones.

Interradial (*in-ter-ra'-de-al*) [*inter*, between; *radius*, a ray]. Situated between two rays.

Interrupted (*in-ter-up'-ted*) [*interrumpere*, to break apart]. Discontinuous; broken; irregular.

Interrupter (*in-ter-up'-ter*) [*interrumpere*, to break apart]. That which interrupts; specifically, a device for breaking an electric current.

Interscapular (*in-ter-skap'-u-lar*) [*inter*, between; *scapula*, the shoulder-blade]. Between the shoulder-blades.

Interstitial (*in-ter-stish'-al*) [*inter*, between; *stere*, to place]. **I.** Situated between im-

portant parts; occupying the interspaces or interstices of a part. **2.** Pertaining to the interstitial or connective tissue. **I. Inflammation**, inflammation of the interstitial or connective tissue. **I. Keratitis**. See *Keratitis*.

Intertransversales (*in-ter-trans-ver-sa'-les*) [*intertransverse*]. Short bundles of muscular fibers extending between the transverse processes of contiguous vertebræ.

Intertransverse (*in-ter-trans-vers'*) [*inter*, between; *transversus*, turned across]. Connecting the transverse processes of contiguous vertebræ.

Intertrigo (*in-ter-tri'-go*) [*inter*, between; *terere*, to rub]. An erythematous eruption of the skin produced by friction of adjacent parts.

Intertrochanteric (*in-ter-tro-kan-ter'-ik*) [*inter*, between; *trochanter*, trochanter]. Between the trochanters. **I. Line**, See *Lines*, Table of.

Intertubular (*in-ter-tu'-bu-lar*) [*inter*, between; *tubulus*, a tube]. Between tubes.

I. Substance, the translucent, granular substance of the dentine of the tooth.

Interureteric (*in-ter-u-re-ter'-ik*) [*inter*, between; *ureter*, ureter]. Situated between the ureters.

Interval (*in'-ter-val*) [*inter*, between; *vallum*, a rampart]. A space or lapse either of time or distance, as the interval between the paroxysms of a fever, or between two organs or parts of the body. **I., Focal**, the distance between the anterior and posterior focal points.

Interventricular (*in-ter-ven-trik'-u-lar*) [*inter*, between; *ventriculum*, a ventricle]. Situated between ventricles. **I. Septum**, the partition between the ventricles of the heart.

Intervertebral (*in-ter-ver'-te-bral*) [*inter*, between; *vertebra*, a bone of the spine]. Between the vertebræ. **I. Discs**, the discs of fibrocartilage between the adjacent surfaces of the bodies of the vertebræ. **I. Foramen**. See *Foramen*. **I. Notch**, the notch at the base of the pedicle on the sides of the body of each vertebra.

Intervillous (*in-ter-vill'-us*) [*inter*, between; *villus*, a tuft of hair]. Situated between villi.

Intestinal (*in-tes'-tin-al*) [*intestinum*, the intestine]. Pertaining to the intestine. **I. Absorption**, the absorption of the products of digestion by the capillaries, veins, and lacteals of the mucous membrane of the intestines. **I. Anastomosis**. See *Anastomosis*. **I. Canal**, the entire intestinal passage from the beginning of the duodenum to the anus. **I. Concretion**. See *Enterolith*. **I. Juice**, succus entericus, the secretion of the intestinal glands, a pale-yellow fluid, alkaline in reaction, having a specific gravity of 1.011,

and possessing diastatic and proteolytic properties. It also, to a certain extent, emulsifies and decomposes fats.

Intestine (*in-tēs'-tin*) [*intestinus*, intestine, from *intus*, within]. The part of the digestive tube extending from the beginning of the pylorus to the anus. It consists of the small and large intestine. The former is about 6¾ meters (20 feet) in length, and extends from the pylorus to its junction with the large intestine at the cecum. Three divisions are described,—the duodenum, 22 cm. long, is the most important; the jejunum, 2.2 meters long, and the ileum, 4 meters long. The large intestine is about 1.6 meters (5 feet) long, and consists of the cecum (with the appendix vermiformis), the colon, and the rectum. The wall of the intestine is made up of four coats,—a serous, muscular, submucous, and mucous. Embedded in the wall are minute glands, and projecting from the surface, in the small intestine, are the villi. The function of the intestine is to continue and complete the changes begun in the mouth and stomach, and to remove the waste-matter or feces.

Intima (*in'-tim-ah*) [*tunica*, membrane, understood]. The innermost of the three coats of an artery.

Intolerance (*in-tol'-er-ans*) [*in*, not; *tolerare*, to bear]. The inability to endure the effect of anything, as *e. g.*, a drug.

Intoxication (*in-toks-ik-a'-shun*) [*in*, in; *τοξικόν*, poison]. 1. Poisoning. 2. The acute state produced by overindulgence in alcohol.

Intra- (*in'-trah-*) [*intra*, within]. A prefix signifying within.

Intraabdominal (*in-trah-ab-dom'-in-al*) [*intra*, within; *abdomen*]. Within the cavity of the abdomen.

Intraarticular (*in-trah-ar-tik'-u-lar*) [*intra*, within; *artculus*, joint]. Within a joint.

Intracapsular (*in-trah-kap'-su-lar*) [*intra*, within; *capsula*, capsule]. Within the capsular ligament of a joint, as I. fracture.

Intracartilaginous (*in-trah-kar-til-aj'-in-us*) [*intra*, within; *cartilago*, cartilage]. Within a cartilage, as I. ossification.

Intracellular (*in-trah-sel'-u-lar*) [*intra*, within; *cellula*, a little cell]. Within the cell.

Intracerebral (*in-trah-ser'-e-bral*) [*intra*, within; *cerebrum*, cerebrum]. Within the cerebrum.

Intracervical (*in-trah-ser'-vik-al*) [*intra*, within; *cervix*, cervix]. Within the cervical canal of the uterus.

Intracranial (*in-trah-kra'-ne-al*) [*intra*, within; *κράνιον*, the skull]. Within the skull.

Intradermic (*in-trah-der'-mik*) [*intra*, within; *δέρμα*, skin]. Within the skin.

Intradural (*in-trah-du'-ral*) [*intra*, within; *durus*, hard]. Situated or occurring within the dura.

Intraligamentous (*in-trah-lig-am-en'-tus*) [*intra*, within; *ligamentum*, a ligament]. Within or between the folds of a ligament, as an I. cyst.

Intralobular (*in-trah-lob'-u-lar*) [*intra*, within; *lobulus*, a little lobe]. Within a lobule, as I. vein of the liver.

Intramembranous (*in-trah-mem'-bran-us*) [*intra*, within; *membrana*, a membrane]. Developed or taking place within a membrane, as I. ossification.

Intrameningeal (*in-trah-men-in'-je-al*) [*intra*, within; *μηνίγξ*, membrane]. Situated within the substance of the membranes of the brain or spinal cord.

Intramural (*in-trah-mu'-ral*) [*intra*, within; *mura*, a wall]. Within the substance of the walls of an organ, as I. fibroid of the uterus.

Intramuscular (*in-trah-mus'-ku-lar*) [*intra*, within; *musculus*, a muscle]. Within the substance of a muscle.

Intranasal (*in-trah-na'-sal*) [*intra*, within; *nasus*, nose]. Within the cavity of the nose.

Intraocular (*in-trah-ok'-u-lar*) [*intra*, within; *oculus*, eye]. Within the globe of the eye, as I. hemorrhage.

Intraorbital (*in-trah-or'-bit-al*) [*intra*, within; *orbita*, orbit]. Within the orbit.

Intraparietal (*in-trah-par-i'-e-tal*) [*intra*, within; *paries*, a wall]. 1. Within the wall of an organ. 2. Within the parietal region of the cerebrum, as the I. fissure.

Intrapelvic (*in-trah-pel'-vik*) [*intra*, within; *pelvis*, basin]. Within the pelvic cavity.

Intraperitoneal (*in-trah-per-it-on-e'-al*) [*intra*, within; *περιτόναιον*, the peritoneum]. Within the peritoneum.

Intrapleural (*in-trah-plu'-ral*) [*intra*, within; *πλευρά*, a rib]. Within the pleural cavity.

Intrapolar (*in-trah-po'-lar*) [*intra*, within; *polus*, pole]. Between two poles.

Intrapulmonary (*in-trah-pul'-mon-a-re*) [*intra*, within; *pulmo*, the lung]. Within the substance of the lung.

Intraspinal (*in-trah-spi'-nal*) [*intra*, within; *spina*, spine]. Within the spinal canal.

Intratubal (*in-trah-tu'-bal*) [*intra*, within; *tuba*, a trumpet]. Within a Fallopian tube.

Intratympanic (*in-trah-tim-pant'-ik*) [*intra*, within; *tympanum*, the ear-drum]. Within the tympanic cavity.

Intrauterine (*in-trah-u'-ter-in*) [*intra*, within; *uterus*, womb]. Within the uterus.

Intravascular (*in-trah-vas'-ku-lar*) [*intra*, within; *vasculum*, a small vessel]. Within the blood-vessels.

Intravenous (*in-trah-ve' nus*) [*intra*, within; *vena*, a vein]. Within, or into the veins.
I. Injection, the introduction of a solution directly into a vein.

Intravesical (*in-trah-vest' ik-al*) [*intra*, within; *vesica*, bladder]. Within the bladder.

Intrinsic (*in-trin' sik*) [*intrinsicus*, on the inside]. Inherent; situated within; peculiar to a part, as the I. muscles of the larynx.

Intro- (*in' tro-*) [*intro*, within]. A prefix signifying within.

Introitus (*in-tro' it-us*) [*intro*, within; *ire*, to go]. An aperture or entrance. **I.** pelvis, the inlet of the pelvis. **I.** vaginae, the entrance to the vagina.

Intromission (*in-tro-mish' un*) [*intro*, within; *mittere*, to send]. The introduction of one body into another.

Introsusception (*in-tro-sus-sep' shun*) [*intro*, within; *suscipere*, to receive]. Intus-susception.

Introversion (*in-tro-ver' shun*) [*intro*, within; *vertere*, to turn]. A turning within, as a sinking within itself of the uterus.

Intubation (*in-tu-ba' shun*) [*in*, in; *tubus*, a pipe]. The introduction of a tube into a part, particularly of a tube into the larynx, to allow the entrance of air into the lungs, as in diphtheria.

Intumescence (*in-tu-mes' ens*) [*intumescere*, to swell]. A swelling.

Intumescencia (*in-tu-mes-ent' she-ah*) [*L.*]. A swelling. **I.** gangliformis, the reddish gangliform swelling of the facial nerve in the aqueductus Fallopii.

Intussusception (*in-tus-sus-ep' shun*) [*intus*, within; *suscipere*, to receive]. Invagination or slipping of one part of the intestine into the part beyond. It is most frequent in the young, occurring as a rule on the right side, the ileum slipping into the ascending colon, carrying the ileocecal valve in front of it. The condition is characterized by pain, tenesmus, frequent small bloody stools, the presence of a sausage-shaped tumor in the flank, and often, on rectal examination, of a mass in the rectum.

Inula (*in' u-lah*) [*L.*]. Elecampane. The root of *I. helenium*, a plant of the natural order Compositae, containing a principle resembling starch and termed inulin ($3C_{12}H_{20}O_{10} + H_2O$), a crystalline body, alantol ($C_{10}H_{16}O$), and helenin (C_8H_8O). Elecampane is tonic, stimulant, diaphoretic, diuretic, emmenagogue, and expectorant, and has been used in amenorrhea, dropsy, and in scaly skin-diseases. Dose gr. xx-3 (1.3-4.0).

Inunction (*in-un' k' shun*) [*inungere*, to anoint]. The act of rubbing an oily or fatty substance into the skin. Also, the substance used.

Invagination (*in-vaj-in-a' shun*) [*in*, in; *vagina*, a sheath]. The act of ensheathing or becoming ensheathed.

Invalid (*in' val-id*) [*in*, not; *valere*, to be well]. 1. Not well. 2. One who is not well, especially one who is chronically ill or whose convalescence is slow. 3. Suitable for an invalid person, as I. diet, I. chair.

Invasion (*in-va' shun*) [*in*, upon; *vadere*, to go]. The onset, especially that of a disease. Also, the manner in which the disease begins its attack.

Inversion (*in-ver' shun*) [*in*, not; *vertere*, to turn]. 1. The act of turning inward. 2. A turning upside down. 3. In chemistry, the conversion of a dextrorotatory compound into one that is levorotatory.

Invert Sugar. A sugar that turns rays of polarized light to the left. The term is usually applied to levulose or to a mixture of dextrose and levulose.

Invertebrata (*in-ver-te-bra' tah*) [*in*, not; *vertebra*, vertebra]. Animals that have no spinal column.

Invertin (*in-ver' tin*) [*in*, not; *vertere*, to turn]. A ferment found in the intestinal juice, and produced by several species of yeast plant; it converts cane-sugar in solution into invert-sugar.

Investing (*in-ves' ting*) [*investire*, to invest]. Ensheathing, surrounding.

Inveterate (*in-vet' er-at*) [*in*, with an intensive force; *vetus*, old]. Long established; obstinate, as an I. skin-disease.

In vitro (*in vit' ro*) [*L.*]. Within glass, especially within test-tubes.

Involucrum (*in-vol-u' krum*) [*involvere*, to enwrap]. The covering of a part. The sheath of bone enveloping a sequestrum.

Involuntary (*in-vol' un-ta-re*) [*in*, not; *velle*, to will]. Performed or acting independently of the will. **I.** Muscles, those that are not governed by the will.

Involution (*in-vo-lu' shun*) [*involvere*, to roll upon]. 1. A turning or rolling inward. 2. The retrogressive change to their normal condition that certain organs undergo after fulfilling their functional purposes. **I.** of the Uterus, the return of the uterus after gestation to its normal weight and condition. **I.** forms, a term applied to microorganisms that have undergone degenerative changes as a result of unfavorable environment.

Iodic Acid (*i-od' ik*). See *Acid*.

Iodid (*i' o-did*) [*iōdos*, violet-colored, from *iov*, a violet; *eidōs*, appearance]. A compound of iodine with a base.

Iodin, **Iodum** (*i' o-din*, *i-od' dum*) [*iōdos*, violet-colored, from *iov*, a violet; *eidōs*, appearance]. Symbol I; atomic weight 126.53; quantivalence 1; specific gravity 4.948 at 17° C. (62.6° F.). A nonmetallic

element with metallic luster, volatilizing at a low temperature, and giving off an irritating crimson-purple vapor. It occurs in most marine plants, in shell-fish, and in cod-liver oil. It is soluble in alcohol, in solutions of potassium iodid and of sodium chlorid. It is a powerful irritant, and is used chiefly as an alterative in scrofula and rickets; as an absorbent in goiter and lymphatic enlargements; as a counterirritant, and to produce inflammatory reaction in hydrocele and other cysts. The long-continued use of iodine and its preparation produces a form of poisoning termed iodism. See *Iodism*. **I.-green**, a green pigment derived from coal-tar, used in histologic work. **Linimentum iodi** (B. P.), used locally. **Liquor iodi compositus**, Lugol's solution. Dose m j-x (0.065-0.65). **Tinct. iodi**. Dose m v-xv (0.32-1.0). It is chiefly used locally. **Unguentum iodi**, is used locally as an absorbent. **Vapor iodi** (B. P.), used for inhalation.

Iodism (*i'-o-dizm*) [*ιώδης*, violet colored, from *iov*, a violet; *eidōs*, appearance]. A condition arising from the prolonged use of iodine or iodine-compounds, marked by frontal headache, coryza, ptialism, and various skin-eruptions, especially acne; rarely by a cachexia with atrophy of the sexual organs and marked nervous symptoms.

Iodized (*i'-o-dizd*) [*ιώδης*, violet-colored, from *iov*, a violet; *eidōs*, appearance]. Impregnated with iodine.

Iodoform (*i'-o'-do-form*) [*iodin*; *forma*, form], CHI_3 . Triiodomethane. Formyl triiodid, a yellow, finely crystalline substance having a peculiar penetrating odor, and containing about 96.7 per cent. iodine by weight. It is readily soluble in chloroform and ether, less readily in alcohol, and but slightly in water. Iodoform is antiseptic and anesthetic, and is used as a dressing to wounds and syphilitic and chancroidal ulcers, either in powder or in the form of iodoform gauze. In tuberculous affections, when it can be directly introduced, it has yielded good results, being in such cases usually employed in the form of an emulsion in olive oil or as an ethereal solution. Internally it has been used as an alterative in goiter, rickets, pulmonary tuberculosis, and syphilis. Dose gr. j-ij (0.065-0.20). The use of large quantities locally has led to the production of toxic symptoms, which resemble those of meningitis, and to fatty degeneration of the internal organs. **I.-gauze**, gauze impregnated with iodoform. **I., Suppositoria** (B. P.), are used after rectal operations and in fissure of the anus. **I., Unguentum** (U. S. P.), is used as a local antiseptic and stimulant.

Iodol (*i'-o-dol*) [*ιώδης*, violet-colored, from *iov*, a violet; *eidōs*, appearance], $\text{C}_4\text{I}_4\text{NH}$.

Tetraiodopyrrol, an odorless, grayish-brown powder, soluble in alcohol and in ether, and used as a substitute for iodoform, and also in the treatment of diabetes mellitus. Dose gr. ss-v (0.032-0.32). Unof.

Iodum (*i'-o'-dum*). Iodine.

Ioduret (*i'-o'-u-ret*) [*iodin*]. An iodid.

Ion (*i'-on*) [*ίων*, *ión*, going]. An element set free by electrolysis, and classified as an anion or kation, according as it is set free at the positive or negative pole.

Iotacism (*i'-o'-tas-izm*) [*ιώτα*, the letter I]. Inability to pronounce distinctly the proper sound of the letter *i*.

Ipecac, **Ipecacuanha** (*ip'-e-kak*, *ip'-e-kak-u-an'-ah*) [Brazil., *ipecanguen*]. Ipecac.

The root of *Cephaelis ipecacuanha*, a plant of the order Rubiaceæ, containing an alkaloid, emetin, $\text{C}_{28}\text{H}_{40}\text{N}_2\text{O}_8$, and ipecacuanbic acid. Ipecac in large doses is emetic, in small doses diaphoretic, expectorant, and in minute doses, a gastric stimulant. It is used as an emetic, especially in narcotic poisoning, and, in children, to dislodge membranes and secretions in croup and capillary bronchitis; as a diaphoretic in acute colds, as an expectorant in bronchitis, as a sedative in vomiting (in minute doses), and in dyspepsia as a stimulant. It is said to be a specific in tropical dysentery. Emetin is emetic, but irritant in large doses. **Emetin**. Dose as an emetic, gr. $\frac{1}{8}$ - $\frac{1}{4}$ (0.008-0.016). **I., Extractum**, Fld. Dose m v-xxx (0.32-2.0). **I. et morphinæ**, **Trochisci**. Dose j-vj. **I. et opii**, **Pulvis** (**Pulvis ipecacuanhæ comp.** B. P.), **Dover's powder**. Dose gr. ij-xv (0.13-1.0). **I. et opii**, **Tinct.** Dose m v-x (0.32-0.65). **I. cum scilla**, **Pilula** (B. P.). Dose gr. v-x (0.32-0.65). **I., Syrupus**. Dose as an emetic, m xxx-f 3 j (2.0-4.0) for a child; f 3 ss-f 3 j (16.0-32.0), for an adult; as an expectorant, m v-f 3 j (0.32-4.0). **I., Trochisci**, each contains about $\frac{1}{3}$ grain (0.021) of ipecac. **I., Vinum**. Dose m j-f 3 j (0.065-4.0).

Iridal (*i'-rid-al*) [*ίρις*, iris]. Relating to the iris.

Iridectomy (*ir-id-ek'-to-me*) [*ίρις*, iris; *ἐκτομή*, excision]. The cutting out of a part of the iris.

Iridenkleisis, or **Iridenkleisis** (*ir-id-en-kli'-sis*) [*ίρις*, iris; *ἐγκλείν*, to lock in]. See *Iridodesis*.

Irideremia (*ir-id-er-e'-me-ah*) [*ίρις*, iris; *ἰσχυρία*, lack]. Absence of one or both irides.

Iridescence (*ir-id-es'-ens*) [*iridescere*, to shine with rainbow-colors]. The property of breaking up light into the spectral colors.

Iridesia (*ir-id-e'-sis*). See *Iridodesis*.

Iridochoroiditis (*ir-id-o-ka-roid-i'-tis*) [*ίρις*, iris; *χόριον* chorion; *εἶδος*, likeness; *ῖτις*, itis].

inflammation]. Inflammation of both the iris and the choroid of the eye.

Iridocyclitis (*ir-id-o-sik-li'-tis*) [*ἰρις*, iris; *κύκλος*, a circle; *τις*, inflammation]. Inflammation of the iris and the ciliary body.

Iridodesis (*ir-id-od'-es-is*) [*ἰρις*, iris; *δέσις*, a binding together]. An operation for the purpose of altering the position of the pupil, by drawing the iris into one or two small openings in the cornea and preventing its return by a loop of silk placed around it.

Iridodialysis (*ir-id-o-di-al'-is-is*). See *Core-dialysis*.

Iridodonesis (*ir-id-o-do-ne'-sis*) [*ἰρις*, iris; *δόνεισις*, a trembling]. Tremulousness of the iris; hippus.

Iridoplegia (*ir-id-o-plé'-je-ah*) [*ἰρις*, iris; *πληγή*, stroke]. Paralysis of the sphincter of the iris.

Iridotomy (*ir-id-ot'-o-me*) [*ἰρις*, iris; *τομή*, section]. An incision into the iris.

Iris (*i'-ris*) [*ἰρις*, a halo or rainbow]. 1. A colored circular membrane, placed between the cornea and the lens, and having a central perforation, the pupil. It is about half an inch in breadth, and consists principally of two sets of unstriped muscular fibers, the sphincter of the iris, or *sphincter pupillæ*, a narrow zone of circular fibers surrounding the pupil, and the dilator of the iris, or *dilator pupillæ*, a radiate band of fibers extending from the pupil to the border of the iris. 1.-**contraction**. See *Reflexes*, *Table of*. 2. Iris, or Blue flag, a plant of the natural order Iridæ. The rhizome of *Iris versicolor* (Iris, U. S. P.) is cathartic, emetic, and diuretic. Dose gr. x-xx (0.65-1.3). **Extractum iridis**. Dose gr. i-ij (0.065-0.13). **Extractum iridis fluidum**. Dose ℥v-x (0.32-0.65). 1., **Florentine**, Orris root, the root of *Iris florentina*, emetocathartic and diuretic. At present it is used chiefly as an ingredient of tooth powders.

Irish Moss. See *Chondrus*.

Iritic (*i-rit'-ik*) [*ἰρις*, iris; *τις*, inflammation]. Of the nature of, pertaining to, or affected with, iritis.

Iritis (*i-rit'-tis*) [*ἰρις*, iris; *τις*, inflammation]. Inflammation of the iris.

Iritomy (*i-rit'-o-me*). See *Iridotomy*.

Iron (*i'-ern*) [AS., *iren*, iron]. See *Ferrum*. **Irradiation** (*ir-ra-de-a'-shun*) [*in*, into; *radiare*, to radiate]. A phenomenon in which, owing to the difference in the illumination of the field of vision, or its background, objects appear much larger than they really are.

Irreducible (*ir-re-du'-se-bl*) [*in*, not; *reducere*, to lead back]. Not reducible; not capable of being replaced in a normal position, as an I. hernia.

Irregular (*ir-reg'-u-lar*) [*in*, not; *regulus*, rule]. Not regular; not normal or accord-

ing to rule; not rhythmic; not recurring at proper intervals, as an I. pulse.

Irrespirable (*ir-res'-pir-a-bl*) [*in*, not; *respirare*, to breathe]. Not capable of being breathed.

Irrigation (*ir-ig-a'-shun*) [*irrigare*, to lead water to]. The act of washing out by a stream of water, as I. of the bladder. 1., **Continuous**, the continuous passage of a stream of water over a surface, in order to reduce or limit inflammation.

Irrigator (*ir'-ig-a-tor*) [*irrigare*, to lead water to]. An apparatus for performing irrigation.

Irritability (*ir-it-ab-il'-it-e*) [*irritare*, to irritate, to tease]. 1. The state of being irritable, or of responding to stimuli. 2. A functional disturbance of a part on account of which it reacts excessively to slight stimulation, as I. of the bladder, a condition in which the urine is voided in small quantities at short intervals. 1., **Faradic**, the state in which the faradic current will cause muscular contraction. 1., **Galvanic**, the state in which the galvanic current will cause muscular contraction. 1., **Muscular**, the inherent contractile quality of a muscle. 1., **Nervous**, the property of a nerve to transmit impulses upon stimulation.

Irritable (*ir'-it-a-bl*) [*irritare*, to irritate, to tease]. 1. Reacting to stimuli. 2. Easily excited. 1. **Bladder**, a condition of the bladder marked by constant desire to void urine. 1. **Heart**, a peculiar condition of the heart characterized by precordial pain, dyspnea on exertion, palpitation, and irregularity of the heart's action.

Irritant (*ir'-it-ant*) [*irritare*, to excite]. 1. Causing or giving rise to irritation. 2. An agent that induces irritation.

Irritation (*ir-it-a'-shun*) [*irritare*, to excite]. 1. A condition of undue excitement. 2. The act of irritating or stimulating. 3. The stimulus necessary to the performance of a function.

Irritative (*ir'-it-a-tiv*) [*irritare*, to excite]. Characterized by or dependent on irritation. 1. **Fever**, a febrile condition dependent upon the presence in the body of irritating substances.

Ischemia (*is-ke'-me-ah*) [*ἰσχεῖν*, to check; *αἷμα*, blood]. Local anemia.

Ischemic (*is-ke-m'-ik*) [*ἰσχεῖν*, to check; *αἷμα*, blood]. Affected with or relating to ischemia.

Ischialgia (*is-ke-al'-je-ah*) [*ἰσχίον*, hip; *ἀλγος*, pain]. Sciatica; neuralgia of the hip.

Ischiatic (*is-ke-al'-ik*) [*ἰσχίον*, hip]. Pertaining to the ischium.

Ischidrosis (*is-kid-ro'-sis*) [*ἰσχεῖν*, to suppress; *ἰδρώς*, sweat]. Suppression of sweat.

Ischio- (*is'-ke-o-*) [*ἰσχίον*, hip]. A prefix indicating relationship to the ischium, or the hip.

Ischioanal (*is-ke-o-a'-nal*) [*ischium*; *anus*]. Pertaining to the ischium and anus.

Ischiobulbar (*is-ke-o-bul'-bar*) [*ischium*; *βάλλος*, a bulb]. Pertaining to the ischium and the bulb of the urethra.

Ischiocapsular (*is-ke-o-cap'-su-lar*) [*ischium*; *capsula*, a capsule]. Pertaining to the ischium and the capsular ligament of the hip.

Ischiocavernosus (*is-ke-o-kav-er-nos'-sus*) [*ischium*; *caverna*, cavern]. The erector penis (or erector clitoridis). The word *Musculus* is understood.

Ischiocele (*is'-ke-o-sel*) [*ischion*, hip; *κήλη*, tumor]. Hernia through the sciatic notch.

Ischiofemoral (*is-ke-o-fem'-o-ral*) [*ischium*; *femur*, femur]. Pertaining to the ischium and the femur.

Ischioneuralgia (*is-ke-o-nu-ral'-je-ah*) [*ischium*; *νεῦρον*, nerve; *ἄλγος*, pain]. Sciatica.

Ischioperineal (*is-ke-o-per-in-e'-al*) [*ischium*; *περίναιον*, perineum]. Pertaining to both ischium and perineum; pertaining to the space between the anus and the scrotum.

Ischiorectal (*is-ke-o-rect'-tal*) [*ischion*, ischium; *rectus*, straight]. Pertaining to both ischium and rectum. **1. Abscess**, an inflammation of the areolar tissue of the ischiorectal fossa. **1. Fossa**, a deep fossa filled with loose areolar tissue situated between the rectum and the ischium.

Ischium (*is'-ke-um*) [*ischion*, hip]. The inferior part of the os innominatum; the bone upon which the body rests in sitting.

Ischomenia (*is-ko-mel'-ne-ah*) [*ischion*, to suppress; *μήν*, month, menses]. Suppression of the menstrual flow.

Ischuretic (*is-ku-ret'-ik*) [*ischion*, to suppress; *urina*, urine]. **1.** Relating to or relieving ischuria. **2.** A remedy or agent that relieves retention or suppression of urine.

Ischuria (*is-ku'-re-ah*) [*ischion*, to suppress; *urina*, urine]. Retention or suppression of urine.

Isinglass (*i'-zing-glas*). See *Ichthyocolla*. **1., Vegetable.** See *Agar-agar*.

Island of Reil (*i'-land; ril*) [J. C. Reil, a German anatomist]. A group of five or six small convolutions (*gyri operi*) situated at the bottom of the fissure of Sylvius.

Iso- (*i'-so-*) [*isos*, equal]. A prefix signifying equal, or, in chemic nomenclature, isomeric.

Isoamylamin (*i-so-am-il'-am-in*) [*isos*, equal; *amylum*, starch; *amin*]. A ptomain obtained in the distillation of horn with potassic hydrate; it also occurs in the putrefaction of yeast.

Isochromatic (*i-so-kro-mat'-ik*) [*isos*, equal; *χρῶμα*, color]. Having the same color throughout.

Isochronous (*i-sak'-ro-nus*) [*isos*, equal; *χρόνος*, time]. Occurring at or occupying equal intervals of time.

Isocoria (*i-so-ko'-re-ah*) [*isos*, equal; *κόρη*,

pupil]. Equality in diameter of the two pupils.

Isodiametric (*i-so-di-am-et'-rik*) [*isos*, equal; *διά*, through; *μέτρον*, measure]. Having equal diameters.

Isodynamic (*i-so-di-nam'-ik*) [*isos*, equal; *δυναμς*, force]. Having or generating equal amounts of force. **1. Foods**, those that produce an equal amount of heat in undergoing the chemic changes of digestion.

Isoelectric (*i-so-e-lek'-trik*) [*isos*, equal; *ἡλεκτρον*, amber]. Having the same electric properties throughout.

Isolate (*is'-o-lat* or *i'-so-lat*) [*isola*, from *insula*, an island]. To separate; to place apart.

Isomer (*i'-som-er*) [*isos*, equal; *μέρος*, a part]. An isomeric body. See *Isomeric*.

Isomeric (*i-so-mer'-ik*) [*isos*, equal; *μέρος*, a part]. Of a chemic substance, composed of the same elements united in the same proportions by weight; in a restricted sense, composed of the same elements and having the same molecular weight as another substance.

Isomerism (*i-som'-er-izm*) [*isos*, equal; *μέρος*, a part]. The quality of being isomeric. **1.** is of two kinds—(a) substances may have the same percentage-composition and the same molecular weights; these are termed *metameric*; (b) they may have the same percentage-composition, but different molecular weights; these are termed *polymeric*. Ammoniumcyanate, CON.NH_4 , and urea, CON_2H_4 , are metameric; acetylene, C_2H_2 , benzene, C_6H_6 , and styrene, C_8H_8 , are polymeric. **1., Physical**, the form in which bodies that are isomeric and not differing chemically present different physical properties, such as their action toward polarized light.

Isometric (*i-so-met'-rik*) [*isos*, equal; *μέτρον*, measure]. Of the same dimensions. **1. Muscular Act**, the preservation of the length of a muscle when stimulated, the muscle only undergoing a change in tension.

Isometropia (*i-so-met-ro'-pe-ah*) [*isos*, equal; *μέτρον*, measure; *ὤψ*, eye]. Equality of kind and degree in the refraction of the two eyes.

Isomorphie (*i-so-mor'-fik*) [*isos*, equal; *μορφή*, form]. Having the same form; of crystals, crystallizing in the same form.

Isomorphism (*i-so-mor'-fizim*) [*isos*, equal; *μορφή*, form]. Similarity in crystalline form.

Isomorphous (*i-so-mor'-fus*). See *Isomorphie*.

Isopathy (*i-sop'-ath-e*) [*isos*, equal; *πάθος*, suffering]. The treatment of disease by the administration of the causative agent or of its products, as the treatment of smallpox by the administration of variolous matter.

Isopelletierin (*i-so-pel-et'-e-er-in*) [*isos*, equal; *pelletierin*]. See *Pelletierin*.

Isopepsin (*i-so-pep'-sin*) [*isos*, equal; *πέψις*, digestion]. **1.** A body formed by heating pepsin to a point between 104° and 140° F.

(40°-60° C.). It changes albumin into para-peptone. 2. Same as *Para-peptone*.

Isophoria (*i-so-fō'-re-ah*) [ἴσος, equal; φόρος, a tending]. A condition in which the eyes lie in the same horizontal plane, the tension of the vertical muscles of each eye being equal, and the visual lines lying in the same plane.

Isothermal (*i-so-ther'-mal*) [ἴσος, equal; θερμῆ, heat]. Of equal or uniform temperature. 1. Lines, lines drawn through places having the same average temperature for a given period of time.

Isotropic, Isotropous (*i-so-trop'-ik, i-sot'-ro-pus*) [ἴσος, equal; τροπή, turning]. 1. Having the same shape and appearance, from whatever point observed. 2. Being singly-refractive.

Issue (*ish'-u*) [Fr., *issue*, from *exire*, to go out]. 1. An ulcer or fistulous passage made and kept up artificially for purposes of counterirritation. 1.-*pea*, a pea-shaped foreign body (as of ivy-wood or orris-root), inserted into an issue to keep up suppuration. 2. Offspring.

Isthmus (*is'-mus*) [ἰσθμός, isthmus]. A narrow, contracted part uniting two larger parts of an organ. 1. of the Fauces, the space between the arches of the palate. 1. of the Thyroid Gland, the narrow part connecting the lobes of the thyroid body.

Italian (*it-al'-yan*) [*Italus*, an Italian]. Of or pertaining to Italy. 1. **Leprosy**. See

Pellagra. 1. **Rhinoplasty**. See *Operation, Taghiacotian*.

Itch (*ich*) [AS., *giccan*, to itch]. 1. An irritating sensation in the skin. 2. A name for various skin-diseases accompanied by itching, particularly scabies. 1., **Barbers'**. See *Tinea sycosis*. 1.-*mite*. See *Acarus scabiei*. -*ite* (*it*). 1. A suffix employed in mineralogy to denote a mineral, or of mineral origin. 2. A suffix employed in chemistry for the salt of an acid that has the suffix -ous.

Iter (*i'-ter*) [*iter*, a journey]. A passage-way. 1. **ad infundibulum**, the passage between the third ventricle of the brain and the infundibulum. 1. **a tertio ad quantum ventriculorum**, the aqueduct of Sylvius, extending from the third ventricle to the fourth. 1. **chordæ anterie**, the aperture through which the chorda tympani nerve leaves the tympanum. 1. **chordæ posterie**, the aperture through which the chorda tympani nerve enters the tympanum. -*itis* (*i'-tis*) [ἰτις, inflammation]. A suffix used to denote inflammation.

Ivory (*i'-vor-e*) [*eboreus*, made of ivory, from *ebur*, ivory]. The hard bone-like substance chiefly obtained from the tusks of elephants. 1.-**black**, animal charcoal. 1., **Dental**, dentine.

Ixodes (*iks'-o'-dex*) [ἰξός, bird-lime; εἶδος, form]. A genus of the order Acaridea, including most of the parasitic ticks.

J

J. Symbol for Joule's equivalent.

Jaborandi (*jab-or-an'-de*) [Braz.]. See *Pilocarpus*.

Jacaranda (*jak-ar-an'-dah*). A genus of bignoniaceous plants of tropical America, several species of which are employed in syphilis in Brazil. **J. caroba**, is antisyphilitic, and is of service in the treatment of urethritis, rheumatism, and skin-diseases. Dose of the fl. ext., gtt. xvj-f3j (1.0-4.0). **J. lancifoliata**, is used by the natives of Brazil in urethritis. Dose of a 1-8 tincture mxxv (1.0); of the fl. ext., gtt. xvj-xxx (1.0-2.0). Unof.

Jacksonian Epilepsy. See *Epilepsy*.

Jacob's Membrane [*Jacob*, an Irish physician]. The layer of rods and cones of the retina. **J.'s Ulcer**. See *Rodent Ulcer*, and *Diseases, Table of*.

Jacobson's Nerve [L. I., *Jacobson*, a Danish anatomist]. The tympanic branch of the glossopharyngeal nerve. **J.'s Organ**, a short, rudimentary canal, extending along the septum of Stensen's duct, and ending in a culdesac.

Jactitation (*jak-tit'-a'-shun*) [*jactitare*, to pour forth]. A tossing about, a condition at times present in grave diseases.

Jadelot's Lines or Furrows [*Jadelot*, a French physician]. Certain furrows of the face observed in conditions of disease. Three sets are distinguished: The **genal** and **nasal** furrows are said to indicate disease of the gastrointestinal tract or abdominal viscera; the former runs from the mouth toward the malar bone, the latter from the nasal alae in a semicircle about the mouth; the **labial** furrow, from the angle of the mouth out-

ward to the lower part of the face, indicates disease of the lungs; the oculozygomatic furrow, beginning at the inner canthus of the eye, and passing outward below the lower lid, to be lost on the cheek; it is said to point to disorders of the nervous system.

Jail-fever. Typhus fever.

v. Jaksch's Disease. Pseudoleukocythemia of infants.

Jalap (*jal'-ap*) [from *Jalapa*, a city of Mexico]. The tuberous root of *Ipomoea jalapa* (U. S. P.), of *Ipomoea purga* (Exogonium purga) (B. P.), a plant of the natural order Convolvulaceae. Its active principle is a resin (*Resina jalape*, U. S. P.), which contains a glucosid, convolvulin, $C_{31}H_{50}O_{16}$. Jalap is an active hydragogue cathartic, and is used to remove dropsical effusions by the bowel. Combined with calomel it is a favorite remedy in bilious fever. Dose of powdered jalap gr. xv-xxx (1.0-2.0). **J., Extractum.** Dose gr. iv-viii (0.26-0.52). **J., Pulvis, Comp.** Dose gr. x-5j (0.65-4.0). **J., Resina.** Dose gr. iv-viii (0.26-0.52).

Jalapin (*jal'-ap-in*) [from *Jalapa*, a city of Mexico]. A purgative glucosid from *Ipomoea* or *Convolvulus orizabensis*.

Jamaica Dogwood. See *Piscidia*.

Janiceps (*jan'-is-eps*) [*Janus*, a two faced divinity; *caput*, head]. A syncephalic monstrosity with two faces.

Jarjavay's Muscle. The depressor urethrae.

Jasmine (*jas'-min*) [Pers., *yasmîn*, jasmine]. See *Gelsemium*.

Jatropha (*jal'-ro-fah*) [*iatriphs*, a physician; τροφή, nourishment]. A genus of euphorbiaceous plants. **J. curcas**, is the source of purging nuts. **J. manihot**, yields tapioca.

Jaundice (*jawn'-dis*) [Fr., *jaunisse*, from *jaune*, yellow]. A yellow discoloration of the skin, mucous membranes, and secretions, due to the presence of bile-pigments in the blood. See *Icterus*. **J., Catarrhal**, that due to swelling of the bile-ducts from catarrh. **J., Hematogenous**, that form due to excessive destruction of blood-corpuscles. **J., Hepatogenous**, that due to obstruction to the flow of bile from the liver. By some all forms of jaundice are considered hepatogenous, since bile is made only in the liver. **J., Malignant**, acute yellow atrophy of the liver. See *Icterus gravis*. **J. of the New-born.** See *Icterus neonatorum*.

Jaw [AS., *crāwan*, to chew]. 1. Either of the two parts of the face (upper or lower jaw) serving the purpose of seizing or masticating the food. 2. Also the bone (jaw-bone or jaw) that forms the framework of the jaw. **J.-jerk**, **J.-clonus**, a reflex contraction of the muscles of mastication produced by suddenly depressing the lower jaw. See *Reflexes*, Table of. **J., Lock**, or **Locked.** See

Trismus. **J., Lumpy**, actinomycosis of cattle.

Jecur (*jel'-ker*) [L.]. The liver.

Jejunal (*jel'-u-nal*) [*jejunus*, empty]. Pertaining to the jejunum.

Jejunocolostomy (*jel'-u-no-ko-los'-to-me*) [*jejunus*, empty; κόλον, colon; στόμα, mouth]. The formation of an artificial passage between the jejunum and the colon.

Jejunoleostomy (*je'-ju-no-il-e-ost'-to-me*) [*jejunus*, empty; ileum; στόμα, mouth]. The formation of an artificial communication between the jejunum and the ileum.

Jejunostomy (*je'-u-nos'-to-me*) [*jejunus*, empty; στόμα, mouth]. The making of an artificial opening through the abdominal wall into the jejunum.

Jejunum (*je'-u-num*) [*jejunus*, empty, because usually found empty after death]. The second division of the small intestine extending between the duodenum and the ileum, and measuring about eight feet (2.2 meters) in length.

Jelly (*jel'-e*) [*gelare*, to freeze]. A soft, gelatinous, tremulous substance. **J., Wharton's**, the gelatinous mucoid connective tissue investing the umbilical cord.

Jennerian (*jen'-e-re-an*) [after Edward Jenner, an English physician]. Pertaining to Edward Jenner, the discoverer of vaccination against smallpox.

Jequirity (*je'-kwir'-it-e*). See *Abrus*.

Jervin (*jer'-vin*). See *Veratrum*.

Jesuits' Bark. Cinchona.

Jigger Flea. See *Pulex*.

Jimson-weed (*jim'-sun-wed*). See *Stramonium*.

Joint (*juingere*, to join). See *Articulation*.

J., Charcot's. See *Diseases*, Table of.

Joule (*joel*) [after J. P. Joule, an English physicist]. 1. A unit of electric energy, equivalent to the work expended when a current of one ampere flows for one second against a resistance of one ohm. 2. A small calorie—the amount of heat required to raise the gram of water 1° C.

Joule's Equivalent (abbreviated J.) (*joelz*) [after J. P. Joule, an English physicist]. The mechanic equivalent of heat or the amount of work that converted into heat will raise the temperature of 1 lb. of water 1° F. It is equivalent to 772 foot-pounds.

Jugal (*ju'-gal*) [*jugum*, a yoke]. Connecting or uniting, as by a yoke. **J. Bone**, the malar bone. **J. Process**, the zygomatic process.

Juglans (*ju'-glans*) [L., walnut]. Butternut. The bark of the root of *J. cinerea*, of the natural order Juglandaceae. It is a mild cathartic, and has also been used in intermittent and remittent fever. Dose of the extract, gr. xx-xxx (1.3-2.0).

Jugular (*ju'gu-lar*) [*jugulum*, throat]. Pertaining to the throat. **J. Foramen**. See *Foramina*, *Table of*. **J. Fossa**, a notch in the posterior border of the petrous portion of the temporal bone, which, with a similar notch in the occipital bone, forms the foramen lacerum posterius. **J. Ganglion**, the superior ganglion of the glossopharyngeal nerve. **J. Process**, a rough process external to the condyle of the occipital bone. **J. Veins**, the internal J. vein collects the blood from the brain, part of the face and neck, and unites with the subclavian vein to form the vena innominata; the external J. vein carries the blood from the exterior of the cranium and parts of the face and empties into the subclavian vein.

Juice (*jūs*) [*jus*, broth]. 1. The liquid contained in vegetal or animal tissues. 2. Any of the secretions of the body, as the intestinal or pancreatic juice.

Jujube (*ju'-jūb*) [*jujuba*, fruit]. The fruit of the jujube tree, *Zizyphus jujuba*. **J.-paste**, a paste containing the pulp of jujubes, and used in pulmonary disorders. It is now made of gum-arabic, or of gelatin, variously flavored.

Julep (*ju'-lep*) [Pers., *jālab*, a sweet drink]. A sweetened drink containing aromatic or medicinal substances.

Jumpers (*ju'm'pers*) [Scand., *gumpa*, to jump]. Persons afflicted with a peculiar neurosis by reason of which they do whatever they are told, and perform sudden leaping or jumping movements.

Juniperus (*ju-nip'us*). 1. A genus of coniferous trees. 2. Juniperus, the fruit or berry of *J. communis*, containing a volatile oil, oleum juniperi, and an amorphous substance, juniperin. *J.* is a stimulant to the genitourinary mucous membrane, and is used

in nephritis, pyelitis, and cystitis. Dose of the oil, $\text{m} \cdot \text{j} \cdot \text{iv}$ (0.065-0.26). **Spiritus juniperi**, dose $\text{m} \cdot \text{xxx} \cdot \text{f} \cdot \text{z} \cdot \text{j}$ (2.0-4.0). **Spiritus juniperi compositus**, is the pharmacopeal representative of the beverage gin; dose $\text{f} \cdot \text{z} \cdot \text{j} \cdot \text{iv}$ (4.0-16.0). *J. sabina*, yields savine (Sabina, U. S. P.). *J. virginiana*, red cedar, the tops of which are used as a substitute for savine.

Junket (*junk'-et*) [*iuncus*, a rush]. "Curds and whey," prepared by coagulating milk with rennet.

Junod's Boot (*ju'-nōr*). A boot-shaped case, usually of stiff leather, made to enclose the leg so that, the air being exhausted, the blood rushes to the enclosed part. It has been employed to relieve inflammation and congestion of the viscera.

Jurisprudence (*ju-ris-pru'-dens*) [*jus*, law; *prudētia*, skill]. The science of the interpretation and application of the law. **J., Medical**, the application of medical knowledge to the principles of common law.

Jury-mast (*ju'-re-mast*) [*jury*, from Dan. *kiōre*, a driving; AS., *mast*, mast]. A steel shaft with curved iron rods attached, employed to support the head in disease of the upper vertebrae.

Justomajor Pelvis (*jus'-to-ma'jor*). See *Pelvis*.

Justomajor Pelvis (*jus'-to-mi'nor*). See *Pelvis*.

Jute (*jūt*) [Beng., *jūt*, matted hair]. The bast fiber of several species of the genus *Corchorus*, grown chiefly in India and Ceylon. Jute is used as an absorbent dressing.

Juxtaposition (*juk'-ta-po-zish'-un*) [*juxta*, near; *positio*, position]. Situation adjacent to another; the act of placing near; apposition.

K

K. The symbol of Potassium (kalium).

K., or Ka. The abbreviation of Kathode, or of Kathodic.

Kairin (*kā'-rin*) [*καίρος*, the right time], $\text{C}_{10}\text{H}_{13}\text{NO} \cdot \text{HCl} \cdot \text{H}_2\text{O}$. The hydrochlorate of oxychinolin-ethyl, is antipyretic, diaphoretic, and emetic, and has been used as a substitute for quinin. Dose gr. v-xv (0.32-1.0).

Kairolin (*kā'-ro-lin*) [*καίρος*, the right time], $\text{C}_{10}\text{H}_{13}\text{N}$. Methylquinolin hydrid. An antipyretic resembling kairin, but less efficient.

Kakké (*kak'-ka*) [Chinese for "leg-disease"]. Epidemic and endemic multiple neuritis, or beriberi.

Kakosmia (*kak-os'-me-ah*). See *Cacosmia*.

Kali (*kā'-li*) [Ar., *qālī*, potash]. Potash.

Kalimeter (*ka-lim'-et-er*). See *Alkalimeter*.

Kalium (*kā'-le-um*). Potassium.

Kamala (*kam'-a'-lah*) [Hind., *kamlā*]. Rottlera. The glands and hairs from the capsules of *Mallotus philippinensis* (Rottlera tinctoria), native to Southern Asia and Abyssinia.

sinia. It is purgative and anthelmintic, and is used for the expulsion of lumbricoid worms and tapeworms. Dose \mathfrak{J} j-ijj (4.0-12.0).

Kangaroo. A marsupial mammal of Australia and the neighboring islands. **K. Tendon,** a tendon derived from the tail of the K. and used for surgical ligatures.

Kaolin (*ka'-o-lin*) [Chin., *kaoling*, "high ridge"]. White clay, China-clay. The silicate of aluminum, obtained from the decomposition of felspar. It is sometimes used as a protective application in eczema and as a coating for pills.

Kaposi's Disease. See *Atrophoderma pigmentosum*, and *Xeroderma pigmentosum*.

Karyokinesis (*kar-e-o-kin-e'-sis*) [*kárvov*, a nut (nucleus); *κίνησις*, movement, change]. Indirect cell-division, the common mode of reproduction of cells. It depends upon complicated changes in the mitome of the cell-nucleus that may be divided into the following steps: 1. The nucleus becomes larger; the mitome-filaments thicken and form a close skein, or *spirem*. 2. The fibrils become less convoluted and more widely separated, forming the loose skein; at the same time the nuclear spindles, two cone-shaped striated bodies, appear in the achromatin. 3. The mitome-fibrils split longitudinally. 4. The segments migrate toward the poles of the new nuclei, constituting daughter-wreaths, or *asters*. 5. Transformation of asters into fully-developed nuclei. 6. Division of the cell-protoplasm.

Karyokinetic (*kar-e-o-kin-e'-ik*) [*kárvov*, nucleus; *κίνησις*, motion]. Pertaining to karyokinesis, as K. figures, the forms assumed by the mitome in karyokinesis.

Karyolysis (*kar-e-ol'-is-is*) [*kárvov*, nucleus; *λύνω*, to loose]. The segmentation of the nucleus of the cell.

Karyomitome (*kar-e-om'-it-ōm*) [*kárvov*, nucleus; *μίτος*, thread]. The mitome-threads of the nucleus.

Karyomitosis (*kar-e-o-mit-ō'-sis*) [*kárvov*, nucleus; *μίτος*, a thread]. Karyokinesis.

Karyomitotic (*kar-e-o-mit-ō'-ik*) [*kárvov*, nucleus; *μίτος*, a thread]. Relating to karyomitosis.

Karyoplasm (*kar'-e-o-plazm*) [*kárvov*, a nut, kernel; *πλάσσειν*, to form]. The nuclear substance of a cell.

Kata- (*kat'-ah-*). For words thus beginning, and not found under K, see *Cata-*.

Katabolic (*kat-ab-ol'-ik*). See *Catabolic*.

Kath-, For words thus beginning see *Cath-*.

Kation (*kat'-e-on*). See *Cation*.

Kava, or Kava-kava (*kah'-vah*) [Hawaiian].

1. An intoxicating beverage prepared in the Sandwich Islands from the root of *Piper methysticum*. 2. The root of *Piper methysticum*, containing a resin, *kawin*, and an

alkaloid, *kavain*. The resin is a motor depressant, locally at first an irritant, later an anesthetic; it is also a cardiac stimulant. Kava-root has been used in gonorrhea, leukorrhea, and incontinence of urine. Dose of fluid extract $\mathfrak{m}\mathfrak{xv}$ - $\mathfrak{f}\mathfrak{J}$ (1.0-4.0).

Kefyr (*kef'-ir*) [Caucasian]. A nutritious substance obtained by a peculiar fermentation of cow's milk produced by certain fungi.

K.-seed, a substance containing the ferment (*Bacillus caucasicus*) of kefir. It is used in preparing the genuine kefir.

Kelectome (*ke'-lek-tōm*) [*κλήη*, a tumor; *ἐκ*, out; *τεμνειν*, to cut]. A cutting instrument introduced into a tumor, by means of a cannula, in order to obtain a part of the substance for examination.

Kelis (*ke'-lis*) [*κηλὶς*, scar]. Keloid.

Keloid (*ke'-loid*) [from *κηλὶς*, a scar, or *χηλὴ*, a claw; *ειδος*, likeness]. Cheloid; Alibert's keloid; kelis. A tumor-like fibrous outgrowth, usually occurring at the site of a scar. It is elevated, whitish or pink in color, and sends prolongations into the surrounding tissues resembling the claws of a crab. By many it is not considered a true tumor, but merely a hyperplastic scar. It affects the colored race more frequently than the white.

K. of Addison, morphea.

Kelotomy (*ke-lot'-o-mē*). Herniotomy.

Keratectasia (*ker-at-ek-ta'-se-ah*) [*κέρας*, horn, cornea; *ἐκτασις*, extension]. A bulging forward of the cornea.

Keratin (*ker'-at-in*) [*κέρας*, horn]. The basis of horny epithelium, hair, nails, feathers, etc. It contains sulphur, and on decomposition yields leucin and tyrosin.

Keratitis (*ker-at-ī'-tis*) [*κέρας*, cornea; *τις*, inflammation]. Inflammation of the cornea.

K. bullosa, the formation of large or small blebs upon the cornea of an eye, the seat of iridocyclitis, interstitial keratitis, or glaucoma.

K., Interstitial, a form of keratitis in which the entire cornea is invested with a diffuse haziness, almost completely hiding the iris. The surface of the cornea presents a ground-glass appearance. Later, from ciliary injection, blood-vessels form in the superficial layers of the cornea, and produce a dull-red color, the "salmon patch" of Hutchinson. The entire cornea may become cherry-red. The disease is most frequent between the ages of five and fifteen, and occurs in syphilitic individuals. **K. neuroparalytica**, keratitis following lesion of the trifacial nerve. Its cause is loss of trophic influence, aided by mechanic irritation and drying of the cornea. **K., Phlyctenular**, a variety characterized by the formation of small papules or pustules, often associated with similar lesions upon the conjunctiva. It is marked by much local congestion, lacrimation, and intense photophobia.

K. punctata, a secondary affection of the cornea in association with affections of the iris, choroid, and vitreous. It is characterized by the formation of opaque dots, generally arranged in a triangular manner upon the posterior elastic lamina of the cornea. It is sometimes designated as descemetitis. **K. purulenta**, that accompanied by the formation of pus. **K., Trachomatous**. See *Pannus*. **K., Traumatic**, that consequent upon wounds or other injury of the cornea.

Keratocele (*ker'-at-o-sel'*) [*κέρας*, cornea; *κῆλη*, tumor]. A hernia of Descemet's membrane through the cornea.

Keratoconus (*ker'-at-o-ko'-nus*) [*κέρας*, cornea; *κῶνος*, cone]. A conical protrusion of the cornea.

Keratogenous (*ker'-at-oj'-en-us*) [*κέρας*, horn; *γενᾶν*, to beget]. Producing a horny or horn-like substance.

Keratoglobus (*ker'-at-o-glo'-bus*) [*κέρας*, cornea; *globus*, a ball]. A globular protrusion of the cornea.

Keratoglossus (*ker'-at-o-glos'-us*) [*κέρας*, horn; *γλῶσσα*, tongue]. See *Muscles Table*.

Keratohyalin (*ker'-at-o-hi'-al-in*) [*κέρας*, horn; *ἵαλις*, glass]. A peculiar substance occurring in granules in the deeper layers of the skin.

Keratoid (*ker'-at-oid*) [*κέρας*, horn; *εἶδος*, like]. Hornlike.

Keratoiditis (*ker'-at-o-i-ri'-tis*) [*κέρας*, cornea; *ίρις*, iris; *ίτις*, inflammation]. Combined inflammation of the cornea and the iris.

Keratoma (*ker'-at-o'-mah*) [*κέρας*, horn; *μα*, tumor]. 1. See *Callositas*. 2. Congenital ichthyosis; the presence of horny plates upon the integument.

Keratomalacia (*ker'-at-o-mal'-se-ah*) [*κέρας*, cornea; *μαλακία*, softness]. A softening of the cornea.

Keratome (*ker'-at-ōm*) [*κέρας*, cornea; *τομή*, a cutting]. A knife with a peculiar trowel-like blade, used for making the incision into the cornea in the operation of iridectomy.

Keratometer (*ker'-at-om'-et-er*) [*κέρας*, cornea; *μέτρον*, measure]. An instrument for measuring the curves of the cornea.

Keratomycosis (*ker'-at-o-mi-ko'-sis*) [*κέρας*, cornea; *μύκης*, fungus]. A fungoid growth of the cornea.

Keratonyxis (*ker'-at-o-niks'-is*) [*κέρας*, cornea; *νύξις*, a pricking]. The needling of a soft cataract by puncture through the cornea; also, the old operation of couching a cataract with the needle.

Keratoplasty (*ker'-at-o-plas-te*) [*κέρας*, cornea; *πλάσσειν*, to form]. Plastic operation upon the cornea, especially the transplantation of a portion of cornea from the eye of a lower animal to that of man.

Keratoscope (*ker'-at-o-skōp*) [*κέρας*, cornea; *σκοπεῖν*, to observe]. An instrument for examining the cornea, and testing the symmetry of its meridians of curvature.

Keratotomy (*ker'-at-o-skōp*) [*κέρας*, cornea; *σκοπεῖν*, to observe]. 1. Examination of the cornea with the keratroscope. 2. Keratotomy, or skiascopy.

Keratosis (*ker'-at-d'-sis*) [*κέρας*, horn]. Any disease of the skin characterized by an overgrowth of the horny epithelium. **K. follicularis**. See *Darier's Disease*. **K. pilaris**, a chronic affection of the skin marked by hard, conical elevations investing the hair-follicles, and somewhat resembling gooseflesh. **K. senilis**, a cornification of the skin of old people, often limited to certain definite regions, as the face and dorsal surfaces of the hands and feet.

Keratotomy (*ker'-at-o-tōm*). See *Keratome*.

Keratotomy (*ker'-at-o'-o-me*) [*κέρας*, cornea; *τέμνειν*, to cut]. Incision of the cornea.

Keraunoneurosis (*ker'-au-no-nu-ro'-sis*) [*καραυνός*, lightning; *νεῖρον*, nerve; *νόσος*, disease]. Nervous disease due to lightning-stroke.

Kerion (*ke'-re-on*) [*κηρίον*, honey-comb]. See *Tinea kerion*.

Kerkring, Valves of. The valvulae conniventes of the small intestine.

Kermes (*ker'-mēs*) [Pers., *qirmis*, crimson]. A red dye-stuff resembling cochineal, made from the bodies of the dried insects, *Coccus ilicis*, found on the Kermes oak. **K. Mineral**, a mixture of the teroxid and tersulphid of antimony.

Ketone (*ke'-tōn*) [an arbitrary variation of *acetone*]. An organic compound consisting of the unsaturated radicle $=C=O$ united to two alcohol-radicles.

Kidney (*kid'-nē*) [ME., *kidnere*, from Icel., *kníðr*, the womb; *nýra*, kidney]. One of the two large glandular organs situated in the upper and posterior portion of the abdominal cavity, and concerned in the excretion of the urine. It consists of an outer cortical substance, and an inner medullary substance. The medulla consists of from 8 to 18 pyramids (pyramids of Malpighi), the apices of which, the papillae, project into the calices of the ureter. The pyramids are striated, and in places send narrow projections into the cortex, the medullary rays, or pyramids of Ferrein. Between the pyramids are extensions from the cortex, the columns of Bertini. The cortex, by the penetration into it of the medullary rays, is divided into medullary rays and the labyrinth. The secreting structure of the kidney consists of long tubes, beginning in an expanded extremity, the capsule of Bowman, which invests a tuft of blood-vessels, the glomerulus, and constitutes, together with this,

a Malpighian body; extending from this is the proximal convoluted tubule; then comes the spiral tubule, then the loop of Henle, consisting of a descending and an ascending limb; then the distal convoluted tubule, which terminates in the collecting tubule. The blood-vessels of the kidney divide into two sets of branches, one supplying the cortex, the other the medulla. The kidney weighs about 150 grams. **K.**, **Amyloid**, a kidney the seat of amyloid degeneration. **K.**, **Fatty**, one the seat of extensive fatty degeneration. **K.**, **Float-**ing, one susceptible of displacement over a considerable extent of the abdomen, a condition most frequent in women, and as a rule affecting the right side. **K.**, **Gouty**, or **K.**, **Granular**, the small kidney resulting from chronic interstitial nephritis. **K.**, **Horse-**shoe. See *Horseshoe-kidney*. **K.**, **Large White**, that of the advanced stage of chronic parenchymatous nephritis. **K.** of **Preg-**nancy, an anemic kidney with fatty infiltration of the epithelial cells, but without any acute or chronic inflammation, occurring in pregnant women. **K.**, **Pigback**, the large congested kidney found in alcoholic subjects. **K.**, **Red Contracted**. See *K.*, *Granular*. **K.**, **Small White**, the final stage of the large white kidney after loss of its substance from atrophy or degeneration. **K.**, **Surgical**, pyelonephritis. **K.**, **Waxy**. Same as *K.*, *Amyloid*.

Kilogram (*kil'-o-gram*) [*χιλιοι*, one thousand; *γράμμα*, an inscription]. One thousand grams, or 2.2 pounds avoirdupois.

Kiloliter (*kil'-o-le-ter*) [*χιλιοι*, thousand; *λίτρον*, a pound]. One thousand liters, or 35.31 cubic feet.

Kilometer (*kil'-o-me-ter*) [*χιλιοι*, thousand; *μέτρον*, measure]. One thousand meters, or 1093.6 yards.

Kilostere (*kil'-o-ster*) [*χιλιοι*, thousand; *στερεός*, solid]. One thousand cubic meters.

Kinæsthesia (*kin-es-the'-ze-ah*). See *Kin-esthesia*.

Kinematics (*kin-em-at'-iks*) [*κινέειν*, to move]. The science of motion.

Kinesiology (*kin-es-e-ol'-o-je*) [*κίνησις*, motion; *λόγος*, discourse]. The science of movements, considered especially as therapeutic or hygienic agencies.

Kinesiometer (*kin-es-e-om'-et-er*) [*κίνησις*, motion; *μέτρον*, measure]. An instrument for determining quantitatively the motion of a part.

Kinesioneurosis (*kin-es-e-o-nu-ro'-sis*) [*κίνησις*, movement; *νεῖρον*, a nerve; *νόσος*, disease]. A functional nervous disease associated with disorders of motion.

Kinesiotherapy (*kin-es-e-o-ther'-ap-e*) [*κίνησις*, movement; *θεραπεία*, treatment]. The

treatment of disease by systematic active or passive movements.

Kinesipathy (*kin-es-ipt'-ath-e*) [*κίνησις*, motion; *πάθος*, disease]. Kinesiotherapy.

Kinesodic (*kin-es-od'-ik*) [*κίνησις*, motion; *ὁδός*, way]. Pertaining to the motor pathways.

Kinesthesia (*kin-es-the'-ze-ah*), **Kinesthesia** (*kin-es-the'-sis*) [*κίνησις*, motion; *αἴσθησις*, sensation]. That quality of sensations whereby we become aware of our position in space, our movements, and that gives us our impression of weight and resistance.

Kinetic (*kin-et'-ik*) [*κινέειν*, to move]. Pertaining to motion; producing motion, as *K.* energy.

King's Evil. Scrofula, on account of a belief that it could be cured by the touch of the king. **King's Yellow**. See *Orpiment*.

Kino (*ki'-no*) [E. Ind.]. The inspissated juice of *Pterocarpus marsupium*, found in India, and similar in action to tannic acid; it is used mainly as a constituent of gargles and diarrhea-mixtures. **K.**, **Tinct.** Dose $\pi x - f 3 ij$ (0.65-8.0). **K.**, **Pulv., Comp.** (B.P.). Dose gr. v-xx (0.32-1.3). **K.-tannic Acid**, a variety of tannic acid found in kino.

Kinone (*kin'-on*). See *Quinone*.

Kissingen Salts (*kis'-ing-en*) [Ger.]. Effervescent salts from the mineral springs of Kissingen. **K. Water**, a laxative tonic mineral-water of Kissingen, in Bavaria.

Klatsch-preparation (*klatch'-prep-ar-a'-shun*) [Ger., *Klatschpräparat*]. A cover-glass preparation made by pressing the cover-glass lightly on a bacterial colony in plate-culture.

Kleptomania (*klep-to-ma'-ne-ah*) [*κλέπτειν*, to steal; *μανία*, madness]. A form of emotional insanity manifested by a morbid desire to commit theft.

Knee (*ne*) [A.S., *cnēd*, knee]. The articulation between the femur and the tibia. **K.-cap**, the patella. **K.**, **Housemaid's**. See *Housemaid's Knee*. **K.**, **In.-** See *Genu valgum*. **K.** of **Internal Capsule**, the angle of junction of the anterior and posterior limbs of the internal capsule. **K.-jerk**, **Patellar Tendon-reflex**, **K.-reflex**, or **K.-phenomenon**, a contraction of the quadriceps extensor femoris muscle as a result of a light blow on the patellar tendon. See also *Reflexes*, *Table of*. **K.-joint**, a hinge-joint consisting of the articulation of the condyles of the femur with the upper extremity of the tibia and the posterior surface of the patella. **K.**, **Knock-.** See *Genu valgum*. **K.**, **Out-.** See *Genu varum*. **K.-pan**, the patella.

Knife (*nif*) [A.S., *cnif*, knife]. An instrument for cutting. In surgery, knives are of various shapes and sizes, according to their use.

Knock-knee (*no-k' ne*). See *Genu valgum*.
Knuckle (*nuk'-l*) [ME., *knokil*, a knuckle or joint]. Any one of the articulations of the phalanges with the metacarpal bones or with each other.

Koch's Lymph [after Robert Koch, a German bacteriologist]. See *Tuberculin*. **K's Method of Sterilization**, a method of interrupted heating. The culture-media are heated for a short time daily for from three to five successive days, usually in the steam-sterilizer. **K's Rules**, rules formulated by Koch that must be complied with before a given microorganism can be accepted as the cause of an infectious disease: 1. The microorganism must be present in every case of the disease, and not in cases of other diseases. 2. It must be isolated and grown on artificial media. 3. Its cultures when injected into an animal-body must produce the disease in question. 4. The microorganism must again be found in the body in which the disease was thus produced.

Kolpo- (*kol'-po-*). See *Colpo-*.

Kombé (*komb'-ba*) [African]. An African arrow-poison (*kombé inée*) extracted from *Strophanthus kombé*.

Kopftetanus (*kôpf'-tel-an-us*) [Ger.]. Cephalic tetanus. See *Tetanus*.

Kopiopia (*kop-e-o'-pe-ah*). See *Copiopia*.

Koroscopy (*kor-ost'-ko-pe*). See *Retinoscopy*.

Kosin (*ko'-sin*) [Abyssinian, *cusso*], $C_{31}H_{50}O_{10}$. Same as *Koussin*. See *Brayera*.

Koumiss (*koo'-mis*) [Tartar, *kumis*, fermented mare's milk]. An alcoholic drink originally made by the fermentation of mare's milk. At present cow's milk is used in making it.

Koussin (*koo'-sin*). See *Brayera*.

Koussou (*koo'-o*). See *Brayera*.

Krameria (*kra-me'-re-ah*) [after J. G. H. Kramer, an Austrian physician]. A genus of polypetalous herbs. *Rhatany*, the root of *K. triandra* and *K. tomentosa*, shrubs native to South America, possessing the same astringent qualities as tannic acid. It is used in serous diarrheas. **K., Ext.** Dose gr. v-x (0.32-0.65). **K., Ext., Fld.** Dose η v- $\bar{\eta}$ ss (0.32-2.0). **K., Infus.** (B. P.)

Dose $\bar{\eta}$ j-ij (32.0-64.0). **K., Syr.**, contains of the fluid extract 35, syrup 65. Dose $\bar{\eta}$ ss- $\bar{\eta}$ ss (2.0-16.0). **K., Trochisci**, contain each gr. j (0.065) of the extract. **K., Tinct.** Dose η v- $\bar{\eta}$ j (0.32-4.0).

Kraurosis (*krau'-ro'-sis*) [*κραῦρος*, dry]. Shriveling and dryness of a part, especially of the vulva.

Krause's Corpuscles (*krau'-ses*). Terminal nerve-corpuscles described by W. Krause (1860) as existing in the conjunctiva, the genitals, and other parts of the human body, and differing from the Pacinian corpuscles only in the absence of a thick laminated investment.

Kreatin (*kre'-at-in*). See *Creatin*.

Kreatinin (*kre-at'-in-in*). See *Creatinin*.

Kreolin (*kre'-o-lin*). See *Croelin*.

Kresol (*kre'-sol*). See *Cresol*.

Kumiss (*koo'-mis*) or **Kumyss** (*koo'-mis*). See *Koumiss*.

Kyestein (*ki-es'-te-in*) [*κίησις*, conception]. A filmy deposit upon decomposing urine, at one time thought to be diagnostic of pregnancy.

Kymograph (*ki'-mo-graf*), **Kymographion** (*ki-mo-graf'-e-on*) [*κῡμα*, wave; *γράφειν*, to write]. An instrument for reproducing graphically the variations in the pressure of the blood.

Kymoscope (*ki'-mo-skôp*) [*κῡμα*, wave; *σκοπεῖν*, to inspect]. A device used in the observation and study of the blood-current.

Kynurin (*ki-nu'-rin*) [*κῡων*, a dog; *urina*, urine], $C_{10}H_{14}N_2O_2$. A crystalline substance obtained from cynurenic acid. It may also be made by oxidizing cinchonic acid and cinchoninic acid with chromic acid.

Kyphoscoliosis (*ki-fo-sko-le-o'-sis*) [*κῑφωσις*, hump-backed; *σκολίωσις*, a bending]. Kyphosis combined with scoliosis.

Kyphosis (*ki-fo'-sis*) [*κῑφωσις*, hump-back]. Hump-back. Angular curvature of the spine, with the prominence or convexity turned dorsad.

Kyphotic (*ki-fo'-ik*) [*κῑφωσις*, hump-back]. Relating to, of the nature of, or affected with, kyphosis.

L

L. The abbreviation of Left, of Lithium, and of *Libra*, a pound.

Labarraque's Solution. See *Chlorin*.

Labial (*la'-be-al*) [*labium*, a lip]. Pertaining to the lips.

Labile (*lab'-il*) [*labi*, to glide]. Gliding to and fro; applied to an electric current when the electrode is moved from place to place over the skin.

Labio- (*la'-be-o-*) [*labium*, lip]. A prefix meaning pertaining to the lip.

Labiochorea (*la-be-o-ko-re'-ah*) [*labium*, lip; *χορεία*, dancing]. A choreic affection of the lips, and the stammering that results from it.

Labioglossopharyngeal (*la-be-o-glos-o-lar-in'-je-al*) [*labium*, lip; *γλῶσσα*, tongue; *λάρυγξ*, larynx]. Pertaining conjointly to lips, tongue, and larynx. **L.** Paralysis. See *Paralysis*, *Bulbar*.

Labioglossopharyngeal (*la-be-o-glos-o-far-in'-je-al*) [*labium*, lip; *γλῶσσα*, tongue; *φάρυγξ*, pharynx]. Pertaining conjointly to lips, tongue, and pharynx.

Labioplasty (*la'-be-o-plas-ty*) [*labium*, lip; *πλάσσειν*, to form]. An operation for repairing an injured or diseased lip; chilo-plasty.

Labium (*la'-be-um*) [*L.*]. A lip. **L. majus**, or **L. pudendi majus**, one of two folds of skin of the female external genital organs, arising just below the mons Veneris, surrounding the vulval entrance, and meeting at the anterior part of the perineum. **L. minus**, **L. pudendi minus**, or *nympha*, one of two folds of mucous membrane at the inner surfaces of the labia majora. **L. tympanicum**, the portion of the lamina spiralis forming the lower border of the sulcus spiralis. **L. urethrae**, the lateral margins of the external urinary meatus. **L. vestibulare**, the overhanging extremity of the lamina spiralis that forms the upper part of the sulcus spiralis.

Labor (*la'-bor*) [*L.*, work]. Parturition; the bringing forth of young. **L.**, Artificial, that effected or aided by other means than the forces of the maternal organism. **L.**, Dry, that in which there is a deficiency of the liquor amnii, or in which there has been a premature rupture of the bag of waters. **L.**, Induced, labor brought on by artificial means. **L.**, Instrumental, one requiring instrumental means to extract the child. **L.**, Missed, retention of the dead fetus *in utero* beyond the period of normal gestation. **L.**, Premature, labor taking place before the normal period of gestation, but when the fetus is viable. **L.**, Stages of, arbitrary divisions

of the period of labor,—the first begins with dilatation of the os, and ends with complete dilatation; the second ends with the expulsion of the child; the third (*placental*) consists in the expulsion of the placenta. **L.**, Mechanism of, the mechanism by which a fetus and its appendages traverse the birth-canal and are expelled. **L.-pains**, the pains consequent upon the contractions of the uterus during labor. **L.**, Precipitate, labor in which the expulsion of the fetus and its appendages takes place with undue celerity. **L.**, Protracted, labor prolonged beyond the usual limit (10-20 hours in primiparae, 2-6 hours in multiparae).

Laboratory (*lab'-or-a-to-re*) [*laborare*, to work]. A room or place designed for experimental scientific work.

Labyrinth (*lab'-ir-inth*) [*λαβύρινθος*, a maze].

1. A name given to the series of cavities of the internal ear comprising the vestibule, cochlea, and the semicircular canals. 2. The parts of the cortex of the kidney between the medullary rays. See *Kidney*. **L.**, Bony. See *L.*, *Osseous*. **L.**, Membranous, the membranous cavity within the osseous labyrinth, from which it is partly separated by the perilymph. **L.**, Osseous, the bony portion of the internal ear.

Labyrinthal (*lab-ir-in'-thal*), **Labyrinthic** (*lab-ir-in'-thik*), **Labyrinthine** (*lab-ir-in'-thin*) [*λαβύρινθος*, a maze]. Pertaining to a labyrinth. **L.**, Vertigo. See *Ménière's Disease*.

Lac (*lak*) [*L.*]. Milk. **L. sulphuris**, milk of sulphur; sulphur precipitatum (U. S. P.).

Laceration (*las'-er-a'-shun*) [*lacerare*, to tear]. A tear. **L. of Perineum**, a tearing through the wall separating the lower extremity of the vagina and rectum, occurring occasionally during childbirth.

Lachrymal (*lak'-rim-al*). See *Lacrimal*.

Lacmus (*lak'-mus*). See *Litmus*.

Lacrimal (*lak'-rim-al*) [*lacrima*, a tear].

Pertaining to the tears or to the organs secreting and conveying the tears. **L. Apparatus**, the lacrimal gland, ducts, canal, sac, and nasal duct. **L. Artery**, the first branch of the ophthalmic artery, supplying the gland. **L. Bone**, a bone upon the nasal side of the orbit, articulating with the frontal, the ethmoid, and superior maxillary bones, in which begin the lacrimal groove and nasal duct. **L. Canals**, or *Canaliculi*, superior and inferior, extend from the lacrimal puncta to the sac, and serve to convey the excess of tears from the eye to the nose. **L. Caruncle**. See *Caruncle*. **L. Ducts**, seven to fourteen ducts,

extending obliquely from the gland to the fornix conjunctivæ, carrying the tears to the conjunctival surface of the eye-ball. **L. Fistula**, a fistula communicating with a lacrimal duct. **L. Gland**, the gland secreting the tears, situated in a depression of the frontal bone, the **L. fossa**, at the upper and outer angle of the orbit. **L. Lake**, the inward prolongation of the palpebral fissure of the eyelids. **L. Papilla**. See *Papilla*. **L. Probe**, a probe for exploring or dilating the canaliculi and nasal duct. **L. Puncta**, the minute orifices of the canaliculi, upon the eyelids near the inner canthus. **L. Sac**, a saccular enlargement of the upper part of the nasal duct, into which the canaliculi empty. **L. Style**, a probe used in stricture of the nasal duct.

Lacrimation (*lak-rim-a'-shun*) [*lacrima*, a tear]. An excessive secretion of tears.

Lactalbumin (*lakt-al'-bu-min*) [*lac*, milk; *albumin*]. A proteid contained in milk; it resembles serum-albumin, and coagulates at a temperature of from 70° to 80° C.

Lactate (*lak'-tat*) [*lac*, milk]. A salt of lactic acid.

Lactation (*lak-ta'-shun*) [*lactare*, to suckle]. Suckling; the period during which the child is nourished from the breast.

Lacteal (*lak'-te-al*) [*lac*, milk]. 1. Pertaining to milk. 2. Any one of the lymphatics of the small intestine that take up the chyle.

Lactic (*lak'-tik*) [*lac*, milk]. Pertaining to milk or its derivatives. **L. Acid**. See *Acid*, *Lactic*.

Lactiferous (*lak-tif'-er-us*) [*lac*, milk; *ferre*, to carry]. Conveying or secreting milk. **L. Ducts**, the ducts of the mammary gland. **L. Glands**, the mammary glands.

Lactifuge (*lak'-tif-uj*) [*lac*, milk; *fugare*, to drive away]. 1. Lessening the secretion of milk. 2. A drug or agent that causes a lessening in the secretion of milk.

Lactigenous (*lak-tif'-en-us*) [*lac*, milk; *γεννᾶν*, to produce]. Milk-producing.

Lactin (*lak'-tin*) [*lac*, milk]. Sugar of milk; lactose.

Lactinated (*lak'-tin-a-ted*) [*lac*, milk]. Containing sugar of milk.

Lactivorous (*lak-tiv'-or-us*) [*lac*, milk; *vorare*, to devour]. Subsisting on milk.

Lacto- (*lak'-to-*) [*lac*, milk]. A prefix denoting relation to milk.

Lactoceles (*lak'-to-sel*) [*lac*, milk; *κυστήρ*, tumor]. See *Galactocoele*.

Lactoglobulin (*lak-to-glob'-u-lin*) [*lac*, milk; *globulin*]. One of the proteids of colostrum.

Lactometer (*lak'-tom'-et-er*) [*lac*, milk; *μετρεῖν*, a measure]. An instrument for determining the specific gravity of milk.

Lactone (*lak'-tōn*) [*lac*, milk], $C_{10}H_{16}O_4$. An

aromatic, colorless, inflammable fluid, obtained in the dry distillation of lactic acid.

Lactophosphate (*lak-to-fos'-fat*) [*lac*, milk; *phosphate*]. A salt composed of a base united to lactic and phosphoric acid.

Lactoprotein (*lak-to-pro'-te-in*) [*lac*, milk; *protein*]. A proteid said to exist in milk.

Lactose (*lak'-tōs*) [*lac*, milk], $C_{12}H_{22}O_{11} + H_2O$. Milk-sugar; a sugar found in the milk of mammals, and at times in the urine of nursing women. It forms white, hard, rhombic crystals, soluble in water, and has a sweetish taste. Under the name of *saccharum lactis* it is official in U. S. P. Its chief use is as a vehicle.

Lactuca (*lak-tu'-kah*) [from *lac*, milk, on account of the milky juice]. A genus of composite-flowered herbs; the lettuces. **L. sativa**, the common garden lettuce. **L. virosa**, is the source of lactucarium.

Lactucarium (*lak-tu-ka'-re-un*). Lettuce. The concrete milky juice of *Lactuca virosa*, a plant of the order Compositæ. It contains a substance, lactucin, to which its properties are thought to be due, is sedative and anodyne, and has been used in cough and nervous irritability. **L., Syrupus**. Dose fʒ ij-ijj (8.0-12.0). **L., Tinct.** Dose fʒj (4.0).

Lacuna (*la-ku'-nah*) [*lacus*, a lake]. A hollow space. **L. of Bone**. See *Bone*. **L., Howship's**. 1. Depressions on the surface of bone beneath the periosteum. 2. Carious excavations in bone filled with granulation-tissue. **L. magna**, the largest of the orifices of the glands of Littre, situated on the upper surface of the fossa navicularis. **L. of Urethra**, follicular depressions in the mucous membrane of the urethra, most abundant along the floor, especially in the region of the bulb. Their mouths are directed forward.

Lacunar (*la-ku'-nar*) [*lacus*, lake]. Pertaining to the lacunæ. **L. Tonsillitis**. See *Tonsillitis*.

Lacunula (*lak-u'-nu-lah*) [dim. of *lacuna*, a lake]. A small lacuna.

Lacus (*la'-kus*) [*L.*, lake]. A small cavity. **L. lacrimalis**, the space at the inner canthus of the eye, near the punctum, in which the tears collect.

Lacustrine (*la-ku'-trin*) [*lacus*, a lake]. Inhabiting lakes or ponds.

Lady Webster Pill. See *Aloe*.

Lady's Slipper. See *Cypripedium*.

Laennec's Cirrhosis (*len-ek's*) [R. T. H. Laennec, a French physician, 1781-1826]. Atrophic cirrhosis of the liver.

Lævo- (*le'-vo-*). For words so commencing see *Levo-*.

Lævulose (*le'-u-lōs*). See *Levulose*.

Lafayette Mixture. A mixture employed in gonorrhea. It contains copaiba, cubets,

liquor potassæ, sweet spirit of niter, and is known also as the compound copaiba mixture.

Lagophthalmos (*lag-off-thal'-mos*) [*λαγός*, hare; *ὀφθαλμός*, eye; from the popular notion that a hare sleeps with open eyes]. A condition in which the eyes cannot be closed.

La Grippe (*lah grip*) [Fr.]. Influenza.

Lake-colored (*lak'-cul-erd*), or **Laky** (*la'-ke*) [Fr., *laque*, rose-colored, from Pers. *lak*]. Applied to blood that is dark-red and transparent from a solution of the hemoglobin in the serum.

Lallation (*la'-al'-shun*) [*lallare*, to babble]. Any unintelligible stammering of speech, such as the prattling of a babe.

Laloneurosis (*la'-a-nu-ro'-sis*) [*λάλος*, prattle; *νεῖρον*, nerve]. An impairment of speech arising from spasmodic action of the muscles. It includes stammering and aphthongia.

Lalopathy (*la'-op'-ath-e*) [*λάλος*, speech; *πάθος*, disease]. Any disorder of speech.

Lambda (*lam'-dah*) [*λάμβδα*, the Greek letter Λ or λ]. The angle of junction of the sagittal and lambdoid sutures.

Lambdacism (*lam'-das-izm*) [*λαμβδακισμός*, lambdacism]. 1. Difficulty in uttering the sound of the letter *l*. 2. Too frequent use of the *l* sound, or its substitution for the *r* sound.

Lambdoid, or **Lambdoidal** (*lam'-doid*, or *lam-doi'-dal*) [*λάμβδα*, the letter λ ; *είδος*, resemblance]. Resembling the Greek letter Λ . **L. Suture**, the suture between the occipital and the two parietal bones.

Lamella (*lam-el'-ah*) [dim. of *lamina*, a plate]. A thin scale or plate. **L. of Bone**, the concentric rings surrounding the Haversian canals. **L., Concentric**, one of the plates of bone surrounding the Haversian canal. **L., Intermediate**, one of the plates filling the spaces between the concentric layers of bone. **L., Periosteal**, or **Peripheral**, a superficial lamella of bone lying under the periosteum.

Lamellar (*lam-el'-ar*) [*lamella*, a thin plate]. Having the nature of or resembling a thin plate; composed of lamellæ or thin plates. **L. Cataract**. See *Cataract*.

Lamina (*lam'-in-ah*) [*L.*, a plate or scale]. A thin plate or layer. **L. cinerea**, the connecting layer of gray matter between the corpus callosum and the optic chiasma. **L., Cribriform**, the cribriform plate of the ethmoid. **L. cribrosa** (*of the choroid*), that portion of the choroid which is perforated for the passage of the optic nerve. **L. cribrosa**. 1. The cribriform fascia covering the sphenous opening. 2. The anterior or posterior perforated space of the brain. 3. The perforated plates of bone through which branches of the

cochlear or auditory nerve pass. **L. fusca**, the pigmentary tissue of the inner layer of the sclera forming the outer layer of the perichoroidal sinus. **L. propria** (*of the tympanic membrane*), the middle or fibrous layer of the tympanic membrane. **L., Reticular**, the hyaline membrane of the inner ear, extending between the conjoined head of Corti's rods and the supporting cells. **L. spiralis**, a thin plate in the ear, osseous in the inner part and membranous in the outer, which divides the spiral tube of the cochlea into the scala tympani and the scala vestibuli. **L. suprachoroidea**, the delicate connective-tissue membrane uniting the choroid and sclerotic coats of the eye. **L., Vitreous**, a homogeneous membrane covering the inner surface of the choroid; it is also called the membrane of Bruch.

Laminar (*lam'-in-ar*) [*lamina*, a layer]. Composed of laminae; having the form of a lamina.

Laminaria (*lam-in-a'-re-ah*) [*lamina*, a thin plate]. 1. A genus of algæ, or seaweeds. 2. The stems of *L. cloustoni*, or *L. digitata*. **L. Tent**, a tent made of the stem of the laminaria plant, for dilatation of the cervix uteri and other canals.

Laminated (*lam'-in-a-ted*). See *Laminar*.

Lamination (*lam-in-a'-shun*) [*lamina*, a layer]. 1. Arrangement in plates or layers. 2. An operation in embryotomy, consisting in cutting the skull in slices.

Laminectomy (*lam-in-ek'-to-me*) [*lamina*, a layer; *ἐκτομή*, excision]. The operation of removing the posterior vertebral arches.

Lamp-black [*λαμπάς*, a lamp]. A fine black substance, almost pure carbon, made by burning coal-oils in an atmosphere deficient in oxygen, or by allowing a gas-flame to impinge on a cold surface.

Lancet (*lan'-set*) [dim. of *lancea*, a lance]. A knife having a double-edged, lance-shaped blade for incising tumors, abscesses, etc. **L., Gum**, a small lancet in which the cutting portion has a convex edge and is at right-angles to the shaft; it is used for cutting the gums. **L., Spring**, one in which the blade is thrust out by means of a spring, controlled by a trigger. **L., Thumb**, one with a double-edged, broad blade.

Lancinating (*lan'-sin-a-ting*) [*lancinare*, to tear]. Tearing; shooting. **L. Pains**, rending, tearing, or sharply-cutting pains, common in posterior spinal sclerosis.

Lancisi, Nerves of. *Striæ longitudinales*. The slight ridges of the corpus callosum on either side of the raphé.

Landry's Disease or **Paralysis**. [J. B. O. Landry, a French physician]. Acute ascending paralysis; a form of paralysis characterized by loss of motor power in

the lower extremities, gradually extending to the upper extremities, and to the centers of circulation and respiration, without sensory manifestations, trophic changes, or variations in electric reaction.

Land-scurvy. *Purpura hæmorrhagica.*

Lanolin (*lan'-o-lin*) [*lana*, wool; *oleum*, oil]. The *Adeps lanae hydrosus* (U. S. P., B. P.). A cholesterol fat obtained from sheeps' wool, and used as a basis for ointments.

Lantermann's Notches or Segments. Cylindric segments in the medullary sheath of a nerve-fiber, marked off from one another by the incisures of Schmidt.

Lanugo (*lan-u'-go*) [*lana*, wool]. The down-like hair that appears upon the fetus at about the fifth month of gestation; also the downy growth often seen upon the face of women and girls.

Laparo- (*lap'-ar-o*) [*λαπάρα*, loin]. A prefix denoting pertaining to the abdomen, properly, referring to the loin or flank. See *Celiotomy*.

Laparocolotomy (*lap'-ar-o-ko-lot'-o-me*) [*λαπάρα*, loin; *κόλον*, colon; *τομή*, a cutting]. Inguinal or abdominal colotomy.

Laparocystectomy (*lap'-ar-o-sis-tek'-to-me*) [*λαπάρα*, loin; *κύστις*, cyst, bladder; *ἐκτομή*, excision]. An operation performed in advanced extrauterine pregnancy for removal of the fetus and the entire gestation-sac.

Laparocystotomy (*lap'-ar-o-sis-tot'-o-me*) [*λαπάρα*, loin; *κύστις*, cyst, bladder; *τομή*, a cutting]. 1. Suprapubic cystotomy. 2. An operation in advanced extrauterine pregnancy for the removal of the fetus, the sac being allowed to remain.

Laparoelytotomy (*lap'-ar-o-el-it-rot'-o-me*) [*λαπάρα*, loin; *ἐλκτρον*, sheath; *τομή*, a cutting]. An operation consisting in an incision over Poupert's ligament, dissecting up the peritoneum until the vagina is reached, incising the latter transversely, dilating the cervix, and extracting the child through the os uteri.

Laparoenterotomy (*lap'-ar-o-en-ter-ot'-o-me*) [*λαπάρα*, loin; *έντερον*, intestine; *τομή*, a cutting]. An opening of the intestine through an abdominal incision.

Laparogastrotomy (*lap'-ar-o-gas-trot'-o-me*) [*λαπάρα*, loin; *γαστήρ*, stomach; *τομή*, a cutting]. The opening of the stomach through an abdominal incision.

Laparohysterectomy (*lap'-ar-o-his-ter-ek'-to-me*) [*λαπάρα*, loin; *ἱστέρα*, womb; *τομή*, a cutting]. The removal of the uterus through an incision in the abdominal walls.

Laparohysteroophorectomy (*lap'-ar-o-his-ter-o-o-off-or-ek'-to-me*) [*λαπάρα*, loin; *ἱστέρα*, womb; *ὄον*, egg; *φέρω*, to bear; *ἐκτομή*, a cutting out]. Removal of the uterus and ovaries through an incision in the abdominal wall.

Laparohysterotomy (*lap'-ar-o-his-ter-ot'-o-me*) [*λαπάρα*, loin; *ἱστέρα*, uterus; *τομή*, a cutting]. The operation of cutting into the uterus through an abdominal incision, as for the purpose of removing a fetus.

Laparoileotomy (*lap'-ar-o-il-e-ot'-o-me*) [*λαπάρα*, loin; *ίλεον*; *τομή*, a cutting]. The operation of cutting into the ileum through an abdominal incision.

Laparonephrectomy (*lap'-ar-o-nef-tek'-to-me*) [*λαπάρα*, loin; *νεφρός*, kidney; *ἐκτομή*, a cutting out]. Nephrectomy by an abdominal incision.

Laparosalpingectomy (*lap'-ar-o-sal-ping-ek'-to-me*) [*λαπάρα*, loin; *σάλπιγξ*, tube; *ἐκτομή*, a cutting out]. Removal of a Fallopian tube through an abdominal incision.

Laparosplenectomy (*lap'-ar-o-splen-ek'-to-me*) [*λαπάρα*, loin; *σπλήν*, spleen; *ἐκτομή*, a cutting out]. Removal of the spleen through an abdominal incision.

Laparotomist (*lap'-ar-ot'-o-mist*) [*λαπάρα*, loin; *τομή*, a cutting]. A surgeon who performs laparotomies.

Laparotomy (*lap'-ar-ot'-o-me*) [*λαπάρα*, loin; *τομή*, a cutting]. 1. An incision through the abdominal wall; celiotomy is the preferable term. 2. The operation of cutting into the abdominal cavity through the loin or flank.

Lapis (*la'-pis*) [L.]. A stone; an alchemic term applied to any nonvolatile substance. L. divinus, aluminated copper. L. imperialis, silver nitrate. L. lazuli, a beautiful blue stone of complex composition, formerly employed as a purgative and emetic, and in epilepsy. L. lunaris, silver nitrate. L. mitigatus, diluted silver nitrate.

Lappa (*lap'-ah*) [L.]. Burdock. The root of the common burdock, *Arctium lappa*, containing a bitter principle, a resin, and tannin. It is aperient, diuretic, and alterative, and has been employed in gout, scorbutus, syphilis, and in various skin-diseases. The dose of the root is ʒj-ij (4.0-8.0); in infusion or tincture ʒx-fʒj (0.65-4.0). L., Ext., Fld. Dose ʒxxx-fʒj (2.0-4.0).

Larch. See *Larix*.

Lard [*lar-da*, lard]. The fat of the interior of the abdominal cavity of the hog, constituting *Adeps*, of U. S. P. Lard is much used in pharmacy as a basis for ointments. See *Adeps*.

Lardacein (*lar-da'-se-in*) [*lar-da*, lard]. Amyloid substance, formed in amyloid degeneration of various organs, particularly the liver, kidney, and spleen. It is a proteid, but insoluble in the ordinary solvents, is not acted upon by the gastric juice, does not readily undergo putrefaction, and gives a mahogany-brown color with iodine, and a blue color with iodine and sulphuric acid.

Lardaceous (*lar-da'-se-us*) [*lar-da*, lard]. Amyloid. **L. Kidney.** See *Bright's Disease*.

Larix (*la'-riks*) [*L.*]. Larch. A genus of coniferous, deciduous trees. **Laricis Cortex** (B. P.), is astringent and stimulant, and has been used in purpura, hemoptysis, bronchitis, and, locally, in skin-diseases. **L., Tinct.,** (B. P.). Dose ℥xx-xxx (1.3-2.0).

Larkspur (*lar-k'-spur*). The seed of *Delphinium consolida*, a diuretic and emmenagogue. Dose of the fld. ext., ℥j-x (0.065-0.65). Unof.

Laryngeal (*lar-in'-je-al*) [*lárnyξ*, larynx]. Pertaining to the larynx. **L. Crisis**, an acute laryngeal spasm, occurring in the course of *tuberculosis dorsalis*. **L. Mirror**, a small circular mirror affixed to a long handle, used in laryngoscopy.

Laryngectomy (*lar-in-jek'-to-me*) [*lárnyξ*, larynx; *ἐκτομή*, a cutting out]. Extirpation of the larynx.

Laryngismus (*lar-in-jis'-mus*) [*lárnyξ*, the larynx]. A spasm of the larynx. **L. stridulus**, a spasmodic affection of the larynx, characterized by sudden arrest of respiration, with increasing cyanosis, followed by long, loud, crowing inspirations. It is most common in rachitic children, but may also occur as a symptom of laryngeal catarrh.

Laryngitis (*lar-in-jis'-tis*) [*lárnyξ*, larynx; *itis*, inflammation]. Inflammation of the larynx. It may be acute or chronic, catarrhal, suppurative, croupous (diphtheric), tuberculous, or syphilitic. Chronic catarrhal **L.** is divided into a hypertrophic and an atrophic stage. The symptoms of acute catarrhal laryngitis, the most common form, are hoarseness, pain, dryness of the throat, dysphagia, and cough.

Laryngocentesis (*lar-in-go-sen-te'-sis*) [*lárnyξ*, larynx; *κέντησις*, puncture]. Puncture of the larynx.

Laryngofissure (*lar-in-go-fish'-ar*) [*lárnyξ*, larynx; *findere*, to cleave]. Division of the larynx for the removal of tumors or foreign bodies.

Laryngology (*lar-in-gol'-o-je*) [*lárnyξ* larynx; *λόγος*, science]. The science of the anatomy, physiology, and diseases of the larynx.

Laryngoparalysis (*lar-in-go-par-al'-is-is*) [*lárnyξ*, larynx; *παράλυσις*, palsy]. Paralysis of the laryngeal muscles.

Laryngopathy (*lar-in-gop'-ath-e*) [*lárnyξ*, larynx; *πάθος*, a suffering]. Any disease of the larynx.

Laryngophantom (*lar-in-go-fan'-tum*) [*lárnyξ*, larynx; *φάντασμα*, an apparition]. An artificial larynx designed for illustrative purposes.

Laryngopharyngeal (*lar-in-go-far-in'-je-al*) [*lárnyξ*, larynx; *φάρυγξ*, pharynx].

Pertaining conjointly to the larynx and pharynx.

Laryngopharynx (*lar-in-go-far'-inks*) [*lárnyξ*, larynx; *φάρυγξ*, pharynx]. The inferior portion of the pharynx. It extends from the greater cornua of the hyoid bone to the inferior border of the cricoid cartilage.

Laryngophony (*lar-in-goff'-o-ne*) [*lárnyξ*, larynx; *φωνή*, voice]. The sound of the voice observed in auscultation of the larynx.

Laryngophthisis (*lar-in-go-tis'-is*, *lar-in-goff'-this-is*) [*lárnyξ*, larynx; *φθίσις*, wasting]. Laryngeal tuberculosis.

Laryngoplasty (*lar-in-go-plas-te*) [*lárnyξ*, larynx; *πλάσσειν*, to shape]. Plastic operation upon the larynx.

Laryngoplegia (*lar-in-go-ple'-je-ah*) [*lárnyξ*, larynx; *σκοπεῖν*, to examine]. Paralysis of one or more muscles of the larynx.

Laryngoscope (*lar-in-go-skop*) [*lárnyξ*, larynx; *σκοπεῖν*, to examine]. A mirror attached to a long handle for examining the interior of the larynx.

Laryngoscopist (*lar-in-gos'-ko-pist*) [*lárnyξ*, larynx; *σκοπεῖν*, to examine]. An expert in laryngoscopy.

Laryngoscopy (*lar-in-gos'-ko-pe*) [*lárnyξ*, larynx; *σκοπεῖν*, to examine]. Examination of the interior of the larynx by means of the laryngoscope.

Laryngospasm (*lar-in-go-spasm*) [*lárnyξ*, larynx; *σπασμός*, spasm]. Spasmodic closure of the glottis.

Laryngostenosis (*lar-in-go-sten-o'-sis*) [*lárnyξ*, larynx; *στένωσις*, contraction]. Contraction or stricture of the larynx.

Laryngotomy (*lar-in-got'-o-me*) [*lárnyξ*, larynx; *τείνειν*, to cut]. The operation of incising the larynx. **L., Complete**, incision of the larynx through its whole length. **L., Median**, incision of the larynx through the thyroid cartilage. **L., Subhyoid**, incision of the larynx through the thyrohyoid membrane. **L., Superior**, and **L., Thyrohyoid**, incision of the larynx through the thyrohyoid membrane.

Laryngotracheal (*lar-in-go-tra'-ke-al*) [*lárnyξ*, larynx; *τραχεία*, the windpipe]. Pertaining conjointly to the larynx and the trachea.

Laryngotracheitis (*lar-in-go-tra-ke-i'-tis*) [*lárnyξ*, larynx; *τραχεία*, windpipe; *itis*, inflammation]. Inflammation of the larynx and the trachea.

Laryngotracheotomy (*lar-in-go-tra-ke-o'-me*) [*lárnyξ*, larynx; *τραχεία*, windpipe; *τομή*, cutting]. That form of tracheotomy in which the cricoid cartilage and one or more of the upper rings of the trachea are divided.

Laryngotypoid (*lar-in-go-ti'-foid*), **Laryngotypus** (*lar-in-go-ti'-fus*) [*lárnyξ*,

larynx; τῖφος, stupor]. Typhoid fever associated with marked laryngeal complications.

Larynx (*lar'-inks*) [*λάρυξ*, larynx]. The organ of the voice situated between the trachea and the base of the tongue. It consists of a series of cartilages, the thyroid, the cricoid, and the epiglottis, and three pairs of cartilages, the arytenoids, and those of Santorini and Wrisberg, which are lined by mucous membrane, and are moved by the muscles of the larynx. The mucous membrane is, on each side, thrown into two transverse folds that constitute the vocal bands, the upper being the false, the lower the true vocal band. By the approximation or separation of the vocal bands the changes in the pitch of the voice are produced. The space between the vocal bands is termed the glottis.

Lash [O. L. G., *laske*, a flap]. An eye-lash.

Lata, or **Latah** (*la'-tah*) [Jav.]. An hysterical neurosis prevalent in Java, and nearly identical with the disorder of the jumpers in Maine and Canada, and with the myriachit of Siberia.

Latent (*la'-tent*) [*latere*, to be hid]. Concealed; not manifest; potential. **L. Heat**, that which apparently disappears when a liquid is vaporized or a solid melted. **L. Period**. 1. The time required for the incubation of a disease. 2. In physiology, the time intervening between the application of a stimulus and the appearance of the resulting phenomenon.

Laterad (*lat'-er-ad*) [*latus*, the side; *ad*, toward]. Toward the lateral aspect.

Lateral (*lat'-er-al*) [*latus*, the side]. At, belonging to, or pertaining to the side; situated on either side of the median vertical plane. **L. Column**, that column of the spinal cord between the anterior and posterior horns. **L. Operation**, that form of lithotomy in which the opening is made on the right or left side of the perineum. **L. Sclerosis**, **Amyotrophic**, a disease of the lateral columns and anterior gray matter of the cord. It is characterized by motor weakness and a spastic condition of the limbs, associated with atrophy of the muscles and final involvement of the nuclei in the medulla oblongata. **L. Sclerosis**, **Primary**, a sclerotic disease of the crossed pyramidal tracts of the cord characterized by paralysis of the limbs, with rigidity, increased tendon-reflexes, and absence of sensory and nutritive disorders. A peculiar characteristic jerking gait is produced, and clonus of the lower limbs may be readily excited. **L. Sinuses**, the two veins of the dura mater situated in the attached margin of the tentorium cerebelli. **L. Ventricles**. See *Ventricle*.

Lateritious, **Lateritious** (*lat'-er-ish'-us*) [*later*, brick]. Resembling brick-dust, as the *L.* sediment of the urine.

Lateroabdominal (*lat'-er-o-ab-dom'-in-al*) [*latus*, side; *abdominalis*, pertaining to the abdomen]. Pertaining both to the side and the abdomen. **L. Posture**, Sims' posture.

Lateroflexion (*lat'-er-o-flek'-shun*) [*latus*, side; *flectere*, to bend]. Flexion or bending to one side.

Lateropulsion (*lat'-er-o-pul'-shun*) [*latus*, side; *pellere*, to drive]. An involuntary motion to one side.

Lateroversion (*lat'-er-o-ver'-shun*) [*latus*, side; *vertere*, to turn]. A turning to one side, as *L.* of the uterus.

Lathyrism (*lath'-ir-izm*) [*λάθρις*, pulse]. Lupinosis; an affection produced by the use of meal from varieties of vetches, chiefly the *Lathyrus sativus* and *L. cicera*. It is a form of spastic paraplegia with tremor, involving chiefly the legs.

Latissimus (*lat'-is'-im-us*) [superl. of *latus*, wide]. An adjective signifying widest. **L. colli**. See *Muscles* (*Platysma myoides*), *Table of*. **L. dorsi**. See *Muscles*, *Table of*.

Laudable (*law'-da-bl'*) [*laus*, praise]. Praiseworthy. **L. Pus**. See *Pus*.

Laudanin (*lod'-an-in*) [*laudanum*]. $C_{20}H_{25}NO_4$. One of the alkaloids of opium.

Laudanum (*lod'-an-um*) [Pers., *lādan*, the gum-herb lada]. *Tinctura opii*. See *Opium*.

L. Sydenham's, *Vinum opii*. See *Opium*. **Laughing**, or **Laughter** (*lahf'-ing*, *lahf'-ter*) [AS., *hlehhhan*, to laugh]. A succession of rhythmic, spasmodic expirations with open glottis and vibration of the vocal bands, and indicating mirth. **L.-gas**, nitrous oxid, or protoxid of nitrogen. See *Nitrous Oxid* and *Anesthetics*.

Laurel (*lor'-el*) [*laurus*]. See *Cherry-laurel*, *Laurus*. **L.-water**, aqua laurocerasi. See *Cherry-laurel*.

Laurocerasus (*lor'-o-ser'-as-us*) [*laurus*, laurel; *cerasus*, cherry-tree]. See *Cherry-laurel*.

Laurus (*lor'-us*) [L.]. A tree of the Lauraceæ. **L. nobilis**, the noble laurel, is indigenous in the south of Europe. Its fragrant oils (one essential, from the leaves, and the other fixed, from the berries) are chiefly used in liniments. Unof.

Lavage (*lav'-ahzh'*) [Fr.]. The irrigation or washing out of an organ, such as the stomach, the bowel, etc.

Lavation (*lav'-a'-shun*) [*lavare*, to wash]. Lavage.

Lavender (*lav'-en-der*) [*lavare*, to wash]. The flowers of *Lavandula vera*, a plant of the order Labiatae. The active principle is a volatile oil (*Oleum lavandulæ florum*), which is used in the preparation of *Spiritus lavandulæ*. Dose mxxx-fj (2.0-4.0). *Tinct. lavandulæ comp.*, formerly termed compound spirit of lavender.

Dose $m_{xxx}-[5]$ (2.0-4.0). Both are stimulant and carminative, and are used as vehicles.

Laveran's Corpuscles, or Plasmodia. See *Plasmodium*.

Law (*laro*) [AS., *lagu*, a law]. A general rule; a constant mode of action of forces, or phenomena. A rule of action prescribed by authority. **L. of Avogadro**, equal volumes of all gases and vapors, at like temperature and like pressure, contain an equal number of molecules. **L., Behring's**, the blood and blood-serum of an individual which has been artificially rendered immune against a certain infectious disease, may be transferred into another individual with the effect of rendering the latter also immune. **L., Berthollet's**, when two salts in solution can by double decomposition produce a salt less soluble than either, this salt will be produced. **L., Boyle's**, at any given temperature the volume of a given mass of gas varies inversely as the pressure that it bears. This is also called *Mariotte's Law*. **L., Charles's**, equal increments of temperature add equal amounts to the product of the volume and pressure of a given mass of gas. The increase is $\frac{1}{273}$ of its volume measured at -273° C., which is the zero of absolute temperature. **L., Colles'**, a child born of a mother who is without obvious venereal symptoms, and which, without being exposed to any infection subsequent to its birth, presents symptoms of syphilis when a few weeks old, will not infect its mother. **L., Dalton's**, or **L., Dalton-Henry's**, although the volume of a gas absorbed by a liquid remains constant, the weight (volume multiplied by the density) of the absorbed gas rises and falls in proportion to its pressure. **L. of Definite Proportions**, when two or more chemic substances unite to form a compound, they do so in a fixed and constant proportion. **L., Fechner's**, with increase of the stimulus the sensation increases only as the logarithm of the stimulus. **L., Gay-Lussac's**. See *L., Charles's*. **L., Graham's**, the rate of diffusion of gases through porous membranes is in inverse ratio to the square-root of their density. **L., Henry's**. See *L., Dalton's*. **L., Hilton's**, a nerve-trunk supplying a given joint also supplies the muscles moving that joint and the skin over the insertion of those muscles. **L., Listing's**, pertains to the movements of the eye-ball; when moved from the position of rest, the angle of rotation in the second position is the same as if the eye were turned about a fixed axis perpendicular to the first and second positions of the visual line. **L., Mariotte's**. See *L., Boyle's*. **L. of Multiple Proportions**, two substances uniting to form a series of chemic compounds

do so in proportions that are simple multiples of one another or of one common proportion. **L., Ohm's**, the current-strength varies directly as the electromotive force, and inversely as the resistance. **L., Profeta's**, a child born of syphilitic parents, though apparently free from syphilis, cannot be infected. **L. of Reciprocal Proportions**, two elements combining with a third do so in proportions that are simple multiples or simple fractions of those in which they combine with each other. **L. of Refraction**, rays of light entering a denser medium are deflected toward a perpendicular drawn through the point of incidence, and those entering a rarer medium are deflected away from the same perpendicular. **L., Weber's**, the variation of stimulus that causes the smallest appreciable variation in sensation maintains, approximately, a fixed ratio to the total stimulus.

Laxative (*laks'-a-tiv*) [*laxare*, to loosen]. 1. Aperient; mildly cathartic. 2. An agent that loosens the bowels; a mild purgative.

Laxator (*laks-a'-tor*) [*laxare*, to loosen]. That which loosens or relaxes. A name applied to various muscles. **L. tympani**. See *Muscles, Table of*.

Layer (*la'-er*) [*legan*, to cause to lie down]. A mass of uniform, or nearly uniform, thickness, spread over a considerable area. **L., Bacillar** (*of the retina*), the layer of rods and cones of the retina. **L., Ganglionic**, a layer of angular cells of the cerebral cortex, best developed in the motor area. **L., Horny** (*of the epidermis*), the superficial layer of the skin. **L., Osteogenetic**, the lower layer of periosteum, connected with the formation of bone. **L. of Rods and Cones**. See *L., Bacillar*.

Layman (*la'-man*) [*laos*, the people]. A member of the laity; a person not a physician, or not professionally educated.

Lazaretto (*laz-ar-et'-o*) [Ital., a pest-house, from *lazar*, a leper]. A quarantine establishment; a pest-house. Also, a place for fumigation and disinfection.

Lead (*led*) [AS., *ledd*, lead]. See *Plumbum*. **L., Black**, graphite. **L.-colic**. See *Colic*. **L.-encephalopathy**, the cerebral manifestations of chronic lead-poisoning. They consist of epilepsy, acute delirium, and hallucinations, and may terminate in insanity. **L.-line**, blue line; the line of discoloration on the gums in cases of chronic lead-poisoning. **L.-pipe Contraction**, the condition of the limbs in the cataleptic state, in which they maintain any position that is given them. **L.-poisoning**, plumbism; saturnism,—a form of poisoning due to the introduction of lead into the system. The symptoms are disturbed nutrition, anemia, a blue line on the

gums, lead-colic, constipation, pains in the limbs, local muscular paralysis (wrist-drop) and wasting, saturnine encephalopathy, etc.

Leader (*le'-der*) [AS., *lād*, a way or path]. A sinew or tendon.

Leathery (*leth'-er-e*) [AS., *lēder*, leather]. Resembling leather, as L. arteries, arteries thickened and feeling like cords of leather.

Lecithin (*les'-ith-in*) [*lekithos*, yolk of egg], $C_{44}H_{90}NPO_9$. A complex nitrogenous fatty substance occurring widely spread throughout the animal body. It is found in the blood, bile, serous fluids, brain, nerves, yolk of egg, semen, pus, and white blood-corpuscles. It is a colorless, slightly crystalline substance, soluble in alcohol, and chemically is looked upon as the glycerophosphate of neurin in which two atoms of hydrogen are replaced by two atoms of stearic, palmitic, or oleic acid.

Leech (*leech*) [AS., *lave*, physician]. A blood-sucking worm, the Sanguisuga, or Hirudo medicinalis, found in Europe; Hirudo decora, the American leech. Leeches are used for the local abstraction of blood. A European leech draws from 13 ss-j (16.0-32.0) of blood. L., **Artificial**, an apparatus for cupping.

Lees (*lez*) [Fr., *lie*, dregs]. The dregs of vinous liquors.

Leg [Icel., *legg*, leg]. The lower extremity, especially that part from the knee to the ankle. L., **Barbadoes**, elephantiasis of the leg. L., **Black**. See *Anthrax*, *Symptomatic*, and *Black-leg*. L., **Bow**, a curving outward of the legs. L., **Milk-**, phlegmasia alba dolens.

Legitimacy (*le-jit'-im-as-e*) [*legitimus*, lawful]. 1. The condition of being within the bounds of the law. 2. The state of a child born within wedlock, or within a period of time necessary to gestation, which may elapse after the death of the father.

Legumin (*leg-ul'-min*) [*legumen*, pulse]. A proteid found in the seeds of many plants belonging to the natural order of Leguminosæ.

Leiodermia (*li-o'-der'-me-ah*). See *Liodermia*.

Leliomyoma (*li-o-mi-o'-mah*). See *Lio-myoma*.

Leiter's Coil, or Tubes. Tubes of soft, flexible metal designed for application about any part of the body. Cold water is passed through the tubes, thereby reducing the temperature of the parts encased.

Lembert's Suture. See *Suture*.

Lemniscus (*lem-nis'-kus*) [*λεμνισκος*, fillet]. See *Fillet*.

Lemon (*lem'-on*). See *Limo*.

Lemonade (*lem-on-ad'*) [*limon*, lemon]. An acidulous, refrigerant drink made from the juice of lemon. L., **Sulphuric Acid**, a solution of sugar in water containing sul-

phuric acid. It is used in lead manufactories to prevent lead-poisoning.

Lens (*lenz*) [L., a lentil]. 1. A piece of glass or crystal for the refraction of rays of light. 2. The crystalline lens of the eye. L., **Achromatic**. See *Achromatic*. L., **Apo-chromatic**. See *Apochromatic*. L., **Biconcave** (negative or minus (—) lens), a thick-edged lens having concave spheric surfaces upon its opposite sides; it is used in spectacles to correct myopia. L., **Biconvex** (positive or plus (+) lens), a thin-edged lens; it has two convex surfaces, and is used to correct hyperopia. L., **Bifocal**. See *Bifocal*. L., **Convergent or Converging**, a double convex or planoconvex lens that focuses rays of light. L., **Convexoconcave**, a lens having a convex and a concave surface, which would not meet if continued. Its properties are those of a convex lens of the same focal distance. L., **Crystalline**, the lens of the eye, situated immediately behind the iris. L., **Cylindric** (either minus or plus), one with a plane surface in one axis and a concave or convex surface in the axis at right angles to the first. L., **Dispersing**, a concave lens. L., **Periscopic**, one with concavoconvex or convexoconcave surfaces, the opposite sides being of different curvatures; such lenses are called meniscus lenses. L., **Spheric**, one the curved surface of which, either concave or convex, is a segment of a sphere.

Lenticonus (*len-tik-o'-nus*) [*lens*, a lens; *conus*, a cone]. A rare, usually congenital, anomaly of the lens in which there is a conical prominence upon its anterior or, more rarely, upon its posterior, surface.

Lenticular (*len-tik'-u-lar*) [*lenticula*, a lentil]. 1. Pertaining to, or resembling, a lens. 2. Pertaining to the crystalline lens. 3. Pertaining to the lenticular nucleus of the brain. L., **Nucleus**, a mass of gray matter, the extraventricular portion of the corpus striatum, situated to the outer side of the internal capsule of the brain.

Lenticulostriate (*len-tik-u-lo-strī'-āt*) [*lenticula*, a lentil; *striatus*, striated]. Pertaining to the lenticular nucleus of the corpus striatum, as L. artery.

Lentigo (*len-ti'-go*) [L., a lentil-shaped spot; *ph.*, *Lentigines*]. A freckle; a circumscribed patch of pigment, small in size, occurring mainly on face and hands, and due to exposure to the sun. L., *æstiva*, summer-freckles.

Leontiasis (*le-on-ti'-as-is*) [*leōn*, a lion]. A lion-like appearance of the face seen in leprosy, elephantiasis, and L. ossea. L., **ossea**, L. ossium, an overgrowth of the bones of the face through which the features acquire a lion-like appearance.

Leontodin (*le-on'-to-din*) [*leōn*, a lion; *δόνος*, a tooth]. The precipitate from a tinc-

ture of the root of dandelion, *Leontodon taraxacum*; it is a tonic, diuretic, and aperient, and an hepatic stimulant. Dose 2-4 grains (0.13-0.26). Unof.

Leontodon (*le-on'-to-don*). See *Taraxacum*.

Leper (*lep'-er*) [*λεπρός*, scaly]. One affected with leprosy.

Lepra (*lep'-rah*) [*λέπρα*, leprosy]. 1. Leprosy. 2. A form of psoriasis. **L. anæsthetica**. See *Leprosy*. **L. maculosa**, the stage of true leprosy characterized by the presence of pigment-spots. **L. mutilans**, the final stage of true leprosy, in which there is a marked loss of tissue.

Leprosy (*lep'-ro-se*) [*λέπρα*, leprosy]. An endemic, chronic, infectious disease, due to the bacillus lepræ. Two forms of leprosy are described—a tubercular and an anæsthetic form. The first begins with a well-defined erythema (Macular L.), which is succeeded by the formation of papules, and later of nodules, although in some cases the erythematous stage is followed by a disappearance of the pigment without nodulation (*Lepra alba*). The nodules eventually break down and ulcerate. The anæsthetic form begins with pains and hyperæsthesia; a macular eruption appears, but again subsides, leaving spots of anæsthesia; trophic lesions develop and lead to the loss of the fingers or toes, with the production of marked deformity.

Leptandra (*lep'-tan'-drah*) [*λεπτός*, thin; *άνδρ*, male]. Culver's Root. The rhizome and rootlets of *L. virginica*, now called *Veronica virginica*. Its properties are thought to be due to a glucosid, leptandrin. It is tonic, laxative, and cholagogue, and is used in indigestion and chronic constipation. Dose of the extract, gr. j-ij (0.065-0.2); of the fld. ext., ℥xx-f 3j (1.3-4.0).

Leptomeningitis (*lep-to-men-in-jit'-tis*) [*λεπτός*, slight; *μήνιξ*, membrane; *ιτις*, inflammation]. Inflammation of the pia and arachnoid of the brain or the spinal cord.

Leptorhine (*lep'-tor-in*) [*λεπτός*, thin; *ῥίς*, nose]. Having a slender nose or proboscis.

Leptothrix (*lep'-to-thriks*) [*λεπτός*, thin; *θρίξ*, hair]. A genus of bacteria, the elements of which form straight filaments, often of great length. See *Bacteria*, Table of.

Leptus (*lep'-tus*) [*λεπτός*, thin]. A genus of beetles. **L. autumnalis**, harvest-bug; mower's mite. A parasite that burrows under the skin, causing lesions similar to those of itch.

Lesion (*le'-zhun*) [*lesio*; *laedere*, to hurt]. An injury, wound, or morbid structural change. **L., Focal**, in the nervous system, a circumscribed lesion giving rise to distinctive and localizing symptoms. **L., Initial**, of Syphilis, the chancre. **L., Irritative**, in the nervous system, a lesion exciting the

functions of the part wherein it is situated.

L., Peripheral, a lesion of the nerve-trunks or of their terminations. **L., Primary**, of the Skin, the change in the skin occurring in the developing stage of a skin-disease.

L., Secondary. 1. In the skin, the change occurring in the primary lesion, due to irritation or other causes. It comprises erosions, ulcers, rhagades, squamæ, cicatrices or scars, crusts, and pigmentation. 2. One of the secondary manifestations of syphilis.

Lethal (*le'-thal*) [*λήθη*, forgetfulness]. Deadly; pertaining to or producing death.

Lethargy (*leth'-ar-jy*) [*λήθη*, forgetfulness]. A condition of drowsiness or stupor that cannot be overcome by the will; also, a stage of hypnotism.

Lettuce (*let'-us*). See *Lactucarium*.

Leucemia (*lu-se'-me-ah*). See *Leukemia*.

Leucin (*lu'-sin*) [*λευκός*, white]. $C_6H_{13}NO_2$. A substance formed during pancreatic digestion, and also found in the urine, together with tyrosin, in acute yellow atrophy of the liver. **L.** crystallizes from the urine in the form of yellowish-brown balls.

Leuco- (*lu'-ko-*). For words not found under this form see *Leuko-*.

Leukemia (*lu'-ke'-me-ah*) [*λευκός*, white; *αἷμα*, blood]. A disease of the blood and the blood-making organs, characterized by a permanent increase in the number of white blood-corpuscles and by enlargement of the spleen, the lymphatic glands, and the marrow of bone, together or separately. The etiology is obscure; by some the disease is considered infectious. **L., Lymphatic**, that form associated with enlargement of the lymphatic glands. **L., Myelogenic**, that in which the bone-marrow is involved. **L., Splenic**, that associated with enlargement of the spleen.

Leukemic (*lu'-kem'-ik*, *lu'-ke'-mik*) [*λευκός*, white; *αἷμα*, blood]. Pertaining to leukemia.

Leukocyte (*lu'-ko-sit*) [*λευκός*, white; *κύτος*, cell]. The colorless or white corpuscle of the blood. Leukocytes have ameboid movement and are formed in the lymphadenoid tissue of the spleen, lymphatic glands, intestinal tract, bone-marrow, etc., and probably also in the lymph and blood. Their average diameter is .01 mm. ($\frac{1}{100}$ inch). In normal blood several forms are distinguished: lymphocytes, large uninuclear leukocytes, transitional forms, and multinuclear cells; according to the granules contained in their protoplasm, three varieties are described—the neutrophile, the eosinophile, and the basophile.

Leukocythemia (*lu'-ko-sit'-he'-me-ah*) [*λευκός*, white; *κύτος*, cell; *αἷμα*, blood]. See *Leukemia*.

Leukocytic (*lu'-ko-sit'-ik*) [*λευκός*, white; *κύτος*, cell]. Relating to or characterized by leukocytes.

Leukocytogenesis (*lu·ko·si·to·jen'-es-is*) [*λευκός*, white; *κύτος*, cell; *γενᾶν*, to beget]. The formation of leukocytes.

Leukocytolysis (*lu·ko·si·tol'-is-is*) [*λευκός*, white; *κύτος*, cell; *λύσις*, solution]. The destruction of leukocytes.

Leukocytoma (*lu·ko·si·to'-mah*) [*λευκός*, white; *κύτος*, cell; *δῆμα*, tumor]. A tumor-like mass composed of leukocytes, as the tubercle, the gumma.

Leukocytopenia (*lu·ko·si·to·pe'-ne-ah*) [*λευκός*, white; *κύτος*, cell; *πενία*, poverty]. Diminution of the number of leukocytes in the blood.

Leukocytosis (*lu·ko·si·to'-sis*) [*λευκός*, white; *κύτος*, cell]. An increase in the number of colorless blood-corpuscles in the blood. It is physiologic during digestion and pregnancy; pathologic in certain anemias, especially leukemia, in some of the infectious fevers, in cachexias, and after hemorrhage. **L., Pure**, a leukocytosis in which the increase of white cells affects the multinuclear form.

Leukoderma (*lu·ko·der'-mah*) [*λευκός*, white; *δέρμα*, skin]. A condition of defective pigmentation of the skin, especially a congenital absence of pigment in patches or bands.

Leukolin (*lu·ko'-lin*). See *Quinolin*.

Leukolysis (*lu·kol'-is-is*) [*λευκός*, white; *λύσις*, solution]. The destruction of the colorless corpuscles of the blood.

Leukoma (*lu·ko'-mah*) [*λέκωμα*, whiteness, from *λευκός*, white]. 1. An opacity of the cornea the result of an ulcer, wound, or inflammation, and presenting an appearance of ground glass. 2. The term has been used also for albumin. See *Leukoplakia*.

Leukomain (*lu·ko'-mah-in*) [*λέκωμα*, whiteness, from *λευκός*, white]. The name applied to any one of the nitrogenous bases or alkaloids normally developed by the metabolic activity of living organisms, as distinguished from the alkaloidal bodies developed in dead bodies, and called ptomains. From their chemie affinities leukomains may be divided into two groups: the uric-acid group, comprising adenin, carnin, guanin, heteroxanthin, hypoxanthin, paraxanthin, pseudoxanthin, spermin, xanthin; and the kreatinin group, in which are classed amphikreatinin, crusokreatinin, xanthokreatinin, and others. The appended table is modified from the work of Vaughan and Novy.

TABLE OF LEUKOMAINS.

NAME.	FORMULA.	DISCOVERER.	SOURCE.	PHYSIOLOGIC ACTION.
Amphicreatin.	$C_6H_{10}N_4O_4$	Gautier.	Muscle.	
Carnin.	$C_7H_9N_4O_3$	Weidel.	Meat-extract.	Nonpoisonous.
Crusocreatin.	$C_5H_8N_4O$	Gautier.	Muscle.	
Gerontin.	$C_6H_{14}N_8$	Grandis.	Liver of dog.	Poisonous; paralyzes nerve-centers.
Guanin.	$C_5H_7N_5O$	Unger.	Animal and vegetal tissues; guano.	Nonpoisonous.
Heteroxanthin.	$C_6H_8N_4O_7$	Salomon.	Urine.	
Hypoxanthin.	$C_5H_8N_4O$	Scherer.	Brain, muscle, kidney, liver, spleen, certain plants, etc.	Nonpoisonous; excitomotor.
Methyl-hydan-toin.	$C_4H_8N_4O_2$	Guaratchi and Mosso.	Fresh meat.	
Parareducin.	$C_6H_9N_4O$	Thudichum.	Urine.	
Paraxanthin.	$C_7H_8N_4O_2$	Thudichum, Salomon.	Urine.	Poisonous.
Protarnin.		Miescher.	Spawn of salmon.	
Pseudoxanthin (?)	$C_4H_8N_4O$	Gautier.	Muscle.	
Reducin (?)	$C_{12}H_{24}N_8O_9$	Thudichum.	Urine.	
Salamandarin.	$C_{21}H_{40}N_2O_5$	Zalesky.	Salamander.	Poisonous.
Sarcin or Sarkin.	See <i>Hypoxanthin</i> .			

TABLE OF LEUKOMAINS.—*Continued.*

NAME.	FORMULA.	DISCOVERER.	SOURCE.	PHYSIOLOGIC ACTION.
Spermin.	C_3H_8N (?)	Schreiner.	Semen, testicles; Charcot-Leyden crystals; tissues in leukemia.	Nonpoisonous.
Unnamed.	$C_3H_8NO_2$	Pouchet.	Urine.	
Unnamed.	$C_7H_{12}N_4O_2$	Pouchet.	Urine.	
Unnamed.	$C_{11}H_{24}N_{10}O_4$	Gautier.	Muscle.	
Unnamed.	$C_{12}H_{25}N_{11}O_5$	Gautier.	Muscle.	
Unnamed.	$C_{12}H_{24}N_2O_2(C_{12}H_{24}N_2O_2PS)$	Eliacheff.	Normal urine.	Slightly poisonous.
Unnamed.	$C_{14}H_{28}N_2O_2$	Eliacheff.	Urine in tuberculosis.	Poisonous.
Urethrobromin.	See <i>Paraxanthin</i> .			
Xanthin.	$C_8H_8N_4O_2$	Marcet.	Most animal tissues; certain plants; certain calculi.	Slightly poisonous; muscle stimulant.
Xanthocreatin.	$C_8H_{10}N_4O$	Gautier.	Muscle.	Poisonous.

Leukomatous (*lu - ko' - mat - us*) [*λεῦκωμα*, whiteness]. Having the nature of or affected with leukoma.

Leukomyelitis (*lu - ko - mi - el - i' - tis*) [*λευκός*, white; *μυελός*, marrow]. Inflammation of the white substance of the spinal cord.

Leukonecrosis (*lu - ko - ne - kro' - sis*) [*λευκός*, white; *νέκρωσις*, death]. A form of dry gangrene, the slough having a light color.

Leukonostoc (*lu - ko - nos' - tok*) [*λευκός*, white, and *nostoc*]. A genus of schizomycetes, having its elements disposed in chains and enveloped in a sheath of tough jelly.

Leukopathy (*lu - kop' - ath - e*) **Leukopathia** (*lu - ko - path' - e - ah*) [*λευκός*, white; *πάθος*, suffering]. Any deficiency of coloring-matter; albinism. See *Leukoderma*.

Leukophlegmasia (*lu - ko - fleg - ma' - ze - ah*) [*λευκός*, white; *φλέγμα*, phlegm]. 1. A condition marked by a tendency to dropsy, accompanied by a pale, flabby skin, and general edema of the whole body; solid edema. 2. Phlegmasia alba dolens. L. dolens puerperarum. Phlegmasia alba dolens.

Leukoplakia (*lu - ko - pla' - ke - ah*) [*λευκός*, white; *πλάξ*, surface]. Whitening of a surface. L. buccalis, L. lingualis, a disease characterized by the presence of pearly-white or bluish-white patches on the surface of the tongue, or the mucous membrane of the cheeks, due to a hyperplasia of the epithelium.

Leukorrhea (*lu - kor - e' - ah*) [*λευκός*, white; *ροία*, a flow]. A whitish, mucopurulent discharge from the female genital canal, popularly called "the whites."

Leukorrheal (*lu - kor - e' - al*) [*λευκός*, white; *ροία*, flow]. Of the nature of or pertaining to leukorrhea.

Levant Wormseed. See *Santonica*.

Levator (*le - va' - tor*) [L., a lifter]. 1. That which raises or elevates, as certain muscles having such a function. See *Muscles*, *Table of*.

Levorotatory (*le - vo - ro' - tat - o - re*) [*laevus*, left; *rotare*, to turn]. Causing to turn toward the left hand; applied to substances that turn the rays of polarized light to the left.

Levulinic Acid (*lev - u - lin' - ik*). See *Acid*. **Levulosan** (*lev - u - lo' - san*) [*laevus*, left], $C_6H_{10}O_5$. A carbohydrate prepared by heating levulose.

Levulose (*lev' - u - lōs*) [*laevus*, left], $C_6H_{12}O_6$. Fruit-sugar, the natural sugar of fruits. It is a colorless, syrupy liquid, and rotates the plane of polarized light to the left. It occurs normally in the intestine, and rarely in the urine in disease.

Leyden Jar (*li' - den*) [*Leyden*, a town in Holland]. A glass jar coated within and without with tinfoil, reaching nearly to the neck, and surmounted by a knobbed conductor in connection with the inner coating. It is designed for the temporary accumulation of electricity, with which the inner foil may be charged.

Leyden's Crystals. See *Charcot-Leyden Crystals*.

Lichen (*li' - ken*) [*λεῖχην*, a lichen]. A generic term for a group of inflammatory affections of the skin, in which the lesions consist of solid papules. L. *acuminatus*, a variety

of *L. ruber* in which the papules are acuminate. It is usually very acute and is accompanied by grave constitutional symptoms (rigors, pyrexia, sweats, prostration), and by itching. *L. pilaris*, an inflammatory disease of the hair-follicles, in which a spinous epidermic peg occupies the center of the papule. *L. planus*, an inflammatory skin-disease, with an eruption made up of papules that are broad and angular at the base, flat and apparently glazed on the summit, slightly umbilicated, and of a dull purplish-red color. The papules may be discrete or may coalesce, and itching may be slight or severe. *L. ruber*, a rare skin-disease, with lesions consisting, in the beginning, of discrete, miliary, conical papules, but, as the disease advances, becoming aggregated and forming continuous red, infiltrated, and scaly patches. The whole surface may eventually become involved. The nails of the fingers and toes become affected, being of a dirty-brown color, rough, flaky, and breaking off short. The etiology and pathology are obscure. *L. scrofulosus*, a form occurring in strumous children. The eruption is situated on the trunk, especially upon the back, either diffusely or in patches. The papules are very small, pale, conical, and surmounted by fine scales; they cause no itching, and on fading leave a rather persistent yellowish pigmentation. *L. tropicus*. Prickly heat. Licorice (*lik'-or-is*) [*γλυκίρριζα*, licorice]. See *Glycyrrhiza*.

Lid [*AS.*, *hlid*, lid]. See *Eyelid*.

Lieberkühn, Crypts of. See *Crypt*.

Lien (*li'-en*) [*L.*]. The spleen.

Lienal (*li'-en-al*) [*lien*, the spleen]. Relating to the spleen.

Lienitis (*li-en-i'-tis*) [*lien*, spleen; *itis*, inflammation]. Splenitis.

Lientery (*li'-en-ter-e*) [*λεῖος*, smooth; *έντερον*, intestine]. A form of diarrhea in which the food passes rapidly through the bowels without undergoing digestion.

Life (*lif*) [*AS.*, *lif*, life]. 1. The sum of properties that enables an organism to adapt itself to surrounding conditions. 2. The characteristic phenomena manifested by living beings. 3. The force or principle underlying or causing the phenomena presented by organized beings. *L.*, *Animal*, the manifestations depending directly on the cerebrospinal nervous system and the voluntary muscles, as distinguished from *L.*, *Vegetative*, that is, the functions of digestion, respiration, reproduction, etc. *L.*, *Change of*, that period in the life of a woman at which menstruation ceases; the climacteric period. *L.*, *Expectation of*, the average number of years a person has to live, as calculated from life-tables.

Ligament (*lig'-am-ent*), Ligamentum (*lig-am-en'-tum*) [*ligare*, to bind]. A band of flexible, compact connective tissue connecting the articular ends of the bones, and sometimes enveloping them in a capsule. *L.*, *Acromioclavicular*, a ligament covering the acromioclavicular articulation and extending from the clavicle to the acromial process of the scapula. *L.*, *Alar*. 1. One of the two folds of synovial membrane on either side of the mucous ligament of the knee-joint. 2. See *L.*, *Odontoid*. *L.*, *Annular (of ankle)*, the broad ligament covering the anterior surface of the ankle-joint. *L.*, *Annular (of wrist)*, a strong ligament extending from the trapezium to the ulnar bone, confining the flexor tendons. *L.*, *Arcuate*, one of the arched ligaments extending from the body of the diaphragm to the last rib, and to the transverse process of the first lumbar vertebra. *L.*, *Atlaxoid*, that joining the atlas and the axis. *L.*, *Atlaooccipital*, that joining the atlas and the occiput. *L.* of *Bigelow*. See *L.*, *Iliofemoral*. *L.*, *Broad (of uterus)*, a fold of peritoneum extending laterally from the uterus to the pelvic wall. *L.*, *Capsular*, a heavy fibrous structure surrounding an articulation, and lined by synovial membrane. *L.*, *Check*. See *L.*, *Odontoid*. *L.*, *Conoid*, the inner portion of the coracoclavicular ligament. It is attached to the coracoid process of the scapula and the conoid tubercle of the clavicle. *L.*, *Coracoclavicular*, one extending from the coracoid process of the scapula to the clavicle. *L.*, *Coronary (of liver)*, a peritoneal fold extending from the posterior edge of the liver to the diaphragm. *L.*, *Costocoracoid*, that joining the first rib and the coracoid process of the scapula. *L.*, *Cotyloid*, a ring of fibrocartilaginous tissue at the margin of the acetabulum. *L.*, *Crucial*, *Anterior*, the smaller crucial ligament of the knee, extending from the upper surface of the tibia to the inner surface of the external condyle of the femur. *L.*, *Crucial*, *Posterior*, one attached below to the back part of the depression behind the spine of the tibia, to the popliteal notch, and to the external semilunar fibrocartilage, and above to the inner condyle of the femur. *L.*, *Crural*. See *L.*, *Poupart's*. *L.*, *Flood's*, the glenohumeral ligament. *L.*, *Gimbernat's*, a triangular expansion of the aponeurosis of the external oblique muscle, connected anteriorly to Poupart's ligament and attached to the iliopectineal line. *L.*, *Glenohumeral*, a portion of the coracohumeral ligament, attached to the inner and upper portion of the bicipital groove. *L.*, *Glenoid (of scapula)*, a ring of fibrocartilaginous tissue attached to the rim of the glenoid fossa. *L.*, *Hey's*, also called the femoral ligament; a sickle-shaped expansion

of the fascia lata. **L., Iliofemoral**, a strong ligament extending from the anterior inferior iliac spine to the lesser trochanter and the intertrochanteric line. **L., Interclavicular**, one joining the sternal extremities of the clavicles and the sternum. **L., Lateral**, (*of liver*), one of the peritoneal folds between the sides of the liver and the inferior surface of the diaphragm. **L., Odontoid**, any one of the broad, strong ligaments arising on either side of the apex of the odontoid process and connecting the atlas with the skull. **L. patellæ**, a strong fibrous structure, extending from the tubercle of the tibia upward to become the tendon of the quadriceps extensor muscle; it embraces the patella. **L. pectinatum**, the spongy tissue at the junction of the cornea and sclera in the sinus of the anterior chamber of the eye. It forms the root of the iris. **L., Poupart's**, the ligament extending from the anterior superior spine of the ilium to the spine of the pubis and the pectineal line. It is the lower portion of the aponeurosis of the external oblique muscle. **L., Pterygomaxillary**, one joining the apex of the internal pterygoid plate of the sphenoid bone and the posterior extremity of the internal oblique line of the lower jaw. **L., Pubic, Inferior**, a triangular ligament, extending from the symphysis pubis to the rami of the pubic bones. **L., Rhomboid**, one joining the cartilage of the first rib and the tuberosity of the clavicle. It is a part of the costoclavicular ligament. **L., Round** (*of hip*). See **L. teres**. **L., Round** (*of liver*), a fibrous cord running from the umbilicus to the notch in the anterior border of the liver. It represents the remains of the obliterated umbilical vein. **L., Round** (*of uterus*), a ligament running from the anterior surface of the cornu of the uterus through the inguinal canal to the mons veneris. **L., Sacrosciatic, Great**, a ligament extending from the sacrum, coccyx, and inferior iliac spine to the tuberosity of the ischium. **L., Stylohyoid**, a fibrous cord extending from the apex of the styloid process of the temporal bone to the lesser cornu of the hyoid bone. **L., Stylomaxillary**, a ligament joining the styloid process of the temporal bone and the inferior surface of the posterior margin of the ramus of the inferior maxilla. **L., Subpubic**. See **L., Pubic, Inferior**. **L., Suspensory** (*of crystalline lens*), the zonule of Zinn. **L. teres**, a rounded fibrous cord attached to the center of the articular surface of the head of the femur, and extending to the margin of the cotyloid notch of the acetabulum. **L., Transverse** (*of atlas*), one attached to two small tubercles on the inner surface of the atlas, and surrounding the odontoid process of the axis. **L., Trans-**

verse (*of hip joint*), one extending across the cotyloid notch of the acetabulum. **L., Transverse** (*of knee joint*), one extending from the anterior margin of the external semilunar fibrocartilage to the anterior extremity of the internal fibrocartilage. **L., Trapezoid**, the anterior or external portion of the coracoclavicular ligament, extending from the upper surface of the coracoid process of the scapula, to the under surface of the clavicle. **L., Triangular** (*of the urethra*), a tendinous band of triangular shape, attached by its apex to the reflected portion of Poupart's ligament, and passing inward beneath the spermatic cord and behind the inner pillar of the external abdominal ring, to join the tendon of the opposite side. **L., Winslow's**, the posterior ligament of the knee-joint. **L., Y-shaped** (*of Bigelow*), the iliofemoral ligament.

Ligamentous (*lig-am-en'-tus*) [*ligare*, to bind]. Of the nature of, or pertaining to, a ligament.

Ligation (*li-ga'-shun*) [*ligare*, to tie]. The operation of tying, especially of tying arteries.

Ligature (*lig'-at-ür*) [*ligare*, to tie]. 1. A cord or thread used for tying about arteries or other parts. 2. **Ligation**. **L., Double**, the application of two ligatures to a vessel, between which it is divided. **L., Elastic**, a narrow band or thread of india-rubber applied tightly to a part so as to destroy the tissues and by compression to lead to separation. It is used in the treatment of hemorrhoids, anal fistula, and in the removal of pedunculated growths. **L., Erichsen's**, one consisting of a double thread, one-half of which is white, the other half black; it is used in the ligation of nævi. **L., Intermittent**, a tourniquet applied above a poisoned wound to interrupt the blood-current; it is occasionally relaxed to allow of renewal of the circulation. **L., Lateral**, partial occlusion of the lumen of a vessel by a loose ligature. **L., Provisional**, a ligature applied during an operation, with the intention of removing it before the completion of the operation.

Light (*ht*) [*AS., leóht, light*]. Ethereal waves of the vibrational frequency that gives rise to the sensation of vision when the rays impinge upon the retina. **L., Axial**, light-rays that are parallel to each other and to the optic axis. **L., Central**. See **L., Axial**. **L., Diffused**, that reflected simultaneously from an infinite number of surfaces, or that which has been scattered by means of a concave mirror or lens. **L., Oblique**, light falling obliquely on a surface. **L., Reflected**, light thrown back from an illuminated object. **L., Refracted**, light-rays that have passed through an object and have been bent from their original course. **L.-sense**, the faculty by which different

degrees of light or brightness may be distinguished. **L.**, **Transmitted**, the light passing through an object.

Lighterman's Bottom. Inflammation of the bursa over the tuberosity of the ischium, from much sitting.

Lightning (*lit'-ning*) [AS., *leóht*, light]. A luminous atmospheric discharge of electricity.

L. Pains, the lancinating pains of locomotor ataxia, coming on and disappearing with lightning-like rapidity.

Lignum (*lig'-num*) [L.]. Wood. **L. benedictum**, guaiac-wood. **L. cedrium**, cedar-wood. **L. vitæ**. See *Guaiacum*.

Ligusticum (*li-gus'-tik-um*) [*λιγυστικός*, Ligurian]. See *Lorage*.

Lily of the Valley. See *Convallaria majalis*.

Limb (*lim*) [AS., *lim*, a limb]. 1. One of the extremities attached to the sides of the trunk and used for prehension or locomotion. 2. An elongated structure resembling a limb, as the limbs of the internal capsule.

Limbus (*lim'-bus*) [L.]. A border; the circumferential edge of any flat organ or part. **L. corneæ**, the edge of the cornea at its junction with the sclerotic coat. **L. laminæ spiralis**, the spiral membranous cushion, at the border of the osseous spiral lamina of the cochlea. **L. luteus**. See *Macula lutea*.

Lime (*lim*) [Pers., *limû*, a lemon]. The fruit of several species of Citrus, as *C. limetta*. **L.-juice**, the juice of the lime.

Lime (*lim*) [AS., *lim*, cement]. Calcium oxid, CaO (quicklime), calcium hydrate, Ca(OH). Calcium oxid (quicklime) has a great affinity for water and for CO₂. On contact with the former, slaked lime is formed, with the evolution of heat. On living tissues it acts as a caustic. **L.**, **Chlorinated**, calx chlorata (U. S. P., B. P.), the chlorid of lime of commerce. It is not a distinct chemic compound; its chief constituent, and the one on which its disinfectant properties depend, is calcium hypochlorite,

which liberates chlorin. **L.**, **Milk of**, a milky fluid consisting of calcium hydrate suspended in water. **L.**, **Quick**, **L.**, **Slaked**, common terms for lime. **L.-water**, a solution of calcium hydrate in water. It is used as an antacid.

Liminal (*lim'-in-al*) [*λυμην*, threshold]. Pertaining to the threshold, especially pertaining to the lowest limit of perception.

Limitans, Limiting (*lim'-it-ans*, *lim'-it-ing*) [*λυμην*, threshold]. Bounding. **L. Membrane**, External, the thin layer between the outer nuclear layer of the retina and that of the rods and cones. **L. Membrane**, Internal, in the eye, the inner layer of the retina.

Limo (*li'-mo*) [L.]. Lemon. The fruit of Citrus limonum, a tree of the order Rutaceæ. The rind (**Limonis cortex**) yields an essential oil (**Oleum limonis**, C₁₀H₁₆), and a glucosid, hesperidin (C₂₇H₃₆O₁₃). The pulp contains a large amount of citric acid. The juice (**Limonis succus**) is refrigerant and antiscorbutic. Locally it has been used in pruritus, sunburn, and as a gargle, in diphtheria. **L.**, **Oleum**. Dose m-j-v (0.065-0.32). **L.**, **Syrupus** (B. P.) is used as a refrigerant and vehicle.

Limosis (*lim'-o'-sis*) [*λυμός*, hunger]. 1. Unnatural appetite. 2. A disease distinguished by depraved appetite.

Limotherapy (*lim'-o-ther'-ap-e*) [*λυμός*, hunger; *θεραπεία*, treatment]. The treatment of disease by partial or total deprivation of food. It has been used in the treatment of aneurysm.

Lincture (*link'-tūr*) [*lingere*, to lick]. A medicine to be taken by licking; an electrolytic.

Line (*lin*) [*linea*, a line]. 1. Extension of dimension having length, but neither breadth nor thickness. 2. The $\frac{1}{2}$ part of an inch. 3. In anatomy, anything resembling a mathematic line in having length without breadth or thickness; a boundary or guide-mark. A table of lines is appended.

TABLE OF LINES OR LINEÆ.

NAME.	DESCRIPTION.
Alba.	1. A tendinous raphe extending in the median line of the abdomen from the pubes to the ensiform cartilage. It is formed by the blending of the aponeuroses of the oblique and transversalis muscles. 2. Hunter's line, the anterior peduncles of the pineal gland.
Aibicantes.	Glistening white lines in either iliac region of the abdomen seen in distention of the abdomen from pregnancy, ascites, or tumors.
Alveolobasilar.	A line joining the basion and the alveolar point.
Alveolonasal.	A line joining the nasal and alveolar points.

TABLE OF LINES OR LINEÆ.—*Continued.*

NAME.	DESCRIPTION.
Aspera.	A rough longitudinal ridge on the posterior surface of the middle third of the femur, dividing below into two and above into three ridges.
Auriculobregmatic.	A line passing from the auricular point to the bregma, and dividing the preauricular from the postauricular part of the cranium.
Axillary, anterior and posterior.	Vertical lines extending downward from the axilla on the side of the trunk.
Base-line.	A line running backward from the infraorbital ridge through the middle of the external auditory meatus, and prolonged to the middle line of the head posteriorly.
Basiobregmatic.	The line joining the basion and the bregma.
Baudelocque's.	The external conjugate diameter of the pelvis.
Biauricular.	The line separating the anterior from the posterior portion of the skull. It extends from one auditory foramen over the vertex to the other.
Blue.	The blue line at the dental margin of the gums in chronic lead-poisoning.
Bryant's.	See <i>L., Test.</i>
Camper's.	A line running from the external auditory meatus to a point just below the nasal spine.
Clapton's.	A green line at the margin of the gums, with a similar green stain extending for some distance on the teeth, in chronic copper-poisoning.
Corrigan's.	A purple line at the junction of the teeth and gums in chronic copper-poisoning.
Costoarticular.	A line drawn between the sternoclavicular articulation and the point of the eleventh rib.
Costoclavicular.	See <i>L., Parasternal.</i>
Curved, inferior (of the ilium).	A line extending from the upper part of the anterior inferior spinous process of the ilium, and terminating at the middle of the great sciatic notch.
Curved, inferior (of the occipital bone).	A ridge extending transversely across the outer surface of the occipital bone a short distance below the superior curved line.
Curved, middle (of the ilium).	A line commencing about an inch or an inch and a half behind the anterior superior spine of the ilium and arching backward and downward to the upper margin of the great sciatic notch.
Curved, superior (of the ilium).	A line commencing about two inches in front of the posterior extremity of the crest of the ilium and curving downward and forward toward the posterior part of the great sciatic notch.
Curved, superior (of the occipital bone).	A semicircular line, passing outward and forward from the external occipital protuberance.
Of demarcation.	A line of division between healthy and gangrenous tissues.
Ellis's curved line.	The curved line followed by the upper border of a pleuritic effusion or a hydrothorax.
Embryonic.	The primitive trace in the center of the germinal area of the ovum.
Eminens (of the cricoid cartilage).	A mesal ridge on the dorsal half of the cricoid cartilage.
Eminens (of the patella).	A ridge on the posterior surface of the patella, dividing that surface into two unequal parts, the outer of which is the larger.
Facial.	1. A straight line tangential to the glabella and some point at the lower portion of the face. 2. See <i>L., Camper's.</i>
Of fixation.	An imaginary line drawn from the object viewed through the center of rotation of the eye.

TABLE OF LINES OR LINEÆ.—*Continued.*

NAME.	DESCRIPTION.
Fraunhofer's.	The black lines that cross the solar spectrum. They are produced by the light from the lower portions of the solar surface passing through certain incandescent metallic vapors in the luminiferous envelop of the sun, and through the aqueous vapor and gases of the earth's atmosphere.
Frohmann's.	Transverse lines or striæ appearing on the axis-cylinder of medullated nerve-fibers, near the nodes of Ranvier, after the fibers have been stained with silver nitrate.
Genal (<i>Jadelot's</i>).	A line seen in the faces of children, in certain diseases, running downward from the region of the malar bone to join the nasal line. See <i>Jadelot's Lines</i> .
Gingival (<i>Burton's</i>).	A reddish streak or margin at the reflected edge of the gums.
Of Haller.	See <i>L. splendens</i> .
Hilton's.	A white line marking the point of junction of the skin of the perineum with the mucosa of the anus, and also the point of separation of the external from the internal sphincter.
Holden's.	A sulcus below the fold of the groin, starting from the femoroscretal furrow, and fading away between the great trochanter and the anterior superior iliac spine. It crosses the middle of the capsule of the hip.
Iliopectineal.	The bony ridge marking the brim of the true pelvis, situated partly on the ilium and partly on the pubis.
Incremental (<i>Saller's</i>).	Curved line in dentine, supposed to indicate the laminar structure, and to correspond to the successive laminæ or strata of dentine.
Intertrochanteric, anterior.	A line upon the anterior surface of the femur, separating the neck and shaft, extending between the tubercle and a point close to and in front of the lesser trochanter.
Intertrochanteric, posterior.	A ridge on the posterior surface of the femur, extending between the greater and lesser trochanters.
Jadelot's.	See <i>Jadelot's Lines</i> .
Kirchoff's.	See <i>L., Fraunhofer's</i> .
Ligar's.	Two lines, one joining the posterior iliac spine at a point midway between the tuber ischii and the great trochanter; the other from the posterior iliac spine to the inner point of trisection of a line between the tuber ischii and the trochanter; the upper point of trisection of the first indicates the emergence of the gluteal artery; the middle of the second, the spot where the sciatic artery leaves the pelvis.
Mamillary.	A vertical line passing through the center of the nipple.
Mylohyoidean.	See <i>L., Oblique, Internal (of the inferior maxilla)</i> .
Nasobasilar.	The line drawn through the basion and the nasal point.
Nélaton's.	A line drawn from the anterior superior spine of the ilium to the most prominent part of the tuberosity of the ischium. In dislocation of the femur backward the trochanter is always found above this line.
Nuchal, inferior.	The inferior curved line of the occiput.
Nuchal, median.	The external occipital protuberance.
Nuchal, superior.	The superior curved line of the occiput.
Oblique (<i>of the fibula</i>).	A prominent ridge on the internal surface of the shaft of the fibula, commencing above at the inner side of the head, and terminating in the interosseous ridge at the lower fourth of the bone.
Oblique (<i>of the radius</i>).	A prominent ridge running from the lower part of the bicipital tuberosity, downward and outward, to form the anterior border of the bone.
Oblique (<i>of the thyroid cartilage</i>).	A line extending downward and outward from the tubercle of the thyroid cartilage.

TABLE OF LINES OR LINEÆ.—*Continued.*

NAME.	DESCRIPTION.
Oblique (<i>of the tibia</i>).	A rough ridge that crosses the posterior surface of the tibia obliquely downward from the back part of the articular facet for the fibula to the internal border.
Oblique, external (<i>of the inferior maxilla</i>).	A prominent ridge on the external surface of the inferior maxilla just below the mental foramen, from which it runs outward, upward, and backward to the anterior margin of the ramus.
Oblique, internal (<i>of the inferior maxilla</i>).	A ridge on the internal surface of the lower jaw, commencing at the posterior portion of the sublingual fossa, continuing upward and outward so as to pass just below the last two molar teeth.
Ogston's.	The line for the tendon of the adductor magnus, running from the tubercle of the femur to the intercondyloid notch.
Parasternal.	A line midway between the nipple-line and the border of the sternum.
Pectineal.	The portion of the iliopectineal line that is formed by the pubic bone.
Primitive.	The primitive streak of the embryo.
Profile (<i>of Camper</i>).	See <i>L., Camper's</i> .
Quadrato.	An eminence on the femur commencing about the middle of the posterior intertrochanteric line, and descending vertically for about two inches along the posterior surface of the shaft.
Respiratory.	The line connecting the bases of the upward strokes in a tracing of the pulse.
Roser's.	Same as <i>L., Nélaton's</i> .
Salter's.	See <i>L., Incremental</i> .
Scapular.	A vertical line downward from the lower angle of the scapula.
Semicircular (<i>Douglas's</i>).	The curved lower edge of the internal layer of the aponeurosis of the internal oblique muscle of the abdomen, where it ceases to cover the posterior surface of the rectus muscle.
Semilunar (<i>of Spigelius</i>).	A curved tendinous condensation of the aponeurosis of the external oblique muscle of the abdomen, running along the outer border of the rectus abdominis.
Of sight.	An imaginary line drawn from the object viewed to the center of the pupil.
Splendens (<i>of Haller</i>).	A longitudinal fibrous band extending along the middle line of the anterior surface of the spinal pia mater.
Sternal.	The median line of the sternum.
Sternomastoid.	A line drawn from a point between the two heads of the sternomastoid muscle to the mastoid process.
Supraorbital.	A line extending horizontally across the forehead immediately above the root of the external angular process of the frontal bone.
Test (<i>Bryant's line</i>).	A line for detecting shortening of the neck of the femur. If two lines are drawn to meet at right angles, one of them backward from the anterior superior spinous process of the ilium, and the other upward from the top of the trochanter major, the latter is the test-line. Its length is to be compared with the same line on the uninjured side.
Thompson's.	A red line of vascular tissue along the margin of the gums frequently noticeable in pulmonary tuberculosis.
Transverse (<i>of the abdomen</i>).	The tendinous intersections in the course of the rectus abdominis muscle.
Trapezoid.	The line of attachment of the trapezoid ligament on the inferior surface of the outer portion of the clavicle.
Virchow's.	The line extending from the root of the nose to the lambda.
Visual.	An imaginary line, drawn from a point looked at, through the nodal point of the eye, to the macula lutea.

Linea (*lin'-e-ah*) [L.]. A line. See *Line*.
Linear (*lin'-e-ar*) [*linea*, a line]. Resembling or pertaining to a line.

Ling's System. A method of treatment of disease by gymnastic and other rhythmic movements of the body, employed by Ling, a Swedish physician; kinesiotherapy.

Lingua (*ling'-wah*) [L.]. The tongue. *L. frænata*, tongue-tie. *L. geographica*, the geographic tongue. See *Tongue*.

Lingual (*ling'-wal*) [*lingua*, tongue]. Pertaining to or shaped like the tongue. *L. Artery*. See *Arteries*, *Table of*. *L. Lobule*. See *Subcalcarine Convolution*. *L. Nerve*. See *Nerves*, *Table of*.

Lingula (*lin'-gu-lah*) [dim. of *lingua*, a tongue]. A small lobule between the valve of Vieussens and the central lobule of the cerebellum. It is also called the linguetta laminosa. *L. mandibularis*, the prominent, thin scale of bone partly surrounding the inferior dental foramen of the lower jaw. *L. of Wrisberg*, the connecting fibers of the motor and sensory roots of the trifacial nerve.

Lingulate (*ling'-gu-lat*) [*lingula*, dim. of *lingua*, tongue]. Tongue-shaped.

Linimentum (*lin-im-en'-tum*) [*linere*, to smear]. A liquid intended for application to the skin by gentle friction. The following are official in the U. S. P.: *L. ammoniac*, or volatile liniment, *L. belladonnæ*, *L. calcis*, or carron-oil, *L. camphoræ*, or camphorated oil, *L. chloroformi*, *L. saponis*, *L. saponis mollis*, or tinctura saponis viridis, *L. sinapis compositum*, *L. terebinthinæ*. *L.*, St. John Long's, linimentum terebinthinæ aceticum (B. P.).

Linolein (*lin-o'-le-in*) [*linum*, flax; *oleum*, oil]. The neutral fat contained in linseed oil, and to which its drying property is due.

Linseed (*lin'-sed*). See *Linum*.

Lint [*linum*, flax]. A loosely woven or partly felted mass of broken linen-fibers, made by scraping and picking linen cloth. It is used as a dressing for wounds. *L.*, Common, lint that is twilled on one side and woolly on the other. In the spreading of an ointment the twilled side is used. *L.*, Patent, English charpie; lint that is scraped on both sides, a soft finish being thus given to the two surfaces.

Linum (*li'-num*) [L.]. Flaxseed; linseed. The seed of *L. usitatissimum*, a plant of the order *Lineæ*, containing a fatty substance, linolein, which is the glycerid of linoleic acid. Flaxseed is a demulcent, emollient, and expectorant, useful in inflammations of mucous membranes. *L.*, *Cataplasma* (B. P.), a poultice made from linseed meal. *L. catharticum*, is an active purgative and vermifuge. *L. Farina* (B. P.), flaxseed-meal, used as a poultice. *L.*, *Infusum*, flaxseed-

tea, unof. Dose indefinite. *L.*, *Oleum*, the fixed oil of flaxseed, a glycerid of linoleic acid. Dose f ʒ ss-ij (16.0-64.0). *L.*, *Semen*, linseed or flaxseed.

Liodermia (*li-o-der'-me-ah*) [*λειος*, smooth; *δέρμα*, skin]. A condition of abnormal smoothness and glossiness of the skin.

Liomyoma (*li-o-mi-o'-mah*) [*λειος*, smooth; *μῦς*, muscle; *δῶμα*, tumor]. A tumor composed of unstriped muscular tissue.

Lip [AS., *lippa*, lip]. 1. One of the two fleshy folds surrounding the orifice of the mouth. 2. One of the labia majora. See *Labium*.

Lipacidemia (*lip-as-id-e'-me-ah*) [*λίπος*, fat; *acidus*, sour; *αἷμα*, blood]. The presence of fatty acids in the blood.

Lipaciduria (*lip-as-id-u'-re-ah*) [*λίπος*, fat; *acidus*, acid; *urina*, urine]. The presence of fatty acids in the urine.

Lipæmia (*lip-e'-me-ah*). See *Lipemia*.

Liparin (*lip'-an-in*) [*λίπος*, fat]. A substitute for cod-liver oil, consisting of pure olive-oil and six per cent. of oleic acid. The dose is from 2 to 6 tablespoonfuls daily.

Liparocele (*lip'-ar-o-sel*) [*λίπαρος*, fat; *κήλη*, a tumor]. A fatty tumor or cyst; a hernia containing fatty tissue.

Lipemia (*lip-e'-me-ah*) [*λίπος*, fat; *αἷμα*, blood]. The presence of an emulsion of fine oil-globules in the blood, a condition sometimes noted in diabetes.

Lipoma (*lip-o'-mah*) [*λίπος*, fat; *δῶμα*, a tumor]. A fatty tumor.

Lipomatosis (*lip-o-mat-o'-sis*) [*λίπος*, fat; *δῶμα*, tumor]. A general deposition of fat; obesity.

Lipomatous (*lip-o'-mat-us*) [*λίπος*, fat; *δῶμα*, tumor]. Of the nature of a lipoma.

Lipomyxoma (*lip-o-mix-o'-mah*) [*λίπος*, fat; *μῖξα*, mucus; *δῶμα*, tumor]. A myxoma combined with fatty tissue.

Lipothymia (*lip-o-thi'-me-ah*) [*λείπειν*, to leave; *θῦμις*, mind]. Faintness.

Lippitude (*lip'-e-tud*). **Lippitudo** (*lip-e-tu'-do*) [*lippus*, bleary-eyed]. The state of being bleary-eyed, a condition marked by ulcerative marginal blepharitis.

Lipuria (*lip-u'-re-ah*) [*λίπος*, fat; *urina*, urine]. The presence of fat in the urine.

Liquefaction (*lik-we-fak'-shun*) [*liquidus*, liquid; *facere*, to make]. The process of changing or being changed into a liquid. *L.-necrosis*. See *Necrosis*.

Liquefactive (*lik-we-fak'-tiv*) [*liquefacere*, to render liquid]. Pertaining to, causing, or characterized by liquefaction.

Liquescent (*lik-wes'-ent*) [*liqueescere*, to become liquid]. Becoming, or tending to become, liquid.

Liqueur (*lik-ur*) [Fr.]. An aromatic alcoholic drink.

Liquid (*lik'-wid*) [*liquere*, to melt]. 1. Fluid; flowing. 2. A substance that flows readily and takes the shape of the containing vessel.

Liquidambar (*lik'-wid-am'-bar*) [*liquidus*, liquid; *ambar*, from Ar., *ambar*, ambergris]. A genus of trees of the Hamamelaceæ. *L. altingia* and *L. orientalis*, afford a portion of commercial styrax. *L. styraciflua*, of North America (sweet-gum, bilsted, copalm) contains a stimulant gum, and is useful in diarrheas, coughs, and coids.

Liquor (*li'-kuor* or *lik'-or*) [L.]. 1. Any liquid. 2. An aqueous solution of a non-volatile substance. *L. amnii*, the liquid contained in the amniotic sac. *L. cotunnii*, the perilymph of the internal ear. *L. folliculi*, the fluid filling the follicle or space about the developing ovum in the ovary. *L. sanguinis*, the blood-plasma.

Liquorice (*lik'-or-is*). See *Glycyrrhiza*.

Listranc's Amputation [*Listranc*, a French surgeon]. A disarticulation of the metatarsal bones from the tarsus. *L.'s Tubercle*, a rough spot on the anterior surface of the first rib near the superior border. It serves for the attachment of the scalenus anticus muscle.

Lisping (*lisp'-ing*) [AS., *zispian*, to lisp]. A defect of speech, in which sibilant letters are sounded like linguals, especially *s* as *th*.

Lissauer's Tract [*Lissauer*, a German scientist]. A group of fine nerve-fibers in the spinal cord lying ventrad and dorsad of the entrance of the dorsal roots.

Listerine (*lis'-ter-in*) [*Lister*, an English physician]. A proprietary antiseptic preparation said to contain thymol, eucalyptus, baptisia, gaultheria, mentha arvensis, benzoic and boric acids.

Listerism (*lis'-ter-izm*). A general name for the antiseptic and aseptic treatment of wounds according to the principles first enunciated by Lister, an English physician.

Listing's Law [*Listing*, a German scientist]. See *Law*. *L.'s Reduced Eye*. See *Eye*.

Liter (*le'-ter*) [*litra*, a pound]. The unit of capacity in the metric system, equal to 0.88036 of an imperial quart, or 1.056 U. S. quarts; it is the volume of one kilogram of water at its maximum density.

Lithagogue (*lith'-ag-og*) [*lithos*, stone; *agwōs*, leading]. 1. Expelling calculi. 2. Any agent tending to expel calculi from the bladder.

Litharge (*lith'-arj*). See *Plumbum*.

Lithate (*lith'-at*) [*lithos*, stone]. A salt of lithic (uric) acid; a urate.

Lithectasy (*lith'-ek'-tas-e*) [*lithos*, stone; *ektrasis*, a stretching out]. Dilatation of the urethra and neck of the bladder for the removal of calculi.

Lithemia (*lith'-el-me-ah*) [*lithos*, stone; *aima*,

blood]. A condition in which, owing to defective metabolism of the nitrogenous elements, the blood becomes charged with dete-rious substances, principally, perhaps, of the uric-acid group, although their exact chem-ic nature is not determined.

Lithemic (*lith-em'-ik*, *lith-el'-mik*) [*lithos*, stone; *aima*, blood]. Pertaining to or suffer-ing from lithemia.

Lithia (*lith'-e-ah*) [*lithos*, a stone]. Li_2O . Lithium oxid. *L. Water*, mineral water containing lithium salts in solution.

Lithiasis (*lith'-e-az*) [*lithos*, stone]. The formation of calculi in the body.

Lithic (*lith'-ik*) [*lithos*, stone]. 1. Pertain-ing to calculi. 2. Pertaining to lithium. *L. Acid*. See *Acid*, *Uric*.

Lithium (*lith'-e-um*) [*lithos*, stone]. Symbol Li; atomic weight 7; quantivalence 1. A soft, silver-white metal belonging to the group of alkalis. It is the lightest solid element, having a specific gravity of 0.535. The salts of L. are used in medicine for their solvent power of uric acid, with which they form easily soluble salts. They are, there-fore, employed in rheumatic and gouty affec-tions. *L. benzoas*. Dose gr. v-xxx (0.32-2.0). *L. bromidum* has the action of the bromids. Dose gr. xv-xxx (1.0-2.0). *L. carbonas*. Dose gr. v-xx (0.32-1.0). *L.-carmin*, a solution of carmin in lith-ium carbonate, used as a stain for tissue. *L. citras*. Dose gr. x-xxx (0.65-2.0). *L. citras effervescens*. Dose \mathfrak{z} (4.0). *L. salicylas*. Dose gr. xxx-xl (1.3-2.0).

Lithocenosia (*lith'-e-sen'-e-sis*) [*lithos*, stone; *kenosis*, evacuation]. The extraction of the fragments of calculi that have been crushed.

Lithoclast (*lith'-o-klast*). See *Lithotrite*.

Lithoclysmia (*lith'-o-klic'-me-ah*) [*lithos*, stone; *κλύσμα*, clyster]. An injection of solvent liquids into the bladder for the re-moval of calculi.

Lithodialysis (*lith'-o-di-ah'-is-is*) [*lithos*, stone; *διαλύειν*, to dissolve]. 1. The solu-tion of calculi in the bladder. 2. The opera-tion of breaking a vesical calculus, previous to its removal.

Litholapaxy (*lith'-o-ap-ak'-s*) [*lithos*, stone; *λάπαξις*, removal]. An operation for crush-ing a stone in the bladder and removing the fragments at the same sitting.

Lithology (*lith'-o-l-og-ee*) [*lithos*, stone; *λογία*, treatise]. The science of the nature and treatment of calculi.

Litholysis (*lith'-o-l-is-is*) [*lithos*, stone; *λύσις*, solution]. See *Lithodialysis*.

Lithonephrotomy (*lith'-o-nel'-o-l'-o-mē*) [*lithos*, stone; *νεφρος*, kidney; *τομή*, a cut-ting]. Incision of the kidney for the re-moval of a renal calculus.

Lithopedion (*lith'-o-pel'-de-on*) [*lithos*, stone;

παιδιον, child]. A retained fetus that has undergone calcareous infiltration.

Lithophone (*lith'-o-fon*) [*λίθος*, stone; *φωνή*, sound]. An instrument for detecting by sound the presence of calculi in the bladder.

Lithoscope (*lith'-o-skop*) [*λίθος*, stone; *σκοπεῖν*, to examine]. An instrument for the detection and examination of calculi in the bladder.

Lithosis (*lith'-o-sis*) [*λίθος*, stone]. Grinder's lung, a diseased condition of the lung caused by the inhalation and deposition in the lung-tissue of particles of silica or aluminium silicate.

Lithotome (*lith'-o-tom*) [*λίθος*, stone; *τομή*, a cutting]. A cutting-instrument for use in lithotomy.

Lithotomist (*lith'-o'-o-mist*) [*λίθος*, stone; *τομή*, a cutting]. A surgeon who performs lithotomy.

Lithotomy (*lith'-o'-o-me*) [*λίθος*, stone; *τομή*, a cutting]. Incision into the bladder to remove a calculus. **L.**, **Bilateral**, a lithotomy performed by a curved transverse incision just in front of the rectum. **L.**, **Lateral**, one in which the incision is made in front of the rectum and to the left of the raphe. **L.**, **Median** or **Marian**, one in which the incision is made in the median line in front of the anus. **L.**, **Mediolateral**, the perineal incision is made in the median line, and the prostatic incision laterally. **L.** **Position**, a position in which the patient rests on his back with the thighs flexed on the abdomen and the legs flexed on the thighs, the knees being widely abducted. **L.**, **Suprapubic**, lithotomy in which the incision is made above the pubis, at a point where the bladder is not covered by peritoneum. **L.**, **Vaginal**, one in which the incision is through the vaginal wall.

Lithotripsy (*lith'-o-trip-se*) [*λίθος*, stone; *τριβεῖν*, to rub]. The operation of crushing calculi in the bladder.

Lithotripter (*lith'-o-trip'-tor*) [*λίθος*, stone; *τριβεῖν*, to rub]. An instrument for crushing calculi in the bladder.

Lithotrite (*lith'-o-trit*) [*λίθος*, stone; *τερερε*, to rub]. An instrument for crushing a vesical calculus.

Lithotritry (*lith'-o'-rit-e*) [*λίθος*, stone; *τερερε*, to rub]. The process of crushing a stone in the bladder, with the lithotrite, into fragments small enough to pass through the urethral canal.

Lithous (*lith'-us*) [*λίθος*, stone]. Having the nature of a stone.

Lithuresis (*lith'-u-re'-sis*) [*λίθος*, stone; *ὀρρηξις*, urination]. The voiding of small calculi with the urine.

Lithuria (*lith'-u'-re-ah*) [*λίθος*, stone; *urina*, urine]. A condition marked by excess of lithic acid, or its salts, in the urine.

Litmus (*lit'-mus*) [Dutch, *lak*, lac; *more*, pulp. A blue pigment obtained from *Roccella tinctoria*, a lichen. It is employed for determining the presence of acids and alkalies. **L. Paper, Blue**, unsized paper steeped in a solution of litmus; it turns red on contact with acid solutions. **L. Paper, Red**, unsized paper steeped in a solution of litmus colored red with acid; it turns blue on contact with alkaline solutions.

Litter (*lit'-er*) [*lectus*, a couch]. A stretcher or couch with handles for carrying the sick or wounded.

Litten's Sign. See *Diaphragmatic Phenomenon*.

Littre's Colotomy, the making of an opening into the colon through the left iliac region. **L.'s Glands.** See *Gland.* **L.'s Hernia.** See *Hernia*.

Livedo (*liv'-e'-do*) [*livere*, to grow black]. Same as *Lividity*.

Liver (*liv'-er*) [*AS.*, *lifer*, liver]. The largest gland in the body, situated on the right side of the abdominal cavity, just below the diaphragm, and forming an appendage of the digestive tract. Its functions are: the secretion of bile; the formation and storage of glycogen; the production, at least at a certain period of development, of blood-corpuscles; the destruction of blood-corpuscles; the formation of a large quantity of urea; the retention and destruction of certain poisonous substances absorbed from the intestinal tract. Anatomically, it consists of five lobes—the right, the left, the lobus Spigelii, the lobus quadratus, and the lobus caudatus. These lobes are made up of lobules or acini, and these again of hepatic cells, capillaries, arteries, veins, lymphatics, and biliary channels, each lobule being surrounded by connective tissue. The weight of the liver is between 50 and 60 ounces. **L.**, **Albuminoid** or **Amyloid**, one the seat of amyloid degeneration. **L.**, **Beavertail**, one the left lobe of which in form resembles a beaver's tail. **L.**, **Biliary Cirrhotic**, one the seat of chronic inflammation, the result of obstruction and distention of the bile-ducts. **L.**, **Cirrhotic**, one the seat of chronic inflammation, with overgrowth of the connective tissue and atrophy of the parenchyma. **L.**, **Fatty**, one with marked fatty infiltration and degeneration. **L.**, **Hobnail.** See *Hobnail Liver*. **L.**, **Gin-drinkers'**, the liver of atrophic cirrhosis. **L.**, **Nutmeg**, a condition of the liver occurring in heart-disease, fatty infiltration, and amyloid disease. The surface of a section has a peculiar mottled appearance, the center of the lobules being dark, the periphery light in color. **L.**-**spot**, chloasma. **L.**, **Syphilitic**, one the seat of gummata, which on healing, leave scars, or of a diffuse inflammation that

may lead to cirrhosis, with atrophy or, especially in hereditary syphilis, enlargement. **L., Tightlace**, one in which the right lobe is thickened vertically from compression, marked by the ribs, and atrophic from constant pressure by tight stays. **L., Waxy**. See **L., Amyloid**. **L.-wort**. See **Hepatica**.

Livid (*liv'-id*) [*livere*, to be dark]. Discolored from the effects of congestion or contusion; black and blue; pale lead-color.

Lividity (*liv-id'-i-tē*) [*livere*, to be dark]. The state of being livid. **L., Cadaveric**, or **Post-mortem**, the reddish or bluish discoloration in the dependent parts of a corpse, due to the gravitation of the blood.

Livor (*liv'-or*) [*livere*, to be dark]. Lividity.

Lixivation (*liks-iv'-e-a'-shun*) [*lixivia*, lye].

The process of leaching ashes. Also, the process of separating by solution any alkaline salt from the insoluble impurities with which it is mixed.

Lixivium (*liks-iv'-e-nūm*) [*lixivia*, lye]. The filtrate obtained by leaching ashes; practically a solution of an impure potassium hydrate.

Lobar (*lo'-bar*) [*lobus*, a lobe]. Pertaining to a lobe. **L. Pneumonia**. See **Pneumonia**.

Lobe (*lob*) [*lobus*, a lobe]. A more or less rounded part or projection of an organ, separated from neighboring parts by fissures and constrictions, as the lobes of the liver, of the brain, etc. **L. of the Cerebellum**, each cerebellar hemisphere is divided into the following lobes:—On the upper surface, the anterior or square lobe, and the posterior or semilunar lobe; on the under surface, the flocculus or subpeduncular lobe, the amygdala or tonsil, the digastric lobe, the slender lobe, and the inferior posterior lobe. **L. of the Cerebrum**, the primary lobes into which each cerebral hemisphere is divided; they are, the frontal, parietal, occipital, temporosphenoidal, and central, or island of Reil. See **Brain** and **Convolution**. **L. of the Liver**. See **Liver**. **L., Marginal**, the first frontal convolution running along the margin of the longitudinal fissure. **Lobes**, **Optic**, the corpora quadrigemina.

Lobelia (*lo-be'-le-ah*) [from *de Lobel*, a French botanist]. The leaves and tops of *L. inflata*, of the order *Lobeliaceæ*. It contains a liquid alkaloid, lobelin; an acid, lobelic acid; an acrid body, lobelacrin; and a crystalline substance, inflatin. Lobelia is expectorant, antispasmodic, and emetic, and has been used in asthma, whooping-cough, and other bronchial affections. **L., Ext.**, **Fl.** Dose $\text{m}\text{j}-\text{v}$ (0.065–0.32). **L., Tinct.** Dose $\text{m}\text{x}-\text{xxx}$ (0.65–2.0). **L., Tinct., Ætherea** (B. P.). Dose $\text{m}\text{x}-\text{xxx}$ (0.65–2.0). **L. syphilitica**, a North American species that has been used as an antisyphilitic.

Lobular (*lob'-u-lar*) [*lobulus*, a little lobé].

Pertaining to, resembling, or composed of, lobules. **L. Pneumonia**. See **Pneumonia**.

Lobulated (*lob'-u-la-ted*) [*lobulatus*, from *lobulus*, a little lobe]. Consisting of lobes or lobules.

Lobule (*lob'-ul*) [*lobus*, a lobe]. See **Lobulus**. **L., Paracentral**, the superior connecting convolution of the ascending frontal and ascending parietal gyri.

Lobulus (*lob'-u-lus*) [dim. of *lobus*, a lobe]. A small lobe. **L. caudatus**, the tailed lobe of the liver that separates the right extremity of the transverse fissure from the commencement of the fissure for the inferior vena cava. **L. quadratus**, the square lobe upon the inferior surface of the right lobe of the liver. **L. spigelii**, the lobule projecting from the posterior portion of the inferior surface of the liver.

Lobus (*lo'-bus*) [*L.*]. A lobe.

Local (*lo'-kl*) [*locus*, place]. Limited to a part or place; not general.

Localization (*lo-kal'-iz-a'-shun*) [*locus*, a place]. 1. The determination of the seat of a lesion. 2. The limitation of a process to a particular place; the opposite of generalization. 3. The faculty of locating sensory impressions. **L., Cerebral**, the determination of the position of the centers in the brain that preside over certain physiologic acts, or of the seat of pathologic conditions interfering with the proper function of these centers.

Localized (*lo'-kal-ized*) [*locus*, a place]. Confined to a particular place.

Lochia (*lo'-ke-ah*) [*λόχιος*, pertaining to childbirth]. The discharge from the genital organs during the first few weeks (from 2 to 4) after labor. **L. alba**, the whitish flow that takes place from about the seventh day. **L. cruenta**, **L. rubra**, the sanguineous flow of the first few days. **L. serosa**, the serous discharge taking place about the fifth day.

Lochial (*lo'-ke-al*) [*λόχιος*, pertaining to childbirth]. Pertaining to the lochia.

Lochiometra (*lo-ke-o-me'-trah*) [*λόχιος*, pertaining to childbirth; *μήτρα*, uterus]. A collection of lochia in the uterus.

Lochiorrhea (*lo-ke-or-el-ah*) [*λόχιος*, pertaining to childbirth; *ῥοία*, a flow]. An abnormal flow of the lochia.

Lock-finger. A peculiar affection of the fingers in which they suddenly become fixed in a flexed position, due to the presence of a small fibrous growth in the sheath of the extensor tendon.

Lock-jaw. Tetanus; trismus.

Lock-spasm. A spasm of the fingers in which they become firmly flexed upon the object in their grasp, as upon the pen in writing. It is sometimes seen in writers' cramp.

- Locomotive** (*lo-ko-mo'tiv*) [*locus*, a place; *movere*, to move]. Moving from place to place; able to change its place; pertaining to locomotion. **L. Pulse.** See *Pulse*.
- Locomotor** (*lo-ko-mo'tor*) [*locus*, a place; *motor*, a mover]. Pertaining to locomotion. **L. Ataxia**, or **L. Ataxy.** See *Ataxy* and *Tubes*.
- Locular, Loculated** (*lok'-u-lar*, *lok'-u-lat-ed*) [*loculus*, a cell]. Divided into loculi.
- Loculus** (*lok'-u-lus*) [dim. of *locus*, a place]. A small space or compartment.
- Locus** (*lo'-kus*) [L.]. A place. **L. cinereus**, **L. cæruleus**, **L. ferrugineus**, a bluish-tinted eminence on the fasciculi teretes of the fourth ventricle of the brain. **L. minoris resistentiæ**, a spot of diminished resistance. **L. niger**, a dark area in the center of a section of the crus cerebri. **L. perforatus**, a name given to the anterior and the posterior perforated space at the base of the brain through which blood-vessels pass. **L. ruber**, the red nucleus of the tegmentum.
- Löffler's Alkaline Solution.** A mixture of 30 parts of a concentrated alcoholic solution of methylene-blue and 100 parts of a 1:10,000 aqueous solution of potassium hydroxid. **L.'s Bacillus**, the bacillus of diphtheria. See *Bacteria*, *Table of*. **L.'s Blood-serum Mixture**, a mixture consisting of one part of neutral meat-infusion bouillon containing 1 per cent. of glucose, and 3 parts of blood-serum; it is used as a culture-medium. **L.'s Toluol-solution**, a solution recommended by Löffler for the local treatment of diphtheria. It consists of menthol 10 grams, toluol, sufficient to make 36 c.c., absolute alcohol 60 c.c., liquor ferri chloridi 4 c.c. Another formula is: menthol 10 grams, toluol, sufficient to make 36 c.c., creolin 2 c.c., absolute alcohol 65 c.c.
- Læmology** (*lem-ol'-o-je*). See *Loimology*.
- Logoneurosis** (*log-o-nu-ro'sis*) [*λόγος*, word; *νεῖρον*, nerve]. 1. A neurosis marked by a speech-defect. 2. A neurosis attended with impairment of the mental powers.
- Logopathy** (*log-op'-ath-e*) [*λόγος*, word; *πάθος*, disease]. A disease affecting the speech.
- Logoplegia** (*log-o-pl'e-je-ah*) [*λόγος*, word; *πῆγμα*, stroke]. Loss of the power of uttering articulate speech.
- Logorrhea** (*log-or-e'-ah*) [*λόγος*, word; *ῥοία*, a flow]. Excessive loquacity.
- Logwood** (*log'-wood*). See *Hematoxylon*.
- Loimology** (*loi-mol'-o-je*) [*λοιμός*, plague; *λόγος*, a treatise]. The science of contagious epidemic diseases.
- Loin** (*loin*) [*lumbus*, loin]. The lateral and posterior region of the body between the false ribs and the top of the pelvis.
- Lombardy Leprosy.** Pellagra.
- London Paste.** A caustic paste containing equal parts of sodium hydroxid and unslaked lime.
- Longevity** (*lon-jev'-it-e*) [*longa*, long; *vita*, life]. Long life.
- Longissimus** (*lon-jis'-im-us*) [super. of *longus*, long]. Longest. **L. capitis**, **L. cervicis**, **L. dorsi.** See *Muscles*, *Table of*.
- Longitudinal** (*lon-je-tu'-din-al*) [*longitudo*, length]. Lengthwise; in the direction of the long axis of a body. **L. Fissure**, the fissure dividing the cerebrum.
- Longus** (*larong'-gus*) [L.]. Long. **L. capitis**, the rectus capitis anticus major muscle. **L. colli.** See *Muscles*, *Table of*.
- Loop** [Irish and Gael, *lub*, a loop]. A bend in a cord or cord-like structure. **L. of Henle.** See *Kidney*.
- Lordoma, Lordosis** (*lor-do'-mah*, *lor do'-sis*) [*λорδοζω*, to bend inward]. A curvature of the spine with a forward convexity.
- Loreta's Operation.** 1. The forcible dilatation of the pylorus for the relief of stricture. 2. The treatment of aneurysm by the introduction of metal wire into the sac, through which an electric current is then passed.
- Lotio** (*lo'-she-o*) [L.]. See *Lotion*. **L. hydrargyri flava** (B. P.), yellow mercurial lotion; yellow wash. **L. hydrargyri nigra** (B. P.), black mercurial lotion; black wash.
- Lotion** (*lo'-shun*) [*lotio*, a wash]. A medicinal solution for bathing a part; a wash.
- Louse.** See *Pediculus*.
- Lovage** (*lu'-aj*) [OF., *levesche*, from *ligusticum*, lovage]. The root of *Ligusticum levisticum* and *Levisticum officinale*, plants of the order Umbelliferae. **L.** is stimulant, aromatic, carminative, and emmenagogue. Dose of a fld. ext. f 3j-ij (4.0-8.0). Unof.
- Lower, Tubercle of.** See *Tubercle*.
- Loxa Bark** (*loks'-ah*). Pale cinchona; the bark of *Cinchona officinalis*.
- Loxopterygium** (*loks-o-ter-ij'-e-um*) [*λοξός*, oblique; *πτέρυξ*, wing]. A genus of anacardiaceous trees. **L. lorentzii**, **L. sagotii**, red quebracho, the bark of which is used as a substitute for cinchona.
- Lozenge** (*lo'-enz*) [OF., *losenge*, lozenge]. A medicated tablet with sugar as a basis.
- Lucid** (*lu'-sid*) [*lucidus*, clear]. Clear, shining, not obscure. **L. Interval**, the transitory return of the normal mental faculties in insane or delirious conditions.
- Lucidification** (*lu-sid-ij'-ik-a'-shun*) [*lucidus*, clear; *facere*, to make]. A clearing-up, especially a clearing-up of the protoplasm of cells.
- Ludwig's Angina.** See *Diseases*, *Table of*.
- Lues** (*lu'-es*) [L.] Formerly a pestilential disease; at present used as a euphemism for syphilis.
- Luetic** (*lu-el'-ik*) [*lues*, a plague]. Affected with or relating to lues; syphilitic.

Lugol's Caustic. A solution of iodine and potassium iodide, of each one part, in water two parts. **L.'s Solution.** See *Iodin*.

Lumbago (*lum-ba'-go*) [*lumbus*, the loin]. Pain in the loins.

Lumbar (*lum'-bar*) [*lumbus*, the loin]. Pertaining to the loins. **L. Colotomy**, colotomy performed in the lumbar region. **L. Puncture.** See *Puncture*. **L. Region.** See *Abdomen*. **L. Vertebra.** See *Vertebra*.

Lumbricales (*lum-brik-a'-lez*). See *Muscles*, *Table of*.

Lumbricoid (*lum'-brik-oid*) [*lumbricus*, earth-worm; *eidos*, likeness]. Pertaining to, or resembling, a lumbricus.

Lumbricus (*lum'-brik-us*) [*L.*]. A genus of worms, including the common earth-worm and certain intestinal worms. The latter are now termed *Ascarides*.

Lumbus [*L.*]. The loin.

Lumen (*lu'-men*) [*lumen*, light]. The cavity surrounded by the walls of a tubular vessel.

Luminiferous (*lu-min-if'-er-us*) [*lumen*, light; *ferre*, to bear]. Conveying or bearing light.

Lumpy Jaw. See *Actinomycosis*.

Lunacy (*lu'-nas-e*) [*luna*, the moon]. Insanity, from the superstitious belief that it was influenced by the moon.

Lunar (*lu'-nar*) [*luna*, moon]. Pertaining to the moon or to silver (*luna* of the alchemists). **L. Caustic**, silver nitrate.

Lunatic (*lu'-nat-ik*) [See *Lunacy*]. 1. Pertaining to or affected with insanity. 2. An insane person.

Lung [*AS.*, *lunge*, lung]. The organ of respiration, in which the impure venous blood is oxidized by the air drawn through the trachea and bronchi into the air-vesicles. There are two lungs, a right and a left, the former usually consisting of three, the latter of two lobes. The lungs are situated in the thoracic cavity, and are enveloped by the pleura. At the root or hilum the bronchus and its arteries, and the pulmonary artery and nerves enter, and the pulmonary and bronchial veins and lymphatics leave. The lung proper consists of minute air-vesicles held in place by connective-tissue trabeculae. Capillaries traverse the walls of the air-vesicles and bring the circulating blood in close proximity to the air. The average weight of the adult right lung is 22 ounces; that of the left 20 ounces. **L.-fever**, croupous pneumonia.

Lunula (*lu'-nu-lah*) [dim. of *luna*, moon].

1. The white semilunar area of a nail near the root. 2. A structure resembling the lunula of a nail.

Lupiform (*lu'-pif-orm*) [*lupus*, a wolf; *forma*, form]. Resembling lupus.

Lupine (*lu'-pin*) [*lupus*, a wolf]. *Lupinus*,

a genus of leguminous plants. *Lupinus albus* and others contain a bitter glucoside, lupinin ($C_{29}H_{57}O_{16}$); while from *L. luteus* an alkaloid, arginin ($C_8H_{14}N_4O_2$), is obtained. The bruised seeds of *L. albus* have been used as an external application to ulcers.

Lupinidin (*lu-pin'-id-in*) [*lupus*, a wolf], $C_8H_{16}N_4$. A liquid alkaloid obtained from *Lupinus luteus*.

Lupoid (*lu'-poid*). See *Lupiform*.

Lupulin (*lu'-pu-lin*) [*lupulus*, hop]. The glandular powder obtained from the strobiles of *Humulus lupulus*. See *Humulus*. It is antispasmodic and sedative, and is used in sexual excitement, delirium tremens, renal and vesical irritation, and spermatorrhea. Dose gr. ij-v (0.13-0.32). **Oleoresina lupulini**. Dose \mathfrak{m} x-xl (0.65-2.6). **Ext. lupulini fluidum**. Dose \mathfrak{f} 3 ss-ij (2.0-8.0).

Lupulus (*lu'-pu-lus*). See *Humulus*.

Lupus (*lu'-pus*) [*L.*, a wolf]. **Lupus vulgaris**; a chronic disease of the skin and mucous membranes, characterized by the formation of nodules of granulation-tissue. It passes through a number of phases, and terminates by ulceration or atrophy, with scarification. The cause of the disease is the tubercle-bacillus. **L., Disseminated Follicular**, a variety of lupus confined to the face, especially in the situations usually occupied by acne. The papules are from a large pin's head to a pea in size, conical and deep-red. **L. erythematosus**, **L. erythematodes**; Cazenave's lupus; a form not due to the tubercle-bacillus. It occurs, as a rule, in multiple patches, with a tendency to symmetric arrangement, chiefly about the face and head, occasionally on the extremities, and rarely on the trunk. The patches are sharply defined at the border, flat, very slightly raised, and with a tendency to the formation of crusts. The color is bright-red, and there are no nodules. It is most common in women of adult or middle age. **L. erythematosus sebaceus**, a form with special involvement of the sebaceous glands. **L. exedens**. Synonym of *L. vulgaris*. **L. hypertrophicus**, that variety of *L. vulgaris*, in which new connective-tissue formation predominates over the destructive process, and markedly raised, thick patches result. **L. maculosus**, a variety of *L. vulgaris*, characterized by the eruption of very soft, smooth, brownish-red, semitranslucent milium nodules that develop in the connective tissue of otherwise healthy skin without subjective sensations. **L. non-exedens**, lupus without ulceration. **L. vegetans**, **L. verrucosus**, the formation in the lupus process of a wart-looking patch liable to become inflamed.

Luschka's Gland. See *Gland*. **L's Tonsil**, the adenoid tissue normally existing between the orifices of the Eustachian tubes, analogous in structure to the lymphoid constituents of the tonsil.

Lusus naturæ (*lu'-sus*) [*ludere*, to play]. A freak of nature.

Lutein (*lu'-te-in*) [*luteus*, yellow]. A yellow pigment obtained from corpora lutea by extraction with chloroform.

Luxation (*luks'-a'-shun*). See *Dislocation*.

Luxus (*luks'-us*) [*L.*]. Excess. **L.-consumption**, a term applied to the metabolism of certain surplus proteid material, which, though inside the body, does not form a component part of any of its tissues, but constitutes a kind of reservoir of force upon which the organism may draw.

Lycanthropy (*li-kan'-thro-pe*) [*λύκος*, a wolf; *άνθρωπος*, man]. A form of mania in which the patient imagines himself a wild beast.

Lycoperdon (*li-ko-per'-don*) [*λύκος*, wolf; *πέπτεσθαι*, to break wind]. A genus of fungi. **L. bovista**, fist-ball, puff-ball, devil's snuff-box. This has been used as a styptic, and is now to some extent employed in nervous diseases.

Lycopodium (*li-ko-po'-de-um*) [*λύκος*, wolf; *πούς*, foot]. Club-moss, witch-meal, wolf's-claw. The sporules of *L. clavatum* and other varieties, occurring in the form of a light, fine, yellowish powder, are used as a desiccant and absorbent on moist and excoriated surfaces, and as an inert powder in which to imbed pills to prevent their adhering to each other.

Lye (*li*) [*AS.*, *leah*, lye]. 1. The solution of alkaline hydrates obtained by leaching ashes. 2. Any alkaline solution.

Lying-in. 1. Being in confinement. 2. The puerperal state.

Lymph (*limf*) [*lymp̄ha*, water]. 1. The fluid in the lymphatic vessels, the product of the filtration of the liquid portion of the blood through the walls of the capillaries. 2. The coagulable exudate on an inflamed surface. 3. The liquid material used for vaccination; vaccine-lymph. **L., Animal**, vaccine-lymph obtained from an animal. **L.-cell**, **L.-corpuscle**, a leukocyte occurring in the lymph. **L.-follicles**, small collections of lymphadenoid tissue occurring in mucous membranes. **L.-hearts**, certain organs found in the frog and in some fishes, which are to the lymph-stream what the blood-heart is to the blood-stream. **L., Humanized**, vaccine from a human being. **L., Inflammatory**, that thrown out as a product of inflammation in wounds, etc. **L., Koch's**, tuberculin. **L., Plastic**, fibrinous lymph; that forming embryonic tissue. **L.-scrotum**, an enlargement of the scrotum due to distention

of the lymphatic vessels and hyperplasia of the tissues. **L.-spaces**, the lacunæ occurring in connective tissue and containing lymph.

Lymphadenectasis (*limf-ad-en-ek'-tas-is*) [*lymp̄ha*, lymph; *ἀδην*, gland; *ἐκτασις*, distention]. Dilatation of the sinuses of a lymph-gland producing a tumor-like mass.

Lymphadenitis (*limf-ad-en-i'-tis*) [*lymp̄ha*, lymph; *ἀδην*, gland; *ιτις*, inflammation]. Inflammation of a lymphatic gland.

Lymphadenoid (*limf-ad'-en-oid*) [*lymp̄ha*, lymph; *ἀδην*, gland; *ειδος*, like]. Resembling, or of the nature of, a lymphatic gland, or of lymphatic tissue.

Lymphadenoma (*limf-ad-en-oid'-mah*) [*lymp̄ha*, lymph; *ἀδην*, gland; *ωμα*, tumor]. Hyperplasia of the lymphatic glands. See *Hodgkin's Disease*, in *Diseases*, Table of.

Lymphæmia (*limf-e'-me-ah*). See *Lymphæmia*.

Lymphagogue (*limf'-ag-og*) [*lymp̄ha*, lymph; *ἀγᾱγός*, leading]. 1. Stimulating the flow of lymph. 2. An agent that stimulates the flow of lymph.

Lymphangiectasis (*limf-an-je-ek'-tas-is*) [*lymp̄ha*, lymph; *ἀγγειον*, vessel; *ἐκτασις*, widening]. Dilatation of the lymphatic vessels.

Lymphangiology (*limf-an-je-ol'-o-je*) [*lymp̄ha*, lymph; *ἀγγειον*, vessel; *λόγος*, treatise]. The anatomy, physiology, and pathology of the lymphatics.

Lymphangioma (*limf-an-je-oid'-mah*) [*lymp̄ha*, lymph; *ἀγγειον*, vessel; *ωμα*, tumor]. A tumor made up of lymphatic vessels. **L. tuberosum multiplex**, a very rare disease of the skin, probably congenital, characterized by the formation of large, brownish-red papules or tubercles, the size of lentils, scattered indiscriminately over the trunk. On section they prove to be made up of dilated lymphatics.

Lymphangitis (*limf-an-ji'-tis*) [*lymp̄ha*, lymph; *ἀγγειον*, vessel; *ιτις*, inflammation]. Inflammation of a lymphatic vessel.

Lymphatic (*limf-at'-ik*) [*lymp̄ha*, lymph]. Pertaining to lymph; containing or characterized by lymph. **L. System**, a system of vessels and glands, accessory to the blood-vascular system, conveying lymph. It begins as innumerable capillaries in interspaces of tissues. These form plexuses studded with lymph-glands that act as filters and finally all those below the diaphragm unite in the receptaculum chyli on the second lumbar vertebra. From this the thoracic duct leads upward to empty into the junction of the left subclavian and internal jugular veins. The lymph from the upper right half of the body and head enters the right lymphatic duct, which empties into the junction of the right internal jugular and subclavian veins.

Lymphemia (*limf-e'-me-ah*) [*lymp̄ha*, lymph; *aīma*, blood]. Leukemia characterized by enlargement of the lymphatic glands.

Lymphocyte (*limf'-o-sit*) [*lymp̄ha*, lymph; *kýtos*, a cell]. 1. A lymph-cell. 2. One of Ehrlich's classes of leukocytes, comprising those small cells having large nuclei and a very small amount of protoplasm. See *Leukocyte*.

Lymphodermia (*limf-o-der'-me-ah*) [*lymp̄ha*, lymph; *dé̄rma*, skin]. An affection of the lymphatics of the skin. L. *perniciosa*, leukemic enlargement of the glands.

Lymphoid (*limf'-oid*) [*lymp̄ha*, lymph; *el̄dos*, like]. Having the appearance or character of lymph.

Lymphoma (*limf-o'-mah*) [*lymp̄ha*, lymph; *ô̄ma*, tumor]. A tumor composed of lymphadenoid tissue. The term includes also formations not strictly tumors, as hyperplasias of the tissues proper to lymphatic glands. L., **Malignant**. Synonym of *Hodgkin's Disease*.

Lymphomatous (*limf-o'-mat-us*) [*lymp̄ha*, lymph; *ô̄ma*, tumor]. Of the nature of, or affected with, lymphoma.

Lymphorrhagia (*limf-or-a'-je-ah*) [*lymp̄ha*,

lymph; *rh̄ḡn̄v̄nai*, to burst forth]. A flow of lymph from a ruptured lymphatic vessel.

Lymphosarcoma (*limf-o-sar-ko'-mah*) [*lymp̄ha*, lymph; *sār̄x*, flesh; *ô̄ma*, tumor]. A sarcoma having some of the structural elements of a lymphatic gland.

Lypomania (*li-pe-ma'-ne-ah*) [*λύπη*, sadness; *μανία*, madness]. A form of dementia accompanied by profound mental depression and refusal to take food.

Lyra (*li'-rah*) [*λίρα*, a lyre]. A lyre. L. of **Fornix**, certain longitudinal, transverse, and oblique lines on the inferior surface of the fornix, the arrangement of which bears a fanciful resemblance to a lyre.

Lysis (*li'-sis*) [*λύειν*, to loose]. The gradual decline of a disease, especially of a fever.

Lysol (*li'-sol*) [*λύειν*, to loose]. A brown liquid substance obtained by boiling tar-oils with alkalis and fats. It is used as an antiseptic in surgery; in lupus, gonorrhea, or as a gargle in sore throat.

Lyssa (*lis'-ah*) [*λύσσα*, madness]. A synonym of *Hydrophobia* or *Rabies*.

Lysophobia (*lis-o-fô'-be-ah*) [*λύσσα*, madness; *φ̄βος*, fear]. Morbid dread of rabies; pseudohydrophobia.

M

M. The abbreviation of *Musculus*, *Meter*, *Myopia*, *Mille*, a thousand, *Minim* and *Misce*, mix; it also represents the number 1000.

Mace (*mās*). See *Myristica*.

Maceration (*mas-er-a'-shun*) [*macerate*, to make soft]. The process of softening a solid substance, or of converting into a soft mass by soaking in a liquid, as, e. g., M. of the fetus.

Macies (*ma'-se-tz*) [*macies*, a wasting]. Atrophy, leanness, wasting.

Macrobiosis (*mak-ro-bi-o'-sis*) [*μακρός*, long; *βίος*, life]. Longevity.

Macrocephalia (*mak-ro-sef-a'-le-ah*) [*μακρός*, large; *κεφαλή*, head]. Abnormal largeness of the head.

Macrocephalous (*mak-ro-sef'-al-us*) [*μακρός*, large; *κεφαλή*, head]. Characterized by an abnormally large head.

Macrocheilia (*mak-ro-ki'-le-ah*) [*μακρός*, large; *χείλος*, lip]. Excessive development of the lips, a characteristic of certain negro tribes. It occurs also in cretinoid states, when the lips and cheeks are the seat of lymphangioma.

Macrocheiria (*mak-ro-ki'-re-ah*) [*μακρός*,

large; *χείρ*, hand]. Great enlargement of the hands.

Macrocyte (*mak'-ro-sit*) [*μακρός*, large; *κύτος*, cell]. A giant blood-corpuscle found in the blood in certain anemias, especially pernicious anemia.

Macroductily (*mak-ro-dak'-til-e*) [*μακρός*, large; *δάκτυλος*, finger]. Abnormally large size of the fingers or toes.

Macroglossia (*mak-ro-glos'-e-ah*) [*μακρός*, great; *γλῶσσα*, tongue]. Enlargement of the tongue, a condition seen in cretins, in whom it is probably due to lymphangioma.

Macromania (*mak-ro-ma'-ne-ah*) [*μακρός*, large; *μανία*, madness]. A form of mania characterized by the delusion that objects are larger than they really are; or the impression that one's own body or members are much larger than they are.

Macromelus (*mak-rom'-el-us*) [*μακρός*, long; *μέλος*, organ or member]. 1. Having excessively large limbs. 2. One having excessively large limbs.

Macrophage (*mak'-ro-fāj*) [*μακρός*, large; *φαγεῖν*, to devour]. A large phagocyte.

Macropodia (*mak-ro-pō'-de-ah*) [*μακρός*, large; *πούς*, foot]. Excessive size of the feet.

Macropsia (*mak-ro-p'-se-ah*). See *Megalopsia*.

Macroscopic (*mak-ro-skop'-ik*) [*μακρός*, large; *σκοπεῖν*, to see]. Large enough to be seen by the naked eye; gross; not microscopic.

Macrosomatia (*mak-ro-so-ma'-she-ah*), **Macrosomia** (*mak-ro-so'-me-ah*) [*μακρός*, large; *σῶμα*, body]. Excessive size of the body.

Macrostoma (*mak-ras'-to-mah*) [*μακρός*, large; *στόμα*, mouth]. Congenital fissure of the angle of the mouth, producing a large opening.

Macrotin (*mak'-ro-tin*) [*macrotys*, *cimicifuga*]. A resin obtained by the precipitation of tincture of *cimicifuga* with water.

Macula (*mak'-u-lah*) [L.]. A spot; especially one upon the skin, not elevated above the surrounding level. **Maculæ acusticæ**, the terminations of the auditory nerve in the sacculæ and utricle. **M. lutea**, the yellow spot of the retina. It is the point of clearest vision. **M. solaris**, a freckle.

Macular (*mak'-u-lar*) [*macula*, spot]. Characterized by or resembling maculæ.

Maculate (*mak'-u-lat*) [*macula*, a spot]. Spotted.

Maculation (*mak'-u-lat'-shun*) [*macula*, a spot]. The state or quality of being spotted; the formation of maculæ.

Macule (*mak'-ul*) [*macula*, a spot]. Macula.

Mad [AS., *mad*, mad]. 1. Insane. 2. Affected with rabies; rabid.

Madder (*mad'-er*) [AS., *maderu*, madder]. See *Rubia*.

Madura Foot (*ma-dur'-rah*). A disease occurring chiefly in India, and characterized by the formation on the foot (sometimes on the hand) of a tender purplish swelling, which in time suppurates, the pus being evacuated through one or more sinuses. These sinuses discharge a seropurulent liquid containing peculiar bodies resembling gunpowder-grains or fish-roe. The disease is chronic, but remains local, and is believed to be due to an organism allied to the actinomyces-fungus. It is also termed *Mycetoma*.

Magendie's Solution. See *Morphin*.

Magenta (*ma-jen'-tah*) [from a town in Italy]. A coal-tar dye from which fuchsin and a large number of other dyes are prepared.

Magnesia (*mag-ne'-se-ah*) [*Μαγνησιος*, belonging to Magnesia, in Thessaly; *λίθος Μαγνησιος*, Magnesian stone.]. Magnesium oxid, MgO. **M. White**, magnesium carbonate. See *Magnesium*.

Magnesian (*mag-ne'-zih*) [*magnesium*]. Pertaining to, or containing magnesium.

Magnesium (*mag-ne'-ze-um*) [*Μαγνησιος*, belonging to Magnesia, in Thessaly; *λίθος Μαγνησιος*, Magnesian stone]. Symbol Mg; atomic weight 24.3; quantivalence 12, specific gravity 1.75. A bluish-white metal of the group to which calcium and barium belong. It is abundantly distributed throughout inorganic and organic nature; its salts are used in the arts and in medicine. The source of magnesium and its salts is chiefly the minerals dolomite and kieserite. The following compounds are used in medicine. **Magnesium carbonas** $MgCO_3$, $Mg(OH)_2 + 5H_2O$, exists in two forms,—as light (*M. carbonas levis*), and as heavy magnesium carbonate (*M. carbonas ponderosa*). It is antacid, laxative, and antilithic. Dose $\overline{3}$ ss- $\overline{3}$ ij (2.0-8.0). **Liq. magnesi carbonatis** (B. P.). Dose f $\overline{3}$ j-ij (32.-64.0). **M. citras effervescens**, is cathartic. Dose $\overline{3}$ j-ij (4.0-12.0). **Liq. magnesi citratis**. Dose f $\overline{3}$ iv-vij (128.0-256.0). **M. oxid**, MgO (Magnesia, U. S. P.), is obtained by calcining magnesium carbonate, and exists in two forms,—as light magnesia, and as heavy magnesia (*Magnesia ponderosa*, U. S. P.). It is used as an antacid and laxative, as a dusting powder, and as an antidote to arsenic. Dose gr. x- $\overline{3}$ j (0.65-4.0). **M. sulphas**, $MgSO_4 + 7H_2O$, Epsom salt, is an active cathartic, especially useful in inflammatory affections. Dose $\overline{3}$ j- $\overline{3}$ j (4.0-32.0). An enema magnesi sulphatis is official in B. P. **M. sulphas effervescens** (B. P.). Dose $\overline{3}$ ij- $\overline{3}$ j (8.0-32.0).

Magnet (*mag'-net*) [*Μαγνησιος*, belonging to Magnesia, in Thessaly, where loadstone was first found]. 1. Loadstone, a magnetic oxid of iron. 2. A body having the power to attract iron bodies. **M., Electro-**, a piece of iron rendered magnetic by a current of electricity passing through a coil surrounding the magnet. **M., Horse-shoe**, an iron magnet having the shape of a horse-shoe. **M., Permanent**, one the magnetic properties of which are permanent, in contradistinction to **M., Temporary**, which derives its magnetism from another magnet or from a galvanic current.

Magnetic (*mag-net'-ik*) [*Μαγνησιος*, pertaining to Magnesia, in Thessaly]. Pertaining or belonging to a magnet. Possessing the property of magnetism.

Magnetism (*mag-net'-izm*) [*magnet*]. The power possessed by a magnet to attract or repel other masses. **M., Animal**, hypnotism.

Magnetization (*mag-net-iz-a'-shun*) [*magnet*]. The process of rendering a substance magnetic.

Magnetolectricity (*mag-net-o-e-lek-tris'*).

it-e) [*magnet*; ἤλεκτρον, amber]. Electricity produced by means of a magnet.

Magnetoinduction (*mag-net'-o-in-duk'-shun*) [*magnet*; *inductio*, induction]. The production of an induced current by the insertion of a magnet within a coil of wire.

Magnetotherapy (*mag-net-o-ther'-ap-e*) [*magnet*; *θεραπεία*, treatment]. The treatment of diseases by magnets.

Magnification (*mag-nif-ik-a'-shun*) [*magnus*, large; *facere*, to make]. Enlargement, especially the enlargement of the image of an object by means of lenses.

Magnifying (*mag'-nif-i-ing*) [*magnus*, large; *facere*, to make]. Enlarging; making greater.

M. Power, the power of a lens to increase the diameters of the image of an object.

Magnum, or **Os magnum** [*L.*, great bone]. The largest bone in the distal row of the carpus situated between the unciform and the trapezoid bones.

Maid, Maiden (*mād, ma'-den*) [*AS.*, *mægdēn*, a maiden]. A young unmarried woman; a virgin.

Maidenhead. 1. Virginity. 2. The hymen.

Maim (*mām*) [*OF.*, *mehaigner*, to maim]. To cripple by injury or removal of a limb.

Main-en-griffe (*mang-on-grif'*) [*Fr.*]. See *Claw-hand*.

Maize (*māz*) [*W. Ind.*, *mahiz*, maize]. Indian corn. See *Zea mays*.

Make (*māk*) [*AS.*, *macian*, to make]. In electricity, to establish the flow of an electric current.

Mal [*Fr.*, from *malum*, evil, disease]. Disease. **M. de mer**, sea-sickness. **M.**, Grand, epilepsy. **M.**, Petit. See *Petit Mal*.

Mala (*ma'-lah*) [*L.*]. The cheek-bone or the cheek.

Malacia (*mal-a'-se-ah*) [*μαλακία*, a softening]. A morbid softening of tissue.

Malacotomy (*mal-ak-o'-o-me*) [*μαλακός*, soft; *τομή*, a cutting]. Incision of the abdomen; celiotomy.

Malady (*mal'-ad-e*) [*malum*, evil]. Disease.

Malaise (*mal-āz'*) [*Fr.*]. A general feeling of illness, accompanied by restlessness and discomfort.

Malar (*mal'-lar*) [*mala*, cheek]. Pertaining to the cheek-bone. **M. Bones**, the two cheek-bones.

Malaria (*mal-a'-re-ah*) [*It.*, *mala aria*, bad air]. See *Malarial Fever*.

Malarial (*mal-a'-re-al*) [*It.*, *mala aria*, bad air]. Pertaining to malaria. **M. Cachexia**, a chronic form of malaria characterized by anemia, general failure of health, a sallow complexion, and enlargement of the spleen. **M. Fever**, a disease associated with the presence in the blood of the *Plasmodium malarie*, and characterized by periodicity, enlargement of the spleen, and the presence in the

blood, free or within the red corpuscles, of parasites (plasmodia) that exert a deleterious influence upon the red cells. The paroxysms may be intermittent, remittent, or irregular. If repeated daily the fever is designated quotidian; if on alternate days, tertian; if with an interval of two days, quartan. If two paroxysms occur daily the fever is designated a double quotidian. There may be a double tertian form, a paroxysm occurring daily, but only those of alternate days being alike; a double quartan form, and others. A typical malarial paroxysm consists of a cold stage, a hot stage, and a sweating stage, occurring in the sequence given. Intermittent fever is characterized by the occurrence of a complete intermission of the symptoms in the interval between two paroxysms, the temperature becoming normal or subnormal. In remittent fever there is only an amelioration of the symptoms in the intervals. In certain localities in which the malarial organisms are exceedingly numerous or intensely virulent, the attack displays a pernicious tendency. Of this type there may be a cerebral form, characterized either by delirium and excitement, or by coma and depression; a thoracic form, in which the respiration is accelerated and there is an urgent sense of the need of air; a gastrointestinal form, attended with nausea, vomiting, jaundice, and diarrhea; or an asthenic or algid form, in which there is a condition of marked debility and a striking coldness of the surface and of the breath. To the irregular manifestations of malarial poisoning, which do not at any time present the classical association of chill, fever, and sweat, the designation of "dumb ague" is given. The enlargement of the spleen in chronic malaria is sometimes designated "ague cake." **M. Hematuria**, the presence of blood in the urine as a result of malarial poisoning. **M. Neuralgia**, neuralgia due to malarial intoxication.

Malassimilation (*mal-as-im-il-a'-shun*) [*malus*, bad; *assimilatio*, likeness]. Defective assimilation.

Malate (*mal'-at*) [*μαλον*, an apple]. A salt of malic acid.

Malaxation (*mal-aks-a'-shun*) [*μαλάσσειν*, to soften]. 1. The act of kneading. 2. A form of massage.

Male (*māl*) [*masculus*, a male]. 1. Pertaining to the male sex, that which impregnates the female. 2. A member of the male sex. 3. Of a double-bladed instrument, the blade which is received into a hollow of the other (female) blade. **M. Fern**, the *Aspidium filix mas*. **M. Organ**, the penis.

Malformation (*mal-for-mal'-shun*) [*malus*, ill; *formatio*, a forming]. An abnormal development or formation of a part of the body.

Malgaigne's Hooks (*mahlgane*) [*Malgaigne*, a French surgeon]. An instrument for holding the parts of a fractured patella in apposition.

Malic Acid (*ma'-lik*). See *Acid*.

Malignancy (*mal-ig'-nan-se*) [*malus*, bad; *gignere*, to produce]. The quality of being malignant.

Malignant (*mal-ig'-nant*) [*malus*, bad; *gignere*, to produce]. Bad; compromising or threatening life. **M. Edema.** See *Edema*. **M. Pustule**, anthrax. **M. Tumor**, a tumor that destroys life. Malignant tumors recur and give rise to metastasis.

Malingering (*mal-in'-jer-er*) [*Fr. malingre*, sickly, from *malus*, bad; *ager*, ill, sick]. One who feigns illness or defect.

Malingering, Malingery (*mal-in'-jer-ing*, *mal-in'-jer-e*) [*Fr. malingre*, from *malus*, bad; *ager*, ill, sick]. The feigning of disease.

Malleable (*mal'-e-a-bl*) [*malleus*, hammer]. Capable of being beaten or rolled into thin sheets.

Malleation (*mal-e-a'-shun*) [*malleus*, hammer]. A spasmodic action of the hands, consisting in continuously striking any near object.

Mallein (*mal'-e-in*) [*malleus*, farcy]. A fluid obtained from cultures of the *Bacillus mallei*, the microorganism of glanders. When injected into the circulation of a glandered animal, it causes an elevation of temperature, and has been recommended for use in the early diagnosis of farcy or glanders.

Malleoincudal (*mal-e-a-ine'-ku-dal*) [*malleus*, hammer; *incus*, anvil]. Relating to the malleus and the incus.

Malleolar (*mal'-e-a-lar*) [*malleolus*, little hammer]. Relating to a malleolus.

Malleolus (*mal'-e-o-lus*) [*dim. of malleus*, hammer]. A part or process of bone having a hammer-head shape. **M., External**, the lower extremity of the fibula. **M., Internal**, a process on the internal surface of the lower extremity of the tibia.

Malleus (*mal'-e-us*) [*malleus*, hammer]. 1. One of the ossicles of the internal ear having the shape of a hammer. 2. Glanders.

Mallow (*mal'-o*). See *Malva*.

Malpighian Body (*mal-pig'-e-an*) [from *Malpighi*, an Italian anatomist]. The commencement of a uriniferous tubule, consisting of the glomerulus of vessels (the Malpighian tuft) and the membranous envelop (Bowman's capsule). **M. Corpuscle**, any one of the minute whitish nodules of lymphadenoid tissue in the red substance of the spleen along the course of the blood-vessels.

Malposition (*mal-po-zish'-un*) [*malus*, bad; *ponere*, to place]. An abnormal position of any part or organ, especially of the fetus.

Malpractice (*mal-prak'-tis*) [*malus*, bad; *πράσσειν*, to do]. Improper treatment; treatment of a disease by a method contrary to that taught by experience; also, the unlawful production of an abortion.

Malpresentation (*mal'-pre-sen-ta'-shun*) [*malus*, bad; *presentare*, to place before]. In obstetrics, such a position of the child at birth that delivery is difficult or impossible.

Malt, Maltum (*mawlt*, *mawlt'-tum*) [*L.*]. The seeds of common barley, *Hordeum distichum*, made to germinate by warmth and moisture, and then baked so as to stop the germinating process. The germinated grains contain diastase, dextrin, and maltose, as well as proteids. Malt is used as a nutrient in wasting diseases. **M., Ext.** Dose ʒj-iv (4.0-16.0). **Unof. M., Ext., Fld. (N. F.)** is a solution of the principles of malt in water and alcohol. **M.-liquors**, infusions of malt fermented so as to contain alcohol. Those in common use are beer, ale, and porter. Beer is made by a comparatively slow fermentation, and contains about 2.5 per cent. of alcohol. Ale and porter are fermented more rapidly, and contain about 4.7 per cent. of alcohol. The malt used in making porter is browned, giving the liquor a darker color.

Malta Fever. A febrile disease prevalent on the coasts bordering on the Mediterranean. It is characterized by malaise, an irregular fever, rigors, headache, pains in the limbs, and diarrhea; and frequent relapses. The etiology is not definitely known; some of the cases are probably forms of typhoid fever. It is also termed Rock fever, Mediterranean fever and Neapolitan fever.

Maltine (*mawlt'-tin*) [*maltum*, malt]. A name given to various proprietary preparations of malted wheat or barley, useful as food for invalids.

Maltose (*mawlt'-sôs*) [*maltum*, malt]. $C_{12}H_{22}O_{11} + H_2O$. A variety of sugar formed, together with dextrin, by the action of malt-diastase upon starch.

Malva (*mal'-vah*) [*L., mallow*]. The mallow; a genus of malvaceous plants. The leaves of *M. alcea*, *M. rotundifolia*, and *M. silvestris* are used as demulcents.

Mamma (*mam'-ah*) [*L.*]. The breast; the milk-secreting gland of the mother.

Mammalgia (*mam'-al'-je-ah*) [*mamma*, breast; *ἀλγος*, pain]. Pain in the mamma.

Mammalia (*mam-a'-le-ah*) [*mamma*, breast]. A division of the class of vertebrates including all animals that suckle their young.

Mammary (*mam'-a-re*) [*mamma*, breast]. Pertaining to the mammae. **M. Artery.** See *Arteries*, Table of. **M. Gland.** See *Gland*.

M. Line, the vertical line passing through the nipple.

Mammilla (*mam-ill'-ah*) [*dim. of mamma*,

breast]. A small prominence or papilla. **M. of Breast**, the nipple or teat.

Mammillaplasty (*mam-il-ap-las'-te*) [*mammilla*, nipple; *πλάσσειν*, to mold]. A plastic operation for the purpose of elevating a depressed nipple.

Mammitis (*mam-i'-tis*). See *Mastitis*.

Mammose (*mam'-ds*) [*mamma*, breast]. Having full or abnormally large breasts.

Mandible, Mandibula (*man'-dib-l, man-dib'-u-lah*) [*mandere*, to chew]. The inferior maxillary bone.

Mandibular (*man-dib'-u-lar*) [*mandere*, to chew]. Pertaining to the mandible, or lower jaw.

Mandragora (*man-drag'-o-rah*) [*μανδραγόρας*, mandrake]. A genus of solanaceous plants; the mandrake. *M. officinalis* has been used as a narcotic and hypnotic. It and other species contain an alkaloid, mandragorin, $C_{17}H_{23}NO_3$, resembling atropin in action.

Mandrake (*man'-drak*). See *Mandragora* and *Podophyllum*.

Mandrin (*man'-drin*) [Ger.]. The firm guide or stylet (usually of metal) that gives rigidity to a flexible catheter whilst it is being inserted.

Manducation (*man-du-ka'-shun*) [*manducatio*, a chewing]. The chewing or mastication of food.

Manganese, Manganum (*man'-gan-er, man-gan'-um*) [an altered form of *Magnesium*]. Symbol Mn; atomic weight 54.8; quantivalence II, IV, VI, VII. A brittle, hard, grayish-white metal, having a specific gravity of 7.2, and resembling iron in properties. It forms several oxides, the highest of which, Mn_2O_7 , forms an acid, $HMnO_4$, from which salts, the permanganates, are produced. **Mangani dioxidum**, MnO_2 , black oxide of manganese, is tonic and alterative, and has been used in syphilis, chlorosis, in various skin-diseases, and in certain forms of dyspepsia. Dose gr. liij-xx (0.2-1.3). It is employed in the arts, and in laboratories for the purpose of obtaining chlorine and oxygen.

Mangani sulphas, $MnSO_4 + 4H_2O$, has been used as a substitute for iron in anemia, and as a cholagogue. Dose gr. v-xx (0.32-1.3). Potassium permanganate is a salt of permanganic acid, the latter being derived from manganese heptoxid, Mn_2O_7 . See *Potassium*.

Mange (*mánj*) [Fr., *manger*, to eat]. A parasitic skin-disease of horses, cattle, and dogs, resembling scabies, and due to various species of *Acarus*.

Mania (*ma'-ne-ah*) [*uavía, uavéσθαι*, to rage]. A form of insanity marked by great mental and emotional excitement, by hallucinations, delusions, physical excitement, and often a

tendency to violence. **M., Alcoholic**, acute mania of alcoholic origin. It is to be distinguished from delirium tremens, although sometimes used synonymously with it. **M. a potu**. See *Delirium tremens*. **M., Bell's**, an acute delirium running a rapidly fatal course, with slight fever, and in which post-mortem no lesions are found sufficient to account for the symptoms. There are the wildest hallucinations, insomnia, and intense excitement, followed by a condition called typhomania, with elevation of temperature, dry tongue, and rapid, feeble pulse. **M., Epileptic**, a maniacal outburst in an epileptic, often associated with a destructive tendency. **M., Puerperal**, a form of mania or abnormal mental action sometimes following childbirth. **M., Religious**, mania in which the central idea is religious in character, or in which a powerful religious emotion has been the exciting cause.

Maniac (*ma'-ne-ah*) [*uavía*, madness]. An insane person; one affected with mania.

Manicure (*man'-ik-ur*) [*manus*, the hand; *cura*, care]. 1. The processes employed in caring for and beautifying the hand. 2. One who professionally attends to the care of the hands and nails.

Manihot (*man'-e-hot*) [L.]. A genus of euphorbiaceous plants, yielding cassava and tapioca.

Manikin (*man'-ik-in*) [OF., *manequin*, a puppet]. A model of the body, made of plaster, papier maché, or other material, and showing by means of movable parts the relations of the organs.

Manioc (*man'-e-ok*). See *Manihot*.

Maniple (*man'-ip-l*) [*manipulus*, from *manus*, hand]. A handful.

Manipulation (*man-ip-u-la'-shun*) [*manipulus*, a handful]. A handling; the use of the hands for the purpose of performing some work in a skillful manner, such as reducing a dislocation, returning a hernia into its cavity, or changing the position of a fetus.

Manipulus (*man-ip'-u-lus*). See *Maniple*.

Manna (*man'-ah*) [*uávva*, manna]. The concrete, saccharine exudation of the flowering ash, *Fraxinus ornus*, and other trees. Manna contains a sweet principle, mannite or mannitol, $C_8H_{14}O_6$, a sugar, a purgative principle, and a mucilage. Some specimens contain also a glucosid, fraxin. Manna is a mild laxative. Dose $\frac{3}{4}$ -ij (32.0-64.0).

Manometer (*man-om'-et-er*) [*uavós*, rare; *μέτρον*, measure]. An instrument for measuring the tension of liquids and gases, consisting either of a bent tube filled with mercury (mercurial M.), or of a spring (spring M.), connected with a writing-style.

Manometric (*man-o-met'-rik*) [*uavós*, rare; *μέτρον*, measure]. Pertaining to a manom-

eter; pertaining to tracings obtained by means of a manometer. **M. Flames**, flames of different heights and characters seen in a rotating mirror and due to the reflection of a pulsating gas-flame when the supplying gas is set in motion by sound-waves. They are also known as Koenig's flames.

Manual (*man'-u-al*) [*manus*, hand]. Pertaining to the hands; performed by the hands.

Manubrium (*man-u'-bre-um*) [L.]. A handle. **M. of Malleus**, the handle-shaped process of the malleus of the ear. **M. manus**, the radius. **M. of Sternum**, the upper piece of the sternum.

Manus (*ma'-nus*) [L.]. The hand.

Manustupration (*man-u-stu-pra'-shun*) [*manus*, hand; *stuprare*, to ravish]. Masturbation.

Manyplices (*man'-ip-li-z*) [AS., *manig*, many; *plicare*, to fold]. The third compartment in the stomach of ruminants. It is also called the omasum or psalterium.

Maranta (*mar-an'-tah*). See *Arrowroot*.

Marantic (*mar-an'-tik*) [*maraineiv*, to make lean]. Pertaining to marasmus. **M. Clot**, a blood-clot produced by slowing of the circulation in depressed states of the system. **M. Thrombosis**, thrombosis due to general malnutrition.

Maraschino (*mar-as-ke'-no*) [Sp.]. A liquor made from morello cherries.

Marasmatic (*mar-as-mat'-ik*). Synonym of *Marasmic*.

Marasmic (*mar-as'-mik*) [*maraineiv*, to grow lean]. Affected with marasmus.

Marasmus (*mar-as'-mus*) [*maraineiv*, to grow lean]. A gradual wasting of the tissues of the body from insufficient or imperfect food-supply. There is either no organic lesion, or gastrointestinal catarrh.

Marc (*mark*) [Fr., dregs]. 1. A by-product in the manufacture of wines, consisting of the stems, skins, and stones of the grapes. 2. The residue remaining after the expression of the oil from certain fruits.

Margaric Acid (*mar-gar'-ik*). See *Acid*. **Margaric Acid Crystals**, needle-shaped crystals consisting of compounds of the fatty acid, found in foci of fatty degeneration, in the urine, etc.

Margarin (*mar'-gar-in*) [*μαργαρος*, the pearl-oyster]. 1. A margarate of glyceryl, especially glyceryl trimargarate, $C_3H_5(C_{17}H_{33}O_2)_3$, found in butter. 2. An artificial substitute for butter.

Marginal (*mar'-jin-al*) [*margo*, margin]. Pertaining to the margin or border. **M. Convolution**. See *Convolutions*, *Table of*.

Mariotte's Law. See *Law*.

Marjoram (*mar'-jo-ram*). See *Origanum*.

Marrow (*mar'-o*) [AS., *marh*, marrow].

The fatty substance contained in the medullary canal of long bones, and in the interstices of cancellous bone. In early life the marrow of all bones is red (red marrow), but later that within the shafts of long bones assumes a light color (yellow marrow). Red marrow is composed of a delicate reticulum of connective tissue, containing blood-vessels, large connective-tissue cells, some of which in growing bone become osteoblasts (marrow-cells), giant-cells (myeloplaxes) and red corpuscles in various stages of formation. In yellow marrow most of the cells have been transformed into fat-cells. The function of bone-marrow is probably the formation of red corpuscles. In certain forms of anemia the marrow undergoes profound changes; that of the shafts of the long bones may return to its embryonical condition. Bone-marrow has been used in the treatment of pernicious anemia.

Marrubium (*mar-u'-be-um*) [L.]. Horehound. The leaves and tops of *M. vulgare*, of the order Labiatae. It contains a volatile oil, a bitter principle, marrubin, tannin, resin, and lignin. At present horehound is mainly employed in catarrhal affections of the respiratory tract. Dose $\bar{5}$ ss-j (2.0-4.0). Unof.

Marsh-fever. Malaria.

Marsh-gas. Methane.

Marsh-mallow. See *Althaea*.

Marsh's Test. A test for arsenic. Hydrogen is generated by the action of an acid on zinc, the suspected liquid is added, and if arsenic is present, it combines with the hydrogen to form arsine, AsH_3 , which may be ignited at the end of the delivery-tube and the flame allowed to impinge against cold porcelain. Metallic arsenic is deposited and is recognized by the character of the stain and chemic tests.

Martin's Bandages. India-rubber bandages, from 5 to 21 feet in length, used for making compression of a limb for the cure of ulcers, varicose veins, etc.

Maruta (*ma-ru'-tah*) [L.]. A genus of herbs of the order Compositae. **M. cotula**, may-weed, or dog-fennel, is used as a substitute for chamomile.

Mask (*mask*) [Fr., *masque*, a mask]. 1. A bandage applied to the face in case of erysipelas, burns or scalds, eczema, etc. 2. Synonym of *Chloasma*.

Masked (*maskt*) [Fr., *masque*, a mask]. Covered with a mask; concealed. **M. Disease**, one that is concealed by concomitant symptoms.

Mass, **Massa** (*mas*, *mas'-ah*) [*massa*, a mass]. 1. An aggregation of particles of matter. 2. A cohesive substance that can be made into pills. **M., Blue**. See *Mercury*.

M., Vallet's. *Massa ferri carbonatis*. See *Ferrum*.

Massage (*mas-ahsh'*) [Fr., from *μάσσειν*, to knead]. A method of rubbing, kneading, or stroking of the superficial parts of the body by the hand or an instrument, for the purpose of modifying nutrition, restoring power of movement, breaking up adhesions, etc. A male person performing M. is termed a *masseur*; a female person, a *masseuse*.

Masseter (*mas'-e-ter*) [*μασητήρ*, chewer]. One of the muscles of mastication. See *Muscles*, *Table of*.

Masseteric (*mas-et-er'-ik*) [*μασητήρ*, chewer]. Pertaining to the masseter-muscle.

Massicot (*mas'-ik-ot*) [Fr.]. PbO. Lead oxid; litharge.

Mast-cells [Ger., *Mastzellen*]. Cells filled with basophile granules, found in the connective tissue and in foci of chronic inflammation.

Mastalgia (*mas-tal'-je-ah*) [*μαστός*, breast; *ἀλγος*, pain]. Pain in the breast.

Mastic, Mastiche (*mas'-tik*, *mas'-ti-ke*) [*μαστήριον*, mastic]. The resin flowing from the incised bark of the *Pistacia lentiscus*, a tree of the *Terebinthaceæ*. It is used as a styptic, as a filling for teeth, and as a microscopic varnish.

Mastication (*mas-tik-a'-shun*) [*masticare*, to chew]. The act of chewing.

Masticatory (*mas'-tik-a-to-re*) [*masticare*, to chew]. 1. Pertaining to mastication, or to the muscles of mastication. 2. A remedy to be chewed but not swallowed, used for its local action on the mouth. **M. Spasm**, spasm of the muscles of mastication; *trismus*.

Mastitis (*mas-ti'-tis*) [*μαστός*, breast; *ίτις*, inflammation]. Inflammation of the breast.

M., Interstitial, inflammation of the connective tissue of the breast. **M., Parenchymatous**, inflammation of the proper glandular substance of the breast.

Mastodynia (*mas-to-din'-e-ah*) [*μαστός*, breast; *ὀδύνη*, pain]. Pain in the breast.

Mastoid (*mas'-toid*) [*μαστός*, breast, nipple; *είδος*, like]. 1. Nipple-shaped, as the M. process of the temporal bone. 2. The mastoid process. 3. Pertaining to the mastoid process, as M. foramen, M. operation. **M. Abscess**, an abscess of the mastoid cells.

M. Antrum, a cavity in the mastoid portion of the temporal bone. **M. Cells**, the hollow air-spaces in the mastoid process communicating with the middle ear. **M. Disease**, inflammation of the mastoid cells, *mastoiditis*. **M. Foramen**. See *Foramina*, *Table of*.

Mastoiditis (*mas-toid-i'-tis*) [*μαστός*, nipple; *ίτις*, like; *ίτις*, inflammation]. Inflammation of the mastoid cells.

Mastooccipital (*mas-to-ok-sip'-it-al*) [*μαστός*, nipple; *occiput*, occiput]. Pertaining to the mastoid process and the occipital bone.

Mastoparietal (*mas-to-par-i'-et-al*) [*μαστός*, nipple; *paries*, wall]. Pertaining to the mastoid process and the parietal bone.

Masturbation (*mas-tur-ba'-shun*) [*masturbare*, to pollute one's self]. Production of the venereal orgasm by friction of the genitals.

Matè (*mah'-ta*) [Sp., *mate*, a vessel]. Paraguayan tea. The leaves of *Ilex paraguayensis*, which are used in South America as a substitute for tea and coffee. Its properties are due to them.

Materia medica (*mat-e'-re-ah med'-ik-ah*) [L., medical matter]. The science that treats of the sources and preparations of the drugs and agents used in medicine.

Materies morbi (*mat-e'-re-tz mor'-bi*) [L., matter of disease]. The material that is the cause of a disease.

Maternal (*mat-ur'-nal*) [*mater*, mother]. Pertaining to the mother. **M. Impressions**. See *Impressions*.

Maternity (*mat-ur'-nit-e*) [*mater*, mother]. 1. Motherhood. 2. A lying-in hospital.

Matico (*mat-e'-ko*) [Sp.]. The leaves of *Piper angustifolium*, of the order *Piperaceæ*. It is aromatic and stimulant, and has been used as a local and general hemostatic, and as an alterative stimulant to mucous membranes. Dose of the powder \mathfrak{ss} – \mathfrak{ij} (2.0–4.0). **Ext. matico fld.** Dose \mathfrak{ss} – \mathfrak{ij} (2.0–4.0). **Tinctura matico.** Dose \mathfrak{ss} (4.0).

Matricaria (*mat-rik-a'-re-ah*) [*matrix*, mold]. German chamomile; the flower-tops of *Matricaria chamomilla*, of the order *Compositæ*. *Matricaria* contains a volatile oil and a bitter extractive principle, and is a mild tonic, and in large doses emetic and antispasmodic.

Matrix (*ma'-tri-ks*) [L., a mold in which anything is cast]. 1. A mold; the cavity in which anything is formed. 2. That part or tissue into which any organ or process is set, as the matrix of a tooth or of a nail. 3. The intercellular substance of a tissue, as of cartilage. 4. The uterus.

Maturation (*ma-tu-ra'-shun*) [*maturare*, to ripen]. Ripening, as the ripening of the ovum or of a cataract.

Mature (*ma-tu'-r*) [*maturare*, to ripen]. 1. To ripen. 2. Ripe.

Matutinal (*ma-tu'-tin-al*) [*Matuta*, goddess of the morning]. Occurring in the morning, as M. nausea.

Matzoon (*mat'-zun*). Milk fermented with a peculiar ferment obtained from Asiatic Turkey. It is used like koumiss in irritated states of the gastrointestinal tract.

Maxilla (*maks-il'-ah*) [L.]. The bone of the upper or lower jaw.

Maxillary (*maks'-il-a-re*) [*maxilla*, jaw-

bone]. Pertaining to the maxillæ or jaws. **M. Bones**, the bones of the jaws, consisting of the lower and upper jaw. **M. Nerve, Inferior.** See *Nerves, Table of*. **M. Nerve, Superior.** See *Nerves, Table of*. **M. Sinus**, the antrum of Highmore in the superior maxilla.

Maximal (*maks'-im-al*) [*maximus*, the greatest]. Pertaining to the maximum; highest; largest. **M. Thermometer**, one registering the highest point reached by the temperature.

Maximum (*maks'-im-um*) [L., neuter of *maximus*, the greatest]. The greatest or highest degree or amount of anything; the highest point attained or attainable by anything. **M. Dose**, the largest dose of a medicament that may be safely given.

May-apple. See *Podophyllum*.

Mayhem (*ma'-hem*) [OF., *mehaigner*, to hurt]. Maiming.

McBurney's Point. A point two inches above the right anterior superior spine of the ilium on a line drawn from this spine to the umbilicus, at which there is tenderness to pressure in many cases of appendicitis.

Meadow Saffron. See *Colchicum*.

Measles (*mèz'-els*) [Du., *maselen*, measles].

1. An acute, infectious disease, characterized by a peculiar eruption and by catarrhal inflammation of the mucosæ of the conjunctiva and the air-passages. After a period of incubation of nearly two weeks the disease begins with a chill, fever, coryza, cough, and conjunctivitis; on the third or fourth day a dusky-red, papular eruption appears, arranged in the form of crescentic groups. After having reached its maximum, in three or four days, the eruption gradually fades, and is followed by a branny desquamation. The disease affects principally the young, is exceedingly contagious, and one attack of it confers almost perpetual immunity. Its cause is thought to be a bacillus (*Bacillus* of Canon and Pielticke). 2. A disease of hogs, cattle, and sheep, due to the presence in the body of the *Cysticercus cellulose* and larvae of other tapeworms. 3. The cysticerci themselves. **M., Black., M., Hemorrhagic**, a grave variety of measles (1st def.), in which the eruption is hemorrhagic and the constitutional symptoms profound. **M., German.** See *Rötheln*.

Measly (*mèz'-le*) [Du., *maselen*, measles]. Containing measles (cysticerci).

Meat (*mè*) [AS., *mete*, meat]. The muscular tissues of an animal, used as food.

Meatus (*me-a'-tus*) [*meare*, to flow or pass]. An opening or passage. **M. auditorius externus**, the canal extending from the concha to the membrana tympani. **M. auditorius internus**, the internal auditory canal. **M. urethræ**, **M. urinarius**, the orifice of the urethra. **M. of Nose**, one of the three pass-

ages into which the turbinal bones divide the nasal cavity.

Mecca balsam. See *Balm of Gilead*.

Mechanic, Mechanical (*me - kan'-ik*, *me-kan'-ik-al*) [*μηχανή*, a machine]. Pertaining to mechanics, or to physical forces, not to chemic or vital forces. **M. Theory**, Virchow's theory of tumor-formation, according to which tumors are due primarily to local irritation.

Mechanics (*me - kan'-iks*) [*μηχανή*, a machine]. The science that treats of the influence and effects of force upon matter, and that may be divided into statics, the science treating of matter at rest, and dynamics, that treating of matter in motion.

Mechanism (*mek'-an-izm*) [*μηχανή*, a machine]. 1. An aggregation of parts arranged in a mechanic way to perform the functions of a machine. 2. The manner in which a mechanic act is performed, as the M. of labor.

Meckel's Diverticulum [I. F. Meckel, a German anatomist]. See *Diverticulum, M.'s Ganglion.* See *Ganglia, Table of*.

Meconarcein (*mek - o - nar'-se - in*) [*μήκων*, opium; *ναρκοίν*, to benumb]. A mixture of alkaloids of opium, free from morphin, having sedative properties.

Meconate (*mek'-an-ât*) [*μήκων*, poppy]. A salt of meconic acid.

Meconic (*mek-on'-ik*) [*μήκων*, poppy]. Pertaining to opium. **M. Acid.** See *Acid*.

Meconin (*mek' - on - in*) [*μήκων*, poppy]. $C_{10}H_{10}O_4$. A crystalline substance that is obtained on boiling narcotin with water. It is hypnotic. Dose 1 gr. (0.06).

Meconium (*mek-a'-ne-um*) [*μήκων*, poppy]. The first fecal discharges of the newborn, a dark-green, viscid substance, composed of the secretion of the liver with exfoliated epithelium from the bowel.

Mediad (*me'-de-ad*) [*medius*, middle]. Toward the median plane or line.

Medial (*me'-de-al*). See *Median*.

Median (*me'-de-an*) [*medius*, the middle]. Situated or placed in the middle; mesal, or mesial. **M. Artery.** See *Arteries, Table of*. **M. Nerve.** See *Nerves, Table of*.

Mediastinal (*me - de - as - ti - nal*) [*mediastinum*]. Pertaining to the mediastinum.

Mediastinitis (*me de-as-tin-i'-tis*) [*mediastinum*; *ετις*, inflammation]. Inflammation of the cellular tissue of the mediastinum.

Mediastinopericarditis (*me-de-as'-tin-o-per-ik-ar-di'-tis*) [*mediastinum*; *περι*, around; *καρδια*, heart; *ετις*, inflammation]. Combined inflammation of the mediastinum and the pericardium.

Mediastinum (*me-de-as-ti'-num*) [*in medio stare*, to stand in the middle]. 1. A partition separating adjacent parts. 2. The space left in the middle of the chest between the two pleuræ, divided into the anterior, middle,

posterior, and superior mediastinum. The anterior M. contains the origins of the triangularis sterni muscles, the internal mammary vessels of the left side, loose areolar tissue, lymphatic vessels, and a few lymphatic glands. The middle M. contains the heart and pericardium, the ascending aorta, the superior vena cava, the bifurcation of the trachea, the pulmonary arteries and veins, and the phrenic nerves. The posterior M. contains a part of the aorta, the greater and lesser azygos veins, the pneumogastric and splanchnic nerves, the esophagus, the thoracic duct, and some lymphatic glands. The superior mediastinum, that part lying above the pericardium, contains the origins of the sternohyoid and sternothyroid muscles, and part of the longus colli muscles, the transverse portion of the aortic arch, the innominate, left carotid, and subclavian arteries, the superior vena cava and the innominate veins, the left superior intercostal vein, the pneumogastric, cardiac, phrenic, and left recurrent laryngeal nerves, the trachea, esophagus, thoracic duct, the remains of the thymus gland, and lymphatics. **M. testis**, a septum in the posterior portion of the testicle formed by a projection inward of the tunica albuginea.

Mediate (*med'-de-ät*) [*medius*, the middle]. Indirect; performed through something interposed, as M. percussion, percussion on a pleximeter.

Medical (*med'-ik-al*) [*mederi*, to heal]. Pertaining to medicine. **M. Diseases**, diseases treated by the physician, as distinguished from surgical diseases. **M. Ethics**, those principles of justice, honor, and courtesy that regulate the intercourse and conduct of physicians. **M. Jurisprudence**. See *Jurisprudence*.

Medicament (*med-ik'-am-ent*) [*mederi*, to heal]. A medicinal substance.

Medicated (*med'-ik-a-ted*) [*mederi*, to heal]. Impregnated with a medicinal substance.

Medication (*med'-ik-a'-shun*) [*mederi*, to heal]. 1. Impregnation with a medicine. 2. Treatment by medicines; the administration of medicines. **M., Hypodermic**, treatment by the introduction of medicines beneath the skin, usually by means of a hypodermic syringe.

Medicinal (*med'-is'-in-al*) [*mederi*, to heal]. Pertaining to, or having the nature of, a medicine. **M. Rashes**, eruptions on the skin following the internal administration of certain drugs.

Medicine (*med'-is-in*) [*medicina*, medicine; from *mederi*, to heal]. 1. Any substance given for the cure of disease. 2. The science of the treatment of disease; the healing art. In a restricted sense, that branch of the healing art dealing with internal diseases. **M.,**

Clinic, the study of disease by the bedside of the patient. **M., Forensic**, or **M., Legal**, medical jurisprudence, or medicine in its relation to questions of law. **M., Practice of**, the practical application of the principles taught by the **Theory of M. M., Preventive**, that which aims at the prevention of disease.

Medicochirurgical (*med-ik-o-ki-rur'-jik-al*) [*medicus*, a physician; *χειρουργος*, a surgeon]. Pertaining conjointly to medicine and surgery.

Medicolegal (*med-ik-o-le'-gal*) [*medicus*, physician; *lex*, law]. Relating both to medicine and the law.

Medicus (*med'-ik-us*) [*L.*]. A physician.

Medina Worm (*me-dit'-nah*). The *Filaria medinensis*.

Mediolateral (*me-de-o-lat'-er-al*) [*medius*, middle; *latus*, side]. Pertaining to the middle and to a side.

Mediopontine (*me-de-o-pont'-tin*) [*medius*, middle; *pons*, bridge]. Pertaining to the central portion of the pons.

Mediterranean Fever (*med-it-er-a'-ne-an*). See *Malta Fever*.

Medium (*me'-de-um*) [*medius*, middle.]. 1. That in which anything moves or through which it acts. 2. The soil upon which anything grows, especially a substance used for cultivating bacteria; culture or nutrient medium.

Medulla (*me-dul'-ä*) [*L.*, marrow]. 1. The marrow. 2. The medulla oblongata. 3. Anything resembling marrow in structure or in its relations to other parts—as a fatty substance or marrow occupying certain cavities. Also the central parts of certain organs, as distinguished from the cortex. **M. of Nerve-fiber**, the white substance of Schwann. See *Nerve*. **M. oblongata**, the upper enlarged part of the spinal cord, extending from the cord opposite the foramen magnum to the pons Varolii. **M. spinalis**, the spinal cord or marrow. **M. of Kidney**. See *Kidney*.

Medullary (*med'-ul-a-re*) [*medulla*, marrow]. 1. Pertaining to the marrow; resembling marrow. 2. Pertaining to any medulla, as that of the brain. 3. Pertaining to the medulla oblongata. **M. Canal**, the hollow interior of long bones in which the marrow lies. **M. Carcinoma**, a soft carcinoma very rich in cells. **M. Groove**, a longitudinal groove at the anterior part of the embryonal shield of the blastoderm. **M. Rays**. See *Kidney*.

Medullated (*med'-ul-a-ted*) [*medulla*, marrow]. Containing or covered by medulla or marrow. **M. Nerve-fibers**, nerve-fibers provided with a medullary sheath, the white substance of Schwann.

Medullitis (*med-ul-it'-tis*) [*medulla*, marrow; *itis*, inflammation]. 1. Inflammation of marrow. 2. Myelitis.

Medullization (*med-ul-iz-ot-shun*) [*medulla*, marrow]. Conversion into marrow, as the softening of bone-tissue in the course of osteitis.

Mega-, Megalo- (*meg'-ah-, meg'-al-o-*) [*μεγας*, large]. A prefix signifying large; also, indicating a unit one million times as great as the unit to which it is prefixed.

Megacephalic, Megacephalous (*meg-ah-sef-al'-ik, meg-as-ef'-al-us*) [*μεγας*, large; *κεφαλη*, head]. Large-headed; applied to a skull the capacity of which exceeds 1450 c.c.

Megaloblast (*meg'-al-o-blast*) [*μεγας*, large; *βλαστος*, germ]. A giant-corpuscle of the blood.

Megalocephalic (*meg-al-o-sef-al'-ik*). Same as *Megacephalic*.

Megalocephaly (*meg-al-o-sef'-al-e*) [*μεγας*, large; *κεφαλη*, head]. 1. The condition of having a very large head. 2. A disease characterized by progressive enlargement of the head, face, and neck, involving both the bony and the soft tissues. It is also termed *leontiasis ossæ*.

Megalocornea (*meg-al-o-kor'-ne-ah*) [*μεγας*, large; *cornea*, cornea]. An enlarged condition of the cornea.

Megalocyte (*meg'-al-o-sit*) [*μεγας*, large; *κυτος*, cell]. An abnormally large red blood-corpuscle.

Megalodactylous (*meg-al-o-dak'-til-us*) [*μεγας*, great; *δακτυλος*, digit]. Having abnormally large fingers and toes.

Megaloglossia (*meg-al-o-glos'-e-ah*). See *Macroglossia*.

Megalomania (*meg-al-o-ma'-ne-ah*) [*μεγας*, great; *μανια*, madness]. 1. Mania characterized by delusions of grandeur. 2. The delirium of grandeur.

Megalopsia (*meg-al-op'-se-ah*) [*μεγας*, large; *οψας*, sight]. A disturbance of vision in which objects seem larger than they are.

Megavolt (*meg'-ah-volt*) [*μεγας*, large; *volt*]. A million volts.

Megohm (*meg'-ohm*) [*μεγας*, great; *ohm*]. A million ohms.

Megrim (*me'-grim*) [Fr., *migraine*]. See *Migraine*.

Meibomian (*mi-bo'-me-an*). Described by Heinrich Meibom, or Meibomius, a German physician, 1638-1700. **M.** Cyst, a cyst of the Meibomian gland. **M.** Glands, the sebaceous glands of the eyelids, each having an excretory duct opening at the free border.

Meio- (*mi'-o-*). For words thus beginning see *Mio-*.

Meissner (*mis'-ner*) [a German scientist]. **M.**, Corpuscles of. Tactile corpuscles situated at the apices of the papillæ of the corium. **M.**, Plexus of, a plexus of nerves found in the submucous layer of the small intestine.

Mel [L.]. Honey. The saccharine substance deposited in the honeycomb by the common honey bee, *Apis mellifica*, and a few other hymenopterous insects. It contains a large amount of dextrose and levulose, and has the same properties as sugar. In medicine it is used as a vehicle, especially in gargles, and as an application to foul ulcers. **M. boracis** (B. P.), is used as a mouth-wash in thrush and aphthæ. **M. despumatum**, clarified honey. **M. rosæ**, is used as an addition to gargles in ulcerated conditions of the mouth and throat.

Melæna (*mel-e'-nah*). See *Melena*.

Melanæmia (*mel-an-e'-me-ah*). See *Melanemia*.

Melancholia, Melancholy (*mel-an-ko'-le-ah, mel'-an-ko'-le*) [*μυλας*, black; *χολη*, bile]. A disorder of the mind characterized by a profound emotional depression, and a tendency toward introspection, impairment of the mental and physical faculties, with or without delusions. **M. agitata**, a form associated with excessive motor excitement. The patient rushes about, wringing his hands and lamenting loudly. **M. attonita**, or **M. with Stupor**, or **M., Stuporous**; a form in which the patient is perfectly motionless, lies in bed or sits up with his eyes open and fixed, and is absolutely indifferent to everything about him. **M., Simple, M. without Delirium, Hypomelancholia**, a mild form without delusions. **M., Stuporous**. See *M. attonita*.

Melancholiac (*mel-an-ko'-le-ah*) [*μυλας*, black; *χολη*, bile]. 1. Suffering from melancholia. 2. A person affected with melancholia.

Melanemia (*mel-an-e'-me-ah*) [*μυλας*, black; *αιμα*, blood]. The presence in the blood-plasma or in the corpuscles, or in both, of dark pigment-granules due to the disintegration of the hemoglobin. It is also termed *hemachromatosis*. Its most common cause is malaria.

Melanephrosis (*mel-an-ef-id-ro'-sia*) [*μυλας*, black; *επιρρωσις*, excessive perspiration]. Black perspiration.

Melanin (*mel'-an-in*) [*μυλας*, black]. A black pigmentary matter occurring naturally in the choroid coat of the eye, the skin, the hair, the muscles, and, pathologically, in the skin in Addison's disease and in melanotic tumors. Melanin usually contains sulphur, rarely iron. It is a product of cell-activity and belongs to the so-called metabolic pigments.

Melanism (*mel'-an-izm*) [*μυλας*, black]. The abnormal deposition of dark pigment in an organ or organism.

Melanocarcinoma (*mel-an-o-kar-sin-o'-mah*) [*μυλας*, black; *καρκινος*, crab; *ωμα*, tumor]. A carcinoma containing melanin.

Melanoderma, Melanodermia (*mel-an-o-der'-mah*, *mel-an-o-der'-me-ah*) [*μέλας*, black; *δέρμα*, skin]. Black pigmentation of the skin. **M.**, Parasitic. See *Vagabond's Disease*.

Melanoid (*mel'-an-oid*) [*μέλας*, black; *εἶδος*, like]. Dark-colored; resembling melanosis.

Melanoma (*mel-an-o'-mah*) [*μέλας*, black; *δύμα*, tumor]. A tumor containing melanin.

Melanopathy (*mel-an-op'-ath-e*) [*μέλας*, black; *πάθος*, illness]. A disease attended with a deposit of dark pigment.

Melanorrhagia (*mel-an-or-a'-je-ah*) [*μέλας*, black; *ῥήγναι*, to burst forth]. The copious discharge of blackened feces.

Melanosarcoma (*mel-an-o-sar'-ko'-mah*) [*μέλας*, black; *σάρξ*, flesh; *δύμα*, tumor]. A sarcoma containing melanin.

Melanosis (*mel-an-o'-sis*) [*μέλας*, black; *νόσος*, disease]. A general tendency to the formation in the blood, and the deposition in organs, of a dark granular pigment which is usually derived from the hemoglobin of the blood. **M. lenticularis progressiva**. See *Xeroderma pigmentosum*.

Melanotic (*mel-an-o'-ik*) [*μέλας*, black; *νόσος*, disease]. Pertaining to or characterized by melanosis or by a deposit of melanin.

Melanuria (*mel-an-u'-re-ah*) [*μέλας*, black; *urina*, urine]. The presence of black pigment in the urine.

Melasma (*mel-as'-mah*) [*μέλας*, black]. A deposit of dark pigment in the skin. **M. suprarenale**, Addison's disease.

Melena (*mel-e'-nah*) [*μέλας*, black]. The discharge of stools colored black by altered blood. It is quite common in the new-born. **M. neonatorum**, an extravasation of blood into the stomach and intestines of the new-born infant, occurring most often in the first few hours of life.

Melezitose (*mel-e'-it-ös*) [Fr., *mélèze*, larch]. $C_{18}H_{32}O_{16} + 2H_2O$. A sugar found in European false manna, or Briançon manna.

Melicera, Meliceris (*mel-is-e'-rah*, *mel-is-e'-ris*) [*μέλι*, honey; *κηρίδιον*, wax]. A cyst containing a substance having a honey-like appearance.

Melilotus (*mel-il-o'-tus*) [*μέλι*, honey; *λωτός*, lotus]. A genus of leguminous herbs. **M. officinalis**, the sweet clover, is official in the G. P. It contains coumarin, $C_9H_6O_2$, and melilotic acid, $C_{15}H_{14}O_6$, coumaric acid, $C_9H_7O_4$, of which coumarin is the anhydride. Melilot is used but little at the present day.

Melissa (*mel-is'-ah*) [*μέλισσα*, a bee]. A genus of labiate plants. The leaves and tops of **M. officinalis**, balm or lemon balm, is the *Melisa* of the U. S. P., which contains tannin, a bitter extractive, and an essential oil. Balm is used as a tonic in febrile affections, and as a flavoring agent.

Melitagra (*mel-it-a'-grah*) [*μέλι*, honey; *ἀγρᾶ*, seizure]. Eczema associated with the formation of honey-like crusts.

Melitemia (*mel-it-e'-me-ah*) [*μέλι*, honey; *αἷμα*, blood]. The presence of an excess of sugar in the blood.

Melitose (*mel-it-ös*) [*μέλι*, honey], $C_{18}H_{32}O_{16} + 5H_2O$. A crystalline sugar occurring in Australian manna, flour of cotton seeds, sugar-beets, and in the molasses obtained in the manufacture of sugar.

Melituria (*mel-it-u'-re-ah*) [*μέλι*, honey; *urina*, urine]. Diabetes mellitus. **M. inosita**, the presence of inosite in the urine.

Mellite (*mel'-it*). See *Mellitum*.

Mellithemia (*mel-ith-e'-me-ah*). See *Mellitemia*.

Mellitum (*mel-i'-tum*) [*mel*, honey]. In pharmacy, a honey, a preparation in which honey is the menstruum. **M. rosæ**. See *Mel rosæ*.

Melon-seed Bodies. Fibrous bodies, resembling melon-seeds in size, sometimes found in joints and cysts of tendon-sheaths.

Meloplasty (*mel'-o-plas-te*) [*μήλον*, cheek; *πλάσσειν*, to form]. A plastic operation on the cheek.

Melting Point. The degree of temperature at which solids pass into the liquid state.

Member (*mem'-ber*) [*membrum*, a limb]. A part of the body, especially a projecting part, as the leg, the arm.

Membrana (*mem-brā'-nah*) [L.]. A membrane. **M. adventitia**, the adventitia of blood vessels; also, the decidua reflexa. **M. agnina**, the amnion. **M. basilaris**. See *Lamina basilaris*. **M. caduca**, the decidua. **M. capsularis**, a capsular ligament. **M. decidua**. See *Decidua*. **M. flaccida**. See *Membrane, Shrapnell's*. **M. germinativa**, the blastoderm. **M. granulosa**, the layer of small polyhedral cells within the theca folliculi of the Graafian follicle. **M. limitans**, the limiting layer of the retina. There are two, the internal and external. **M. propria**, the delicate membrane upon which the epithelium of mucous membranes rests; the basement membrane. **M. ruychiana**, the middle or capillary layer of the choroid. **M. serotina**. The part of the decidua entering into the formation of the placenta. **M. tectoria**. A delicate membrane of the internal ear, also known as *Corti's membrane*. See *Ear*. **M. tensa**, the tympanic membrane proper, exclusive of Shrapnell's membrane. **M. tenuis**, the arachnoid. **M. tympani**. See *Membrane, Tympani*.

Membranaceous (*mem-bran-ä'-sh-us*) [*membrana*, membrane]. Pertaining to, consisting of, or of the nature of, a membrane.

Membrane (*mem'-bran*) [*membrana*, Latin, *membrum*, member]. A thin, spread-out

surrounding a part or separating adjacent cavities. **M.**, **Animal**, a membrane made from animal tissues, used in dialyzing. **M.**, **Basement-**, a delicate membrane, made up of flattened cells, underlying the epithelium of mucous surfaces. **M.-bone**, any bone that originates, not in cartilage, but in membrane, as, *e.g.*, some of the cranial bones. **M.** **of Bruch**, a thin transparent membrane, the external layer of the choroid of the eye. **M.** **of Corti**. See *Membrana tectoria*. **M.**, **Costocoracoid**, a dense layer of fascia extending between the subclavius muscle and the pectoralis minor, and forming the anterior portion of the sheath of the axillary vessels. **M.**, **Cricothyroid**, the membrane connecting the thyroid and cricoid cartilages of the larynx. **M.**, **Croupous**. See *M.*, *Diphtheric*. **M.**, **Débove's**, a layer of flattened connective tissue cells beneath the epithelium of the bronchial, vesical, and intestinal mucous membranes. **M.** **of Descemet**. See *Descemet's Membrane*. **M.**, **Diphtheric**, a fibrinous layer formed on a mucous membrane or cutaneous surface and extending downward for a variable depth. When not firmly adherent, as in the larynx, it is termed a croupous membrane. It is the result of coagulation-necrosis, generally, but not always, brought about by the bacillus of diphtheria. **M.**, **Drum**, the tympanic membrane. **M.**, **False**. See *M.*, *Diphtheric*. **M.**, **Fenestrated**, the elastic membrane of the intima of arteries, also known as the fenestrated membrane of Henle. **M.**, **Fetal**, name given to the chorion, amnion, or allantois. **M.**, **Hyaloid**, a delicate membrane investing the vitreous humor of the eye. **M.**, **Jacob's**, the layer of rods and cones of the retina. **M.**, **Krause's**, a thin, transverse, dark disc bisecting the light substance of the striped muscle-fibers. **M.**, **Mucous**, the membrane lining those cavities and canals communicating with the air. It is kept moist by the mucus secreted by the goblet-cells and mucous glands. **M.** **of Nasmyth**, a delicate firm membrane covering the external surface of the enamel for a short time after birth. **M.**, **Nictitating**, the winking membrane of the lower animals, represented in the human eye by the plica semilunaris. **M.**, **Obturator**, the fibrous membrane closing the obturator foramen. **M.**, **Persistent Pupillary**. See *M.*, *Pupillary*. **M.**, **Pupillary**, a delicate, transparent membrane closing the pupil in the fetus. It disappears between the seventh and eighth months; when it persists after birth it is termed persistent pupillary membrane. **M.**, **Pyogenic**, the lining of an abscess-cavity or a fistulous tract. The term should be restricted to the lining of an abscess that is spreading and in

which the membrane produces pus. **M.** **of Reissner**, a delicate membrane stretching across the outer wall of the cochlea. **M.**, **Reticular**, the membrane covering the space of the outer hair-cells of the cochlea. **M.** **of Ruysch**. See *Membrana ruyschiana*. **M.**, **Schneiderian**, the mucosa lining the nasal fossæ. **M.**, **Secondary Tympanic**, the membrane closing the fenestra rotunda. **M.**, **Serous**, a delicate membrane covered with flat endothelial cells lining closed cavities of the body, *e.g.*, the peritoneum, the pleura. **M.**, **Shrapnell's**, a small portion of the drum-membrane filling the notch of Rivini. **M.**, **Synovial**, a membrane covering the articular extremities of bones and the inner surface of ligaments entering into the formation of a joint. **M.**, **Tenon's**. See *Tenon's Capsule*. **M.**, **Thyrohyoid**, the membrane joining the thyroid cartilage and hyoid bone. **M.**, **Tympanic**, the drum-membrane; the membrane separating the external from the middle ear. It consists of three layers: an outer or skin-layer, a fibrous layer, and an inner mucous layer.

Membranous (*mem'-bran-us*) [*membrana*, membrane]. Pertaining to, having the nature of, or consisting of a membrane. **M.** **Labyrinth**. See *Labyrinth*. **M.** **Urethra**, the part of the urethra between the two layers of the triangular ligament.

Memory (*mem'-o-ry*) [*memor*, mindful]. That faculty of the mind by which ideas and sensations are recalled.

Menarche (*men'-ar-ka*) [*μήν*, month; *ἀρχή*, beginning]. The period at which menstruation is inaugurated.

Menhidrosis, Menidrosis (*men-hid-ro'-sis*, *men-id-ro'-sis*) [*μήν*, month; *ἰδρωσις*, sweat]. The replacement of the menstrual flow by a bloody sweat.

Menière's Disease (*men-e-är'*). See *Diseases*, *Table of*.

Meningeal (*men-in'-je-al'*) [*μήνινξ*, membrane]. Pertaining to the meninges.

Meninges (*men-in'-jēs*) [pl. of *μήνινξ*, membrane]. The membranes covering the brain and spinal cord, and consisting of the dura, pia, and arachnoid.

Meningitic (*men-in-jit'-ik*) [*μήνινξ*, membrane; *τις*, inflammation]. Pertaining to, or affected with, meningitis.

Meningitis (*men-in-jit'-tis*) [*μήνινξ*, membrane; *τις*, inflammation]. Inflammation of the membranes of the brain or cord; that of the dura is termed pachymeningitis, that of the pia-arachnoid, leptomeningitis, or simply meningitis. Meningitis of the membranes of the brain is classified into acute and chronic, the former being subdivided into serous and purulent, the latter into fibrous,

ossifying, and deep, or encephalomeningitis. According to location, two varieties are spoken of—that of the vertex and that of the base. **M.**, Acute Cerebral, is due to traumatism, to extension of inflammation from adjacent structures, especially from the middle ear, the orbit, the nasal sinuses, or to tuberculosis (Tuberculous **M.**); it may be secondary to acute infectious processes elsewhere in the body, as pneumonia, erysipelas, typhoid fever, influenza, small-pox, or it may be a primary disease, as in Cerebrospinal **M.** The symptoms are fever, slow pulse, later rapid pulse, headache, delirium, rigidity and retraction of the neck, convulsions, vomiting, a scaphoid abdomen, constipation, optic neuritis; in advanced stages various palsies, such as ptosis, squint, facial paralysis. In epidemic cerebrospinal **M.** there is usually a characteristic eruption. See *Exanthemata*, Table of. **M.**, Acute Spinal, may occur in tuberculosis, as a secondary process in acute infectious diseases, such as small-pox, scarlet-fever, pneumonia, as a part of epidemic cerebrospinal meningitis, as the result of extension of inflammation from neighboring parts, and as the result of exposure to cold and wet. The symptoms are chill, fever, pain in the back and limbs, rigidity of the muscles, dyspnea, exaggerated reflexes, later paralysis. **M.**, Chronic Cerebral, is due to syphilis, tuberculosis, or may be associated with disease of the brain in encephalomeningitis (paretic dementia). The main symptoms are, in the first two, headache, convulsions, and rigidity of the muscles of the neck. **M.**, Chronic Spinal, is due to syphilis, traumatism, and the excessive use of alcohol. It is also frequently an accompaniment of the sclerosis of the spinal cord. The symptoms are pain, hyperesthesia along the spinal nerves, increased reflexes, paralysis.

Meningocele (*men-in'-go-sel'*) [*μήνινξ*, membrane; *κήλη*, tumor]. A protrusion of the cerebral or spinal meninges through a defect in the skull or vertebral column. It forms a cyst filled with cerebrospinal fluid.

Meningocerebritis (*men-in-go-ser-e-brit'-tis*). See *Meningoencephalitis*.

Meningoencephalitis (*men-in-go-en-sef-al-i'-tis*) [*μήνινξ*, membrane; *encephalitis*]. Inflammation of the brain and its membranes.

Meningomyelitis (*men-in-go-mi-el-i'-tis*) [*μήνινξ*, membrane; *myelitis*]. Inflammation of the spinal cord and its meninges.

Meningomyelocoele (*men-in-go-mi-el-o-sel'*) [*μήνινξ*, membrane; *μυελίς*, marrow; *κήλη*, tumor]. A protrusion from the spinal column of a portion of the cord and membranes.

Meningorrhachidian (*men-in-go-rak-id'-e-an*) [*μήνινξ*, membrane; *ραχίς*, spine]. Relating to the spinal cord.

Meningosis (*men-in-go'-sis*) [*μήνινξ*, membrane]. The union of bones by a membranous attachment.

Meninx (*men'-ing-ks*) [*μήνινξ*, membrane]. A membrane, especially one of the brain or spinal cord.

Meniscus (*men-is'-kus*) [*μήνισκος*, a crescent]. A crescent or crescentic body, especially an interarticular fibrocartilage. Also a concavoconvex (Positive **M.**) or convexoconcave lens (Negative **M.**).

Menispermum (*men-is-per'-mum*) [*μήνη*, moon; *σπέρμα*, seed]. Yellow parilla; Canadian moonseed. The rhizome and roots of *M. canadense*, of the order Menispermaceae. It contains an alkaloid, menisperm, and berberin. It is alterative and is used as a substitute for sarsaparilla. Ext. menispermis fid. Dose ʒss-j (2.0-4.0).

Menocelis (*men-o-sel'-lis*) [*μήν*, month; *κέλις*, spot]. Dark erythematous or hemorrhagic spots occurring upon the skin in failure of menstruation.

Menopause (*men-o-pauz*) [*μήν*, month; *παύω*, cessation]. The physiologic cessation of menstruation, usually occurring between the 45th and 50th year.

Menoplasia (*men-o-pla'-ne-ah*) [*μήν*, month; *πλάω*, deviation]. A discharge of blood occurring at the menstrual period, but derived from some other part of the body than the uterus.

Menorrhagia (*men-or-a'-je-ah*) [*μήν*, month; *μή*, rivat, to burst forth]. An excessive menstrual flow.

Menorrhea (*men-or-e'-ah*) [*μήν*, month; *ῥοία*, a flow]. The normal flow of the menses. Also, excessive menstruation.

Menostasia, Menostasis (*men-os-ta'-ze-ah*, *men-os-tas-is*) [*μήν*, month; *στάσις*, standing]. A suppression of the menstrual flow.

Menses (*men'-sez*) [*mensis*, a month]. The recurrent monthly discharge of blood from the genital canal of a woman during sexual life.

Menstrual (*men'-stru-al*) [*menstruus*, monthly, from *mensis*, month]. Pertaining to menstruation.

Menstruation (*men-stru-a'-shun*) [*menstruus*, monthly, from *mensis*, month]. A periodic discharge of a sanguineous fluid from the uterus, occurring during the period of a woman's sexual activity, from puberty to the menopause. **M.**, Vicarious, the discharge of blood at the time of menstruation from some other organ or part than the vagina.

Menstruum (*men'-stru-um*) [from *menstruus*, monthly—so called because its action was assisted by a moderate fire—because chemic solvents under the influence of the

u-ra'-

care,

to measure]. The act of measuring; one of the methods of physical diagnosis.

Mentagra (*men-ta'-grah*). See *Sycosis*.

Mentagrophyton (*men-tag-roff'-it-on*) [*men-tagra*, sycosis; *φυτόν*, plant]. A fungus, microsporon mentagrophytes, thought to be the cause of sycosis.

Mental (*men'-tal*) [*mens*, the mind; *mentum*, the chin]. 1. Pertaining to the mind. 2. Pertaining to the chin.

Mentha (*men'-thah*) [L.]. A genus of labiate plants, the mints. *M. piperita*, peppermint, the leaves and tops of *M. piperita*. It is an aromatic stimulant used to relieve nausea, flatulence, and spasmodic pains in the stomach and bowels. **Aqua menthæ piperitæ**. Dose indefinite. **Oleum menthæ piperitæ**. Dose m-j-v (0.065-0.32). **Spiritus menthæ piperitæ**. Dose m-x-xxx (0.65-2.0). **Trochisci menthæ piperitæ**. Dose indefinite. **Mentha viridis**, spearmint, the leaves and tops of *M. viridis*. It has properties and uses similar to those of peppermint. **Aqua menthæ viridis**. Dose indefinite. **Oleum menthæ viridis**. Dose m-j-v (0.13-0.32). **Spiritus menthæ viridis**. Dose m-x-xl (0.65-2.3).

Menthene (*men'-then*) [*mentha*, mint], $C_{10}H_{18}$. A liquid hydrocarbon produced when menthol is distilled with phosphorus pentoxid.

Menthol (*men'-thol*) [*mentha*, mint], $C_{10}H_{18}O$. Mint-camphor, a crystalline stearopten derived from oil of peppermint, and used as an anodyne and rubefacient in neuralgia, in skin-diseases associated with itching and in rhinitis.

Mentoanterior (*men-to-an-te'-re-or*) [*mentum*, chin; *anterior*, before]. Having the chin toward the front.

Mentoposterior (*men-to-pos-te'-re-or*) [*mentum*, chin; *posterior*, after, behind]. Having the chin toward the back.

Mentum (*men'-tum*) [L.]. The chin.

Menyanthes (*men-e-an'-thes*) [*μήν*, month; *άνθος*, flower, from its reputed emmenagogue properties]. A genus of plants of the Gentianaceæ. *M. trifoliata*, buckbean. It contains a bitter principle, menyanthin ($C_{23}H_{36}O_{11}$), and has been used in malaria, scrofula, dropsy, jaundice, rheumatism, etc. Dose of the powdered leaves, gr. xx-xxx (1.3-2.0).

Mephitic (*mef-it'-ik*) [*mephitis*, a pestilential exhalation]. Foul or noxious; stifling; noisome. *M. Gangrene*, necrosis of bone, associated with the evolution of offensive odors.

Meralgia (*mer-al'-je-ak*) [*μυρῶς*, thigh; *ἀλγος*, pain]. Neuralgic pain in the thigh.

Mercaptan (*mer-kap'-tan*) [*mercurius*, mercury; *captans*, seizing, on account of combining readily with mercury]. A derivative of an alcohol in which the oxygen of the latter is replaced by sulphur.

Mercurial (*mer-ku'-re-al*) [*mercury*]. 1. Pertaining to or caused by mercury. 2. Any preparation of mercury or its salts.

Mercurialism (*mer-ku'-re-al-izm*) [*Mercurius*, a Latin divinity, the god of traffic]. A state of poisoning due to the absorption of mercury.

Mercurialization (*mer-ku'-re-al-iz-a'-shun*) [*Mercurius*, a Latin divinity, the god of traffic]. The act of bringing under the influence of mercury.

Mercuric (*mer-ku'-rik*) [*Mercurius*, a Latin divinity, the god of traffic]. Pertaining to mercury as a bivalent element.

Mercuricum (*mer-ku'-rik-um*) [*mercury*]. Mercury when acting as a bivalent radicle.

Mercurosum (*mer-ku'-ro-sum*) [*mercury*]. Mercury when acting as a univalent radicle.

Mercury (*mur'-ku-re*) [*Mercurius*, a Latin divinity, the god of traffic]. Symbol, Hg; atomic weight, 199.8; quantivalence, II. A shining, silver-white, liquid, volatile metal, having a specific gravity of 13.55. It is insoluble in the ordinary solvents, in hydrochloric acid, and in sulphuric acid in the cold; it dissolves in the last when boiled with it, and is readily soluble in nitric acid. It boils at 357.25°C., and solidifies at -39.40°C. Mercury is found pure, but is chiefly obtained as the native sulphid, or cinnabar. It forms two classes of compounds—the **Mercurous**, those in which two atoms of the metal, and the **Mercuric**, those in which one atom, is combined with a bivalent radicle. The mercuric salts are more soluble and more poisonous than the mercurous. The uses of mercury and its salts are as follows: As a purgative and cholagogue (calomel, blue mass, mercury with chalk), as an alterative in chronic inflammations, as an antisyphilitic, an antiphlogistic, an intestinal antiseptic, a disinfectant, a parasiticide, a caustic, and an astringent. The absorption of mercury in sufficient quantity causes poisoning, characterized by a coppery taste in the mouth, ptyalism, loosening of the teeth, sponginess of the gums; in severer cases, ulceration of the cheeks, necrosis of the jaws, marked emaciation; at times neuritis develops, and a peculiar tremor. The soluble salts when taken in excess act as intense gastrointestinal irritants. Metallic mercury is present in the following preparations: **Emplastrum ammoniaci cum hydrargyro**, is used to disperse indolent swellings, and is applied over the liver in chronic hepatitis. **Emplastrum hydrargyri**, mercurial plaster, has the same uses as the preceding. **Hydrargyrum cum creta**, mercury with chalk. Dose gr. v-ʒ ss (0.32-2.0). **Massa hydrargyri**, blue mass, blue pill. Dose gr. iij (0.20). **Unguentum hydrargyri**, mercurial ointment, blue ointment. **Supposi-**

toria hydrargyri (B. P.). Each contains 5 grains of unguentum hydrargyri. **Hydrargyri benzoas**, $\text{Hg}(\text{C}_6\text{H}_5\text{COO})_2 + \text{H}_2\text{O}$, used in injection in gonorrhea. **Hydrargyri bromidum**, Hg_2Br_2 , mercurous bromid. Dose gr. j (0.065) daily. **Hydrargyri carbolas**, $\text{Hg}(\text{C}_6\text{H}_5\text{O})_2$. Dose, gr. $\frac{1}{4}$ – $\frac{1}{2}$ (0.02–0.032). **Hydrargyri chloridum corrosivum**, HgCl_2 , mercuric chlorid; corrosive sublimate. Dose, as a tonic, gr. $\frac{1}{80}$ – $\frac{1}{60}$ (0.0006–0.0011); as an antisyphilitic, gr. $\frac{1}{80}$ – $\frac{1}{60}$ (0.0011–0.003). It is also a valuable antiseptic. **Hydrargyri chloridum mite**, Hg_2Cl_2 , mild mercurous chlorid; calomel. Dose, as an alternative, gr. $\frac{1}{2}$ –j (0.032–0.065); as a sedative to the stomach and bowels, gr. $\frac{1}{8}$ – $\frac{1}{4}$ (0.008–0.016); as a purgative, gr. $\frac{1}{4}$ – $\frac{1}{2}$ (0.016–0.032) every hour; or gr. v–xv (0.32–1.0). Added to lime-water, 3j to Oj, it forms lotio nigra, or black wash. **Hydrargyri cyanidum**, $\text{Hg}(\text{CN})_2$. Dose, gr. $\frac{1}{2}$ – $\frac{1}{8}$ (0.004–0.008). **Hydrargyri et zinci cyanidum**, $\text{Zn}_2\text{Hg}(\text{CN})_{10}$, is used locally as an antiseptic. **Hydrargyri iodidum flavum**, Hg_2I_2 , yellow mercurous iodid; yellow (or green) iodid of mercury; protoiodid of mercury. Dose, gr. ss (0.033) daily. **Hydrargyri iodidum rubrum**, HgI_2 , red mercuric iodid; biniodid of mercury. Dose, gr. $\frac{1}{8}$ – $\frac{1}{4}$ (0.004–0.016). It is used also in preparing **Liquor arseni et hydrargyri iodidi** (Donovan's solution). Dose $\text{m} \cdot \text{v}$ (0.32). **Hydrargyri iodotannas**, is used hypodermically. **Hydrargyri oxycyanidum**, $\text{Hg}_2\text{O}(\text{CN})_2$, is a local anesthetic. **Hydrargyri oxidum flavum**, HgO , yellow mercuric oxid, is used locally in eye-diseases. It enters into **Unguentum hydrargyri oxidi flavi** and **Oleatum hydrargyri**. **Hydrargyri oxidum rubrum**, HgO , red mercuric oxid, red precipitate, is used locally. It enters into **Unguentum hydrargyri oxidi rubri**, and is also used in preparing **Liquor hydrargyri nitratis**. **Unguentum hydrargyri nitratis**, citrine ointment, is used in chronic skin-diseases. **Unguentum hydrargyri nitratis dilutum** (B. P.), is less irritant. **Hydrargyri peptonas**, is used hypodermically. **Hydrargyri persulphas** (B. P.), HgSO_4 , has no medical uses. **Hydrargyri salicylas**. Dose, gr. $\frac{1}{6}$ (0.001). **Hydrargyri sozodolicum** ($\text{C}_6\text{H}_5\text{I}_2(\text{OH})\text{SO}_3$), Hg , is used locally as an alternative. **Hydrargyri subsulphas flavus**, $\text{Hg}(\text{HgO})_2\text{SO}_4$, yellow mercuric subsulphate; basic mercuric sulphate; turpeth mineral. Dose, as an alternative, gr. $\frac{1}{4}$ – $\frac{1}{2}$ (0.016–0.032); as an emetic, gr. ij–v (0.13–0.32); as an errhine, gr. j (0.065). **Hydrargyri succinimidum** ($\text{C}_4\text{H}_4\text{O}_2\text{N}_2$), Hg , is used hypodermically. **Hydrargyri tannas**. Dose, gr. ij (0.20) daily. **Hydrargyrum ammoniatum**, NH_2HgCl ,

ammoniated mercury; white precipitate; mercuric ammonium chlorid. It is used chiefly locally, in **Unguentum hydrargyri ammoniati**.

Meridian (*mer-id'-e-an*) [*meridies*, midday, from *medius*, middle; *dies*, day]. A great circle surrounding a sphere and intersecting the poles. **M. of the eye**, a line drawn around the globe of the eye and passing through the poles of the vertical axis (**Vertical M.**), or through the poles of the transverse axis (**Horizontal M.**).

Merismopedia (*mer-is-mo-pe'-de-ah*) [*μερισμός*, division; *παῖς*, child]. A bacterium multiplying by two rectangular divisions, thus forming a tablet-like group of four cells in one plane.

Merispore (*mer'-is-por*) [*μέρος*, part; *σπός*, seed]. A spore resulting from division of another spore.

Meroblastic (*mer-o-blas'-tik*) [*μέρος*, a part; *βλαστός*, a germ]. Dividing only in part; as **M. ova**, those in which the process of segmentation is confined to one portion of the ovum.

Merocèle (*mer'-o-sel*) [*μηρός*, thigh; *ἄλῃ*, hernia]. Femoral hernia.

Merycism (*mer'-is-tim*) [*μυρικός*, rumination]. Rumination; chewing the cud,—a normal process in the ruminating animals, and sometimes occurring in man.

Mesad (*mes'-ad*) [*μέσος*, middle; *ad*, to]. Toward the median line or plane.

Mesal (*mes'-al*) [*μέσος*, the middle]. Pertaining to or situated in the middle line or plane.

Mesameboids (*mes-am-e'-hoids*) [*μέσος*, middle; *ἀμοιβή*, a change; *εἶδος*, shape]. 1. Nonepithelial ameboid cells derived from the mesoderm. 2. The leukocytes.

Mesaraic (*mes-ar'-al'-ik*) [*μέσος*, middle; *ἄραια*, belly]. Mesenteric.

Mesarteritis (*mes-ar-ter-i'-tis*) [*μέσος*, middle; *ἀρτηρία*, artery; *τις*, inflammation]. Inflammation of the middle coat of an artery.

Mesaticephalic (*mes-at-e-sef-al'-ik*). See *Index*.

Mesaticephalus (*mes-at-e-sef'-al-us*) [*μεσότης*, median; *κεφαλή*, head]. In craniometry, a term applied to a skull having a cephalic index of between 75 and 79.

Mesencephalon (*mes-en-sef'-al-on*) [*μέσος*, middle; *ἐγκέφαλος*, the brain]. The mid-brain; that part of the brain developed from the middle cerebral vesicle; the corpora quadrigemina, the crura cerebri, and the aqueduct of Sylvius.

Mesenchyma (*mes-enj'-him-ah*) [*μέσος*, middle; *ἐκχρυσ*, an infusion]. The portion of the mesoderm that produces all the connective tissues of the body, the bloodvessels, and the blood, the entire lymphatic system

proper, and the heart; the nonepithelial portions of the mesoderm.

Mesenteric (*mes-en-ter'-ik*) [*μεσεντερικός*, from *μέσος*, middle; *έντερον*, intestine]. Pertaining to the mesentery.

Mesenteritis (*mes-en-ter-it'-is*) [*μέσος*, middle; *έντερον*, an intestine; *ιτις*, inflammation]. Inflammation of the mesentery.

Mesenteron (*mes-en-ter-on*) [*μέσος*, middle; *έντερον*, intestine]. The middle portion of the primitive digestive tube, lined by entoderm, and giving rise to the part of the alimentary tract between the pharynx and the lower third of the rectum.

Mesentery (*mes'-en-ter-e*) [*μέσος*, middle; *έντερον*, bowel]. A fold of the peritoneum that connects the intestine with the posterior abdominal wall; that of the small intestine is termed mesentery proper; that of the colon, cecum, and rectum, mesocolon, mesocecum, mesorectum, respectively.

Mesiad (*mes'-e-ad*) [*μέσος*, middle]. Same as *Mesad*.

Mesial (*mes'-e-al*) [*μέσος*, middle]. Same as *Median*.

Mesmeric (*mes-mer'-ik*) [from *Mesmer*, a German physician]. Pertaining to or induced by mesmerism, as mesmeric sleep.

Mesmerism (*mes'-mer-izm*) [*Mesmer*, a German physician]. Hypnotism.

Meso- (*mes'-o*) [*μέσος*, middle]. A prefix signifying middle.

Mesoblast (*mes'-o-blast*) [*μέσος*, middle; *βλαστός*, a germ]. The middle layer of the blastoderm, probably derived from both the ectoderm and entoderm, and giving rise to the vascular, muscular, and skeletal systems, the generative glands, and the kidneys.

Mesocæcum or Mesocecum (*mes-o-se'-kum*) [*μέσος*, middle; *cæcus*, blind]. The mesentery that in some cases connects the cecum with the right iliac fossa.

Mesocèle (*mes'-o-sel'*) [*μέσος*, middle; *κοιλία*, cavity]. The aqueduct of Sylvius or iter a tertio ad quartum ventriculum.

Mesoccephalon (*mes-o-sef'-al-on*). See *Pons Varolii*.

Mesocolic (*mes-o-kol'-ik*) [*μέσος*, middle; *κόλον*, colon]. Pertaining to the mesocolon.

Mesocolon (*mes-o-ko'-lon*) [*μέσος*, middle; *κόλον*, colon]. The mesentery connecting the colon with the posterior abdominal wall. It is divided into ascending, descending, and transverse portions.

Mesoderm (*mes-o-derm*). See *Mesoblast*.

Mesogaster (*mes-o-gas'-ter*) [*μέσος*, middle; *γαστήρ*, stomach]. The part of the primitive gut giving rise to the duodenum, the liver, the pancreas, the jejunum, and the ileum.

Mesogastric (*mes-o-gas'-trik*) [*μέσος*, middle; *γαστήρ*, stomach]. Pertaining to the umbilical region.

Mesogastrium (*mes-o-gas'-tre-um*) [*μέσος*, middle; *γαστήρ*, stomach]. 1. The umbilical region of the abdomen. 2. A fold of mesentery that in early fetal life connects the stomach with the posterior abdominal wall.

Mesognathic (*mes-og-na'-thik*) [*μέσος*, middle; *γνάθος*, jaw]. 1. Relating to the mesognathion. 2. See *Index*.

Mesognathion (*mes-og-na'-the-on*) [*μέσος*, middle; *γνάθος*, jaw]. The intermaxillary bone.

Mesolobe, Mesolobus (*mes'-o-lób, mes-ol'-o-bus*) [*μέσος*, middle; *λόβος*, lobe]. The corpus callosum.

Meson (*mes'-en*) [*μέσος*, the middle]. The imaginary plane dividing the body into the right and left halves.

Mesonephric (*mes-o-nef'-rik*) [*μέσος*, middle; *νεφρός*, kidney]. Pertaining to the mesonephron.

Mesonephros (*mes-o-nef'-ros*) [*μέσος*, middle; *νεφρός*, kidney]. The Wolffian body, the middle division of the segmental organs. It precedes in the embryo the development of the permanent kidney.

Mesonephron (*mes-o-nef'-ron*) [*μέσος*, middle; *νεφρός*, kidney]. A fold of peritoneum by which a floating kidney is attached to the abdominal wall.

Mesophryon (*mes-off'-re-on*) [*μέσος*, middle; *ὄφρις*, eyebrow]. The glabella.

Mesorchis (*mes-or'-ke-um*) [*μέσος*, middle; *ὄρχις*, testicle]. A fold of the peritoneum containing the fetal testes at about the fifth month of embryonic life.

Mesorectum (*mes-o-rek'-tum*) [*μέσος*, middle; *rectus*, straight]. The narrow fold of the peritoneum connecting the upper part of the rectum with the sacrum.

Mesoretina (*mes-o-ret'-in-ah*) [*μέσος*, middle; *retina*, retina]. The middle layer of the retina, composed of the nuclear and the rod-and-cone layer.

Mesosigmoid (*mes-o-sig'-moid*) [*μέσος*, middle; *σίγμα*, sigma; *εἶδος*, form]. The mesentery of the sigmoid flexure of the colon.

Mesosternum (*mes-o-ster'-num*) [*μέσος*, middle; *στέρνον*, the sternum]. The gladiolus, or second piece of the sternum.

Mesothelium (*mes-o-thel'-le-um*) [*μέσος*, middle; *θηλή*, the nipple]. The lining of the wall of the primitive body-cavity situated between the somatopleure and splanchnopleure. It is the precursor of the endothelium.

Mesovarium (*mes-o-va'-re-um*) [*μέσος*, middle; *ovarium*, ovary]. A peritoneal fold connecting the ovary and the broad ligament.

Meta- (*met'-ah-*) [*μετά*, among or between]. A prefix signifying over, beyond, among, between, change, or transformation.

Metabasis (*met-ab'-as-is*) [*μετά*, over; *βαίνειν*, to go]. Change.

Metabolic (*met-ah-bol'-ik*) [*μεταβολή*, change]. Pertaining to metabolism. **M.** Equilibrium, the equality between the absorption and assimilation of food and the excretion of end-products.

Metabolism (*met-ab'-o-lizm*) [*μεταβολή*, change]. The group of phenomena whereby organic beings transform food-stuffs into complex tissue-elements (Constructive M., assimilation, anabolism) and convert complex substances into simple ones in the production of energy (Destructive M., disassimilation, katabolism).

Metabolite (*met-ab'-o-lit*) [*μεταβολή*, change]. A product of metabolic change.

Metacarpal (*met-ah-kar'-pal*) [*μετά*, beyond; *καρπός*, wrist]. Pertaining to the metacarpus, or to a bone of the metacarpus.

Metacarpophalangeal (*met-ah-kar'-po-fa-lan'-je-al*) [*μετά*, beyond; *καρπός*, wrist; *φάλαγξ*, phalanx]. Belonging to the metacarpus and the phalanges.

Metacarpus (*met-ah-kar'-pus*) [*μετά*, beyond; *καρπός*, wrist]. That part of the hand between the carpus and the phalanges and consisting of five bones.

Metacele, Metacelia (*met'-ah-sel, met-ah-se'-le-ah*) [*μετά*, beyond; *κοιλία*, cavity]. The fourth ventricle.

Metachromatism (*met-ah-krom'-at-ism*) [*μετά*, beyond; *χρώμα*, color]. The quality of being different in color from other parts.

Metakinesis (*met-ah-sin-e'-sis*). Synonym of *Metakinesis*.

Metagaster (*met-ah-gas'-ter*) [*μετά*, beyond; *γαστήρ*, stomach]. The permanent intestinal canal, succeeding the primitive canal, or protogaster.

Metagastrula (*met-ah-gas'-tru-lah*) [*μετά*, beyond; *γαστήρ*, belly]. A modification of segmentation, producing a form of gastrula differing from the simple gastrula of the amphioxus.

Metakinesis (*met-ah-kin-e'-sis*) [*μετά*, toward; *κίνησις*, movement, change]. The term applied to that stage of cell-division, in which the secondary threads or loops tend to pass toward the two poles of the nuclear spindle.

Metal (*met'-al*) [*μέταλλον*, a metal]. An elementary substance characterized by malleability, ductility, fusibility, luster, its electric affinities, and the basic character of its oxid.

Metalbumin (*met-al-but'-min*). See *Paralbumin*.

Metallic (*met-al'-ik*) [*μέταλλον*, metal]. Resembling a metal. **M.** Tinkling, peculiar metallic or bell-like sounds heard over a pneumothorax or large pulmonary cavity. The sounds are produced by coughing, speaking, or deep breathing.

Metalloid (*met'-al-oid*) [*μετάλλον*, metal; *εἶδος*, form]. 1. Resembling a metal. 2. Any nonmetallic element.

Metallioscopy (*met-al-oi'-ko-pe*) [*μέταλλον*, a metal; *σκοπεῖν*, to examine]. The determination of the effects produced by the application of metals to the surface of the body. See also *Metallotherapy*.

Metallotherapy (*met-al-o-ther'-ap-e*) [*μέταλλον*, metal; *θεραπεία*, therapy]. The treatment of certain nervous diseases, particularly hysteria, by the application of different metals to the affected part.

Metameric (*met-am-er'-ik*) [*μετά*, after; *μέρος*, part]. See *Isomeric*.

Metamerid (*met-am'-er-id*) [*μετά*, after; *μέρος*, part]. A metameric substance; a group of metameric bodies.

Metamerism (*met-am'-er-izm*) [*μετά*, after; *μέρος*, a part]. A variety of isomerism. See *Isomeric*.

Metamorphopsia (*met-am-or-fop'-se-ah*) [*μεταμορφόειν*, to change shape; *ὄψις*, sight]. A defect of vision, in which, owing to disease of the retina or imperfection of the media, objects appear distorted.

Metamorphosing (*met-am-or-fop'-zing*) [*μεταμορφόειν*, to change shape]. Altering; changing. **M. Breath-sound**. See *Respiration*.

Metamorphosis (*met-am-or-fop'-sis*) [*μετά*, across; *μορφή*, to change]. A structural change, or transformation. In pathology, a degeneration. **M.** Fatty, fatty degeneration. **M.** Regressive, **M.** Retrograde, a disintegrating change; a degeneration. **M.** Viscous, the agglutination of blood-plaques in the process of thrombosis.

Metanephros, Metanephron (*met-ah-nef'-ros, met-ah-nef'-ron*) [*μετά*, after; *νεφρός*, kidney]. The posterior of the three segmental bodies of the fetus, which is transformed into the permanent kidney and ureter.

Metaphosphoric Acid (*met-ah-fos-for'-ik*) [*μετά*, beyond; *phosphorus*]. See *Acid, Phosphoric*.

Metaplasia (*met-ah-pla'-ze-ah*) [*μετά*, beyond; *πλάσσειν*, to form]. A transformation of a tissue into another without the intervention of an embryonal tissue, as the conversion of cartilage into bone.

Metaplastic (*met-ah-plas'-tik*) [*μετά*, beyond; *πλάσσειν*, to form]. Pertaining to metaplasia.

Metapneumonic (*met-ah-nu-mon'-ik*) [*μετά*, after; *pneumonia*]. Secondary to, or consequent upon, pneumonia.

Metapophysis (*met-ah-poff'-is-is*) [*μετά*, after; *ἀπόφυσις*, offshoot]. A mammillary process, such as is seen upon the lumbar vertebrae.

Metastasis (*met-as'-tas-is*) [*μετάστασις*, from *μεθίσταμαι*, to transpose]. The transfer of a diseased process from a primary focus to a distant one by the conveyance of the causal

agents through the bloodvessels or lymph-channels.

Metastatic (*met'-ah-stat'-ik*) [*μετάστασις*, from *μεθίσταται*, to transpose]. Characterized by or pertaining to metastasis. **M. Abscess**, the secondary abscess in pyemia. **M. Calcification**, calcareous infiltration due to an excess of lime-salts in the blood in diseases associated with rapid disintegration of bone.

Metasternum (*met-ah-ster'-num*) [*μετά*, behind; *στέρνον*, sternum]. The xiphoid cartilage of the sternum.

Metasyphilis (*me-ta-sif'-il-is*) [*μετά*, beyond; *syphilis*, syphilis]. That form of inherited syphilis presenting only the syphilitic diathesis, i. e., the degenerations and general diffuse changes in which localized lesions are absent.

Metatarsal (*met-ah-tar'-sal*) [*μετά*, beyond; *ταρσός*, tarsus]. Pertaining to the metatarsus.

Metatarsophalangeal (*met-ah-tar-so-fa-lan'-je-al*) [*μετά*, beyond; *ταρσός*, tarsus; *φάλαγξ*, phalanx]. Pertaining to the metatarsus and the phalanges.

Metatarsus (*met-ah-tar'-sus*) [*μετά*, beyond; *ταρσός*, tarsus]. The bones of the foot, five in number, situated between the tarsus and the phalanges.

Metathesis (*met-ath'-es-is*) [*μετά*, beyond; *τίθειναι*, to place]. 1. The act of changing the seat of a diseased process from one part to another. 2. In chemistry, decomposition. It is termed single M., when the compound breaks up into its constituents, or exchanges one of them for a molecule of another compound, and double M., when two compounds interchange some of their constituents.

Metazoa (*met-ah-zo'-ah*) [*μετά*, beyond; *ζῷον*, animal]. Animals the development of which is characterized by segmentation of the ovum. They comprise all animals except the Protozoa.

Metencephalon (*met-en-sef'-al-on*) [*μετά*, beyond; *ἐνκέφαλος*, brain]. 1. The after-brain; the postoblongata, or most caudal portion of the brain. 2. Of Huxley, the cerebellum and the pons.

Meteorism (*me'-te-or-izm*) [*μετεωρίζειν*, to elevate]. Distention of the abdomen with gas; tympanites.

Meter (*me'-ter*) [*μέτρον*, measure]. See *Metrie System*.

Methal (*meth'-al*) [*μετά*, with; *ἔλη*, matter], $C_{14}H_{30}O$. An alcohol, not yet isolated, occurring in spermaceti.

Methane (*meth'-an*) [*μετά*, with; *ἔλη*, matter], CH_4 . Marsh-gas. The first member of the homologous series of paraffins, C_nH_{2n+2} . It occurs wherever decomposition of organic matter is going on, especially in marshes, and

is also found at times in the stomach and intestines. It is a colorless, odorless, inflammable gas.

Methemoglobin (*met-hem-o-glo-bin*) [*μετά*, with; *αἷμα*, blood; *globus*, a ball]. A body similar in composition to hemoglobin, but having its oxygen more firmly united with it. It is prepared from hemoglobin by the action of potassium ferricyanid, potassium chlorate, sodium nitrite, etc. In poisoning with potassium chlorate, the nitrites, acetanilid, and some other bodies, the blood contains methemoglobin.

Methemoglobinemia (*met-hem-o-glo-bin-e'-me-ah*) [*μετά*, with; *αἷμα*, blood; *globus*, a ball; *αἷμα*, blood]. The presence of methemoglobin in the blood.

Methemoglobinuria (*met-hem-o-glo-bin-u'-re-ah*) [*μετά*, with; *αἷμα*, blood; *globus*, a ball; *ουρία*, urine]. The presence of methemoglobin in the urine.

Methomania (*meth-o-mal'-ne-ah*) [*μέθυ*, wine; *μανία*, madness]. Insanity due to the excessive use of strong drink.

Methyl (*meth'-il*) [*μετά*, with; *ἔλη*, wood], CH_3 . A univalent hydrocarbon radicle, the first of the univalent hydrocarbons of the marsh-gas series; the radicle of methyl alcohol. **M. Alcohol**, CH_3OH , wood-alcohol, wood-spirit, a liquid similar to ordinary alcohol, obtained in the destructive distillation of wood. **M.-blue**. See *Methylene-blue*. **M. Ether**, C_2H_5O ; dimethyl oxid, an inflammable gas; also, a salt of methyl. **M.-green**, an anilin dye, used in staining tissues; also, as an antiperiodic. **M.-guanidin**, $C_2H_5N_3$; a colorless, crystalline, strongly alkaline base, formed by the oxidation of kreatin and kreatinin, and also found in decomposing horse-flesh, and in cultures of the comma-bacillus and the bacillus of anthrax. It is highly poisonous. **M. hydrid**, CH_4 ; methane, or marsh-gas. **M. salicylas** (U. S. P.), $C_8H_8O_3$; artificial oil of wintergreen; an oily liquid of a peculiar odor, identical with the essential constituent of the oil of wintergreen; it is used in rheumatism like the natural oil of gaultheria. **M.-theobromin**, caffeine. **M.-uramin**. See *M. guanidin*. **M.-violet**, an anilin dye used for staining bacteria. Under the name of pyoktanin it is used as an antiseptic.

Methylal (*meth'-il-al*) [*μετά*, with; *ἔλη*, wood; *alcohol*], $C_3H_8O_2$. A substance prepared by distilling methyl alcohol with sulphuric acid. It is hypnotic and antispasmodic. Dose f ʒj (4.0). Unof.

Methylamin (*meth'-il-am-in*) [*μετά*, with; *ἔλη*, wood; *amin*]. $N(CH_3)H_2$. A colorless basic gas occurring in herring-brine and in cultures of the comma-bacillus.

Methylate (*meth'-il-at*) [*μετά*, with; *ἔλη*, wood]. A compound formed from methyl

alcohol by the substitution of the hydrogen of the hydroxyl by a base.

Methylated (*meth'-il-ated*) [*μετά*, with; *ύλη*, wood]. Containing methyl-alcohol. **M.** Spirit, methyl-alcohol, wood-alcohol.

Methylene (*meth'-il-en*) [*μεθ*, with; *ύλη*, wood], CH_2 . A bivalent hydrocarbon radicle. **M.-blue**, $\text{C}_{16}\text{H}_{10}\text{N}_2\text{S}\text{Cl}$; a blue anilin dye used as a stain in microscopy. It has also been employed as a local application in diphtheria, tonsillitis, scarlatinal sore-throat, and other inflammatory conditions, and internally in malaria and neuralgia. Dose gr. jss (0.1). **M. Dichlorid**, CH_2Cl_2 ; a general anesthetic, used instead of chloroform.

Methylic (*meth'-il'-ik*). Containing methyl. **Metopagus** (*met'-op'-ag-us*) [*μετωπον*, front; *πάγος*, joined]. A twin monstrosity with united foreheads.

Metra (*me'-trah*) [*μήτρα*, uterus]. The uterus.

Metre (*me'-ter*). See *Meter*.

Metrectopia, **Metrectopy** (*met-rek to'-pe-ah*, *met-rek'-to-pe*) [*μήτρα*, womb; *εκτοπος*, displaced]. Displacement of the womb.

Metric (*met'-rik*) [*μετρον*, a measure]. Pertaining to the system of weights and measures, of which the meter is the basis. **M. System**, a decimal system of weights and measures employed in France, Germany, and other countries, and used generally in the sciences. The standard is the meter, the ten-millionth part of the distance from the equator to the north-pole. The actual standard unit is the distance between two lines on a platinum-iridium rod preserved in the archives of the International Metric Commission at Paris, and is equivalent to 39.37079 inches; in the U. S. the length of the meter is assumed as 39.37 inches. The standard of capacity is the liter, a cubical volume $\frac{1}{10}$ meter in each dimension. The standard of weight is the gram, the weight of $\frac{1}{1000000}$ liter (one cubic centimeter) of distilled water at its maximum density. As the unit of microscopic measurement the thousandth part of a millimeter has been adopted. It is called micromillimeter, or micron; its symbol is μ . The multiples in the metric system are expressed by the prefixes deca, hecto, and kilo; the subdivisions by the prefixes deci, centi, and milli.

1000	meters	= 1 kilometer.
100	meters	= 1 hectometer.
10	meters	= 1 decameter.
.1	meter	= 1 decimeter.
.01	meter	= 1 centimeter.
.001	meter	= 1 millimeter.

1000	liters	= 1 kiloliter.
100	liters	= 1 hectoliter.
10	liters	= 1 decaliter.
.1	liter	= 1 deciliter.
.01	liter	= 1 centiliter.
.001	liter	= 1 milliliter.

1000	grams	= 1 kilogram.
100	grams	= 1 hectogram.
10	grams	= 1 decagram.
.1	gram	= 1 decigram.
.01	gram	= 1 centigram.
.001	gram	= 1 milligram.

In common practice, however, the following divisions only are used, the others being expressed in figures:—

10	millimeters	= 1 centimeter.
100	centimeters	= 1 meter.
1000	meters	= 1 kilometer.
1000	cubic centimeters	= 1 liter.
1000	milligrams	= 1 gram.
1000	grams	= 1 kilogram.

The following are the equivalent values:—

1	meter	= 39.37 inches.
1	liter	= 1 quart $\frac{1}{2}$ gill, U. S. measure.
1	gram	= 15.43 grains.
1	minim	= 0.061 cubic centimeter.

See, also, *Weights and Measures*.

Metritis (*met-rit'-tis*) [*μήτρα*, womb; *ιτις*, inflammation]. Inflammation of the uterus.

Metrocele (*met'-ro-sel*) [*μήτρα*, womb; *κήλη*, hernia]. Hernia of the uterus.

Metrocyst (*met'-ro-klist*) [*μήτρα*, womb; *κύστις*, to wash out]. An instrument for giving uterine douches.

Metrocolpocele (*met-ro-kol'-po-sel*) [*μήτρα*, womb; *κόλπος*, vagina; *κήλη*, tumor]. Protrusion of the uterus into the vagina, the wall of the latter being pushed in advance.

Metrocyte (*met'-ro-sit*) [*μήτρα*, womb; *κύτος*, cell]. A large uninuclear spheroidal cell the protoplasm of which contains hemoglobin, and which is supposed to be the source of the red corpuscles of the blood.

Metrodynia (*met-ro-din'-e-ah*) [*μήτρα*, womb; *δύνη*, pain]. Pain in the uterus.

Metropathy (*met-rop'-ath-e*) [*μήτρα*, womb; *πάθος*, suffering]. Any uterine disease.

Metroperitonitis (*met-ro-per-it-on-i'-tis*) [*μήτρα*, womb; *περιτόνιον*, peritoneum; *ιτις*, inflammation]. 1. Combined inflammation of the uterus and the peritoneum. 2. Peritonitis secondary to inflammation of the uterus. 3. Inflammation of the peritoneum about the uterus.

Metrophlebitis (*met-ro-fleb-i'-tis*) [*μήτρα*, womb; *φλέψ*, vein; *ιτις*, inflammation]. Inflammation of the veins of the uterus.

Metrorrhagia (*met-ror-a'-je-ah*) [*μήτρα*, womb; *ῥήγνιναι*, to burst forth]. Uterine hemorrhage independent of the menstrual period.

Metrorrhea (*met-ror-e'-ah*) [*μήτρα*, womb; *ῥοία*, a flow]. Any morbid discharge from the uterus.

Metrorrhexis (*met-ror-ek'-is*) [*μήτρα*, womb; *ῥήξις*, rupture]. Rupture of the uterus.

Metrosalpingitis (*met-ro-sal-ping-jil'-tis*) [*μήτρα*, womb; *σαλπιγξ*, tube; *ιτις*, inflammation]. Inflammation of the uterus and oviducts.

Metroscope (*met'-ro-skōp*) [*μήτρα*, womb; *σκοπεῖν*, to observe]. An instrument for examining the uterus.

Metrostaxis (*met-ro-staks'-is*) [*μήτρα*, womb; *στάξις*, a dropping]. Slight but persistent uterine hemorrhage.

Mezereon (*mez-e'-re-on*) [*Ar.*, *māzarīyūn*, the camellia]. The bark of *Daphne mezereum*, and other species of *Daphne*, of the natural order Thymelacæ. It contains a glucosid, daphnin, and an acrid resin. Locally applied, *M.* is an irritant and vesicant, and has been used to stimulate indolent ulcers. Internally, it has been employed in syphilis, scrofula, chronic rheumatism, and various skin-diseases. Dose of the bark, gr. x (0.65). **Extractum mezerei æthereum** (B. P.) and **Extractum mezerei fluidum** (U. S. P.). Dose mij-v (0.13-0.12). It is used chiefly in ointments.

Miasm (*mi'-azm*) [*μᾶινειν*, to pollute]. Anything harmful contaminating the air; a noxious exhalation affecting individuals directly.

Miasmatic (*mi-as-mat'-ik*) [*μᾶινειν*, to pollute]. Pertaining to or having the nature of miasm. **M. Diseases**, diseases produced by miasms.

Mica (*mi'-kah*) [*L.*]. 1. A crumb. 2. A mineral occurring in the form of thin, shining, transparent scales. **M. panis**, a crumb of bread.

Microacoustic (*mi'-krah-keu'-stik*) [*μικρός*, small; *ἀκουστικός*, pertaining to hearing]. 1. Assisting in hearing very faint sounds. 2. An instrument possessing this property.

Micrencephalon (*mi-kren-sef'-al-on*) [*μικρός*, small; *ἐγκέφαλος*, brain]. 1. A small brain, as in cretinism. 2. The cerebellum.

Micro- (*mi'-kro*) [*μικρός*, small]. 1. A prefix signifying minute. 2. A prefix generally used to signify a unit one-thousandth, sometimes one-millionth part of the unit to which it is prefixed.

Microbe (*mi'-krōd*) [*μικρός*, small; *βίος*, life]. A living organism of very small size. The term is generally used synonymously with bacterium.

Microbian (*mi-kro'-be-an*) [*μικρός*, small; *βίος*, life]. Pertaining to or of the nature of a microbe.

Microbicide (*mi-kro'-bis-id*) [*μικρός*, small; *βίος*, life; *καῖναι*, to kill]. 1. Destructive to microbes. 2. An agent that destroys microbes.

Microbiohemia (*mi-kro-bi-o-hem'-e-ah*) [*μικρός*, small; *βίος*, life; *αἷμα*, blood]. A diseased condition resulting from the presence of microorganisms in the blood.

Microbiology (*mi'-kro-bi-ol'-o-je*) [*μικρός*, small; *βίος*, life; *λόγος*, science]. The science of the nature, life, and actions of microbes.

Microblast (*mi'-kro-blast*) [*μικρός*, small; *βλαστός*, germ]. An immature blood-corpuscle.

Microblepharia, **Microblepharon** (*mi-kro-blef'-a'-re-ah*, *mi-kro-blef'-ar-on*) [*μικρός*, small; *βλέφαρον*, lid]. Smallness of the eyelids.

Microbrachius (*mi-kro-brā'-ke-us*) [*μικρός*, small; *βραχίον*, arm]. Smallness of the arms.

Microcephalic (*mi-kro-sef'-al'-ik*) [*μικρός*, small; *κεφαλή*, head]. Having a small head.

Microcephalon (*mi-kro-sef'-al-on*) [*μικρός*, small; *κεφαλή*, head]. An abnormally small head.

Microchemistry (*mi-kro-kem'-is-tre*) [*μικρός*, small; *chemistry*]. 1. The chemistry of the minute organisms and substances of nature. 2. The study of chemic reactions with the aid of the microscope.

Micrococcus (*mi-kro-kok'-us*) [*μικρός*, small; *κόκκος*, kernel]. A genus of bacteria the individuals of which have a spheric shape. When united in such a way as to resemble a bunch of grapes they are called staphylococci; when united in couples they are called diplococci; when arranged in strings they are called streptococci.

Microcornea (*mi-kro-kor'-ne-ah*) [*μικρός*, small; *cornea*]. Abnormal smallness of the cornea.

Microcosm (*mi'-kro-kōsm*) [*μικρός*, small; *κόσμος*, world]. Man in contradistinction to the universe, or the macrocosm.

Microcosmic (*mi-kro-kōs'-mik*) [*μικρός*, small; *κόσμος*, world]. Pertaining to the microcosm. **M. Salt**, sodium ammonium phosphate; so called because formerly derived from the urine of man, "the microcosm."

Microcoulomb (*mi-kro-kool'-lōm*) [*μικρός*, small; *Coulomb*, a French physician]. One millionth of a coulomb. See *Coulomb*.

Microcoustic (*mi-kro-kōws'-tik*). Synonym of *Microacoustic*.

Microcrith (*mi'-kro-krit*) [*μικρός*, small; *κριθή*, barley]. A unit of molecular weight, equivalent to the weight of an atom of hydrogen.

Microcrystalline (*mi-kro-kris'-tal-in*) [*μικρός*, small; *crystalline*]. Composed of crystals of microscopic size.

Microcyst (*mi'-kro-sist*) [*μικρός*, small; *κύστος*, cyst]. A cyst of very small size.

Microcyte (*mi'-kro-sit*) [*μικρός*, small; *κύτος*, cell]. A small red blood-corpuscle.

Microcythemia (*mi-kro-si-the'-me-ah*) [*μικρός*, small; *κύτος*, cell; *αἷμα*, blood]. A condition of the blood characterized by the presence of abnormally small red corpuscles.

Microcytosis (mi-kro-si-to'-sis). Same as *Microcythemia*.

Microdont (mi'-kro-dont) [μικρός, small; ὀδούς, tooth]. Having small teeth.

Microglossia (mi-kro-glos'-e-ah) [μικρός, small; γλῶσσα, tongue]. Abnormal smallness of the tongue.

Micrognathia (mi-krog-na'-the-ah) [μικρός, small; γνάθος, jaw]. Abnormal smallness of the jaws, especially of the lower jaw.

Microgram (mi'-kro-gram) [μικρός, small; γραμ]. A millionth part of a gram.

Micrography (mi-krog'-ra-fe) [μικρός, small; γράφειν, to write]. A description of bodies that are studied under the microscope.

Microhm (mi'-krôm) [μικρός, small; ὄhm]. The millionth part of an ohm.

Microliter (mi'-kro-le-ter) [μικρός, small; λίτρον, liter]. The millionth part of a liter.

Micromania (mi-kro-ma'-ne-ah) [μικρός, small; mania]. A form of insanity in which the patient believes himself diminutive in size and mentally inferior.

Micromelus (mi-kram'-el-us) [μικρός, small; μέλος, limb]. A single autistic monster of the species ectromelus, characterized by the presence of abnormally small limbs.

Micrometer (mi-kram'-et-er) [μικρός, small; μέτρον, measure]. An instrument designed for measuring minute objects seen through the microscope. **M., Eye-piece**, **M., Ocular**, a micrometer to be used with the eye-piece of a microscope. **M.-screw**, a fine screw with a scale attached showing the distance passed at each fraction of a revolution. **M., Stage-**, a micrometer attached to the stage of a microscope.

Micrometry (mi-kram'-et-re) [μικρός, small; μέτρον, measure]. The measurement of objects by the aid of a micrometer.

Micromillimeter (mi-kro-mil'-im-e-ter) [μικρός, small; millimeter]. 1. The one-thousandth part of a millimeter or the one-millionth part of a meter. It is the unit of microscopic measurements. It is called also **micron**, and is symbolized by μ .

Micron (mi'-kron) [μικρός, small]. The millionth part of a meter or a thousandth part of a millimeter. It is the equivalent of $\frac{1}{25400}$ of an English inch, and its symbol is μ .

Microorganism (mi-kro-or'-gan-izm) [μικρός, small; organism]. A microscopic being of the animal or vegetal kingdom; in a special sense it is restricted to that vegetal group known as bacteria.

Micropathology (mi-kro-path-al'-o-je) [μικρός, small; pathology]. 1. The study of minute pathologic changes. 2. The study of microorganisms in their relation to disease.

Microphage (mi'-kro-faj) [μικρός, small; φάγειν, to devour]. A small phagocyte.

Microphone (mi'-kro-fôn) [μικρός, small;

φωνή, sound]. An instrument that amplifies feeble sounds and renders them audible.

Microphotograph (mi-kro-fa'-to-graf) [μικρός, small; φῶς, light; γράφειν, to write]. 1. A photograph of microscopic size. 2. See *Photomicrograph*.

Microphthalmus (mi-krof'-thal'-mus) [μικρός, small; ὀφθαλμός, eye]. 1. The condition of having an abnormally small eye. 2. A person having such an eye.

Microphyte (mi'-kro-fu) [μικρός, small; φυτόν, plant]. Any microscopic plant, especially one that is parasitic.

Micropsia (mi-krop'-se-ah) [μικρός, small; ὄψις, sight]. A defective state of vision in which objects appear very small.

Micropyle (mi'-kro-pil) [μικρός, small; πύλη, gate, orifice]. The small opening in an ovum through which the spermatozoon may penetrate.

Microscope (mi'-kro-skôp) [μικρός, small; σκοπεῖν, to view]. An apparatus through which minute objects are rendered visible.

It consists of a lens or group of lenses by which a magnified image of the object is produced.

M., Binocular, a microscope having divergent oculars, one for each eye, so that the object is seen with both eyes. **M., Compound**, one that consists of two or more lenses or lens-systems, of which one, the objective, placed near the object, gives an enlarged and inverted real image; the other, the ocular, acting like a simple microscope, gives an enlarged virtual image of the real image. **M., Simple**, one consisting of one or more lenses or lens-systems acting as a single lens. The rays of light that enter the eye of the observer, after refraction through these lenses, proceed directly from the object itself.

Microscopic (mi-kro-skop'-ik) [μικρός, small; σκοπεῖν, to view]. 1. Pertaining to the microscope. 2. Visible only with the aid of a microscope.

Microscopist (mi-kros'-ko-pist) [μικρός, small; σκοπεῖν, to view]. One who is skilled in the use of the microscope.

Microscopy (mi-kros'-ko-pe) [μικρός, small; σκοπεῖν, to view]. The use of the microscope; examination with the microscope.

Microseme (mi'-kro-stem) [μικρός, small; σημα, sign, index]. Having the orbital index less than 83.

Microsomia (mi-kro-sol'-me-ah) [μικρός, small; σῶμα, body]. Abnormal smallness of the whole body.

Microspectroscope (mi-kro-spek'-tro-skôp) [μικρός, small; spectrum, spectrum; σκοπεῖν, to view]. A spectroscope used in connection with the ocular of a microscope, and by means of which the spectra of microscopic objects can be examined.

Microsporidia (*mī-kro-spor-id' -e-ah*) [*μικρός*, small; *σπόρος*, seed]. A genus of the class of sporozoa, occurring as parasites in the muscles of the frog, the marsh-tortoise, worms, and insects.

Microsporion (*mī-kro-spo' -ron*) [*μικρός*, small; *σπόρος*, seed]. A fungus to which several diseases of the skin and hair are believed to be due. **M. furfur**. See *Tinea versicolor*.

Microstomia (*mī-kro-sto' -me -ah*) [*μικρός*, small; *στόμα*, mouth]. Abnormal smallness of the mouth.

Microtome (*mī' -kro-tōm*) [*μικρός*, small; *τέμνειν*, to cut]. An instrument for making thin sections for microscopic examination. **M.**, **Freezing**, one in which the tissue is frozen, in order to secure the hardness required for properly cutting sections.

Microtomy (*mī-kro' -o-me*) [*μικρός*, small; *τέμνειν*, to cut]. Section-cutting.

Microvolt (*mī' -kro-vōlt*). One millionth of a volt.

Microzyme (*mī' -kro-zīm*) [*μικρός*, small; *ζῆν*, to live]. One of certain minute particles of living matter that are by some supposed to be living organisms capable of an independent existence, and which are the cause of normal and pathologic fermentation; the real agents of the functions of the organism, the perversion of whose function constitutes disease.

Micturition (*mīk -tu -rish' -un*) [*micturire*, to pass water]. The act of passing urine. **M.**-**center**. The center governing the act of micturition; it is situated in the lumbar region of the spinal cord.

Midbrain [**AS.**, *mid*, middle; *brain*]. The mesencephalon.

Midfrontal [**AS.**, *mid*, middle; *frons*, forehead]. Pertaining to the middle of the forehead.

Midgut [**AS.**, *mid*, middle; *gut*, gut]. See *Mesogaster*.

Midriff [**AS.**, *mid*, middle; *hrif*, belly]. Diaphragm.

Midwife [**AS.**, *mid*, with; *wif*, a woman]. A female obstetrician; a female nurse who attends women in childbirth.

Midwifery [**AS.**, *mid*, with; *wif*, a woman]. Obstetrics.

Migraine (*mē' -grān*) [**Fr.**, from *ἡμι*, half; *κρᾶνιον*, skull]. A paroxysmal affection characterized by headache, usually unilateral, and by gastric, vasomotor, and visual disturbances.

Migrainin (*mīg' -ra -nin*) [*migraine*]. A name given to a mixture of antipyrin, citric acid, and caffeine. It is used for the treatment of migraine, of the headache of influenza, and of that due to alcohol, tobacco, and morphin. The dose is from 17 to 45 grains (1.0-3.0) in 24 hours.

Migration (*mī-gra' -shun*) [*migrare*, to wander]. A wandering. **M.**, **External** (*of the Ovum*), the passage of the ovum from an ovary to the tube of the opposite side. **M.**, **Internal** (*of the Ovum*), the passage of the ovum through the tube related to the ovary from which the ovule was discharged, into the uterus and across into the opposite tube. **M. of Ovum**, the passage of the ovum from the ovary to the Fallopian tube. **M. of White Corpuscles**, one of the phenomena of inflammation, consisting in the passage of the white corpuscles of the blood through the vessel-wall.

Mildew (*mī' -du*) [**AS.**, *meledēw*, honey-dew]. A common name for minute fungi parasitic on plants, and also found on dead vegetable substances.

Milfoil (*mī' -fōil*). See *Achillea*.

Miliaria (*mī' -e-a' -re-ah*) [*milium*, millet]. An acute inflammatory disease of the sweat-glands, the lesions consisting of vesicles and papules, accompanied by a pricking or tingling sensation. It occurs especially in summer, is due to excessive sweating, runs an acute or subacute course, and is followed by slight desquamation. Relapses are common. **M. alba** or **M. crystallina**, a variety of **M.** in which the sweat accumulates under the superficial horny layers of the epidermis to form small, clear, transparent vesicles. It is also called sudamina crystallina. **M. papulosa**, the well-known "prickly heat." **M. rubra**. See *Miliaria*.

Miliary (*mī' -e-a-re*) [*milium*, millet]. 1. Of the size of a millet-seed, as **M. aneurysm**, **M. tubercle**. 2. Attended or characterized by the formation of numerous lesions the size of a millet-seed, as **M. tuberculosis**. **M. Tuberculosis**. See *Tuberculosis*.

Milium (*mī' -e-um*) [**L.** millet-seed]. 1. A disease of the skin characterized by the formation of small, pearly, noninflammatory elevations (milia) situated mainly on the face. It is due to the occlusion of the ducts of sebaceous follicles, the secretion of which accumulates and distends the follicles. 2. One of the elevations characteristic of milium.

Milk [**AS.**, *mīlc*, milk]. The opaque white secretion of the mammary glands of the female of mammalia. The relative proportion of the constituents in milk of different animals may be seen from the appended table of analyses from Wynter Blyth. **M.**, **Butter-**, cream from which the fatty matter has been removed. **M.**, **Condensed**, milk of cows from which a large part of the water has been evaporated, a syrupy liquid remaining which is preserved with or without the addition of sugar. **M.**-**cure**, the method of treating certain diseases by an exclusive

diet of milk. **M.-fever**, a slight rise of temperature attending the establishment of the secretion of milk. It is due to a mild degree of septic intoxication. **M.-leg**. See *Phlegmasia alba dolens*. **M.-punch**, a preparation made by adding brandy, whisky, or rum, to milk in the proportion of about one to four or six parts, and flavoring with sugar and nutmeg. **M.**, Skim-, is the residue left after the removal of the cream. **M.-sugar**. Lactose. **M.-teeth**, the teeth of the first dentition. **M.-tester**, a lactometer.

	FAT.	CASEIN.	ALBUMIN.	MILK-SUGAR.	ASH.	TOTAL SOLIDS.	WATER.
Human milk,	3.90	2.40	0.57	5.87	0.16	12.00	88.00
Cow's milk,	3.50	3.98	0.77	4.00	0.17	13.13	86.87
Camel's milk,	2.90	3.84		5.66	0.66	13.06	86.94
Goat's milk,	4.20	3.00	0.62	4.00	0.56	12.46	87.54
Ass's milk,	1.02	1.09	0.70	5.50	0.42	8.83	91.17
Mare's milk,	2.50	2.19	0.42	5.50	0.50	11.20	88.80
Sheep's milk,	5.30	6.10	1.00	4.20	1.00	17.73	82.27

Milliampere (*mil-e-am-pär'*) [*mille*, thousand; *Ampère*, a French scientist]. One-thousandth of an ampere, = 1 volt divided by 1000.

Milliamperemeter (*mil-e-am-pär'-me-ter*) [*mille*, thousand; *Ampère*, a French scientist; *μέτρον*, measure]. An instrument for measuring the strength of an electric current.

Milligram (*mil'-ig-ram*) [*mille*, a thousand; *gram*]. A thousandth part of a gram.

Milliliter (*mil'-il-e-ter*) [*mille*, a thousand; *liter*]. A thousandth part of a liter.

Millimeter (*mil'-im-e-ter*) [*mille*, a thousand; *meter*]. A thousandth part of a meter.

Millon's Reagent. A reagent made by dissolving 10 grams of mercury in 20 grams of nitric acid, diluting the solution with an equal volume of water, and decanting in twenty-four hours. With proteids and with derivatives of benzene and naphthalene it gives a red color (**Millon's Test**).

Millstone-makers' Phthisis. A form of pneumonokoniosis due to the inhalation of fine particles in the manufacture of millstones.

Milzbrand (*milts'-brand*) [Ger.]. Anthrax.

Mimetic (*mim-et'-ik*) [*μῖμος*, an actor]. Imitative; mimic. **M.** Paralysis, paralysis of the facial muscles. **M.** Spasm, spasm of the facial muscles.

Mimic (*mim'-ik*). See *Mimetic*.

Mimicry (*mim'-ik-re*) [*μῖμος*, an actor]. Imitation.

Mind-blindness. A form of aphasia in which, although the patient is able to see, no intellectual impression is conveyed to his mind by the object seen. The condition is also called visual amnesia.

Mind-cure. The alleged cure of disease through mental operations.

Mind-deafness. A form of aphasia in which sounds, though heard and perceived as such, awaken no intelligent conception.

Mindererus, Spirit of. Liquor ammonii acetatis. See *Ammonia*.

Miner's Elbow. Enlargement of the bursa over the olecranon, common in miners, due to irritation while working and lying on the side.

Miner's Phthisis. Anthracosis; a chronic affection of the lungs due to the constant inhalation of coal-dust.

Mineral (*min'-er-al*) [*minera*, a mine]. An inorganic chemic compound found in nature, especially one that is solid. **M.**, Kermes, antimony oxysulphid. **M.** Oil, petroleum. **M.** Pitch, bitumen. **M.** Water, water naturally or artificially impregnated with inorganic salts in sufficient quantity to give it special properties.

Mineralization (*min-er-al-e-za'-shun*) [*minera*, mine]. 1. The addition of mineral substances to a body. 2. The relative amount of mineral substances dissolved in a mineral water.

Minim (*min'-im*) [*minimus*, least]. The one-sixtieth of a fluid dram. Symbol \mathfrak{m} .

Minimal (*min'-im-al*) [*minimus*, least]. Least, lowest. Of doses, the least quantity that is yet effective.

Minimum (*min'-im-um*) [*minimus*, least]. The least, the lowest; the lowest intensity, or level. **M.** Thermometer. See *Thermometer*.

Minium (*min'-e-um*) [L., *minium*, red lead]. Red lead oxid, Pb_2O_3 , used formerly in plasters.

Mint. See *Mentha*.

Miosis (*mi-o'-sis*) [*μειωσις*, less]. Contraction, especially of the pupil.

Miotic (*mi-ot'-ik*) [from *μειωσις*, less]. 1. Pertaining to, or characterized by, miosis. 2. Causing contraction of the pupil. 3. An agent that contracts the pupil.

Mirror (*mir'-or*) [*mirari*, to admire]. 1. A polished surface for reflecting light or forming images of objects placed in front of it. **M.**, Concave, one the reflecting surface of which is concave. **M.**, Convex, one with a convex reflecting surface. **M.**, Frontal, **M.**, Head-, a circular mirror with a central perforation, strapped to the head by a band, and used to throw light on parts to be examined.

M., Laryngoscopic, **M.**, Ophthalmoscopic, **M.**, Rhinoscopic; a mirror used respectively in laryngoscopy, ophthalmoscopy, and rhinoscopy, to illuminate the cavity to be inspected. **M.**, Plane, one the reflecting surface of which is flat. **M.**-writing, a peculiar form of writing at times observed in left-handed persons, and in cases of aphasia, and characterized by a reversal of the form and arrangement of the letters, which appear as if seen in a mirror.

Miryachit (*me-re-ach'-it*) [Russian word]. A peculiar disease in which the patient mimics or imitates everything said or done by another.

Miscarriage (*mis-kar'-aj*) [A S., prefix *mis-*, OF., *carrier*, to carry]. 1. The expulsion of the fetus during the fourth and the sixth month of pregnancy. 2. Abortion.

Misce (*mis'-e*) [*miscere*, to mix]. Mix, a direction placed on prescriptions; abbreviated, **M**.

Miscegenation (*mis-je-en-a'-shun*) [*miscere*, to mix; *genus*, race]. Mixture of different races by intermarriage.

Miscible (*mis'-ib-l*) [*miscere*, to mix]. Capable of being mixed.

Miserere mei (*miz-er-a'-ra me'-e*) [L., have mercy on me]. An old name for volvulus, or intestinal colic; also for stercoraceous vomiting.

Misogamy (*mis-og'-am-e*) [*μισος*, to hate; *γάμος*, marriage]. Aversion to marriage.

Misogyny (*mis-og'-in-e*) [*μισος*, to hate; *γυνή*, woman]. Hatred of women.

Missed (*mist*) [A S., *missan*, to miss]. Passed; failed of completion. **M.** Abortion, the retention of the product of conception in the uterine cavity after its death and with the appearance of some of the symptoms of abortion. **M.** Labor, the retention of the product of conception in the uterus beyond term, and after the occurrence of a few ineffectual labor-pains.

Mistletoe (*mis'-l-to*). See *Viscum*.

Mistura (*mis-tu'-rah*) [L.]. A mixture. A preparation made by suspending an insoluble substance in watery fluids, by means of gum-arabic, sugar, yolk of egg, or other cohesive substance. When the suspended substance is of an oily nature, the preparation is termed an emulsion (*emulsum*). In pharmacy, an aqueous preparation of an insoluble substance held in suspension, usually by the addition of soluble gum, egg-emulsion, etc. The following mixtures are official in U. S. P. and B. P. **M.** Creasoti (B. P.). Dose $\mathfrak{f}\overline{\mathfrak{z}}$ j (32.0). **M.** Cretæ (U. S. P., B. P.). Dose $\mathfrak{f}\overline{\mathfrak{z}}$ ss (16.0). **M.** ferri aromatica (B. P.). Dose $\mathfrak{f}\overline{\mathfrak{z}}$ j-ij (32.0-64.0). **M.** ferri composita, Griffith's mixture (U. S. P. and B. P.). Dose $\mathfrak{f}\overline{\mathfrak{z}}$ j-ij (32.0-64.0). **M.** glycyrrhizæ

composita (U. S. P.). Brown mixture. Dose $\mathfrak{f}\overline{\mathfrak{z}}$ j-ij (32.0-64.0). **M.** guaiaci (B. P.). Dose $\mathfrak{f}\overline{\mathfrak{z}}$ ss-ij (16.0-64.0). **M.** olei ricini (B. P.). Dose $\mathfrak{f}\overline{\mathfrak{z}}$ ss-ij (16.0-64.0). **M.** rhei et sodæ (U. S. P.). Dose, for children, $\mathfrak{f}\overline{\mathfrak{z}}$ ss- $\mathfrak{f}\overline{\mathfrak{z}}$ j (2.0-4.0). **M.** scammonii (B. P.). Dose $\mathfrak{f}\overline{\mathfrak{z}}$ j (64.0). **M.** sennæ composita (B. P.). Dose $\mathfrak{f}\overline{\mathfrak{z}}$ j-iss (32.0-48.0). **M.** spiritus vini gallici (B. P.). Dose $\mathfrak{f}\overline{\mathfrak{z}}$ j-ij (32.0-64.0).

Mite (*mit*) [A S., *mlte*, mite]. A name applied to several Acari.

Mithridatism (*mith-rid'-at-izm*) [*Μιθριδάτης*, Mithridates, king of Pontus, who was said to have become so charged with the poisons with which he experimented that he acquired an immunity to them all]. Immunity from the effects of a poison induced by the administration of gradually increased doses.

Mitigate (*mit'-ig-at*) [*mitigare*, to soften]. To allay; to make milder; to moderate.

Mitigated Stick, or Mitigated Caustic (*mit'-ig-a-ted*) [*mitigare*, to soften]. See *Argentum*.

Mitome (*mi'-tōm*) [*μῑτός*, a thread]. The threads of the protoplasmic reticulum of a cell (cytomitome) or of the nucleus (karyomitome).

Mitosis (*mi-to'-sis*) [*μῑτός*, a thread]. Karyokinesis.

Mitotic (*mi-tol'-ik*) [*μῑτός*, thread]. Pertaining to mitosis.

Mitral (*mi'-tral*) [*μῑτρα*, a belt, a turban].

1. Resembling a miter, as the **M.** valve. 2. Pertaining to the auriculoventricular valve of the left side of the heart. **M.** Disease, disease of the mitral valve of the heart.

M. Incompetence or **M.** Insufficiency.

See *M. Regurgitation*. **M.** Murmur. See *Murmur*.

M. Obstruction, disease of the mitral valve causing obstruction to the flow of blood through the left auriculoventricular opening. **M.** Regurgitation, imperfect closure of the mitral valve during the cardiac systole, permitting blood to be forced back into the left auricle. **M.** Stenosis. See *M. Obstruction*. **M.** Valve. See *Valve*.

Mixture (*miks'-tūr*). See *Mistura*.

Mnemonics (*ne-mon'-iks*) [*μνῑσμός*, I remember]. The science of cultivation of the memory by systematic methods.

Mobile (*mo'-bil*) [*movere*, to move]. Movable.

M. Spasm, a slow, irregular movement gradually taking place in different muscles, occurring at times in the paralyzed parts in hemiplegia.

Mobility (*mo-bil'-it-e*) [*movere*, to move]. The condition of being movable.

Modal (*mo'-dal*) [*modus*, mode]. A term applied to the order of response of muscles and nerves to the galvanic current.

Moderator Band. See *Band*.

Modiolus (*mo-di' o-lus*) [L., nave]. 1. The central pillar or axis of the cochlea, around which the spiral canal makes two and one-half turns. 2. The crown of a trephine.

Modus (*mo'-dus*) [L.]. A mode or method. **M. operandi**, the method of the performance of an action.

Mogigraphia (*mog-ig-raf'-e-ah*) [*μῳγῖς*, with difficulty; *γράφειν*, to write]. Writers' cramp.

Mogilalia (*mog-il-a'-le-ah*) [*μῳγῖς*, with difficulty; *λαλῶν*, talk]. Stammering, stuttering.

Mogiphonia (*mog-if-o'-ne-ah*) [*μῳγῖς*, with difficulty; *φωνή*, sound]. Difficulty in speaking, excited by an effort of singing or speaking loudly.

Mohrenheim's Fossa. 1. The space between the deltoid and pectoralis major muscles. 2. The supraclavicular fossa.

Moist [*mustus*, young, fresh, new]. Damp; slightly wet; characterized by the presence of fluid. **M. Chamber**, a large circular glass with a lid, used in bacteriologic work, especially for growing potato-cultures. **M. Filter**, a filter paper that is moistened with water. **M. Gangrene**, the form of gangrene that occurs in a part filled with blood. See *Gangrene*.

Molar (*mol'-lar*) [*moles*, mass]. Pertaining to masses, in contradistinction to molecular.

Molar (*mol'-lar*) [*mola*, a millstone]. 1. Grinding; used for grinding. 2. A grinding tooth, a grinder.

Molasses (*mo-las'-see*) [*melleo*, made with honey]. The syrupy liquid remaining after the refining of sugar. It contains a considerable quantity of uncrystallizable sugar, some cane-sugar, and gummy and coloring-matter. There are two kinds: West India M., from which rum is prepared, and sugar-house M., which is somewhat thicker than the first. M. is used for making pills, and combined with sulphur as a domestic remedy for constipation.

Mold (*mol'd*) [AS., *molde*, dust]. A variously colored deposit produced by the growth of different forms of fungi on moist surfaces. The principal molds are *Penicillium*, the *Mucorini*, and *Aspergillus*.

Mole (*mol*) [*mola*, a mass]. 1. A mass formed in the uterus by an ovum, the growth of which has become arrested, or which has undergone degeneration. 2. **Nevus**. **M., Blood-**, a mass of coagulated blood and retained fetal membranes and placenta, sometimes found in the uterus after an abortion. **M., Carneous**. See *M., Fleshy*. **M., Cystic**. Same as *M., Hydatidiform*. **M., False**, one not containing any tissues derived from the ovum. **M., Fleshy**, a blood-mole which has become more solid and has assumed a fleshy appearance; the body formed in the uterine cavity when an ovum that has died is retained with-

in the uterus for some time. **M., Hydatid**, **M., Hydatidiform**, one formed by a proliferation and cystic degeneration of the chorion villi; it is a form of myxoma, and has a tendency to involve the uterine wall. **M., True**, one which is the remains of an ovum. **M., Vesicular**. Same as *M., Hydatidiform*. **Molecular** (*mo'-lek'-u-lar*) [*moles*, mass].

Pertaining to, or composed of, molecules.

M. Death, death of a part in minute invisible particles, as ulceration, caries. **M., Force**, a force acting between molecules, as cohesion.

M. Motion, the movements of the molecules of a substance. **M. Volume**, the volume of a molecule of a substance in the gaseous state; under the same conditions of temper-

ature and pressure the M. volumes of all substances are equal. **M. Weight**, the weight of a molecule of any substance as compared with the weight of an atom of hydrogen. It is equal to the sum of the weights of its constituent atoms.

Molecule (*mol'-e-kul*) [dim. of *moles*, mass].

1. A minute portion of matter. 2. In physics, the smallest quantity into which a substance can be divided and retain its characteristic properties; or the smallest quantity that can exist in a free state.

Molimen (*mo-li'-men*) [L., *molimen*, endeavor]. An effort or attempt. **M., Menstrual**, any of the symptoms attendant upon the menstrual act or function.

Mollin (*mol'-in*) [*mollis*, soft]. A soft soap of potassium hydroxid and coconut-oil, used as a basis for ointments.

Mollities (*mo-lish'-e-ee*) [*mollis*, soft]. Softness. **M. ossium**. See *Osteomalacia*.

Molluscous (*mol-us'-kus*) [*molluscum*, a shell-fish]. 1. Pertaining to the Mollusca. 2. Pertaining to the disease Molluscum.

Molluscum (*mol'-us'-kum*) [L., *mollusca*, shell-fish, from *mollis*, soft]. A term applied to several diseases of the skin. **M. contagiosum**, a disease of the skin characterized by the formation of pinhead-sized to pea-sized, rounded, sessile, or pedunculated pearl like elevations of a yellowish-white or pinkish color. The lesions may be single or multiple, are usually situated upon the face, and are due to a hyperplasia of the rete mucosum, the growth probably beginning in the hair-follicles. The lesions on microscopic examination are found to contain peculiar ovoidal, sharply-defined bodies—*Molluscum-bodies*—which are by some considered as forms of epithelial degeneration, by others as protozoan parasites. **M. fibrosum**, a disease of the skin characterized by the formation of multiple fibromata, which may be sessile or pedunculated, and grow from the deeper layers of the corium and the subcutaneous tissue.

Molybdenum (*mol'-ib'-de'-num*) [*μόλυβδος*,

lead]. A metallic element, found in nature chiefly as the sulphid molybdenite (MoS_2). Atomic weight, 96; symbol, Mo; quantivalence, II, IV, VI, VIII; specific gravity, 8.6. Its principal oxid, MoO_3 , forms molybdic acid, H_2MoO_4 , the ammonium salt of which is used as a reagent in metallurgy, etc. Molybdic acid combines with phosphoric acid to form phosphomolybdic acid.

Momentum (*mo-men'-tum*) [*movere*, to move]. Quantity of motion. The M. of a body depends upon its mass and velocity. Also, the quantity of potential energy possessed by a body in motion. It is usually expressed by the formula $m = wz$, i. e., the momentum equals the weight multiplied by the velocity.

Momordica (*mo-mor'-dik-ah*). See *Elaeterium*.

Monad (*mon'-ad*) [*μόνος*, single]. A univalent element or radicle.

Monamin (*mon'-am-in*) [*μόνος*, single; *amin*]. An amin formed by the replacement of the hydrogen in one atom of ammonia by basic radicles.

Monarda (*mo-nar'-dah*) [after N. Monardès, a Spanish physician]. A genus of labiate plants, comprising *M. bistulosa*, wild bergamot, *M. didyma*, bee-balm, and *M. punctata*, horse-mint. The last is diaphoretic, carminative, and stimulant. It is also a source of thymol.

Monas (*mon'-as*) [*μονάς*, unit]. A genus of infusorians.

Monaster (*mon-as'-ter*) [*μόνος*, single; *ἀστήρ*, star]. Mother-star. See *Karyokinesis*.

Monatomic (*mon-at-om'-ik*) [*μόνος*, single; *ἄτομος*, atom]. 1. Having but one atom of replaceable hydrogen, as a M. acid. 2. Having only one atom, as a M. molecule. 3. Having the combining power of one atom of hydrogen, as a M. radicle. 4. Formed by the replacement of one hydrogen-atom in a compound by a radicle, as a M. alcohol.

Monerula (*mon-er'-u-lah*) [*μονήρης*, single]. The impregnated ovum at a stage when it has no nucleus.

Monilethrix (*mo-nil'-eth-riks*) [*monile*, a necklace; *ῥιζή*, hair]. An affection of the hair in which nodes are strung regularly or irregularly along the hair-shaft, giving it a beaded appearance.

Monilliform (*mon-ill'-if-orm*) [*monile*, necklace; *forma*, form]. Shaped like a necklace; beaded or bead-like, resembling a string of beads.

Monilithrix (*mo-nil'-ith-riks*). See *Monilethrix*.

Monkshood (*munk'-hood*). See *Aconite*.

Mono- (*mon'-o-*) [*μόνος*, single]. A prefix signifying one or single.

Monobasic (*mon-o-ba'-sik*) [*μόνος*, single; *βάσις*, foundation]. Of an acid, acid salt, or alcohol, having one hydrogen-atom replaceable by a base.

Monoblepsia, **Monoblepsis** (*mon-o-blep'-se-ah*, *mon-o-blep'-sis*) [*μόνος*, single; *βλέψω*, sight]. 1. A condition in which either eye has a better visual power than both together. 2. The form of color-blindness in which but one color can be perceived.

Monobrachius (*mon-o-bra'-he-us*) [*μόνος*, single; *βραχίον*, arm]. A monster having but one arm.

Monobromated (*mon-o-bro'-ma-ted*) [*μόνος*, one; *bromate*]. Containing one atom of bromin in the molecule.

Monobromid (*mon-o-bro'-mid*) [*μόνος*, one; *bromid*]. A compound having one atom of bromin in the molecule, or containing an amount of bromin which, when compared with the amount of bromin in other bromids of the same base, may be regarded as unity.

Monocephalus (*mon-o-sef'-al-us*) [*μόνος*, single; *κεφαλή*, head]. A monster consisting of a single head with two bodies more or less completely fused.

Monochlorid (*mon-o-klo'-rid*). See *Monobromid*.

Monochorea (*mon-o-ko-re'-ah*) [*μόνος*, single; *chorea*]. Chorea confined to a single member or part of the body.

Monochroic (*mon-o-kro'-ik*) [*μόνος*, single; *χρῶς*, color]. Having only one color. Arterial blood is monochroic.

Monochromatic (*mon-o-kro-mat'-ik*). See *Monochroic*.

Monocle (*mon'-o-kl*) [*μόνος*, single; *oculus*, eye]. 1. A lens for one eye only. 2. A bandage for one eye.

Monococcus (*mon-o-kok'-us*) [*μόνος*, single; *κόκκος*, grain]. A coccus occurring in single individuals; one not united in chains or pairs.

Monocranus (*mon-ok'-ran-us*) [*μόνος*, single; *κρανίον*, cranium]. A double monster having a single cranium.

Monocular (*mon-ok'-u-lar*) [*μόνος*, single; *oculus*, eye]. 1. Pertaining to or affecting only one eye, as M. diplopia; performed with one eye only, as M. vision. 2. Having a single ocular or eye-piece, as a M. microscope.

Monogastric (*mon-o-gas'-trik*) [*μόνος*, single; *γαστήρ*, stomach]. Having one stomach or one belly.

Monogenesis (*mon-o-jen'-es-is*) [*μόνος*, single; *γενεσις*, origin]. 1. Development of offspring resembling the parent, as distinguished from metagenesis. 2. Development from a single hermaphroditic parent; asexual reproduction. 3. Origin of all organisms from a single cell.

Monohydric (*mon-o-hi'-drik*) [*μόνος*, single; *ὕδωρ*, water]. Containing one atom of replaceable hydrogen, as M. acid, M. alcohol.

Monoiodid (*mon-o-i'-o-did*). See *Monobromid*.

Monolocular (*mon-o-lok'-u-lar*). See *Unilocular*.

Monomania (*mon-o-ma'-ne-ah*) [*μόνος*, single; *μανία*, madness]. A form of insanity characterized by a limited disturbance of the mental functions that dominates the person's thought and actions.

Monomaniac (*mon-o-ma'-ne-ak*) [*μόνος*, single; *μανία*, madness]. A person affected with monomania.

Monomphalus (*mon-om'-fal-us*) [*μόνος*, single; *ὀμφαλός*, navel]. A double monster united by a common umbilicus.

Mononuclear (*mon-o-nu'-kle-ar*). See *Uninuclear*.

Monoparesis (*mon-o-par'-es-is*) [*μόνος*, single; *πάρεσις*, paresis]. Paralysis of a single part of the body, as of one limb.

Monophasia (*mon-o-fa'-se-ah*) [*μόνος*, single; *φάναι*, to speak]. A form of aphasia, in which speech is limited to a single syllable, word, or phrase.

Monophobia (*mon-o-fo'-be-ah*) [*μόνος*, single; *φόβος*, fear]. Morbid dread of being alone.

Monoplasmatic (*mon-o-plas-mat'-ik*). See *Monoplastic*.

Monoplast (*mon'-o-plast*) [*μόνος*, single; *πλάσσειν*, to form]. A simple cell.

Monoplastic (*mon-o-plas'-tik*) [*μόνος*, single; *πλάσσειν*, to form]. Composed of only one substance.

Monoplegia (*mon-o-ple'-je-ah*) [*μόνος*, single; *πληγή*, stroke]. Paralysis of a single limb or of a single muscle or group of muscles. It is designated as Brachial, Crural, or Facial, when affecting the arm, the leg, or the face, respectively, and as Central (Cerebral), or Peripheral, according to the seat of the causal lesion.

Monops (*mon'-ops*). See *Cyclops*.

Monorchid, **Monorchis** (*mon-or'-kid*, *mon-or'-kis*) [*μόνος*, single; *ὄρχις*, testis]. A person who has but one testicle, or in whom one testicle only has descended into the scrotum.

Monosomus, **Monosomia** (*mon-o-so'-mus*, *mon-o-so'-me-ah*) [*μόνος*, single; *σῶμα*, body]. A double monster with a single body and two heads.

Monospasm (*mon'-o-spasm*) [*μόνος*, single; *σπασμός*, spasm]. Spasm affecting limited areas, as one side of the face, a single limb, or a single muscle or muscle-group. It is designated as Brachial, Crural, or Facial, according to the part affected, and as Central (Cerebral) or Peripheral, according to the seat of the causal lesion.

Monoxid (*mon-oks'-id*). See *Monobromid*.

Monro, **Foramen of**. See *Foramina*, *Table of*.

Mons (*mons*) [L.]. A mountain. **M. veneris**, the eminence in front of the symphysis pubis of the woman.

Monssel's Salt. Ferric subsulphate, a solution of which, **Monssel's Solution**, or **Liquor ferri subsulphatis** (U. S. P.), is used as a styptic.

Monster (*mon'-ster*) [*monstrum*]. An individual that by reason of congenital faulty development is incapable of properly performing the vital functions, or that owing to an excess or deficiency of parts differs in a marked degree from the normal type of the species.

Monstrosity (*mon-stros'-it-e*) [*monstrum*, a monster]. 1. The condition of a monster. 2. A monster.

TABLE OF MONSTROSITIES.*

ACCORDING TO GEOFFROY SAINT-HILAIRE, ALTERED BY HIRST AND PIERSON.

HEMITERATA.

I. ANOMALIES OF VOLUME.

A. OF STATURE.

1. *General Diminution*, as in a dwarf—delayed growth.
2. *General Increase*, as in a giant—precocious development.

B. OF VOLUME, strictly speaking.

1. *Local Diminution*. Affecting—
 - a. *Regions*, as a limb.
 - b. *Systems*, as undeveloped muscles.
 - c. *Organs*, as small breasts, stenosis of canals, etc.
2. *Local increase*, affecting—
 - a. *Regions*, as the head.
 - b. *Systems*, as the adipose tissue.
 - c. *Organs*, as large breasts in women, lactiferous breasts in men.

II. ANOMALIES OF FORM. *Single Order*, including—deformed heads; anomalies of shape in the stomach; deformed pelves, etc.

III. ANOMALIES OF COLOR.

- A. *Deficiency*, complete, partial, or imperfect, as in albinism.
- B. *Excess*, complete, partial, or imperfect, as in melanism.
- C. *Alteration*, as in unusual color of the iris.

IV. ANOMALIES OF STRUCTURE.

- A. *Deficiency in Consistency*, as cartilaginous conditions of bones.
- B. *Excess in Consistency*, as anomalous ossification.

V. ANOMALIES OF DISPOSITION.

A. BY DISPLACEMENT.

1. *Of the splanchnic organs*, as anomalous direction of heart or stomach, hernias, exstrophy of the bladder, etc.

* Reproduced, with the kind consent of the Publishers, from "Human Monstrosities," by Barton Cooke Hirst, M.D., and George A. Piersol, M.D. Philadelphia: Lea Brothers & Co., 1892.

2. *Of the nonsplanchnic organs*, as club-foot, curvature of the spine, misplaced teeth, misplaced blood-vessels, etc.
- B. BY CHANGE OF CONNECTION.
1. *Anomalous articulations*.
 2. *Anomalous implantations*, as teeth out of line.
 3. *Anomalous attachments*, as of muscles and ligaments.
 4. *Anomalous branches*, as of arteries and nerves.
 5. *Anomalous openings*, as of veins into the left auricle, of the ductus choledochus in an unusual situation, of the vagina into the rectum, of the rectum into the male urethra, of the rectum at the umbilicus, etc.
- C. IN CONTINUITY.
1. *Anomalous imperforations*, as of rectum, vulva, vagina, mouth, esophagus.
 2. *Anomalous union of organs*, as of kidneys, testicles, digits, teeth, ribs; adhesion of the tongue to the palate.
- D. BY CLOSURE, as in complete transverse septum in the vagina.
- E. BY DISJUNCTION.
1. *Anomalous perforations*, as persistence of foramen ovale, ductus arteriosus, urachus.
 2. *Anomalous divisions*, as splits, fissures in various organs, hare-lip, hypospadias, fissured tongue, cleft palate, fissured cheek.

VI. ANOMALIES OF NUMBER AND EXISTENCE.

1. *By numeric defect*, as absence of muscles, vertebrae, ribs, digits, teeth, a lung, a kidney, the uterus, the bladder, etc.
2. *By numeric excess*, as supernumerary digits, ribs, teeth, breasts, a double uterus.

HETEROTAXIS.

- I. SPLANCHNIC INVERSION.
- II. GENERAL INVERSION.

HERMAPHRODITES.*

- I. TRUE HERMAPHRODITES.
 - a. *Bilateral hermaphrodites*.
 - b. *Unilateral hermaphrodites*.
 - c. *Lateral hermaphrodites*.
- II. PSEUDOHERMAPHRODITES, with double sexual formation of the external genitals, but with unisexual development of the reproductive glands (ovaries and testicles).
 - a. *Male pseudohermaphrodites* (with testicles).
 1. *Internal pseudohermaphrodites*. Development of uterus masculinus.
 2. *External pseudohermaphrodites*. External genitals approach the female type; the monstrosity presents a feminine appearance and build.
 3. *Complete pseudohermaphrodites* (internal and external). Uterus masculinus with tubes; separate efferent canals for bladder and uterus.
 - b. *Female pseudohermaphrodites* (with ovaries). Persistence of male sexual parts.

* According to Klebs.

1. *Internal hermaphrodites*. Formation of vas deferens and tubes.
2. *External hermaphrodites*. Approach of the external genitals to the male type.
3. *Complete hermaphrodites* (internal and external). Masculine formation of the external genitals and of a part of the sexual tract.

MONSTERS.

CLASS I.—SINGLE MONSTERS.

ORDER I.—AUTOSITIC MONSTERS.

Genus I,	Species 1. <i>Ectromelus</i> , . .	Phocomelus Hemimelus Micromelus Ectromelus
	Species 2, . .	Symelus Uromelus Sirenomelus
Genus II,	Single species, <i>Celosoma</i> , .	Aspalasoma Agnosoma Cyllosoma Schistosoma Pleurisoma Celosoma
	Species 1. <i>Exencephalus</i> , .	Notencephalus Proencephalus Podencephalus Hyperencephalus Iniencephalus Exencephalus
Genus III,	Species 2. <i>Pseudencephalus</i> , .	Nosencephalus Thlipsencephalus Pseudencephalus
	Species 3. <i>Anencephalus</i> , .	Derencephalus Anencephalus
Genus IV,	Species 1. <i>Cyclocephalus</i> , .	Ethmocephalus Cecocephalus Rhinocephalus Cyclocephalus Stomiocephalus
	Species 2. <i>Otocephalus</i> , . .	Sphenocephalus Otocephalus Edocephalus Opococephalus Triocephalus

ORDER II.—OMPHALOSITIC MONSTERS.

Genus I,	Species 1. <i>Paracephalus</i> , .	Paracephalus Omacephalus Hemiacephalus
	Species 2. <i>Acephalus</i> , .	Acephalus Peracephalus Mylacephalus
	Species 3. <i>Asomata</i> , . . .	

Genus II, Single species, *Anideus*.

CLASS II.—COMPOSITE MONSTERS.

ORDER I.—DOUBLE AUTOSITIC MONSTERS.

- A. *Terata katadidyma*.
 Genus I, Diprosopus
 Genus II, Dicephalus
 Genus III, Ischiopagus
 Genus IV, Pygopagus
- B. *Terata anadidyma*.
 Genus I, Dipygus
 Genus II, Syncephalus
 Genus III, Craniopagus

C. *Terata anakatadidyma.*

Genus I, Prosopothoracopagus
Genus II, Omphalopagus
Genus III, Rachipagus

ORDER II.—DOUBLE PARASITIC MONSTERS.

Genus I,	Species 1. <i>Heterotyphus</i> , .	Heteropagus Heterodelphus Heterodymus Heterotyphus Heteromorphus
		Epicomus
	Species 2. <i>Heteralius</i> , . .	Epignathus Hypognathus Paragnathus Angnathus
Genus II,	Species 1. <i>Polygnathus</i> , .	Pygomelus Gastromelus Notomelus Cephalomelus Melomelus
		Dermocyma Endocyma
	Species 2. <i>Polytelus</i> , .	
Genus III,	Endocyma, .	

ORDER III.—TRIPLE MONSTERS.

Montgomery's Glands. See *Gland*.

Monthly Courses, M. Sickness, or Monthlies, the menses. **M. Nurse,** a nurse who attends after childbirth.

Monticulus (*mon-tik'-u-lus*) [L.]. A small elevation. **M. cerebelli,** the prominent central portion of the superior vermiciform process of the cerebellum.

Moore's Test. A test for glucose, consisting in the development of a black color when the solution is heated with potassium or sodium hydroxid.

Morbid (*mor'-bid*) [*morbus*, disease]. Pertaining to disease or diseased parts. **M. Anatomy,** the anatomy of organs or tissues in a state of disease.

Morbidity (*mor'-bid'-it-e*) [*morbus*, disease]. 1. The quality of disease or of being diseased. 2. The conditions inducing disease. 3. The ratio of the number of sick individuals to the total population of a place.

Morbific (*mor'-bif'-ik*) [*morbus*, disease; *facere*, to make]. Producing disease.

Morbilli (*mor-bil'-i*) [pl. : dim. of *morbus*, disease]. Measles.

Morbus (*mor'-bus*) [L.]. Disease. **M. Addisonii,** Addison's Disease. **M. anglicus,** rachitis. **M. basedowii.** See *Exophthalmic Goiter*. **M. brightii.** See *Bright's Disease*. **M. caducus,** epilepsy. **M. celsi,** catalepsy. **M. ceruleus,** congenital cyanosis. **M. coxarius,** coxalgia. **M. divinus,** epilepsy. **M. gallicus,** syphilis. **M. maculosus werlhofii,** purpura hemorrhagica. **M. magnus,** epilepsy. **M. medicorum,** the mania of those who seek the advice of physicians for imaginary diseases. **M. regius,** jaundice. **M. sacer,** epilepsy.

Morcellation (*mor-sel-a'-shun*) [Fr. *morceler*, to cut up or parcel out]. The art of reduc-

ing to fragments, as *e. g.*, the fetus in embryotomy.

Mordant (*mor'-dant*) [*mordere*, to bite]. A substance, such as alum, phenol anilin-oil, that fixes the dyes used in coloring textures or in staining tissues and bacteria.

Morgagni (*mor-gan'-ye*). An Italian anatomist of the eighteenth century. **M., Cataract** of. See *Cataract*. **Hydatid** of. See *Hydatid*.

Morgue (*morg*) [Fr.]. A place where unknown dead are exposed for identification.

Moria (*mo'-re-ah*) [*μωρία*, folly]. A form of dementia characterized by talkativeness and silliness.

Moribund (*mor'-ib-und*) [*moribundus*, from *moriri*, to die]. In a dying condition.

Morioplasty (*mor'-e-o-plas-te*) [*μόριον*, a small piece; *πλασσειν*, to mold]. Plastic surgery.

Morning-sickness. The nausea of pregnant women, occurring chiefly in the early months of gestation; also, the imitative or sympathetic nausea sometimes experienced by the husband during the wife's early pregnancy.

Morphea, Morphœa (*mor-fe'-ah*) [*μορφή*, a blotch]. A disease of the skin characterized by the presence of rounded or oval, pinkish, or ivory-white patches, due to an excess of fibrous tissue, with atrophy of the skin-structures proper. Morphea is believed to be a trophoneurosis, and is considered a circumscribed form of scleroderma. It is also termed Addison's keloid, or circumscribed scleroderma. **M., Acroteric,** the form in which the beginning and the greatest intensity of the disease are at the extremities. **M., Herpetiform,** that in which the lesions follow those of herpes in their distribution.

Morphia (*mor'-fe-ah*). See *Morphin*.

Morphin, Morphina (*mor'-fin, mor'-fin-nah*) [*Morpheus*, god of sleep], $C_{17}H_{19}NO_3 + H_2O$. A colorless or white crystalline alkaloid obtained from opium, to which the chief effects of the latter are due. It differs from opium in being less stimulant, less constipating, and less likely to produce disagreeable after-effects. On account of its insolubility in water, morphin is used principally in the form of its salts. The dose of the salts of morphin is gr. $\frac{1}{4}$ – $\frac{1}{2}$ (0.008–0.032). **M. acetate,** morphine acetate, $C_{17}H_{19}NO_3 \cdot C_2H_3O_2 + 3H_2O$. From it are prepared *Liquor morphine acetatis* (B. P.) (gr. ss to the fluidram), dose \mathfrak{mxx} –xl (1.3–2.6), and *Injectio morphine hypodermica* (gr. j in 10 minims). **M. hydrochlorate,** morphine hydrochlorate, $C_{17}H_{19}NO_3 \cdot HCl + 3H_2O$. From it are prepared *Liquor morphine hydrochloratis* (B. P.) (gr. ss to the fluidram), dose \mathfrak{mxx} –xxx (1.0–2.0); *Suppositoria morphine* (B. P.)

(gr. ss each); *Tinctura chloroformi et morphinæ* (gr. $\frac{1}{4}$ to the fluidram); *Trochisci morphinæ* (B. P.) (gr. $\frac{1}{2}$), and *Trochisci morphinæ et ipecacuanhæ* (B. P.) (gr. $\frac{3}{8}$). **M. phthalate** (unofficial) is employed hypodermically. **M. sulphate**, morphinæ sulphas ($C_{17}H_{19}NO_3 \cdot H_2SO_4 + 5H_2O$). From it are prepared: *Liquor morphinæ sulphatis* (B. P.), dose $m_{xx}-xl$ (0.65-4.0); *Pulvis morphinæ compositus*, Tully's powder, dose gr. x (0.65), containing gr. $\frac{1}{6}$ (0.01) of morphin sulphate; *Trochisci morphinæ compositi* (U. S. P.), each containing gr. $\frac{1}{16}$ (0.0016), and *Liquor morphinæ hypodermicus* (N. F.), Magendie's solution, containing 16 grains to the fluidounce. **M. tartrate** (unofficial) is employed for hypodermic use.

Morphinism (*mor'-fin-izm*) [*morphin*]. 1. The condition caused by the habitual use of morphin. 2. The morphin-habit.

Morphinomania, **Morphiomania** (*mor'-fin-o-ma'-ne-ah*, *mor'-fe-o-ma'-ne-ah*) [*morphin*; *mania*, madness]. 1. A morbid craving for morphin. 2. Insanity due to the morphin-habit.

Morphography (*mor'-fog'-ra-fe*) [*μωρφηγία*, form; *γραφειν*, to write]. See *Morphology*.

Morphologic (*mor'-fo-loj'-ik*) [*μωρφηγία*, form; *λογος*, science]. Pertaining to morphology.

Morphology (*mor'-fol'-o-je*) [*μωρφηγία*, form; *λογος*, science]. The science that treats of the form and structure of organized beings.

Morphometry (*mor'-fom'-et-re*) [*μωρφηγία*, form; *μετρον*, measure]. The measurement of the forms of organisms.

Morphon (*mor'-fon*) [*μωρφηγία*, form]. An individual element of an organism, characterized by a definite form, as a cell or a segment of a vertebrate.

Morphosis (*mor'-fo'-sis*) [*μωρφηγία*, form]. The act, mode, or order of formation of an organism.

Morphotic (*mor'-fet'-ik*) [*μωρφηγεω*, to form]. Pertaining to morphosis; entering into the formation of the framework of an organism.

M. Proteids, those that enter into the structure of the tissues.

Morpio (*mor'-pe-o*) [L.]. The crab-louse.

Morrhua (*mor'-u-ah*) [L.]. The cod. *Oleum morrhuæ*, cod-liver oil, a fixed oil obtained from the fresh livers of *Gadus morrhua*, or of other species of *Gadus*. Three varieties of oil are known in commerce—a white or pale-yellow, a brownish-yellow, and a dark-brown. The oil contains gaduin ($C_{25}H_{48}O_9$), oleic, palmitic, stearic, myristic, and physsetolic acids, glycerol, butyric and acetic acids, biliary pigments, iodine, and bromine. A crystalline substance, morrhinol, containing phosphorus, iodine, and bromine, has also been isolated, as well as several leukomains and the fixed bases, asellin ($C_{28}H_{52}N_4$) and mor-

ruin ($C_{19}H_{27}N_3$). Cod-liver oil is used in pulmonary and other forms of tuberculosis, and in wasting conditions due to other causes.

Dose $f\overline{3}j-f\overline{3}ss$ (4.0-16.0).

Morrhuin (*mor'-u-in*). See *Morrhua*.

Morrhual (*mor'-u-ol*). See *Morrhua*.

Mors (*morz*) [L.]. Death.

Morsus (*mor'-sus*) [L.]. A bite. **M. diaboli**, the fimbriated extremity of the oviduct.

Mortal (*mor'-tal*) [*mortalis*, from *mors*, death]. Liable to death or dissolution; terminating in death; causing death; deadly.

Mortality (*mor'-tal'-it-e*) [*mors*, death]. 1. The quality of being mortal. 2. The death-rate.

Mortar (*mor'-tar*) [*mortarium*, an urn]. An urn-shaped vessel of porcelain, iron, or glass, for pulverizing substances by means of a pestle.

Mortification (*mor'-tif-ik-a'-shun*). See *Gangrene*.

Morton's Foot [T. G. Morton, an American surgeon]. A painful affection of the metatarsophalangeal joint of the fourth toe. See *Diseases, Table of*.

Mortuary (*mor'-tu-a-re*) [*mortuarium*, a tomb; from *mors*, death]. 1. A house for temporary burial. 2. Relating to death or burial.

Morula (*mor'-u-lah*) [dim. of *morum*, a mulberry]. The solid mass of cells resulting from the complete segmentation of the vitellus of an ovum.

Morulation (*mor'-u-la'-shun*) [*morula*, a little mulberry]. The formation of the morula during the process of the segmentation of the egg.

Morus (*mo'-rus*) [L.]. See *Mulberry*.

Morvan's Disease. See *Diseases, Table of*.

Moschus (*mos'-kus*) [*μωσχος*, musk]. See *Musk*.

Mosquito (*mus-ke'-to*) [Sp., a little gnat]. An insect, the *Culex* mosquito, the sting of which causes the formation of a wheal that itches intensely.

Moss [AS., *mebs*, moss]. 1. A small cryptogamic plant of the natural order Musci.

M., Ceylon. See *Agar-agar*. **M., Club**. See *Lycopodium*. **M., Corsican**. See *Corsican Moss*. **M., Iceland**. See *Chondrus*.

M., Irish. See *Carragheen*.

Moth [AS., *modde*, moth]. Chloasma.

Mother (*mu-th'-er*) [AS., *mōder*, mother]. 1. A female parent. 2. The source of anything. **M.-cell**, a cell from which other cells are formed, especially one the nucleus of which is undergoing karyokinetic changes preparatory to dividing into daughter-cells. **M.-liquor**, the liquid remaining after distillation of substances have separated by crystallization. **M.'s Mark**, a birth-mark. See *Nevus*.

Mother (*mutʰ-er*) [allied to AS., *mud*]. A slimy film formed on the surface of fermenting liquid, as, *e. g.*, on vinegar.

Motile (*moʹ-til*) [*movere*, to move]. Able to move; capable of spontaneous motion, as a motile flagellum.

Motility (*mo-tilʹ-it-e*) [*movere*, to move]. Ability to move spontaneously.

Motion (*moʹ-shun*) [*movere*, to move]. 1. The act of changing place. 2. An evacuation of the bowels; the matter evacuated.

Motor (*moʹ-tor*) [*movere*, to move]. 1. Moving or causing motion. 2. Concerned in or pertaining to motion, as M. cell, M. center, M. nerve. **M. Aphasia.** See *Aphasia*. **M. Area**, the portion of the cerebral hemisphere presiding over voluntary motion, including the precentral and postcentral gyri, the posterior part of the three frontal gyri and the paracentral lobule on the median surface of the hemisphere. **M. oculi**, the third cranial or oculomotor nerve supplying all the muscles of the eye, except the superior oblique and external rectus. **M. Points**, the points on the surface of the body where the various branches of the motor nerves supplying the muscles may be stimulated by electricity.

Motorial (*mo-toʹ-re-al*) [*movere*, to move]. Of or pertaining to motion. **M. End-plate**, an eminence of protoplasm within the sarcolemma of a muscular fiber, representing the termination of the motor nerve-fiber.

Motorium (*mo-toʹ-re-um*) [*movere*, to move]. 1. A motor center. 2. The motor apparatus of the body, both nervous and muscular, considered as a unit.

Mottling (*motʹ-ling*) [OF., *matellé*, clotted, curdled]. A spotted condition.

Mould (*mold*). See *Mold*.

Mountain (*maunʹ-ten*) [*mons*, mountain]. A high hill. **M. Anemia**, ankylostomiasis. **M. Fever**, **M. Sickness**. 1. A condition characterized by dyspnea, rapid pulse, headache, nausea, and vomiting, depending upon the rarefied state of the air at high altitudes. This is properly called mountain-sickness. 2. A form of typhoid fever occurring in mountainous districts.

Mounting (*maunʹ-ting*) [*mons*, mountain]. The act of arranging objects, especially anatomic specimens, on a suitable support and in a proper medium for ready examination. For macroscopic specimens the medium is usually alcohol, for microscopic specimens, Canada balsam or glycerol.

Mouth (*moʊth*) [AS., *mūð*, mouth]. The commencement of the alimentary canal, the cavity in which mastication takes place. In a restricted sense, the aperture between the lips. 2. The entrance to any cavity or canal. **M.-breather**, a person who habitually breathes through the mouth. **M.-breathing**,

respiration through the mouth instead of, as normally, through the nose.

Movement (*moonʹ-ment*) [*movere*, to move].

The act of moving. **M., Ameboid**, a movement produced in certain cells, as the white corpuscles, by the protrusion of processes of the protoplasm into which the whole cell then seems to flow; so-called from the resemblance of the movement to that of the ameba.

M., Associated, an involuntary movement in one part when another is moved voluntarily. **M., Brownian**, a physical phenomenon, a form of communicated motion observed in aggregations of minute particles, and consisting of a rapid, oscillating movement without change of the relative position of the moving particles. **M., Ciliary**, a lashing movement produced by delicate hair-like processes termed cilia, as on the epithelium of the respiratory tract and in certain microorganisms. **M., Circus-**, rapid circular movements or somersaults, produced by injury of the corpus striatum, of the optic thalamus, or of the crus cerebri of one side.

M., Communicated, that produced by a force acting from without; opposed to spontaneous movement. **M., Petal**, the movements of the fetus in utero. **M., Forced**, movement of the body from injury of the motor centers or the conducting paths, as, *e. g.*, **index M.**, when the cephalic part of the body is moved about the stationary caudal part; **rolling M.**, when the animal rolls on its long axis. **M., Molecular**. Synonym of *M., Brownian*.

Moxa (*moksʹ-ah*) [from the Japanese]. A combustible material which is applied to the skin and ignited for the purpose of producing an eschar. It is prepared from several species of *Artemisia*; artificial moxa is made from cotton saturated with niter. **M.-bearer**, or **Porte-moxa**, an instrument for applying the moxa.

Mucedin (*muʹ-se-din*) [*mucus*, mucus]. A nitrogenous substance obtained from gluten.

Mucic Acid (*muʹ-sik*), $C_6H_{10}O_6$. A crystalline dibasic acid produced by the oxidation of gums and certain sugars.

Mucigen (*muʹ-sij-en*) [*mucus*, mucus; *γεννᾶν*, to produce]. A substance producing mucin; it is contained in epithelial cells that form mucus.

Mucigenous (*muʹ-sij-en-us*) [*mucus*, mucus; *γεννᾶν*, to produce]. Producing mucus.

Mucilage (*muʹ-sil-ij*) [*mucilago*, moldy moisture, from *mucus*]. In pharmacy, a solution of a gum in water. Mucilages (mucilagines) are employed as applications to irritated surfaces, particularly mucous membranes, as excipients for pills, and to suspend insoluble substances. The following are employed: *Mucilago acaciæ*, *M. amyli* (B. P.), *M. sassafras medullæ*, *M. tragacanthæ*, *M. ulni*.

Mucilaginous (*mu-sil-aj'-in-us*) [*mucilago*, from *mucus*, mucus]. Pertaining to or of the nature of mucilage.

Mucilago (*mu-sil-aj'-go*). See *Mucilage*.

Mucin (*mu'-sin*) [*mucus*, mucus]. An albuminoid substance, the characteristic constituent of mucus. It is supposed to be produced by the union of an albuminous body and a colloid carbohydrate, the animal gum of Landwehr. Mucin occurs in saliva, bile, secretions of mucous membranes, synovia, in mucous tissue, in certain cysts, etc. It is insoluble in water, and is precipitated by alcohol and acetic acid.

Mucinogen (*mu-sin'-o-jen*) [*mucus*, mucus; γεννᾶν, to produce]. The antecedent principle from which mucin is derived.

Mucinoid (*mu'-sin'-oid*) [*mucus*, mucus; εἶδος, like]. Resembling mucin.

Mucinuria (*mu-sin-u'-re-ah*) [*mucus*, mucus; *urina*, urine]. The presence of mucin in the urine.

Muciparous (*mu-sip'-ar-us*) [*mucus*, mucus; *parere*, to bring forth]. Secreting or producing mucus.

Mucocele (*mu'-ko-sel*) [*mucus*, mucus; κήλη, tumor]. A tumor formed from the distention of the lacrimal sac by mucus.

Mucocutaneous (*mu-ko-ku-ta'-ne-us*) [*mucus*, mucus; *cutis*, skin]. Pertaining to a mucous membrane and the skin; pertaining to the lines where these join.

Mucoid (*mu'-koid*) [*mucus*, mucus; εἶδος, like]. Resembling mucus.

Mucopurulent (*mu-ko-pu'-ru-lent*) [*mucus*; *pus*, pus]. Containing mucus mingled with pus.

Mucopus (*mu'-ko-pus*) [*mucus*, mucus; *pus*, pus]. A mixture of mucus and pus.

Mucor (*mu'-kor*) [*muere*, to be moldy]. A genus of hyphomycetes. **M. corymbifer**, one found in the cerumen of the external auditory meatus. **M. mucedo**, a species found on fecal matter and nitrogenous organic substances.

Mucosa (*mu-ko'-sah*) [*mucosus*, mucous—membrana understood]. A mucous membrane.

Mucosin (*mu'-ko-sin*) [*mucus*, mucus]. The form of mucus to which the nasal, uterine, and bronchial mucus owe their viscosity.

Mucous (*mu'-kus*) [*mucus*, mucus]. Containing or having the nature of mucus; secreting mucus, as *M. membrane*; depending on the presence of mucus, as *M. rales*. **M. Casts**, a term given to the casts found in the feces in cases of membranous enteritis. **M. Catarrh**, catarrhal inflammation of a mucous membrane. **M. Disease**, enterocolitis, especially of children. **M. Patch**, a flattened grayish-white exudate, occurring in secondary syphilis on mucous membranes and at mucocutaneous junctions. **M. Polyp**, a soft,

gelatinous outgrowth from a mucous membrane; it may be a true myxoma, but usually is a hyperplasia due to chronic inflammation. **M. Tissue**, a form of connective tissue in which the intercellular substance is of a soft, gelatinous character and contains mucin. The cells from pressure assume a stellate or spindle shape.

Mucuna (*mu-ku'-nah*) [Braz.]. A genus of leguminous herbs. **M. pruriens**, cowhage, the hairs of the pods of which were formerly used as a vermifuge and counterirritant.

Mucus (*mu'-kus*) [L.]. The viscid liquid secreted by mucous membranes. It consists of water, mucin, and inorganic salts, together with epithelial cells, leukocytes, etc., held in suspension.

Mud-bath. See *Bath*.

Muguet (*mu-gwa'*) [Fr.]. Thrush.

Mulberry (*mul'-ber-ee*) [*morus*, mulberry-tree; *berry*, from AS., *berige*]. A tree of the natural order Urticarie. **Morus nigra** is the source of *Mori succus* of the B. P., the latter being used to make *Syrupus mori* (B. P.). Both are employed as drinks in fevers and as additions to gargles in pharyngitis. The fruit of *Morus alba* is used as food for silk-worms. **Mulberry Calculus**. See *Calculus*. **Mulberry Mass**. See *Morula*.

Mulder's Test. A test for glucose, consisting in the addition of indigo (sodium sulph-indigotate) to the alkalized solution and warming. The blue color changes to violet-red, then to yellow or white, if glucose is present.

Mullein (*mul'-en*). See *Verbascum*.

Müller's Fibers. The supporting fibers of the retina running transversely to its layers.

Müller's Fluid. A fluid used for hardening tissues. Its composition is as follows: Potassium dichromate, 2 to 2.5 parts; sodium sulphate, 1 part; water, 100 parts.

Müller's Muscle. 1. Small bundles of non-striped muscle in the upper and lower lids, assisting in elevating the former and depressing the latter. 2. The circular fibers of the ciliary muscle. 3. The orbital muscle. See *Muscle*, *Müller's*, in *Muscles*, *Table of*.

Müller's Ring. A constriction in the cervix uteri below the true os internum.

Müllerian Duct, *Müller's Duct*. See *Duct of Müller*.

Multi- (*mul'-te*) [pl. of *multus*, much]. A prefix signifying many.

Multicellular (*mul'-te-sel'-u-lar*) [*multus*, many; *cellula*, cell]. Many-celled.

Multigravida (*mul'-te-grav'-id-ah*) [*multus*, many; *gravidus*, pregnant]. A pregnant woman who has passed through one or more pregnancies.

Multilobular (*mul'-te-lob'-u-lar*) [*multus*, many; *lobus*, a lobe]. Many-lobed.

Multilocular (*mul-te-lok'-u-lar*) [*multus*, many; *loculus*, a locule or cell]. Many-celled; polycystic.

Multinuclear (*mul-te-nu'-kle-ar*) [*multus*, many; *nucleus*, kernel]. Having several or many nuclei.

Multipara (*mul-tip'-ar-ah*) [*multus*, many; *parere*, to bring forth]. 1. A pregnant woman who has already borne one or more children. It has been customary to designate the number of the pregnancy of a multipara by the unpronounceable terms II-para, III-para, IV-para, etc. More commendable would be the following terms: Secundipara, Tertipara, Quartipara, Quintipara, Sextipara, Septimipara, Octavipara, Nonipara, Decimipara, etc., to designate respectively a woman in her second, third, fourth, etc., pregnancy. 2. A woman bearing several offspring at a birth.

Multiparous (*mul-tip'-ar-us*) [*multus*, many; *parere*, to bring forth]. Having borne several children.

Multiple (*mul'-tip-l*) [*multus*, many; *pliare*, to fold]. Manifold; affecting many parts at the same time. **M. Neuritis.** See *Neuritis*. **M. Sclerosis.** See *Sclerosis*.

Multipolar (*mul-te-pol'-lar*) [*multus*, many; *polus*, a pole]. Having more than one pole; as *M. nerve-cells*, those having more than one process.

Multivalent (*mul-tiv'-al-ent*) [*multus*, many; *valere*, to be worth]. In chemistry, combining with more than two atoms of a univalent element.

Mummification (*mum-if-ik-a'-shun*) [*mummy*, from Pers., *mum*, wax; *facere*, to make]. The change of a part into a hard, dry mass; dry gangrene.

Mumps [*Du.*, *mompen*, to mumble]. An acute infectious disease characterized by swelling of the parotid and at times of the other salivary glands. After a period of incubation of from two to three weeks, the disease begins with fever and pain below the ear; soon a tense, painful swelling forms in the region of the parotid gland, rendering mastication and deglutition difficult and painful. In the course of a week the swelling subsides without suppuration. The most frequent complication is orchitis; in rare cases the ovaries are affected.

Mural (*mu'-ral*) [*murus*, wall]. Pertaining to a wall, as a *M. fibroid*. **M. Gestation.** **M. Pregnancy**, pregnancy in the uterine extremity of a Fallopian tube.

Murexid (*mu-rek'-id*) [*murex*, the purple-fish], $C_{12}H_8N_4O_6 + H_2O$. Ammonium purpurate, a dichroic crystalline salt obtained from guano and used as a dye. It is also formed when a solution of uric acid or urates is evaporated with nitric acid and ammonia

added to the residue. This is the **Murexid Test**.

Muriate (*mu'-re-at*) [*muria*, brine]. An old name for a chlorid.

Muriated (*mu'-re-a-ted*) [*muria*, brine]. Containing chlorin or a chlorid.

Muriatic (*mu-re-at'-ik*) [*muria*, brine]. Pertaining to brine. **M. Acid.** See *Acid, Hydrochloric*.

Murmur (*mer'-mer*) [*L.*, a murmur]. A blowing or rasping sound heard on auscultation. **M., Accidental**, a murmur dependent on an accidental circumstance, as *e.g.*, on compression of an artery by the stethoscope. **M., Anemic.** See *M., Hemic*.

M., Aneurysmal, or aneurysmal bruit, the murmur or bruit heard over an aneurysm. See *Bruit*. **M., Arterial**, the sound made by the arterial current. **M., Blood-.** See *M., Hemic*. **M., Cardiac**, any adventitious sound heard over the region of the heart.

In relation to their seat of generation, cardiac murmurs are designated as mitral, aortic, tricuspid, and pulmonary; according to the period of the heart's cycle at which they occur they are divided into systolic, those occurring during the systole; diastolic, those occurring in diastole; presystolic and prediastolic, those occurring just before systole and diastole respectively. **M., Cardiopulmonary**, one produced by the impact of the heart against the lung. **M., Direct**, a murmur produced by obstruction to the blood-current as it is passing in its normal direction. **M., Duroziez's**, the double murmur sometimes heard in the femoral artery in aortic regurgitation.

M., Dynamic, one resulting from tumultuous and irregular action of the heart. **M., Endocardial**, a murmur produced within the cavities of the heart. **M., Exocardial**, a murmur connected with the heart, but produced outside of its cavities. **M., Flint's**, a murmur sometimes heard at the apex of the heart in aortic regurgitation. It is generally presystolic in time and is probably due to the fact that on account of the extreme ventricular dilatation the valves cannot be forced back against the walls and produce a relative narrowing of the auriculoventricular orifice.

M., Friction-, a sound produced by the rubbing of two inflamed serous surfaces upon each other. **M., Functional**, a cardiac murmur occurring from excited action of the heart or anemic condition of the individual, without any structural change in the valves or orifices. **M., Hemic**, a sound believed to be due to changes in the quality or amount of the blood, and not to lesions of the vessels or valves. It is heard especially in anemic conditions.

M., Indirect, one produced by the blood flowing in a direction contrary to the normal current. **M., Inorganic**, a murmur not due

to valvular lesions; a hemic or a functional murmur. *M.*, Musical, a cardiac murmur having a musical quality. *M.*, Organic, a murmur due to structural changes in the heart.

muscaria. It causes depression of the action of the heart and the respiration, increases the excretions of the salivary and lacrimal glands and of the intestines, and produces contraction of the pupil.

TABLE OF ENDOCARDIAL MURMURS.

TIME.	POINT OF MAXIMUM INTENSITY.	LINE OF CONDUCTION.	LESION.	QUALITY.
Systolic.	Center of mitral area, above and to left of apex.	At sixth rib opposite apex, a line drawn from the anterior fold of axilla to lower angle of left scapula.	Mitral insufficiency or incompetence.	Variable; usually soft, blowing; may be distinctly musical.
Systolic.	Midsternum or to right of it, opposite third rib or second interspace.	Toward top of sternum, and along aorta and its large branches.	Aortic obstruction.	Usually loud and harsh. Harshness is one of its distinguishing characteristics.
Diastolic.	Midsternum opposite upper border of cartilage of third rib.	Down sternum to ensiform cartilage.	Aortic insufficiency or incompetence.	Soft, blowing, sometimes rough, frequently musical. It has the greatest area of diffusion of all the cardiac murmurs.
Presystolic.	Over mitral area around the apex.	Usually not transmitted.	Mitral obstruction.	Generally low-pitched, rough, churning, grinding, or bubbling. Subject to great variation of pitch and quality.
Systolic.	Midsternum just above the ensiform cartilage.	Toward the epigastrium.	Tricuspid insufficiency or incompetence.	Low-pitched, superficial, blowing, soft, faint.
Presystolic.	Midsternum opposite the cartilage of fourth rib.	Not transmitted.	Tricuspid obstruction.	Undetermined.
Systolic.	Second interspace to the left of sternum or at the level of third rib.	Upward a short distance and to left of sternum, stopping abruptly.	Pulmonary obstruction.	Often harsh and audible over the whole precordia; may be very faint.
Diastolic.	Second left interspace.	Down left edge of sternum to ensiform cartilage.	Pulmonary insufficiency or incompetence.	Soft and blowing.

Murphy's Button. A mechanic device for bringing together the visceral surfaces of the intestines in intestinal anastomosis.

Murr, Murrain (*mur*, *mur'-an*) [Fr., from Lat., *mori*, to die]. 1. Any fatal disease of cattle and sheep. 2. Foot-and-mouth disease.

Muscæ (*mus'-ke*) [L.]. Plural of *Musca*, a fly. *M. hispaniolæ*, cantharides. *M. volitantes*, floating specks in the field of vision due to opacities in the media of the eye.

Muscardine (*mus'-kahr-din*) [Fr., a peculiar fungus]. A disease of silk-worms caused by the fungus *Botrytis bassiana*.

Muscarin (*mus'-kar-in*), $C_8H_{12}NO_2 \cdot H_2O$. A poisonous alkaloid obtained from *Amanita*

Muscle (*mus'-l*) [*musculus*, a muscle]. A structure composed chiefly of muscular tissue, and having the property of contracting. Muscles are of two kinds, the striped, or striated, and the smooth, or unstriated, the first being voluntary, the second involuntary, *i. e.*, not subject to the will. *M.-casket*, a muscle-compartment. *M.-column*. See *Muscular Tissue*. *M.-compartment*, one of the divisions of a muscle-fiber produced by the extension of Krause's membrane from the sarcolemma. *M.-fiber*, the ultimate element of which muscular tissue is made up. Voluntary muscles consist of transversely striated fibers, involuntary muscles of spindle-

shaped fibers or cells. *M.*, *Involuntary*, one not under the control of the will, as the non-striated muscles. *M.*, *Nonstriated*, one composed of spindle-shaped muscle-fibers. See *Muscular Tissue*. *M.*, *Papillary*, the muscular eminences in the ventricles of the heart, from which the cordæ tendineæ arise. *M.-plasma*, the fluid portion of the muscle-tissue. It is neutral or alkaline and spontaneously coagulable, and contains myosinogen (the coagulable substance), paramyo-

sinogen, myoglobulin, myoalbumoses, and albumin. *M.-rod*, one of the ultimate divisions of the dim band of a muscle-compartment. *M.*, *Skeletal*, any one of the muscles attached to and acting on the skeleton. *M.*, *Striated*, *M.*, *Striped*, a muscle constituted of striped muscle-fibers. See *Muscular Tissue*. *M.-sugar*. See *Inosite*. *M.*, *Unstriated*, *M.*, *Unstriped*. See *M.*, *Nonstriated*. *M.*, *Voluntary*. See *M.*, *Striped*.

TABLE OF MUSCLES (ARRANGED ALPHABETICALLY).

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
Abductor hallucis.	Outer head, os calcis, plantar fascia, inter-muscular septum; inner head, internal annular ligament, and tendon tibialis posterior.	Inner portion of lower surface of base of great toe and inner side of internal sesamoid bone.	Internal plantar division of posterior tibial nerve.	Flexes and abducts first phalanx of great toe.
Abductor longus pollicis.	See <i>Extensor ossis</i>	<i>metacarpi pollicis</i> .		
Abductor minimi digiti.	Pisiform bone.	First phalanx of the little finger.	Ulnar.	Abducts little finger.
Abductor minimi digiti.	Outer tuberosity of the os calcis and plantar fascia.	First phalanx of the little toe.	External plantar.	Abducts little toe.
Abductor pollicis.	Trapezium, scaphoid, annular ligament, palmar fascia.	First phalanx of thumb.	Median.	Abducts and flexes first phalanx of thumb.
Abductor hallucis.	Inner tuberosity of the os calcis.	First phalanx of great toe.	Internal plantar.	Abducts great toe.
Accelerator urinæ.	Central tendon of perineum and median raphe.	Bulb, spongy and cavernous parts of the penis.	Perineal.	Ejects urine.
Adductor brevis.	Ramus of pubes.	Upper part of the linea aspera of femur.	Obturator.	Adducts, rotates externally, and flexes thigh.
Adductor hallucis.	Tarsal ends of the three middle metatarsal bones.	Base of the first phalanx of great toe.	External plantar.	Adducts great toe.
Adductor longus.	Front of pubes.	Middle of linea aspera of femur.	Obturator.	Adducts, rotates outward, and flexes thigh.
Adductor magnus.	Rami of pubes and ischium.	Along the linea aspera of femur.	Obturator and great sciatic.	Adducts thigh and rotates it outward.
Adductor minimus.	A name given to the upper portion of the adductor magnus.			
Adductor pollicis.	Third metacarpal.	First phalanx of thumb.	Ulnar.	Draws thumb to median line.
Anconeus.	Back of external condyle of humerus.	Olecranon process and shaft of ulna.	Musculospiral.	Extends forearm.
Arrectores pili.	Pars papillaris of the skin.	Hair-follicles.	Sympathetic.	Elevate the hairs of the skin.

TABLE OF MUSCLES.—*Continued.*

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
Arytenoepiglottideus inferior.	Arytenoid (anteriorly).	Epiglottis.	Recurrent laryngeal.	Compresses sacculus of larynx.
Arytenoepiglottideus superior.	Apex of arytenoid cartilage.	Arytenoepiglottidean folds.	Recurrent laryngeal.	Constricts aperture of larynx.
Arytenoidens.	Posterior and outer border of one arytenoid.	Back of other arytenoid.	Superior and recurrent laryngeal.	Closes back part of glottis.
Attollens aurem.	Occipitofrontalis aponeurosis.	Pinna.	Temporal branch of facial.	Elevates pinna.
Attrahens aurem.	Lateral cranial aponeurosis.	Helix.	Facial.	Advances pinna.
Asygos uvulae.	Posterior nasal spine of palate bone.	Uvula.	Facial through sphenopalatine ganglion.	Raises uvula.
Biceps.	1. Long head—glenoid cavity. 2. Short head—coracoid process.	Tuberosity of radius.	Musculocutaneous.	Flexes and supinates forearm.
Biceps.	Ichial tuberosity and linea aspera.	Head of fibula and outer tuberosity of head of fibula.	Great sciatic and external popliteal.	Flexes and rotates leg outward.
Biventer cervicis.	Transverse processes, 2-4 upper dorsal.	Superior curved line of occipital bone.	It is a portion of the complexus.	Retracts and rotates head.
Bowman's.	See <i>Ciliary</i> .			
Brachialis anticus.	Lower half of shaft of humerus.	Coronoid process of ulna.	Musculocutaneous, musculospiral.	Flexes forearm.
Brücke's.	See <i>Ciliary</i> .			
Buccinator.	Alveolar process of maxillary bones and pterygomaxillary ligament.	Orbicularis oris.	Buccal branch of facial.	Compresses cheeks, retracts angle of mouth.
Bulbocavernosus.	See <i>Accelerator urinae</i> .			
Cephalopharyngeus.	See <i>Constrictor of pharynx, superior</i> .			
Cervicalis ascendens.	Angles of 5 upper ribs.	Transverse processes of 4th, 5th, and 6th cervical vertebrae.	Branches of cervical.	Keeps head erect.
Ciliary.	1. <i>Longitudinal</i> portion (Brücke's m.): junction of cornea and sclera; 2. <i>Circular</i> portion (Müller's m.): the fibers form a circle.	1. Outer layers of choroid. 2. Ciliary processes.	Ciliary.	The muscle of visual accommodation.
Coccygeus.	Ischial spine.	Coccyx, sacrum, and sacrococcygeal ligament.	Sacral.	Supports coccyx and closes pelvic outlet.
Complexus.	Transverse processes 7th cervical and 6 upper dorsal, and articular processes of 3d to 6th cervical vertebrae.	Occipital bone.	Suboccipital, great occipital, and branches of cervical.	Retracts and rotates head.
Compressor narium.	Nasal aponeurosis.	Fellow muscle and canine fossa.	Facial.	Compresses nostril.

TABLE OF MUSCLES.—*Continued.*

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
Compressor na- rii minor.	Alar cartilage.	Skin at end of nose.	Facial.	Dilates nostril.
Compressor sac- culi laryngis.	Fibers of the aryteno- epiglottideus.		Recurrent laryn- geal.	Compressor of sac- cule of larynx.
Compressor ure- thræ.	Ramus of pubes.	Fellow muscle.	Perineal.	Compresses mem- branous urethra.
Compressor vagi- næ.	The analogue of the two bulbocavernosi of the male.		Perineal.	Compresses vagi- na.
Compressor venæ dorsalis penis.	Fibers of the ischio- cavernosus.	Fascial sheath of penis, over dor- sal vessels.		Compressor of dor- sal vein.
Constrictor of pha- ryn timer (inferior).	Cricoid and thyroid cartilages.	Pharyngeal raphe.	Glossopharyngeal, pharyngeal plex- us, and external laryngeal.	Contracts caliber of pharynx.
Constrictor of pha- ryn timer (middle).	Cornua of hyoid bone and stylohyoid liga- ment.	Pharyngeal raphe.	Glossopharyngeal, and pharyngeal plexus.	Contracts caliber of pharynx.
Constrictor of pha- ryn timer (superior).	Internal pterygoid plate, pterygomaxil- lary ligament, jaw, and side of tongue.	Pharyngeal raphe.	Glossopharyngeal, and pharyngeal plexus.	Contracts caliber of pharynx.
Coracobrachialis.	Coracoid process of scapula.	Inner surface of shaft of humerus.	Musculocutane- ous.	Adducts and flexes arm.
Corrugator cutis ani.	Submucous tissue on interior of anus.	Subcutaneous tis- sue on opposite side of anus.	Sympathetic.	Corrugates skin about anus.
Corrugator super- cilii.	Superciliary ridge of frontal bone.	Orbicularis palpe- brarum.	Facial.	Draws eyebrow downward and inward.
Crampton's.	See <i>Ciliary</i> .			
Cre master.	Upper and deep sur- face of middle of Pou- part's ligament.	Spine and crest of pubic bone and fascia propria.	Genital branch of genitocrural.	Elevates testicle.
Cricoar ytenoideus lateralis.	Side of cricoid carti- lage.	Angle and external surface of aryte- noid.	Recurrent laryn- geal.	Closes glottis.
Cricoar ytenoideus posticus.	Back of cricoid carti- lage.	Base of arytenoid cartilage.	Recurrent laryn- geal.	Opens glottis.
Cricothyroid.	Cricoid cartilage.	Thyroid cartilage (lower inner bor- der).	Superior laryn- geal.	Makes vocal bands tense.
Crureus.	See <i>Vastus internus</i> .			
Deltoid.	Clavicle, acromion, and spine of scapula.	Shaft of humerus.	Circumflex.	Abducts humerus.
Depressor alæ nasi.	Incisive fossa of supe- rior maxillary bone.	Septum and ala of nose.	Facial: buccal branch.	Contracts nostril.
Depressor anguli oris.	External oblique line of inferior maxillary bone.	Angle of mouth.	Facial: supra- maxillary branch.	Depresses angle of mouth.
Depressor epiglott- idis.	Those fibers of the thyroepiglottideus that are continued forward to the margin of the epiglottis.			

TABLE OF MUSCLES.—Continued.

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
Depressor labii inferioris.	External oblique line of the inferior maxillary bone.	Lower lip.	Facial: supramaxillary branch.	Depresses lip.
Depressor urethrae.	Ramus of ischium near deep transversus perinei.	Fibers of constrictor vaginae muscle.		
Diaphragm.	Ensiform cartilage, 6 or 7 lower ribs, ligamenta arcuata, bodies of lumbar vertebrae.	Central tendon.	Phrenic and sympathetic.	Respiration.
Digastric (anterior belly).	Inner surface of inferior maxillary bone, near symphysis.	Hyoid bone.	Inferior dental.	Elevates hyoid and tongue.
Digastric (posterior belly).	Digastric groove of mastoid process.	Hyoid bone.	Facial.	Elevates hyoid and tongue.
Dilator naris anterioris.	Alar cartilage.	Border of ala of nose.	Facial: infraorbital branch.	Dilates nostril.
Dilator naris posterioris.	Nasal notch of superior maxillary bone.	Skin at margin of nostril.	Facial: infraorbital branch.	Dilates nostril.
Dorsal interossei, 4.	Sides of metacarpal bones.	Bases of corresponding phalanges.	Ulnar.	Abduct fingers from median line.
Dorsal interossei, 4.	Sides of metatarsal bones.	Base of first phalanx of corresponding toe.	External plantar.	Abduct toes.
Erector clitoridis.	Tuberosity of ischium.	Each side of crus of clitoris.		Erects clitoris.
Erector penis.	Ischial tuberosity, crus penis, and pubic ramus.	Tunica albuginea of corpus cavernosum.	Perineal.	To maintain erection.
Erector pili.	See <i>Arrectores pili</i> .			
Erector spinæ.	Iliac crest, back of sacrum, lumbar and three lower dorsal spines.	Divides into sacro-lumbalis, longissimus dorsi, and spinalis dorsi.	Lumbar nerves, posterior division.	Extension of lumbar spines on pelvis.
Extensor brevis digitorum (pedis).	Os calcis, externally.	First phalanx of great toe and tendons of extensor longus.	Anterior tibial.	Extends toes.
Extensor brevis hallucis.	A name applied to that portion of the extensor brevis digitorum that goes to the great toe.			
Extensor brevis pollicis.	See <i>Extensor primi inter nodii pollicis</i> .			
Extensor carpi radialis brevis.	External condyloid ridge of humerus.	Base second and third metacarpal.	Posterior interosseous.	Extends wrist.
Extensor carpi radialis longior.	Lower $\frac{1}{2}$ external condyloid ridge of humerus.	Base of second metacarpal.	Musculospiral.	Extends wrist.
Extensor carpi ulnaris.	1st head, external condyle of humerus. 2d head, posterior border of ulna.	Base of fifth metacarpal.	Posterior interosseous.	Extends wrist.
Extensor coccygis.	Last bone of sacrum or first of coccyx.	Lower part of coccyx.	Sacral branches.	Extends coccyx.

TABLE OF MUSCLES.—*Continued.*

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
Extensor communis digitorum.	External condyle of humerus.	All of the second and third phalanges.	Posterior interosseous.	Extends fingers.
Extensor indicis.	Back of ulna.	Second and third phalanges of index finger.	Posterior interosseous.	Extends index-finger.
Extensor longus digitorum pedis.	Outer tuberosity of tibia and shaft of fibula.	Second and third phalanges of toes.	Anterior tibial.	Extends toes.
Extensor longus pollicis.	See <i>Extensor secundi internodii pollicis</i> .			
Extensor minimi digiti.	External condyle of humerus.	Second and third phalanges of little finger.	Posterior interosseous.	Extensor of little finger.
Extensor ossis metacarpi pollicis.	Back of radius and ulna and interosseous membrane.	Base of metacarpal of thumb and fascia.	Posterior interosseous.	Extends thumb.
Extensor ossis metatarsi hallucis (anomalous).	Extensor proprius hallucis, or extensor communis digitorum, or tibialis anticus.	Metatarsal bone of hallux.		
Extensor primi internodii pollicis.	Back of radius.	Base of first phalanx of thumb.	Posterior interosseous.	Extends thumb.
Extensor proprius minimi digiti.	Lower part of ulna, or posterior ligament of wrist-joint.	Base of first phalanx of little finger.		Extends little finger.
Extensor proprius hallucis.	Middle of fibula.	Base of last phalanx of great toe.	Anterior tibial.	Extends great toe.
Extensor secundi internodii pollicis.	Back of ulna.	Base of last phalanx of thumb.	Posterior interosseous.	Extends thumb.
Flexor accessorius digitorum (of foot) (2 heads).	Inner and outer surface of os calcis.	Tendon of flexor longus digitorum.	External plantar.	Accessory flexor of toes.
Flexor accessorius longus digitorum (of foot).	Shaft of tibia or fibula.	Tubercle of os calcis, and joins tendon of long flexor.	External plantar.	Assists in flexing toes.
Flexor brevis digitorum (of foot).	Inner tuberosity of os calcis and plantar fascia.	Second phalanges of the lesser toes.	Internal plantar.	Flexes lesser toes.
Flexor brevis hallucis.	Under surface of cuboid, plantar ligaments, and external cuneiform.	Base of first phalanx of great toe.	Internal plantar.	Flexes and slightly adducts first phalanx of great toe.
Flexor brevis minimi digiti (of hand).	Unciform bone and annular ligament.	First phalanx of little finger.	Ulnar.	Flexes little finger.
Flexor brevis minimi digiti (of foot).	Base of fifth metatarsal.	Base of first phalanx of little toe.	External plantar.	Flexes little toe.
Flexor brevis pollicis (of hand).	2 heads— <i>outer</i> : lower border of anterior annular ligament; ridge of trapezium; <i>inner</i> : os magnum, and bases of first, second, and third metacarpal bones.	Base of first phalanx of thumb.	<i>Outer head</i> —median: palmar branch. <i>Inner head</i> —deep ulnar.	Flexes metacarpal bone of thumb.
Flexor brevis pollicis (of foot).	See <i>Flexor brevis hallucis</i> .	<i>lucis</i> .		

TABLE OF MUSCLES.—*Continued.*

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
Flexor carpi radi- alis.	Internal condyle of hu- merus.	Metacarpal bone of index finger.	Median.	Flexes wrist.
Flexor carpi ulna- ris (<i>2 heads</i>).	1. Internal condyle. 2. Olecranon and ulna.	5th metacarpal, annular lig. and pisiform bone.	Ulnar.	Flexes wrist.
Flexor indicis.	The name given to the indicial portion of the flexor profundus digi- torum, when it is dis- tinct.			
Flexor longus digi- torum (of foot).	Shaft of tibia.	Last phalanges of toes.	Posterior tibial.	Flexes phalanges and extends ankle.
Flexor longus hal- lucis.	Lower two-thirds of shaft of fibula.	Last phalanx of great toe.	Posterior tibial.	Flexes great toe.
Flexor longus pol- licis.	Shaft of radius and co- ronoid process of ulna.	Last phalanx of thumb.	Anterior interos- seous.	Flexes the thumb.
Flexor profundus digitorum.	Shaft of ulna.	Last phalanges by four tendons.	Ulnar and anterior interosseous.	Flexes the pha- langes.
Flexor sublimis digitorum (<i>3 heads</i>).	1. Inner condyle. 2. Coronoid process. 3. Oblique line of radius.	Second phalanges by four tendons.	Median.	Flexes second pha- langes.
Gastrocnemius (<i>2 heads</i>).	Condyles of femur.	Os calcis by tendo Achillis.	Internal popliteal.	Extends foot.
Gemellus inferior.	Tuberosity of ischium and lesser sacrospinous notch.	Great trochanter.	Sacral.	External rotator of thigh.
Gemellus superior.	Ichial spine and lesser sacrospinous notch.	Great trochanter.	Sacral.	External rotator of thigh.
Geniohyoglossus.	Superior genial tuberc- le of inferior maxil- lary bone.	Hyoid and inferior surface of tongue.	Hypoglossal.	Retracts and pro- trudes tongue.
Geniohyoid.	Inferior genial tubercle of inferior maxillary bone.	Body of hyoid.	Hypoglossal.	Elevates and ad- vances hyoid.
Gluteus maximus.	Superior curved iliac line and crest, sacrum, and coccyx.	Fascia and femur below great tro- chanter.	Inferior gluteal and sacral plexus.	Extends, abducts, and rotates thigh outward.
Gluteus medius.	Ilium between superior and middle curved lines.	Oblique line of great trochanter.	Superior gluteal.	Rotates, abducts, and advances thigh.
Gluteus minimus.	Ilium between middle and inferior curved lines.	Great trochanter.	Superior gluteal.	Rotates, abducts, and draws thigh forward.
Gracilis.	Rami of pubes and ischium.	Tibia, upper and inner part.	Obturator.	Flexes and abducts leg.
Gubernaculum tes- tis.	See <i>Cremaster</i> .			
Guthrie's.	See <i>Transversus peri-</i> <i>nei, deep</i> .			
Helicis major et minor.	Tubercle on helix.	Rim of helix near summit.	Auriculotemporal and posterior au- ricular.	
Hilton's.	See <i>Compressor sacculi</i> <i>laryngis</i> .			
Horner's.	See <i>Ciliary and Tensor</i> <i>tarsi</i> .			

TABLE OF MUSCLES.—*Continued.*

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
Hyoglossus.	Cornua of hyoid bone.	Side of tongue.	Hypoglossal.	Depresses side of tongue and retracts tongue.
Iliacus.	Iliac fossa and crest, base of sacrum.	Lesser trochanter, upper part shaft of femur.	Anterior crural.	Flexes and rotates thigh outward.
Iliocostal.	See <i>Sacro-lumbar</i> .			
Iliopsoas.	The iliacus and psoas muscles considered as one muscle.			
Infracostals, 10.	Inner surface of ribs.	Inner surface of two or three ribs above.	Intercostal.	Expiration, by depressing ribs.
Infraspinatus.	Infraspinous fossa.	Great tuberosity of humerus.	Suprascapular.	Rotates humerus outward.
Interarytenoid.	One arytenoid cartilage.	The other arytenoid cartilage.	Recurrent laryngeal.	Approximates arytenoid cartilages.
Intercostals, external, 11.	Outer lip of inferior costal border.	Superior border of ribs above.	Intercostal.	Raise ribs in inspiration.
Intercostals, internal, 11.	Inner lip of inferior costal border.	Superior border of rib below.	Intercostal.	Depress ribs in expiration.
Interossei of foot, dorsal (4).	Adjacent surfaces of metatarsal bones.	Bases of first phalanges.	External plantar.	Abduct from the middle line of the second toe.
Interossei of foot, plantar (3).	Inner lower surface of 3 outer metatarsal bones.	Bases of first phalanges of three outer toes.	External plantar.	Adduct the outer three toes.
Interossei of hand, dorsal (4).	Five metacarpal bones.	Sides of aponeurosis of extensor communis and adjacent parts of first phalanges.	Ulnar.	Abduct index, middle, and ring fingers, aid in flexing first phalanges and extending second and third.
Interossei of hand, palmar (3).	Sides of metacarpal bones.	Aponeurosis of extensor tendons, adjacent part of first phalanges.	Ulnar.	Adduct index, ring, and little fingers, aid in flexing first phalanges and in extending second and third.
Interspinales.	Upper surface of spines of vertebræ, near tip.	Posterior part of lower surface of spine above.	Internal divisions of posterior branches of spinal nerves.	Extend the vertebræ next above.
Intertransversales.	Between transverse processes of contiguous vertebræ.		Spinal nerves.	Flex the spinal column laterally.
Ischiocavernosus.	See <i>Erector penis</i> .			
Jarjavay's.	See <i>Depressor urethræ</i> .			
Latissimus dorsi.	Spines of 6 lower dorsal and lumbar and sacral vertebræ, crest of ilium, and 3 or 4 lower ribs.	Bicipital groove of humerus.	Long subscapular.	Draws arm backward and downward and rotates it inward.
Laxator tympani.	Spinous process of sphenoid bone and Eustachian tube.	Neck of malleus.	Facial.	Relaxes membrana tympani.

TABLE OF MUSCLES.—*Continued.*

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
Levator anguli oris.	Canine fossa of superior maxillary bone.	Angle of mouth.	Infraorbital branch of facial.	Elevates angle of mouth.
Levator anguli scapulae.	Transverse processes of four upper cervical vertebrae.	Posterior border of scapula.	Fifth cervical and cervical plexus.	Elevates upper angle of scapula.
Levator ani.	Posterior portion of body and ramus of pubes, pelvic fascia, ischial spine.	Rectum, coccyx, and fibrous raphe.	Sacral and perineal.	Supports rectum and vagina.
Levator labii inferioris.	Incisive fossa of inferior maxillary bone.	Skin of lower lip.	Supramaxillary branch of facial.	Elevates lower lip.
Levator labii superioris.	Lower margin of orbit.	Upper lip.	Infraorbital branch of facial.	Elevates upper lip.
Levator labii superioris alaeque nasae.	Nasal process of superior maxillary bone.	Alar cartilage and upper lip.	Infraorbital branch of facial.	Elevates upper lip, dilates nostril.
Levator menti.	See <i>Levator labii inferioris</i> .			
Levator palati.	Petrous portion of temporal bone.	Soft palate.	Sphenopalatine ganglion (facial).	Elevates soft palate.
Levator palpebrae superioris.	Lesser wing of sphenoid.	Upper tarsal cartilage.	Third cranial.	Lifts upper lid.
Levatores costarum, 12.	Transverse processes of last cervical and dorsal vertebrae.	Each to the rib below.	Intercostal.	Raise ribs.
Lingualis.	Under surface of tongue.		Chorda tympani and hypoglossal.	Elevates middle of tongue.
Longissimus dorsi.	Erector spinae.	Transverse processes of lumbar and dorsal vertebrae and 7th-11th ribs.	Branches of lumbar and dorsal.	Erects spine and bends trunk backward.
Longus colli:— 1. Superior oblique portion. 2. Inferior oblique portion. 3. Vertical portion.	Transverse processes 3d-5th cervical vertebrae. Bodies of 1st-3d dorsal vertebrae. Bodies of three dorsal and two cervical vertebrae.	Anterior tubercle of atlas. Transverse processes 5th-6th cervical vertebrae Bodies of 2d-4th cervical vertebrae	Lower cervical.	Flexes cervical vertebrae.
Lumbricales, 4, of foot.	Tendons of flexor longus digitorum.	First phalanges of the lesser toes.	Internal and external plantar.	Accessory flexors.
Lumbricales, 4, of hand.	Tendons of flexor profundus digitorum.	Tendons of common extensor.	Median and ulnar.	Flex first phalanges.
Masseter.	Zygomatic arch.	Angle and ramus of jaw.	Inferior maxillary.	Muscle of mastication.
Midriff.	See <i>Diaphragm</i> .			
Müller's.	See <i>Ciliary</i> .			
Müller's (orbital).	Connected with the orbital periosteum; it crosses the sphenomaxillary fissure.		Sympathetic.	
Müller's (superior palpebral).	Connected with the levator palpebrae superioris.	Along the upper border of the tarsus.	Sympathetic.	Assists in raising upper lid.

TABLE OF MUSCLES.—*Continued.*

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
Multiſidus ſpinæ.	Sacrum, iliac ſpine, articular proceſſes lumbar and cervical vertebræ, and tranſverſe proceſſes of dorsal and 7th cervical.	Laminæ and ſpines from laſt lumbar to ſecond cervical vertebræ.	Posterior ſpinal branches.	Erects and rotates ſpinal column.
Muſculus accēſſorius ad ſacrolumbalem.	Angles of ſix lower ribs.	Angles of ſix upper ribs.	Branches of dorsal.	Erects ſpine and bends trunk backward.
Mylohyoid.	Mylohyoid ridge of inferior maxillary bone.	Body of hyoid and raphe.	Inferior dental.	Elevates and advances hyoid bone.
Nasolabialis.	Nasal ſeptum.	Upper lip.	Facial.	Connects upper lip to ſeptum of noſe.
Obliquus auris.	Concha of ear.	Foſſa of antihelix.	Temporal and poſterior auricular.	
Obliquus capitis inferior.	Spinous proceſſus of axis.	Tranſverſe proceſſus of atlas.	Suboccipital and great occipital.	Rotates atlas and cranium.
Obliquus capitis ſuperior.	Tranſverſe proceſſus of atlas.	Occipital bone.	Suboccipital and great occipital.	Draws head backward.
Obliquus externus.	Eight lower ribs.	Middle line, iliac creſt, Poupart's ligament.	Intercostal, iliohypogaſtric, ilioinguinal.	Compreſſes viſcera and flexes thorax.
Obliquus inferior.	Orbital plate of ſuperior maxillary bone.	Sclerotic.	Third cranial.	Rotates eyeball upward and outward.
Obliquus internus.	Lumbar fascia, iliac creſt, Poupart's ligament.	Three lower ribs, linea alba, pubic creſt, pectineal line.	Intercostal, iliohypogaſtric, ilioinguinal.	Compreſſes viſcera, flexes thorax, and aſſiſts in expiration.
Obliquus ſuperior.	Above optic foramen, through pulley.	Sclerotic.	Fourth cranial.	Rotates eyeball downward and inward.
Obturator externus.	Obturator foramen and membrane.	Digital foſſa at baſe of great trochanter.	Obturator.	External rotator of thigh.
Obturator internus.	Obturator foramen and membrane.	Great trochanter.	Sacral.	External rotator of thigh.
Occipitalis.	See <i>Occipitofrontalis</i> .			
Occipitofrontalis.	Superior curved line of occiput and angular proceſſus of frontal bone.	Aponeuroſis.	Posterior auricular, ſmall occipital, facial.	Moves ſcalp.
Omohyoid.	Upper border of ſcapula.	Body of hyoid bone.	Deſcendens and communicans noni.	Depreſſes and retracts hyoid bone.
Opponens minimi digiti.	Unciform bone.	Fifth metacarpal.	Ulnar.	Flexes little finger.
Opponens pollicis.	Trapezium, anterior annular ligament.	Metacarpal bone of thumb.	Median, palmar diſiſion.	Flexes thumb.
Orbicularis oris.	Nasal ſeptum and canine foſſa of inferior maxilla, by accēſſory fibers.	Forms lips and ſphincter of mouth.	Buccal and ſupramaxillary branches of facial.	Cloſes mouth.
Orbicularis palpebrarum.	Meſal margin of orbit.	Lateral margin of orbit.	Facial.	Cloſes eyelids.

TABLE OF MUSCLES.—*Continued.*

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
Palatoglossus.	Soft palate.	Side and dorsum of tongue.	Sphenopalatine ganglion.	Constricts the fauces.
Palatopharyngeus.	Soft palate.	Thyroid cartilage and pharynx.	Sphenopalatine ganglion.	Closes posterior nares.
Palmaris brevis.	Annular ligament and palmar fascia.	Skin of palm of hand.	Ulnar.	Corrugates skin of palm.
Palmaris interossei.	Palmar surfaces second, fourth, and fifth metacarpal bones.	Bases of first phalanges of corresponding fingers.	Ulnar.	Adductors of fingers.
Palmaris longus.	Internal condyle of humerus.	Annular ligament and palmar fascia.	Median.	Renders palmar fascia tense.
Pectineus.	Iliopectineal line and pubes.	Femur below lesser trochanter.	Anterior crural, obturator.	Flexes and rotates the thigh outward.
Pectoralis major.	Clavicle, sternum, and costal cartilages.	External bicipital ridge of humerus.	Anterior thoracic, external and internal.	Draws arm downward and forward.
Pectoralis minor.	Third, fourth, and fifth ribs.	Coracoid process.	Anterior thoracic.	Depresses point of shoulder.
Peroneus brevis.	Middle third of shaft of fibula, externally.	Base of fifth metatarsal bone.	Musculocutaneous.	Extends foot.
Peroneus longus.	Head and shaft of fibula.	First metatarsal of great toe.	Musculocutaneous.	Extends and everts foot.
Peroneus tertius.	Lower fourth of fibula.	Fifth metatarsal bone.	Anterior tibial.	Flexes tarsus.
Plantaris.	Outer bifurcation of linea aspera and posterior ligament of knee-joint.	Os calcis by means of the tendo Achillis.	Internal popliteal.	Extends foot.
Plantaris interossei.	Shafts of 3d, 4th, and 5th metatarsal bones.	Bases of first phalanges of corresponding toes.	External plantar.	Adduct toes.
Platysma myoides.	Clavicle, acromion, and fascia.	Inferior maxillary bone, angle of mouth.	Facial and superficial cervical.	Wrinkles skin and depresses mouth.
Popliteus.	External condyle of femur.	Shaft of tibia above oblique line.	Internal popliteal.	Flexes leg.
Pronator quadratus.	Lower fourth of ulna.	Lower fourth of shaft of radius.	Anterior interosseous.	Pronates hand.
Pronator radii teres.	Internal condyle of humerus and coronoid process of ulna.	Outer side of shaft of radius.	Median.	Pronates hand.
Psoas magnus.	Bodies and transverse processes of last dorsal and all lumbar vertebræ.	Lesser trochanter.	Lumbar.	Flexes and rotates thigh outward, and flexes trunk on pelvis.
Psoas parvus.	Bodies of last dorsal and first lumbar vertebræ.	Iliopectineal eminence and iliac fascia.	Lumbar.	Flexes pelvis upon abdomen.
Pterygoid (external).	Two heads: 1, external pterygoid plate of sphenoid bone; 2, great wing of sphenoid bone.	Neck of condyle of lower jaw.	Inferior maxillary.	Draws inferior maxillary bone forward.
Pterygoid (internal).	Pterygoid fossa of sphenoid bone.	Inner surface of angle of jaw.	Inferior maxillary.	Raises and draws inferior maxilla forward.

TABLE OF MUSCLES.—*Continued.*

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
Pyramidalis.	Pubes.	Linea alba.	Iliohypogastric.	Renders linea alba tense.
Pyramidalis nasi.	Occipitofrontalis.	Compressor naris.	Infraorbital branch of facial.	Depresses inner angle of eyebrow.
Pyriformis.	Front of sacrum, through great sciatic foramen.	Great trochanter.	Branch of sacral plexus.	External rotator of thigh.
Quadratus femoris.	Tuberosity of the ischium.	Quadrate line of femur.	Fifth lumbar and first sacral.	External rotator of thigh.
Quadratus lumborum.	Crest of ilium, transverse processes of lower three lumbar vertebræ.	Last rib, transverse processes of upper three lumbar vertebræ.	Twelfth thoracic and upper lumbar.	Flexes thorax laterally.
Quadriceps extensor femoris.	Includes the rectus, vastus internus and externus, and crureus muscles. Their common tendon surrounds the patella.			
Rectus abdominis.	Pubic crest and fibrous tissues in front of symphysis.	Cartilages of the ribs, from the fifth to the seventh.	Intercostal, iliohypogastric, ilioinguinal.	Compresses viscera and flexes thorax.
Rectus capitis anticus major.	Transverse processes third to sixth cervical vertebræ.	Basilar process of occipital bone.	First and second cervical.	Flexes head and slightly rotates it.
Rectus capitis anticus minor.	Transverse process and lateral mass of atlas.	Basilar process of occipital bone.	First cervical.	Flexes head.
Rectus capitis lateralis.	Ventral cephalic surface of lateral mass of atlas.	Jugular process of occipital bone.	First cervical.	Flexes head laterally.
Rectus capitis posticus major.	Spine of axis.	Inferior curved line of occipital bone.	Suboccipital and great occipital.	Rotates head.
Rectus capitis posticus minor.	Dorsal arch of atlas.	Below inferior curved line of occipital bone.	Suboccipital and great occipital.	Draws head backward.
Rectus externus.	Two heads, outer margin of optic foramen.	Sclera.	Sixth cranial.	Rotates eyeball outward.
Rectus femoris.	Anterior inferior iliac spine, brim of acetabulum.	Proximal border of patella.	Anterior crural.	Extends leg.
Rectus inferior.	Lower margin of optic foramen.	Sclera.	Third cranial.	Rotates eyeball downward.
Rectus internus.	Inner margin of optic foramen.	Sclera.	Third cranial.	Rotates eyeball inward.
Rectus superior.	Upper margin of optic foramen.	Sclera.	Third cranial.	Rotates eyeball upward.
Retrahens aurem.	Mastoid process.	Concha.	Posterior auricular.	Retracts pinna.
Rhomboideus major.	Spines of first five thoracic vertebræ.	Root of spine of scapula.	Fifth cervical.	Elevates and retracts scapula.
Rhomboideus minor.	Spines of seventh cervical and first dorsal vertebræ.	Root of spine of scapula.	Fifth cervical.	Retracts and elevates scapula.
Risorius.	Fascia over masseter.	Angle of mouth.	Buccal branch of facial.	Draws angle laterally.
Rotatores spinæ.	Transverse processes of from second to twelfth thoracic vertebræ.	Lamina of next vertebra above.	Dorsal branches of spinal.	Rotate spinal column.

TABLE OF MUSCLES.—Continued.

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
* Sacrolumbalis.	Erector spine.	Angle of six lower ribs.	Branches of dorsal.	Erects spine and bends trunk backward.
Sartorius.	Anterior superior spine of ilium.	Upper internal portion of shaft of tibia.	Anterior crural.	Flexes leg upon thigh; thigh upon pelvis; rotates thigh outward.
Scalenus anticus.	Scalene tubercle on first rib.	Transverse processes 3d-6th cervical vertebrae.	Lower cervical.	Flexes neck laterally.
Scalenus medius.	First rib.	Transverse processes of six lower cervical vertebrae.	Lower cervical.	Flexes neck laterally.
Scalenus posticus.	Second rib.	Transverse processes of three lower cervical vertebrae.	Lower cervical.	Bends neck laterally.
Semimembranosus.	Tuberosity of ischium.	Inner tuberosity of tibia.	Great sciatic.	Flexes leg and rotates it inward.
Semispinalis colli.	Transverse processes four upper dorsal and articular processes four lower cervical vertebrae.	Spines of second to fifth cervical vertebrae.	Cervical branches.	Erects spinal column.
Semispinalis dorsal.	Transverse processes sixth to tenth dorsal vertebrae.	Spines last two cervical and first four thoracic.	Branches of dorsal.	Erects spinal column.
Semitendinosus.	Tuberosity of ischium.	Upper and inner surface of tibia.	Great sciatic.	Flexes leg on thigh.
Serratus magnus.	Eight upper ribs.	Inner margin of dorsal border of scapula.	Posterior thoracic.	Elevates ribs in inspiration.
Serratus posticus inferior.	Spines of last two thoracic and first three lumbar.	Four lower ribs.	Tenth and eleventh intercostal.	Depresses ribs in expiration.
Serratus posticus superior.	Spines of seventh cervical and first two thoracic vertebrae.	Second, third, fourth, and fifth ribs.	Second and third intercostal.	Raises ribs in inspiration.
Soleus.	Shaft of fibula, oblique line of tibia.	Os calcis by tendo Achillis.	Internal popliteal and posterior tibial.	Extends foot.
Sphincter ani, external.	Tip of coccyx.	Tendinous center of perineum.	Perineal, pudic, and fourth sacral.	Closes anus.
Sphincter ani, internal.	A thickening of the circular fibers of the intestine an inch above the anus.		Hemorrhoidal nerves.	Constricts rectum.
Sphincter vaginae.	Central tendon of perineum.	Corpora cavernosa and clitoris.	Homologue of accelerator urinae in the man.	
Sphincter vesicae internus.	Near the urethral orifice of the bladder.		Vesical nerves.	Constricts internal orifice of urethra.
Spinalis cervicis (normal, but inconstant).	Spines fifth, sixth, and seventh cervical and first two thoracic vertebrae.	Spine of axis, sometimes spines of 3d and 4th cervical vertebrae.		
Spinalis colli.	Spines of fifth and sixth cervical vertebrae.	Spine of axis, or third and fourth cervical spines.	Cervical branches.	Steadies neck.

TABLE OF MUSCLES.—*Continued.*

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
Spinalis dorsi.	Last two thoracic and first two lumbar spines.	Remaining thoracic spines.	Dorsal branches.	Erects spinal column.
Splenius capitis.	Lower two-thirds ligamentum nuchæ, spines of seventh cervical and first two thoracic vertebræ.	Outer third of middle oblique line of occiput and outer surface of mastoid process.	Middle cervical, posterior branches.	Extends head and neck and rotates and flexes laterally.
Splenius colli.	Spines of third to sixth thoracic vertebræ.	Dorsal tubercles of transverse processes of upper three or four cervical vertebræ.	Posterior divisions of lower cervical.	Extends, flexes laterally, and rotates neck.
Stapedius.	Interior of pyramid.	Neck of stapes.	Facial.	Depresses base of stapes.
Sternocleidomastoid.	Two heads, sternum and clavicle.	Mastoid process and outer half of superior oblique line of occiput.	Spinal accessory and cervical plexus.	Depresses and rotates head.
Sternohyoid.	Sternum and clavicle.	Hyoid bone.	Descending and communicating branches of the hypoglossal.	Depresses hyoid bone.
Sternothyroid.	Sternum and cartilage of first rib.	Side of thyroid cartilage.	Descendens and communicans hypoglossi.	Depresses larynx.
Styloglossus.	Styloid process.	Side of tongue.	Hypoglossal.	Elevates and retracts tongue.
Stylohyoid.	Styloid process.	Body of hyoid.	Facial.	Draws hyoid upward and backward.
Stylopharyngeus.	Styloid process.	Thyroid cartilage.	Glossopharyngeal, and pharyngeal plexus.	Elevates pharynx.
Subanconeus.	Humerus above olecranon fossa.	Posterior ligament of elbow.	Musculospiral.	Tensor of ligament.
Subclavius.	Cartilage of first rib.	Inferior surface of clavicle.	Fifth and sixth cervical.	Draws clavicle downward.
Subcrureus.	Anterior distal part of femur.	Synovial sac behind patella.	Anterior crural.	Draws up synovial sac.
Subscapularis.	Under surface of scapula.	Humerus, lesser tuberosity and shaft.	Subscapular.	Chief internal rotator of humerus.
Supinator longus.	External condyloid ridge of humerus.	Styloid process of radius.	Musculospiral.	Flexes and supinates forearm.
Supinator radii brevis.	External condyle of humerus, oblique line of ulna.	Neck of radius and its bicipital tuberosity.	Posterior interosseous.	Supinates hand.
Supraspinales.	Lie on spinous processes in cervical region.			
Supraspinatus.	Supraspinous fossa.	Great tuberosity of humerus.	Suprascapular.	Supports shoulder-joint, raises arm.
Tailors'.	See <i>Sartorius</i> .			
Temporal.	Temporal fossa and fascia.	Coronoid process of mandible.	Inferior maxillary.	Elevates mandible.
Tensor palati.	Scaphoid fossa and alar spine of sphenoid.	Soft palate; winds about hamular process.	Otic ganglion.	Renders palate tense.

TABLE OF MUSCLES.—*Continued.*

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
Tensor tarsi.	Crest of lacrimal bone.	Tarsal cartilages.	Infraorbital branch of facial.	Compresses puncta and lacrimal sac.
Tensor tympani.	Temporal bone, Eustachian tube and canal, sphenoid bone.	Handle of malleus.	Otic ganglion.	Renders tense the membrana tympani.
Tensor vaginae femoris.	Iliac crest and anterior superior spinous process.	Fascia lata.	Superior gluteal.	Tensor of fascia lata.
Teres major.	Inferior angle of scapula.	Internal bicipital ridge of humerus.	Subscapular.	Draws arm downward and backward.
Teres minor.	Axillary border of scapula.	Great tuberosity of humerus.	Circumflex.	Rotates humerus outward and adducts it.
Thyroarytenoid-eus.	Thyroid cartilage and cricothyroid membrane.	Arytenoid, inferior and anterior surface.	Recurrent laryngeal.	Relaxes vocal bands.
Thyroepiglottideus.	Inner surface of thyroid cartilage.	Epiglottis and arytenoepiglottidean fold.	Recurrent laryngeal.	Depresses epiglottis.
Thyrohyoid.	Side of thyroid cartilage.	Body and greater cornu of hyoid bone.	Hypoglossal.	Elevates larynx.
Tibialis anticus.	Outer tuberosity and upper part of shaft of tibia.	Internal cuneiform and first metatarsal bone.	Anterior tibial.	Flexes tarsus and elevates inner border of foot.
Tibialis posticus.	Shaft of fibula and tibia, interosseous membrane.	Tuberosity of scaphoid, internal cuneiform, and bases of second to fourth metatarsal.	Posterior tibial.	Extends tarsus and inverts foot.
Trachelomastoid.	Transverse processes of 3d-6th thoracic, and articular processes of last three or four cervical vertebrae.	Mastoid process.	Branches of cervical.	Steadies head.
Tragicus.	Tragus.	Tragus.	Temporal and posterior auricular.	
Transversalis abdominis.	Poupart's ligament, iliac crest, six lower ribs, lumbar vertebrae.	Linea alba, pubic crest, pectineal line.	Intercostal, iliohypogastric, ilioinguinal.	Compresses viscera and flexes thorax.
Transversalis colli.	Transverse processes of third to sixth thoracic vertebrae.	Transverse processes of five lower cervical vertebrae.	Cervical branches.	Keeps neck erect.
Transversus auris.	Convexity of concha.	Convexity over groove of helix.	Temporal and posterior auricular.	Retracts helix.
Transversus pedis.	Head fifth metatarsal and plantar ligaments of metatarsophalangeal joints.	First phalanx of great toe.	External plantar.	Adducts great toe.
Transversus perinei.	Ramus of ischium.	Central tendon.	Perineal.	Tensor of central tendon.
Transversus perinei, deep.	See <i>Compressor urethrae</i> .			

TABLE OF MUSCLES.—*Continued.*

NAME.	ORIGIN.	INSERTION.	INNERVATION.	FUNCTION.
Trapezius.	Superior curved line of occipital bone, spinous processes of last cervical and all the dorsal vertebræ.	Clavicle, spine of scapula, and acromion.	Spinal accessory and cervical plexus.	Draws head backward.
Triangularis sterni.	Ensiform cartilage, costal cartilages of three or four lower true ribs, and sternum.	Border of inner surfaces 2d, 3d, 4th, and 5th costal cartilages.	Intercostal.	Expiration.
Triceps (3 heads).	<i>External</i> and <i>internal</i> near musculospiral groove, shaft of humerus; <i>middle</i> or <i>long</i> , lower margin of glenoid cavity.	Olecranon process of ulna.	Musculospiral.	Extends forearm.
Vastus externus.	Anterior border great trochanter and linea aspera of femur.	Tuberosity of tibia.	Anterior crural.	Extends leg.
Vastus internus.	Inner lip of linea aspera of femur.	Tuberosity of tibia.	Anterior crural.	Extends leg.
Zygomaticus major et minor.	Malar bone.	Angle of mouth.	Infraorbital branch of facial.	Draw lip upward and outward.

Muscular (*mus'-ku-lar*) [*musculus*, muscle].

1. Pertaining to or affecting muscles. 2. Having well-developed muscles. **M. Rheumatism**, rheumatism affecting the muscles. **M. Sense**, the sense of motion, weight, and position upon which the adjustment of the body to its surroundings depends. **M. Tissue**, the tissue of which muscles are composed; it is of two principal varieties—the striped, striated, or voluntary, and the unstriped, nonstriated, or involuntary muscular tissue. The striped muscular tissue is composed of muscle-fibers, the unstriped of elongated, spindle-shaped cells. A muscle-fiber consists of sarcolemma, muscle-nuclei, and muscle-substance. The sarcolemma is an elastic, homogeneous connective-tissue sheath which lightly invests the muscle-fiber. The nuclei are fusiform in shape, are situated immediately beneath the sarcolemma, and run parallel with the axis of the fiber. The muscle-fiber itself consists of two substances—a dark, doubly-refracting, or anisotropic contractile substance, and a lighter, semifluid, isotropic substance termed sarcoplasm. The contractile substance consists of delicate spindles, the apposition of the thicker portions of which produces the dark transverse disc. The spindles terminate in minute beads, the juxtaposition of which gives rise to the intermediate disc or membrane of Krause. The neutral sarcoplasm fills out the spaces left between the spindles, and, as ordinarily the tapering ends of the spindles are too delicate

to be seen, the space between the intermediate and transverse discs—the lateral disc—looks homogeneous. The contractile fibrillæ into which the contractile substance is divided, formed by the end-to-end union of the spindles, are aggregated into bundles—the muscle-columns—surrounded by a layer of sarcolemma. On transverse section these muscle-columns give rise to the appearance known as Cohnheim's fields. The muscle-fibers are held together by delicate areolar tissue, the endomysium. Several grouped together form a primary bundle, which is surrounded by a sheath termed the perimysium. Fasciculi are aggregations of primary bundles, and are the units of which the complete muscle is composed, the latter being surrounded by the epimysium. **M. Tone**. See *Tone*. **M. Tumor**. See *Myoma*.

Muscularis mucosæ (*mus-ku-la'-ris mu-ko'-se*). The layer of unstriped muscular tissue separating the mucosa (of mucous membranes) from the submucosa.

Musculature (*mus'-ku-la-târ*) [*musculus*, muscle]. The muscular system of the body, or a part of it.

Musculin (*mus'-ku-lin*) [*musculus*, a muscle]. Extract of muscle-tissue. See *Organotherapy*.

Musculocutaneous (*mus-ku-lo-ku-ta'-ne-us*) [*musculus*, muscle; *cutis*, skin]. Pertaining to or supplying the muscles and skin, as the M. nerve of the arm or leg.

Musculomembranous (*mus-ku-lo-mem'-bran-us*) [*musculus*, muscle; *membrana*,

membrane]. Pertaining to or consisting of both muscles and membrane.

Musculophrenic (*mus-ku-lo-fren'-ik*) [*musculus*, muscle; *φρήν*, diaphragm]. Pertaining to or supplying the muscles and the diaphragm, as the M. nerve.

Musculospiral (*mus-ku-lo-spi'-ral*) [*musculus*, muscle]. 1. Supplying muscles and having a spiral course. 2. Pertaining to the musculospiral nerve.

Mushroom. See *Agaric*.

Musk [Skt., *mushka*, a testicle]. The dried secretions from the preputial follicles of *Moschus moschiferus*, a mammal of the order of Ruminantia. M. (*Moschus*, U. S. P.), occurs in grains or lumps, possessing a peculiar, penetrating odor. Its composition is complex, and among its constituents are ammonia, stearin, olein, cholesterolin, a volatile oil, gelatin, albumin, and certain salts. M. is stimulant and antispasmodic, and is used in typhoid fever and other low fevers, in adynamic pneumonia, in laryngismus stridulus, and in hiccough. Dose gr. x (0.65).

Mussitation (*mus-it-a'-shun*) [*mussitare*, to mutter]. Movement of the lips without the production of articulate speech.

Must [*mustum*, new wine]. The juice freshly expressed from grapes.

Mustard (*mus'-terd*) [*mustum*, must, new wine, with Teut. suffix]. 1. A plant of the genus *Brassica* (*Sinapis*), of the natural order Crucifere. 2. The seed of the mustard-plant. The seeds of two species are chiefly employed, those of *Brassica nigra* yielding black mustard, or *Sinapis nigra* (U. S. P.), and those of *Brassica alba* yielding white mustard, or *Sinapis alba* (U. S. P.). The seeds contain a fixed oil consisting of the glycerol-compounds of stearic, oleic, and erucic or brassic acid, and of sinapoleic and behenic acids, sinalbin (in white mustard), and (in black mustard) a volatile oil, oleum sinapis volatile (U. S. P.). The volatile oil contains allyl isosulphocyanid, the active principle, which does not exist preformed in the seeds, but is produced by the decomposition of potassium myronate, or sinigrin, under the influence of a ferment, myrosin. Mustard is used in dyspepsia, as an emetic (dose ʒj-ij (4.0-8.0)), as a rubefacient and counter-irritant, especially in the form of mustard-plaster, and in baths. Linimentum sinapis compositus is used locally. Mustard-plaster (sinapism) is made by mixing mustard and flour with water. M.-paper (*charta sinapis*, B. P.) is used locally. M.-poultice (*cataplasma sinapis*, B. P.) is made by mixing mustard and linseed-meal with water.

Mutacism (*mu'-tas-izm*) [*μετακταμός*, mutacism]. The too frequent use of the m sound, and its substitution for other sounds.

Mute (*mūt*) [*mutus*, dumb]. Dumb; unable to use articulate speech.

Mutilation (*mu-til-a'-shun*) [*mutilare*, to cut]. 1. The act of maiming or disfiguring. 2. The state thereby produced. Also, the condition of the parts remaining after the excision or amputation of a member or part.

Mutism (*mu'-tizm*) [*mutus*, dumb]. Dumbness.

Myalgia (*mi-al'-je-ah*) [*μῦς*, muscle; *ἀλγος*, pain]. Pain in the muscles; muscular rheumatism.

Myasthenia (*mi-as-then-i'-ah*, *mi-as-the'-ne-ah*) [*μῦς*, muscle; *ἀσθένεια*, weakness]. Muscular debility. M. *gravis pseudo-paralytica*, a disease characterized by an abnormal exhaustibility of the voluntary muscles, manifesting itself in a rapid diminution of contractility both when the muscle is innervated by the will and when stimulated by the electric current.

Myasthenic (*mi-as-then'-ik*) [*μῦς*, muscle; *ἀσθένεια*, weakness]. Characterized by myasthenia.

Mycelium (*mi-se'-le-um*) [*μύκης*, a fungus; *ἕλος*, nail, wart, excrescence]. The vegetative filaments of fungi, usually forming interwoven masses.

Mycetoma (*mi-set-o'-mah*). See *Fungus-foot*.

Mycoderma (*mi-ko-der'-mah*) [*μύκης*, fungus; *δέρμα*, skin]. A genus of fungi forming membranes upon or in fermenting liquids. M. *aceti*, the microorganism of acetous fermentation.

Mycobemia (*mi-ko-hem'-e-ah*) [*μύκης*, a fungus; *αἷμα*, blood]. A condition in which fungi are in the blood.

Mycology (*mi-kol'-o-je*) [*μύκης*, a fungus; *λόγος*, science]. The science of fungi.

Myconostoc (*mi-kon-as'-tok*) [*μύκης*, fungus; *νόστος*]. A genus of schizomycetes.

Mycophylaxin (*mi-ko-fil-ak'-in*) [*μύκης*, a fungus; *φύλαξ*, a protector]. See *Proteids*, *Defensive*.

Mycoprotein (*mi-ko-pro'-te-in*) [*μύκης*, a fungus; *protein*, from *πρώτος*, first]. The albuminoid substance obtained from the bodies of bacteria.

Mycosis (*mi-ko'-sis*) [*μύκης*, fungus]. 1. A growth of fungi within the body. 2. A disease caused by the presence of microorganisms in the body. M. *fungoides*, an affection of the skin characterized by the appearance, chiefly on the scalp, face, and chest, of pinkish or reddish, nodular or flattened tumors, which may go on to ulceration. Itching is often present. The disease is believed to be microorganismal in origin. M. *intestinalis*. Synonym of *Anthrax*.

Mycosozin (*mi-ko-so'-zin*) [*μύκης*, a fungus; *σώζω*, to save]. See *Sporei*.

Mycotic (*mi-kot'-ik*) [*μύκης*, fungus]. Characterized by mycosis; due to microorganisms. **M.** Endocarditis, that due to microorganismal infection. A synonym of ulcerative or malignant endocarditis.

Mydaine (*mid-a'-le-in*) [*μυδαίνος*, putrid]. A ptomain obtained from putrefying cadaveric organs. It produces salivation, dilatation of the pupil, elevation of temperature, peristaltic action of the intestines, first a rise, then a fall of temperature, and diastolic arrest of the heart. See *Ptomains*, *Table of*.

Mydatoxin (*mid-at-oks'-in*) [*μυδάειν*, to be wet; *τοξικόν*, poison], $C_8H_{11}NO_2$. A ptomain obtained from horse-flesh and human flesh. It produces symptoms resembling those of curara-poisoning.

Mydin (*mid'-in*) [*μυδίν*, to be wet], $C_8H_{11}NO$. A nonpoisonous ptomain produced in the putrefaction of human internal organs. The free base is strongly alkaline, has an ammoniacal odor, and is characterized by its strong reducing properties. See *Ptomains*, *Table of*.

Mydriasis (*mid-ri'-as-is*) [*μυδρίασις*]. Dilatation of the pupil of the eye. **M.** Paralytic, that due to paralysis of the oculomotor nerve. **M.** Spasmodic, **M.** Spastic, that caused by overaction of the sympathetic or dilator nerve of the iris. **M.** Spinal, that produced by irritation of the ciliospinal center of the spinal cord.

Mydriatic (*mid-re-at'-ik*) [*μυδρίασις*, dilatation of the pupil]. 1. Causing mydriasis, or dilatation of the pupil. 2. An agent causing mydriasis, or dilatation of the pupil.

Myel (*mi'-el*) [*μυελός*, marrow]. The spinal cord.

Myelencephalon (*mi-el-en-sef'-al-on*) [*μυελός*, marrow; *ἐγκεφαλος*, brain]. 1. The cerebrospinal axis. 2. See *Metencephalon*.

Myelin (*mi'-el-in*) [*μυελός*, marrow]. 1. The white substance of Schwann—the sheath of a medullated nerve. 2. A substance obtained from nerve-tissue. See *Organotherapy*.

Myelinic (*mi-el-in'-ik*) [*μυελός*, marrow]. Relating to myelin, or to myelinic nerve-fibers.

Myelitis (*mi-el-i'-tis*) [*μυελός*, marrow; *ιτις*, inflammation]. 1. Inflammation of the spinal cord. If it affects the gray matter, it is called poliomyelitis; if the white matter, leukomyelitis; if it extends entirely across the cord, transverse myelitis. **M.** Acute, that due to exposure to cold, injuries, or to acute general diseases; it is attended with sensory disturbances and motor paralysis, complete or incomplete, according to the part of the cord involved and the degree of the inflammation. **M.** Ascending, a form in which the inflammation

travels up the cord. **M.** Bulbar, that affecting the medulla and manifesting itself in disturbances of the cardiorespiratory apparatus, dysphagia, vomiting, and other symptoms. **M.** Central, myelitis limited chiefly to the gray matter of the cord. **M.** Chronic, a slowly advancing myelitis, presenting generally the same features as acute myelitis, but extended over a longer period.

M. Compression-, an inflammation of the spinal cord, secondary to compressing lesions outside of the cord (caries, carcinoma, exostosis of the vertebrae, aneurysm) or to tumors of the cord itself. **M.** Cornual, inflammation affecting the anterior or posterior cornua. **M.** Descending, a form in which the inflammation extends downward, following the pyramidal tracts. **M.** Diffuse, widely distributed inflammation of the cord involving large areas. **M.** Disseminated, one in which there are several foci. **M.** Focal, one in which a small area is affected. **M.** Hemorrhagic, myelitis associated with or dependent upon hemorrhage. **M.** Pressure-. See *M.* Compression-. **M.** Sclerotic, a form characterized by overgrowth of the interstitial tissue, which undergoes contraction, producing an increase of hardness of the cord. **M.** Systemic, a variety in which the inflammation is confined to distinct systems or tracts of the cord. **M.** Traumatic, myelitis following direct injury.

The symptoms of myelitis vary with the character and the seat of the lesion. The sensory disturbances consist of hyperesthesia or anesthesia, girdle-pains, and usually a hyperesthetic zone at the level of the lesion. The reflexes are generally lost in the beginning, but later become exaggerated; sometimes they do not return. Paralysis is present in the parts below the lesion; the muscles are flaccid at first, later rigid. The sphincters are generally paralyzed. Bedsores are common in certain forms of myelitis. 2. Inflammation of bone-marrow; osteomyelitis.

Myelo- (*mi'-el-o-*) [*μυελός*, marrow]. A prefix denoting reference to or connection with the spinal cord, or with the bone-marrow.

Myelocoele (*mi'-el-o-sel*) [*μυελός*, marrow; *κοιλία*, cavity]. The central canal of the spinal cord.

Myelocoele (*mi'-el-o-sel*) [*μυελός*, marrow; *κῆλη*, tumor]. Spina bifida.

Myelocyte (*mi'-el-o-sit*) [*μυελός*, marrow; *κύτος*, cell]. 1. A large uninuclear leukocyte containing neutrophile granules, and supposed to be derived from the bone-marrow. 2. A free nucleus of a central or ganglionic nerve-cell.

Myelogenic (*mi-el-o-jen'-ik*) [*μυελός*, marrow; *γενᾶν*, to produce]. Produced in or

by bone marrow. **M. Leukemia**, leukemia due to disease of the bone-marrow.

Myeloid (*mi'-el-oid*) [*μυελός*, marrow; *εἶδος*, like]. Resembling marrow. **M. Cell**, myeloplax. **M. Sarcoma**, a form of sarcoma distinguished by the presence of multi-nucleated cells.

Myeloma (*mi-el-o'-mah*) [*μυελός*, marrow; *δῆμα*, tumor]. 1. An encephaloid tumor. 2. A giant-celled sarcoma.

Myelomalacia (*mi-el-o-mal-a'-se-ah*) [*μυελός*, marrow; *μαλακία*, softening]. A softening of the spinal cord.

Myelomeningitis (*mi-el-o-men-in-jit'-tis*) [*μυελός*, marrow; *μῆνιξ*, a membrane; *ιτις*, inflammation]. Inflammation of the membranes of the spinal cord; spinal meningitis.

Myelomeningocele (*mi-el-o-men-in'-go-sel*) [*μυελός*, marrow; *μῆνιξ*, membrane; *κῆλη*, tumor]. Spina bifida.

Myelomyces (*mi-el-om'-is-es*) [*μυελός*, marrow; *μύκης*, fungus]. An encephaloid tumor.

Myelon (*mi'-el-on*) [*μυελός*, marrow]. The spinal cord.

Myelopathy (*mi-el-op'-ath-e*) [*μυελός*, marrow; *πάθος*, disease]. Any disease of the spinal cord.

Myeloplast (*mi'-el-o-plast*) [*μυελός*, marrow; *πλαστός*, shaped]. The peculiar cells of the bone-marrow resembling leukocytes.

Myeloplax (*mi'-el-o-plaks*) [*μυελός*, marrow; *πλάξ*, a plaque]. One of the large multi-nucleated cells found upon the inner surface of bone, and concerned in its absorption.

Myelosclerosis (*mi-el-o-skle-ro'-sis*) [*μυελός*, marrow; *σκληρώς*, hard]. Sclerosis of the spinal cord or of the marrow of bones.

Myelospongium (*mi-el-o-spon'-je-um*) [*μυελός*, marrow; *σπόγγος*, sponge]. A network in the wall of the neural canal of the embryo, composed of processes given off by the outer extremities of the spongioblasts.

Myelosyphilosis (*mi-el-o-sif-il-o'-sis*) [*μυελός*, marrow; *syphilis*, syphilis]. Syphilitic disease of the spinal cord.

Myiodeopsia, Myiodesopsia (*mi-e-o-de-op'-se-ah, mi-i-o-des-op'-se-ah*) [*μυιόδωψ*, fly-like; *ὄψις*, vision]. The subjective appearance of muscæ volitantes.

Myiosis (*mi'-i-o-sis*) [*μύια*, a fly]. The presence of larvae of flies, or maggots, in or on the body, as in the nose or ear, in a wound, or upon mucous membranes.

Myitis (*mi'-i-tis*). See *Myositis*.

Mylo- (*mi'-lo-*) [*μύλος*, mill]. A prefix denoting connection with the lower jaw or the molar teeth.

Mylohyoid, Mylohyoidean (*mi-lo-hi'-oid, mi-lo-hi-oid-el-an*) [*μύλη*, mill; *ὑοειδής*, hyoid]. Pertaining to the region of the lower molar teeth and the hyoid bone. **M. Muscle**. See *Muscles, Table of*.

Myo- (*mi'-o-*) [*μῦς*, a muscle]. A prefix signifying pertaining to muscle.

Myoblast (*mi'-o-blast*) [*μῦς*, muscle; *βλαστός*, germ]. A cell developing into a muscle fiber.

Myocarditis (*mi-o-kar-di'-tis*) [*μῦς*, muscle; *καρδία*, heart; *ιτις*, inflammation]. Inflammation of the myocardium, the muscular tissue of the heart.

Myocardium (*mi-o-kar'-de-um*) [*μῦς*, muscle; *καρδία*, heart]. The muscular tissue of the heart.

Myocele (*mi'-o-sel*) [*μῦς*, muscle; *κῆλη*, tumor]. The protrusion of a muscle through its ruptured sheath.

Myochrome (*mi'-o-krom*) [*μῦς*, muscle; *χρῶμα*, color]. A reddish pigment found in muscles, and believed to be identical with hemoglobin.

Myochronoscope (*mi-o-kro'-no-skōp*) [*μῦς*, muscle; *χρόνος*, time; *σκοπεῖν*, to examine]. An instrument for measuring the rapidity of muscular contraction or the propagation of a nerve-stimulus through a muscle.

Myoclonus (*mi-ok'-lo-nus*) [*μῦς*, muscle; *κλόνος*, clonus]. Clonic spasm of a muscle, or of various muscles. **M. multiplex**. See *Paramyoclonus multiplex*.

Myocomma (*mi-o-kom'-ah*) [*μῦς*, muscle; *κόμμα*, segment]. Any one of the transverse segments into which embryonic muscle-tissue is divided.

Myodes (*mi-o'-des*) [*μῦς*, muscle; *εἶδος*, like]. Myoid.

Myodesopsia (*mi-o-des-ops'-e-ah*). See *Myodesopsia*.

Myodynamometer (*mi-o-di-nam-om'-et-er*) [*μῦς*, muscle; *δύναμις*, power; *μέτρον*, measure]. See *Dynamometer*.

Myodinia (*mi-o-din'-e-ah*) [*μῦς*, muscle; *δύνη*, pain]. Pain in the muscles.

Myofibroma (*mi-o-fi-bro'-mah*) [*μῦς*, muscle; *φίβρα*, fiber; *δῆμα*, tumor]. A tumor containing muscular and fibrous tissue.

Myoglobulin (*mi-o-glob'-u-lin*) [*μῦς*, muscle; *globulin*]. A substance obtained from muscles, closely resembling serum-globulin in its properties.

Myogram (*mi'-o-gram*) [*μῦς*, muscle; *γράμμα*, an inscription]. The tracing made by the myograph.

Myograph (*mi'-o-graf*) [*μῦς*, muscle; *γράφειν*, to write]. An instrument for recording the phases of a muscular contraction.

Myographic (*mi-o-graf'-ik*) [*μῦς*, muscle; *γράφειν*, to write]. Pertaining to a myograph.

Myography (*mi-og'-ra-fe*) [*μῦς*, muscle; *γράφειν*, to write]. 1. A description of the muscles. 2. The employment of the myograph.

Myohematin (*mi-o-hem'-at-in*) [*μῦς*, muscle; *αἷμα*, blood]. A red pigment found in muscles.

Myoid (*mi'-oid*) [*μῦς*, muscle; *εἶδος*, form]. Like a muscle or muscular tissue.

Myoidema (*mi-oid-e'-mah*) [*μῦς*, muscle; *οἰδημα*, a swelling]. The wheal brought out by a sharp tap on a muscle in certain conditions of exhaustion.

Myolemma (*mi-o-lem'-ah*) [*μῦς*, muscle; *λεμμα*, covering]. The sarcolemma.

Myolin (*mi'-o-lin*) [*μῦς*, muscle]. The substance formerly supposed to form the contents of a muscular fibrilla.

Myolipoma (*mi-o-lip-o'-mah*) [*μῦς*, muscle; *λίπος*, fat; *δυμα*, tumor]. A myoma containing fatty tissue.

Myology (*mi-ol'-o-je*) [*μῦς*, muscle; *λόγος*, science]. The science of the nature, structure, functions, and diseases of muscles.

Myoma (*mi-o'-mah*) [*μῦς*, muscle; *δυμα*, tumor]. A muscular tumor. If composed of nonstriated muscular tissue it is called liomyoma; if of striated muscular tissue, rhabdomyoma.

Myomalacia (*mi-o-mal-a'-se-ah*) [*μῦς*, muscle; *μαλακία*, softening]. Softening of muscles. **M. cordis**, softening of a portion of the heart-muscle, usually resulting from thrombosis or embolism.

Myomatous (*mi-o'-mat-us*) [*μῦς*, muscle; *δυμα*, tumor]. Of the nature of a myoma.

Myomectomy (*mi-o-mek'-to-me*) [*μῦς*, muscle; *δυμα*, tumor; *ἐκτομή*, excision]. Excision of a uterine or other myoma.

Myometritis (*mi-o-met-ri'-tis*) [*μῦς*, muscle; *μητρα*, uterus; *τις*, inflammation]. Inflammation of the uterine muscular tissue.

Myomotomy (*mi-o-mot'-o-me*) [*μῦς*, muscle; *δυμα*, tumor; *τομή*, a cutting]. See *Myomectomy*.

Myoneuralgia (*mi-o-nu-ral'-je-ah*) [*μῦς*, muscle; *νεῦρον*, nerve; *ἄλγος*, pain]. Neuralgic pain in the muscles.

Myoparalysis (*mi-o-par-al'-is-is*) [*μῦς*, muscle; *παράλυσις*, paralysis]. Paralysis of a muscle or muscles.

Myopathic (*mi-o-path'-ik*) [*μῦς*, muscle; *πάθος*, disease]. Depending upon or relating to disease of the muscles. **M. Facies**, "facies myopathique;" a peculiar form of expression seen in infantile forms of myopathic muscular atrophy. It is characterized by imperfect movement of the facial muscles, sinking in of the cheeks, and drooping of the lower lip.

Myopathy (*mi-op'-ath-e*) [*μῦς*, muscle; *πάθος*, suffering]. A disease of the muscles.

Myope (*mi'-op*) [*μῦς*, to close; *ὤψ*, eye; see *Myopia*]. A person affected with myopia.

Myopericarditis (*mi-o-per-ik-ar-di'-tis*) [*μῦς*, muscle; *περί*, around; *καρδία*, heart; *τις*, inflammation]. A combination of pericarditis with myocarditis.

Myopia (*mi-o'-pe-ah*) [*μῦς*, to close; *ὤψ*, eye, myopes having the habit of partially

closing the lids to avoid spheric aberration]. Near-sightedness; an optic defect, usually due to an abnormal length of the anteroposterior diameter of the globe, whereby the focal image is formed in front of the retina.

Myopic (*mi-op'-ik*) [*μῦς*, to close; *ὤψ*, eye]. Pertaining to or having the nature of myopia; near-sighted. **M. Crescent**, a yellowish-white crescentic area about the papilla, due to atrophy or breaking away of the choroid and exposure of the sclerotic.

Myosarcoma (*mi-o-sar-ko'-mah*) [*μῦς*, muscle; *σῆρξ*, flesh; *δυμα*, tumor]. A sarcoma containing muscular tissue.

Myosclerosis (*mi-o-skle-ro'-sis*) [*μῦς*, muscle; *σκληρός*, hard]. A hardening or induration of a muscle.

Myosin (*mi'-o-sin*) [*μῦς*, muscle]. A proteid of the globulin class, found in coagulated muscle-plasma, and formed from the antecedent globulin, myosinogen. It is also found in the cornea and in some vegetables. **M.** is soluble in strong saline solutions, and is changed into syntonin by the action of dilute hydrochloric acid and heat.

Myosinogen (*mi-o-sin'-o-je*) [*μῦς*, muscle; *γεννᾶν*, to beget]. One of the proteids of muscle-plasma. It is the antecedent of myosin. Its coagulation after death is the cause of rigor mortis.

Myosis (*mi-o'-sis*) [*μῦς*, to close]. See *Miosis*.

Myositis (*mi-o-si'-tis*) [*μῦς*, muscle; *τις*, inflammation]. Inflammation of the muscles.

M., **Interstitial**, an inflammation of the connective tissue of muscle. **M. ossificans**, a variety characterized by the formation of osseous deposits in the muscles. **M.**, **Specific**, **M.**, **Syphilitic**, that due to syphilis. **M. trichinosa**, that due to the presence of trichina spiralis in the muscles.

Myospasm (*mi'-o-spazm*) [*μῦς*, muscle; *σπασμός*, spasm]. Muscular spasm; a cramp.

Myosuture (*mi-o-su'-tūr*) [*μῦς*, muscle; *sutura*, a suture]. Suture of a muscle.

Myotatic (*mi-o-tat'-ik*) [*μῦς*, muscle; *τάσσειν*, a stretching]. Produced by or depending upon passive tension of the muscles. **M. Irritability**, an increased irritability of muscles produced by passive extension, and giving rise, when the muscle is stimulated, to a **M. Contraction**, or tendon reflex.

Myotic (*mi-o'-ik*). See *Miotic*.

Myotome (*mi'-o-tōm*) [*μῦς*, muscle; *τομή*, a cutting]. 1. An instrument for performing myotomy. 2. See *Myocomma*.

Myotomy (*mi-ol'-o-me*) [*μῦς*, muscle; *τομή*, a cutting]. 1. Division of a muscle, particularly through its belly. 2. The dissection of muscles.

Myotonia (*mi-o-to'-ne-ah*) [*μῦς*, muscle; *τόνος*, tension]. 1. Muscular spasm. 2.

The stretching of a muscle. **M. congenita**, **M. hereditaria**, Thomsen's disease; an hereditary disease characterized by tonic spasms in the voluntarily moved muscles.

Myrcia (*mur'-se-ah*). A genus of aromatic shrubs of the natural order Myrtaceæ. **M. acris** is the source of oil of bay, *Oleum myrciæ* (U. S. P.), from which is prepared bay-rum, or *Spiritus myrciæ* (U. S. P.). Bay-rum is used as a local application in headache.

Myriachit (*mir'-e-ah-it*). See *Palmas*.

Myricin (*mir'-is-in*) [*μυρίκη*, the tamarisk]. A constituent of wax (*Cera flava*), insoluble in boiling alcohol. It consists chiefly of myricyl palmitate, $C_{50}H_{61} \cdot C_{16}H_{31}O_2$, which is a compound of palmitic acid and myricyl alcohol.

Myricyl (*mir'-is-il*). See *Myricin*.

Myringitis (*mir-in-jit'-tis*) [*myringa*, drum-membrane; *itis*, inflammation]. Inflammation of the membrana tympani.

Myringodectomy (*mir-in-go-dekt'-to-me*) [*myringa*, drum-membrane; *ἐκτομή*, excision]. Excision of a part or of the whole of the membrana tympani.

Myringomycosis (*mir-in-go-mi-ko'-sis*) [*myringa*, drum-membrane; *μύκωσις*, fungus]. Disease of the drum-membrane, due to parasitic fungi, especially the *Aspergillus*.

Myringoplastic (*mir-in-go-plas-tik*) [*myringa*, drum-membrane; *πλαστικόν*, to form]. Pertaining to myringoplasty.

Myringoplasty (*mir-in-go-plas-te*) [*myringa*, drum-membrane; *πλαστικόν*, to shape]. A plastic operation on the membrana tympani.

Myringotome (*mir-in-go-tōm*) [*myringa*, drum-membrane; *τομή*, a cutting]. An instrument used in incising the membrana tympani.

Myringotomy (*mir-in-got'-o-me*) [*myringa*, drum-membrane; *τομή*, a cutting]. Incision of the tympanic membrane.

Myristic Acid (*mir'-is'-tik*) [*μυρίστις*, to anoint]. $C_{14}H_{26}O_2$. A monobasic acid found in *Myristica fragrans*, in cocoanut-oil, in spermaceti, etc.

Myristica (*mir'-is'-tik-ah*) [*μυρίστις*, to anoint]. See *Nutmeg*.

Myristicin (*mir'-is'-tis-in*) [*μυρίστις*, to anoint]. A peculiar fatty body contained in nutmeg.

Myristicol (*mir'-is'-tik-el*) [*μυρίστις*, to anoint; *oleum*, oil]. $C_{19}H_{32}O$. An oily substance extracted from oil of nutmeg.

Myronic Acid (*mi-ron'-ik*). See *Acid*.

Myrosin (*mi'-ro-sin*) [*μυρσίνη*, ointment]. An albuminous ferment occurring in mustard-seed, which liberates the oil of mustard from potassium myronate.

Myrrh (*mur*) [*μύρρα*, myrrh]. A gum-resin obtained from *Commiphora myrrha* (U. S. P.) or *Balsamodendron myrrha* (B. P.), a tree of

the natural order Burseraceæ. Myrrh contains a volatile oil, a resin (myrrhin), and a gum, and is used as a stimulant tonic in dyspepsia, pulmonary affections, chlorosis, and amenorrhea. It is employed as a local application in various forms of stomatitis. Dose gr. x-xxx (0.65-2.0). **Tinctura myrrhæ**. Dose m x-xxx (0.65-2.0). Myrrh is also an ingredient of *Mistura ferri composita*, *Pilulæ aloes et myrrhæ*, and *Tinctura aloes et myrrhæ*.

Myrtiform (*mur'-tif-orm*) [*μύρτιος*, the myrtle; *forma*, form]. Shaped like a myrtle-berry or myrtle-leaf, as the myrtiform caruncles (caruncule myrtiformes), and the myrtiform fossa of the superior maxilla.

Myrtle (*mur'-tl*) [*μύρτος*, the myrtle]. A genus of plants of the natural order Myrtaceæ. The leaves of *Myrtus chekan* are used in bronchitis. Dose of fluid extract, f 3 j-ij (4.0-12.0). The leaves of *Myrtus communis* yield myrtol, which distils between 160° and 180° C. The leaves are used as an antiseptic dressing for wounds. Myrtol is employed as an antiseptic in bronchitis, cystitis, and pyelitis. Dose gr. j-ij (0.065-0.13).

Myrtol (*mur'-tol*). See *Myrtle*.

Myrtus (*mur'-tus*). See *Myrtle*.

Mysophobia (*mi-so-fo'-be-ah*) [*μύσος*, filth; *φόβος*, dread]. An abnormal dread of contact or of dirt.

Mytacism (*mit'-as-izm*). See *Mutacism*.

Mytilotoxin (*mit-il-o-toks'-in*) [*μυτίλος*, mussel; *τοξικόν*, poison], $C_6H_{15}NO_2$. A poisonous ptomain found in poisonous mussel; it is similar in action to curara. See *Ptomain*, Table of.

Myxedema (*miks-e-de'-mah*) [*μύξα*, mucus; *οίδημα*, edema]. A disorder of nutrition in which the subcutaneous tissue, especially of the face and hands, becomes infiltrated with a mucin-like substance, giving rise to a pale, edematous swelling, which does not pit on pressure. It is associated with dullness of the intellect, slow monotonous speech, muscular weakness, tremors, and absence of sweating. It is thought to be due to atrophy or degeneration of the thyroid gland, a view that gains strength from the facts that a similar condition to that of myxedema is produced in animals by the removal of the thyroid gland (cachexia strumipriva), and that the administration of thyroid gland in myxedema produces a marked amelioration of symptoms. In individuals in whom the thyroid gland is congenitally diseased or absent, a state resembling myxedema is apt to develop. It is known as cretinism.

Myxedematous (*miks-e-dem'-at-us*) [*μύξα*, mucus; *οίδημα*, edema]. Pertaining to, affected with, or of the nature of, myxedema.

Myxochondroma (*miks-o-kon-dro'-mah*) [*μύξα*, mucus; *χόνδρος*, cartilage; *δμα*, tumor]. A tumor containing mucous and cartilaginous tissue.

Myxodes (*miks-o'-dez*) [*μύξα*, mucus; *ειδος*, like]. Resembling mucus.

Myxofibroma (*miks-o-fi-bro'-mah*) [*μύξα*, mucus; *fibra*, fiber; *δμα*, tumor]. A fibroma that has in part undergone myxomatous degeneration.

Myxoglioma (*miks-o-gli-o'-mah*) [*μύξα*, mucus; *γλία*, glue; *δμα*, tumor]. A glioma with myxomatous degeneration.

Myxolipoma (*miks-o-lip-o'-mah*) [*μύξα*, mucus; *λίπος*, fat; *δμα*, tumor]. A fatty tumor that has in part undergone myxomatous change.

Myxoma (*miks-o'-mah*) [*μύξα*, mucus; *δμα*, tumor]. A connective-tissue tumor after the type of the jelly of Wharton of the umbilical cord. It consists of a gelatinous, mucin-con-

taining, intercellular substance, in which are scattered peculiar branched or stellate cells.

Myxomatous (*miks-o'-mat-us*) [*μύξα*, mucus; *δμα*, tumor]. Of the nature of, or pertaining to, myxoma. **M.**, Degeneration, mucoid degeneration. It attacks epithelium and connective tissue, giving rise to the production of a gelatinous substance containing mucin. In epithelial tissue the cells are affected; in connective tissue the intercellular substance.

Myxosarcoma (*miks-o-sar-ko'-mah*) [*μύξα*, mucus; *σάρξ*, flesh; *δμα*, tumor]. A sarcoma that has in part undergone myxomatous degeneration.

Myxosporidia (*miks-o-spor-id'-e-ah*) [*μύξα*, mucus; *σπόρος*, seed]. A group or order of Sporozoa, found as parasitic bodies in the muscles and epithelia of fishes; they produce a psorospermiosis which is frequently wide-spread and destructive.

N

N. The symbol of nitrogen; **n**, the sign of the index of refraction.

N. A. Abbreviation for numerical aperture.

N. F. National Formulary; a collection of formulas issued by the American Pharmaceutical Association as a supplement to the U. S. Pharmacopeia.

Na. Symbol for sodium.

Nævus (*ne'-vus*). See *Nevus*.

Nail (*nal*) [*AS.*, *naegel*, a nail]. The horny structure covering the dorsal aspect of the terminal phalanx of each finger and toe. It consists of intimately united horny epithelial cells derived from the stratum lucidum of the epidermis. A nail rests in the **N.-bed**, a vascular tissue corresponding to the corium and the stratum Malpighii of the skin. **N.-culture**, a term applied in bacteriology to a stab-culture showing a growth along the needle-track, and on the surface a button-like projection, giving the appearance of a nail driven into the gelatin. **N.-fold**, the portion of epidermis that covers the root and edges of the nail. **N., Hang-**, a little shred of skin hanging from the finger at the side of the nail. **N., Ingrowing or Ingrown**, an overlapping of the nail by the flesh, from pressure, attended with ulceration. **N.-matrix**, the proximal end of the *N.-bed*; the structure from which the nail grows. **N., Turtle-back**, a nail curved in all directions;

a condition seen in certain trophic disturbances.

Nanism (*na'-nizm*) [*vānos*, dwarf]. Dwarfishness.

Nanocephalous (*na-no-sef'-al-us*) [*vānos*, dwarf; *κεφαλή*, head]. Possessing a dwarfed head.

Nanoid (*na'-noid*) [*vānos*, dwarf; *ειδος*, like]. Dwarf-like.

Nanomelia, Nanomelus (*na-no-me'-le-ah, na-nom'-el-us*) [*vānos*, dwarf; *μέλος*, limb]. A monster characterized by undersized limbs.

Nanus (*na'-nus*) [*vānos*, a dwarf]. 1. A dwarf. 2. Dwarfed, stunted.

Nape (*nāp*) [*ME.*, *knappe*, from *AS.*, *cnap*, a knob]. The back part of the neck; the nucha.

Napellin (*na-pe'll-in*) [*napellus*, dim. of *napius*, a turnip], $C_{10}H_8NO_{11}$. An alkaloid of *Aconitum napellus*. It is an anodyne and antineuralgic. Dose, gr. $\frac{1}{6}$ - $\frac{1}{2}$.

Naphtha (*naf'-thah*) [*vāpha*, from *Arab.*, *naft*, naphtha]. 1. Formerly, any strong-smelling, inflammable, volatile liquid, as naphtha vitrioli (sulphuric ether), wood-naphtha (methyl-alcohol). 2. A colorless, inflammable oil distilled from petroleum, bituminous shale, etc.

Naphthalene, Naphthalin, Naphthalin (*naf'-thal-en, naf'-thal-in, naf'-tal-in*)

[*νάφα*, from Arab., *naft*, naphtha], $C_{10}H_8$. A hydrocarbon crystallizing in large, silvery, rhombic plates, slightly soluble in hot, but insoluble in cold water, though easily soluble in methyl and ethyl alcohols, chloroform, ether, and benzene. It is an antiseptic; is used in intestinal putrefaction, in typhoid fever, etc.; locally, in scabies and pruritus. Dose, gr. v-x (0.32-0.65).

Naphthalol (*naft'-thal-ol*) [*naphtha*], $C_{10}H_7OH.COOC_{10}H_7$. A salicylic-acid derivative of β -naphthol.

Naphthol, **Naphtol** (*naft'-thol*, *naft'-tol*), $C_{10}H_7OH$. A substance found in coal-tar and prepared artificially from naphthalin. It exists in two isomeric forms, α -naphthol and β -naphthol, and occurs in the form of pale, buff-colored crystals. β -naphthol (Naphtol, U. S. P.) is employed in dyspepsia, and as an intestinal antiseptic in diarrheas, typhoid fever, etc. Locally, it is used in eczema, prurigo, herpes, favus, etc. Dose, gr. v-x (0.32-0.65).

Narcein (*nar'-se-in*) [*νάρκη*, numbness], $C_{23}H_{29}NO_9$. An alkaloid contained in opium. It is sparingly soluble in water and alcohol, and forms fine, silky, inodorous, bitter crystals. It is used as a substitute for morphin. Dose, gr. $\frac{1}{4}$ (0.016).

Narcophyria (*nar-ko-hip'-ne-ah*) [*νάρκη*, stupor; *ἵπνος*, sleep]. Waking numbness; a peculiar state in which the patient has a sense of numbness on awaking.

Narcolepsy (*nar'-ko-lep-se*) [*νάρκη*, stupor; *καταβαίνει*, to seize]. An uncontrollable tendency to attacks of deep sleep of short duration. It has been observed in epilepsy and other affections.

Narcosis (*nar-ko'-sis*) [*ναρκόω*, to benumb]. The state of complete unconsciousness produced by a narcotic drug or an anesthetic.

Narcotic (*nar-ko'-ik*) [*ναρκόω*, to benumb]. 1. Producing stupor. 2. A drug that produces narcosis.

Narcotin (*nar'-kol-in*) [*ναρκόω*, to benumb], $C_{22}H_{23}NO_7$. An alkaloid of opium, separated from morphin by potassium hydroxid. It crystallizes from alcohol in shining prisms, and melts at $176^{\circ}C$. It is sudorific and antipyretic, but has no narcotic effects. Dose, gr. i-iii (0.06-0.20).

Narcotism (*nar'-kol-izm*) [*ναρκόω*, to benumb]. The condition resulting from the use of a narcotic.

Narcotize (*nar'-ko-tiz*) [*ναρκόω*, to benumb]. To put under the influence of a narcotic; to render unconscious by means of a narcotic.

Naris (*na'-ris*) [L.]. One of a pair of openings at the anterior part (**Anterior nares**), or at the posterior part (**Posterior nares**) of the nasal fossæ.

Nasal (*na'-zal*) [*nasus*, the nose]. Pertain-

ing to the nose. **N. Artery**. See *Arteries*, *Table of*. **N. Bones**, the two small bones forming the arch of the nose. **N. Catarrh**, catarrh of the nasal mucous membrane; coryza. **N. Duct**. See *Lacrimal Duct*. **N. Eminence**. See *Eminence*. **N. Fossæ**, the cavities of the nose. **N. Ganglion**, the sphenopalatine ganglion. **N. Labyrinth**, the irregular cavity formed by the turbinal bones in the nasal passages. **N. Line**. See *Lines*, *Table of*. **N. Nerve**. See *Nerves*, *Table of*. **N. Voice**, a peculiar muffled timbre of the voice, especially marked in cases of perforation of the palate.

Nascent (*nas'-ent*) [*nasci*, to be born]. A term applied to gaseous substances at the moment of their liberation from chemic union.

Nasion (*na'-se-on*) [*nasus*, nose]. The median point of the nasofrontal suture.

Nasmyth's Membrane. The epithelial membrane enveloping the enamel of the tooth during its development and for a short time after birth; cuticula dentis.

Naso- (*na'-zo-*) [*nasus*, nose]. A prefix denoting connection with or relation to the nose.

Nasofrontal (*na-zo-fron'-tal*) [*nasus*, nose; *frons*, forehead]. Pertaining to the nasal and the frontal bones.

Nasolabial (*na-zo-lab'-e-al*) [*nasus*, nose; *labium*, lip]. Pertaining to the nose and lip.

Nasolacrimal (*na-zo-lak'-rim-al*) [*nasus*, nose; *lacrima*, tear]. Pertaining to the nose and the lacrimal apparatus, as the nasolacrimal duct.

Nasopalatine (*na-zo-pal'-at-en*) [*nasus*, nose; *palatine*]. 1. Pertaining to both the nose and the palate, as the N. nerve. 2. Giving passage to the N. nerve, as the N. canal.

Nasopharyngeal (*na-zo-far-in'-je-al*) [*nasus*, nose; *φάρυγξ*, pharynx]. Pertaining to both the nose and the pharynx, or to the nasopharynx.

Nasopharynx (*na-zo-far'-inkz*) [*nasus*, nose; *φάρυγξ*, pharynx]. The space between the posterior nares and a horizontal plane through the lower margin of the soft palate.

Nataloin (*na-tal'-o-in*) [*Natal*, an African state; *ἀλόη*, the aloë], $C_{25}H_{25}O_{11}$. The aloin derived from Natal aloes.

Natal Sore. See *Furunculus orientalis*.

Nates (*na'-tez*) [*natis*, the buttock]. The buttocks. **N. of Brain**, the anterior pair of the corpora quadrigemina.

Native (*na'-tiv*) [*nativus*, born]. Of indigenous origin or growth; occurring in its natural state; not artificial. **N. Albumins**, a class of proteids occurring ready-formed in the tissues.

Natrium (*na'-tre-um*). See *Sodium*.

Natron (*na'-trun*) [Ar., *natrūn*, native car-

bonate of sodium]. 1. Native sodium carbonate. 2. Soda. 3. Sodium or potassium nitrate.

Natural (*nat'-u-ral*) [*natura*, nature]. Pertaining to nature. **N.** History, a term including a description of all the products and phenomena of nature, but at present generally restricted to the sciences of zoology and botany. **N.** Philosophy, the science treating of the physical properties of matter at rest and in motion; now usually called physics.

Nauheim Treatment (*now'-hem*). See *Schott's Treatment of Heart-disease*.

Nausea (*now'-she-ah*) [*nausia*, seasickness]. Sickening at the stomach, with inclination to vomit. **N. marina**, **N. navalis**, seasickness.

Nauseant (*now'-she-ant*) [*nausia*, seasickness]. 1. Nauseating; producing nausea. 2. Any agent that produces nausea.

Nauseating, **Nauseous** (*now'-she-a-ting*, *now'-she-us*) [*nausia*, seasickness]. Producing nausea or loathing.

Navel (*na'-vel*) [*AS.*, *nafela*, navel]. The umbilicus. **N.-string**, the umbilical cord.

Navicular (*na-vik'-u-lar*) [*navicula*, a little ship]. Boat shaped. **N. Bone**, the scaphoid bone of the foot. **N. Fossa**. 1. A depression between the vaginal aperture and the fourchet. 2. A dilatation of the urethra near the glans penis.

Near-point. The *punctum proximum*, the point nearest the eye at which an object can be seen distinctly. **N.** Absolute, that near-point for either eye alone at which no effort at accommodation is made. **N.** Relative, that near-point for both eyes at which accommodation is brought into play.

Near-sight. See *Myopia*.

Nearthrosis (*ne-ar-thro'-sis*) [*νῆος*, new; *ἄρθρον*, joint]. A false articulation.

Nebula (*neb'-u-lah*) [*L.*]. A cloud; a faint, grayish opacity of the cornea.

Nebulizer (*neb'-u-li-zer*) [*nebula*, a cloud]. See *Atomizer*.

Neck (*nek*) [*AS.*, *hnecca*, neck]. The constricted portion of the body connecting the head with the trunk; also, the narrow portion of any structure serving to join its parts. See *Cervix*. **N.** Anatomic, the constricted portion of the humerus, just below the articular surface, serving for the attachment of the capsular ligament. **N.-berg**, the vulgar name of the cervical portion of the thymus gland of animals. See *Heart-berg*. **N.** Derbyshire. Synonym of *Gaiter*. **N. Surgical**, the constricted part of the humerus just below the condyles. **N.** Wry-. Synonym of *Torticollis*.

Necrobiosis (*nek-ro-bi-o'-sis*) [*νεκρός*, corpse; *βίος*, life]. Molecular death of tissue.

Necrogenic (*nek-ro-jen'-ik*) [*νεκρός*, corpse;

γεννῆν, to beget]. Originating from dead substances.

Necrometer (*nek-rom'-et-er*) [*νεκρός*, corpse; *μέτρον*, measure]. An instrument for weighing organs at an autopsy.

Necropsy (*nek'-rop-se*) [*νεκρός*, corpse; *δύσις*, seeing]. The examination of a dead body; autopsy; postmortem.

Necrosis (*nek-ro'-sis*) [*νεκρός*, corpse]. The death of cells surrounded by living tissue. Necrosis proper refers to death in mass, necrobiosis to death of individual cells. Among the causes of necrosis are: direct injury, obstruction of the circulation, loss of trophic influence. The varieties of necrosis are: coagulative necrosis, liquefactive necrosis, cheesy necrosis, dry, and moist gangrene. Necrosed tissues may be absorbed, retained, or thrown off. The dead tissue is called sequestrum in case of bone, and sphacelus in case of soft parts. In surgery the term necrosis is often applied specifically to the death of bone. Necrosis of bone is divided into **N.** Superficial, when the portion of bone just beneath the periosteum is affected; **N.** Central, when the internal portions are involved, and **N.** Total, when the bone through its entire thickness is necrotic. **N.** Balser's Fatty. See *Diseases*, *Table of*.

N. Cheesy, necrosis characterized by the formation of a cheese-like material. It is usually due to the presence of the tubercle bacillus, and is most often seen in the lungs. **N.** Coagulation-, **N.** Coagulative, a variety characterized by the formation of fibrin. It is most common in the blood, in thrombosis. It also occurs on mucous membranes, where it produces a false membrane, chiefly under the influence of the bacillus of diphtheria, but also of other irritant agents. Occasionally it occurs in the interior of organs as the result of embolism or thrombosis, leading to the formation of an anemic infarct. **N.** Colliquative. See *N.* Liquefactive. **N.** Embolic, coagulative necrosis in an anemic infarct following embolism. **N.** Liquefactive, a process analogous to coagulative necrosis, but instead of fibrin the peculiar reaction of fluids and cells gives rise to the formation of a liquid. **N.** Mercurial, a necrosis of bones due to chronic poisoning with mercury. **N.** Phosphorus-, a necrosis of bone, especially of the lower jaw, occurring in those exposed to the fumes of phosphorus. **N.** ustilaginea, dry gangrene from ergotism.

Necrotic (*nek-ro'-ik*) [*νεκρός*, corpse]. Pertaining to or characterized by necrosis.

Necrotomy (*nek-ro'-o-me*) [*νεκρός*, a corpse; *τομή*, a cutting]. The dissection of a dead body.

Nectandra (*nek-tan'-drah*) [*νεκταρ*, nectar;

ἀνήρ, a male]. A tree of the order Lauraceae. The bark of *Nectandra rodiei* (Nectandra cortex, B.P.), contains tannic acid, resin, sugar, albumin, various salts, and two alkaloids, beberin and sipirin. It is tonic, astringent, and febrifuge, and has been used in malarial fevers, but is much less antiperiodic than cinchona. Dose, gr. xx-3], (1.3-4.0); of beberin sulphate, gr. ij-v (0.13-0.32).

Needle (*ne'-dl*) [AS., *nađl*, needle]. A sharp-pointed steel instrument used for sewing, and for penetrating tissues for the purpose of carrying a ligature through. **N.**, **Aneurysm**-, one fixed on a handle, and with the eye at the point, especially adapted for ligating vessels. **N.**, **Cataract**-, one for operating upon the cataractous lens or its capsule. **N.**, **Discission**-, one for insertion through the cornea, and breaking the capsule and substance of the crystalline lens. **N.**, **Exploring**, a grooved sharp-pointed rod introduced into a cavity or a part for the purpose of determining the presence of fluid. **N.**, **Hagedorn's**, a flat suture-needle curved on its edge, with the eye perforating the side. **N.**-**holder**, a handle for claspings a needle. **N.**, **Hypodermic**, the fine needle-pointed metallic tube attached to the barrel of the hypodermic syringe. **N.**-**knife**, one that has a sharp cutting edge; it is used in the discission of cataracts.

Needling (*ne'-dlng*) [AS., *nađl*, needle]. The process of lacerating a cataract with a needle, to afford entrance to the aqueous humor and cause absorption of the lens.

Negative (*neg'-at-iv*) [*negare*, to deny]. Denying; contradicting; opposing. Of quantities, less than nothing; in physics, opposed to a quality termed positive. **N.** **Accommodation**, the absence of active accommodation; the state of the eye at rest, or when looking at an object at an infinite distance. **N.** **Blood-pressure**, pressure which is less than that of the atmosphere. It exists in the large veins near the heart, owing to the aspirating action of the thorax. **N.** **Chemotaxis**, the absence of the power of attracting leukocytes and wandering cells, or their actual repulsion. **N.** **Electrode**, **N.** **Pole**, the electrode or pole of an electric battery connected with the positive plate and conducting N. electricity. **N.** **Variation of the Muscle-current**, a diminution in the strength of the muscle-current during tetanic contraction.

Neisseria (*ni-se'-re-ah*) [after *Neisser*, a German physician]. A group of schizomycetes. See *Bacteria*, Table of.

Nélaton [*Nélaton*, a French surgeon]. **N.'s Catheter**, a soft rubber catheter. **N.'s Line**. See *Lines*, Table of. **N.'s Probe**. See *Probe*.

Nelavan (*nel'-ev-an*). See *Lathargy*, *African*.

Nematoblast (*nem'-at-o-blast*) [*nēma*, thread; *βλαστός*, germ]. A spermatoblast.

Nematoda (*nem-at-o'-dah*) [*nēma*, thread; *ειδός*, form]. A genus of worms, the thread-worms, some of which are parasitic in man and the lower animals. The most important of these are *Anguillula*, *Ankylostoma*, *Ascaris*, *Eustrongylus*, *Filaria*, *Oxyuris*, *Strongylus*, *Trichina*, *Trichocephalus*.

Nematode, **Nematoid** (*nem'-at-od*, *nem'-at-oid*) [*nēma*, thread; *ειδός*, form]. Thread-like; belonging to or resembling the Nematoda, or thread-worms; applied to thread-worms, hair-worms, round worms, pin-worms.

Neoarthritis (*ne-o-ar-thro'-sis*) [*νέος*, new; *άρθρον*, a joint]. A false joint.

Neonatal (*ne-o-na'-tal*) [*νέος*, new; *natus*, born]. Pertaining to the new-born.

Neoplasm (*ne'-o-plasm*) [*νέος*, new; *πλάσσειν*, to mold]. A circumscribed new growth of tissue, characterized by abnormality of structure or location. As generally used, the term includes all true tumors, as well as tumor-like growths due to microorganisms, as the gumma and tuberculous tumor. **N.**, **Inflammatory Fungoid**, mycosis fungoides.

Neoplastic (*ne-o-plas'-tik*) [*νέος*, new; *πλάσσειν*, to mold]. Pertaining to, or of the nature of a neoplasm.

Neoplasty (*ne'-o-plas-te*) [*νέος*, new; *πλάσσειν*, to mold]. The restoration of lost tissue by a plastic operation.

Nepeta (*nept'-et-ah*). See *Catnip*.

Nephralgia (*nef-rall'-je-ah*) [*νεφρός*, kidney; *ἄλγος*, a pain]. Neuralgic pain of the kidney.

Nephrectomy (*nef-rek'-to-me*) [*νεφρός*, kidney; *ἐκτομή*, excision]. Excision of the kidney. **N.** **Abdominal**, **N.** through an abdominal incision. **N.**, **Lumbar**, **N.** through an incision in the loin.

Nephria (*neft'-re-ah*). See *Bright's Disease*.

Nephric (*neft'-rik*) [*νεφρός*, kidney]. Pertaining to the kidney.

Nephritic (*neft-ril'-ik*) [from *νεφρός*, kidney; *ιτις*, inflammation]. 1. Pertaining to nephritis. 2. Improperly, pertaining to the kidney; the correct term is nephric.

Nephritis (*neft-ril'-tis*) [*νεφρός*, the kidney; *ιτις*, inflammation]. Inflammation of the kidney. The inflammation when affecting the epithelium is termed **Parenchymatous N.**, which may be acute or chronic. It is also known as **Tubular**, **Catarrhal**, or **Desquamative N.**, or, when affecting especially the glomerule, as **glomerulonephritis**. That involving the connective tissue chiefly is called **Interstitial N.**, and may also be acute or chronic. The acute gives rise to abscess

of the kidney. A nephritis in which both elements are involved is termed **Diffuse N.** In **Acute Parenchymatous N.**, or acute Bright's disease, the kidney is enlarged, congested, its structural markings are obscured, the epithelium is in a state of cloudy swelling or fatty degeneration, and many tubules contain casts; in others the epithelium is desquamated. The symptoms are fever, lumbar pains, and often dropsy; the urine is scanty and contains albumin, blood, casts, and epithelial cells; uremia is common. In **Chronic Parenchymatous N.** (large white kidney) the kidney is enlarged, pale or yellow, and soft; the epithelium presents an advanced stage of fatty degeneration. Casts are often present. The symptoms are anemia, dropsy, gastrointestinal disturbances, dyspnea, and retinitis; the urine is scanty and contains a large amount of albumin, granular, fatty, and waxy casts, and compound granule-cells. **Acute interstitial N.** is due to septic infection either through the blood, as in pyemia, or through extension along the ureter or from neighboring structures. In **chronic interstitial N.** (granular or gouty kidney), the kidney is small, hard, the capsule is adherent, the surface is granular and marked by cysts. The cortex is diminished in thickness. Microscopically there is a marked increase in the connective tissue about the tubules, the glomeruli, and the blood-vessels. The glomeruli show degenerative changes, the epithelium is atrophic and degenerated. The symptoms are anemia, loss of flesh and strength, disturbances of digestion, dyspnea, and asthmatic attacks; polyuria, the urine being of a low specific gravity, pale, and containing only a small amount of albumin and but few hyaline casts. There is also hypertrophy of the heart and sclerosis of the arteries; the pulse is of high tension, and the second aortic sound is accentuated. Uremia is common. **N., Saturnine**, that due to chronic lead-poisoning. **N., Scarlatinal**, an acute nephritis due to scarlet fever. **N., Tuberculous**, that due to the presence of tubercle-bacilli. It presents itself either in the form of caseating masses or cavities in the substance of the kidney, or as miliary tuberculosis of the organ.

Nephrogenous (*nef-roj'-en-us*) [*νεφρός*, kidney; *γεννάν*, to beget]. Originating in the kidney; of renal origin.

Nephrolith (*nef'-ro-lith*) [*νεφρός*, kidney; *λίθος*, stone]. A calculus of the kidney.

Nephrolithiasis (*nef-ro-lith-i'-as-is*) [*νεφρός*, kidney; *λίθος*, stone]. The formation of renal calculi, or the diseased state that leads to their formation.

Nephrolithotomy (*nef-ro-lith-ot'-o-me*)

[*νεφρός*, kidney; *λίθος*, stone; *τομή*, a cutting]. An incision of the kidney for the removal of a calculus.

Nephropoysis (*nef-ro-pi-o'-sis*). Same as *Pyonephrosis*.

Nephrorrhagia (*nef-ror-a'-je-ah*) [*νεφρός*, kidney; *ρρύναι*, to burst forth]. Renal hemorrhage.

Nephrorrhaphy (*nef-ror'-a-fe*) [*νεφρός*, kidney; *ῥαφή*, suture]. The stitching of a floating kidney to the posterior wall of the abdomen or to the loin.

Nephrotomy (*nef-rot'-o-me*) [*νεφρός*, kidney; *τομή*, a cutting]. Incision of the kidney.

N., Abdominal, one through an abdominal incision. **N., Lumbar**, one through an incision in the loin.

Nephrotiphoid, Nephrotiphus (*nef-ro-ti'-foid, nef-ro-ti'-fus*) [*νεφρός*, kidney; *τίφος*, stupor]. Enteric fever with prominent renal complications.

Neroli (*ner'-o-le*) [Fr.]. Oil of orange-flowers.

Nerve (*nurv*) [*nervus*, nerve]. An elongated, cord-like structure made up of aggregations of N.-fibers and having the property of transmitting nervous impulses. **N., Accelerator**, the cardiac sympathetic N., stimulation of which causes acceleration of the heart's action. **N., Afferent**, one that transmits impulses from the periphery to the central nervous system. **N., Calorific**, a N., stimulation of which increases the heat of the parts to which it is distributed. The term is applied to the dilator nerves and to those stimulating functional activity of organs.

N.-cell, a mass of protoplasm containing a large vesicular nucleus within which lies a well-marked nucleolus. N.-cells have one or more elongated processes, and in accordance with the number of these are designated unipolar, bipolar, or multipolar. The processes are of two kinds: the axis-cylinder process and the protoplasmic processes. The axis-cylinder (Deiter's) process either becomes an axis-cylinder of a N.-fiber, or divides within the gray matter into delicate filaments; it gives off minute branches termed collaterals; the other processes are supposed to have nutritive and conducting functions. **N.-center**, a group of ganglion-cells acting together in the performance of a function. **N., Centrifugal**. See *N., Efferent*. **N., Centripetal**. See *N., Afferent*. **N.-corpuscles**, nucleated corpuscles lying between the neurilemma and the medullary sheath of medullated N.-fibers. **N., Cranial**, a N. arising directly from the brain and making its exit through one of the foramina in the skull. **N., Depressor**, an afferent N., irritation of which depresses or inhibits the vasomotor center. **N., Efferent**,

one carrying impulses from the central nervous system to the periphery. **N.-endings**, the terminations of nerves at the periphery or in the nerve-centers. **N. of Expression**, the facial nerve. **N.-fiber**, a fiber having the property of conducting invisible or molecular waves of stimulation from one part of an organism to another, and so establishing physiologic continuity between such parts without the necessary passage of waves of contraction (Romanes). There are two kinds of nerve-fibers: the medullated, or myelinic, and the nonmedullated, or amyelinic. A typical medullated fiber consists of the axis-cylinder, which may be surrounded by a sheath, the axilemma; the medullary sheath, or white substance of Schwann; the neurilemma, or sheath of Schwann. The axis-cylinder is made up of fine fibrillæ cemented together by a granular interstitial substance, the neuroplasm. The medullary sheath consists of a reticulated framework composed of a material resembling neurokeratin, and a semifluid substance filling the interstices, called myelin. At regular intervals the medullated sheath is marked by interruptions, the nodes of Ranvier. Each internode, as the portions of fiber included between two Ranvier's nodes are termed, possesses a nerve-corpuscle. The neurilemma, sheath of Schwann, or primitive sheath, is a delicate elastic membrane, closely investing the medullary substance. On its inner surface are placed the nerve-corpuscles. The nonmedullated, pale, or Remak's fibers do not possess a medullary sheath, but consist only of axis-cylinder and neurilemma. The nerve-corpuscles are more abundant than in medullated nerve-fibers. Medullated nerve-fibers are found in the cerebrospinal nerves, while nonmedullated fibers occur in the sympathetic nerves and tend to form plexuses. Nerve-fibers are aggregated into fasciculi; these in turn unite to form nerve-trunks. The nerve-fibers in a fasciculus are held

together by the endoneurium; while each fasciculus is surrounded by a sheath termed the perineurium. The epineurium holds together the several fasciculi composing a nerve-trunk. **N., Frigorific**, a sympathetic nerve, stimulation of which causes a fall of temperature; the vasoconstrictor nerves are frigoric nerves. **N.-grafting**, the transplanting of a portion of healthy nerve from an animal to man, to reestablish the continuity of a divided nerve; neuroplasty. **N.-head**, the optic disc or papilla. **N., Inhibitory**, one the stimulation of which inhibits or lessens the activity of an organ. **N., Motor**, one containing only or chiefly motor fibers. **N.-plexus**, a grouping of nerves. **N., Pressor**, an afferent nerve, irritation of which stimulates the vasomotor center. **N., Secretary**, an efferent nerve, stimulation of which causes increased activity of the gland to which it is distributed. **N., Spinal**, one of those arising from the spinal cord and making its exit through an intervertebral foramen. There are 31 pairs of spinal nerves. **N.-storm**, a sudden outburst or paroxysm of nervous disturbance. **N.-stretching**, mechanic elongation or tension of a nerve, for the relief of neuralgia, spasmodic contraction, and other pathologic conditions. **N., Sympathetic**, one of a system of nerves distributed chiefly to the blood-vessels and to the viscera. See *Sympathetic*. **N., Thermic**. A synonym of *N., Calorific*. **N.-tire**, neurasthenia. **N., Trophic**, a nerve, the function of which is to preside over the nutrition of the part to which it is distributed. **N., Vasoconstrictor**. See *N., Vasomotor*. **N., Vasodilator**. See *N., Vasomotor*. **N., Vasomotor**, any one of the nerves controlling the caliber of the blood-vessels; they are of two kinds—those stimulation of which causes contraction of the vessels, vasoconstrictor nerves, and those stimulation of which causes active dilatation, vasodilator nerves. Ordinarily vasomotor is synonymous with vasoconstrictor.

TABLE OF NERVES (ALPHABETIC).

NAME.	FUNCTION.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Abducens (sixth cranial).	Motion.	Fourth ventricle.	External rectus of eye.	
Arnold's.	See <i>Auricular</i> .			
Auditory (eighth cranial, portio mollis of seventh).	Hearing.	Restiform body.	Internal ear.	Vestibular, cochlear,
Auricular.	Sensation.	Lesser occipital.	Integument of posterior and upper portion of pinna.	

TABLE OF NERVES.—*Continued.*

NAME.	FUNCTION.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Auricular, anterior.	Sensation.	Inferior maxillary.	Integument of external ear.	
Auricular (Arnold's).	Sensation.	Vagus.	External ear.	
Auricular, posterior.	Motion.	Facial.	Retrahens aurem, attollens aurem, occipitofrontalis.	Auricular, occipital.
Auricularis magnus.	Sensation.	Cervical plexus, second and third cervical.	Parotid gland, face, ear.	Facial, mastoid and auricular.
Auriculotemporal.	Sensation.	Inferior maxillary.	Pinna and temple.	Articular, two branches to meatus, parotid, anterior auricular, superficial temporal.
Buccal.	Motion.	Facial.	Buccinator and orbicularis oris muscles.	
Buccal, long.	Sensation, motion?	Inferior maxillary.	Cheek.	Superior and inferior buccinator and external pterygoid.
Calcanean, internal.	Sensation.	Posterior tibial.	Fascia and integument of heel and sole.	
Cardiac (cervical and thoracic).	Inhibition.	Vagus.	Heart.	Branches to cardiac plexuses.
Of Casser.	See <i>Perforating</i> , and <i>Musculo-cutaneous</i> .			
Cervical, first (anterior division).	Motion and sensation.	Cord.	Rectus lateralis and two anterior recti.	Filaments to vagus, hypoglossal, sympathetic.
Cervical, first (posterior division).	Motion and sensation.	Cord.	Recti, obliqui, complexus.	Communicating and cutaneous filaments.
Cervical, fourth (anterior division).	Motion and sensation.	Cord.	Shoulder.	Communicating filaments, muscular, etc.
Cervical, second (anterior division).	Motion and sensation.	Cord.	Communicating.	Ascending, descending, communicating.
Cervical, second (posterior division).	Motion and sensation.	Cord.	Obliquus inferior, scalp, ear, complexus, splenius, trachelomastoid.	Internal or occipitalis major, and external.
Cervical, third (anterior division).	Motion and sensation.	Cord.	Communicating.	Ascending, descending, and communicating filaments.
Cervical, third (posterior division).	Motion and sensation.	Cord.	Occiput, splenius, complexus.	Internal and external.
Cervicals, fifth to eighth (anterior divisions).	Motion and sensation.	Cord.	Brachial plexus.	Communicating.
Cervicals, fourth to eighth (posterior divisions).	Motion and sensation.	Cord.	Muscles and skin of neck.	Internal and external branches.
Cervicofacial.	Motion.	Facial.	Lower part of face and part of neck.	Buccal, supramaxillary, inframaxillary.
Chorda tympani.	Motion.	Facial.	Tongue, tympanum, submaxillary gland.	

TABLE OF NERVES.—*Continued.*

NAME.	FUNCTION.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Ciliary.	Sensation, nutrition, motion.	Ciliary ganglion.	Eye-ball.	
Circumflex.	Motion and sensation.	Brachial plexus.	Teres minor, deltoid, and skin.	Anterior, posterior, and articular.
Coccygeal.	Motion.	Coccygeal plexus.	Coccygeus and gluteus maximus.	
Cochlear.	Hearing.	Auditory.	Cochlea.	
Colli, superficialis.	Sensation.	Cervical plexus.	Platysma myoides and anterolateral parts of neck.	Ascending and descending branches.
Communicans noni.	Motion and sensation.	Second cervical, third cervical.	Descendens noni, depressor muscles of hyoid bone.	Omohyoid, ansa hypoglossi.
Communicating.	Motion and sensation.	Cervical plexus.	Spinal accessory.	
Communicating.	Sensation and motion.	First and second cervical.	Vagus, hypoglossal, sympathetic.	
Of Cotunnus.	See <i>Nasopala</i> .	<i>tine nerve.</i>		
Crural.	Sensation.	Genitocrural.	Shin, upper and central part anterior aspect of thigh.	
Crural, anterior.	Motion and sensation.	Lumbar plexus, second, third, and fourth lumbar nerves.	Thigh.	Middle and internal cutaneous, long saphenous, muscular, articular.
Cutaneous.	Sensation.	Musculospiral.	Skin of arm, radial side of forearm.	One internal, two external.
Cutaneous.	Sensation.	Ulnar.	Wrist and palm.	First and palmar cutaneous.
Cutaneous, dorsal.	Sensation.	Ulnar.	Little and ring fingers.	Communicating branches.
Cutaneous, external.	Sensation.	Second and third lumbar.	Skin of thigh.	Anterior and posterior.
Cutaneous, internal.	Sensation.	Brachial plexus.	Forearm.	Anterior and posterior branches.
Cutaneous, lesser internal (of Wrisberg).	Sensation.	Brachial plexus.	Inner side of arm.	
Cutaneous, middle and internal.	Sensation; motion?	Anterior crural.	Skin of thigh.	Communicating.
Cutaneous, perforating.	Sensation.	Fourth sacral.	Integument covering gluteus maximus.	
Cyon's.	See <i>Depressor.</i>			
Dental, inferior or mandibular.	Sensation.	Inferior maxillary.	Teeth, muscles.	Mylohyoid, incisive, mental, dental.
Dentals, anterior and posterior.	Sensation.	Superior maxillary.	Teeth.	
Depressor.	Lowering of the blood-pressure.	In the rabbit, from the vagus.	Heart.	

TABLE OF NERVES.—*Continued.*

NAME.	FUNCTION.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Descendens hypoglossi.	Motor.	Cervical plexus.	Omohyoid, sternohyoid, sternothyroid, thyrohyoid, geniohyoid, hyoglossus, and muscles of the tongue.	Muscular, lingual.
Descendens noni.	See <i>Descendens hypoglossi.</i>			
Digastric.	Motion.	Facial.	Posterior belly of digastric.	
Dorsal, 12 (anterior and posterior divisions).	Motion and sensation.	Cord.	Muscles and skin of trunk.	External, internal, cutaneous.
Dorsal (of penis).	Sensation.	Pudic.	Penis.	
Esophageal.	Motion.	Vagus.	Mucous and muscular coats of esophagus.	Esophageal plexus.
Facial.	Sensation.	Great auricular.	Skin over parotid.	
Facial (seventh cranial, portio dura).	Motion.	Floor of fourth ventricle.	Face, ear, palate, tongue.	Petrosals, tympanic, chorda tympani, communicating, posterior auricular, digastric, stylohyoid, lingual, temporal, malar, infra-orbital, buccal, superior and inferior maxillary.
Frontal.	Sensation.	Ophthalmic.	Forehead and eyelids.	Supraorbital, supratrochlear.
Gastric.	Motion.	Vagus.	Stomach.	
Genital.	Motion and sensation.	Genitocrural.	Cremaster muscle.	
Genitocrural.	Motion and sensation.	First and second lumbar.	Cremaster and thigh.	Genital, crural, communicating.
Glossopharyngeal (ninth cranial).	Sensation and taste.	Fourth ventricle.	Tongue, middle ear, tonsils, pharynx, meninges.	Tympanic, carotid, pharyngeal, muscular, tonsillar, lingual.
Gluteal, inferior.	Motion.	Sacral plexus (second and third sacral nerves).	Gluteus maximus.	
Gluteal, superior.	Motion.	Sacral plexus.	Glutei, tensor vaginæ femoris.	
Gustatory.	See <i>Lingual.</i>			
Hemorrhoidal (inferior).	Sensation and motion.	Pudic.	External sphincter ani, and adjacent integument.	
Hepatic.	(?)	Vagus.	Liver.	Hepatic plexus.
Hirschfeld's.	Motion.	Facial.	Styloglossus and palatoglossus.	
Hypogastric.	Sensation.	Iliohypogastric.	Skin about external abdominal ring.	
Hypoglossal (twelfth cranial).	Motion.	Floor of fourth ventricle.	Hypoglossus and hyoid muscles.	Descendens noni or hypoglossi, muscular, thyrohyoid, geniohyoid, and meningeal.

TABLE OF NERVES.—*Continued.*

NAME.	FUNCTION.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Iliac.	Sensation.	Iliohypogastric.	Integument covering forepart of gluteal region.	
Iliac.	Sensation.	Last dorsal.	Integument covering forepart of gluteal region.	
Iliohypogastric.	Motion and sensation.	First lumbar.	Abdominal and gluteal regions.	Iliac, hypogastric, communicating.
Ilioinguinal.	Motion and sensation.	First lumbar.	Inguinal region and scrotum.	Muscular, cutaneous, and communicating.
Incisive.	Sensation.	Inferior dental.	Canine and incisor teeth and corresponding portion of gums.	
Inframaxillary.	See <i>Inframaxillary</i> .			
Inframaxillary.	Motion.	Facial.	Platysma myoides.	
Infraorbital.	Sensation and motion.	Facial.	Nose and lip.	Palpebral, nasal, labial.
Infratrochlear.	Sensation.	Nasal.	Skin and conjunctiva of inner part of eye, lacrimal sac.	
Intercostal.	Motion and sensation.	Spinal cord.	Muscles and integument of thorax.	Muscular, anterior and lateral cutaneous.
Intercostohumeral.	Sensation.	Second intercostal.	Integument of upper two-thirds of inner and posterior part of arm.	
Interosseous, anterior.	Motion.	Median.	Deep muscles of forearm.	
Interosseous, posterior.	Motion and sensation.	Musculospiral.	Carpus and radial and posterior brachial regions.	
Jacobson's.	See <i>Tympanic</i> .	<i>of the glossopharyngeal.</i>		
Labial.	Motion and sensation.	Superior maxillary.	Muscles and mucous membrane of lips.	
Lacrimal.	Sensation.	Ophthalmic.	Lacrimal gland and conjunctiva.	
Of Lancisi.	Longitudinal striations on upper surface of corpus callosum.			
Laryngeal, recurrent or inferior.	Motion.	Vagus.	Larynx.	Branches to all laryngeal muscles except cricothyroid.
Laryngeal, superior.	Sensation and motion.	Vagus.	Larynx.	External—cricothyroid muscle and thyroid gland; internal—mucous membrane of larynx.
Lingual.	Motion and sensation.	Facial.	Mucous membrane of tongue, palatoglossus and styloglossus muscles.	

TABLE OF NERVES.—*Continued.*

NAME.	FUNCTION.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Lingual.	Sensation.	Glossopharyngeal.	Circumvallate papillæ and glands of tongue.	
Lingual.	Taste and sensation.	Inferior maxillary.	Tongue and mouth.	
Lumbar (5).	Motion and sensation.	Cord.	Lumbar and genital regions.	Anterior and posterior divisions, lumbar plexus.
Malar.	Motion.	Facial.	Lower part of orbicularis palpebrarum, and eyelids.	
Malar.	Sensation.	Orbital.	Skin over malar bone.	
Mandibular.	See <i>Maxillary</i> ,	<i>inferior.</i>		
Masseteric.	Motor.	Inferior maxillary.	Masseter muscle (and temporal?).	
Mastoid.	Sensation.	Great auricular.	Skin over mastoid process.	
Mastoid.	Motion.	Lesser occipital.	Skin over mastoid process.	
Maxillary, inferior.	Sensation, motion, and taste.	Trigeminus.	Muscles of mastication, ear, cheek, tongue, teeth.	Masseteric, auriculotemporal, buccal, gustatory, inferior dental.
Maxillary, superior.	Sensation.	Trigeminus.	Cheek, face, teeth.	Orbital, sphenopalatine, dentals, infraorbital.
Median.	Motion and sensation.	Brachial plexus.	Pronator radii teres, flexors, two lumbricales, fingers, palm.	Muscular, anterior interosseous, palmar cutaneous.
Meningeal.	Sensation.	Glossopharyngeal.	Pia and arachnoid.	
Meningeal.	Sensation.	Hypoglossal.	Dura mater.	
Meningeal.	Sensation.	Vagus.	Dura mater around lateral sinus.	
Meningeal, recurrent.	Sensation.	Inferior maxillary.	Dura mater and mastoid cells.	
Mental.	Motion and sensation.	Inferior maxillary.	Mucous membrane of lower lip and chin.	
Motor oculi (third cranial).	Motion.	Floor of aqueduct of Sylvius.	All muscles of the eye, except external rectus, superior oblique, and orbicularis palpebrarum.	
Musculocutaneous (of Casser).	Motion and sensation.	Brachial plexus.	Coracobrachialis, biceps, brachialis anticus, forearm.	Anterior and posterior.
Musculocutaneous.	Motion and sensation.	External popliteal.	Muscles of fibular side of leg, skin of dorsum of foot.	Internal, external.
Musculospiral.	Motion and sensation.	Brachial plexus.	Back of arm and forearm, skin of back of hand.	Musculocutaneous, radial, posterior interosseous.
Mylohyoid.	Motion.	Inferior maxillary.	Mylohyoid and digastric muscles.	
Nasal.	Sensation.	Dental, anterior.	Mucous membrane of inferior meatus.	

TABLE OF NERVES.—*Continued.*

NAME.	FUNCTION.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Nasal.	Sensation.	Maxillary, superior.	Integument of lateral aspect of nose.	
Nasal.	Sensation.	Ophthalmic.	Iris, ciliary ganglion, nose.	Ganglionic, ciliary, infraorbital.
Nasal, inferior.	Sensation.	Anterior palatine.	Mucous membrane of nose.	
Nasal, superior.	Sensation.	Meckel's ganglion.	Mucous membrane of nose and posterior ethmoid cells.	
Nasopalatine.	Sensation.	Meckel's ganglion.	Nasal septum.	
Obturator.	Motion and sensation.	Lumbar plexus, third and fourth nerves.	Obturator externus, adductors, hip-joint, and skin.	Anterior and posterior articular and communicating.
Obturator, accessory.	Motion and sensation.	Lumbar plexus.	Pectineus and hip-joint.	
Occipital, smallest or third.	Sensation.	Third cervical.	Integument of occiput.	
Occipitalis minor.	Sensation.	Second cervical.	Occipitofrontalis, ear, etc.	Communicating, auricular.
Occipitalis magnus.	Motion and sensation.	Second cervical.	Complexus, trapezius, and scalp.	
Oculomotor.	See <i>Motor oculi</i> .			
Olfactory (first cranial).	Smell.	Frontal lobe, optic thalamus, island of Reil.	Schneiderian membrane of nose.	
Ophthalmic.	Sensation.	Trigeminal.	Forehead, eyes, nose.	Frontal, lacrimal, nasal.
Optic (second cranial).	Sight.	Cortical center in occipital lobe.	Retina.	
Orbital.	Sensation.	Meckel's ganglion.	Mucosa of posterior ethmoid cells, and sphenoidal sinus.	
Orbital.	Sensation.	Superior maxillary.	Temple and cheek.	Temporal and malar.
Palatine, anterior or great.	Sensation.	Meckel's ganglion.	Hard palate, gums, and nose.	Two inferior nasal.
Palatine, external.	Sensation.	Meckel's ganglion.	Tonsil and soft palate.	
Palatine, posterior or small.	Motor.	Meckel's ganglion.	Levator palati and azygos uvulae.	
Palmar, cutaneous.	Sensation.	Median.	Thumb and palm.	
Palmar, deep.	Motion.	Ulnar.	Little finger, dorsal and palmar interossei, two inner lumbricales, abductor pollicis.	
Palmar, superficial.	Sensation and motion.	Ulnar.	Palmaris brevis, inner side of hand, and little finger.	
Palpebral.	Motor.	Superior maxillary.	Integument of lower lid.	
Parotid.	Sensation.	Auriculotemporal.	Parotid gland.	

TABLE OF NERVES.—*Continued.*

NAME.	FUNCTION.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Patellar.	Sensation.	Long saphenous.	Integument over patella and plexus patellæ.	
Patheticus (fourth cranial).	Motion.	Valve of Vieussens.	Superior oblique of eye.	
Pectineus.	Motion.	Anterior crural.	Pectineus muscle.	
Perforating (of Casser).	See <i>Musculo-cutaneous.</i>			
Perineal.	Motion and sensation.	Pudic.	Perineum, genitalia, and skin of perineal region.	Cutaneous and muscular.
Perineal.	Motion and sensation.	Fourth sacral.	External sphincter ani and integument of anus.	
Petrosals.	Motion.	Facial.	Ganglia and plexus.	Great, small, external to Meckel's ganglion, otic ganglion, and meningeal plexus, respectively.
Pharyngeal.	Motion and sensation.	Glossopharyngeal.	Pharynx.	Enters into formation of pharyngeal plexus.
Pharyngeal.	Sensation.	Meckel's ganglion.	Upper part of pharynx, posterior nares, and sphenoidal sinus.	
Pharyngeal.	Motion.	Vagus.	Pharynx.	Pharyngeal plexus, muscles and mucosa.
Pharyngeal.	Sensation.	Sympathetic.	Pharynx.	Helps to form the pharyngeal plexus.
Phrenic.	Motion and sensation.	Third, fourth, and fifth cervical.	Diaphragm, pericardium, pleura.	
Plantar, external.	Motion and sensation.	Posterior tibial.	Little toe and deep muscles of foot.	Superficial and deep.
Plantar, internal.	Sensation and motion.	Posterior tibial.	Sole of foot, adductor pollicis, flexor brevis digitorum, toes.	Cutaneous, muscular, articular, digital.
Pneumogastric (tenth cranial, par vagum or vagus).	Sensation and motion.	Floor of fourth ventricle.	Ear, pharynx, larynx, heart, lungs, esophagus, stomach.	Auricular, pharyngeal, superior and inferior laryngeal, recurrent laryngeal, cardiac, pulmonary, esophageal, gastric, hepatic, communicating, meningeal.
Popliteal, external.	Sensation and motion.	Great sciatic.	Extensors of foot, skin, and fascia.	Anterior tibial, musculocutaneous, articular, cutaneous.
Popliteal, internal.	Motion and sensation.	Great sciatic.	Knee, gastrocnemius, tibia posterior, plantaris, soleus, popliteus, skin of foot.	Articular, muscular, cutaneous, external saphenous, plantar.
Pterygoid, external.	Motion.	Inferior maxillary.	External pterygoid muscle.	
Pterygoid, internal.	Motion.	Inferior maxillary.	Internal pterygoid muscle.	
Pudendal, long (nerve of Sæmmering).	Sensation.	Small sciatic.	Integument of genitalia and inner and proximal part of thigh.	

TABLE OF NERVES.—*Continued.*

NAME.	FUNCTION.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Podic.	Motion and sensation.	Sacral plexus.	Perineum, anus, genitalia.	Inferior hemorrhoidal, perineal, cutaneous, dorsal of penis.
Pulmonary, anterior and posterior.	(7)	Vagus.	Lungs.	Branches to pulmonary plexuses.
Radial.	Sensation.	Musculospiral.	Skin of radial side and ball of thumb; skin on posterior surface of ulnar side of thumb; skin of index finger, middle finger, and radial side of ring-finger.	External and internal.
Sacral.	Motion and sensation.	Cord.	Multifidus spine, skin of gluteal region.	Branches to sacral plexus.
Saphenous, external or short.	Sensation.	Internal popliteal.	Integument of foot and little toe.	
Saphenous, long or internal.	Sensation.	Anterior crural.	Knee, ankle.	Cutaneous, patellar, communicating.
Of Scarpa.	See <i>Nasopalatine</i> .			
Sciatic, great.	Motion and sensation.	Sacral plexus.	Skin of leg, muscles of back of thigh, and those of leg and foot.	Articular, muscular, popliteal.
Sciatic, small.	Sensation and motion.	Sacral plexus.	Perineum, back of thigh and leg, gluteus maximus.	Muscular, cutaneous, long pudendal.
Of Sommering.	See <i>Podradial</i> .	<i>long.</i>		
Sphenopalatine.	Sensation.	Superior maxillary.	Meckel's ganglion.	
Spinal.	Motion and sensation.	Spinal cord.	Trunk.	
Spinal accessory (eleventh cranial).	Motion.	Floor of fourth ventricle.	Sternocleidomastoid, trapezius.	
Splanchnic, great.	Sympathetic.	Thoracic ganglia.	Semilunar ganglion, renal and suprarenal plexuses.	
Splanchnic, lesser.	Sympathetic.	Tenth and eleventh thoracic ganglia, great splanchnic.	Celiac plexus and great splanchnic.	
Splanchnic, renal or smallest.	Sympathetic.	Last thoracic ganglion.	Renal and celiac plexuses.	
Stapedial.	Motion.	Facial.	Stapedius muscle.	
Stylohyoid.	Motion.	Facial.	Stylohyoid muscle.	
Subscapular.	Motion.	Brachial plexus.	Subscapular, teres major, and latissimus dorsi.	
Supraacromial.	Sensation.	Cervical plexus.	Skin over deltoid.	
Supraclavicular, descending.	Sensation.	Third and fourth cervical.	Skin of neck, breast, and shoulder.	Sternal, claviclar, acromial.
Supramandibular.	See <i>Maxillary</i> .	<i>superior.</i>		

TABLE OF NERVES.—*Continued.*

NAME.	FUNCTION.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Supramaxillary.	See <i>Maxillary</i> .	<i>superior.</i>		
Supraorbital.	Sensation.	Ophthalmic.	Upper lid, forehead.	Muscular, cutaneous, and pericranial.
Suprascapular.	Motion and sensation.	Brachial plexus.	Scapular muscles.	
Suprasternal.	Sensation.	Cervical plexus.	Integument over upper part of sternum.	
Supratrochlear.	Sensation.	Ophthalmic.	Forehead and upper eyelid.	Muscular and cutaneous.
Sympathetic.	See <i>Sympathetic.</i>			
Temporal.	Motion.	Inferior maxillary.	Temporal muscle.	
Temporal.	Sensation.	Orbital.	Integument over temporal muscle.	
Temporal.	Motion.	Temporofacial.	Orbicularis palpebrarum, occipitofrontalis, attrahens and attollens aurem, corrugator supercilii.	Muscular.
Temporal, superficial.	Sensation.	Auriculotemporal.	Integument over temporal fascia.	
Temporofacial.	Motion.	Facial.	Upper part of face.	Temporal, malar, infra-orbital.
Temporomalar.	See <i>Orbital.</i>			
Thoracic, posterior or long. (External respiratory nerve of Bell).	Motion.	Brachial plexus.	Serratus magnus.	
Thoracic, spinal.	Motion and sensation.	Cord.	Muscles and skin of thorax.	
Thoracic, anterior and exterior.	Motion.	Brachial plexus.	Pectoralis major and minor.	
Tibial, anterior.	Motion and sensation.	External popliteal.	Tibialis anticus, extensor longus digitorum, peroneus tertius, joints of foot, skin of great toe.	Muscular, external, internal.
Tibial, posterior.	Motion and sensation.	Internal popliteal.	Tibialis posticus, flexor longus digitorum, flexor longus pollicis, skin of heel and sole, knee-joint.	Plantar, muscular, calcaneoplantar, cutaneous or internal calcanean, articular.
Tonsillar.	Sensation.	Glossopharyngeal.	Tonsil, soft palate, and fauces.	
Trigeminal or trifacial (5th cranial).	Motion and sensation (taste).	Floor of fourth ventricle.	Skin and structures of face, tongue, and teeth.	Ophthalmic, superior and inferior maxillary.
Trochlear.	See <i>Pathetic.</i>			
Tympanic.	Motion.	Facial.	Stapedius and laxator tympani muscles.	
Tympanic.	Sensation.	Sympathetic.	Tympanum.	

TABLE OF NERVES.—*Continued.*

NAME.	FUNCTION.	ORIGIN.	DISTRIBUTION.	BRANCHES.
Tympanic (Jacobson's nerve).	Motion.	Glossopharyngeal.	Tympanum.	Tympanic plexus, and communicating.
Ulnar.	Motion and sensation.	Brachial plexus.	Muscles, shoulder-joint and wrist-joint, and skin of little finger.	Articular, muscular, palmar cutaneous, dorsal, superior palmar, deep palmar.
Vagus.	See <i>Pneumo</i>	<i>gastric.</i>		
Vestibular.	Sensation.	Auditory.	Utricle and ampullæ of the semicircular canals.	
Vidian.	Sensation.	Union of large superficial and deep petrosal.	Sphenomaxillary fossa, and posterior part of upper nasal meatus.	
Of Willis.	See <i>Spinal ac</i>	<i>cessory.</i>		
Wrisberg's.	See <i>Cutaneous,</i>	<i>lesser internal.</i>		

Nervi (*nur'-vi*) [L.]. Plural of *Nervus*, a nerve. **N. nervorum**, the small nerves distributed to the nerve-sheaths.

Nervine (*nur'-vin*) [*nervus*, nerve]. 1. Acting upon the nerves. 2. A remedy that calms nervous excitement or acts favorably on nervous diseases.

Nervous (*nur'-vus*) [*nervus*, a nerve]. 1. Pertaining to or composed of nerves or nerve-structures. 2. Characterized by excessive irritability of the nervous system. **N. Fluid**, a hypothetic fluid supposed to traverse the nerves from the nerve-centers to the periphery.

Nervousness (*nur'-vus-nes*) [*nervus*, nerve]. A condition of excessive excitability of the nervous system, characterized by great mental and physical unrest.

Nessler's Reagent. A solution of potassium iodid, mercuric chlorid, and sodium hydroxid, used in estimating the amount of ammonia in water.

Nettle-rash. See *Urticaria*.

Neumann's Disease (*noi'-mans*). See *Pemphigus vegetans*.

Neura (*nu'-rah*) [*νεῦρον*, nerve]. Synonym of *Neuron*.

Neurad (*nu'-rad*) [*νεῦρον*, a nerve; *ad*, to]. Toward the neural aspect or axis.

Neuradynamia (*nu'-rah-din-a'-me-ah*). See *Neurasthenia*.

Neural (*nu'-ral*) [*νεῦρον*, nerve]. Pertaining to nerves or nervous tissue. **N. Arch**, the part of the vertebra that encloses the spinal cord, formed by the two neurapophyses. **N. Axis**, the spinal cord. **N. Canal**. 1. The dorsal tube of the embryo, formed

by the union of the dorsal folds and constituting the earliest traces of the nervous system.

2. The bony canal comprising the cavity of the cranium and vertebral column, which in the vertebrate animals contains the central nervous system. It is situated dorsad, the hemal canal, enclosing the heart, etc., being ventrad. See *Canal*, *Hemal*. **N. Groove**, the medullary groove. **N. Lamina**, the lateral portion of the neural arch of a vertebra.

Neuralgia (*nu'-ral'-je-ah*) [*νεῦρον*, nerve; *ἄλγος*, pain]. Severe paroxysmal pain along the course of a nerve and not associated with demonstrable structural changes in the nerve. According to their anatomic situation the following forms of neuralgia are described: trigeminal N., tic douloureux, or prosopalgia; supraorbital N.; cervicoccipital N.; cervicobrachial and brachial N.; intercostal N.; sciatica or ischias; coccygodynia; visceral N. (as hepatic, gastric, intestinal, uterine, ovarian N.). According to their causes, neuralgias are classed as anemic, malarial, gouty, rheumatic, syphilitic, diabetic, toxic (*e. g.*, alcoholic, saturnine), hysteric, and reflex. The pain of N. is sharp, stabbing, and paroxysmal, lasting usually but a short time; tenderness is often present at the points of exit of the nerve (*points douloureux*). Intercostal neuralgia is at times associated with herpes zoster.

Neuralgic (*nu'-ral'-jik*) [*νεῦρον*, nerve; *ἄλγος*, pain]. Pertaining to, or affected with, neuralgia.

Neurapophysis (*nu'-rap-off'-is-is*) [*νεῦρον*, nerve; *ἀπόφυσις*, offshoot]. Either one of the two apophyses on each vertebra which

blend and form the neural arch, or the dorsal wall of the spinal foramen.

Neurasthenia (*nu-ras-then-i'-ah*, *nu-ras-the'-ne-ah*) [*νεῦρον*, nerve; *ἀσθένεια*, weakness]. A group of symptoms resulting from debility or exhaustion of the nerve-centers. Among the more common symptoms are a lack of energy, undue readiness of fatigue, disinclination to activity, a sense of fulness or pressure at the top of the head, pain in the back, impaired memory, and disturbed sleep; gastrointestinal symptoms, such as anorexia, constipation, fulness after eating; amenorrhea and dysmenorrhea in women, and spermatorrhea and impotence in men. **N.**, **Cerebral**, **N.** marked by depression, inability to concentrate the mind, insomnia, irritability, headache, visual disturbances, etc. **N. gastrica**, nervous dyspepsia, a condition of disturbed functional activity of the stomach, as a rule without retardation of digestion; flatulence, pain, palpitation of the heart, and constipation are prominent symptoms. **N.**, **Sexual**, a depressed state of the nervous system associated with disturbance of the sexual function; it is characterized by pain in the back, tender points along the spine, weakness of the extremities, great prostration on slight exertion, neuralgic pains, and other nervous manifestations.

Neurasthenic (*nu-ras-then'-ik*) [*νεῦρον*, nerve; *ἀσθένεια*, weakness]. Relating to, or characterized by neurasthenia.

Neuraxis (*nu-raks'-is*) [*νεῦρον*, nerve; *ἄξις*, axis]. The neural axis.

Neuraxon (*nu-raks'-un*) [*νεῦρον*, nerve; *ἄξις*, axis]. The axis-cylinder process of a nerve-cell.

Neure (*nūr*) [*νεῦρον*]. Synonym of *Neuron*.

Neurectasis, **Neurectasy** (*nu-rek'-tas-is*, *nu-rek'-tas-e*) [*νεῦρον*, nerve; *ἐκτασις*, stretching]. Nerve-stretching.

Neurectomy (*nu-rek'-to-me*) [*νεῦρον*, nerve; *ἐκτομή*, excision]. Excision of a part of a nerve.

Neurenteric (*nu-ren-ter'-ik*) [*νεῦρον*, nerve; *έντερον*, intestine]. Pertaining to the embryonic neural canal and the intestinal tube. **N. Canal**, a temporary communication existing between the neural canal and the intestinal tube of the embryo. It is also called Kowalevsky's Canal.

Neuridin (*nu'-rid-in*) [*νεῦρον*, a nerve], $C_8H_{14}N_2$. A ptomain produced in the putrefaction of horseflesh, beef, human muscle, fish, cheese, etc. It has a repulsive odor and is nonpoisonous. See *Ptomains*, Table of.

Neurilemma (*nu-ril-em'-ah*) [*νεῦρον*, nerve; *λέμμα*, bark]. The sheath encasing a nerve-fiber; the sheath of Schwann. See *Nerve-fiber*.

Neurin (*nu'-rin*) [*νεῦρον*, nerve]. 1. The

albuminous substance forming the basis of nerve-tissue. 2. $C_8H_{12}NO$; a poisonous ptomain obtained from decomposing flesh and in the decomposition of protagon by barium hydroxid. Its action is similar to that of muscarin. 3. An extract of nerve-tissue employed therapeutically. See *Organotherapy*.

Neurit (*nu'-rit*) [*νεῦρον*, nerve]. Synonym of *Neurite*.

Neurite (*nu'-rit*) [*νεῦρον*, nerve]. The axis-cylinder process of a nerve-cell.

Neuritic (*nu-ril'-ik*) [*νεῦρον*, nerve; *τις*, inflammation]. Pertaining to neuritis.

Neuritis (*nu-ril'-tis*) [*νεῦρον*, nerve; *τις*, inflammation]. Inflammation of a nerve. **N.**, **Alcoholic**. See *N.*, *Multiple*. **N.**, **Ascending**, inflammation extending from the periphery of a nerve toward the spinal cord or brain. **N.**, **Degenerative**, neuritis in which the nerve-substance proper degenerates. It is the same as *N.*, *Parenchymatous*. **N.**, **Descending**, neuritis the result of extension of disease from the spinal cord or the brain toward the periphery. **N.**, **Diabetic**, a polyn neuritis sometimes seen in diabetes, and probably the result of autointoxication with the products of faulty metabolism. **N.**, **Interstitial**, inflammation of the interstitial connective tissue of a nerve-trunk. **N.**, **Lipomatous**, that form in which the nerve-fibers are completely destroyed and replaced by a fibrous connective tissue in which much fat is deposited. **N.**, **Malarial**, neuritis due to the malarial poison. **N.**, **Multiple**, the simultaneous inflammation of several nerve-trunks, usually symmetrically situated on both sides of the body. Its most common cause is alcoholic poisoning; it may be due to arsenic, malaria, mercury, lead; to diphtheria, pneumonia, typhoid fever, and other infectious diseases. Beriberi is a form of multiple neuritis. The symptoms depend somewhat upon the cause: in alcoholic neuritis, the most frequent variety, there are paresthesias and pain, followed by paralysis of the legs and arms, wrist-drop and foot-drop, with edema, loss of reflexes, and delirium. Anesthesia is slight; there may be hyperesthesia or pain; in some cases ataxia is present. The sphincters are generally not affected. **N.**, **Optic**, inflammation of the optic nerve. See *Papillitis*. **N.**, **Parenchymatous**, a form in which the medullary substance and the axis-cylinders are chiefly involved, the interstitial tissue being but little altered or only affected secondarily. It is also called degenerative neuritis. **N.**, **Pressure-**, inflammation of a nerve resulting from compression.

Neuro- (*nu'-ro-*) [*νεῦρον*, nerve]. A prefix signifying connection with or relation to a nerve.

Neuroblast (*nu'-ro-blast*) [*νεῦρον*, nerve; *βλαστός*, germ]. A cell derived from the primitive ectoderm, and giving rise to nerve-fibers and nerve-cells.

Neurocardiac (*nu-ro-kar'-de-ak*) [*νεῦρον*, nerve; *καρδία*, heart]. Pertaining to the nervous system and the heart. **N. Disease**, exophthalmic goiter.

Neurocyte (*nu'-ro-sit*) [*νεῦρον*, nerve; *κύτος*, cell]. Synonym of *Neuron*.

Neurodendron (*nu-ro-den'-dron*) [*νεῦρον*, nerve, *δένδρον*, ball]. Synonym of *Neuron*.

Neurodermatitis (*nu-ro-der-mat-i'-tis*) [*νεῦρον*, nerve; *δέρμα*, skin; *ιτις*, inflammation]. A neurotic affection of the skin associated with itching.

Neurodin (*nu'-ro-din*) [*νεῦρον*, nerve]. Acetylparaoxyphenylurethane, a crystalline substance used as an antineuralgic and antipyretic. Dose gr. xv-xxiiss (1.0-1.5).

Neuroepithelium (*nu-ro-ep'-e-the'-le-um*) [*νεῦρον*, nerve; *epithelium*]. The highly specialized epithelial structures constituting the terminations of the nerves of special sense, as the rod-cells and cone-cells of the retina, the olfactory cells of the nose, the hair-cells of the internal ear, the gustatory cells of the taste-buds.

Neurofibroma (*nu-ro-fi-brō'-mah*) [*νεῦρον*, nerve; *fibra*, fiber; *ωμα*, tumor]. A tumor of a nerve, composed of fibrous tissue.

Neurogenesis (*nu-ro-jen'-es-is*) [*νεῦρον*, nerve; *γενᾶν*, to produce]. The formation of nerves or nerve-tissue.

Neurogenetic (*nu-ro-gen'-et-ik*) [*νεῦρον*, nerve; *γενᾶν*, to produce]. Pertaining to neurogenesis.

Neuroglia (*nu-ro-gli'-le-ah*) [*νεῦρον*, nerve; *γλία*, glue]. The tissue, probably of ectodermic origin, forming the basis of the supporting framework of the nervous tissue of the cerebrospinal axis. It consists of peculiar cells, the glia-cells, having many fine branching processes.

Neurogliar (*nu-ro-gli'-le-ar*) [*νεῦρον*, nerve; *γλία*, glue]. Pertaining to or resembling neuroglia.

Neuroglioma (*nu-ro-gli'-o'-mah*) [*νεῦρον*, nerve; *γλία*, glue; *ωμα*, tumor]. A tumor composed of neurogliar tissue; a glioma. **N.**, Ganglionar, **N.**, Ganglionare, a glioma containing ganglion-cells.

Neurography (*nu-ro-gli'-ra-fe*) [*νεῦρον*, nerve; *γράφειν*, to write]. A treatise on the anatomy and physiology of the nerves and the nervous system.

Neurokeratin (*nu-ro-ker'-at-in*) [*νεῦρον*, nerve; *κέρας*, horn]. The form of keratin found in nerve-sheaths and the white substance of Schwann.

Neurolemma (*nu-ro-lem'-ah*). Synonym of *Retina*.

Neurologic (*nu-ro-loj'-ik*) [*νεῦρον*, nerve; *λόγος*, science]. Pertaining to neurology.

Neurologist (*nu-ro-l'-o-jist*) [*νεῦρον*, nerve; *λόγος*, science]. One versed in neurology.

Neurology (*nu-ro-l'-o-je*) [*νεῦρον*, nerve; *λόγος*, science]. The branch of medicine dealing with the anatomy, physiology, and pathology of the nervous system.

Neurolysis (*nu-ro-l'-is-is*) [*νεῦρον*, nerve; *λῦσις*, solution]. Exhaustion of a nerve in consequence of overstimulation.

Neuroma (*nu-ro'-mah*) [*νεῦρον*, nerve; *ωμα*, tumor]. A tumor springing from a nerve. It is a true **N.** when made up of nervous tissue proper, and a false **N.** when consisting of other than nervous tissue. False neuromas may be fibromas, myxomas, gliomas, etc. **N.**, **Amputation-**, the neuroma of a stump, forming at the end of a divided nerve. **N.**, **Amyelinic**, a neuroma made up of nonmedullated nerve-fibers. **N.**, **Ganglionic**, **N.**, **Ganglionated**, a neuroma made up of nerve-cells. **N.**, **Myelinic**, one made up of medullated nerve-fibers. **N.**, **Plexiform**, one characterized by the formation of multiple enlargements along the course of the cutaneous nerves, giving rise to the appearance of an intertwining network.

Neuromatous (*nu-ro'-mat-us*) [*νεῦρον*, nerve; *ωμα*, tumor]. Of the nature of a neuroma.

Neuromere (*nu'-ro-mēr*) [*νεῦρον*, nerve; *μέρος*, part]. A natural segment of the cerebrospinal axis.

Neuromimesis (*nu-ro-mi-mē'-sis*) [*νεῦρον*, nerve; *μιμῆσις*, imitation]. Hysteric phenomena resembling true organic disease.

Neuromuscular (*nu-ro-mus'-ku-lar*) [*νεῦρον*, nerve; *musculus*, a muscle]. Pertaining conjointly to nerves and muscles.

Neuromyositis (*nu-ro-mi-o-si'-tis*) [*νεῦρον*, nerve; *μῦς*, muscle; *ιτις*, inflammation]. Myositis associated with neuritis.

Neuron (*nu'-ron*) [*νεῦρον*, nerve]. 1. The cerebrospinal axis taken as a whole. 2. One of the countless number of units of which the nervous system is composed. Each neuron consists of a cell and a series of processes. One of these processes is as a rule an axis-cylinder and terminates free, generally breaking up into a bundle of fine fibrillae. The other processes branch dichotomously, but also terminate in free extremities; they are known as protoplasmic processes or dendrites. In the gray matter of the brain and spinal cord the terminal processes of the neuron form a dense felt-work, but not a communicating reticulum. In every physiologic act involving the nervous system at least two, usually more, neurons participate. The neuron at which the impulse starts is termed archineuron; the one at the termination, the teloneuron. See also *Nerve-cell*.

Neuroparalysis (*nu-ro-par-all'-is-is*) [*νεῦρον*, nerve; *παράλυσις*, paralysis]. Paralysis due to disease of a nerve.

Neuropathic (*nu-ro-path'-ik*) [*νεῦρον*, nerve; *πάθος*, disease]. 1. Characterized by a diseased or imperfect nervous system. 2. Depending upon or pertaining to nervous disease. **N. Eschar**, a bed sore following disease of the spinal cord.

Neuropathology (*nu-ro-path-ol'-o-jē*) [*νεῦρον*, nerve; *πάθος*, disease; *λόγος*, science]. The pathology of diseases of the nervous system.

Neuropathy (*nu-ro-path'-e*) [*νεῦρον*, nerve; *πάθος*, disease]. Any nervous disease.

Neurophysiology (*nu-ro-fis-e-ol'-o-jē*) [*νεῦρον*, nerve; *φύσις*, nature; *λόγος*, science]. The physiology of the nervous system.

Neuroplasm (*nu'-ro-plazm*) [*νεῦρον*, nerve; *πλάσσειν*, to mold]. The protoplasm filling the interstices of the fibrillae of nerve-cells.

Neuroplasty (*nu'-ro-plas-te*) [*νεῦρον*, nerve; *πλάσσειν*, to mold]. A plastic operation on the nerves.

Neuroretinitis (*nu-ro-ret-in'-it-is*) [*νεῦρον*, nerve; *retina*, retina; *ιτις*, inflammation]. Inflammation of both the optic nerve and the retina.

Neurorrhaphy (*nu-ror'-a-fe*) [*νεῦρον*, nerve; *ῥαφή*, suture]. The operation of suturing a divided nerve.

Neurosin (*nu-ro'-sin*) [*νεῦρον*, nerve]. A trade-name for several preparations containing calcium, glycerol, and phosphates.

Neurosis (*nu-ro'-sis*) [*νεῦρον*, nerve]. A functional disease of the nervous system—a disturbance of the nerve-centers or peripheral nerves not due to any demonstrable structural change. **N., Occupation-, N., Professional**, a functional disorder affecting groups of muscles used in the performance of special movements.

Neuroskeleton (*nu-ro-skel'-et-on*) [*νεῦρον*, nerve; *σκελετόν*, skeleton]. The vertebrate endoskeleton, or true skeleton; so called from being made up of parts that correspond with and largely serve to protect portions of the central nervous system.

Neurostearic (*nu-ro-ste-ar'-ik*) [*νεῦρον*, nerve; *στέαρ*, fat]. Pertaining to nervous tissue and fat. **N. Acid**, $C_{18}H_{36}O_2$. An acid isomeric with stearic acid, occurring in the brain.

Neurotabes (*nu-ro-ta'-bēs*) [*νεῦρον*, nerve; *tabes*, wasting]. A form of multiple neuritis resembling posterior sclerosis.

Neurotic (*nu-ro'-tik*) [*νεῦρον*, nerve]. 1. Pertaining to the nerves; nervous. 2. Pertaining to neuroses. 3. Having a disordered nervous system; suffering from a neurosis.

Neurotomy (*nu-ro'-o-me*) [*νεῦρον*, nerve; *τομεν*, to cut]. The division of a nerve.

Neutral (*nu'-tral*) [*neuter*, neither]. Neither

alkaline nor acid; bland and soothing; inactive. **N. Mixture**, liquor potassii citratis.

Neutralize (*nu'-tral-iz*) [*neuter*, neither]. To render neutral; to render inert; to counterbalance an action or influence.

Neutrophile (*nu'-tro-fil*) [*neuter*, neither; *φιλος*, loving]. Stained readily by neutral dyes; applied to certain cells. **N. Leukocytes**, leukocytes the protoplasm of which contains granules colored by neutral stains.

Nevus, Nævus (*ne'-vus*) [L.]. 1. A circumscribed area of pigmentation; a mole.

2. An angioma of the skin, usually congenital; mother's mark. **N. araneus**. See *Acne rosacea*. **N., Capillary**, one that involves the capillaries of the skin. **N., Cutaneous**, a nevus of the skin. **N. lipomatodes**, a large, soft mole, containing a quantity of fat and loose connective tissue.

N. maternus. See *Nevus*, 2d definition.

N. pigmentosus, a mole; a circumscribed, congenital pigmentary deposit in the skin, varying in color from a light fawn to a blackish tint, and often associated with hypertrophy of the hairs. **N. vascularis**. See *Nevus*, 2d definition.

Niccolium, Nickel (*nik-ol'-e-um*, *nik'-el*) [G., *nickel*, nickel]. Symbol Ni; atomic weight 58; quantivalence II, IV. A metal of silver-white luster, resembling iron in physical properties. See *Elements, Table of*. **N. Bromid** ($NiBr_2 + 3H_2O$) has been used in epilepsy. Dose gr. v-x. (0.32-0.65). **N. Chlorid**, $Ni Cl_2$, has been used as a tonic in anemia. Dose gr. ij (0.13). **N. Sulphate** ($NiSO_4 + 7H_2O$), has been used as a tonic. Dose gr. ss-j (0.032-0.065).

Nicol's Prism [after William Nicol, of Edinburgh]. A polished prism of Iceland-spar, cut diagonally across the principal axis, the sections being joined together by means of Canada balsam. It has the property of reflecting the ordinary ray of light out of the field, while the so-called polarized ray is transmitted.

Nicotiana (*nik'-o-she-a'-nah*). See *Tabacum*.

Nicotianin (*nik'-o-she-a'-nin*) [after Jean Nicot, a French ambassador to Portugal]. The volatile principle to which tobacco owes its flavor.

Nicotin (*nik'-o-tin*) [after Jean Nicot, a French ambassador to Portugal], $C_{10}H_{14}N_2$. A liquid poisonous alkaloid found in the leaves of the tobacco-plant.

Nictation (*nik-ta'-shun*). Same as *Nictitation*.

Nictitating (*nik'-tit-a-ting*) [*nictitare*, to wink]. Winking. **N. Membrane**. See *Membrane*. **N. Spasm**, blepharospasm.

Nictitation (*nik-tit-a'-shun*) [*nictitare*, to wink]. Abnormal frequency of winking.

Nidus (*ni'-dus*) [L., nest]. 1. A central

point or focus of infection; a place in which an organism finds conditions suitable for growth and development. 2. A collection of ganglion-cells at the deep origin of a cranial nerve; a nucleus. **N. hirundinis**, a deep fossa in the cerebellum situated between the velum medullare posterius in front and the nodulus and uvula behind.

Niemeyer's Pill (*ni'-mi-er*) [*Niemeyer*, a German physician]. A pill of quinin, grain 1, digitalis, grain one-half, and opium, grain one-quarter. It is used in pulmonary tuberculosis, and is taken every six hours.

Night-blindness. See *Hemeralopia*.

Night-blooming Cereus. See *Cactus grandiflorus*.

Nightmare [AS., *neah*, *niht*, night; *mara*, mare]. A dream characterized by great distress and a sense of oppression or suffocation.

Nightshade. A name applied to plants of the genus *Solanum*. **N.**, Deadly, the *Atropa belladonna*.

Night-terrors. Distressing dreams occurring in children and causing them to wake up with cries of fear.

Nightingale (*ni'-ting-gal*) [after Florence *Nightingale*, a nurse]. A short cape used in hospitals to protect the shoulders and chest of nurses and patients.

Nigrosin (*ni'-gro-sin*) [*niger*, black]. A blue-black anilin dye, used in staining brain-tissue.

Nipple (*ni'-l*) [allied to *neb*, the beak of a bird, from AS., *nebb*, the face]. The conic projection in the center of the mamma, containing the outlets of the milk-ducts. **N.-line**, a vertical line drawn on the surface of the chest through the nipple.

Nisus (*ni'-sus*) [*nissus*, effort, from *niti*, to endeavor]. The periodic desire for procreation manifested in the spring-season by certain species of animals. Also, the contraction of the diaphragm and abdominal muscles for the expulsion of the feces or the urine.

Nit [AS., *hnitu*, a nit]. The egg or larva of the louse.

Niter (*ni'-ter*) [Arab., *nitran*, natron]. Potassium nitrate or saltpeter. **N.**, Cubic, sodium nitrate. **N.**, Sweet Spirit of, spiritus ætheris nitrosi.

Nitrate (*ni'-trat*) [*niter*]. A salt of nitric acid.

Nitrated (*ni'-tra-ted*) [*niter*]. Combined with nitric acid.

Nitration (*ni'-tra'-shun*) [*niter*]. The process of combining or treating with nitric acid.

Nitric (*ni'-trik*) [*niter*]. Pertaining to or containing niter. **N. Acid.** See *Acid, Nitric*.

N.-acid Test, a test for albumin, consisting in the addition of nitric acid to the suspected fluid—if albumin is present a precipitate is formed. The test is usually applied by superimposing the suspected fluid over the acid (Heller's test).

Nitrification (*ni-trif-ik-a'-shun*) [*niter*; *facere*, to make]. The conversion of the nitrogen of ammonia and organic compounds into nitrous and nitric acids, a process constantly going on in nature under the influence of certain bacteria and other agencies.

Nitrifying (*ni-trif'-i-ing*) [*niter*; *facere*, to make]. Converting into nitrous and nitric acids. **N. Bacillus.** See *Bacteria, Table of*.

Nitril (*ni'-tril*) [*niter*]. A compound of cyanogen with an alcohol-radicle in which the nitrogen is trivalent and the radicle is united to the remaining carbon-atom. The nitrils are readily converted into acids.

Nitrite (*ni'-trit*) [*niter*]. A salt of nitrous acid. See *Amyl nitrite*, *Sodium nitrite*, *Potassium nitrite*. The nitrites produce dilatation of the blood-vessels, diminution of the blood-pressure, increased rapidity of the pulse, and depression of the motor centers in the spinal cord. They are used as antispasmodics in asthma and angina pectoris, in spasmodic dysmenorrhea, tetanus, epileptic and hysteric convulsions, and in cases of arteriosclerosis with high arterial tension. Full doses in man give rise to flushing of the face, throbbing, and headache.

Nitro- (*ni'-tro-*) [*niter*]. A prefix denoting combination with nitrogen, or with the radicle NO_2 .

Nitrobacteria (*ni-tro-bak-ter-ia*) [*niter*; *bacterium*]. Bacteria that convert ammonia into nitric acid.

Nitrobenzol, **Nitrobenzene** (*ni-tro-benz'-zol*, *ni-tro-benz'-zene*) $\text{C}_6\text{H}_5\text{NO}_2$. An oily, sweetish liquid made by the action of strong nitric acid on benzol. It is an intermediate product in the manufacture of anilin-oil, and is employed as a flavoring agent under the name of artificial oil of bitter almonds, or oil of mirbane. It is a powerful poison, resembling hydrocyanic acid in action. Persons engaged in its manufacture often suffer from headache and drowsiness.

Nitrocellulose (*ni-tro-sell'-u-lis*). See *Pyraxylin*.

Nitrogen (*ni'-tro-jen*) [Arab., *nitran*, natron; *yemân*, to produce]. Symbol **N**; atomic weight 14; quantivalence 1, III, V. A non-metallic element existing free in the atmosphere, of which it constitutes about 77 per cent. by weight. It is a colorless, odorless gas, incapable of sustaining life. Chemically it is very inert, and combines directly with but few elements. It is an important constituent of all animal and vegetal tissues. With hydrogen it combines to form ammonia, from which a multitude of important compounds are derived. It forms several oxygen-compounds: Nitrogen monoxid, N_2O ; nitrogen dioxid, NO ; nitrogen trioxid,

N_2O_3 ; nitrogen tetroxid, NO_2 ; nitrogen pentoxid, N_2O_5 . The oxids combine with water to form acids. With the halogens nitrogen forms very explosive compounds; nearly all explosive substances are compounds of nitrogen.

Nitrogenous (*ni-troj'-en-us*) [*nitrogen*]. Containing nitrogen.

Nitroglycerin (*ni-tro-gliss'-er-in*) [*niter*; glycerin], $C_3H_5(NO_3)_3$. Glonoin, a colorless oily liquid produced by the action of sulphuric and nitric acids upon glycerol. It is a powerful explosive; physiologically it has the actions of the nitrites, but is more persistent than amyl nitrite, which it most resembles. Dose gr. $\frac{1}{100}$ – $\frac{1}{50}$ (0.0003–0.0013). Spiritus glonoini (U. S. P.), is a 1 per cent. alcoholic solution. Dose m-j (0.065–0.13). Tablets nitroglycerini (B. P.) contain each gr. $\frac{1}{100}$ (0.0006) of nitroglycerin.

Nitrohydrochloric Acid, Nitromuriatic Acid (*ni-tro-hi-dro-klo'-rik*, *ni-tro-mu-reat'-ik*). See *Acid*, *Nitrohydrochloric*.

Nitroso- (*ni-tro'-so-*) [*niter*]. A prefix signifying combination with nitrosyl, the univalent radicle, NO.

Nitrosnitric Acid (*ni-tro-so-ni'-trik*). Fuming nitric acid containing nitrous acid gas.

Nitrous (*ni'-trus*) [*niter*]. 1. Containing nitrogen as a univalent or trivalent element. 2. Pertaining to or derived from nitrous acid HNO_2 , an acid having one atom of oxygen less than nitric acid. **N. Ether**, C_2H_5NO , ethyl nitrite, a very volatile liquid, having properties similar to those of amyl nitrite. **N. Oxid**, N_2O , nitrogen protoxid, hyponitrous oxid, laughing gas. It is used as a general anesthetic in dentistry and in minor surgery. See *Anesthetic*.

Nitrosyl (*ni-tro'-sil*). The univalent radicle NO.

Nocturnal (*noh'-tur-nal*) [*nox*, night]. Pertaining to the night. **N. Emission**, **N. Pollution**, the discharge of semen without coitus during sleep. **N. Enuresis**, incontinence of urine at night during sleep. **N. Epilepsy**, epilepsy in which the convulsions occur at night.

Nodal (*no'-dal*) [*nodus*, a node]. Pertaining to a node. **N. Point**, the point of intersection of convergent rays of light with the visual axis of the eye. The first nodal point is 6.9685 mm. behind the summit of the cornea. The second nodal point is 7.3254 mm. behind the summit of the cornea, or 0.1254 mm. behind the lens.

Node (*nod*) [*nodus*, a swelling or node]. 1. A knob, swelling, or protuberance. 2. A point of narrowing or constriction. **N.**, **Parrot's**, one of the bony prominences on the outer table of the skull, the result of syphilitic periostitis. **N.**, **Ranvier's**, one of the constrictions occurring in the course of

medullated nerve-fibers, where the sheath of Schwann lies upon the axis-cylinder. **N.**, **Syphilitic**, the localized swelling on bones due to syphilitic periostitis.

Nodose (*no'-dos*) [*nodus*, node]. Characterized by nodes; jointed or swollen at intervals.

Nodosity (*no-dos'-it-e*) [*nodus*, node]. 1. The state of having nodes. 2. A node. **N.**, **Bouchard's**, a nodose condition of the second joints of the fingers, supposed to be indicative of gastrectasis. **N.**, **Hagarth's**, **N.**, **Heberden's**, one of the firm nodules on the sides of the distal phalanges of the fingers in rheumatoid arthritis.

Nodular (*nod'-u-lar*) [*nodus*, node]. Composed of or covered by nodules; resembling a nodule.

Nodule (*nod'-ul*) [*nodulus*, dim. of *nodus*, a node]. A small node. **N. of Arantius**. See *Corpus Arantii*. **N. of Cerebellum**, the anterior termination of the inferior vermiciform process of the cerebellum. **N. of Morgagni**. Same as *N. of Arantius*.

Noematachograph (*no-e-ma-tak'-o-graf*). See *Nematachograph*.

Noematachometer (*no-e-ma-tak'-em'-et-er*). See *Nematachometer*.

Noli-me-tangere (*no'-li-me-tan'-jer-e*) [L.]. See *Rodent ulcer*.

Noma (*no'-mah*) [*ropah*, a feeding, from *napiv*, to eat]. Cancrum oris, gangrena oris, gangrenous stomatitis. A grave, usually fatal, form of stomatitis, occurring in debilitated children, generally during the convalescence from one of the exanthemata. It is characterized by the formation of a rapidly spreading ulcer involving the cheek and soon becoming gangrenous. The disease is micro-organismal in origin. **N. pudendi**, **N. vulvæ**, a similar ulceration occurring about the genital region of female children.

Nonan (*no'-nan*) [*nonus*, ninth]. Occurring every ninth day, or having an exacerbation every ninth day.

Nonipara (*non-ip'-ar-ah*) [*nonus*, ninth; *parere*, to bring forth]. See *Multipara*.

Normal (*nor'-mal*) [*norma*, rule]. 1. Conforming to natural order or law. 2. Having the typic structure. **N. Antitoxic Serum**, **N. Therapeutic Serum**, an antitoxic blood-serum of which 0.1 gram is sufficient to neutralize ten times the fatal dose of toxin for a guinea-pig weighing 300 grams. **N. Salt-solution** or **N. Saline Solution**, an aqueous solution of sodium chlorid of a strength similar to that of the body-fluids—usually 0.6–0.75 per cent. **N. Solution**, a solution containing in one liter a quantity of the reagent equal to the molecular weight in grams, and reduced so that the replaceable hydrogen or its equivalent

represents one gram. A decinormal solution is one of one-tenth the strength, and a centinormal solution one of one-hundredth the strength of the normal solution.

Normoblast (*nor'-mo-blast*) [*norma*, rule; *βλαστός*, germ]. A nucleated red corpuscle of the same size as an ordinary red corpuscle.

Norris's Invisible Corpuscles. See *Corpuscle*.

Nose (*nōs*) [A.S., *nōsu*, nose]. The prominent organ occupying the center of the face, the upper part of which (*regio olfactoria*) constitutes the organ of smell, the lower part of which (*regio respiratoria*) represents the commencement of the respiratory tract, in which the inspired air is warmed, moistened, and deprived of impurities. The nose consists of two symmetric cavities, separated by a septum, and is lined internally by mucous membrane (Schneiderian membrane). **N.-bleed**, epistaxis. **N.-piece**, a mechanic device to be attached to the tube of a microscope for holding two, three, or four objectives.

Nosegay, Riolan's. The entire group of muscles arising from the styloid process of the temporal bone.

Nosema (*nas-e'-mah*) [*nosēiv*, to be ill]. 1. Illness, disease. 2. A genus of Schizomycetes. **N. bombycis.** See *Bacteria*, *Table of*.

Nosogeny (*nos-oj'-en-e*) [*νόσος*, disease; *γεννέω*, to beget]. The development of diseases.

Nosography (*nos-og'-ra-fe*) [*νόσος*, disease; *γράφειν*, to write]. A treatise on diseases.

Nosology (*nos-ol'-o-je*) [*νόσος*, disease; *λόγος*, science]. The science of the classification of diseases.

Nosomania (*nos-o-ma'-ne-ah*) [*νόσος*, disease; *μανία*, mania]. 1. A morbid dread of disease. 2. A delusion that one is suffering from disease.

Nosophobia (*nos-o-fo'-be-ah*) [*νόσος*, disease; *φόβος*, fear]. A morbid fear of disease.

Nostalgia (*nos-tal'-je-ah*) [*νόστος*, return; *ἄλγος*, pain]. Homesickness.

Nostoc (*nos'-tok*) [Ger., *nostoch*, nostoc]. A genus of algae having a gelatinous nature.

Nostril (*nos'-tril*) [A.S., *nōsu*, nose; *ðyrel*, orifice]. One of the external orifices of the nose.

Nostrum (*nos'-trum*) [*noster*, ours]. A quack medicine; a secret medicine.

Notal (*no'-tal*) [*νώτον*, back]. Pertaining to the back; dorsal.

Notanencephalia (*no-tan-en-sef'-a'-le-ah*) [*νώτον*, back; *ἄν* priv.; *ἐγκέφαλος*, brain]. Congenital absence of the dorsal part of the cranium.

Notch [O. Du., *noek*, notch]. A deep indentation. **N.**, Intervertebral, one of the depressions on the vertebral pedicles, either

on the upper or lower surface. The apposition of two notches of contiguous vertebrae forms the intervertebral foramen. **N.**, Ischiatic. See *N.*, *Sacrosciatic*. **N.**, Jugular, a notch forming the posterior boundary of the jugular foramen. **N.**, Nasal, an uneven interval between the internal angular processes of the frontal bone, which articulates with the nasal bone and the nasal process of the superior maxillary bone. **N.**, Popliteal, the depression on the posterior surface of the head of the tibia, separating the two tuberosities. **N. of Rivinus**, the deficiency in the osseous tympanic ring, filled in by Shrapnell's membrane. **N.**, *Sacrosciatic*, one of two notches on the posterior edge of the innominate bone. The greater **N.** is just above the spine of the ischium, and is converted into a foramen by the lesser sacrosciatic ligament; the lesser **N.** is below the spine of the ischium, and is converted into a foramen by the sacrosciatic ligaments. **N.**, Sigmoid, a deep semilunar depression separating the coronoid and condyloid processes of the inferior maxillary bone. **N.**, *Sphenopalatine*, the notch that separates the orbital and sphenoid processes of the palate bone. **N.**, *Suprascapular*, a notch in the superior border of the scapula at the base of the coracoid process, for the passage of the suprascapular nerve. **N.**, *Suprasternal*, the depression at the top of the manubrium, between the two sternoclavicular articulations.

Notencephalus (*no-ten-sef'-al-us*) [*νώτον*, back; *ἐγκέφαλος*, brain]. A variety of monster in which the cranial contents are in large part outside the skull, resting upon the back of the neck.

Notochord (*no'-to-kord*) [*νώτον*, back; *χορδή*, a cord]. An elongated cord of cells enclosed in a structureless sheath, which in the embryo represents the vertebral column; the chorda dorsalis, or primitive backbone.

Notomelus (*no-tom'-el-us*) [*νώτον*, back; *μῆλος*, limb]. A form of double monster in which the rudimentary limbs are attached to the back.

Noxious (*nok'-shus*) [*noxius*, harmful]. Harmful; poisonous or deleterious.

Nubecula (*nu-bek'-u-lah*) [dim. of *nubes*, a cloud]. 1. The cloudiness caused by the suspension of insoluble matter in the urine. 2. A cloudiness of the cornea.

Nubility (*nu-bil'-it-e*) [*nubere*, to marry]. The state of sexual development when marriage may be consummated.

Nucha (*nu'-kah*) [L.]. The nape of the neck.

Nuchal (*nu'-kal*) [*nucha*, nape of neck]. Pertaining to the nape of the neck.

Nuck, Canal of. See *Canal*.

Nuclear (*nu'-kle-ar*) [*nucleus*, nucleus]. Pertaining to or resembling a nucleus. **N. Figures**, the peculiar arrangement of the mitome during karyokinesis. **N. Layer**, a stratum of gray matter in the cortex of the brain. **N. Paralysis**, paralysis from lesions of the nuclei of origin of the nerves. **N. Spindle**, delicate striae appearing in the nucleus undergoing mitosis, arranged with the apices pointing toward the poles of the future nuclei. Its function probably is to guide the movements of the mitome threads. **N. Stain**, a pigment showing a strong affinity for nuclei.

Nucleated (*nu'-kle-a-ted*) [*nucleus*, nucleus]. Possessing a nucleus.

Nuclein (*nu'-kle-in*) [*nucleus*, nucleus], $C_{12}H_{19}N_5P_3O_{22}$. An amorphous substance resembling the proteids, and forming the essential chemie constituent of all living cells. It is composed of **Nucleinic Acid** and a base; the former seems to be the same for all nucleins, but the base varies. Nucleins are generally insoluble in dilute acids and soluble in dilute alkalis. They are supposed to represent the germicidal constituent of blood serum.

Nucleoalbumin (*nu-kle-o-al-bu'-min*) [*nucleus*, nucleus; *albumin*]. A compound of a proteid and nuclein, occurring in cell-protoplasm.

Nucleohiston (*nu-kle-o-his'-ton*) [*nucleus*, nucleus; *histon*, tissue]. A substance composed of nuclein and histon found in the leukocytes of the blood. Nuclein induces coagulation of the blood; histon prevents it. The liquid state of the blood is supposed to be dependent on the integrity of the compound formed by these two bodies.

Nucleolus (*nu-kle'-o-lus*) [dim. of *nucleus*, a nucleus]. The small spheric body within the cell-nucleus. Its true function has not as yet been established.

Nucleopetal (*nu-kle-o-pet'-al*) [*nucleus*, nucleus; *petere*, to seek]. Seeking the nucleus (said of the movement of the male pronucleus toward the female pronucleus).

Nucleoplasm (*nu'-kle-o-plazm*) [*nucleus*, nucleus; *πλασσω*, to mold]. 1. The protoplasm of the nucleus. 2. Chromatin.

Nucleus (*nu'-kle-us*) [*nucleus*, from *nux*, nut]. 1. The essential part of a typic cell, usually round in outline, and situated near the center. 2. The center around which the mass of a crystal aggregates. 3. The central element in a compound, as the carbon in hydrocarbons. 4. A collection of gray matter in the central nervous system having a distinct function. 5. The deep origin of a nerve. **N. ambiguus**, **N., Anterior**, a collection of nerve-cells near the N. of the vagus. **N. amygdalæ**, an irregular aggregation of gray matter situated at the apex of

the temporal lobe of the brain, between it and the apex of the middle ventricular horn. **N. of Bechterew**, an ill-defined group of ganglion-cells lying dorsad of Deiters' nucleus, and supposed to give origin to some of the fibers of the middle root of the auditory nerve. It is also called the N. vestibularis. **N., Burdach's**. See *N. funiculi cuneati*. **N., Caudal**. See *N., Oculomotor*. **N., Caudate**, the intraventricular part of the corpus striatum. **N. cinereus**, a term sometimes used to designate the gray substance of the restiform bodies. **N., Cleavage-**. Synonym of *N., Segmentation-*. **N., Daughter-**. See *Mitosis*. **N. emboliformis**, a small mass of gray matter situated in the interval between the N. dentatus and N. fastigii, and lying nearer the former. **N. fastigii**, a flat expanse of gray matter on each side of the inferior vermiform process of the cerebellum directly over the roof of the fourth ventricle. **N. fimbriatus**. Synonym of *Corpus dentatum*. **N. funiculi cuneati**, a mass of gray matter of the posterior column of the medulla, lying beneath the funiculus cuneatus. **N. funiculi gracilis**, an elongated club-shaped mass of gray matter in the mesal portion of the posterior column of the medulla. **N., Germinal**, the N. resulting from the union of the male and female pronuclei. **N., Gingival**, a part of the cerebellum in the fetus (between the third and fourth months), which bears some resemblance to the gums. **N., Globic**, **N. globosus**, the globulus, a number of small round or oval masses of gray matter situated in the interval between the N. dentatus and N. fastigii, lying near to, and probably an accessory detachment of the latter. **N. gracilis**, a column of gray matter in the posterior pyramid of the medulla oblongata. **N., Gray**, the gray matter of the spinal cord. **N., Intraventricular**, the caudate nucleus. **N., Kölliker's**, the gray matter surrounding the canal of the spinal cord. **N., Laryngeal**, the nucleus of origin of the nerve-fibers of the larynx. **N., Lenticular**, **N. lentiformis**, the extraventricular portion of the corpus striatum, lying between the internal and external capsules. **N. of Luys**, an almond-shaped mass of gray matter with pigmented ganglion-cells in the subthalamic region. **N. magnocellularis**, Deiters' nucleus; a nucleus of gray matter in the medulla oblongata, situated mesad of the restiform column at the level of entry of the auditory nerve-roots. **N., Mother-**. See *Mother-star*, under *Karyokinesis*. **N., Motor**, a collection of nerve-cells in the central nervous system giving origin to a motor nerve. **N., Oculomotor**, the nucleus of the oculomotor nerve, lying

dorsad of the posterior longitudinal bundle, under the aqueduct of Sylvius. **N., Olivary**, a folded mass of gray matter in the medulla oblongata, producing a swelling on the surface, the olivary body. **N. of Origin**, the collection of ganglion-cells in the central nervous system giving origin to a nerve. **N. pyramidalis**, the inner accessory olivary nucleus. **N., Red**. See *N. tegmenti*. **N., Restiform**, the gray matter of the restiform body. **N., Segmentation-**, the N. that appears shortly after the fusion of the male and female pronuclei; the last step in the process of fertilization; it is so called because within it cleavage is first established. **N., Spermatic**. Synonym of the male pronucleus. **N., Still-ing's**. Synonym of *N. tegmenti*. **N., Subependymal**, the internal nucleus of the eighth or auditory nerve, lying just beneath the ependyma of the fourth ventricle. **N., Tegmen-tal**, **N. tegmenti**, red N.; a mass of reticular substance in the tegmentum of the crus cerebri, to the inner side of the substantia nigra. **N., Vesicular**, a rather large cell-nucleus, the membrane of which stains deeply, while the central portion remains relatively pale. **N. vestibularis**. See *N. of Bechterew*. **N., Vitelline**, a nucleus resulting from the fusion of the male and female pronuclei within the vitellus. **N., West-phal's**, the N. of origin of a part of the trochlear fibers; it is situated posteriorly to the trochlear nucleus proper. **N., White**, the white substance of the dentate body of the olive.

Nuel's Space. A triangular space between the outer hair-cells and the outer rods of Corti of the internal ear.

Nullipara (*nul-ip'-ar-ah*) [*nullus*, none; *parere*, to bring forth]. A woman who has never borne a child.

Nulliparity (*nul-ip-ar'-it-e*) [*nullus*, none; *parere*, to bring forth]. The condition of being nulliparous.

Nulliparous (*nul-ip'-ar-us*) [*nullus*, none; *parere*, to bring forth]. Having never borne children.

Nummular (*num'-u-lar*) [*nummus*, coin]. Resembling a coin in form, as N. sputum; resembling rouleaux, or rolls of coin.

Nurse's Contracture. Tetany occurring in nursing women.

Nutation (*nu-ta'-shun*) [*nutare*, to nod]. Nodding or oscillation. **N. of Sacrum**, a partial rotation of the sacrum on its transverse axis, whereby the distance between the upper extremity or the lower extremity and the anterior pelvic wall is increased.

Nutmeg [AS., *hnutu*, nut; Pers., *musk*, musk]. The seed of various species of *Myristica*, of the order *Myristicaceae*. *Myris-*

tica (U. S. P., B. P.) is the seed of *M. fragrans*, and is the source of a volatile oil (*Oleum myristicæ*, U. S. P., B. P.). The latter contains a stearopten, myristin, which is the glycerid of myristic acid, $C_{18}H_{38}O_2$. N. is employed as a condiment, as a corrective and mild flavoring agent, and has also slight narcotic properties. Dose, gr. v-xx (0.32-1.3). *Oleum myristicæ*. Dose 2 to 3 drops (0.13-0.2). *Oleum myristicæ expressum* (B. P.) is also known as oil of mace.

Nutrient (*nu'-tre-ent*) [*nutrire*, to nourish]. 1. Affording nutrition. 2. A substance that nourishes; a food.

Nutrimet (*nu'-trim-ent*) [*nutrire*, to nourish]. Anything that nourishes.

Nutrition (*nu'-trish'-un*) [*nutrire*, to nourish]. The process by which tissue is built up and waste repaired.

Nux vomica (*nuks vom'-ik-ah*) [L., pestilential or vomiting nut]. The seed of *Strychnos Nux vomica*, an Indian tree of the order *Loganiaceæ*. It contains several alkaloids, the most important being strychnin and brucin (*q. v.*), which are united with a peculiar acid called igasuric acid. In small doses it is a bitter tonic, stimulating gastric digestion; it raises blood-pressure by stimulating the heart and the vasomotor center, and stimulates the respiratory center, and the motor centers of the spinal cord. In overdoses it produces tetanic convulsions, and risus sardonicus; the reflex excitability is enormously increased, and the slightest stimulus serves to bring on a convulsion. Death usually occurs from asphyxia (cramp-asphyxia), more rarely from exhaustion. *Nux vomica*, or strychnin, is employed in dyspepsia, in convalescence from acute diseases, in acute infectious diseases, in shock, in poisoning by chloroform and opium, in emphysema, phthisis, and other conditions associated with dyspnea, in chronic bronchitis, in constipation, in atony of the bladder, in lead-palsy, and in amaurosis from tobacco or alcohol. Dose of *nux vomica* gr. j-v (0.065-0.32). *Extractum nucis vomica*. Dose gr. $\frac{1}{6}$ - $\frac{1}{4}$ (0.01-0.016). *Extractum nucis vomicae fluidum*. Dose \mathfrak{m} j-v (0.065-0.32). *Tinctura nucis vomicae*. Dose \mathfrak{m} v-xxx (0.32-2.0).

Nyctalopia (*nik-tal'-o'-pe-ah*) [*nyx*, night; *ōps*, the eye]. 1. Night-vision; the condition in which the sight is better by night or in semidarkness than by daylight. It is a symptom of central scotoma, the more dilated pupil at night allowing a better illumination of the peripheral portions of the retina. 2. According to the usage of some writers, this term designates night-blindness. See *Hemeralopia*.

Nympha (*nim'-fah*) [*νύμφα*, nymph]. A labium minus of the vulva.

Nymphomania (*nim-fo-ma'-ne-ak*) [*νύμφη*, nymph; *μανία*, madness]. Excessive sexual desire on the part of a woman.

Nymphomaniac (*nim-fo-ma'-ne-ak*) [*νύμφη*, nymph; *μανία*, madness]. One affected with nymphomania.

Nystagmus (*ni-stag'-mus*) [*νυστάζειν*, to nod in sleep]. An oscillatory movement

of the eyeballs. It may be congenital or dependent on intracranial disease, especially meningitis, on multiple sclerosis, etc. **N.**, Lateral, oscillation of the eyes in the horizontal meridian. **N.**, Rotatory, an oscillatory, partial rolling of the eyeball around the visual axis. **N.**, Vertical, oscillatory movement in the vertical meridian.

O

O. The symbol of Oxygen; of *oculus*, eye; of *octarius*, a pint.

O. D. Abbreviation for *oculus dexter*, right eye.

Oak (*ak*) [AS., *ac*, oak]. A genus of trees, *Quercus*, of the order *Cupuliferæ*. The bark of *Quercus alba* (U. S. P., *Quercus cortex* and of *Quercus robur*, B. P.) and other species, contains a peculiar tannic acid known as quercitannic acid, and a bitter principle, quercin, and is used as an astringent tonic, especially in the form of the oak-bath. It has also been employed in leukorrhea, hemorrhoids, and prolapse of the rectum. **O.-bark** (*Quercus alba*, U. S. P.; *Quercus cortex*, B. P.) is an astringent tonic containing tannic acid, and is mostly employed in washes and decoctions for external use.

Oakum (*o'-kum*) [AS., *acumba*, tow]. A material made by picking old rope into pieces. It has been used as a dressing for wounds, and in the form of pads to absorb lochial discharges.

Oat (*ot*) [AS., *atas*, oats]. A cereal plant, *Avena sativa*, or other species of *Avena*, and its seed. **O.-meal**, the meal made from oats. It is used in the form of a gruel, as a food, as a demulcent and laxative, and as an emollient poultice.

Ob- [L.]. A prefix signifying on, against, in front of, or toward.

Obdormition (*ob-dor-mish'-un*) [*obdormitio*; *obdormire*, to fall asleep]. Numbness of a part due to interference with nervous function; the state of a part when it is "asleep."

Obduction (*ob-duk'-shun*) [*ob*, against; *ducere*, to lead]. A necropsy.

O'Beirne's Tube. A long, flexible tube used in making rectal injections.

Obelion (*o-be'-le-on*) [*ὀβελός*, a spit]. See *Cranimetric Points*.

Obermaier, Spirillum of. A spirillum found in the blood in relapsing fever. See *Bacteria*, Table of.

Obesity (*o-bes'-it-e*) [*obesitas*, from; *obedere*, to eat]. An excessive development of fat throughout the body; corpulence; polysarcia.

Obex (*o'-beks*) [L., a barrier]. A band of white nervous matter at the point of the calamus scriptorius.

Obfuscation (*ob-fus-ka'-shun*) [*ob*, in front of; *fuscus*, dusky]. The act of darkening or clouding, as O. of the cornea.

Object-glass (*ob'-jekt-glas*). See *Objective* (3d definition).

Objective (*ob-jekt'-tiv*) [*ob*, against; *jacere*, to throw]. 1. Pertaining to an object or to that which is contemplated or perceived as distinguished from that which contemplates or perceives. 2. Pertaining to those relations and conditions of the body perceived by another, as O. signs of disease. 3. The lens of a microscope nearest the object.

Obligate (*ob'-lig-at*) [*obligare*, to bind]. Constrained, bound. **O. Aerobic**, of a microorganism, one that can only live as an aerobe.

O. Anaerobic, of a microorganism, one that can only live as an anaerobe. **O. Parasite**, a parasite that can only live as a parasite.

Oblique (*ob-lek' or ob-lik'*) [*obliquus*, slanting]. Slanting.

Obliquity (*ob-lik'-wit-e*) [*obliquus*, slanting]. The state of being oblique.

Obliteration (*ob-lit-er-a'-shun*) [*obliterare*, to efface, from *ob*, over; *littera*, a letter]. Removal of a part; extirpation; complete closure of a lumen.

Obsession (*ob-sesh'-un*) [*ob*, over; *sedere*, to sit]. Possession by evil spirits.

Obsolescence (*ob-so-les'-ens*) [*obsolescere*, to grow old]. The state of becoming old or going out of use.

Obstetric (*ob-stet'-rik*) [*obstare*, to stand before]. Pertaining to the practice of obstetrics.

Obstetrician (*ob-stet-rish'-an*) [*obstare*, to stand before]. One who practices obstetrics.

Obstetrics (*ob-stet'-riks*) [*obstare*, to stand before]. The branch of medicine that deals with the care of women in pregnancy, labor, and the puerperium.

Obstipation (*ob-stip-a'-shun*) [*obstipare*, to close tightly]. Constipation.

Obstruction (*ob-struk'-shun*) [*ob*, over against; *struere*, to build]. 1. The state of being stopped or blocked up. 2. The act of stopping up. 3. An impediment or obstacle.

Obstructive (*ob-struk'-tiv*) [*ob*, over against; *struere*, to build]. 1. Stopping or blocking up. 2. Due to an obstruction, as O. jaundice.

Obtund (*ob-tund'*) [*ob*, against, upon; *tundere*, to beat]. To blunt or dull; to lessen, as to O. sensibility.

Obturator (*ob'-tu-ra-tor*) [*obturare*, to stop up]. 1. Closing an opening. 2. That which closes an opening. 3. Pertaining to the obturator membrane, muscles, etc. **O. Foramen**, a foramen in the anterior part of the os innominatum. See *Foramina, Table of*. **O. Membrane**, the membrane closing the obturator foramen. **O. Muscle**. See *Muscles, Table of*. **O. Nerve**. See *Nerves, Table of*.

Occipital (*ok-sip'-it-al*) [*occiput*]. Pertaining to or in relation with the occiput. **O. Artery**. See *Arteries, Table of*. **O. Lobe**. See *Lobe*. **O. Nerve**. See *Nerves, Table of*. **O. Protuberance**, the prominence on the inner surface (internal) or on the outer surface (external) of the occipital bone. **O. Triangle**. See *Triangles, Table of*.

Occipito- (*ok-sip'-it-o-*) [*occiput*, occiput]. A prefix denoting connection with or relation to the occipital bone or the occiput.

Occipitoanterior (*ok-sip'-it-o-an-te'-re-or*) [*occiput*; *anterior*]. Having the occiput directed toward the front, as the O. position of the fetus in utero.

Occipitoatloid (*ok-sip'-it-o-at'-loid*) [*occiput*; *atlas*; *εἶδος*, form]. Pertaining to the occipital bone and the atlas.

Occipitoaxoid (*ok-sip'-it-o-aks'-oid*) [*occiput*; *axis*; *εἶδος*, form]. Pertaining to the occipital bone and the axis.

Occipitobregmatic (*ok-sip'-it-o-breg-mat'-ik*) [*occiput*; *bregma*]. Pertaining to the occiput and the bregma.

Occipitofrontal (*ok-sip'-it-o-fron'-tal*) [*occiput*; *frons*, forehead]. Pertaining to the occiput and forehead, or to the O. muscle (*Occipitofrontalis*).

Occipitomastoid (*ok-sip'-it-o-mas'-toid*) [*occiput*; *μαστός*, nipple; *εἶδος*, form]. Pertaining to the occipital bone and the mastoid process.

Occipitomenal (*ok-sip'-it-o-men'-tal*) [*occiput*; *mentum*, the chin]. Pertaining to the occiput and the chin.

Occipitoparietal (*ok-sip'-it-o-par'-i'-et-al*) [*occiput*; *paries*, wall]. Pertaining to the occipital and parietal bones, or to the occipital and parietal lobes of the brain.

Occipitoposterior (*ok-sip'-it-o-pos-te'-re-or*) [*occiput*; *posterior*]. Having the occiput directed backward, as the O. position of the fetus in utero.

Occipitotemporal (*ok-sip'-it-o-tem'-per-al*) [*occiput*; *tempus*, temple]. Pertaining to the occipital and temporal bones.

Occiput (*ok'-sip-ut*) [*ob*, against; *caput*, the head]. The back part of the head.

Occlusion (*ok-lu'-shun*) [*ob*, against; *claudere*, to shut]. 1. A closing or shutting up. 2. The state of being closed or shut. 3. The absorption by a metal of gas in large quantities, as of hydrogen by platinum.

Occlusive (*ok-lu'-siv*) [*ob*, against; *claudere*, to shut]. Closing or shutting up; as an occlusive surgical dressing.

Occupation-disease. One caused by the occupation of the patient. The most common variety, the *Occupation-neurosis*, is a functional disturbance of the part used in carrying on a certain occupation, as, e.g., writer's cramp, telegrapher's cramp, etc.

Ochlesis (*ok-le'-sis*) [*ὄχλος*, crowd]. Any disease due to over-crowding.

Octan (*ok'-tan*) [*octo*, eight]. Returning every eighth day, as an octan fever.

Octarius (*ok-ta'-re-us*) [*octo*, eight]. An eighth part of a gallon; a pint. Symbol O.

Octavipara (*ok-tav-ip'-ar-ah*). See *Multi-para*.

Ocular (*ok'-u-lar*) [*oculus*, the eye]. 1. Pertaining to or in relation with the eye. 2. The lens of a microscope that is turned toward the eye. **O., Compensating**, one that compensates for axial aberrations of the objective.

Oculist (*ok'-u-list*). Synonym of *Ophthalmologist*.

Oculo- (*ok'-u-lo-*) [*oculus*, eye]. A prefix signifying pertaining to the eye.

Oculomotor (*ok-u-lo-mo'-tor*) [*oculus*, eye; *movere*, to move]. 1. Pertaining to the movement of the eye, as the oculomotor nerve. 2. Pertaining to the oculomotor nerve, as the oculomotor nucleus.

Oculus (*ok'-u-lus*) [L.]. An eye.

Oculozygomatic (*ok-u-lo-zi-go-mat'-ik*) [*oculus*, eye; *zygoma*]. Pertaining to the eye and the zygoma. See *Jadot's Lines*.

Odontagra (*o-don-ta'-grah*) [*ὀδούς*, tooth; *ἄγρα*, seizure]. Toothache, especially a form due to gout.

Odontalgia (*o-don-tal'-je-ah*) [*ὀδούς*, tooth; *ἄλγος*, pain]. Toothache.

Odontiasis (*o-don-ti'-as-is*) [*ὀδούς*, tooth]. Dentition; the cutting of teeth.

Odontitis (*o-don-ti'-tis*) [*ὀδούς*, tooth; *τις*, inflammation]. Inflammation of the teeth.

Odonto- (*o-don'-to-*) [*ὀδούς*, tooth]. A prefix signifying pertaining to a tooth.

Odontoblast (*o-don'-to-blast*) [*ὀδούς*, tooth; *βλαστός*, germ]. One of the cells covering the dental papilla and forming the dentine.

Odontogeny (*o-don-toj'-en-e*) [*ὀδούς*, tooth; *γενᾶν*, to beget]. The origin and development of teeth.

Odontography (*o-don-tog'-ra-fe*) [*ὀδούς*, tooth; *γράφειν*, to write]. The descriptive anatomy of the teeth.

Odontoid (*o-don'-toid*) [*ὀδούς*, tooth; *εἶδος*, like]. Resembling a tooth. **O. Ligament.** See *Ligament*. **O. Process**, the dentate process of the second cervical vertebra.

Odontolith (*o-don'-to-lith*) [*ὀδούς*, tooth; *λίθος*, a stone]. The calcareous accretion on the teeth, popularly known as tartar.

Odontology (*o-don-tol'-o-je*) [*ὀδούς*, tooth; *λόγος*, science]. The branch of knowledge dealing with the anatomy and diseases of the teeth.

Odontoma (*o-don-to'-mah*) [*ὀδούς*, tooth; *μα*, tumor]. A tumor arising in connection with the teeth and presenting a tooth-like structure.

Odontonecrosis (*o-don-to-nek-ro'-sis*) [*ὀδούς*, tooth; *νεκρός*, corpse]. Necrosis or decay of the tissues of the teeth.

Odontopathy (*o-don-top'-ath-e*) [*ὀδούς*, tooth; *πάθος*, suffering]. Any disease of the teeth.

Odontorthosia (*o-don-tor-tho'-ze-ah*) [*ὀδούς*, tooth; *ὀρθός*, straight]. The operation of straightening irregularly growing teeth.

Odontotherapy (*o-don-to-ther'-ap-e*) [*ὀδούς*, tooth; *θεραπεία*, treatment]. The treatment of diseases of the teeth.

Odoriferous (*o-dor-if'-er-us*) [*odor*, odor; *ferre*, to carry]. Fragrant.

O'Dwyer's Tubes (*o-dwi'-era*) [*O'Dwyer*, an American physician]. Tubes used for intubation of the larynx.

-odyne, -odynia (*o-din, -o-din'-e-ah*) [*ὀδύνη*, pain]. A suffix denoting pain.

Edema, Edematous (*e-de'-mah, e-dem'-at-us*). See *Edema*, *Edematous*.

Oese [Ger.]. An instrument consisting of a platinum-wire affixed to a glass handle and employed in bacteriologic investigation.

Esophagism (*e-sof'-aj-izm*). See *Esophagismus*.

Esophagus (*e-sof'-ag-us*). See *Esophagus*.

Official (*off-ish'-al*) [*officium*, duty, service]. Of medicines, sanctioned by the recognized authority, *i. e.*, the Pharmacopeia.

Offical (*of-is'-in-al*) [*officina*, a workshop]. For sale in the shops; kept on sale in apothecaries' shops.

Ohm (*om*) [Dr. G. S. Ohm, a German phy-

sicist]. The unit of electric resistance. The ohm adopted as a standard varies; the British Association O. is the resistance of a column of mercury 1 square millimeter in section and 1.049318 meters long. The legal O. is similar to the B. A. O., except that the column of mercury is 1.06 meters in length. The international O. adopted 1893, is the resistance of a column of mercury 1.063 centimeters long and weighing 14.4521 grams.

Oidium (*o-id'-e-um*) [dim. of *ὄν*, egg]. A genus of parasitic fungi. **O. albicans**, the thrush-fungus.

Oikoid (*oi'-koid*) [*οἶκος*, house; *εἶδος*, like]. The stroma of red corpuscles.

Oil [*oleum*, from *ἐλαίον*, oil]. A liquid of animal or vegetal, sometimes of mineral, origin, having a peculiar feel, and not miscible with water. **Animal and Vegetal Oils** are either volatile or fixed. **Oil of Amber.** See *Succinum*. **O. of Cade**, an oil derived from the wood of *Juniperus communis*, and used in the treatment of diseases of the skin. **O., Camphorated**, a solution of camphor in olive-oil. **O. of Caraway**, a volatile oil from the fruit of *Carum carui*. It is a carminative and flavoring agent. Dose m. j- ($0.065-0.32$). See *Carum*. **O., Carron-**, an application to burns, consisting of equal or nearly equal parts of linseed oil and lime-water. It is named after the Carron iron-works in Scotland, where it was first employed. **O., Essential**, a volatile oil, so called because it contains the essence or active principle of a plant. **O., Ethereal**, heavy oil of wine. See *Oleum athericum*. **O., Fatty**, fatty oils are salt-like bodies, composed of characteristic acids (oleic, palmitic, and stearic), known as fatty acids, and a base. In most cases the base is glyceryl, the radicle of the triatomic alcohol, glycerol, so that the oils are said to be glycerids of the several fatty acids. See *O., Fixed*. **O., Fixed**, one not volatilizing on the application of heat. Fixed oils are also called fatty oils, because they in part constitute the animal and vegetable fats. Some are liquid, as olive-oil, cotton-seed oil, linseed-oil, castor-oil, etc., others are solid, as tallow and beeswax, which chemically belong to the group of oils. See *O., Fatty*. **O., Heavy** (*of wine*), the product obtained when alcohol is treated with an excess of sulphuric acid. See *Oleum athericum*. **O., Mineral**, petroleum and certain of its derivatives. **O. of Neroli**, an essential oil from the flowers of bitter orange, used as a perfume. **O., Rock-**, petroleum. **O., Straits-**, the first oil obtained by the exposure of the livers of cod-fish to the sun in casks, on board the ships. **O., Sweet.** See *Olive*. **O., Volatile, O., Essential**, one which is

volatile at ordinary temperatures. Volatile oils are odoriferous, and are generally obtained by distillation.

Oinomania (*oi-no-ma'-ne-ah*) [*οἶνος*, wine; *μανία*, madness]. 1. A form of insanity characterized by an irresistible craving for, and consequent indulgence in, drink. 2. Delirium tremens.

Ointment (*oint'-ment*) [*unguentum*]. A fatty material of the consistence of butter, generally impregnated with a medicinal substance, and used for application to the skin. **O.**, Maury's, one composed of one dram of mercury nitrate and half a dram each of powdered opium and rhubarb to an ounce of simple ointment. It is used as an application to ulcers.

Oleate (*ol'-le-at*) [*oleum*, oil]. 1. A salt of oleic acid. 2. A mixture of oleic acid with certain medicinal principles.

Olecranon (*o-lek'-ran-on*) [*ὀλῆνη*, elbow; *κρῆνιον*, skull]. The large concave process at the upper extremity of the ulna.

Olefiant Gas (*o-lef'-e-ant*) [*oleum*, oil; *facere*, to make]. See *Ethylene*.

Olefin (*ol'-lef-in*) [*oleum*, oil; *facere*, to make]. Olefiant gas; also any one of a series of unsaturated hydrocarbons having the formula C_nH_{2n} .

Oleic Acid (*o-le'-ik*). See *Acid*.

Olein (*ol'-le-in*) [*oleum*, oil], $C_{55}H_{104}O_8$. A neutral fat, glyceryl trioleate, occurring in olive oil, butter, and other animal and vegetable fats. It is a colorless oil with a faint sweetish taste, insoluble in water, readily soluble in alcohol and ether.

Oleo- (*ol'-le-o*) [*oleum*, oil]. A prefix to denote connection with or relation to an oil.

Oleobalsamic Mixture (*o-le-o-bal-sam'-ik*). *Mistura oleobalsamica*. A mixture of the oils of lavender, thyme, lemon, mace, orange-flowers, cloves, and cinnamon, with balsam of Peru and alcohol. It is used as a nervine.

Oleocresote (*o-le-o-cres'-o-sot*) [*oleum*, oil; *cresote*]. A yellowish neutral liquid composed of cresote, 33 per cent., and oleic acid. It is used in bronchial and pulmonary diseases. Dose $\text{m}x-xv$ (0.65-1.0).

Oleomargarin (*o-le-o-mar'-gar-in*) [*oleum*, oil; *μάργαρος*, the pearl-oyster]. An artificial butter made by removing the excess of stearin from tallow or suet.

Oleoresin (*o-le-o-res'-in*) [*oleum*, oil; *resina*, resin]. A substance consisting chiefly of a mixture of an essential oil and a resin extracted from plants with ether.

Oleum (*ol'-le-um*) [L.]. See *Oil*. **O.** æthereum, a volatile, yellowish liquid consisting of equal volumes of heavy oil of wine and ether. Heavy oil of wine is produced when alcohol and sulphuric acid are distilled, and is a mixture of ethyl sulphate, ethyl sulphite,

and several polymeric forms of ethylene. **O.** fixum, a fixed oil. See *Oil*.

Olfactometer (*ol-fak-tom'-et-er*) [*olfacere*, to smell; *μέτρον*, measure]. An instrument for determining the power of smell.

Olfactory (*ol-fak'-to-re*) [*olfacere*, to smell]. Pertaining to the sense of smell. **O.** Bulb. See *Bulb*.

O. Cells, the cells of the nasal fossæ forming the peripheral end-organs of the olfactory nerve.

O. Center, the cerebral center for the sense of smell, supposed to be in the hippocampal gyrus.

O. Groove. See *O. Sulcus*.

O. Lobe, the olfactory tubercle, olfactory tract, and olfactory bulb considered together.

O. Nerve. See *Nerves*, *Table of*.

O. Region, the area of distribution of the olfactory nerve in the upper part of the nose.

O. Sulcus, the furrow for the olfactory tract and bulb on the cribriform plate of the ethmoid bone, and on the orbital surface of the cerebral hemispheres.

O. Tract, the central portion of the olfactory lobe terminating anteriorly in the olfactory bulb and posteriorly in the olfactory tubercle.

O. Vesicle, a part budded off from the cerebral vesicle and forming the olfactory lobe.

Olibanum (*o-lib'-an-um*) [Ar., *al-luban*, frankincense]. Frankincense, a gum-resin produced by various species of *Boswellia*. It has been used as a substitute for the balsams of Peru and tolu, as an inhalation in laryngeal and bronchial inflammations, for fumigation, and in plasters.

Oligæmia (*ol-ig'-e'-me-ah*). See *Oligemia*.

Oligemia (*ol-ig'-e'-me-ah*) [*ὀλίγος*, few; *αἷμα*, blood]. A state in which the total quantity of the blood is diminished.

Oligo- (*ol'-ig-o*) [*ὀλίγος*, few]. A prefix signifying want or deficiency.

Oligochromemia (*ol-ig-o-kro'-me'-me-ah*) [*ὀλίγος*, few; *χρῶμα*, color; *αἷμα*, blood]. Deficiency of hemoglobin in the blood.

Oligocythemia (*ol-ig-o-si-the'-me-ah*) [*ὀλίγος*, few; *κύτος*, cell; *αἷμα*, blood]. A deficiency of red corpuscles in the blood.

Oligohydramnios (*ol-ig-o-hy-dram'-ne-oi*) [*ὀλίγος*, few; *ὕδωρ*, water; *ἄμνιον*, the amnion]. A deficiency in the quantity of the amniotic fluid.

Oligomania (*ol-ig-o-ma'-ne-ah*) [*ὀλίγος*, few; *μανία*, madness]. Insanity in which only a few of the mental faculties are deranged.

Oligoplasmia (*ol-ig-o-plas'-me-ah*) [*ὀλίγος*, scanty; *πλάσσειν*, to mold]. A deficient amount of plasma in the blood.

Oligospermia (*ol-ig-o-sper'-me-ah*) [*ὀλίγος*, few; *σπέρμα*, seed]. A deficiency in the secretion of semen.

Oligozoospermia (*ol-ig-o-zo-o-sper'-me-ah*) [*ὀλίγος*, few; *ζῷον*, animal; *σπέρμα*, seed]. Deficiency of the spermatozoa in the spermatogenic fluid.

Oliguria (*ol-ig-u'-re-ah*) [*ὀλίγος*, scanty; *urina*, urine]. A diminution in the quantity of urine excreted.

Oligydria (*ol-ig-id'-re-ah*) [*ὀλίγος*, scanty; *ιδρώς*, sweat]. A deficiency in the sweat.

Olivary (*ol'-iv-a-re*) [*oliva*, olive]. Resembling an olive in shape. **O. Body**, an oval mass of gray matter situated behind the anterior pyramid of the medulla. **O. Fasciculus**. See *Fillet, Olivary*. **O. Nucleus**, the corpus dentatum or central gray matter of the olivary body. **O. Peduncle**, the mass of fibers entering the hilus of the olivary body. **O. Process**. See *Process*.

Olive (*ol'-iv*) [*oliva*, olive]. 1. The olive-tree, *Oliva europæa*, of the natural order Oleaceæ, and its fruit. The fixed oil expressed from the fruit is the *Oleum olivæ* of the U. S. P. and B. P. It consists chiefly of olein and palmitin, and is used as a food and condiment; in medicine as a laxative; in the treatment of gall-stones; as an anthelmintic; as an emollient external application to wounds, burns, etc.; and as an ingredient of liniments, ointments, and plasters. 2. The olivary body.

-oma (*-o'-mah*) [*ὄμα*]. A termination denoting a tumor.

Omagra (*om'-a-grah*) [*ὄμος*, shoulder; *ἄγρα*, seizure]. Gout in the shoulder.

Malgia (*o-mal'-je-ah*) [*ὄμος*, shoulder; *ἄλγος*, pain]. Pain in the shoulder.

Omarthritis (*o-mar-thrit'-tis*) [*ὄμος*, shoulder; *ἀρθρον*, joint; *τις*, inflammation]. Inflammation of the shoulder-joint.

Omental (*o-men'-tal*) [*omentum*, the caul]. Pertaining to the omentum. **O. Hernia**. See *Epiplotele*.

Omentum (*o-men'-tum*) [L.]. A fold of the peritoneum connecting the abdominal viscera with the stomach. **O., Gastrocolic**, **O., Great**, a fold of peritoneum attached to the greater curvature of the stomach above and after dipping down over the intestines returning to enclose the transverse colon. Between the ascending and descending folds is the cavity of the great omentum. **O., Gastrohepatic**, **O., Lesser**, a double fold of peritoneum passing from the lesser curvature of the stomach to the transverse fissure of the liver. On the left side it includes the esophagus; on the right its edges are free and enclose all the structures issuing from or entering the transverse fissure of the liver: the hepatic vessels and nerves, and the bile-duct. Behind it is the foramen of Winslow. **O., Gastrosplenic**, the fold of peritoneum passing from the stomach to the spleen.

Omnivorous (*om-niv'-o-rus*) [*omnis*, all; *vorare*, to devour]. Subsisting on all kinds of food.

Omo- (*o'-mo-*) [*ὄμος*, shoulder]. A prefix

denoting connection with or relation to the scapula or shoulder.

Omodynia (*o-mo-din'-e-ah*) [*ὄμος*, shoulder; *ὀδύνη*, pain]. Pain in the shoulder.

Omothyoid (*o-mo-hi'-oid*) [*ὄμος*, shoulder; *ἰσοειδής*, like the Greek letter upsilon]. Pertaining conjointly to the scapula and the hyoid bone. **O. Muscle**. See *Muscles, Table of*.

Omphalic (*om-fal'-ik*) [*ὀμφαλός*, navel]. Pertaining to the umbilicus. **O. Duct**, the vitelline duct; the duct connecting the umbilical vesicle with the fetal intestines during the first three months of intrauterine life.

Omphalitis (*om-fal'-i'-tis*) [*ὀμφαλός*, navel; *τις*, inflammation]. Inflammation of the navel.

Omphalo- (*om-fal'-o-*) [*ὀμφαλός*, navel]. A prefix denoting relation to the umbilicus.

Omphalocele (*om-fal'-o-vel*) [*ὀμφαλός*, navel; *κήλη*, tumor]. Umbilical hernia.

Omphalomesaraic, **Omphalomesenteric** (*om-fal-o-mez-ar-a'-ik*, *om-fal-o-mez-en-ter'-ik*) [*ὀμφαλός*, navel; *mesentery*]. Pertaining conjointly to the umbilicus and the mesentery. **O. Arteries**. See *Arteries, Table of*.

O. Duct, a duct connecting the intestinal canal of the embryo with the umbilical vesicle.

Omphalopagus (*om-fal-op'-ag-us*) [*ὀμφαλός*, navel; *πηγνύναι*, to make fast]. A double monster united at the umbilicus.

Omphalophlebitis (*om-fal-o-fleb-i'-tis*) [*ὀμφαλός*, navel; *φλέψ*, vein; *τις*, inflammation]. Inflammation of the umbilical vein.

Omphalorrhagia (*om-fal-or-a'-je-ah*) [*ὀμφαλός*, navel; *ρηνύναι*, to burst forth]. Hemorrhage from the umbilicus.

Omphalos (*om'-fal-os*) [*ὀμφαλός*, navel]. The umbilicus.

Omphalosite (*om-fal'-o-sit*) [*ὀμφαλός*, navel; *σιτος*, nourishment]. A single monster, which, lacking the heart, receives its blood-supply through the umbilical vessels, and is, therefore, incapable of extrauterine existence.

Omphalotomy (*om-fal-ot'-o-me*) [*ὀμφαλός*, navel; *τομή*, a cutting]. The cutting of the umbilical cord.

Onanism (*o'-nan-ism*) [from *Onan*, the son of Judah]. 1. Incomplete coitus. 2. Masturbation.

Oncograph (*ong'-ko-graf*) [*ὄγκος*, swelling; *γράφειν*, to record]. An instrument registering the changes of volume of an organ placed in an oncometer.

Oncology (*ong-kol'-o-je*) [*ὄγκος*, tumor; *λογία*, science]. The branch of surgery and pathology relating to tumors.

Oncometer (*ong-koni'-et-er*) [*ὄγκος*, mass; *μετρον*, measure]. An instrument for measuring variations in the volume of an organ, especially of the kidney or spleen.

Oncotomy (*ong-kol'-o-me*) [*ὄγκος*, tumor; *τομή*, a cutting]. The operation of incising a tumor or other swelling.

Onion (*un'-yun*) [*L.*, *unia*, an onion]. The *Allium cepa* and its bulb. The latter contains an oil resembling oil of garlic, and consisting largely of allyl sulphid (C_3H_5S). The onion is diuretic, expectorant, and rubefacient, and is at times used in dropsy, bronchitis, etc.; locally, as an emollient poultice.

Ontogenesis, Ontogeny (*on-to-jen'-es-is*, *on-toj'-en-e*) [*ὄν*, *ὄντος*, existing; *γεννᾶν*, to beget]. The development of the individual organism. See also *Phylogeny*.

Onychatrophia (*on-ik-at-ro'-fe-ah*) [*ὄνυξ*, nail; *ἀτροφία*, atrophy]. Atrophy of the nails.

Onychauxis (*on-ik-awks'-is*) [*ὄνυξ*, the nail; *αἰξή*, increase]. Hypertrophy of the nail.

Onychia, Onychitis (*on-ik'-e-ah*, *on-ik-i'-tis*) [*ὄνυξ*, nail]. Inflammation of the matrix of the nail. **O. maligna**. A form occurring in debilitated persons, and characterized by an unhealthy ulcer in the matrix of the nail, the latter becoming discolored and thrown off. **O. simplex**, **O.** without much ulceration, with loss of the nail and its replacement by a new one.

Onychogryposis (*on-ik-o-gri-po'-sis*) [*ὄνυξ*, nail; *γρίπσις*, curvature]. A thickened, ridged, and curved condition of the nail.

Onychomycosis (*on-ik-o-mi-ko'-sis*) [*ὄνυξ*, nail; *μύκης*, fungus]. A disease of the nails due to parasitic fungi, as the trichophyton, achorion, etc.

Onychophagy (*on-ik-off'-aj-e*) [*ὄνυξ*, nail; *φαγεῖν*, to eat]. The practice of biting the nails.

Onychosis (*on-ik-o'-sis*) [*ὄνυξ*, nail]. Any disease of the nails.

Onyx (*on'-iks*) [*ὄνυξ*, nail]. 1. A nail of the fingers or toes. 2. A collection of pus between the corneal lamellæ at the most dependent part.

Onyxitis (*on-iks-i'-tis*). Onychia.

Ooblast (*o'-o-blast*) [*ὄβιον*, an egg; *βλαστός*, a germ]. A cell of the germinal epithelium giving rise to an ovum.

Oophorectomy (*o-off-or-ek'-to-me*) [*ὄβιον*, egg; *φέρειν*, to bear; *ἐκτομή*, excision]. Excision of the ovary.

Oophoritis (*o-off-or-i'-tis*) [*ὄβιον*, egg; *φέρειν*, to bear; *ιτις*, inflammation]. Inflammation of the ovary.

Oophoromania (*o-off-or-o-ma'-ne-ah*) [*ὄβιον*, egg; *φέρειν*, to bear; *μανία*, mania]. Insanity due to ovarian disorder.

Oophorrhapy (*o-off-or'-a-fe*) [*ὄβιον*, egg; *φέρειν*, to bear; *ραφή*, suture]. The operation of suturing an ovary to the pelvic wall.

Oosperm (*o'-o-sperm*) [*ὄβιον*, egg; *σπέρμα*, seed]. The cell formed by union of the ovum and the spermatozoon.

Opacity (*o-pas'-it-e*) [*opacus*, dull]. 1. The condition of being impervious to light. 2. An opaque spot, as opacity of the cornea or lens.

Opaline (*o'-pal-en*) [*ὀπάλινος*, an opal]. Having the appearance of an opal.

O. Patch, the mucous patch of syphilis, forming, in the mouth, a whitish pellicle.

Open [*AS.*, *open*]. Exposed to the air, as an open wound; interrupted, as an open circuit, one that is interrupted so that the electric current cannot pass.

Operation (*op-er-a'-shun*) [*operatio*, from *operari*, to labor, to do]. 1. Anything done or performed, especially anything done with instruments; a surgical procedure. 2. The mode of action of anything. **O., Capital**, one involving a risk of life. **O., High**. 1. Suprapubic lithotomy. 2. The application of the forceps to the fetal head at the superior strait. **O., Indian**, for rhinoplasty; a flap is taken from the forehead, with its pedicle at the root of the nose; hollow plugs are inserted into the nostrils, and the flap is secured. **O., Italian**, for rhinoplasty; the skin is taken from the arm over the biceps; the flap is cut on three sides, and after the skin has shrunk it is fitted to the fresh margins of the defect, the arm being bandaged in position for at least eight days. **O., Major**, an important and serious operation. **O., Minor**, a comparatively trivial operation. **O., Plastic**, one for the purpose of restoring a lost part or repairing a deformity. **O., Radical**, one removing the cause of the disease or the diseased part itself. **O., Surgical**, one performed by the surgeon by means of the hands or instruments. **O., Tagliacotian**. See *O., Tagliacozzi's*, or *O., Italian*. An eponymic table of the more important operations is appended; for a more extended list see *Gould's Illustrated Dictionary of Medicine*.

EPONYMIC TABLE OF SURGICAL OPERATIONS.

Abbé's, lateral anastomosis of the intestines with cat gut rings. **Adams'**, osteotomy for ankylosis of the hip-joint, the neck of the femur being divided subcutaneously within the capsule. **Alexander's**: 1. For the relief of epilepsy: ligation of the vertebral ar-

tery to diminish the blood-supply to the brain. 2. For prolapse and retroflexion of the uterus: shortening of the round ligaments for the purpose of holding the uterus in its normal position. **Allingham's** (*H.*), for inguinal colotomy: the incision is made parallel with the

outer third of Poupart's ligament, and about one-half inch above. **Amussat's**, lumbar colotomy: a transverse incision is made, crossing the outer border of the quadratus lumborum muscle. **Anel's**: 1. For aneurysm: ligation on the cardiac side close to the aneurysm. 2. For stricture of the lacrimal duct: dilatation by means of a gold probe, followed by the injection of an astringent fluid into the sac. **Annandale's**: 1. For dislocated cartilages: incision into the knee-joint and stitching of the dislocated cartilages into their proper position. 2. For genu valgum: partial excision of both condyles of the femur. **Antyllus'**, for aneurysm: it consists in ligation above and below the sac, followed by opening of the aneurysm and evacuation of its contents. **Appolito's**, enterorrhaphy by means of a form of right-angle continuous suture. **Arlt-Jaesche's**, for distichiasis: the edge of the lid and the contained ciliary bulbs are dissected from the tarsus, a crescentic-shaped piece of skin is removed from the lid above the flap, and the edges of the wound are united, thus transplanting the ciliary bulbs further away from the edge of the lids. **Ball's**: 1. Iliac colotomy, in which the incision is made in the left linea semilunaris, and the bowel secured above and below the future artificial opening by means of two special clamps, which are removed after suturing and opening of the bowel. 2. For inguinal hernia: the sac is separated up to the abdominal portion of the neck, and then twisted around its own axis, after which the fundus is cut away and the stump is secured in the ring. **Barker's**: 1. For excision of the astragalus: the incision runs from just above the tip of the external malleolus forward and a little inward, curving toward the dorsum of the foot. 2. For excision of the hip: by an anterior incision, commencing on the front of the thigh, one-half inch below the anterior superior iliac spine, and running downward and inward for three inches. **Barwell's**, osteotomy for genu valgum: division of the lower end and upper end of the tibia above and below their respective epiphyses. **Bassini's**, for the radical cure of inguinal hernia: the sac is exposed, twisted, and the neck ligated and removed; the spermatic cord is lifted, and the border of the rectus is stitched to the edge of the internal oblique, the transversalis muscle, and the transversalis fascia to Poupart's ligament under the cord. The cord is placed upon this layer and the border of the external oblique is stitched to Poupart's ligament over the cord. **Batley's**, also called normal ovariectomy: a removal of the ovaries for the purpose of inducing the menopause. **Beer's**, for cataract: extraction by the flap-method. **Billroth's**,

pylorectomy: a parietal incision is made in almost a transverse direction; the divided walls of the stomach and bowel are brought together and united by sutures on the side of the greater curvature of the stomach. **Bras-dor's**, for aneurysm: ligation immediately beyond the aneurysm. **Bryant's**, lumbar colotomy: an oblique incision is made midway between the last rib and the iliac crest. The bowel is fixed in position and opened. **Carnochan's**: 1. For elephantiasis: ligation of the main artery of the limb. 2. Removal of the second division of the fifth nerve, together with the sphenopalatine ganglion as far back as the foramen rotundum, for the relief of neuralgia. **Cheyne's**, for the radical cure of femoral hernia: after reducing the hernia, a flap of the pectineus muscle is raised and made to cover the hernial orifice. **Chopart's**, a method of amputation through the foot: the bony structure is divided at the articulation between the astragalus and the calcaneum behind, and the scaphoid and cuboid bones anteriorly. **Cock's**, a method of external urethrotomy: the urethra is opened behind the stricture without a guide, the knife being carried into the median line of the perineum and the incision extended vertically so far as is necessary. **Cooper's**, for ligation of the external iliac artery: an incision four or five inches long is made parallel with Poupart's ligament, and nearly an inch above it, commencing just outside the center of the ligament and extending outward and upward beyond the anterior superior iliac spine. **Cripps'**, iliac colotomy: an imaginary line from the anterior superior iliac spine to the umbilicus is crossed at right angles, $1\frac{1}{2}$ inches from the superior spine, by an incision $2\frac{1}{2}$ inches long. The bowel is fixed in position and opened. **Czerny's**, for the radical cure of inguinal hernia: the sac is exposed and isolated; the neck is tied with a strong catgut-ligature, and cut off below this point; the stump is pushed into the abdominal cavity; the borders of the opening are freshened and united by continuous catgut-sutures. **Dallas'**, for the radical cure of inguinal and femoral hernia: after a transverse incision through the integument a special instrument is introduced to produce abrasion of the hernial canal; the instrument is then withdrawn, and the external wound sealed with iodoform-collodion, and a compress applied. The canal is obliterated by the resulting inflammatory action. **Diefenbach's**, for amputation at the hip-joint: an elastic ligature is applied around the limb, a circular incision is made down to the bone, the vessels are secured and the ligature removed; a knife is inserted two inches above the greater trochanter and the in-

cision is carried down the outer aspect of the bone to meet the circular incision; the joint is then disarticulated. **Dittel's**, for enlarged prostate: enucleation of the lateral lobes of the prostate by an external incision. **Dudley's**, for retroversion of the uterus: the abdomen is opened, and a strip is denuded from the anterior surface of the uterus, and likewise a strip along the inner side of each round ligament, followed by suturing of the three together. **Duplay's**: 1. For epispadias: the urethra is formed at the expense of the corpus spongiosum and corpora cavernosa instead of by flaps. 2. For hypospadias: it is performed in three stages: (1) straightening of the penis and the formation of a meatus; (2) the formation of a canal from the meatus to the hypospadiac opening; (3) junction of the old and new canals. **Dupuytren's**, for amputation at the shoulder-joint: two rounded flaps are taken from the outer and inner aspects of the arm; the outer flap is made first by transfixion from behind at a point two inches below the acromion; the bone is then disarticulated and a short inner flap cut from within outward. **Emmet's**, for lacerated perineum: the crest of the rectocle and the two lower caruncles are drawn together by tenacula, and the tissue thus folded together is denuded as far into the sulci as the folds extend. V-shaped stitches are passed from the center toward the circumference in the sulci, and the perineum is closed by two or three superficial sutures. **Esmarch's**: 1. For amputation at the hip-joint: the soft parts of the thigh are divided to the bone by a single sweep of the knife five inches below the tip of the trochanter; the bone is then sawn across and a second incision is made to join the first from a point two inches above the trochanter, when the bone is shelled out. 2. For ankylosis of the lower jaw: an incision about two inches long is made along the lower border of the jaw, and a wedge-shaped piece of bone is removed from the horizontal portion. **Estlander's**, for empyema: resection of several ribs for the escape of the pus and to allow the chest-wall to come in contact with the lung. **Gant's**, for ankylosis of the hip-joint: division of the shaft of the femur just below the lesser trochanter. **Von Graefe's**, for cataract: extraction of the cataract through a scleral incision, with iridectomy and laceration of the capsule. **Gritti's**, for amputation above the knee-joint: the patella is preserved in a long anterior flap, and, having had a thin slice removed from its deep surface, is secured in apposition with the femur, the latter having been deprived of its articular surface by being sawn through the condyles. **Halsted's**, for the radical cure of

inguinal hernia: the same as Bassini's operation. **Hancock's**, for amputation through the foot: a modification of Pirogoff's operation, in which the sawn surface of the os calcis is brought in contact with the transverse section of the astragalus. **Heath's**, for fixity of the lower jaw: division of the ascending ramus beneath the masseter with a saw introduced through the mouth by means of a small incision above the last molar tooth. **Hey's**: 1. For amputation through the foot: the same as Lisfranc's operation, except that the internal cuneiform bone is sawn through in a line with the articulation of the second metatarsal bone instead of being disarticulated. 2. For amputation of the leg: the amputation is made in the middle of the leg by a long posterior flap, cut by transfixion, and a slightly shorter anterior one. **Holmes'**, for excision of the os calcis: an incision is made from the inner edge of the tendo Achillis along the upper border of the os calcis and the outer border of the foot to the calcaneocuboid joint, and this is joined by another incision running across the sole, the perineal tendons being divided. **Huguier's**, a method of performing colotomy: the right lumbar operation. **Hunter's**, for aneurysm: ligation of the artery on the cardiac side of the aneurysm at some distance from it. **Jordan's**: 1. For amputation at the hip-joint: the soft parts are divided down to the bone by the circular method, as low down as possible, and the femur is shelled out by a longitudinal incision along the outer side of the thigh. 2. For amputation at the shoulder-joint: the soft parts are divided down to the bone by the circular method, three or four inches below the axilla, and the humerus is shelled out by a longitudinal incision along the outer and posterior aspect of the limb, meeting the circular incision at right angles. **Knapp's**, for cataract-extraction: a broad iridectomy and peripheral opening of the capsule; the lens is expelled by gentle pressure on the lower part of the cornea. **Kocher's**: 1. For excision of the ankle-joint: the incision is made beneath the external malleolus, and is followed by division of the peroneal tendons after being secured with threads, and opening of the joint with removal of the diseased parts; the foot is replaced and the tendons sutured. 2. For excision of the thyroid gland: an incision is made in the median line from the sternal notch to the upper limit of the bronchocle; from this point two lateral incisions are made upward and outward to a point a little below the angle of the jaw. **Kocher's** method of reducing subcoracoid dislocation of the humerus: The patient is placed upon the back and anesthetized if necessary. The surgeon grasps the elbow of

the dislocated arm with his opposite hand, and with the other hand seizes the patient's wrist; the elbow is brought to the side; the forearm is next flexed to a right angle with the arm and then carried outward to a right angle with the patient's body by the surgeon's hand grasping the wrist; while in this position the elbow is raised by the hand grasping it and carried upward and inward across the patient's chest, the olecranon being kept close to the chest-wall until firm resistance is met; the arm is then quickly rotated upward and the patient's hand placed upon his opposite shoulder. **Kraske's**, for carcinoma of the rectum: resection of the coccyx and of a portion of the sacrum to give access to the rectum high up. **Laborde's Method** of artificial respiration: rhythmic traction of the tongue to stimulate the superior laryngeal nerve. **Lannelongue's**, for craniotomy: (a) the vertex of the skull is exposed by means of a single incision in its long axis and a narrow strip of parietal bone is removed close to the sagittal suture; (b) the bones of the skull are so cut as to produce osseous flaps of various shapes. **Larrey's**, for amputation at the shoulder-joint: an incision is made from the acromion to a point $2\frac{1}{4}$ inches below, from which point anterior and posterior spiral incisions are carried down the arm, meeting on the inner aspect; after disarticulation of the bone the flap is brought up and secured in the space made by the gaping of the first incision. **Lisfranc's**, for amputation through the foot: disarticulation through the tarsometatarsal joints. **Littre's**, inguinal colotomy: it is performed on the left side for opening of the sigmoid flexure; an incision from $1\frac{1}{2}$ to 3 inches long is made parallel with the outer part of Poupart's ligament, and about $\frac{1}{4}$ inch above it. **Loreta's**: 1. For aneurysm: the insertion of a small wire into the sac, and the use of electrolysis as well. 2. For stenosis of the orifices of the stomach: digital or instrumental division of the pyloric or cardiac orifice, following gastrotomy. **Macewen's**: 1. For aneurysm: a long pin with a tapering point is passed into the aneurysm and the point is made to touch the opposite wall of the sac; the point is then moved over the inner surface of the aneurysm for ten minutes, for the purpose of causing irritation. 2. For the radical cure of inguinal hernia: the sac is dissected out and separated from its connections, and then so drawn upward by means of a suture as to form a pad upon the inner surface of the circumference of the ring; the ring and canal are then closed by means of sutures. 3. Osteotomy for genu valgum: supracondyloid division of the femur from the inner side. **Marian's**, the old median perineal operation for stone

in the bladder. **McBurney's**, for the radical cure of inguinal hernia: after exposing the sac the anterior wall of the canal is incised and the sac ligated and excised at the internal ring; the skin is inverted and stitched to the ligamentous and tendinous structures, the wound healing by granulation. **McDowell's**, oophorectomy. **McGill's**, prostatectomy by a suprapubic incision, as in suprapubic lithotomy. **Mikulicz's**, a tarsectomy; the same as *Wladimiroff's* operation. **Müller's**: 1. For Cesarean section: a modification of Porro's operation, whereby the whole uterus is turned out of the abdomen before it is opened. 2. For vaginal hysterectomy: the uterus is divided into halves longitudinally; one part is brought down at a time, and the broad ligament is ligated in sections. **Nélaton's**, for amputation through the foot: subastragaloid disarticulation by dorsal and plantar flaps, larger on the inner than on the outer side. **Panas'**: 1. For proctotomy: linear proctotomy. 2. For ptosis: the tarsal portion of the lid is raised by sutures and the occipitofrontalis muscle is caused to assume, to a great extent, the function of the levator palpebræ. **Phelps'**, for club foot: a direct open incision is made through the inner and plantar surfaces of the foot. **Pirogoff's**: 1. For amputation through the foot: a partial osteoplastic operation in which the os calcis is sawn through obliquely from above downward and forward, and the posterior portion is brought up and secured against the surface made by sawing off the lower ends of the tibia and fibula. **Pollock's**, for amputation of the knee-joint: it is done by a long anterior and a short posterior skin-flap, somewhat rectangular in outline, the patella being left. **Porro's**, for puerperal hysterectomy: Cesarean section, followed by removal of the uterus at the cervical junction, together with the ovaries and oviducts. **Porro-Müller's**, for otherwise impossible labor: a modification of the Porro-operation in which the uterus is brought out of the abdomen before extracting the fetus. **Porro-Veit's**, for otherwise impossible labor: a modification of the Porro-operation: the stump is ligated and dropped. **Pratt's**, the practice of orificial surgery, based on the belief that many chronic diseases are due to morbid conditions of the orifices of the body, particularly the anus and the urogenital canal. The operation consists in the dilatation of either or both of these orifices and the removal of any irritating condition that may be present. **Ramsden's**, for ligation of the third part of the subclavian artery: a transverse incision three inches in length is made across the base of the posterior triangle of the neck, about one-half inch

above the clavicle. **Reverdin's**, for skin-grafting: a point of skin is raised on an ordinary sewing needle, and shaved off with a scalpel or scissors; the graft is then transferred to the fresh surface next to the healthy granulations. **Roberts'**, for deflected nasal septum: a linear incision is made with a bistoury along a prominent line of the deflection, so as to enable it to be pressed into its proper position, after which a long steel pin is thrust into the septum in such a manner as to maintain this position. **Rose's**, for the relief of neuralgia of the fifth nerve: removal of the Gasserian ganglion. **Sæmisch's**, for hypopyon-ulcer: the cornea is transfixed and the intervening tissue, including the base of the ulcer, is divided by cutting outward. **Sänger's**, a method of performing Cesarean section: a modification of the usual operation in which the uterus is brought out through a long abdominal incision before extraction of the fetus. **Scarpa's**, for ligation of the femoral artery: it is done at the middle of Scarpa's triangle. **Schmalz's**, for stricture of the lacrimal duct: the introduction of a thread through the sac, and as far into the duct as possible. **Schücking's**, for prolapse of the uterus: vaginal hysteropexia by means of a special needle, the thread being passed through the uterine cavity, out at the fundus, and through the anterior vaginal vault; the two ends are tied together and allowed to slough out. **Sédillot's**: 1. A method of performing amputations: the combination of the flap and circular methods; the superficial flaps are formed from within outward, and the deep muscles are divided circularly. 2. For amputation at the ankle-joint: this method resembles Syme's operation, except that the flap is made from the inner and plantar surfaces of the foot and ankle. 3. For amputation of the leg: it is done at the "place of election" by a large external flap cut by transfixion. 4. For amputation of the thigh: it is done by a single long anterior flap, the structures at the back of the limb being divided to the bone by one transverse incision. **Senn's**, for intestinal anastomosis: the diseased tissue is excised and the ends of the bowel entirely and permanently closed, the continuity of the canal being restored by means of "intestinal anastomosis by lateral approximation" without the use of bone-plates. **Sigault's**, for otherwise impossible labor: symphysiotomy. **Simon's**, for exploration of the rectum and adjacent parts: it consists in introducing the hand or the hand and arm of the operator into the bowel. **Smith's**, for hemorrhoids: crushing by means of a clamp, and applying the Paquelin-cautery to the stump after cutting

away the projecting part. **Syme's**: 1. For amputation at the ankle joint: a single plantar flap is made, from which the os calcis is dissected out, after disarticulation has been accomplished; the malleoli and lower end of the tibia are then removed. 2. For external urethrotomy: the stricture is divided through the perineum upon a grooved director. **Tagliacozzi's**, for rhinoplasty: the forming of a new nose from a flap from the arm secured in place, and left attached to the arm until union has occurred. **Tait's**, for lacerated perineum: the flap-splitting operation; the rectovaginal septum is split transversely and from the extremities of this incision two lateral incisions are made, running up toward the lower terminus of each nymphæ and down to a little above the anal orifice; these two flaps are dissected up; the upper one is drawn toward the urethra, the lower one down toward the anus; sutures are passed laterally and the edges are brought together. **Teale's**: 1. For amputation of the arm: the long flap is placed upon the anteroexternal aspect of the arm; the brachial artery and the median and ulnar nerves are divided with the posterior flap. 2. For amputation of the leg: a rectangular flap-operation, in which a long anterior and a short posterior flap are made, each consisting of both integument and muscle; the length of the anterior flap is equal to half the circumference of the limb, and the posterior flap is one-quarter of the length. **Thiersch's**, for skin-grafting: the graft is made by a to-and-fro motion with a razor, shaving off as long and as broad a section of the upper layers of the skin as is desired. **Thomas's**, laparoclytrotomy: removal of the fetus through incisions in the abdominal wall and in the vagina, without opening the peritoneum or wounding the uterus. **Van Buren's**, for prolapsus ani: a linear cauterization of the mucosa with the Paquelin-cautery. **Verneuil's**, an iliac colotomy: a nearly vertical incision is made, and a knuckle of gut is transfixed and kept in place by two needles which lie on the parietes; the exposed bowel is then stitched to the margins of the parietal wound and the protruding knuckle is excised. **Wardrop's**, for aneurysm: ligation of a main branch of the artery distal to the aneurysm, leaving a circulation, however, through another branch. **White's**, for hypertrophy of the prostate: removal of the testicles. **Whitehead's**: 1. For excision of hemorrhoids: removal of a circular strip of mucous membrane around the anus, including the tumors. 2. For excision of the tongue: removal through the mouth, using only scissors. **Wilde's**, for mastoid or cerebral abscess: the bone is exposed from the base to the apex of the mastoid process,

$\frac{1}{2}$ inch behind the auricle, and if necessary the bone is opened with a drill, gouge, or trephine. **Winiwarter's**, cholecystenterostomy: it is performed in two stages; in the first the gall-bladder is united to the upper portion of the jejunum and the parts are fixed to the parietal peritoneum; in the second, after about five days, the bowel is incised and a communication is established between the latter and the gall-bladder.

Wladimiroff's, a form of tarsectomy: the astragalus, os calcis, and the soft parts covering them, are removed; the articular surfaces of the tibia, fibula, cuboid, and scaphoid bones are sawn off and the foot brought into a straight line with the leg. **Wültzer's**, for the radical cure of inguinal hernia: plugging of the hernial canal by an invagination of the scrotum, and its retention by a special instrument, thus exciting adhesive inflammation in the neck of the sac. **Wyeth's**, for amputation at the hip-joint; the bloodless method: two steel mattress-needles are inserted in the tissues and rubber tubing is passed several times around the limb above the needles.

Operative (*op'-er-a-tiv*) [*operari*, to labor]. 1. Able to act; effective. 2. Pertaining to operations.

Operculum (*o-per'-ku-lum*) [L.]. 1. A lid or cover, as O. ilei, the ileocecal valve. 2. The convolutions covering the island of Reil.

Ophryon (*off'-re-on*) [*ὀφρύς*, eyebrow]. In craniometry, the middle of a line drawn across the forehead at the level of the upper margin of the orbits.

Ophthalmia (*off'-thal'-me-ah*) [*ὀφθαλμός*, eye]. Inflammation of the eye, especially one in which the conjunctiva is involved. **O.**, **Catarrhal**, simple conjunctivitis; a hyperemia of the conjunctiva, with a mucopurulent secretion. **O.**, **Caterpillar-**, inflammation of the conjunctiva or of the cornea, the result of penetration of the tissues by the hairs of caterpillars. **O.**, **Egyptian**. See *Trachoma*. **O.**, **Gonorrheal**, an acute and severe form of purulent conjunctivitis, caused by infection from urethral discharges containing the gonococcus of Neisser. **O.**, **Granular**. See *Trachoma*. **O.** **neonatorum**, a gonorrheal or purulent ophthalmia of the new-born, the eyes having been infected by the mother's vaginal discharges. **O.**, **Neuroparalytic**, disease of the eye from lesion of the Gasserian ganglion or of branches of the fifth nerve supplying the eyeball. **O.**, **Phlyctenular**, conjunctivitis characterized by phlyctenules or small vesicles situated in the epithelial layer of the conjunctiva or cornea. **O.**, **Purulent**, conjunctivitis with a purulent discharge. **O.**, **Sympathetic**, a severe destructive inflammation, a form of iridocy-

clitis secondary to injury or disease of the fellow-eye.

Ophthalmic (*off'-thal'-mik*) [*ὀφθαλμός*, eye]. Pertaining to the eye. **O.** **Artery**. See *Arteries*, *Table of*. **O.** **Ganglion**, the ciliary ganglion. See *Ganglia*, *Table of*. **O.** **Nerve**. See *Nerves*, *Table of*.

Ophthalmitis (*off'-thal'-mi'-tis*) [*ὀφθαλμός*, eye; *ιτις*, inflammation]. Inflammation of the eye.

Ophthalmo- (*off'-thal'-mo-*) [*ὀφθαλμός*, eye]. A prefix signifying connection with or relation to the eye.

Ophthalmoblenorrhoea (*off'-thal'-mo-blen-or-e'-a*) [*ὀφθαλμός*, eye; *βλέννα*, mucus]. Blenorrhoea of the conjunctiva.

Ophthalmocoele (*off'-thal'-mo-sel*) [*ὀφθαλμός*, eye; *κήλη*, tumor]. See *Exophthalmus*.

Ophthalmocopia (*off'-thal'-mo-ko'-pe-ah*) [*ὀφθαλμός*, eye; *κόπος*, fatigue]. Fatigue of visual power; asthenopia.

Ophthalmodynia (*off'-thal'-mo-din'-e-ah*) [*ὀφθαλμός*, eye; *ὀδύνη*, pain]. Neuralgic pain in the eye.

Ophthalmologist (*off'-thal'-mol'-o-jist*) [*ὀφθαλμός*, eye; *λόγος*, science]. One versed in ophthalmology.

Ophthalmology (*off'-thal'-mol'-o-je*) [*ὀφθαλμός*, eye; *λόγος*, science]. The science of the anatomy, physiology, and diseases of the eye.

Ophthalmomalacia (*off'-thal'-mo-mal-a'-se-ah*) [*ὀφθαλμός*, eye; *μαλακία*, softness]. Abnormal softness or subnormal tension of the eye.

Ophthalmometer (*off'-thal'-mom'-et-er*) [*ὀφθαλμός*, eye; *μέτρον*, measure]. 1. An instrument for measuring the capacity of the chambers of the eye. 2. An instrument for measuring refractive errors, especially astigmatism. 3. An instrument for measuring the eye as a whole.

Ophthalmometry (*off'-thal'-mom'-et-re*) [*ὀφθαλμός*, eye; *μέτρον*, measure]. The determination of refractive errors by means of the ophthalmometer.

Ophthalmopathy (*off'-thal'-mop'-a-the*) [*ὀφθαλμός*, eye; *πάθος*, disease]. Any disease of the eye.

Ophthalmophthisis (*off'-thal'-moff'-this-is*). See *Phthisis bulbi*.

Ophthalmoplegia (*off'-thal'-mo-ple'-je-ah*) [*ὀφθαλμός*, eye; *πληγή*, stroke]. Paralysis of the ocular muscles. **O.** **externa**, paralysis of the external ocular muscles. **O.** **interna**, paralysis of the internal muscles of the eye, those of the iris and ciliary body. **O.**, **Nuclear**, due to a lesion of the nuclei of origin of the motor nerves of the eyeball. **O.**, **Partial**, a form in which only some of the muscles are paralyzed. **O.**, **Progressive**, a form in which all of the muscles of both eyes gradually become paralyzed. **O.**, **Total**,

that form involving the iris and ciliary muscle as well as the external muscles of the eyeball.

Ophthalmoplegic (*off-thal-mo-ple'-jik*) [*ὀφθαλμικός*, eye; *πληγή*, stroke]. Pertaining to ophthalmoplegia.

Ophthalmoscope (*off-thal'-mo-skōp*) [*ὀφθαλμικός*, eye; *σκοπεῖν*, to see]. An instrument for examining the interior of the eye. It consists essentially of a mirror with a hole in it, through which the observer looks, the concavity of the eye being illuminated by light reflected from the mirror into the eye and seen by means of the rays reflected from the eye-ground back through the hole in the mirror. The ophthalmoscope is fitted with lenses of different powers that may be revolved in front of the observing eye, and these neutralize the ametropia of either the patient's or the observer's eye, thus rendering the details of the fundus oculi clear.

Ophthalmoscopic (*off-thal-mo-skōp'-ik*) [*ὀφθαλμικός*, eye; *σκοπεῖν*, to see]. Pertaining to the ophthalmoscope or its use.

Ophthalmoscopy (*off-thal-mo-skō-pe*) [*ὀφθαλμικός*, eye; *σκοπεῖν*, to see]. The examination of the interior of the eye by means of the ophthalmoscope. **O.**, **Direct**, the method of the erect or upright image, the observer's eye and the ophthalmoscope being brought close to the eye of the patient. **O.**, **Indirect**, the method of the inverted image; the observer's eye is placed about 16 inches from that of the patient and a 20 D. biconvex lens is held about two inches in front of the observed eye, thereby forming an aerial inverted image of the fundus.

Ophthalmostat (*off-thal'-mo-stat*) [*ὀφθαλμικός*, eye; *στάνααι*, to cause to stand]. An instrument used in fixing the eye in any position during an operation on it.

Ophthalmostatometer (*off-thal-mo-stat-om'-et-er*) [*ὀφθαλμικός*, eye; *στάνααι*, to cause to stand; *μετρον*, measure]. An instrument for determining the position of the eyes.

Ophthalmotonometer (*off-thal-mo-ton-om'-et-er*) [*ὀφθαλμικός*, eye; *τόνος*, tone; *μετρον*, measure]. An instrument for measuring intraocular tension.

Ophthalmotrope (*off-thal'-mo-trōp*) [*ὀφθαλμικός*, eye; *τρόπος*, a turn]. An instrument used for the demonstration of the direction and the position that the eye takes under the influence of each of its muscles, and the position of the false image in the case of paralysis of a given muscle.

Opiate (*o'-pe-at*) [*ὄπιον*, poppy-juice]. A preparation of opium.

Opiophagism, Opiophagy (*o-pe-ōff'-aj-izm, o-pe-ōff'-aj-e*) [*ὄπιον*, opium; *φαγῖν*, to eat]. Opium-eating.

Opisthopia (*o-pis-tho-po-ri'-ah*) [*ὀπισθεν*,

behind; *πορεία*, going]. Involuntary backward-walking in an attempt to go forward.

Opisthotonus (*o-pis-thoi'-on-us*) [*ὀπισθεν*, behind; *τόνος*, tone]. A condition in which from a tetanic spasm of the muscles of the back the head and lower limbs are bent backward, and the body arched forward.

Opium (*o'-pe-um*) [*ὄπιον*, from *ὀπός*, juice].

The inspissated juice obtained by incising the unripe capsules of *Papaver somniferum*, of the order *Papaveraceæ*, occurring in commerce in the form of brownish cakes having a narcotic odor and a bitter taste. **O.** contains a large number of alkaloids, of which morphin is the most important since it represents the chief properties of the drug. Other alkaloids are narcotin, $C_{22}H_{25}NO_7$, codein, $C_{16}H_{21}NO_3$, thebain or paramorphin, $C_{16}H_{21}NO_3$, papaverin, $C_{20}H_{21}NO_4$, narcein, $C_{22}H_{29}NO_5$, pseudomorphin, $C_{34}H_{48}N_2O_6$, laudanin, $C_{20}H_{25}NO_4$. These bases occur in **O.** combined with meconic and thebolactic acids. Moist **O.** should contain not less than seven per cent. of crystallized morphin (U. S. P.) **O.** acts as a narcotic, producing deep sleep, which, however, is often preceded by a stage of mental excitement and exhilaration; on awakening there may be headache, nausea, or vomiting. It slows the pulse and increases its force, and raises blood-pressure; small doses do not depress the respiration, large doses do so in a marked degree. **O.** checks the motor activity of the stomach and intestines, and lessens all secretions except that of the skin; it produces contraction of the pupil. When taken in poisonous doses it causes unconquerable drowsiness, passing into deep sleep, with slow, full respiration, slow pulse, contracted pupils; later cyanosis develops, the respiration becomes exceedingly slow, and the pulse rapid and feeble; death takes place from failure of the respiration. See *Poisons, Table of*. There is a chronic form of **O.**-poisoning produced by the habitual use of **O.** or morphin, and characterized by mental depression, a deterioration of the moral sense, and attacks of diarrhea. **O.** is used for the relief of pain of all forms except that due to cerebral inflammation; in insomnia; in inflammation of serous membranes; in spasmodic conditions; in acute colds; cough, retention of urine, vomiting, diarrhea, certain forms of dyspnea, particularly that from heart-disease; and locally as an application to sprains and inflamed surfaces. Dose gr. j (0.065).

PREPARATIONS AND DOSES.—*Acetum opii* (U. S. P.), vinegar of opium, black drop; *πλv-xxx* (0.32-2.0). *Confectio opii* (B. P.), *πλv-xx* (0.32-1.3). *Emplastrum opii* (U. S. P., B. P.). *Enema opii* (B. P.). *Extractum opii* (U. S. P., B. P.), gr

0.032). *Extractum opii liquidum* (B. P.), ηv -xx (0.32-1.3). *Opium pulvis*, gr. $\frac{1}{2}$ -j (0.016-0.065). *Opium deodoratum* (U. S. P.), gr. j (0.065). *Pilulae opii* (U. S. P.), each containing gr. j (0.065). *Pilula saponis composita* (B. P.), contains about 20 per cent. of opium. *Pulvis ipecacuanhae et opii*, Dover's powder, contains ten per cent. each of opium and ipecac. *Pulvis opii compositus* (B. P.), contains ten per cent. of opium. *Tinctura ipecacuanhae et opii* (U. S. P.), ηv -xv (0.32-1.0); *Tinctura opii, laudanum* (U. S. P., B. P.), ηv -xv (0.32-1.0). *Tinctura opii ammoniata* (B. P.), ηxxx -fzj (2.0-4.0). *Tinctura opii camphorata* (U. S. P.), (*Tinctura camphorae composita*, B. P.), paregoric, fzj-iv (4.0-16.0). *Tinctura opii deodorati* (U. S. P.), ηv -xv (0.32-1.0). *Trochisci glycyrrhizae et opii* (U. S. P., B. P.), each containing 0.005 gram of opium. *Vinum opii* (U. S. P., B. P.). Sydenham's laudanum, ηv -xv (0.32-1.0). See also *Morphin*.

Opiumism (*o'-pe-um-izm*) [*opium*]. The condition produced by the action of opium on the system.

Opobalsamum (*o-po-bawl-sam'-um*) [*ὀπός*, juice; *βάλσαμον*, balsam]. A resin from *Balsamodendron opobalsamum*, and *Balsamodendron gileadense*. See *Balm of Gilead*.

Opocephalus (*o-po-sef'-al-us*) [*ὤψ*, eye; *κεφαλή*, head]. A monster characterized by fusion of the ears, one orbit, and absence of mouth and nose.

Opodeldoc (*op-o-del'-dok*) [origin obscure]. See under *Soap*.

Opodymus (*o-pod'-im-us*) [*ὤψ*, eye; *δίδυμος*, twin]. A monster with a single body and skull, but with two distinct faces.

Opothrapy (*op-o-ther'-ap-e*) [*ὀπός*, juice; *therapy*]. Synonym of *Organotherapy*.

Opponents (*op-o'-nens*) [*ob*, against; *ponere*, to place]. Opposing. A term applied to certain muscles that bring one part opposite another, as *Opponens minimi digiti*, a muscle placing the little finger opposite the thumb.

Optic (*op'-tik*) [*ὀπτικός*, from the base, *οπ-*, to see]. Pertaining to vision or to the science of optics. **O. Atrophy**, atrophy of the optic nerve. **O. Axis**, the axis of the eye.

O. Center. 1. The point in the main axis of the crystalline lens at which the rays of light meet. 2. The nerve center concerned in the visual function. **O. Chiasm**, **O. Commis-sure**. See *Commissure*. **O. Cup**, the concave area formed by the involution of the distal extremity of the primary optic vesicle. **O. Disc**, the optic papilla. **O. Foramen**. See *Foramina*, *Table of*. **O. Groove**, the groove on the sphenoid bone for the optic chiasm. **O. Lobes**, the corpora quadrigemina. **O. Nerve**. See *Nerves*, *Table of*. **O.**

Neuritis. See *Papillitis*. **O. Papilla**, the circular prominence formed by the optic nerve after its entrance into the eyeball. **O. Radiations**, a large bundle of nerve-fibers joining the optic thalamus and the occipital lobe of the cerebrum. **O. Thalamus**. See *Thalamus*. **O. Tract**. See *Tract*. **O. Vesicle**, a diverticulum from each side of the primary anterior vesicle of the embryo, forming the basis of the future eye.

Optical (*op'-tik-al*) [*ὀπτικός*, pertaining to sight, from the base *οπ-*, to see]. Pertaining to sight.

Optician (*op'-tish'-un*) [*ὀπτικός*, from the base *οπ-*, to see]. A maker of optical instruments.

Opticociliary (*op'-tik-o-sil'-e-a-re*) [*ὀπτικός*, pertaining to vision, from the base *οπ-*, to see; *ciliary*]. Pertaining to the optic and the ciliary nerves.

Opticopupillary (*op'-tik-o-pu'-pil-a-re*) [*ὀπτικός*, pertaining to vision, from the base *οπ-*, to see; *pupillary*]. Pertaining to the optic nerve and the pupil.

Optics (*op' tiks*) [*ὀπτικός*, from the base *οπ-*, to see]. The science treating of light and vision.

Optogram (*op'-to-gram*) [*ὀπτός*, visible; *γράφειν*, to write]. A faint image stamped on the retina for a brief period after death, believed to be that of the last object seen before death.

Optometer (*op'-tom'-et-er*) [*ὀπτός*, visible; *μέτρον*, measure]. An instrument for determining the strength of vision, especially the degree of refractive error that is to be corrected to render vision normal.

Optostriate (*op'-to-stri'-at*) [*ὀπτός*, visible; *striatum*, striped]. Pertaining to the optic thalamus and the corpus striatum.

Ora (*o'-rah*) [L.]. Margin. **O. serrata**, the jagged anterior margin of the retina.

Oral (*o'-ral*) [*os*, the mouth]. Pertaining to the mouth.

Orange (*or'-anj*) [Pers., *nāranj*, orange]. See *Aurantium*.

Orbicular (*or-bik'-u-lar*) [*orbicularis*, dim. of *orbis*, circle]. Circular. A term applied to circular muscles, as the orbicular muscle of the eye or of the mouth (*Orbicularis palpebrarum*, *Orbicularis oris*).

Orbicularis (*or-bik-u-la'-ris*). See *Muscles*, *Table of*.

Orbit (*or'-bit*) [*orbita*, from *orbis*, circle]. The bony pyramidal cavity containing the eye, and formed by the frontal, sphenoid, ethmoid, nasal, lacrimal, superior maxillary, and palate bones.

Orbital (*or'-bit-al*) [*orbita*, from *orbis*, circle]. Pertaining to the orbit.

Orchi-, **Orchid-**, **Orchio-** (*or'-ke-*, *or'-kid-*, *or'-ke-o-*) [*ὄρχις*, testicle]. A prefix signifying connection with or relation to the testicle.

Orchialgia (*or-ke-al'-je-ah*) [*ὄρχις*, testicle; *ἄλγος*, pain]. Neuralgia of the testicle.

Orchichorea (*or-ke-ko-re'-ah*) [*ὄρχις*, testicle; *chorea*]. Irregular movements of the testicle due to contraction of the cremaster muscle.

Orchidectomy (*or-kid-ek'-to-me*) [*ὄρχις*, testicle; *ἐκτομή*, excision]. Castration.

Orchidotomy (*or-kid-ot'-o-me*) [*ὄρχις*, testicle; *τομή*, a cutting]. Incision of the testicle.

Orchiocoele (*or'-ke-o-sel'*) [*ὄρχις*, testicle; *κύλη*, tumor]. 1. A tumor of the testicle. 2. Scrotal hernia.

Orchiotomy (*or-ke-ot'-o-me*). Synonym of *Orchidotomy*.

Orchis (*or'-kis*) [*ὄρχις*, testicle]. 1. A genus of plants furnishing salep and vanilla. 2. The testicle.

Orchitis (*or-ki'-tis*) [*ὄρχις*, testicle; *ιτις*, inflammation]. Inflammation of the testicle.

Orcin (*or'-sin*) [*Orcus*, Pluto, from its dark color], $C_7H_6(OH)_2$. A substance found in many lichens of the genera *Rocella* and *Lecanora*. It is an antiseptic, and has been used instead of resorcin.

Ordeal-bark (*or'-de-al-bark*). See *Erythrophloeum*. **Ordeal-bean**, **Ordeal-nut**. See *Physostigma*.

Orexin (*o-regs'-in*) [*ὄρεξις*, appetite], $C_{14}H_{17}N_7HCl \cdot 2H_2O$. Hydrochlorate of phenylidihydroquinazolin,—used as a stomachic. Dose gr. iv-vij (0.26-0.45).

Organ (*or'-gan*) [*ὄργανον*, organ]. A part of the body having a definite function to perform. **O. of Corti**. See *Ear*. **O., Enamel-**, a club-shaped process of epithelium growing from the dental ridge and forming a cap over the dental papilla. From it the enamel of the tooth is developed. **O. of Giralès**, the paradidymis. **O., Jacobson's**. See *Jacobson's Organ*. **O.'s of Reproduction**, the testicles and penis and its glands in the male, the uterus and its appendages and the vagina and its glands, in the female. **O. of Rosenmüller**. See *Parovarium*. **O., Segmental**, a mesoblastic embryonic structure consisting of three parts—the pronephros, the mesonephros, or Wolffian body, and the metanephros.

Organic (*or-gan'-ik*) [*ὄργανον*, an organ]. Having, pertaining to, or characterized by organs; pertaining to the animal and vegetal worlds; affecting the structure of organs. **O. Chemistry**, the chemistry of the carbon-compounds, carbon being the central element of compounds occurring in organized beings. **O. Disease**, disease of an organ attended with structural changes.

Organism (*or'-gan-izm*) [*organ*]. A body consisting of an aggregation of organs having a definite function; a living being.

Organization (*or-gan-iz-a'-shun*) [*ὄργανον*, organ]. 1. The orderly arrangement of or-

gans or parts. 2. An organism. 3. The conversion into an organ, or into something resembling an organ, or into living tissue.

Organogenesis (*or-gan-o-jen'-es-is*) [*ὄργανον*, organ; *γενᾶν*, to beget]. The process of the development of an organ.

Organogenetic (*or-gan-o-jen-et'-ik*) [*ὄργανον*, organ; *γενᾶν*, to beget]. Pertaining to organogenesis.

Organography (*or-gan-og'-ra-fe*) [*ὄργανον*, organ; *γράφειν*, to write]. A descriptive treatise of the organs of an animal or plant.

Organoid (*or'-gan-oid*) [*ὄργανον*, organ; *εἶδος*, like]. Resembling an organ.

Organoleptic (*or-gan-o-lep'-tik*) [*ὄργανον*, organ; *ληπτικός*, taking]. Affecting organs, particularly those of special sense.

Organology (*or-gan-ol'-o-je*) [*ὄργανον*, organ; *λόγος*, science]. The science that treats of the organs of plants and animals.

Organotherapy (*or-gan-o-ther'-ap-e*) [*ὄργανον*, organ; *θεραπεία*, treatment]. The treatment of diseases by the administration of animal organs, or extracts prepared from them. The organs chiefly employed are: 1. The testicles (testicular extract), for various nervous affections. From testicular juice a crystalline substance, spermin, has been isolated and its use advised in nervous depression, debility, and asthma. 2. Thyroid gland (in the form of the gland itself, glycerol-extracts, or the desiccated gland). It has been found of much value in myxedema, cretinism, various chronic skin-diseases, in obesity, colloid goiter, and of questionable value in exophthalmic goiter. 3. Pancreas (extract of pancreas), for diabetes mellitus. 4. Suprarenal bodies (extract of suprarenal bodies), for Addison's disease. 5. Bone-marrow, for pernicious anemia and other anemias. Other extracts proposed are those of brain, spinal marrow, muscles, kidney, ovary, heart-muscle, spleen, thymus gland, and stomach.

Orgasm (*or'-gasm*) [*ὄργασμα*, to swell]. Intense excitement, especially that occurring during sexual intercourse.

Oriental (*o-re-ent'-tal*) [*oriens*, rising]. Pertaining to the Orient or east. **O. Boil**. See *Furunculus orientalis*.

Orientation (*o-ri-en-ta'-shun*) [*oriri*, to rise (in the east)]. The act of determining one's position in space (subjective O.) or the position of surrounding objects with reference to each other (objective O.).

Orifice (*or'-if-is*) [*orificium*; *os*, mouth; *facere*, to make]. An opening.

Orificial (*or-if-ish'-al*) [*orificium*, an orifice]. Pertaining to an orifice. **O. Surgery**. See *Pratt's Operation*, in *Operations*, *Table of*.

Origanum (*o-rig'-an-um*) [*ὄριγανον*, marjoram]. A genus of plants of the order Labi-

atæ. **O. majorana**, sweet marjoram, is used as a condiment. **O. vulgare**, wild marjoram, contains a pungent oil (Oleum origani), consisting chiefly of terpene, $C_{10}H_{16}$. **O.** is tonic, excitant, diaphoretic, and emmenagogue, and locally anodyne. It is also employed as a clearing-agent in microscopy.

Origin (*or'-ij-in*) [*origo*, from *oriri*, to arise]. The beginning or starting-point of anything. **O.**, Deep, of a nerve, its **O.** in the cells of the nerve-center. **O.**, Superficial, of a nerve, the point at which it emerges from the brain or cord. **O.** of a muscle, the point of attachment of a muscle which remains relatively fixed during contraction of the muscle.

Orinasal (*o-re-na'-zal*) [*os*, mouth; *nasus*, nose]. Pertaining to the mouth and the nose.

Orolingual (*o-ro-lin'-gw'al*) [*os*, mouth; *lingua*, tongue]. Pertaining to the mouth and the tongue.

Oronasal (*o-ro-na'-zal*). See *Orinasal*.

Oropharynx (*o-ro-far'-inks*) [*os*, mouth; *φάρυγξ*, pharynx]. The pharynx proper, situated below the level of the lower border of the soft palate, as distinguished from the nasopharynx.

Orpiment (*or'-pim-ent*). Arsenic trisulphid.

Irris (*or'-is*). See *Iris*, 2d definition.

Ortho- (*or'-tho-*) [*ὀρθός*, right, straight]. 1. A prefix denoting straight, normal, or true. 2. In chemistry, a prefix denoting that one among several compounds of the same elements which is considered the normal compound. Among derivatives of the benzol-ring it refers to those formed by the substitution of two adjacent hydrogen-atoms.

Orthocephalous (*or-tho-sef'-al-us*) [*ὀρθός*, straight; *κεφαλή*, head]. Having a skull with a vertical index of from 70.1 to 75.

Orthodontia (*or-tho-don'-she-ah*) [*ὀρθός*, straight; *ὀδούς*, tooth]. The correction of irregularities of the teeth.

Orthognathous (*or-thog'-na-thus*) [*ὀρθός*, straight; *γνάθος*, the jaw]. Straight-jawed; having a gnathic angle of from 83° to 90°.

Orthometer (*or-thom'-et-er*) [*ὀρθός*, straight; *μέτρον*, measure]. An instrument for measuring the relative degree of protrusion of the eyes.

Orthopedic (*or-tho-pe'-dik*) [*ὀρθός*, straight; *παῖς*, child]. Pertaining to the correction of deformities, especially in children. **O. Surgery**, the branch of surgery devoted to the correction of deformities.

Orthopedics (*or-tho-pe'-diks*). See *Orthopedic Surgery*.

Orthopedist (*or-tho-pe'-dist*) [*ὀρθός*, straight; *παῖς*, child]. One who practices orthopedic surgery.

Orthophoria (*or-tho-fó'-re-ah*) [*ὀρθός*, straight; *φορός*, a tending]. A tending of the visual lines in parallelism.

Orthophosphoric Acid (*or-tho-fos-for'-ik*). See *Acid*.

Orthopnea (*or-thop-ne'-ah*) [*ὀρθός*, straight; *πνοή*, breathing]. A condition marked by quick and labored breathing, in which the patient finds relief only by maintaining an upright position.

Orthopneic (*or-thop-ne'-ik*) [*ὀρθός*, straight; *πνοή*, breathing]. Characterized by orthopnea.

Orthoptic (*or-thop'-tik*) [*ὀρθός*, straight; *ὀπτικός*, seeing]. Pertaining to normal binocular vision. **O. Training**, a method of correcting the defective vision of those having strabismus or muscular insufficiency, by stereoscopic and other ocular exercises of a gymnastic kind.

Orthoscope (*or'-tho-skōp*) [*ὀρθός*, straight; *σκοπεῖν*, to see]. An instrument for examination of the eye through a layer of water, whereby the curvature, and hence the refraction of the cornea, is neutralized and the cornea acts as a plane medium.

Orthoscopic (*or-tho-skop'-ik*) [*ὀρθός*, straight; *σκοπεῖν*, to see]. 1. Pertaining to an orthoscope or to orthoscopy. 2. Applied to lenses cut from the periphery of a large lens. 3. Having normal vision.

Orthoscopy (*or-thos'-ko-pe*) [*ὀρθός*, straight; *σκοπεῖν*, to see]. The examination of the eye with the orthoscope.

Orthotonus (*or-thot'-o-nus*) [*ὀρθός*, straight; *τόνος*, tension]. Tetanic cramp in which the body lies rigid and straight.

Oryza (*o-ri'-zah*) [*ὄρυζα*, rice]. See *Rice*.

Os [*L.*, gen., *oris*]. The mouth. **O. externum**, the external orifice of the cervical canal of the uterus opening into the vagina. **O. internum**, the internal orifice of the cervical canal opening into the cavity of the body of the uterus. **O. uteri**, either orifice of the cervical canal of the uterus.

Os [*L.*, gen., *ossis*]. A bone. **O. calcis**, the bone of the heel. **O. capitatum**. See *O. magnum*. **O. coxæ**. See *O. innominatum*. **O. femoris**, the femur. **O. hamatum**, the unciform bone. **O. humeri**, the humerus. **O. ilii**, the ilium. **O. innominatum**, the innominate bone. **O. ischii**, the ischium. **O. magnum**, the third bone of the second row of the carpus. **O. pubis**, the pubis. **Ossa suturarum**, **Ossa triquetra**, **Ossa wormiana**, the Wormian bones. **O. unguis**, the lacrimal bone.

Os. Symbol for osmium.

Oscedo (*os-se'-do*) [*L.*]. Yawning.

Oscheitis (*os-ke-i'-tis*) [*ὀσχεον*, scrotum; *itis*, inflammation]. Inflammation of the scrotum.

Oscheocele (*os'-ke-o-sel*) [*ὀσχεον*, scrotum; *κίλη*, tumor]. Scrotal hernia.

Oscheohydrocele (*os-ke-o-hi'-dro-sel*) [*ὀσ-*

yeon, scrotum; *ιδωρ*, water; *κήλη*, tumor]. A hydrocele occupying the sac of a scrotal hernia after the return of the bowel to the peritoneal cavity and the shutting off of the sac from the latter.

Oscheoplasty (*os'-ke-o-plas-te*) [*ὀσχεον*, scrotum; *πλασσειν*, to form]. Plastic surgery of the scrotum.

Oscitation (*os - it - a' - shun*) [*oscitare*, to yawn]. The act of yawning.

Osculation (*os-ku-la'-shun*) [*osculari*, to kiss]. The union of vessels by their mouths.

-ose (*-ōz*) [*-osis*]. A suffix denoting a member of the carbohydrate group.

-osis (*-ō'-sis*). A suffix signifying condition of, or state caused by.

Osmate (*os'-mat*) [*osmium*]. A salt of osmic acid.

Osmazome (*os'-maz-ōm*) [*ὀσμῆ*, smell; *ζωμός*, broth]. A brownish-yellow substance developed by heat in muscular fibers, and formerly supposed to give to cooked meats their peculiar flavor.

Osmic (*os'-mik*) [*ὀσμῆ*, smell]. Pertaining to or containing osmium. **O. Acid**, strictly, a dibasic acid, H_2OsO_4 ; as generally used, however, osmium tetroxid, OsO_4 , also called perosmic acid. It is a crystalline substance, freely soluble in water, and giving off irritant vapors; locally, it acts as a caustic. It is employed in microscopy as a stain for fat and as a fixing agent. In medicine it has been used in neuralgia.

Osmidrosis (*os-mid-ro'-sis*) [*ὀσμῆ*, smell; *ιδρωσις*, sweat]. The secretion of a malodorous perspiration; bromidrosis.

Osmium (*os'-me-um*) [*ὀσμῆ*, smell]. A heavy metallic element belonging to the platinum-group. Symbol *Os*, specific gravity 22.48, atomic weight 190.3, quantivalence II, IV, VI, VIII. See *Osmic Acid*.

Osmometer (*os-mom'-et-er*) [*ὀσμῆ*, smell; *μετρον*, measure]. 1. An instrument for testing the sense of smell. 2. An apparatus for measuring osmosis.

Osmosis (*os-mo'-sis*) [*ὀσμός*, impulse]. The passage of liquids and substances in solution through porous septa. See *Endosmosis* and *Exosmosis*.

Osmotic (*os-mo'-tic*) [*ὀσμός*, impulse]. Pertaining to osmosis.

Ossa (*os'-ah*) [*Os*, *pl.* of *os*, a bone]. Bones. See *Os*.

Ossein (*os'-e-in*) [*os*, a bone]. The organic base of osseous tissue.

Osseoponeurotic (*os-e-o-pō-nē-u-ro'-tik*) [*os*, bone; *aponeurosis*]. Bounded by bone and the aponeurosis of a muscle.

Osseous (*os'-e-us*) [*os*, bone]. Bony; composed of or resembling bone.

Ossicle (*os'-ik-l*) [*ossiculum*, dim. of *os*, a bone]. A small bone. **O.**, Auditory, one

of a chain of small bones found in the tympanic cavity of the ear.

Ossiferous (*os-iff'-er-us*) [*os*, a bone; *ferre*, to bear]. Containing or producing bone-tissue.

Ossific (*os-iff'-ik*) [*os*, bone; *facere*, to make]. Producing bone.

Ossification (*os-iff-ik-a'-shun*) [*os*, a bone; *facere*, to make]. The formation of bone.

Ossifying (*os-iff-i-ing*) [*os*, bone; *facere*, to make]. Changing into bone.

O. Chondroma, a chondroma that is undergoing ossification. **O. Myositis**, inflammation of muscle, attended with, or followed by, deposition of bone-like masses.

Ostalgia (*os-tal'-je-ah*) [*ὀστέον*, bone; *ἀλγος*, pain]. Pain in a bone.

Osteectomy (*os-te-ek'-to-me*) [*ὀστέον*, bone; *ἐκτομή*, excision]. Excision of a portion of a bone.

Osteitis (*os-te-īt'-tis*) [*ὀστέον*, bone; *ιτις*, inflammation]. Inflammation of bone.

O. Condensing. See *Osteosclerosis*. **O. deformans**, a rare form characterized by the production of deformity.

O. Gummatous, a chronic form due to syphilis and characterized by the formation of gummata in the cancellous tissue of the epiphysis or in the shaft of a bone, or in the periosteum. **O. Rarefying**. See *Osteoporosis*. **O. Sclerosing**. See *Osteosclerosis*.

Osteo- (*os'-te-o-*) [*ὀστέον*, bone]. A prefix signifying connection with or relation to bone.

Osteoarthritis (*os-te-o-ar-thri'-tis*) [*ὀστέον*, bone; *ἀρθρον*, joint; *ιτις*, inflammation].

An inflammation of the bones forming a joint.

Osteoarthropathy (*os-te-o-ar-throp'-ath-e*) [*ὀστέον*, bone; *ἀρθρον*, joint; *πάθος*, disease].

Any disease of bony articulations.

O. Hypertrophic Pulmonary, a disease characterized by a bulbous enlargement of the terminal phalanges of the fingers and toes, a thickening of the articular ends of the bones, and a peculiar curvature of the nails. The condition is usually associated with disease of the lungs or pleura, hence the name *osteoarthropathie pncumique hypertrophique* given to it by Marie, and results from the absorption of toxic products from the diseased foci.

Osteoblast (*os'-te-o-blast*) [*ὀστέον*, bone; *βλαστός*, a germ]. Any one of the cells of mesoblastic origin concerned in the formation of bony tissue.

Osteocarcinoma (*os-te-o-kar-sin-o'-mah*) [*ὀστέον*, bone; *carcinoma*]. 1. Ossifying carcinoma. 2. Carcinoma of bone.

Osteochondritis (*os-te-o-kon-dri'-tis*) [*ὀστέον*, bone; *χόνδρος*, cartilage; *ιτις*, inflammation]. Inflammation involving both bone and cartilage.

Osteochondroma (*os-te-o-kon-dro'-mah*) [*ὀστέον*, bone; *χόνδρος*, cartilage; *δμα*, tumor]. A tumor that is in part bony and in part cartilaginous.

Osteoclasia (*os-te-ok'-lu-sis*) [*ὀστέον*, bone; *κλάειν*, to break]. 1. Fracture of bones for purposes of remedying deformity. 2. The destruction of bony tissue by osteoclasts.

Osteoclast (*os'-te-o-klast*) [*ὀστέον*, bone; *κλάειν*, to break]. 1. An instrument for performing osteoclasia. 2. One of the large multinuclear cells found against the surface of bone in little eroded depressions (Howship's lacunæ), and concerned in the removal of bone.

Osteocope (*os'-te-o-kōp*). See *Osteocopic Pain*.

Osteocopic Pain (*os-te-o-kop'-ik*) [*ὀστέον*, bone; *κόπος*, a beating]. A severe pain in a bone, usually worse at night. It is a symptom of osteitis or periostitis, especially of syphilitic origin.

Osteogenesis (*os-te-o-jen'-es-is*) [*ὀστέον*, bone; *γενᾶν*, to beget]. The development of bony tissue.

Osteogenetic (*os-te-o-jen-et'-ik*) [*ὀστέον*, bone; *γενᾶν*, to beget]. Pertaining to, or concerned in, osteogenesis. **O.** Cell, an osteoblast. **O.** Layer, the deep layer of periosteum from which bone is formed.

Osteohalisteresis (*os-te-o-hal-is-ter-et'-sis*) [*ὀστέον*, bone; *ἅλς*, salt; *στερεῖν*, to deprive]. A loss of the mineral constituents of bone.

Osteoid (*os'-te-oid*) [*ὀστέον*, bone; *εἶδος*, like]. Resembling bone.

Osteology (*os-te-ol'-o-je*) [*ὀστέον*, bone; *λόγος*, science]. The science of the anatomy and structure of bones.

Osteolysis (*os-te-ol'-is-is*) [*ὀστέον*, bone; *λύσις*, dissolution]. Absorption of bone.

Osteoma (*os-te-o'-mah*) [*ὀστέον*, bone; *δμα*, tumor]. A bony tumor. **O.** durum, a tumor consisting of hard bony tissue. **O.** eburneum. Synonym of *O. durum*. **O.** medullare, an *O.* containing marrow-spaces. **O.** spongiosum, an *O.* containing cancellated bony tissue.

Osteomalacia (*os-te-o-mal-a'-se-ah*) [*ὀστέον*, bone; *μαλακία*, softness]. Softening of bone from loss of its earthy constituents. It occurs chiefly in adults, especially in women in the course of pregnancy.

Osteomyelitis (*os-te-o-mi-el-i'-tis*) [*ὀστέον*, bone; *μυελός*, marrow; *ιτις*, inflammation]. Inflammation of the marrow of bone.

Osteonecrosis (*os-te-o-ne-kro'-sis*) [*ὀστέον*, bone; *νεκρός*, corpse]. Necrosis of bone.

Osteopathy (*os-te-op'-ath-e*) [*ὀστέον*, bone; *πάθος*, suffering]. Any disease of bone.

Osteoperiostitis (*os-te-o-per-e-os-ti'-tis*) [*ὀστέον*, bone; *περί*, around; *ὀστέον*, bone; *ιτις*, inflammation]. Combined inflammation of the bone and periosteum.

Osteophyte (*os'-te-o-flt*) [*ὀστέον*, bone; *φυτόν*, plant]. A bony outgrowth of dendritic character.

Osteoplastic (*os-te-o-plas'-tik*) [*ὀστέον*, bone; *πλάσσειν*, to form]. 1. Pertaining to the formation of bone. 2. Pertaining to plastic operations upon bone. **O.** Resection, an operation in which a portion of bone, cut loose from its attachments except at one point, is laid back, the underlying diseased structure removed, and the bone replaced. It is also known as the Wagner-Wolff method.

Osteoplasty (*os'-te-o-plus-te*) [*ὀστέον*, bone; *πλάσσειν*, to form]. Plastic operations performed upon bone.

Osteoporosis (*os-te-o-por-o'-sis*) [*ὀστέον*, bone; *πόρος*, a pore]. An enlargement of the spaces of bone whereby a porous appearance is produced.

Osteosarcoma (*os-te-o-sar-ko'-mah*) [*ὀστέον*, bone; *σάρξ*, flesh; *δμα*, tumor]. A sarcoma containing bone.

Osteosclerosis (*os-te-o-skle-ro'-sis*) [*ὀστέον*, bone; *σκληρός*, hard]. A condition in which the bone becomes hard and heavy; it is seen in sclerosing or condensing osteitis.

Osteotome (*os'-te-o-tōm*) [*ὀστέον*, bone; *τομή*, a cutting]. 1. An instrument for cutting bone. 2. An instrument used in cutting the bones of the fetal head in embryotomy.

Osteotomy (*os-te-ot'-o-me*) [*ὀστέον*, bone; *τομή*, a cutting]. The division of a bone. **O.**, Cuneiform, an osteotomy in which a wedge of bone is removed. **O.**, Linear, a simple division of a bone. **O.**, Macewen's. See *Operations*, Table of.

Osteotrite (*os'-te-o-trit*) [*ὀστέον*, bone; *τρίβειν*, to rub]. An instrument for scraping away carious bone.

Ostitis (*os-ti'-tis*). See *Osteitis*.

Ostium (*os'-te-um*) [*L.*]. A mouth or aperture. **O.** abdominale, the orifice of the oviduct communicating with the peritoneal cavity. **O.** internum, the uterine opening of the oviduct. **O.** pharyngeum, the pharyngeal opening of the Eustachian tube. **O.** tympanicum, the tympanic opening of the Eustachian tube. **O.** vaginæ, the external orifice of the vagina.

Otacoustic (*o-ta-koos'-tik*) [*οἰς*, ear; *ἀκοῦεν*, to hear]. 1. Pertaining to, or aiding hearing. 2. An ear-trumpet.

Otalgia (*o-tal'-je-ah*) [*οἰς*, ear; *ἄλγος*, pain]. Earache.

Otaphone (*o'-taf-on*) [*οἰς*, ear; *φωνή*, sound]. A clamp for throwing the pinnæ forward and outward, and thus assisting the hearing.

Othematoma (*ot-hem-at-o'-mah*) [*οἰς*, ear; *αἷμα*, blood; *δμα*, tumor]. Hematoma of the external ear, usually the pinna; hematoma auris. From its comparative frequency in the insane, it is also called insane ear.

Otiatrics (*o-te-at'-riks*) [*oĩs*, ear; *iathp*, surgeon]. The study of diseases of the ear and their treatment.

Otic (*o'-tik*) [*otĩkĩs*, from *oĩs*, ear]. Pertaining to the ear. **O. Ganglion**. See *Ganglia*, Table of.

Otitis (*o-ti'-tis*) [*oĩs*, ear; *itis*, inflammation]. Inflammation of the ear. It may affect the external ear, **O. externa**; the middle ear, **O. media**; or the internal ear, **O. interna**. When confined to the mastoid cells it is called **O. mastoidea**, or mastoid disease.

Oto- (*o'-to-*) [*oĩs*, ear]. A prefix signifying connection with, or relation to the ear.

Otocephalus (*o-to-sef'-al-us*) [*oĩs*, ear; *kefalĩ*, head]. A monster characterized by a union or close approach of the ears, by absence of the lower jaw, and an ill-developed mouth.

Otocleisis (*o-to-kli'-sis*) [*oĩs*, ear; *kleĩsis*, closure]. Occlusion of the ear.

Otoconite (*o-to-k'-o-nit*). See *Otoconium*.

Otoconium (*o-to-ko'-ne-um*) [*oĩs*, ear; *konĩs*, dust]. An otolith.

Otocrane, **Otocranium** (*o'-to-kran*, *o-to-kran'-ne-um*) [*oĩs*, ear; *kravĩn*, skull]. The cavity of the petrous portion of the skull holding the organ of hearing.

Otodynia (*o-to-din'-e-ah*) [*oĩs*, ear; *odĩnĩ*, pain]. Pain in the ear.

Otography (*o-tug'-ra-fe*) [*oĩs*, ear; *grafeĩn*, to write]. Descriptive anatomy of the ear.

Otolith (*o'-to-lith*) [*oĩs*, ear; *lĩthĩs*, stone]. One of the calcareous concretions within the membranous labyrinth of the ear.

Otology (*o-tol'-o-je*) [*oĩs*, ear; *lĩgĩs*, science]. The science of the ear, its anatomy, functions, and diseases.

Otomassage (*o-to-mas'-ah-zĩ*) [*oĩs*, ear; *massage*]. The application of passive motion to the tympanic membrane and auditory ossicles.

Otopathy (*o-top'-ath-e*) [*oĩs*, ear; *patĩs*, disease]. Any affection of the ear.

Otopharyngeal (*o-to-far'-in'-je-al*) [*oĩs*, ear; *pharĩnĩs*, pharynx]. Pertaining to the ear and the pharynx. **O. Tube**, the Eustachian tube.

Otophone (*o'-to-fĩn*) [*oĩs*, ear; *phonĩ*, voice]. 1. An ear-trumpet, or other device for gathering and intensifying sound-waves. 2. An auscultating tube used in ear-diseases.

Otorrhea (*o-tor'-ah*) [*oĩs*, ear; *rhoĩa*, a flow]. A discharge from the external auditory meatus.

Otoscope (*o'-to-skĩp*) [*oĩs*, ear; *skĩpĩn*, to examine]. An instrument for examining the ear, especially a rubber tube, one extremity of which is inserted into the ear of the subject, and the other extremity into the ear of the examiner, a current of air being passed by means of a Politzer bag and a Eustachian catheter through the middle ear. In case of

tympanic perforation the rushing sound made by the passing air is audible to the examiner.

Otoscopy (*o-tĩs'-ko-pe*) [*oĩs*, ear; *skĩpĩn*, to examine]. Examination of the ear, especially by means of the otoscope.

Ototomy (*o-tol'-o-me*) [*oĩs*, ear; *tomĩ*, a cutting]. Dissection of the ear.

Ouabain (*ou-a'-ba-in*), $C_{20}H_{40}O_{12}$. A poisonous glucosid from the wood of *Carissa schimperii*. It is a depressant to the heart and the respiration, and has been used in whooping-cough. Dose gr. $\frac{1}{100}$ (0.00013).

Oulitis (*oo-li'-tis*). See *Ullitis*.

Ounce (*ouns*) [*uncia*, a contraction of *undecia*, a twelfth part]. A unit of measure of weight.

Avoirdupois O., the sixteenth part of the avoirdupois pound or 437.5 grains, equal to 31.1 grams. **Troy O.**, the twelfth part of the Troy pound or 480 grains.

Ouro- (*oo'-ro-*). See *Uro-*.

Oval (*o'-val*) [*ovum*, egg]. Egg-shaped.

Ovalbumin (*o-val-bu'-min*) [*ovum*, egg; *albumin*]. The albumin of the egg.

Ovarialgia (*o-va-re-al'-je-ah*) [*ovarium*, an ovary; *algĩs*, pain]. Neuralgic pain in the ovary.

Ovarian (*o-va'-re-an*) [*ovarium*, an ovary]. Pertaining to the ovaries.

Ovario- (*o-va'-re-o-*) [*ovarium*, ovary]. A prefix denoting relation to the ovary.

Ovariocele (*o-va'-re-o-sel*) [*ovarium*, ovary; *kelĩn*, tumor]. Hernia of an ovary.

Ovariocentesis (*o-va-re-o-sen-te'-is*) [*ovarium*, ovary; *kenĩnĩs*, puncture]. Puncture of the ovary or of an ovarian cyst.

Ovariologist (*o-va-re-al'-o-mĩst*) [*ovarium*, ovary; *tomĩ*, a cutting]. One who performs ovariectomy.

Ovariotomy (*o-va-re-al'-o-me*) [*ovarium*, ovary; *tomĩ*, a cutting]. Literally, incision of an ovary. As generally used, removal of an ovary; oophorectomy. **O., Normal**, Battey's operation, the removal of an ovary that is free from disease.

Ovaritis (*o-var-i'-tis*) [*ovarium*, ovary; *itis*, inflammation]. Oophoritis.

Ovary (*o'-var-e*) [*ovarium*, an egg-holder, from *ovum*, egg]. One of a pair of glandular organs giving rise to ova. It consists of a fibrous framework or stroma, in which are imbedded the Graafian follicles, and is surrounded by a serous covering derived from the peritoneum.

Overextension [*AS., ofer, over; extension*]. Excessive extension; extension beyond the normal point or line.

Overtone [*AS., ofer, over; tone*]. An harmonic tone heard above the fundamental tone.

Oviduct (*o'-vid-ukt*) [*ovum*, egg; *ductus*, a canal]. The Fallopian tube; a small tube upon either side of the uterus through which the ovule passes to the uterus.

Oviferous (*o-vif'-er-us*) [*ovum*, an egg; *ferre*, to bear]. Producing or bearing ova.

Ovification (*o-vif-ik-a'-shun*) [*ovum*, egg; *facere*, to make]. The production of ova.

Ovigerum (*o'-vij-er-um*) [*ovum*, an egg; *germen*, sprout, bud]. A cell producing or developing into an ovum.

Ovigerous (*o-vij'-er-us*) [*ovum*, egg; *gerere*, to carry]. Producing or carrying ova.

Ovination (*o-vin-a'-shun*) [*ovis*, a sheep]. Inoculation with the virus of sheep-pox.

Oviparous (*o-vip'-ar-us*) [*ovum*, egg; *parere*, to bring forth]. Laying eggs; bringing forth young in the egg-stage of development.

Ovisac (*o'-vis-ak*) [*ovum*, egg; *saccus*, sack]. The capsule of an ovum; a Graafian follicle.

Ovoid (*o'-void*) [*ovum*, egg; *ειδος*, like]. Egg-shaped. O., Fetal, the fetal ellipse, or the ellipse formed by the bending of the fetal body in utero.

Ovoviviparous (*o-vo-vi-vip'-ar-us*) [*ovum*, egg; *viviparus*, bringing forth alive]. Reproducing by means of eggs hatched within the body.

Ovule (*o'-vul'*) [*ovum*, egg]. 1. The ovum before its escape from the Graafian vesicle. 2. A small egg; especially a small egg-like body, as the O. of Naboth, one of the small cysts resulting from obstruction of the ducts of the glands of the cervix uteri. O., Migration of, the transfer of the ovule from the ovary to the oviduct.

Ovum (*o'-vum*) [*ovum*, an egg]. The reproductive cell of an animal or vegetable; an egg. A human ovum is a cell consisting of a large amount of protoplasm (vitellus) and a large spheric nucleus, the germinal vesicle, within which is a bright spot, the nucleolus, or germinal spot. It is surrounded by an inner zone, the zona pellucida, and an outer, the vitelline membrane. O., Alecithal, one in which the food-yolk is entirely absent, or present only in very small quantity. O., Blighted, an impregnated ovum the development of which has been arrested by disease or by hemorrhage into the chorion or amniotic cavity. O., Centrolecithal, one in which the formative yolk is arranged in a regular layer around the whole ovum, as well as in a mass at the center in which lies the germinal vesicle. O., Holoblastic, one in which the food-yolk is scant and more or less thoroughly intermingled with the formative yolk, and in which germination is accompanied by a practically uniform segmentation. O., Meroblastic, an ovum with a large amount of food-yolk that takes no active part in the development of the embryo. O., Telolecithal, one in which the food-yolk and the formative yolk divide the egg into two hemispheres, or in which the quantity of the nutritive yolk is greatly in excess.

Oxalate (*oks'-a-lat*) [*ὀξαλῖς*, sorrel]. A salt of oxalic acid.

Oxalic Acid (*oks-al'-ik*) [*ὀξαλῖς*, sorrel]. See *Acid*, *Oxalic*.

Oxaluria (*oks-a-lu'-re-ah*) [*ὀξαλῖς*, sorrel; *urina*, urine]. The presence of an excessive amount of calcium oxalate in the urine.

Oxalyl-urea (*oks-a-lyl-u-re'-ah*) [*ὀξαλῖς*, sorrel; *urea*]. Parabanic acid, a substance produced by oxidizing uric acid or alloxan with nitric acid.

Oxid (*oks'-id*) [*ὀξῖς*, sharp]. A binary compound of oxygen and another element or radicle.

Oxidation (*oks-e-da'-shun*) [*ὀξῖς*, sharp]. The act or process of combining with oxygen.

Oxidize (*oks'-e-diz*) [*ὀξῖς*, sharp]. To combine or to cause to combine with oxygen.

Oxy- (*oks'-e-*) [*ὀξῖς*, sour, sharp]. 1. A prefix denoting sharp or acid. 2. A prefix denoting combined with oxygen.

Oxyacid (*oks'-e-as-id*) [*oxy-*; *acid*]. See *Acid*.

Oxyakoa, Oxyacoa (*oks-e-a-koi'-ah, oks-e-a-ko'-ah*) [*ὀξύς*, sharp; *ἀκοή*, hearing]. Increased acuteness of hearing.

Oxysalt (*oks'-e-salt*) [*oxy-*; *salt*]. A salt of an oxyacid.

Oxybutyric Acid (*oks-e-bu-ti'-rik*) [*ὀξύς*, acid; *butyrum*, butter], $C_4H_8O_4$. A fatty acid found in the urine in certain fevers and in diabetes. Its presence in the body in diabetes, simultaneously with that of diacetic acid, is supposed to be the cause of diabetic coma.

Oxychlorid (*oks-e-klo'-rid*) [*ὀξύς*, acid; *χλωρός*, green]. A compound of a basic element or radicle with both oxygen and chlorine.

Oxydum (*oks'-id-um*). See *Oxid*.

Oxyecoia (*oks-e-e-koi'-ah*). Synonym of *Oxyakoa*.

Oxygen (*oks'-e-jen*) [*ὀξῖς*, acid; *γεννάν*, to produce]. A colorless, tasteless, odorless gas, one of the nonmetallic elements, having an atomic weight of 15.96; quantivalence 11; symbol O; and constituting one-fifth of the atmosphere, eight-ninths of water, three-fourths of organized bodies, and about one-half the crust of the globe; it supports combustion, and is essential to the respiration of animals and plants. It combines with most elements, its combination with the nonmetallic substances giving rise to acids (oxyacids). In medicine it has been employed by inhalation in the treatment of pneumonia, pulmonary tuberculosis, grave anemias, asphyxia, and poisoning by opium.

Oxygenated (*oks'-e-jen-a-ted*) [*ὀξύς*, acid; *γεννάν*, to produce]. Containing or impregnated with oxygen.

Oxyhemoglobin (*oks-e-hem-o-glo'-bin*) [*ὀξύς*,

acid; *alma*, blood; *globus*, a round body]. Oxidized hemoglobin; that found in arterial blood.

Oxyiodid (*oks-e-i'-o-did*) [*ὀξίς*, acid; *ιώδης*, like a violet]. A salt formed by the combination of an element with both oxygen and iodine.

Oxymel (*oks-im-el*) [*ὀξύς*, sharp; *μέλι*, honey]. 1. A mixture of honey and vinegar or dilute acetic acid. 2. Any preparation containing honey and vinegar (or acetic acid) as a vehicle, as **O. of Squill**.

Oxyntic (*oks-in'-tik*) [*ὀξύς*, acid]. Secreting acid. **O.** Cells, cells of the fundus-glands of the stomach supposed to secrete hydrochloric acid.

Oxyopia (*oks-e-o'-pe-ah*) [*ὀξύς*, sharp; *ὤψ*, eye]. Increased acuity of vision.

Oxytotic (*oks-e-tos'-ik*) [*ὀξύς*, sharp; *τόκος*, labor]. 1. Hastening parturition. 2. A drug that hastens parturition.

Oxyuris (*oks-e-u'-ris*) [*ὀξύς*, sharp; *οὐρά*, tail]. A genus of nematode worms, the pinworms.

O. vermicularis, the common seat-worm or pinworm infesting the rectum of children.

Ozena (*o-ze'-nah*) [*ὀζή*, a stench]. Chronic disease of the nose accompanied by a fetid discharge, and depending on atrophic rhinitis or syphilitic ulceration or caries.

Ozocerite (*o-zo-se'-rit*) [*ὀζή*, smell; *κηρός*, wax]. A solid paraffin found free in Galicia and Roumania, and used in diseases of the skin.

Ozone (*o'-zōn*) [*ὀζειν*, to smell]. An allotropic form of oxygen, the molecule of which consists of three atoms. It occurs free in the atmosphere, and is a powerful oxidizing agent. In medicine it is employed as a disinfectant.

Ozonized (*o'-zo-nīzd*) [*ὀζειν*, to smell]. Containing ozone.

P

P. 1. The symbol of phosphorus. 2. An abbreviation for (1) Position; (2) Punctum proximum (near-point).

p. Abbreviation for para-.

Pabulum (*pab'-u-lum*) [*L.*]. Food.

Pacchionian (*pak-e-o'-ne-an*). Described by *Pacchioni*, an Italian anatomist (1665-1726). **P. Bodies**, **P. Glands**, small fibrous outgrowths of the arachnoid which perforate the dura and by pressure produce depressions in the cranium (**P. Depressions**). The bodies are found along the superior longitudinal sinus.

Pachy- (*pak'-e*) [*παχύς*, thick]. A prefix meaning thick.

Pachycephaly (*pak-e-sef'-al-e*) [*παχύς*, thick; *κεφαλή*, head]. Abnormal thickness of the skull.

Pachydermatocoele (*pak-e-der-mat'-o-sel*) [*παχύς*, thick; *δέρμα*, skin; *κύλη*, tumor]. A tumor due to thickening of the skin.

Pachydermatous (*pak-e-der'-mat-us*) [*παχύς*, thick; *δέρμα*, skin]. Thick-skinned.

Pachyderma, **Pachydermia** (*pak-e-der-mah*, *pak-e-der'-me-ah*) [*παχύς*, thick; *δέρμα*, skin]. 1. Thickening of the skin. 2. Elephantiasis. **P. laryngis**, extensive thickening of the mucous membrane of the larynx.

Pachylosis (*pak-e-lo'-sis*) [*παχύς*, thick]. A condition of the skin, especially of the

legs, in which it is thick, dry, harsh, and scaly.

Pachymeningitis (*pak-e-men-in-jī'-tis*) [*παχύς*, thick; *μηνίς*, membrane; *ιτις*, inflammation]. Inflammation of the dura of the brain (Cerebral **P.**) or the spinal cord (Spinal **P.**). **P., External**, that affecting the external layer of the dura. **P., Hemorrhagic**, **P. hæmorrhagica interna**, an effusion of blood on the inner surface of the dura. **P., Internal**, that involving the internal layer of the dura. **P., Syphilitic**, that due to syphilis.

Pacini's Fluid (*pah-tshe'-nēz*). A diluting fluid used in counting the red blood-corpuscles. Its composition is: Mercuric chlorid 2 grams; sodium chlorid 4 grams; glycerol 26 c. c.; distilled water 226 c. c. This is diluted with three parts of water before using.

Pacinian (*pa-sin'-e-an*, *pak-tshin'-e-an*). Described by *Pacini*, an Italian anatomist. **P. Corpuscles**, elliptic, semitransparent bodies, that represent one form of termination of the sensory nerves of the skin, the peritonicum, and other parts.

Pack (*pak*). A blanket, either dry or soaked in hot or cold water, and wrapped about the body. **P., Cold**, a blanket wrung out of cold water and wrapped about the body. **P., Hot**, a blanket wrung out of hot water and wrapped about the body. **P., Wet**, a

blanket wrung out of warm or cold water, and wrapped about the body.

Packing (*pak'-ing*). 1. The act of filling a wound or cavity with gauze or other material. 2. The material used for filling the cavity.

Pacquin Cautery (*pah'-kel-ang*). See *Cautery*.

Pagenstecher's Ointment [*Pagenstecher*, a German ophthalmologist]. An ointment of yellow oxid of mercury, used in ophthalmic practice.

Paget's Abscess (*pag'-etz*) [Sir J. Paget, an English surgeon]. A residual abscess. **P. Disease of the Nipple**. See *Diseases*, *Table of*.

Pain (*pān*) [*pāna*, punishment, pain]. 1. Bodily suffering due to irritation of a sensory nerve, or possibly in rare cases to changes in the central nervous system. 2. One of the rhythmic contractions of the uterus during labor. **P., After-**, that following labor, and caused by the uterus contracting to expel clots, etc. **P., Bearing-down**, pain with a sensation of dragging or bearing down of the pelvic organs, occurring in labor and in various inflammatory affections of the female pelvic organs. **P., False**, that occurring in the latter part of pregnancy and resembling labor-pain, although not followed by labor. **P., Fulgurant**, **P., Fulgurating**, the intense shooting pain affecting principally the limbs of patients suffering from locomotor ataxia. **P., Girdle-**, a painful sensation as of a cord tied about the waist; it is a symptom of organic disease of the spinal cord. **P., Growing**, a popular term for the soreness about the joints in young persons at puberty. Some attribute it to increased vascularity of the epiphyses of long bones; others, to rheumatism. **P., Lancinating**. See *P., Fulgurant*. **P., Starting**, pain caused by a spasmodic contraction of the muscles just before the onset of sleep. It occurs in joint-diseases when the cartilages are ulcerated.

Painters' Colic. Lead-colic.

Palatal (*pal'-at-al*) [*palatum*, palate]. Pertaining to the palate.

Palate (*pal'-at*) [*palatum*, palate]. The roof of the mouth. It is composed of the hard palate, formed by the palatal processes of the superior maxillary bones and the palatal bones, and the soft palate, or velum palati, consisting of an aggregation of muscles, the tensor palati, azygos uvulae, palatoglossus, and palatopharyngeus. **P., Artificial**, a plate of hard material used as an obturator to close a fissure in the palate. **P., Cleft**, a congenital deformity characterized by incomplete closure of the lateral halves of the palate.

Palatine (*pal'-a-tin*) [*palatum*, palate]. 1. Pertaining to the palate, as the P. arteries. 2. Conveying the palatine vessels or nerves.

Palatoglossal (*pal'-a-to-glos'-al*) [*palatum*, palate; γλῶσσα, tongue]. Pertaining to the palate and the tongue.

Palatoglossus (*pal'-a-to-glos'-us*). See *Muscles*, *Table of*.

Palatonasal (*pal'-a-to-na'-zal*) [*palatum*, palate; *nasus*, nose]. Pertaining to the palate and the nose.

Palatopharyngeal (*pal'-a-to-far-in'-je-al*) [*palatum*, palate; φάρυγξ, pharynx]. Pertaining conjointly to the palate and the pharynx.

Palatopharyngeus (*pal'-a-to-far-in'-je-us*). See *Muscles*, *Table of*.

Palatorrhaphy (*pal'-a-tor'-a-fe*). See *Staphylorrhaphy*.

Palatosalpingeus (*pal'-a-to-sal-pin'-je-us*). The tensor palati muscle.

Palatoschisis (*pal'-a-tos'-kiss-is*) [*palatum*, palate; σχισμα, cleft]. Cleft palate.

Palatouvularis (*pal'-a-to-u-vu-la'-ris*). The azygos uvulae muscle.

Palaeontology (*pa-le-on-tol'-o-je*) [παλαιός, old; ὄν, being; λόγος, science]. The science of the early life-forms of the earth.

Palingenesis (*pal'-in-jen'-es-is*) [παλιν, again; γένεσις, production]. The form of development of an individual germ in which the development of its ancestors is succinctly repeated.

Palliative (*pal'-e-a-tiv*) [*palliare*, to cloak]. 1. Relieving or alleviating suffering. 2. A drug relieving or soothing the symptoms of a disease without curing it.

Palm (*palm*) [*palma*, palm]. 1. The inner or flexor surface of the hand; the hollow of the hand. 2. A palm-tree. **P.-oil**, oleum palmae, a fixed oil obtained from the fruit of *Elais guineensis*. It is employed in making soap.

Palma (*pal'-mah*) [*L.*]. Palm. **Palmæ plicatæ**, the arborescent rugæ of the interior of the vagina.

Palmar (*pal'-mar*) [*palma*, palm]. Pertaining to the palm of the hand. **P. Abscess**, an abscess in the palm of the hand, usually situated beneath the palmar fascia. **P. Arch**, one of the two curved arches, superficial and deep, formed by the anastomosis of the radial and ulnar arteries in the hand. **P. Fascia**, the sheath investing the muscles of the hand.

Palmaris (*pal'-ma'-ris*). See *Muscles*, *Table of*.

Palmella (*pal-mel'-ah*). A genus of fresh-water algae.

Palmellin (*pal-mel'-in*). A red coloring-principle of a fresh-water alga, the *Palmella cruenta*, resembling hemoglobin.

Palmitic Acid (*pal-mit'-ik*). See *Acid*.

Palmitin (*pal'-mit-in*) [*palme*, palm-tree], $C_{15}H_{31}(C_{15}H_{31}O_2)_2$. Glyceryl tripalmitate, a solid crystallizable substance which, with stearin, constitutes the greater proportion of solid fats.

Palms (*pal'-mss*) [*palme*, a twitch]. Jumpers' disease, lata, or miryachit; a form of convulsive tic.

Palpate (*pal'-pāt*) [*palparr*, to feel]. To examine by touch.

Palpation (*pal-pā'-shun*) [*palparr*, to feel]. In physical diagnosis, the laying of the hand on a part of the body or the manipulation of a part with the hand for the purpose of ascertaining its condition or that of underlying organs. P., Bimanual, the use of the two hands in examining an organ.

Palpebral (*pal'-pe-bral*) [*palpebra*, eyelid]. Pertaining to the eyelid. P. Conjunctiva, the conjunctiva of the eyelid. P. Fissure, the opening between the upper and lower eyelids.

Palpitation (*pal-pit'-e'-shun*) [*palpitarr*, to quiver]. A fluttering or throbbing, especially of the heart, of which the person is conscious.

Palsy (*paol'-se*) [from *paralysis*]. Paralysis. P., Bell's. See *Diseases, Table of*. P., Birth-. See *Birth*. P., Crutch-, that due to pressure of the crutch in the axilla upon the nerves of the brachial plexus. P., Erb's. See *Erb's Palsy*. P., Hammer-, that due to excessive use of the hammer. P., Shaking. Synonym of *Paralysis agitans*. P., Wasting, progressive muscular atrophy.

Paludal (*pal'-u-dal*) [*palus*, marsh]. Pertaining to, or originating in marshes; malarial.

Paludism (*pal'-u-dizm*) [*palus*, marsh]. Malarial poisoning.

Pampiniform (*pam-pin'-e-form*) [*pampinus*, tendril; *forma*, form]. Having the form of a tendril. P. Plexus. See *Plexus, Spermatic*.

Pampinocoele (*pam-pin'-o-sel*) [*pampinus*, tendril; *κήλη*, tumor]. A varicocele of the veins of the pampiniform plexus.

Panacea (*pan-a-se'-ah*) [*πανακεια*, all-healing, from *πᾶς*, all; *ἄκος*, a cure]. A remedy curing all diseases; a cure-all.

Panama-fever. 1. A pernicious form of malarial fever occurring on the isthmus of Panama. 2. Yellow fever. P.-paralysis, beriberi.

Panaris (*pan-a'-ris*) [*panaricium*, a corruption of *paronychium*, paronychia]. See *Paronychia*.

Panaritium (*pan-ar-ish'-e'-um*). See *Paronychia*.

Panarthrit (*pan-ar-thrit'-tis*) [*πᾶς*, all; *ἄρθρον*, joint; *ιτις*, inflammation]. Inflammation of all the structures of a joint.

Panax (*pa'-naks*) [*πᾶς*, all; *ἄκος*, cure]. Ginseng; the root of *P. quinquefolium*, used in China as a remedy for many diseases. It is a demulcent and mild purgative.

Pancreas (*pan'-kre-as*) [*πᾶς*, all; *κρέας*, flesh]. A compound racemose gland, from six to eight inches in length, lying transversely across the posterior wall of the abdomen; the sweetbread of animals; the abdominal salivary gland of the Germans. Its right extremity, the head, lies in contact with the duodenum; its left extremity, the tail, is in close proximity to the spleen. It secretes a limpid, colorless fluid that digests proteins, fats, and carbohydrates. The secretion is conveyed to the duodenum by the pancreatic duct, or duct of Wirsung. P., Lesser, a small, partially detached portion of the gland, lying posteriorly to its head and having occasionally a separate duct that opens into the pancreatic duct proper.

Pancreatic (*pan-kre-at'-ik*) [*πᾶς*, all; *κρέας*, flesh]. Pertaining to the pancreas, as the P. duct; depending upon disease of the pancreas, as P. diabetes. P. Fluid or Juice, the secretion of the pancreas, a thick, transparent, colorless, odorless fluid, of a salty taste, and strongly alkaline.

Pancreaticoduodenal (*pan-kre-at-ik-o-du-od'-en-al*) [*πᾶς*, all; *κρέας*, flesh; *duodenum*, duodenum]. Pertaining to the pancreas and the duodenum, as the P. arteries.

Pancreatin (*pan-kre-at-in*) [*πᾶς*, all; *κρέας*, flesh]. The active elements of the juice of the pancreas; also the commercial extract of the pancreas, supposed to possess a fermentative action similar to that of the pancreatic juice.

Pancreatitis (*pan-kre-at-i'-tis*) [*πᾶς*, all; *κρέας*, flesh; *ιτις*, inflammation]. Inflammation of the pancreas. It may be hemorrhagic, suppurative, or gangrenous. The onset of P. is usually sudden, with severe abdominal pain, vomiting, tympanites, and tenderness of the abdomen. It is usually fatal.

Pancreatotomy (*pan-kre-at-o'-o-me*) [*πᾶς*, all; *κρέας*, flesh; *τομή*, a cutting]. Incision of the pancreas.

Pandemic (*pan-dem'-ik*) [*πᾶς*, all; *δῆμος*, people]. Epidemic over a wide area.

Pandiculation (*pan-dik-u-la'-shun*) [*pandiculari*, from *pandere*, to stretch out]. The act of stretching the limbs, especially on waking from sleep.

Pang, Brow-. Hemisrania.

Pangenesis (*pan-jen'-es-is*) [*πᾶς*, all; *γεννᾶν*, to produce]. Darwin's theory of heredity, which supposes the existence of gemmules or minute particles separated from the body-cells and segregated from the circulation by the reproductive glands. These preformed constituents of all parts of the fully-formed ani-

mal or plant become aggregated in the germ, and give rise by a process of evolution to the new organism.

Panneuritis (*pan-nu-ri'-tis*) [*πᾶς*, all, every; *νεῖρον*, nerve; *ιτις*, inflammation]. Multiple neuritis.

Panniculus (*pan-ik'-u-lus*) [dim. of *pan-nus*, a piece of cloth]. A membrane or layer. *P. adiposus*, the layer of subcutaneous fat. *P. carnosus*, the layer of muscles contained in the superficial fascia. It is well developed in the lower animals, but in man is represented mainly by the platysma.

Pannus (*pan'-us*) [L., a cloth]. Vascularization of the cornea, usually due to the irritation of trachoma-granulations.

Panodic (*pan-od'-ik*). See *Panthodic*.

Panophobia (*pan-o-fō'-be-ah*) [*πᾶς*, all; *φόβος*, fear]. Morbid fear of everything; a symptom present in some cases of neurasthenia.

Panophthalmitis (*pan-off-thal-mi'-tis*) [*πᾶς*, all; *ὀφθαλμός*, eye; *ιτις*, inflammation]. Inflammation of all the tissues of the eyeball.

Panosteitis (*pan-os-te-i'-tis*) [*πᾶς*, all; *ὀστέον*, bone; *ιτις*, inflammation]. An inflammation of all the structures of a bone.

Panotitis (*pan-o-ti'-tis*) [*πᾶς*, all; *οὖς*, ear; *ιτις*, inflammation]. An inflammation involving all the structures of the ear.

Panspermatism (*pan-sper'-mat-ism*) [*πᾶς*, all; *σπέρμα*, seed]. The theory that germs are omnipresent.

Panaphygmograph (*pan-sfig'-mo-graf*) [*πᾶς*, all; *σφυγμός*, pulse; *γράφειν*, to write]. An instrument by means of which tracings can be taken simultaneously of the cardiac movements, the arterial pulse, and the respiration.

Panthodic (*pan-thod'-ik*) [*πᾶς*, all; *ὁδός*, way]. Of nervous impulses, radiating to all parts of the body.

Pantophobia (*pan-to-fō'-be-ah*). See *Panophobia*.

Pantoscopic (*pan-to-skop'-ik*). See *Bifocal*.

Papain (*pa'-pa-in*). See *Papaya*.

Papaver (*pa-pa'-ver*). See *Opium*.

Papaverin (*pa-pa'-ver-in*) [*papaver*, poppy], $C_{21}H_{21}NO_4$. A crystalline alkaloid found in opium and thought to possess narcotic properties. Dose gr. $\frac{1}{4}$ (0.016).

Papaya (*pa-pa'-yah*). Melon-tree; pawpaw—the *Carica papaya*, a tree of the order Passifloræ. The unripe fruit yields a milky juice containing an albuminous substance, papain or papayotin, capable of digesting fibrin and other proteid-bodies. Papain in commerce occurs as a grayish powder, and has been used as a digestant in dyspepsia, as an application to false membranes, warts, epitheliomata, etc. Dose gr. v-x (0.32-0.65).

Papayotin (*pa-pa-yo'-tin*). See *Papaya*.

Papilla (*pap-il'-ah*) [L., a nipple]. 1. A small, nipple-like eminence. 2. Synonym of *Optic Disc*. *P.*, Circumvallate, one of the large papillæ at the root of the tongue, arranged like the letter V opening forward. *P.*, Clavate. Synonym of *P.*, *Fungiform*. *P.*, Conical. See *P.*, *Filiform*. *P.*, Dental. See *Organ*, *Enamel*. *P.*, Duodenal, the elevation at the point where the ductus communis choledochus enters the duodenum. *P.*, Filiform, any one of the papillæ occurring on all parts of the tongue, consisting of an elevation of connective tissue covered by a layer of epithelium. *P.*, Fungiform, any one of the low, broad papillæ found on the surface of the tongue, consisting of a connective-tissue elevation, covered by secondary papillæ. *P.*, Lacrimal, a small conic eminence on the eyelid at the inner canthus, pierced by the lacrimal punctum. *P.*, Renal, the summit of any one of the renal pyramids projecting into the renal pelvis.

Papillary (*pap'-il-a-re*) [*papilla*]. Composed of or containing papillæ; resembling a papilla.

Papilliferous (*pap-il-iff'-er-us*) [*papilla*, papilla; *ferre*, to bear]. Bearing or containing papillæ, as a *P. cyst*.

Papilliform (*pap-il'-if-orm*) [*papilla*, nipple; *forma*, to form]. Shaped like a papilla.

Papillitis (*pap-il-i'-tis*) [*papilla*, papilla; *ιτις*, inflammation]. Inflammation of the optic disc; optic neuritis; "choked disc."

Papilloma (*pap-il-o'-mah*) [*papilla*, papilla; *δμα*, tumor]. A growth on the skin or mucous membrane resembling hypertrophied papillæ. It is a benign tumor, occurring in two forms, the hard *P.*, one growing from squamous epithelium, and the soft *P.*, one developed from columnar epithelium.

Papule (*pap'-ul*) [*papula*, a pimple]. A small circumscribed, solid elevation of the skin. *P.*, Moist, the syphilitic condyloma.

Paquelin's Cautery. See *Cautery*.

Par [L.]. A pair. *P. vagum*, the vagus nerves.

Para- (*par'-ah-*) [*παρά*, beyond, beside]. 1. A prefix signifying beyond, beside, near, the opposite of, etc. 2. In chemistry, prefixed to a derivative of the benzol-ring, it indicates the substitution of two atoms of hydrogen situated opposite each other.

Parabanic Acid (*par-ah-ba'-nik*). See *Oxalyl-urea*.

Parablast (*par'-ah-blast*) [*παρά*, beside; *βλαστός*, germ]. One of the two main divisions of the embryonal tissues, from which the blood-vessels, lymphatic vessels, and other connective tissues are developed.

Parablastic (*par-ah-blas'-tik*) [*παρά*, beside; *βλαστός*, germ]. Pertaining to the parablast.

Parablastoma (*par-ah-blas-to'-mah*) [*παρά*,

beside; βλαστόν, germ; όμα, tumor]. A tumor composed of parablatic tissue.

Paracentesis (*par-ah-sen-te'-sis*) [παρά, beside; κέντησις, puncture]. Puncture; especially puncture of the wall of a cavity of the body, such as the thoracic wall, cornea, tympanic membrane.

Paracentral (*par-ah-sen'-tral*) [παρά, beside; κέντρον, center]. Situated near the center. **P. Lobule**, a convolution on the mesial surface of the cerebral hemisphere uniting the upper ends of the ascending frontal and ascending parietal convolutions.

Paracephalus (*par-ah-sef'-al-us*) [παρά, beside; κεφαλή, head]. A monster characterized by a rudimentary, misshapen head, and defective trunk and limbs.

Parachroma (*par-ah-kro'-mah*) [παρά, beside; χρώμα, color]. Change in color, especially in the natural color of the skin.

Paracolpitis (*par-ah-kol-pi'-tis*) [παρά, by; κόλπος, vagina; ιτις, inflammation]. Inflammation of the connective tissue about the vagina.

Paracoto (*par-ah-ko'-to*) [παρά, beside; Sp., coto, a cubit]. A South American tree the bark of which contains a neutral substance, paracotoin, closely resembling cotoin.

Paracresylol (*par-ah-kres'-il-ol*) [παρά, beside; cresol]. A derivative of cresol.

Paracusia, **Paracusis** (*par-ah-ku'-ze-ah*, *par-ah-ku'-sis*) [παρά, beside; ακούειν, to hear]. Perversion of the sense of hearing.

Paracystitis (*par-ah-sis-ti'-tis*) [παρά, beside; κύστις, bladder; ιτις, inflammation]. Inflammation of the connective tissue surrounding the bladder.

Paradidymis (*par-ah-did'-im-is*) [παρά, beside; διδυμος, testicle]. The organ of Giraldès, the atrophic remains of the tubules of the Wolffian body, lying among the convolutions of the epididymis.

Paradoxic Contraction. A slow tonic contraction occurring in a muscle when suddenly relaxed or when its length is suddenly shortened. **P. Pulse**. See *Pulse*.

Paræsthesia (*par-es-the'-ze-ah*). See *Paræsthesia*.

Paraffin (*par'-af-in*) [*parum*, little; *affinis*, affinity]. 1. Any saturated hydrocarbon of the marsh-gas series, having the formula C_nH_{2n+2} . 2. A white, odorless, translucent hydrocarbon obtained from coal-tar or by the destructive distillation of wood. **P. durum** (B. P.), hard or solid paraffin, a mixture of several of the harder members of the paraffin-series of hydrocarbons. It is usually obtained by distillation from shale. **P., Liquid**, a liquid hydrocarbon of the paraffin-series. **P. molle**, soft paraffin; the Petrolatum molle of the (U. S. P.).

Paragrammacismus (*par-ah-gam-a-sis'-mus*)

[παρά, beside; γάμμα, the letter g]. Inability to pronounce the hard g, and also k, other consonants being substituted, as d or t. **Parageusia** (*par-ah-gu'-ze-ah*) [παρά, beside; γείσις, taste]. Perversion of the sense of taste.

Paraglobulin (*par-ah-glob'-u-lin*) [παρά, beside; globulus, a little ball]. A globulin found in blood-serum and other fluids of the body. It is also called fibroplastin, fibrinoplastin, and serum-globulin.

Paraglobulinuria (*par-ah-glob'-u-lin-u'-re-ah*) [παρά, beside; globulus, a little ball; urina, urine]. The presence of paraglobulin in the urine.

Paragraphia (*par-ah-graf'-e-ah*) [παρά, aside; γράφειν, to write]. A form of aphasia in which the person writes the improper word or misplaces the words.

Paralactic Acid (*par-ah-lak'-tik*). See *Acid, Sarcolactic*.

Paralalia (*par-ah-la'-le-ah*) [παρά, beside; λαλία, speech]. Disturbance of the faculty of speech.

Paralbumin (*par-al-bu'-min*) [παρά, beside; albumin]. A proteid substance found in ovarian cysts.

Paraldehyd (*par-al'-de-hid*) [παρά, beside; aldehyd, $C_6H_{12}O_3$]. A polymeric form of aldehyd, occurring as a colorless liquid of repulsive odor and unpleasant taste, with powerful hypnotic properties. It is used in delirium tremens, mania, tetanus, and other nervous affections. Dose, m 30-60 (2.0-4.0).

Paralexia (*par-ah-lek'-e-ah*) [παρά, beside; λήξις, speech]. Disturbance of the power of reading, consisting in the transposition or substitution of words or syllables.

Paralgnesia (*par-al-je'-ze-ah*) [παρά, beside; άλγος, pain]. An abnormal painful sensation; painful paresthesia.

Parallax (*par'-al-aks*) [παρά, beside; ἀλλος, other]. The apparent displacement of an object due to a change in the position of the observer, or by looking at it alternately with one eye, then with the other. **P., Crossed**, **P., Heteronymous**, that in which the object moves away from the uncovered eye. **P., Homonymous**, that in which the object moves toward the uncovered eye. **P., Vertical**, that in which the object moves upward or downward.

Paralogia (*par-ah-lo'-je-ah*) [παρά, beside; λόγος, reason]. Difficulty in thinking logically. **P., Thematic**, a condition in which the thought is unduly concentrated on one subject.

Paralysis (*par-al'-is-is*) [παρά, beside; λύνειν, to loosen]. A loss of motion or of sensation in a part. **P., Acute Ascending**. See *Landry's Paralysis*, in *Diseases*, Table of. **P., Acute Atrophic**. See *P., Infantile*.

P., Acute Progressive. Synonym of *P., Acute Ascending*. **P., agitans**, shaking palsy, Parkinson's disease. **P., Atrophic Spinal.** See *P., Infantile*. **P., Atrophospastic.** Synonym of *Amyotrophic Lateral Sclerosis*. **P., Bell's.** See *P., Facial*. **P., Brown-Séquard's**, a motor paralysis of one side of the body with sensory paralysis of the other side. **P., Bulbar**, glossolabio-laryngeal paralysis. **P., Central**, a paralysis due to a lesion of the brain or spinal cord. **P., Cerebral**, a paralysis due to a brain-lesion. **P., Crossed**, a paralysis of the arm and leg of one side, associated with either a facial paralysis or a paralysis of the oculomotor nerve of the opposite side. **P., Crutch-**. See *Crutch-paralysis*. **P., Cruveilhier's**, progressive muscular atrophy. **P., Diphtheric**, a motor paralysis due to the action of the diphtheria-poison on the nervous system, chiefly on the peripheral nerves. **P., Duchenne's.** See *Duchenne's Disease in Diseases, Table of*. **P., Erb's**, a partial paralysis of the brachial plexus, involving the nerves supplying the deltoid, biceps, brachialis anticus, and supinator longus, often the supinator brevis, and occasionally the infraspinatus and subscapularis muscles. **P., Facial**, a paralysis of the muscles of the face, usually of one side only, due to central disease or due to a lesion of the facial nerve. **P., General (of the Insane)**, general paresis, paralytic dementia, paretic dementia, progressive paralysis of the insane; an organic disease of the brain, characterized by progressive loss of power and by a deterioration of the mental faculties, ending eventually in dementia and death. The main symptoms may be divided into psychic, motor, and sensory. The psychic symptoms are principally a change of character and delusions of grandeur; the motor are weakness, tremor, disturbance of speech, apoplectic or epileptiform seizures, and finally motor paralysis; there is often inequality of the pupils, with myosis or mydriasis; sensory symptoms are slight and consist chiefly in paresthesias. The causes are obscure—syphilis and severe nervous strain are important factors. **P., Infantile**, acute anterior poliomyelitis, acute atrophic paralysis, atrophic spinal paralysis, essential paralysis; a disease peculiar to childhood, and characterized by sudden paralysis of one or more limbs or of individual muscle-groups, and followed by rapid wasting of the affected parts, with reaction of degeneration and deformity. The paralysis is due to changes in the anterior cornua of the gray matter of the spinal cord, and is probably the result of infection. **P., Ischemic**, paralysis of a part due to stoppage of the circulation, *e. g.*, paralysis of the

lower limb following embolism or thrombosis of the femoral artery. **P., Klumpke's**, a paralysis involving the lower portion of the brachial plexus, the eighth cervical and first dorsal nerves, and characterized by paralysis of the small muscles of the hand, of some of the muscles of the forearm, with anesthesia in the distribution of the ulnar and median nerves. Pupillary changes may be present. **P., Lead-**, a paralysis due to lead, usually of the extensors of the wrist, causing wrist-drop. It is nearly always bilateral and is caused by a peripheral neuritis induced by the lead-poisoning. **P., Little's**, infantile cerebral spastic paraplegia. **P., Motor**, paralysis of the voluntary muscles. **P., Musculospiral**, paralysis of the extensors and supinators of the wrist, due to an injury or to inflammation of the musculospiral nerve. **P., Nuclear**, one due to a lesion of the nuclei of origin of a cranial nerve. **P., Obstetric**, any paralysis of the child resulting from injuries received during delivery. **P., Peripheral**, loss of power due to a lesion of the nervous motor mechanism between the nuclei of origin and the peripheral termination. **P., Postdiphtheric.** See *P., Diphtheric*. **P., Pseudobulbar**, a symmetric lesion of the halves of the cerebrum producing paralysis of the lips, the tongue, and the larynx or the pharynx. **P., Pseudohypertrophic Muscular**, a chronic disease characterized by progressive muscular weakness, associated with an apparent hypertrophy of the affected muscles. The disease usually begins in the muscles of the calf, and spreads over the body, the muscles of the hand almost always escaping. There is marked lordosis and a peculiar gait with wide separation of the legs and swaying of the body from side to side. The characteristic symptom is the manner in which the patient arises from the floor—he "climbs" up on his legs, on account of the weakened state of the extensor muscles of the back. It is most common in young male children. The disease is entirely myopathic, no adequate nerve-lesion having as yet been discovered. The muscles are the seat of hypertrophy and atrophy of muscular fibers, hyperplasia of the connective tissue, and fatty infiltration. **P., Reflex**, the paralysis sometimes following immediately upon a wound of a nerve, or the paraplegia sometimes due to irritation of an adherent prepuce. Other so-called reflex palsies, as that from renal calculus, are probably due to secondary changes in the spinal cord or nerves. **P., Spastic**, a paralysis associated with rigidity of the muscles and heightened tendon-reflexes. **P., Spastic Spinal**, lateral sclerosis.

Paralytic (*par-ah-lit'-ik*) [*παράλυσις*, paralysis]. 1. Of the nature of paralysis; affected with paralysis. 2. A person suffering from paralysis; also, one suffering from general paralysis of the insane.

Paralyzant (*par-ah-li'-zant*) [*παράλυσις*, paralysis]. 1. Causing paralysis. 2. An agent or drug that induces paralysis. **P., Motor**, a drug paralyzing any part of the motor apparatus.

Paralyzing Vertigo. See *Gerlier's Disease*, *Diseases*, *Table of*.

Paramagnetic (*par-ah-mag-net'-ik*) [*παρά*, beside; *magnet*]. Exhibiting a polarity in the same direction as the magnetizing force.

Paramagnetism (*par-ah-mag'-net-izm*) [*παρά*, beside; *magnet*]. The phenomena exhibited by paramagnetic substances.

Paramastitis (*par-ah-mas-tit'-tis*) [*παρά*, near; *μαστός*, mamma; *ιτις*, inflammation]. Inflammation of the connective tissue about the mamma.

Paramenia (*par-ah-me'-ne-ah*) [*παρά*, beside; *μήνεις*, menses]. Difficult or disordered menstruation.

Parametric (*par-ah-met'-rik*) [*παρά*, beside; *μήτρα*, uterus]. Pertaining to the tissues about the uterus.

Parametritis (*par-ah-me-trit'-tis*) [*παρά*, near; *μήτρα*, womb; *ιτις*, inflammation]. Inflammation of the cellular tissue about the uterus; pelvic cellulitis.

Parametrium (*par-ah-me'-tre-um*) [*παρά*, near; *μήτρα*, womb]. The connective tissue surrounding the uterus.

Paramimia (*par-ah-mim'-e-ah*) [*παρά*, beside; *μιμῆσθαι*, to mimic]. A form of aphasia characterized by the faulty use of gestures.

Paramitome (*par-ah-mit'-tōm*) [*παρά*, beside; *μίτος*, thread]. The fluid portion of the cell-substance, contained in the meshes of the mitome.

Paramnesia (*par-am-ne'-ze-ah*) [*παρά*, beside; *μνησκειν*, to remind]. Illusion of memory, especially the illusion of feeling, as if one had already undergone the experience which may be passing.

Paramorphin (*par-ah-mor'-fin*). See *Thebain*.

Paramusia (*par-ah-mu'-ze-ah*) [*παρά*, beside; *μουσική*, music]. A form of aphasia in which there is perversion of the musical sense, resulting in the production of improper notes and intervals.

Paramyoclonus multiplex (*par-ah-mi-ok'-lo-nus mul'-tip-leks*). A neurosis marked by sudden, shock-like muscular contractions, which are bilateral and do not as a rule affect the hands or face. The etiology is unknown, and the disease is believed to be analogous to chronic adult chorea.

Paramyosinogen (*par-ah-mi-o-sin'-o-jen*) [*παρά*, beside; *μύς*, muscle; *γεννάν*, to beget]. One of the proteids of muscle-plasma, coagulating at 47° C.

Paramyotonia (*par-ah-mi-o-to'-ne-ah*) [*παρά*, aside; *μύς*, muscle; *τόνος*, tone]. A perversion of muscular tonicity, characterized by tonic spasms. It is usually congenital (**P. congenita**). See *Thomsen's Disease*, *Diseases*, *Table of*.

Paranephritis (*par-ah-nef-rit'-tis*) [*παρά*, beside; *νεφρός*, kidney; *ιτις*, inflammation]. 1. Inflammation of the paranephros. 2. Inflammation of the connective tissue about the kidney.

Paranephros (*par-ah-nef'-ros*) [*παρά*, beside; *νεφρός*, kidney]. The suprarenal capsule.

Parangi (*par-an'-je*). See *Frambesia*.

Paranea, **Paranoia** (*par-ah-ne'-ah*, *par-ah-noi'-ah*) [*παρά*, beside; *νόος*, mind]. Mental aberration, especially a chronic disease characterized by systematized delusions.

Paranoiac (*par-ah-noi'-ak*) [*παρά*, beside; *νόος*, mind]. 1. Affected with paranoia. 2. A person who is affected with paranoia.

Paranucleus (*par-ah-nu'-kle-us*) [*παρά*, beside; *nucleus*, kernel]. An irregular spherule body lying in the protoplasm of a cell near the nucleus and perhaps extruded by the latter.

Parapeptone (*par-ah-pep'-tōn*). See *Pep-tone*.

Paraphasia (*par-ah-fa'-ze-ah*) [*παρά*, beside; *φάσις*, speech]. A form of aphasia in which there is inability to connect ideas with the proper words to express the ideas.

Paraphimosis (*par-ah-fī-mo'-sis*) [*παρά*, beside; *φιδεα*, to bridle]. Retraction and constriction of the prepuce behind the glans penis.

Paraphonia (*par-ah-fo'-ne-ah*) [*παρά*, beside; *φωνή*, voice]. Any abnormal condition of the voice.

Paraphrasia (*par-ah-fra'-ze-ah*) [*παρά*, beside; *φράσις*, utterance]. A form of aphasia characterized by incoherence of speech.

Paraplast (*par'-ah-plasm*) [*παρά*, beside; *πλάσμα*, a thing formed]. 1. The fluid substance in the meshes of the cell-protoplasm. 2. A heteroplasm or false growth.

Paraplegia (*par-ah-ple'-je-ah*) [*παρά*, beside; *πληγή*, stroke]. Paralysis of the lower half of the body, or of the lower extremities. **P., Ataxic**, a disease characterized clinically by a combination of ataxia and exaggerated tendon-reflexes, and anatomically by sclerosis of the posterior and lateral columns of the cord. **P., Infantile Spasmodic**, spastic cerebral paraplegia, spasmodic tabes dorsalis, birth-palsy; a spastic paralysis coming on in early childhood, and usually dependent on a cerebral lesion with failure of proper de-

velopment or secondary sclerosis of the motor tracts of the spinal cord. The causes are injuries during birth, intrauterine cerebral inflammation, or anomalies of brain-development. **P.**, Spastic, lateral sclerosis.

Paraplegic (*par-ah-ple'-jik*) [*παρά*, beside; *πληγή*, a stroke]. Pertaining to, or affected with, paraplegia.

Paraproctitis (*par-ah-prok-ti'-tis*) [*παρά*, beside; *πρωκτός*, anus; *ιτις*, inflammation]. Inflammation of the connective tissue about the rectum.

Parareducin (*par-ah-re-du'-sin*) [*παρά*, beside; *reducere*, to lead back]. A leukomatin found in conjunction with reducin in the urine.

Pararhotacism (*par-ah-ro'-tas-ism*). See *Rhotacism*.

Pararthria (*par-ar'-thre-ah*) [*παρά*, beside; *ἄρθρον*, articulation]. A disorder of articulate speech.

Parasigmatism (*par-ah-sig'-mat-ism*) [*παρά*, beside; *σίγμα*, the letter *ς*]. The inability to pronounce *s* or *sh*, another letter, as *f*, being substituted.

Parasinoidal (*par-ah-sin-oi'-dal*) [*παρά*, beside; *sinus*, sinus]. Lying near or along a cerebral sinus. **P.** Spaces, the expansion of the cerebral veins just before emptying into the superior longitudinal sinus.

Parasite (*par'-ah-sit*) [*παρά*, near; *σῖτος*, food]. 1. An animal or vegetable living upon or within another organism, termed the host. The most important classes of animal parasites of man are the following: (A) Protozoa, unicellular organisms, as the Plasmodium malarie; the Amoeba coli. (B) Metazoa, multicellular organisms, as (a) The Vermes, or worms, including the Trematodes, the most important of which are the Distomas; the Cestodes, Tæniæ or tape-worms, including Tænia solium, Tænia mediocanellata or saginata, Tænia echinococcus, and Bothriocephalus latus; the Nematodes, comprising Ascaris, Oxyuris, Anchylostomum, Filaria, Trichina, and Trichocephalus; (b) the Arthropoda, including the itch-mite, louse, flea, etc. The vegetable parasites are the bacteria and fungi, *q. v.* 2. In teratology, a fetus or fetal parts attached to or included in another fetus.

Parasitic (*par-ah-sit'-ik*) [*παρά*, beside; *σῖτος*, food]. 1. Of the nature of a parasite; living upon or in an animal or vegetable; as *P. bacteria*, *P. worms*. 2. Caused by parasites; as *P. skin-diseases*.

Parasiticide (*par-ah-sit'-is-id*) [*παρά*, beside; *σῖτος*, food; *cadere*, to kill]. 1. Destructive to parasites. 2. An agent capable of destroying parasites; especially one destroying the parasites living upon or in the skin.

Parasitism (*par'-ah-si-tism*) [*παρά*, beside;

σῖτος, food]. The relation that a parasite bears to its host; infestation by parasites.

Parasitogenetic (*par-ah-si-to-gen-et'-ik*) [*παρά*, beside; *σῖτος*, food; *γεννᾶν*, to beget]. Produced by parasites; depending for its origin upon parasites.

Parasitology (*par-ah-si-tol'-o-je*) [*παρά*, beside; *σῖτος*, food; *λόγος*, science]. The study of parasites.

Paraspadia (*par-ah-spa'-de-ah*) [*παρά*, beside; *σπάειν*, to draw]. A condition in which the urethra opens on one side of the penis.

Parasternal (*par-ah-ster'-nal*) [*παρά*, beside; *στέρνον*, sternum]. Beside or near the sternum. **P.** Line, the imaginary vertical line midway between the margin of the sternum and the line passing through the nipple. **P.** Region, the region between the sternal margin and the parasternal line.

Parathyroid (*par-ah-thi'-roid*) [*παρά*, beside; *θυρεός*, a shield; *εἶδος*, like]. 1. Lying beside the thyroid gland. 2. An accessory thyroid gland.

Paratoloid (*par-ah'-to-loid*). Synonym of *Tuberculin*.

Paratonia (*par-ah-to'-ne-ah*) [*παρά*, beside; *τόνος*, stretching]. Overextension; excessive tension.

Paratrichosis (*par-ah-trik'-o'-sis*) [*παρά*, beside; *τριξ*, hair]. A condition in which the hair is either imperfect in growth or develops in abnormal places.

Paratrimma (*par-ah-trim'-ah*) [*παρά*, along; *τριβεῖν*, to rub]. Intertrigo.

Paratyphlitis (*par-ah-tif-li'-tis*) [*παρά*, beyond; *τύφλος*, blind; *ιτις*, inflammation]. Inflammation of the connective tissue behind the cecum.

Paraxanthin (*par-ah-san'-thin*) [*παρά*, beyond; *ξανθός*, yellow], $C_8H_8N_2O_8$. Dimethylxanthin, a crystalline leukomatin occurring in normal urine, and isomeric with theobromin, which it resembles in its action upon the organism, producing muscular rigidity, dyspnea, and diminution in reflex excitability.

Paraxial (*par-aks'-e-al*) [*παρά*, beside; *ἄξις*, axis]. Lying near the axis of the body.

Paregoric (*par-e-gor'-ik*) [*παρηγορικός*, consoling]. 1. Soothing, or assuaging. 2. A soothing remedy, as *P. Elixir*, or simply paregoric, the tinctura opii camphorata.

Pareira (*par-i'-rah*). The root of Chondrodendron tomentosum, of the natural order Menispermaceæ. It was formerly called *P. brava*. It contains a resin, an alkaloid, pelosin, identical with berberin, a bitter principle, a nitrogenous substance, calcium malate, potassium nitrate, and other salts. *P.* is used as a diuretic in calculous pyelitis; in cystitis; and has also been employed in

leukorrhea, rheumatism, and dropsy, and in Brazil in the treatment of snake-bite. Decoctum pareira fluidum (B. P.). Dose f5j-ij (32.0-64.0). Extractum pareiræ (B. P.). Dose gr. x-xx (0.65-1.3). Extractum pareiræ fluidum (U. S. P.). Dose f3ss-f3j (2.0-4.0). Extractum pareiræ liquidum (B. P.). Dose f3j (4.0).

Parencephalon (*par-en-sef'-al-on*) [παρά, beside; ἐνέφαλος, brain]. The cerebellum.

Parencephalus (*par-en-sef'-al-us*) [παρά, beside; ἐνέφαλος, brain]. A congenital malformation of the brain. See *Porencephalus*.

Parenchyma (*par-eng'-kim-ah*) [παρά, beside; ἔχειν, to pour in]. The essential or specialized part of an organ, as distinguished from the supporting connective tissue.

Parenchymatous (*par-eng'-kim'-at-us*, or *par-eng'-kiv'-mat-us*) [παρά, beside; ἔχειν, to pour in]. Pertaining to, or affecting the parenchyma. **P. Degeneration**, cloudy swelling. **P. Inflammation**, inflammation of the parenchyma, as distinguished from that of the interstitial tissue.

Parepididymis (*par-epi-did'-im-is*). See *Paradidymis*.

Paresis (*par'-es-is*) [παρά, from; ἵσται, to let go]. A slight paralysis; incomplete loss of muscular power. **P., General**. See *Paralysis, General*.

Paresthesia (*par-es-the'-se-ah*) [παρά, beside; αἰσθῆσις, sensation]. 1. Morbid or perverted sensation, as numbness, formication, "pins and needles." 2. See *Acroparesthesia*.

Paresthetic (*par-es-the'-ik*) [παρά, beside; αἰσθῆσις, sensation]. Pertaining to, affected with, or characterized by paresthesia.

Paretic (*par-et'-ik*) [παρά, beside; ἵσται, to let go]. Pertaining to, or affected with paresis. **P. Dement**, a person suffering from paretic dementia. **P. Dementia**. See *Paralysis, General*.

Pareunia (*par-u'-ne-ah*) [παρά, beside; εὐνή, bed]. Coitus.

Paridrosis (*par-id-ro'-sis*) [παρά, beside; ἰδρώς, sweat]. Any abnormal condition of the secretion of sweat.

Parietal (*par-i'-et-al*) [*paries*, wall]. 1. Forming or situated on a wall, as the P. layer of the peritoneum. 2. Pertaining to, or in relation with the parietal bone of the skull, as the P. foramen, P. lobe of the brain. **P. Cells**, cells found in the periphery of the peptic glands of the stomach, immediately beneath the basement-membrane. Their function is supposed to be the secretion of hydrochloric acid.

Parietooccipital (*par-i-et-o-ok-sip'-it-al*) [*paries*, a wall; *occiput*, the back of the head]. Pertaining to the parietal and occipital bones or lobes.

Parietosphenoid (*par-i-et-o-sf'-enoid*) [*paries*, wall; *οἶψ*, wedge; *εἶδος*, like]. Pertaining to the parietal and sphenoid bones.

Parietosquamosal (*par-i-et-o-skwa-mo'-sal*) [*paries*, a wall; *squama*, a scale]. Of or pertaining to the parietal bone and the squamous portion of the temporal bone. **P. Suture**, a suture between the squamous portion of the temporal bone and the parietal bone.

Parietotemporal (*par-i-et-o-tem'-po-ral*) [*paries*, a wall; *tempora*, the temples]. Pertaining to the parietal and temporal bones. **P. Suture**, the suture between the parietal and temporal bones.

Paris Green, $\text{Cu}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 3\text{Cu}(\text{AsO}_2)_2$. Acetoarsenite of copper, a poisonous substance used in the arts and for the destruction of the potato-bug.

Parkinson's Disease. See *Paralysis agitans*.

Paronychia (*par-o-nik'-e-ah*) [παρά, beside; ὀνύξ, nail]. An inflammation of the flexor tendons and tendinous sheaths of the fingers; whitlow.

Paroophoron (*par-o-off'-o-ron*) [παρά, beside; ὄφρ, an egg; φέρειν, to bear]. The persistent tubules of the posterior part of the Wolffian body in the female, corresponding to the organ of Girdaldes in the male.

Parosmia (*par-os-me-ah*) [παρά, aside; ὀσμή, smell]. A perversion of the sense of smell.

Parostitis (*par-os-ti'-tis*) [παρά, near; ὀστέον, bone; ἰτις, inflammation]. Inflammation of the outer surface of periosteum.

Parostosis (*par-os-to'-sis*) [παρά, beside; ὀστέον, bone]. The abnormal formation of bone outside of the periosteum, or in the connective tissue surrounding the periosteum.

Parotid (*par-ot'-id*) [παρά, near; οὖς, ear].

1. Situated near the ear, as the P. Gland. 2. Pertaining to or affecting the parotid gland. **P. Abscess**, an abscess of the parotid gland; The term is sometimes also applied to abscess of the lymphatic gland lying upon the parotid. **P. Gland**, one of the salivary glands situated in front of and below the external ear. It is a compound racemose gland and secretes saliva containing ptyalin, a globulin-like body, potassium sulphocyanid, a trace of urea, and mineral salts. Its duct is known as Stensen's duct.

Parotiditis (*par-ot-id-i'-tis*). See *Parotitis*.

Parotitis (*par-o-ti'-tis*) [παρά, beside; οὖς, ear; ἰτις, inflammation]. Inflammation of the parotid gland, especially the specific infectious disease known as mumps; the name is also given to inflammation of the lymphatic gland overlying the parotid (Parotid bulbo). **P., Metastatic**, that secondary to disease elsewhere; it occurs in infectious fevers, as

typhoid fever, and usually goes on to suppuration.

Parovarian (*par-o-va'-re-an*) [*παρά*, beside; *ovarium*, ovary]. 1. Situated near the ovary. 2. Pertaining to the parovarium.

Parovarium (*par-o-va'-re-um*) [*παρά*, beside; *ovarium*, ovary]. The remnant of the Wolffian body of the female; the organ of Rosenmüller.

Paroxysm (*par'-oks-izm*) [*παρά*, beside; *ὀξύειν*, to sharpen]. The periodic increase or crisis in the progress of a disease; a sudden attack, a sudden reappearance of symptoms, or a sudden increase in the intensity of existing symptoms.

Paroxysmal (*par'-oks-iz'-mal*) [*παρά*, beside; *ὀξύειν*, to sharpen]. Of the nature of or resembling a paroxysm; occurring in paroxysms.

Parrot-beak Nails. Nails that are curved strongly anteroposteriorly, like the beak of the parrot.

Parrot's Disease. Syphilitic pseudoparalysis. See *Diseases*, Table of.

Parrot's Nodes. See *Signs and Symptoms*, Table of.

Parry's Disease. Exophthalmic goiter.

Pars (*parz*) [L.]. A part. **P. ciliaris** retinae, the part of the retina in front of the ora serrata.

Parsley (*pars'-le*) [*πέτρος*, rock; *σέλινον*, a kind of parsley]. The *Apium petroselinum*, a plant of the order Umbelliferae, containing a volatile oil. From the seed a peculiar oily liquid, termed apiol (*q. v.*), is obtained. The root is used in renal diseases and dropsy; the juice of the fresh herbs and the seeds are employed as antiperiodics; apiol is an emmenagogue.

Parson's Disease. Exophthalmic goiter.

Parthenium (*par-the'-ne-um*) [*παρθένος*, a virgin]. A genus of herbs of the order Compositae. **P. hysterophorus** contains parthenic acid and several alkaloids, one of which, called parthenin, seems to be the active principle of the plant and has been used as an antipyretic and antineuralgic. **P. integrifolium**, prairie dock, a perennial plant of the Southern U. S., is used as an antiperiodic.

Parthenogenesis (*par'-then-o-jen'-es-is*) [*παρθένος*, a virgin; *γένεσις*, a begetting]. The development of an organism from an unfertilized ovum.

Partridge-berry. See *Gaultheria procumbens*.

Parturient (*par-tu'-re-ent*) [*parturire*, to bring forth]. 1. Being in labor; giving birth; as a P. woman. 2. Traversed during birth, as the P. canal.

Parturiometer (*par-tu-re-om'-et-er*) [*parturire*, to bring forth; *μέτρον*, measure]. An instrument for determining the progress of

labor by measuring the expulsive force of the uterus.

Parturition (*par-tu-rish'-un*) [*parturire*, from *parturire*, to bring forth]. The act of giving birth to young. See *Labor*.

Partus (*par'-tus*) [*parere*, to bring forth]. The bringing forth of offspring; labor. **P. cæsaricus**, Cesarean section. **P. difficilis**, dystocia. **P. immaturus**, premature labor. **P. maturus**, labor at term. **P. præcipitatus**, precipitate labor. **P. siccus**, dry labor.

Parulis (*par-u'-lis*) [*παρά*, near; *ούλον*, the gum]. Abscess of the gum; gum-boil.

Paruria (*par-u'-re-ah*) [*παρά*, beside; *urina*, urine]. Abnormality in the excretion of the urine.

Parvolin (*par'-vo-lin*), $C_8H_{13}N$. 1. A synthetic liquid base. 2. A ptomain isomeric with it, occurring in decomposing fish and horseflesh.

Parvule (*par'-vul*) [*parvus*, small]. A small pill, or pellet.

Passage (*pas'-aj*) [*passare*, to pass]. 1. A channel. 2. The act of passing from one place to another. 3. The introduction of an instrument into a cavity or channel. 4. An evacuation of the bowels. **P.**, False, a false channel, especially one made by the unskillful introduction of an instrument into the urethra.

Passion (*pass'-un*) [*passio*, from *pati*, to suffer]. 1. Pain; suffering; as iliac P., a synonym of volvulus. 2. An intense emotion of the mind; intense sexual excitement.

Passive (*pas'-iv*) [*pati*, to suffer]. Not active; not performed or produced by active efforts, but by causes coming from without.

P. Congestion, congestion due to retention of blood in a part, and not to an active flow of blood toward the part. **P. Immunity.** See *Immunity*. **P. Motion**, the movement produced by external agency, and not by the person himself.

Passula (*pas'-u-lah*) [L.]. A raisin.

Paste (*pást*) [*pasta*; *πάστη*, mess; *πάσσειν*, to sprinkle]. Any soft, sticky substance, especially a mixture of starch or flour and water. **P.**, Arsenical, a caustic paste containing arsenic. **P.**, Canquoin's. See *Canquoin's Paste*. **P.**, London-, a mixture of equal parts of sodium hydrate and slaked lime, moistened with alcohol. **P.**, Vienna-, a mixture of potassium hydrate and caustic lime moistened with water.

Pasteur's Fluid or **P.'s Liquid** [Louis Pasteur, a French scientist]. An artificial liquid for the cultivation of bacteria, composed of water 100 parts, crystallized sugar 10 parts, carbonate of ammonium and ashes of yeast, each 1 part.

Pasteurization (*pas-tur-i-za'-shun*) [Louis Pasteur, a French scientist]. The process

of checking fermentation in milk, wine, and other organic fluids by heating them to 60° or 70° C.

Pastil, Pastille (*pas'-til, pas-till'*) [dim. of *pasta*, paste]. 1. A small mass composed of aromatic substances and employed in fumigation. 2. A troche.

Patch [Low Ger. *plakke*, a spot]. An irregular spot or area. **P., Mucous.** One of the characteristic lesions of syphilis, occurring in the so-called secondary stage, and appearing as a whitish papule or patch on mucous membranes and at mucocutaneous junctions. It is also called mucous papule and condyloma latum. **P., Opaline.** See *Opaline*. **Peyer's Patches.** See *Glands, Peyer's*.

Patella (*pat-el'-ah*) [dim. of *patina*, a shallow dish]. The knee-pan, a small, round sesamoid bone in front of the knee, developed in the tendon of the quadriceps extensor femoris muscle.

Patellar (*pat-el'-ar*) [*patina*, a shallow dish]. Pertaining to the patella. **P. Reflex,** or **P. Tendon-reflex.** See *Knee-jerk* and *Reflexes, Table of*.

Patency (*pa'-ten-se*) [*patere*, to stand open]. The state of being open; openness.

Pathetic (*path-el'-ik*) [*παθητικός; πάθος*, disease]. Arousing pity; indicating sadness or sorrow; appealing; that which appeals to or stirs the passions; applied to the fourth cranial nerve (P. nerve), which innervates the P. muscle (Patheticus, superior oblique) of the eye, by which the eye is rolled outward and downward.

Pathetism (*path'-et-izm*) [*πάθος*, disease]. Hypnotism.

Pathoanatomy (*path-o-an-at'-o-me*) [*πάθος*, disease; *ἀνατομή*, a cutting up]. Pathologic anatomy.

Pathogenesis (*path-o-jen'-es-is*) [*πάθος*, disease; *γεννᾶν*, to beget]. The origin or development of disease.

Pathogenic, Pathogenetic (*path-o-jen'-ik, path-o-jen-el'-ik*) [*πάθος*, disease; *γεννᾶν*, to beget]. Producing disease. **P. Microorganism,** one that when introduced into the system causes disease.

Pathognomonic (*path-og-no-mon'-ik*) [*πάθος*, disease; *γνώμων*, judge]. Characteristic of a disease, distinguishing it from other diseases.

Pathologic (*path-o-loj'-ik*) [*πάθος*, disease; *λόγος*, science]. Pertaining to pathology; pertaining to disease. **P. Anatomy.** See *Anatomy*. **P. Histology,** the microscopic study of diseased tissues.

Pathologist (*path-ol'-o-jist*) [*πάθος*, disease; *λόγος*, science]. One versed in pathology.

Pathology (*path-ol'-o-je*) [*πάθος*, disease; *λόγος*, science]. The branch of medical

science that treats of the modifications of function and changes in structure caused by disease. **P., Cellular,** pathology that makes the cell the basis of all vital phenomena, and teaches that every cell is derived from a preexisting parent-cell. **P., Comparative,** a study of pathologic processes in lower animals, for purposes of tracing resemblances and differences among them and between them and those of the human body. **P., Experimental,** the study of pathologic processes artificially induced in lower animals. **P., General,** that department of pathology which takes cognizance of those morbid processes that may be observed in various diseases and in any organ, *e. g.*, inflammation, hypertrophy. **P., Geographic,** pathology in its relation to climatic and geographic conditions. **P., Humoral,** the old doctrine that disease is due to abnormal conditions of the blood. It has been revived in recent times in a modified form, and is now based on the theory that both immunity and susceptibility to disease reside in the juices of the body. **P., Special,** that treating of changes in function and structure occurring in special organs, *e. g.*, pneumonia. **P., Surgical,** the pathology of diseases treated by the surgeon.

Pathophobia (*path-o-fō'-be-ah*) [*πάθος*, disease; *φόβος*, fear]. Exaggerated dread of disease.

Patient (*pa'-shent*) [*pati*, to suffer]. A person under the care of a physician; a sick person.

Paullinia (*pa-u-lin'-e-ah*). See *Guarana*.

Paulocardia (*pa-u-lo-kar'-de-ah*) [*παύλα*, pause; *καρδία*, heart]. A subjective sensation of intermission or momentary stoppage of the heart-beat.

Pavement-epithelium. Epithelium consisting of flattened scale-like cells fitted together by their edges like the tiles of a pavement.

Pavilion (*pa-vill'-yon*) [*papilio*, a butterfly; a tent]. The expanded extremity of a canal or tube, as the P. of the ear—the auricle; the P. of the Fallopian tube—the fimbriated extremity of the Fallopian tube.

Pavor (*pa'-vor*) [L.]. Fright; fear. **P. nocturnus,** night-terrors.

Pavy's Disease. Intermitting or recurrent albuminuria.

Pawpaw. See *Papaya*.

Pearl (*perl*) [Fr., *perle*, from L., *pirula*, a little pearl]. In pharmacy, a small hollow glass body containing a dose of a volatile liquid medicine, as a P. of amyl nitrite. **P.-disease,** tuberculosis of serous membranes in the lower animals, especially cattle, so-called on account of the most manifest lesion, the pearly nodules or tumors,

which are often pendulous. **P.**, Epidermic, **P.**, Epithelial, one of the spheroid concentric masses of epithelial cells often seen in hard papillomata, in squamous epitheliomata, and in cholesteatomata; they are also called pearly bodies.

Pearly Body. See *Pearl*, *Epithelial*.

Pebrine (*pek'-rin*) [*Fr.*]. An infectious epidemic disease of silkworms.

Pecquet, Cistern of, or Reservoir of. The receptaculum chyli.

Pecten (*pek'-ten*) [*pecten*, a comb]. The os pubis.

Pectin (*pek'-tin*) [*πηκτός*, congealed]. A white amorphous carbohydrate contained in ripe fleshy fruits and in certain roots, and believed to be formed from the pectose found in unripe fruits by the action of acids. It is also called vegetable jelly.

Pectinate (*pek'-tin-at*) [*pecten*, a comb]. Arranged like the teeth of a comb. **P.** Ligament, fibers of connective tissue at the angle of the anterior chamber of the eye, between the iris and the cornea. **P.** Muscles, the muscoli pectinati, muscular ridges in the auricles of the heart.

Pectineal (*pek'-tin-e'-al*) [*pecten*, a comb]. 1. Comb-shaped. 2. Pertaining to the pecten or os pubis. **P.** Line, that part of the ileopectineal line found on the os pubis.

Pectineus (*pek'-tin-e'-us*). See *Muscles*, *Table of*.

Pectiniform (*pek'-tin'-if-form*) [*pecten*, a comb; *forma*, a form]. Comb-shaped.

Pectoral (*pek'-to-ral*) [*pectus*, breast]. 1. Pertaining to the chest, as the **P.** muscles. 2. Useful in diseases of the chest. 3. A remedy useful in diseases of the chest.

Pectoralis (*pek'-tor-a'-lis*). See *Muscles*, *Table of*.

Pectoriloquy (*pek'-tor-il'-o-kwe*) [*pectus*, breast; *loqui*, to speak]. The distinct transmission of articulate speech to the ear on auscultation. It may be heard over cavities in the lung, over areas of consolidation near a large bronchus, over a pneumothorax when the opening in the lung is patulous, and over some pleural effusions. **P.**, Whispering, the transmission of the whispered words to the auscultating ear. The sounds seem to emanate directly from the spot auscultated.

Pectose (*pek'-tos*) [*πηκτός*, congealed]. A compound occurring in unripe fruits, and giving rise to pectin.

Pectus (*pek'-tus*) [*pectus*, breast]. The chest, or breast. **P.** carinatum, keeled breast, pigeon-breast; a narrow chest projecting anteriorly in the region of the sternum.

Pedal (*pel'-dal*) [*pes*, foot]. Pertaining to the foot.

Pedatrophia, Pedatrophly (*ped-at-ro'-fe-ah*,

ped-at'-ro-fe) [*παῖς*, child; *atrophy*]. 1. Any wasting disease of childhood. 2. *Tabes mesenterica*.

Pederasty (*ped'-er-as-te*) [*παῖς*, boy; *ἐράειν*, to love]. Sexual intercourse through the anus.

Pediatrics, Pediatrics (*pe-de-at'-riks*, *pe'-de-at-re*) [*παῖς*, child; *ιατρία*, therapeutics]. The branch of medicine dealing with the diseases of children.

Pedicle (*ped'-ikl*) [*pediculus*, dim. of *pes*, foot]. 1. A slender process acting as a foot or stem, as the **P.** of a tumor. 2. Of a vertebra, the portion of bone projecting backward from each side of the body and connecting the lamina with the body.

Pediculation (*pe-dik-u-lu'-shun*) [*pediculus*, louse]. The state of one suffering from pediculosis.

Pediculophobia (*pe-dik-u-lo-fo'-be-ah*) [*pediculus*, louse; *φόβος*, fear]. Morbid dread of lice.

Pediculosis (*pe-dik-u-lo'-sis*) [*pediculus*, louse]. Lousiness; a skin-affection characterized by the presence of pediculi or lice.

Pediculus (*pe-dik'-u-lus*) [*pediculus*, a louse]. A small parasitic hemipterous insect, the louse. **P.** capitis, the head-louse. **P.** corporis, the body-louse. **P.** vestimenti. Synonym of *P. corporis*.

Pediluvium (*ped-il'-u'-ve-um*) [*pes*, foot; *lavare*, to wash]. A foot-bath.

Peduncle (*pe'-dung-kl*) [*pedunculus*, dim. of *pes*, foot]. A narrow part acting as a support. **P.**, Cerebral, the crus cerebri. **P.**, Inferior Cerebellar, one of two bands of white matter passing up from the medulla oblongata, connecting the medulla with the cerebellum, and forming the lower lateral wall of the fourth ventricle. **P.**, Middle Cerebellar, one of the bands of white matter joining the pons and the cerebellum. **P.** of the Pineal Gland, a delicate white band passing forward from each side of the pineal gland along the edge of the third ventricle. **P.**, Posterior Cerebellar. Synonym of *P.*, Inferior Cerebellar. **P.**, Superior Cerebellar, one of the two bands of white matter that pass from the cerebellum to the testes of the corpora quadrigemina.

Peduncular (*ped-ung'-ku-lar*) [*pedunculus*, dim. of *pes*, foot]. Pertaining to a peduncle.

Pedunculate, Pedunculated (*ped-ung'-ku-lat*, *ped-ung'-ku-la-ted*) [*pedunculus*, dim. of *pes*, foot]. Having a peduncle; stalked.

Pelada (*pel'-a-dah*) [*Fr.*]. Alopecia of the scalp.

Pelargonic Acid (*pel-ar-gon'-ik*) [*πελαργός*, a stork], $C_{15}H_{31}O_2$. A monobasic crystalline acid obtained from the essential oil of Pelargonium roseum and from other oils. It is employed in the flavoring of wines.

Peliosis (*pel-e-o'-sis*) [*πέλιος*, livid]. Purpura. *P. rheumatica*, purpura rheumatica; a disease characterized by a purpuric rash, with arthritis and fever.

Pellagra (*pel-a'-grah*) [*πέλλα*, skin; *ἀγρα*, seizure]. A disease occurring in Italy, Southern France, and Spain, and attributed to the use of diseased maize. It is characterized in the early stages by debility, spinal pains, and digestive disturbances; later erythema develops, with drying and exfoliation of the skin. In severe cases various nervous manifestations arise, such as spasms, ataxic paraplegia, and mental disturbances. In cases presenting ataxic paraplegia the spinal cord has shown combined posterior and lateral sclerosis. *P.* is also known as Lombardian leprosy.

Pellet (*pel'-et*) [*pila*, ball]. A small pill.

Pelletierin (*pel-et'-er-in*), $C_8H_{13}NO$. A liquid alkaloid obtained together with an isomeric body, isopelletierin, also a liquid alkaloid, from pomegranate-bark. It is used as a teniafuge, chiefly in the form of the tannate. Dose $\frac{1}{2}$ to 2 gr. (0.032-0.13).

Pellicle (*pel'-ikl*) [*pellis*, skin]. A thin membrane, film, scum, or cuticle.

Pellitory (*pel'-it-or-e*). See *Pyrethrum*.

Pelvic (*pel'-vik*) [*pelvis*, a basin]. Pertaining to the pelvis. *P. Abscess*, a suppurative inflammation of the connective tissue of the pelvic cavity, most common in women, and usually associated with puerperal or gonorrheal infection. *P. Arch*. Same as *P. Girdle*. *P. Cellulitis*, inflammation of the connective tissue of the pelvis. *P. Girdle*, the arch formed by the ilium, ischium, and pubis, or in the higher vertebrates by the two innominate bones. *P. Index*, the relation of the anteroposterior to the transverse diameter of the pelvis. *P. Inlet*, the superior strait. *P. Outlet*, the inferior strait. *P. Region*, the region within the true pelvis.

Pelvimeter (*pel-vim'-et-er*) [*pelvis*, pelvis; *μέτρον*, measure]. An instrument for measuring the pelvic dimensions.

Pelvimetry (*pel-vim'-et-re*) [*pelvis*, pelvis; *μέτρον*, measure]. The measurement of the dimensions of the pelvis.

TABLE OF MEASUREMENTS OF THE FEMALE PELVIS COVERED BY THE SOFT PARTS.

Between iliac spines,	26 cm.
Between iliac crests,	29 "
External conjugate diameter,	20½ "
Internal conjugate diagonal,	12¾ "
True conjugate, estimated,	11 "
Right diagonal,	22 "
Left diagonal,	22 "
Between trochanters,	31 "
Circumference of pelvis,	90 "

Pelvipерitonitis (*pel-ve-per-e-ton-i'-tis*) [*pel-vis*; *peritonitis*]. Pelvic peritonitis.

Pelvis (*pel'-vis*) [*pelvis*, a basin]. 1. A

basin or basin-shaped cavity, as the *P.* of the kidney. 2. The bony ring formed by the two innominate bones and the sacrum and coccyx.

3. The cavity bounded by the bony pelvis.

The pelvis consists of two parts, the true *P.* and the false *P.*, which are separated by the ileopectineal line. The entrance of the true pelvis, corresponding to this line, is known as the inlet or superior strait; the outlet or inferior strait, is bounded by the symphysis pubis, the tip of the coccyx, and the two ischia. In measuring the pelvis the *cardinal points of Capuron* are used as landmarks. They are the two ileopectineal eminences and the two sacroiliac joints. *P. æquabiliter justo major*, one equally enlarged in all diameters. *P. æquabiliter justo minor*, a pelvis with all its diameters reduced below the normal. *P.*, *Axis of (of inlet or outlet)*, a perpendicular to the middle of the anteroposterior diameter. *P.*, *Beaked*, one in which the pubic bones are compressed laterally so as to approach each other, and are pushed forward; a condition seen in osteomalacia. *P.*, *Brim of*, the entrance to the pelvic cavity, called the inlet, superior strait, margin, or isthmus. *P.*, *Diameters of*, imaginary lines drawn between certain bony points. (*a*) Of the inlet: the anteroposterior (sacropubic, or conjugate), from the upper edge of the promontory of the sacrum to a point an eighth of an inch below the upper border of the pubic symphysis; it measures 11 cm.; the transverse, from side to side at the widest point, measuring 13½ cm.; the oblique (right and left), measuring 12¾ cm. (*b*) Of the outlet; the anteroposterior, from the tip of the coccyx to the subpubic ligament, measuring 9½ cm.; the transverse, between the ischial tuberosities, measuring 11 cm.; the oblique, from the under surface of the sciatic ligaments to the junction of the ischiopubic rami. *P.*, *Floor of*, the mass of skin, connective tissue, muscles, and fascia forming the inferior boundary of the pelvis. *P.*, *Justo major*. See *P. æquabiliter justo major*. *P.*, *Justo minor*. See *P. æquabiliter justo minor*. *P.*, *Kyphotic*, one characterized by increase of the conjugate diameter of the inlet, but decrease of the transverse diameter of the outlet, through approximation of the tuberosities of the ischium. *P.*, *Nægele's Oblique*, a pelvic deformity with ankylosis of one sacroiliac synchondrosis, lack of development of the associated lateral sacral mass, and other defects that distort the diameters and render the conjugate oblique in direction. *P.*, *Obliquity of*, *P.*, *Inclination of*, the angle between the axis of the pelvis and that of the body. *P.*, *Osteomalacic*, a distorted pelvis characterized by a lessening of the transverse

and oblique diameters, with great increase of the anteroposterior diameter. **P., Planes** of, imaginary surfaces touching all points of the circumference. The plane of pelvic expansion perforates the middle of the symphysis, the tops of the acetabula, and the sacrum between the second and third vertebrae. Its anteroposterior diameter is $12\frac{3}{4}$ cm., its transverse diameter is $12\frac{1}{2}$ cm. The plane of pelvic contraction passes through the tip of the sacrum, the spines of the ischia and the under surface of the symphysis. Its anteroposterior diameter is $11\frac{1}{2}$ cm.; its transverse diameter is $10\frac{1}{2}$ cm. **P., Rachitic**, one characterized by a sinking in and forward of the sacrovertebral angle, with a flaring outward of the iliac crests and increased separation of the iliac spines. **P., Robert's**, one in which there is an ankylosis of both sacroiliac joints, with a rudimentary sacrum, both lateral sacral masses being undeveloped, the oblique and transverse diameters being much narrowed. **P., Simple Flat**, one in which the only deformity consists in a shortening of the anteroposterior diameter. **P. spinosa**, a rachitic pelvis in which the crest of the pubis is very sharp, and presents a spine at the insertion of the psoas parvus. **P., Split**, a form in which there is congenital separation of the pubic bones at the symphysis. It is often associated with exstrophy of the bladder.

Pemphigoid (*pen'-fig-oid*) [*πέμφιξ*, blister; *ζῖδος*, like]. Resembling or having the nature of pemphigus.

Pemphigus (*pen'-fig-us*) [*πέμφιξ*, a blister]. An acute or chronic disease of the skin characterized by the appearance of bullæ or blebs. Two principal varieties are described. **P. vulgaris**, and **P. foliaceus**. **P. vulgaris** is usually chronic, the blebs appearing in successive crops; on healing they leave a pigmented spot. Itching and pain may be present. **P. foliaceus**, is a rare form characterized by crops of flaccid blebs containing a turbid fluid. The disease is usually of long duration but eventually ends fatally. The cause of **P.** is not definitely known. Other forms of **P.** are: **P. neonatorum**, an acute form of **P.** occurring in infants and supposed to be due to a microorganism. **P. syphiliticus**, a bullous eruption due to syphilis. **P. vegetans**, Neumann's disease, an affection characterized by sore mouth, followed by some form of dermatitis attended by vesication and followed by papillary growths, gradual emaciation, and death.

Pendjeh Sore. See *Furunculus orientalis*.

Pendulous (*pen'-du-lus*) [*pendere*, to hang]. Hanging down loosely.

Penetrating (*pen'-e-tra-ting*) [*penetrare*, to pierce]. Entering beyond the surface. **P.**

Wound, one that pierces the wall of a cavity or enters into an organ.

Penetration (*pen'-e-tra'-shun*) [*penetrare*, to pierce]. 1. The act of penetrating or piercing into. 2. Of a microscope, the focal depth. 3. The entrance of the penis into the vagina.

Penicillium (*pen-is-il'-e-um*) [*penicillus*, a brush]. A genus of fungi, of which the *P. glaucum*, or common blue mold, is a familiar example.

Penile (*pe'-nil*) [*penis*]. Pertaining to the penis.

Penis (*pe'-nis*) [L.]. The male organ of copulation. It consists of the corpus spongiosum, enclosing the urethra, the two corpora cavernosa, largely composed of erectile tissue, and the glans.

Penitis (*pe-ni'-tis*) [*penis*, penis; *itis*, inflammation]. Inflammation of the penis.

Penniform (*pen'-if-orm*) [*penna*, feather; *forma*, form]. Shaped like a feather; said of certain muscles.

Pennyroyal (*pen'-e-roi'-al*). See *Hedeoma*.

Pennyweight (*pen'-e-wât*) [AS., *pening*, penny; *wegan*, weigh]. A weight of 24 grains.

Pentad (*pen'-lad*) [*πέντε*, five]. An element or radicle having a valence of five.

Pentamethylendiamin (*pen-tah-meth-il-en-di-am'-in*). See *Cadaverin*.

Pentane (*pen'-tân*) [*πέντε*, five], C_5H_{12} . The fifth member of the paraffin-series of hydrocarbons. It is a liquid and occurs in naphtha.

Pentavalent (*pen-tav'-al-ent*) [*πέντε*, five; *valens*, having power]. Quinivalent.

Pentene (*pen'-ten*) [*πέντε*, five], C_5H_{10} . Amylene, one of the olefin-series of hydrocarbons.

Pentose (*pen'-tôs*) [*πέντε*, five]. Any one of a class of carbohydrates containing five atoms of carbon. The pentoses are not fermentable, and on boiling with dilute hydrochloric acid yield furfural, $C_5H_4O_2$.

Pentosuria (*pen-to-su'-re-ah*) [*pentose*; *urina*, urine]. The presence of pentose in the urine. Urine containing pentose reduces Fehling's solution, but does not ferment.

Pepper [*piper*, from Skt. *pippala*]. The fruit of various species of *Piper*, of the order Piperaceæ. **P., Black**, is the *Piper* of U. S. P. (*Piper nigrum* of B. P.). It contains a neutral principle piperin (Piperinum, U. S. P.), an acrid resin, and a volatile oil, and is used as a condiment and as a carminative stimulant, and to a slight extent is antiperiodic. Dose gr. v-xx (0.32-1.3). **Oleo-resina piperis**. Dose $\mathfrak{m} \frac{1}{4}$ -j (0.016-0.065). **Piperinum**. Dose gr. j-vj (0.065-0.4). **P., White**, resembles black pepper but is less active.

Peppermint. See *Mentha*.

Pepsin (πέψις, digestion). A ferment found in the gastric juice, and capable of digesting proteins in the presence of an acid. It splits albumin into antialbumose and hemialbumose, the former of which it separates into two molecules of antipeptone, while the latter is acted upon by trypsin and split into two molecules of hemipeptone. Pepsin is used in medicine to aid digestion. **Pepsinum** (U. S. P., B. P.) is obtained from the stomach of pigs. Dose gr. x-xv (0.65-1.0). **Pepsinum saccharatum** (U. S. P.), is pepsin mixed with sugar of milk. Dose gr. x-xxx (0.65-2.0). **Vinum pepsini** (Unof.), contains 0.3 per cent. of hydrochloric acid. Dose f 3 ss-j (16.0-32.0).

Pepsinogen (πέψις-ο-γεν) [πέψις, digestion; γεννᾶν, to beget]. The antecedent substance or symogen of pepsin, present in the cells of the gastric glands, and which during digestion is converted into pepsin.

Peptic (πέψις-τικός) [πέψις, digestion]. 1. Pertaining to pepsin. 2. Pertaining to digestion. **P.** Glands, the glands situated in the cardiac and middle thirds of the stomach, and secreting pepsin and hydrochloric acid. **P.** Ulcer, the round ulcer of the stomach, due to erosion of the mucous membrane by the gastric juice.

Peptogenic (πέψις-το-γεν-τικός) [πέπτειν, to cook; γεννᾶν, to produce]. Producing pepsin or peptones.

Peptone (πέψις-τὸν) [πέπτειν, to cook]. A proteid-body formed by the action of ferments on albumins during gastric and pancreatic digestion. It may be considered a hydrated albumin. Before the final formation of peptone, several similar intermediate compounds are produced, as hemipeptone and antipeptone. (See *Pepsin*.) Amphopeptone is a mixture of these two. Propeptone or hemialbumose is a mixture of several intermediate products. Parapeptone is also an intermediate product of digestion and is closely allied to syntonin. Gelatin-peptone is a peptone formed in the digestion of gelatin. Peptones are soluble, readily diffusible, are not precipitated by boiling, by nitric acid, or by potassium ferrocyanid; they are precipitated by mercuric chlorid, by tannic acid, and by phosphomolybdic acid; they give Millon's test, and the xanthoproteic and biuret reactions; they are levorotatory.

Peptonemia (πέψις-το-νέ-μια) [πέπτειν, to cook; αἷμα, blood]. The presence of peptone in the blood.

Peptonization (πέψις-το-νι-σά-σιον) [πέπτειν, to cook]. The process of converting proteids into peptones.

Peptonize (πέψις-το-νίζω) [πέπτειν, to digest]. To digest with pepsin; to predigest.

Peptonuria (πέψις-το-νυ-ρί-α) [πέπτειν, to cook; οὐρίνα, urine]. The presence of peptones in the urine.

Peptotoxin (πέψις-το-τοξί-ν) [πέπτειν, to cook; τοξικόν, poison]. A poisonous ptomain found in peptones, and in putrefying albuminous substances, such as fibrin, casein, brain, liver, and muscle. **P.**, Cholera-, a toxic substance generated by the cholera-bacillus, and chemically allied to peptone.

Peracute (περ-ακ-ού) [περ, through; acutus, sharp]. Very acute.

Perception (περ-σέψις-σιον) [περ, through; capere, to receive]. 1. The act of receiving impressions through the medium of the senses. 2. The faculty receiving such impressions.

Perchloric Acid (περ-κλό-ρικός) [περ, through; χλωρός, green]. HClO₄. The highest oxyacid of chlorine. It is a volatile liquid decomposing in contact with organic substances, and forming salts called Perchlorates.

Percolate (περ-κο-λάω) [percolare, to strain through]. 1. To submit to the process of percolation. 2. The solution obtained by percolation.

Percolation (περ-κο-λά-σιον) [percolare, to strain through]. The process of extracting the soluble constituents of a substance by allowing the solvent to trickle through a powdered mass placed in a long conic vessel, the percolator.

Percussion (περ-κυσίς-σιον) [percutere, to strike through]. A method of physical diagnosis applied by striking upon any part of the body, with a view of ascertaining the conditions of the underlying organs by the character of the sounds elicited. **P.**, Auscultatory, percussion combined with auscultation. It is best performed by placing a double stethoscope at a fixed point and percussing gently all around. **P.**, Immediate, percussion in which the surface is struck directly, without the interposition of a pleximeter. **P.**, Instrumental, the use of a special hammer as a plexor, either alone or with a plate as a pleximeter. **P.**, Mediate, the percussion in which a pleximeter is used. **P.-wave**, the term given to the chief ascending wave of the sphygmographic tracing.

Percutaneous (περ-κυ-τά-νε-ος) [περ, through; cutis, the skin]. Performed through the skin, as **P.** faradization.

Pereirin (περ-ι-ρίν). An amorphous alkaloid found in the bark of Geissospermum laeve. Its hydrochlorate and valerianate are used as antipyretics.

Perflation (περ-φλά-σιον) [perflare, to blow through]. 1. A method of ventilation by which a current of air blowing against a swelling is made to force its way in. 2. The act of forcing air into a cavity for the purpose of evacuating fluid.

Perforans (*per'-for-ans*) [*per*, through; *forare*, to bore]. Penetrating or perforating, a term applied to a muscle or nerve perforating a part.

Perforated (*per'-for-a-ted*) [*per*, through; *forare*, to bore]. Pierced through. **P.** Space, a part of the base of the brain pierced with many small holes for the passage of blood-vessels. The anterior **P.** space is situated on the inner side of the fissure of Sylvius, the posterior **P.** space between the corpora albicantia in front, the pons behind, and the crura cerebri on either side.

Perforation (*per-for-a'-shun*) [*per*, through; *forare*, to bore]. 1. The act of piercing or boring into a part; especially the piercing of the fetal head during labor. 2. A hole made through a part or the wall of a cavity.

Perforating (*per'-for-a-ting*) [*per*, through; *forare*, to bore]. Piercing. **P.** Ulcer. See *Ulcer*.

Perforator (*per'-for-a-tor*) [*per*, through; *forare*, to bore]. An instrument for boring through the fetal skull; also for perforating other bones.

Perforatus (*per-for-a'-tus*) [*per*, through; *forare*, to bore]. Perforated, applied to a muscle which is perforated by another muscle or by a nerve.

Perfusion (*per-fu'-shun*) [*per*, through; *fundere*, to pour]. A pouring of fluid into or through.

Peri- (*per'-e-*) [*περί*, around]. A prefix signifying around, or surrounding.

Perianal (*per-e-a'-nal*) [*περί*, around; *anus*, anus]. Situated or occurring around the anus.

Periarteritis (*per-e-ar-ter-i'-tis*) [*περί*, around; *ἀρτηρία*, artery; *itis*, inflammation]. Inflammation of the external sheath of an artery. **P. nodosa**, a thickening of the arterial coats in localized areas, producing nodular swellings.

Periarthritis (*per-e-ar-thri'-tis*) [*περί*, around; *ἄρθρον*, joint; *itis*, inflammation]. Inflammation of the tissues about a joint.

Periarticular (*per-e-ar-tik'-u-lar*) [*περί*, around; *articulus*, joint]. About a joint.

Periaxial (*per-e-aks'-e-al*) [*περί*, around; *axis*]. Surrounding an axis. **P. Neuritis**, inflammation of the myelin-sheath but not of the axis-cylinder of a nerve.

Periblast (*per'-e-blast*) [*περί*, around; *βλαστός*, a germ]. The protoplasm surrounding the nucleus of a cell.

Peribronchial (*per-e-brong'-ke-al*) [*περί*, around; *βρόγχος*, bronchus]. Surrounding a bronchus; occurring about a bronchus.

Peribronchitis (*per-e-brong-ki'-tis*) [*περί*, around; *βρόγχος*, bronchus; *itis*, inflammation]. Inflammation of the tissue immediately surrounding the bronchi.

Pericardiac, Pericardial (*per-e-kar'-de-ak, per-e-kar'-de-al*) [*περί*, around; *καρδία*, heart]. Pertaining to the pericardium.

Pericarditis (*per-e-kar-di'-tis*) [*περί*, around; *καρδία*, heart; *itis*, inflammation]. Inflammation of the pericardium. The symptoms are slight fever, precordial pain, and tenderness, cough, dyspnea, and rapid pulse. The physical signs vary—in the early stage there is a distinct friction-sound on auscultation, and sometimes a fremitus on palpation. In the stage of effusion there is bulging of the precordia, a triangular area of dullness, the base of which is downward; the heart's sounds are muffled. In chronic pericarditis with adhesions there is often systolic retraction of the precordia. The causes of **P.** are rheumatism, the acute and chronic infectious diseases, Bright's disease, and extension of inflammation from neighboring parts. **P., Adhesive**, **P.** in which the two layers of pericardium tend to adhere. **P., Carcinomatous**, **P.** due to carcinoma of the pericardium. **P., Dry**, **P.** without effusion. **P., Fibrinous**, a form in which the membrane is covered with a fibrinous exudate, first soft and buttery in consistence, but later organizing. **P., Hemorrhagic**, a form in which the fluid is hemorrhagic. This is the case most often in tuberculous pericarditis; also in scorbutus and in cachectic conditions. **P., Localized**, a form giving rise to whitish areas, the so-called milk-spots. **P., Purulent**, **P.** in which the effused fluid becomes purulent. **P., Sero-fibrinous**, a form in which there is but little lymph or fibrin, but a considerable quantity of serous fluid. **P., Tuberculous**, **P.** due to tuberculous infection of the pericardium.

Pericardium (*per-e-kar'-de-um*) [*περί*, around; *καρδία*, heart]. The closed membranous sac enveloping the heart. Its base is attached to the central tendon of the diaphragm; its apex surrounds for a short distance the great vessels arising from the base of the heart. It consists of an outer fibrous coat, derived from the cervical fascia, and an inner serous coat. The sac normally contains from 5–20 grams of clear serous liquid. The part in contact with the heart (visceral **P.**) is termed the epicardium, the other is the parietal **P.** **P., Bread-and-butter**, a peculiar appearance produced in fibrinous pericarditis, by the rubbing of the two surfaces of the membrane over each other. **P., Shaggy**, a pericardium upon which, as the result of fibrinous pericarditis, thick, loose, shaggy layers of fibrin are deposited.

Pericardotomy (*per-e-kar-dot'-o-me*) [*peri-cardium*; *τομή*, a cutting]. The operation of opening the pericardium.

Pericecal (*per-e-se'-kal*) [*περί*, around; *cæcus*, blind]. Surrounding the cecum.

Perichondritis (*per-e-kon-dri'tis*) [*περί, around; χόνδρος, a cartilage; ιτις, inflammation*]. Inflammation of the perichondrium.

Perichondrium (*per-e-kon'-dre-um*) [*περί, around; χόνδρος, gristle; cartilage*]. The fibrous connective tissue covering the surface of cartilage.

Pericorneal (*per-e-kor'-ne-al*) [*περί, around; cornus, horny*]. Surrounding the cornea.

Pericranial (*per-e-kra'-ne-al*) [*περί, around; κρανιον, skull*]. Pertaining to the pericranium.

Pericranium (*per-e-kra'-ne-um*) [*περί, around; κρανιον, skull*]. The periosteum of the skull.

Peridectomy (*per-e-dek'-to-me*). Synonym of *Peritomy*.

Peridiastole (*per-e-di-as'-to-le*) [*περί, around; diastole*]. The pause between the systole and diastole.

Perididymitis (*per-e-did-im-i'-tis*) [*περί, around; διδυμος, testicle; ιτις, inflammation*]. Inflammation of the tunica albuginea testis.

Periencephalitis (*per-e-en-sef-al-i'-tis*) [*περί, around; εγκέφαλον, brain; ιτις, inflammation*]. Inflammation of the pia mater.

Perihepatitis (*per-e-hep-at-i'-tis*) [*περί, around; ήπαρ, liver; ιτις, inflammation*]. Inflammation of the peritoneum surrounding the liver.

Perilymph (*per'-e-limf*) [*περί, around; λυμφα, lymph*]. The fluid separating the membranous from the osseous labyrinth of the ear.

Perilymphatic (*per-e-lim-fat'-ik*) [*περί, around; λυμφα, lymph*]. 1. Pertaining to the perilymph. 2. Situated or occurring about a lymphatic vessel, as P. space.

Perimeningitis (*per-e-men-in-jit'-tis*) [*περί, around; μενινγίς, membrane; ιτις, inflammation*]. Inflammation of the dura mater.

Perimeter (*per-im'-et-er*) [*περί, around; μέτρον, measure*]. An instrument for measuring the extent of the field of vision. It consists ordinarily of a flat, narrow metal plate, bent in a semicircle, graduated in degrees, and fixed to an upright at its center by a pivot, on which it is movable. Various colored discs are moved along the metal plate and the point noted at which the person, looking directly in front of him, distinguishes the color.

Perimetritis (*per-e-met-ri'-tis*) [*περί, around; μήτρα, womb; ιτις, inflammation*]. Inflammation of the peritoneal covering of the uterus.

Perimetrium (*per-e-me'-tre-um*) [*περί, around; μήτρα, womb*]. The serous covering of the uterus.

Perimetry (*per-im'-et-re*) [*περί, around;*

μέτρον, measure]. The measuring of the field of vision.

Perimyelitis (*per-e-mi-el-i'-tis*) [*περί, around; μυελός, marrow; ιτις, inflammation*]. Inflammation of the pia mater of the spinal cord.

Perimysium (*per-e-mis'-e-um*) [*περί, around; μίς, muscle*]. The connective tissue enveloping the primary bundles of muscle-fibers.

Perineal (*per-e-ne'-al*) [*περίναιον, perineum*]. Pertaining to the perineum. **P. Body**, the mass of tissue composed of skin, muscle, and fascia, occupying the interval between the vagina and the rectum of the woman. **P. Cystotomy**, cystotomy performed through a perineal incision. **P. Fossa**, the ischio-rectal fossa. **P. Hernia**, a hernia perforating the perineum by the side of the rectum or between the rectum and the bladder or the vagina. **P. Section**, incision through the perineum for the relief of urethral stricture, the removal of calculi from the bladder, or the relief of other morbid conditions.

Perineocele (*per-e-ne'-o-sel*) [*περίναιον, perineum; κήλη, tumor*]. Perineal hernia.

Perineoplasty (*per-e-ne'-o-plas-te*) [*περίναιον, perineum; πλασσειν, to form*]. Plastic operation upon the perineum.

Perineorrhaphy (*per-e-ne-or'-a-fe*) [*περίναιον, perineum; ραφή, suture*]. Suture of the perineum, usually for the repair of a laceration caused during childbirth.

Perineotomy (*per-e-ne-ot'-o-me*) [*περίναιον, perineum; τομή, a cutting*]. Incision through the perineum.

Perinephric (*per-e-nef'-rik*) [*περί, around; νεφρός, kidney*]. Situated, or occurring around the kidney, as P. Abscess.

Perinephritic (*per-e-nef-rit'-ik*) [*περί, around; νεφρός, kidney; ιτις, inflammation*]. 1. Pertaining to perinephritis. 2. Improperly used instead of perinephric.

Perinephritis (*per-e-nef-rit'-tis*) [*περί, around; νεφρός, the kidney; ιτις, inflammation*]. Inflammation of the tissues surrounding the kidney.

Perineum (*per-e-ne'-um*) [*περίναιον*]. That portion of the body included in the outlet of the pelvis, bounded in front by the pubic arch, behind by the coccyx and great sacro-sciatic ligaments, and at the sides by the tuberosities of the ischium. It is occupied by the terminations of the rectum, the urethra, and the root of the penis, together with their muscles, fasciae, vessels, and nerves.

Perineuritis (*per-e-nu-ri'-tis*) [*περί, around; νευρον, nerve; ιτις, inflammation*]. Inflammation of the perineurium.

Perineurium (*per-e-nu'-re-um*) [*περί, around; νευρον, nerve*]. The connective-tissue sheath investing a funiculus or primary bundle of nerve-fibers.

Perinuclear (*per-e-nu'-kle-ar*) [*περί, around; nucleus*]. Surrounding the nucleus.

Period (*pe'-re-ad*) [*περίοδος, period, from περί, around; ὁδός, way*]. The space of time during which anything is in progress or an event takes place. **P., Incubation-.** See *Incubation*. **P., Menstrual, P., Monthly**, the menses.

Periodic (*pe-re-ad'-ik*) [*περίοδος, period*]. Recurring at more or less regular intervals.

Periodicity (*pe-re-o-dis'-it-e*) [*περίοδος, period, from περί, around; ὁδός, way*]. Recurrence at regular intervals.

Periodontal (*per-e-o-don'-tal*) [*περί, around; ὁδός, tooth*]. Surrounding a tooth, as the P. membrane, that lining the cement of a tooth.

Periodontitis (*per-e-o-don-ti'-tis*) [*περί, around; ὁδός, tooth; ιτις, inflammation*]. Inflammation of the periodontal membrane.

Periodontium (*per-e-o-don'-she-um*) [*περί, around; ὁδός, tooth*]. The membrane surrounding a tooth; the periodontal membrane.

Periodoscope (*per-e-o-d'-o-skōp*) [*περίοδος, period; σκοπεῖν, to inspect*]. A calendar in the form of a movable dial, used in determining the probable date of confinement.

Periophoritis (*per-e-o-off-or-i'-tis*) [*περί, around; ὠόν, egg; φορέω, bearing; ιτις, inflammation*]. Inflammation of the peritoneum and connective tissue covering the ovary.

Periometry (*per-e-op-tom'-et-re*) [*περί, around; ὁπτός, visible; μέτρον, measure*]. The measurement of the limits of the visual field.

Periorbita (*per-e-or'-bit-ah*) [*περί, around; orbita, orbit*]. The periosteum of the eye-socket.

Periorbital (*per-e-or'-bit-al*) [*περί, around; orbita, orbit*]. 1. Surrounding the orbit. 2. Pertaining to the periorbita.

Periosteal (*per-e-os'-te-al*) [*περί, around; ὀστέον, bone*]. Pertaining to the periosteum.

Periosteitis (*per-e-os-te-i'-tis*). See *Periostitis*.

Periosteotome (*per-e-os'-te-o-lōm*) [*περί, around; ὀστέον, bone; τομή, a cutting*]. An instrument for incising the periosteum, and scraping it from the bone.

Periosteotomy (*per-e-os-te-ol'-o-me*) [*περί, around; ὀστέον, bone; τομή, a cutting*]. The operation of incising the periosteum.

Periosteum (*per-e-os'-te-um*) [*περί, around; ὀστέον, bone*]. A fibrous membrane investing the surfaces of bones, except at the points of tendinous and ligamentous attachment and on the articular surfaces, where cartilage is substituted.

Periostitis (*per-e-os-ti'-tis*) [*περί, around; ὀστέον, bone; ιτις, inflammation*]. Inflammation of the periosteum. It may be acute or chronic, the latter being the more frequent

form. Acute P. is either traumatic or the result of infection; the chronic is due to traumatism, syphilis, tuberculosis, or actinomycosis. In the acute there is swelling and diffuse suppuration with fever and other constitutional symptoms; in the chronic, pain, which is usually worse at night, swelling, and tenderness. **P. albuminosa**, a mild form of inflammation characterized by the formation of a clear, ropy, albuminous liquid resembling synovia. It is most common in the young and is unaccompanied by fever. **P., Hemorrhagic, P.** accompanied by bleeding between the periosteum and the bone.

Periostosis (*per-e-os-to'-sis*) [*περί, around; ὀστέον, bone*]. An osseous formation on the exterior of a bone.

Periotic (*per-e-ol'-tik*) [*περί, around; ὄς, ear*]. 1. Situated about the ear. 2. Of or pertaining to the parts immediately about the internal ear. 3. The petrous and mastoid parts of the temporal bone.

Peripachymeningitis (*per-e-pak-e-men-in-jit'-tis*) [*περί, around; παχύς, thick; μῆνιγξ, membrane; ιτις, inflammation*]. Inflammation of the connective tissue between the dura mater and the bone.

Peripancatitis (*per-e-pān-kre-at-i'-tis*) [*περί, around; πάγκρεας, pancreas; ιτις, inflammation*]. Inflammation of the tissues about the pancreas.

Peripherad (*per-if'-er ad*) [*περί, around; φέρειν, to carry; ad, toward*]. Toward the periphery.

Peripheral, Periphereal (*per-if'-er-al, per-e-ser'-ik*) [*periphery*]. Pertaining to or placed near the periphery.

Peripheraphose (*per-if'-er-a-fōs*). See under *Phose*.

Peripherophose (*per-if'-er-o-fōs*). See under *Phose*.

Periphery (*per-if'-er-e*) [*περί, around; φέρειν, to carry*]. Circumference; the external surface.

Periphlebitis (*per-e-φleb-i'-tis*) [*περί, around; φλέψ, vein; ιτις, inflammation*]. Inflammation of the tissues about a vein.

Perioplast (*per'-e-plast*) [*περί, around; πλάσσειν, to mold, form*]. The protoplasm of a cell surrounding the nucleus; cell-protoplasm.

Peripleuritis (*per-e-plu-ri'-tis*) [*περί, around; πλευρά, rib; ιτις, inflammation*]. Inflammation of the tissues surrounding the pleura.

Peripneumonia (*per-e-nu-mo'-ne-ah*) [*περί, around; πνεύμων, lung*]. 1. Pneumonia. 2. Pleuropneumonia.

Peripolar (*per-e-po'-lar*) [*περί, around; πόλος, pivot*]. Surrounding a pole or the poles.

Periportal (*per-e-por'-tal*) [*περί, around; porta, door*]. Surrounding the portal vein. **P. Carcinoma**, a primary carcinoma devel-

oping around the portal vein, beginning at its entrance into the liver, thence extending along the portal vessels to the remotest branches.

Periproctitis (*per-e-prok-tis*) [*περί, around; πρωκτός, anus; ιτις, inflammation*]. Inflammation of the areolar tissue about the rectum or anus.

Perirectal (*per-e-rek'tal*) [*περί, around; rectum, rectum*]. About the rectum.

Perirenal (*per-e-re'nal*) [*περί, around; ren, kidney*]. Around the kidney.

Perisalpingitis (*per-e-sal-pin-jit-is*) [*περί, around; σαλπιγξ, tube; ιτις, inflammation*]. Inflammation of the peritoneal covering of the Fallopian tube.

Periscopic (*per-e-skop'ik*) [*περί, around; σκοπεῖν, to see*]. Designed for looking around, as a P. lens. See *Lens*.

Perisplenitis (*per-e-splen-i'tis*) [*περί, around; σπλήν, spleen; ιτις, inflammation*]. Inflammation of the peritoneal coat of the spleen.

Perissad (*per'-is-ad*) [*περισσός, odd*]. 1. Having an odd quantivalence, as, *e. g.*, nitrogen, the quantivalence of which is three or five. 2. An element having such a quantivalence.

Peristalsis (*per-e-stal'sis*) [*περί, around; στάσις, constriction*]. A peculiar wave-like movement seen in tubes provided with longitudinal and transverse muscular fibers. It consists in a narrowing and shortening of a portion of the tube, which then relaxes while a lower portion becomes shortened and narrowed. By means of this movement the contents of the tube are forced toward the opening. **P., Reversed**, peristaltic movement opposite to the normal direction.

Peristaltic (*per-e-stal'tik*) [*περί, around; στάσις, constriction*]. Pertaining to or resembling peristalsis. **P. Unrest**, a common symptom of neurasthenia, consisting in increased peristaltic movements of the stomach coming on shortly after eating, with borborygmus and gurgling.

Perisystole (*per-e-sis'to-le*) [*περί, around; συστολή, contraction*]. The slight interval between the diastole and systole.

Perithelium (*per-e-the'le-um*) [*περί, around; θηλή, nipple*]. The layer of cells surrounding the capillaries and smaller vessels.

Peritomy (*per-it'o-me*) [*περί, around; τομή, a cutting*]. 1. The removal of a strip of conjunctival and subconjunctival tissue from about the cornea for the relief of pannus. 2. Circumcision.

Peritoneal (*per-e-ton-e'al*) [*peritoneum*]. Pertaining to the peritoneum.

Peritoneum (*per-e-ton-e'um*) [*περί, around; τένειν, to stretch*]. The serous membrane lining the interior of the abdominal cavity

and surrounding the contained viscera. In tracing its extension we may begin at the diaphragm and take two layers. From the diaphragm two layers of peritoneum proceed to the liver; they separate to enclose that organ, meet again on its under aspect, and pass on, as the gastrohepatic omentum, to the lesser curvature of the stomach. They embrace the stomach, and, meeting again at its greater curvature, pass down in front of the small intestine, forming the great omentum. They are then reflected upward as far as the transverse colon, which they enclose, meeting again at the back of the colon, and proceeding to the spine as the transverse mesocolon. Here the two layers diverge; the upper ascends in front of the pancreas to the under surface of the diaphragm, the starting-point. The lower layer is reflected from the spine, over the small intestines, as the mesentery. From the root of the mesentery it passes into the pelvis, invests the upper part of the rectum, and is thence reflected on the bladder, forming between the two the rectovesical pouch. In the woman it is reflected from the rectum to the upper part of the vagina, and thence over the uterus, from which it proceeds to the bladder. From the bladder it passes up the anterior wall of the abdomen to the diaphragm. The following structures are completely invested by peritoneum:—The stomach, liver, spleen, first portion of the duodenum, jejunum, ileum, transverse colon, sigmoid flexure, the upper part of the rectum and of the uterus, and the ovaries.

Peritonitis (*per-e-ton-i'tis*) [*περιτόναιον, peritoneum; ιτις, inflammation*]. Inflammation of the peritoneum. It may be acute or chronic. Acute P. may be due to exposure to cold and wet (Idiopathic P.), traumatism, perforation of an abdominal viscus, extension from neighboring parts, rheumatism, or to Bright's disease. The symptoms are moderate fever, a wiry pulse, abdominal pain, tenderness, and distention; the patient lies on his back with the thighs flexed; there is vomiting and constipation. Chronic P. is due to tuberculosis, syphilis, carcinoma, nephritis, or it may be the sequel of an acute attack. **P., Adhesive**, peritonitis with adhesion between the parietal and visceral layers. **P., Diffuse**, that affecting the entire peritoneum. **P., Puerperal**, that following labor, and usually due to septic infection. **P., Septic**, peritonitis due to the microorganisms of supuration.

Peritonsillar (*per-e-ton'sil-ar*) [*περί, around; tonsilla, tonsil*]. About the tonsil.

Perityphlitis (*per-e-tif-li'tis*) [*περί, around; τυφλός, blind; ιτις, inflammation*]. Inflammation of the peritoneum surrounding the cecum.

- Periurethritis** (*per-e-u-re-thrit-is*) [*περί, around; οὐρήθρα, urethra; ιτις, inflammation*]. Inflammation of the connective tissue about the urethra.
- Periuterine** (*per-e-u'-ter-in*) [*περί, around; uterus, uterus*]. About the uterus.
- Perivascular** (*per-e-vas'-ku-lar*) [*peri, around; vascular*]. About a vessel.
- Perkinsism** (*per'-kins-ism*). A method of empiric treatment devised by Elisha Perkins, a Connecticut physician. It consisted in drawing over the affected part the extremities of two rods (metallic tractors) of different metals; tractoration.
- Perlèche** (*la*) (*lah pār-lash*) [Fr.]. A peculiar contagious disease of the mouth occurring in children. It consists in a thickening and desquamation of the epithelium at the angles of the mouth, with occasionally the formation of small fissures, giving rise to a smarting sensation in the lips. The disease is probably microbic in origin.
- Perles'Anemia-bodies**. Small, club-shaped, actively motile bodies, from 3 to 4 μ in length, found by Perles in the blood in three cases of pernicious anemia.
- Perles or Pearls**, Lænnec's, the rounded gelatinous masses of sputum seen in the early stage of an attack of bronchial asthma.
- Perlsucht** [G.]. See *Pearl-disease*.
- Permanent** (*per'-man-ent*) [*per, through; manere, to remain*]. Lasting; fixed; enduring, as P. teeth.
- Permanganate** (*per-man'-gan-at*). A salt of permanganic acid. See *Manganese*.
- Permanganic Acid** (*per-man'-gan'-ik*), HMnO₄. A monobasic acid known chiefly in its salts.
- Permeable** (*per'-me-a-bl*) [*per, through; meare, to pass*]. Capable of affording passage. P. Stricture, a stricture that permits the passage of an instrument.
- Pernicious** (*per-nish'-us*) [*perniciōsus, destructive*]. Highly destructive; of intense severity; deadly; fatal. P. Anemia, a disease of the blood characterized by a great diminution in the number of red corpuscles, and a relatively smaller diminution of the hemoglobin, by the presence in the blood of poikilocytes, macrocytes, microcytes, and nucleated red corpuscles. A symptomatic pernicious anemia may be produced by parasites in the intestinal canal, by atrophy of the gastric mucous membrane, and by hemorrhage. The nature of primary pernicious anemia is still obscure. Most writers believe that the disease is dependent upon increased hemolysis, *i. e.*, blood-destruction. The disease most common in middle life is usually fatal, although recoveries are reported in several instances. P. Malaria. See *Malaria*. P. Vomiting, persistent, uncontrollable vomiting, occurring in pregnancy.
- Pernio** (*per'-ne-o*) [L.]. Synonym of *Chilblain*.
- Peroneal** (*per-o-ne'-al*) [*περόνη, pin*]. Pertaining to the fibula.
- Peroneus** (*per-o-ne'-us*) [*περόνη, pin*]. Pertaining to the fibula. P. Muscle. See *Muscles, Table of*.
- Peronospora** (*per-o-nos'-po-rah*) [*περόνη, point; σπόρος, spore*]. A genus of fungi producing mildew. P. ferrari, P. barcinonæ, a species that was supposed to cause cholera. P. lutea, a species that has been held to be the cause of yellow fever.
- Per os** [L.]. By the mouth.
- Perosomus** (*pe-ro-so'-mus*) [*πρόος, maimed; σῶμα, body*]. A monster presenting malformation of the entire body.
- Peroxid** (*per-oks'-id*) [*per, through; ὀξύς, sharp*]. That oxid of any base which contains the most oxygen.
- Perplication** (*per-plik'-a'-shun*) [*per, through; plicare, to fold*]. The operation of turning an incised vessel upon itself by drawing its end through an incision in its own wall.
- Per rectum** [L.]. By the rectum.
- Personal** (*per'-son-al*) [*persona, a person*]. Pertaining to a person. P. Equation, the peculiar difference of individuals in their reaction to various orders of stimuli.
- Perspiration** (*per-spir'-a'-shun*) [*perspiratio; perspirare, to perspire*]. 1. The secretion of sweat. 2. The sweat. P. Insensible, that which takes place constantly, the fluid being evaporated as fast as secreted. P., Sensible, that accumulating in visible drops or beads; the sweat.
- Perspire** (*per-spir'*) [*perspirare*]. To sweat.
- Pertussis** (*per-tus'-is*). See *Whooping-cough*.
- Peruvian** (*pe-ru'-ve-an*) [*Peru*]. Pertaining to Peru. P. Bark. See *Cinchona*.
- Pervious** (*per'-ve-us*) [*per, through; via, way*]. Open; permeable.
- Pea** (*pes*) [L.]. A foot, or foot-like structure. P. accessorius, the eminentia collateralis, a smooth white eminence in the brain, situated at the junction of the posterior and descending cornua of the lateral ventricle. P. anserinus, goose's foot; the radiate branching of the facial nerve after its exit at the side of the face. P. hippocampi major, the lower portion of the hippocampus major. P. hippocampi minor. The same as *Hippocampus minor*.
- Pessary** (*pes'-ar-e*) [*πέσος, an oval-shaped stone*]. An instrument placed in the vagina to hold the uterus in position.
- Pestiferous** (*pes-tif'-er-ous*) [*pestis, pest; ferre, to bear*]. Causing pestilence.
- Pestilence** (*pes'-til-ens*) [*pestis, plague*]. Any deadly epidemic disease, especially the plague.

Pestilential (*pes-til-en'-shal*) [*pestis*, plague]. Having the nature of or producing a pestilence.

Pestle (*pes'-l*) [*pistillum*]. The instrument with which substances are rubbed in a mortar.

Petechia (*pet-e'-ke-ah*) [It., *peteche*, a flea-bite]. A small spot beneath the epidermis, due to an effusion of blood.

Petechial (*pet-e'-ke-al*) [It., *peteche*, flea-bite]. Characterized by or of the nature of petechiae. **P. Fever**, typhus fever.

Petit, Canal of (*pet-e'*). See *Canal*.

Petit Mal (*pet-e mahl*) [Fr., little illness]. A slight epileptic seizure characterized by a momentary, scarcely recognizable loss of consciousness, often with an upward staring of the eyes and fibrillary movements of the facial muscles. See *Epilepsy*.

Petit's Triangle. See *Triangles, Table of*.

Petri's Dishes. Shallow, double glass-dishes, for use in cultivating bacteria.

Petrification (*pet-rif-ak'-shun*) [*πίτρα*, a stone; *facere*, to make]. Conversion into stone, as **P. of the fetus**, the formation of a lithopedion.

Petrolatum (*pet-ro-la'-tum*) [*πίτρα*, rock; *oleum*, oil]. A jelly-like preparation obtained from the residuum of petroleum, soluble in ether, insoluble in water and alcohol, and known commercially as vaselin or cosmolin. It is used as a basis for ointments and as an emollient. **P. liquidum** (U. S. P.), liquid petrolatum. **P. molle** (U. S. P., B. P.), soft petroleum-ointment. **P. spissum** (U. S. P.), hard petroleum.

Petroleum (*pet-ro'-le-um*) [*πίτρα*, rock; *oleum*, oil]. An oily liquid issuing from the earth in various places, and consisting of a mixture of hydrocarbons with small amounts of oxidation-products. The hydrocarbons belong chiefly to the paraffin-series.

Petromastoid (*pet-ro-mas'-toid*) [*πίτρα*, stone; *mastoid*]. Pertaining to the petrous and mastoid portions of the temporal bone. **P. Canal**, a short passage connecting the mastoid sinuses and the tympanic cavity. **P. Foramen**, the tympanic orifice of the petromastoid canal.

Petrooccipital (*pet-ro-ok-sip'-it-al*) [*πίτρα*, stone; *occiput*, occiput]. Pertaining to the petrous portion of the temporal bone and to the occipital bone.

Petrosal (*pet-ro'-sal*) [*πίτρα*, rock]. 1. Pertaining to the petrous portion of the temporal bone, as the **P. sinus** (superior and inferior), **P. nerves**. 2. The petrous portion of the temporal bone itself.

Petroselinum (*pet-ro-se-li'-num*) [*πίτρα*, rock; *σέλινον*, parsley]. See *Parsley*.

Petrosomastoid (*pet-ro-so-mas'-toid*). Synonym of *Petromastoid*.

Petrosphenoid (*pet-ro-sfe'-noid*) [*πίτρα*,

rock; *σφην*, wedge; *ειδος*, like]. Pertaining to the petrous portion of the temporal bone and the sphenoid bone. **P. Suture**, the suture between the temporal bone and the great wing of the sphenoid bone.

Petrosquamosal, Petrosquamous (*pet-ro-skwa-mo'-sal*, *pet-ro-skwa'-mus*) [*πίτρα*, rock; *squama*, scale]. Pertaining to the petrous and squamous portions of the temporal bone. **P. Fissure, P. Suture**, the line of juncture of the squamous and petrous portions of the temporal bone. **P. Sinus**, a venous passage formed in the dura mater at the junction of the petrous and squamous portions of the temporal bone. It opens into the lateral sinus.

Petrous (*pe'-trus*) [*πίτρα*, rock]. 1. Stony, of the hardness of stone, as the **P. portion** of the temporal bone. 2. See *Petrosal*.

Pettenkoffer's Test. A test for the presence of biliary acids. A few drops of the suspected liquid are dropped into a fresh solution of sugar and dilute sulphuric acid. If biliary acids be present a purplish-crimson color is produced.

Peyer's Glands or Patches. Aggregations of lymph-follicles situated in the mucous membrane of the lower part of the small intestine, opposite the mesenteric attachment.

Pfeiffer's Phenomenon. When a mixture of cholera-germs and cholera antitoxic serum is injected into the peritoneal cavity of a guinea-pig, the microorganisms are quickly destroyed and dissolved. This does not occur when normal or other serum is employed, nor are other organisms destroyed when injected together with cholera-serum. The same phenomenon has been observed in the case of the typhoid-bacillus and typhoid antitoxic serum, and is a valuable differential sign.

Pflüger's Law. The law that a nerve-trunk is stimulated by the appearance of catelectrotonous and the disappearance of anelectrotonous, but not under the reverse condition.

Phacoidoscope (*fa-koid'-o-skop*). Synonym of *Phacoscope*.

Phacosclerosis (*fa-ko-skle-ro'-sis*) [*φακος*, lens; *σκληρός*, hard]. Hardening of the crystalline lens.

Phacoscope (*fa'-ko-skop*) [*φακος*, lens; *σκοπεῖν*, to inspect]. An instrument for observing the accommodative changes of the lens.

Phagedena (*faj-ed'-nah*) [*φαγάδαινα*, from *φαγεῖν*, to eat]. A rapidly spreading destructive ulceration of soft parts.

Phagedenic (*faj-ed'-en'-ik*) [*φαγάδαινα*; *φαγεῖν*, to eat]. Of the nature of phagedena. **P. Chancroid**, a chancroid that spreads rapidly and destroys a large amount of tissue.

Phagocyte (*fag-o-sit*) [*φαγεῖν*, to eat; *κύτος*, cell]. A cell having the property of englobing and digesting foreign or other particles

harmful to the body. Phagocytes are either fixed—endothelial cells, fixed connective-tissue cells—or free—the wandering cells or leukocytes. A large phagocyte is termed a *macrophage*; a small one, a *microphage*.

Phagocytic (*fag-o-sit'-ik*) [*φαγεῖν*, to eat; *κύτος*, cell]. Of, pertaining to, or caused by phagocytes.

Phagocytosis (*fag-o-si-to'-sis*) [*φαγεῖν*, to eat; *κύτος*, cell]. The ingestion of foreign or other particles, principally bacteria, by certain cells. P. has been claimed to be the cause of immunity against infectious diseases.

Phakitis (*fa-ki'-tis*) [*φακός*, lens; *ιτις*, inflammation]. Inflammation of the crystalline lens of the eye; a condition that has, however, not been observed.

Phako-. See *Phaco-*.

Phalacrosis (*fal-ak-ro'-sis*) [*φαλακρός*, bald]. Baldness.

Phalangeal (*fa-lan'-je-al*) [*φάλαγξ*, phalanx]. Pertaining to a phalanx.

Phalanges (*fa-lan'-jēz*) [*φάλαγξ*, phalanx]. Plural of *Phalanx*.

Phalanx (*fa'-lanks*) [*φάλαγξ*, phalanx]. 1. One of the bones of the fingers or toes. 2. One of the delicate processes of the head-plate of the outer rod of Corti projecting beyond the inner rod.

Phallic (*fal'-ik*) [*φαλλός*, penis]. Pertaining to the penis.

Phallus (*fal'-us*) [*φαλλός*, penis]. Penis.

Phaneroscope (*fan'-er-o-skōp*) [*φανερός*, visible; *σκοπεῖν*, to see]. An instrument for rendering the skin transparent; it is used in examining for diseases of the skin, such as lupus.

Phantasm (*fan'-tazm*) [*φαντάζειν*, to render visible]. An illusive perception of an object that does not exist; an optic illusion; an apparition.

Phantom (*fan'-tum*) [*φαντάζειν*, to make visible]. 1. An apparition. 2. A model of a part or the whole of the human body used in practising various operations and procedures. **P.-tumor**, a tumor-like swelling produced artificially by the contraction of a muscle or by other causes.

Pharmaceutic (*far-ma-su'-tik*) [*φαρμακεύειν*, to administer a drug]. Pertaining to pharmacy.

Pharmaceutics (*far-ma-su'-tiks*) [*φαρμακεύειν*, to administer a drug]. See *Pharmacy* (1st def.).

Pharmacist (*far'-ma-sist*) [*φάρμακον*, a drug]. An apothecary.

Pharmaco- (*far-ma-ko-*) [*φάρμακον*, drug]. A prefix meaning pertaining to drugs.

Pharmacodynamics (*far-ma-ko-di-nam'-iks*) [*φάρμακον*, drug; *δύναμις*, force]. The science of the action of drugs.

Pharmacognosis, Pharmacognosy (*far-*

ma-kog'-no-sis, *far-ma-kog'-no-se*) [*φάρμακον*, drug; *γνώσις*, knowledge]. The science of crude drugs.

Pharmacography (*far-ma-kog'-ra-fē*). See *Pharmacognosy*.

Pharmacologist (*far-ma-kol'-o-jist*) [*φάρμακον*, drug; *λόγος*, science]. One versed in pharmacology.

Pharmacology (*far-ma-kol'-o-jē*) [*φάρμακον*, drug; *λόγος*, science]. The science of the nature and properties of drugs.

Pharmacopeia (*far-ma-ko-pe'-ah*) [*φάρμακον*, drug; *ποιεῖν*, to make]. A collection of formulas and methods for the preparation of drugs, especially a book of such formulas recognized as a standard, as the United States or British P. The former is issued every ten years, under the supervision of a national committee.

Pharmacopeial (*far-ma-ko-pe'-al*) [*φάρμακον*, drug; *ποιεῖν*, to make]. Contained in or sanctioned by the pharmacopeia.

Pharmacy (*far'-ma-se*) [*φαρμακεία*, the use of drugs]. 1. The art of preparing, compounding, and dispensing medicines. 2. A drug-store.

Pharyngeal (*far-in'-je-al*) [*φάρυγξ*, pharynx]. Pertaining to the pharynx. **P. Tonsil**. See *Luschka's Tonsil*. **P. Tubercle**, a small elevation near the middle of the inferior surface of the basilar process of the occipital bone, for the attachment of the pharynx.

Pharyngectomy (*far-in-jek'-to-me*) [*φάρυγξ*, pharynx; *ἐκτομή*, excision]. Excision of the pharynx.

Pharyngismus (*far-in-jis'-mus*) [*φάρυγξ*, pharynx]. Spasm of the pharynx.

Pharyngitis (*far-in-ji'-tis*) [*φάρυγξ*, pharynx; *ιτις*, inflammation]. Inflammation of the pharynx. **P., Acute, P., Catarrhal**, is due to exposure to cold, to the action of irritant substances, or to certain infectious causes, and is characterized by pain on swallowing, by dryness, later by moisture, and by congestion of the mucous membrane. **P., Chronic**, this is generally the result of repeated acute attacks, and is associated either with hypertrophy of the mucous membrane (**P., Hypertrophic**), or with atrophy (**P., Atrophic**). **P., Croupous, P., Diphtheric**, is characterized by the presence of a false membrane, the product of the action of the diphtheria-bacillus. **P., Granular**, a form of chronic pharyngitis in which the mucous membrane has a granular appearance.

Pharyngo- (*far-in'-go-*) [*φάρυγξ*, pharynx]. A prefix signifying pertaining to the pharynx.

Pharyngocele (*far-in'-go-sēl*) [*φάρυγξ*, pharynx; *κήλη*, tumor]. A hernia or pouch of the pharynx projecting through the pharyngeal wall.

Pharyngolaryngeal (*far-in-go-lar-in'-je-al*) [*pharynx*, *larynx*]. Pertaining both to the pharynx and the larynx.

Pharyngolaryngitis (*far-in-go-lar-in-jit-tis*) [*pharynx*, *larynx*; *itis*, inflammation]. Simultaneous inflammation of the larynx and pharynx.

Pharyngomycosis (*far-in-go-mi-ko'-sis*) [*pharynx*; *mycosis*, fungus]. Disease of the pharynx due to the action of fungi.

Pharyngoplegia (*far-in-go-ple'-je-ah*) [*pharynx*; *plegia*, a stroke]. Paralysis of the muscles of the pharynx.

Pharyngoscope (*far-in-go-skop*) [*pharynx*; *σκοπεῖν*, to inspect]. An instrument for use in examining the pharynx.

Pharyngoscopy (*far-in-go-skop-ee*) [*pharynx*; *σκοπεῖν*, to examine]. Examination of the pharynx with the pharyngoscope.

Pharyngospasm (*far-in-go-spazm*) [*pharynx*; *σπασμός*, spasm]. Spasmodic contraction of the pharyngeal muscles.

Pharyngotomy (*far-in-go-tom*) [*pharynx*; *τομή*, a cutting]. An instrument for incising the pharynx.

Pharyngotomy (*far-in-go-tom-ee*) [*pharynx*; *τομή*, a cutting]. Incision into the pharynx. **P., Inferior**, one in which the tissues between the hyoid bone and the cricoid cartilage are divided. **P., Lateral**, incision into one side of the pharynx. **P., Subhyoidan**, that through the thyrohyoid membrane.

Pharyngotonsillitis (*far-in-go-ton-sil-it-tis*) [*pharynx*; *tonsillitis*]. Inflammation of the pharynx and the tonsil.

Pharynx (*far-in-ks*) [*pharynx*, throat]. The musculomembranous pouch situated back of the nose, mouth, and larynx, and extending from the base of the skull to a point opposite the sixth cervical vertebra, where it becomes continuous with the esophagus. It is lined by mucous membrane, covered in its upper part with columnar ciliated epithelium, in its lower part with stratified epithelium. On the outside of this is a layer of fibrous tissue, the pharyngeal aponeurosis. This in turn is surrounded by the muscular coat. The upper portion of the pharynx communicates with the nose through the posterior nares, is known as the nasopharynx, and functionally belongs to the respiratory tract; the lower portion is divided into the oropharynx and laryngopharynx, and is a part of the digestive tract. The pharynx communicates with the middle ear by means of the Eustachian tube.

Phenacetin (*fe-nas'-et-in*) [*phenol*; *acetum*, vinegar], $C_{10}H_{13}NO_2$. Acetphenetidin, a compound derived from phenol, having antipyretic and antineuralgic properties. It is crystalline, tasteless, and almost insoluble in water. Dose gr. iv-xxx (0.26-2.0).

Phenate (*fe-nat*) [*phenol*]. A compound of phenol and a base; a carbolate.

Phenetidin (*fe-net'-id-in*) [*phenol*], $C_8H_{11}NO$. The base from which phenacetin is prepared by substitution.

Phenic Acid [*phenol*]. See *Acid, Carboic*.

Phenocoll (*fe'-no-kol*) [*phenol*], $C_{10}H_{14}N_2O_2$. Amidophenacetin, a substance resembling phenacetin, and the hydrochlorate of which is used as an antipyretic. Dose gr. x-xv (0.65-1.0).

Phenol (*fe'-nol*) [*phoinix*, purple-red]. 1. Carboic acid. 2. Any derivative of benzene homologous with phenol.

Phenolphthalein (*fe-nol-tha'-le-in*) [*phoinix*, purple-red; *phthalic*]. A substance produced by the action of phenol on phthalic acid and used generally in a 1 per cent. solution in 50 per cent. alcohol, as a delicate test for acids and alkalis. It is turned red by alkalis and decolorized by acids.

Phenolsulphonic Acid (*fe-nol-sul-fon'-ik*). *Sulphocarboic Acid*.

Phenyl (*fe'-nil*) [*phoinix*, purple-red; *ῥηγν*, matter]. The univalent radicle, C_6H_5 , of phenol.

Phenyl-glucosazone (*fe-nil-glu-ko'-saz-on*), $C_{16}H_{25}N_3O_4$. A yellow crystalline compound produced in the phenyl-hydrazin test for glucose.

Phenyl-glycuronic Acid (*glik-u-ron'-ik*). A crystalline body, a compound of phenol and glycuronic acid, occurring in the urine after the ingestion of phenol.

Phenyl-hydrazin (*fe-nil-hi'-dra-zin*), $C_6H_5N_2$. A liquid base, crystallizing in plates, the hydrochlorate of which is used as a test for sugar (Phenyl-hydrazin test).

Phenylic (*fe-nil'-ik*) [*phoinix*, purple-red]. Pertaining to or containing phenyl.

Phenyl-urethane (*fe-nil-u'-reth-an*), $C_9H_{11}NO_2$. See *Euphorin*.

Phial (*fi'-al*) [*φιάλη*, a saucer]. A small glass bottle; a vial.

Phimosis (*fi-mo'-sis*) [*φίμωσις*, a muzzle]. Elongation and constriction of the prepuce, so that the latter cannot be retracted over the glans penis.

Phlebectasia, Phlebectasis (*feb-ek-ta'-sah, feb-ek'-tas-is*) [*φλέψ*, vein; *εκτασις*, dilatation]. Dilatation of a vein; varicosity.

Phlebectomy (*feb-ek'-to-me*) [*φλέψ*, vein; *εκτομή*, excision]. Excision of a vein.

Phlebitis (*feb-i'-tis*) [*φλέψ*, vein; *itis*, inflammation]. Inflammation of a vein. This is generally suppurative (suppurative P.), and is the result of the extension of suppuration from adjacent tissues. It leads to the formation of a thrombus within the vein (Thrombophlebitis), which may break down and cause the distribution of septic emboli to various parts of the body. When not due to

a suppurative process the P., called plastic, adhesive, or proliferative, may give rise to obliteration of the vein. The symptoms of P. are pain and edema of the affected part, redness along the course of the vein, the latter appearing as a hard, tender cord. P., Sinus-, phlebitis of the sinuses of the dura mater.

Phlebo- (*feb'-o-*) [*φλέψ*, vein]. A prefix denoting pertaining to a vein.

Phlebogram (*feb'-o-gram*) [*φλέψ*, vein; *γράφειν*, to write]. A tracing of the movements of a vein by the sphygmograph.

Phlebolite, **Phlebolith** (*feb'-o-lith*, *feb'-o-lith*) [*φλέψ*, vein; *λίθος*, stone]. Vein-stone, a hard concretion sometimes found in veins, and produced by calcareous infiltration of a thrombus.

Phlebosclerosis (*feb-o-skle-ro'-sis*) [*φλέψ*, a vein; *σκληρός*, hard]. Sclerosis of a vein.

Phlebotomy (*feb-o'-o-me*) [*φλέψ*, vein; *τομή*, a cutting]. Opening of a vein for the purpose of blood-letting. The vein most often selected is the median cephalic at the bend of the elbow.

Phlegm (*flem*) [*φλέγμα*, phlegm]. 1. A viscid, stringy mucus, secreted by the mucosa of the upper air-passages. 2. One of the four humors of the old writers.

Phlegmasia (*fleg-ma'-ze-ah*) [*φλέγειν*, to burn]. Inflammation. P. alba dolens, milk-leg, a painful swelling of the leg, beginning either at the ankle and ascending, or at the groin and extending down the thigh, its usual cause being septic infection after labor.

Phlegmatic (*fleg-mat'-ik*) [*φλέγμα*, phlegm]. Full of phlegm; hence, indifferent, apathetic.

Phlegmon (*fleg'-mon*) [*φλεγμονή*, inflammation]. An inflammation characterized by the spreading of a purulent or fibrinopurulent exudate within the tissues.

Phlegmonous (*fleg'-mon-us*) [*φλεγμονή*, inflammation]. Of the nature of or pertaining to phlegmon.

Phlogogenic (*flog-o-jen'-ik*) [*φλόξ*, flame; *γεννάν*, to produce]. Causing inflammation.

Phlogosin (*flog'-o-sin*) [*φλόγωσις*, inflammation, from *φλόξ*, flame]. A crystalline body isolated from cultures of pyogenic staphylococci, and causing suppuration when injected beneath the skin or introduced into the eye.

Phloridzin, **Phlorizin** (*flor-id'-zin*, *flor-i'-zin*) [*φλοιός*, bark; *ρίζα*, root], $C_{21}H_{34}O_{10} \cdot 2H_2O$. A bitter crystalline glucosid occurring in the root and trunk of the apple, pear, and other fruit-trees. It is said to possess antipyretic properties. Dose gr. v-x (0.32-0.65). Given to lower animals it produces glycosuria. P.-diabetes, the glycosuria induced in lower animals, especially dogs, by the administration of phloridzin.

Phloroglucin (*flo-ro-gluc'-sin*) [*φλοιός*, bark; *γλυκύς*, sweet], $C_6H_4O_2-C_6H_3(OH)_3$. A crystalline substance found in the bark of the cherry, pear, apple, and other trees, and used as a test for woody tissue (lignin) and hydrochloric acid.

Phlyctena (*flik-te'-nah*) [*φλύκταινα*, blister]. A vesicle.

Phlyctenoid (*flik'-ten-oid*) [*φλύκταινα*, blister; *ειδος*, like]. See *Phlyctenular*.

Phlyctenula, **Phlyctenule** (*flik'-ten'-u-lah*, *flik'-ten'-ul*) [*φλύκταινα*, blister]. A little vesicle or blister.

Phlyctenular (*flik'-ten'-u-lar*) [*φλύκταινα*, blister]. Resembling a phlyctenule; characterized by the formation of phlyctenules, as P. conjunctivitis.

Phlyzacion, **Phlyzacion** (*phi-sa'-se-on*, *phi-sa'-se-um*) [*φλύζειν*, to inflame]. A pustular vesicle on an indurated base. P. acutum. See *Ecthyma*.

Phocomelus (*fo-kom'-el-us*) [*φωκή*, a seal; *μέλος*, limb]. A monster with rudimentary limbs, the hands and feet being attached almost directly to the trunk.

Phonation (*fo-na'-shun*) [*φωνή*, voice]. The production of vocal sound or articulate speech.

Phonatory (*fon'-a-tor-e*) [*φωνή*, voice]. Pertaining to phonation.

Phonautograph (*fo-nauf'-to-graf*) [*φωνή*, voice; *αὐτός*, self; *γράφειν*, to write]. An apparatus for recording automatically the vibrations of the air produced by the voice.

Phonetic (*fo-net'-ik*) [*φωνή*, voice]. 1. Pertaining to or representing sounds. 2. Pertaining to the voice.

Phonetics (*fo-net'-iks*) [*φωνή*, voice]. The science dealing with the mode of production of sounds.

Phonic (*fon'-ik*) [*φωνή*, voice]. Pertaining to the voice. P. Spasm, a spasm of the laryngeal muscles occurring on attempting to speak.

Phonograph (*fo'-no-graf*) [*φωνή*, sound; *γράφειν*, to record]. An instrument consisting of a wax-coated cylinder revolving under a stylus attached to a diaphragm. The vibrations of the diaphragm, set in motion by the voice, cause the cylinder to be indented by the stylus. When the cylinder is again revolved the movement of the stylus upon the cylinder throws the diaphragm into vibration and reproduces the original sounds of the voice.

Phonology (*fo-nol'-o-je*) [*φωνή*, voice; *λόγος*, science]. The science of vocal sounds; phonetics.

Phonometer (*fo-nom'-et-er*) [*φωνή*, voice; *μέτρον*, measure]. An instrument for measuring the intensity of the voice.

Phorometer (*fo-rom'-et-er*) [*φορέειν*, to tend;

μέτρον, measure]. An instrument for measuring the relative strength of the ocular muscles.

Phose (*fōs*) [*φῶς*, light]. A subjective sensation of light or color, as, *e. g.*, scotoma scintillans. **Aphose**, a subjective sensation of shadow or darkness, as, *e. g.*, muscae volitantes. **Centraphoses**, aphoses originating in the optic centers. **Centrophoses**, phoses originating in the optic centers. **Chromophose**, a subjective sensation of color. **Peripheraphoses**, peripheral aphoses. **Peripherophoses**, phoses originating in the peripheral organs of vision (the optic nerve or eyeball).

Phosphate (*fōs'-fāt*) [*φωσφόρος*, phosphorus]. A salt of phosphoric acid. **Normal P.**, one in which the three hydrogen-atoms, or the six of two molecules, are substituted by metals, *e. g.*, Na_3PO_4 , $\text{Ca}_3(\text{PO}_4)_2$. **Acid P.**, one in which one or two of the hydrogen-atoms only have been replaced by metals. **P.**, **Ammoniomagnesium**, a double salt of ammonium and magnesium and phosphoric acid. **P.**, **Earthy**, a phosphate of one of the alkaline earths. **P.**, **Triple**, ammoniomagnesium phosphate. The phosphates are used in medicine as tonics and alteratives in conditions associated with malnutrition of the bones (rickets, scrofula). Sodium phosphate is employed as a cholagogue and laxative.

Phosphatic (*fōs'-fāt'-ik*) [*φωσφόρος*, phosphorus]. Containing phosphates; characterized by the excretion of large amounts of phosphates, as *P. diathesis*.

Phosphatid (*fōs'-fāt'id*) [*φωσφόρος*, phosphorus]. Any one of a large group of phosphorus-compounds, found in brain substance, and resembling the phosphates.

Phosphaturia (*fōs'-fāt'-ur'-ah*) [*φωσφόρος*, phosphorus; *urina*, urine]. A condition in which an excess of phosphates is passed in the urine.

Phosphene (*fōs'-fēn*) [*φῶς*, light; *φαίνω*, to show]. A subjective luminous sensation caused by pressure upon the eyeball.

Phosphid (*fōs'-fid*) [*φωσφόρος*, phosphorus]. A compound of phosphorus and another element or radicle acting as a base. The phosphids are used in medicine as substitutes for phosphorus.

Phosphin (*fōs'-fin*) [*φωσφόρος*, phosphorus]. 1. Hydrogen phosphid, PH_3 , a poisonous gas of alliaceous odor. 2. A substitution-compound of PH_3 , bearing the same relation to it that an amine does to ammonia.

Phosphite (*fōs'-fit*) [*φωσφόρος*, phosphorus]. A salt of phosphorous acid.

Phosphoglyceric Acid (*fōs'-fo-glīs'-ē'-rik*) [*φωσφόρος*, phosphorus; *γλυκύς*, sweet], $\text{C}_3\text{H}_5\text{PO}_6$. A liquid body obtained from lecithin.

Phosphomolybdic Acid (*fōs'-fo-mol'-ib-dik*) [*phosphorus*; *molybdenum*]. A compound of phosphoric acid and molybdenum trioxid, used as a test for alkaloids.

Phosphonium (*fōs'-fō'-ne-um*) [*φωσφόρος*, phosphorus]. The hypothetical univalent radicle PH_4 ; it is analogous to ammonium, NH_4 .

Phosphorated (*fōs'-for'-at-ed*) [*φωσφόρος*, phosphorus]. Containing phosphorus.

Phosphorescence (*fōs'-for'-es'-ens*) [*φωσφόρος*, phosphorus]. The spontaneous luminosity of phosphorus and other substances in the dark.

Phosphorescent (*fōs'-for'-es'-ent*) [*φωσφόρος*, phosphorus]. Possessing the quality of phosphorescence.

Phosphoreted (*fōs'-for'-et-ed*) [*φωσφόρος*, phosphorus]. Combined with phosphorus.

Phosphoric Acid (*fōs'-for'-ik*). See *Acids*, *Table of*.

Phosphoridrosis (*fōs'-for-id-ro'-sis*) [*phosphorus*; *ιδρώς*, sweat]. The secretion of phosphorescent sweat.

Phosphorism (*fōs'-for-izm*) [*phosphorus*]. Chronic phosphorus poisoning.

Phosphorized (*fōs'-for'-izd*) [*phosphorus*]. Containing phosphorus.

Phosphorous Acid (*fōs'-for'-us*). See *Acids*, *Table of*.

Phosphorus (*fōs'-for-us*) [*φῶς*, light; *φορέω*, to bear]. A nonmetallic element, having a quantivalence of 3 or 5, and an atomic weight of 31. Symbol *P*. In commerce it is prepared from bone-ash or from sombrerite, an impure calcium phosphate found in West Indian guano. *P* may be obtained in several allotropic forms. *Ordinary P.* is a yellowish-white, waxy solid, of a specific gravity of 1.837. *Red or amorphous P.* is a dark-red powder, having a specific gravity of 2.11, insoluble in carbon disulphid, nonflammable, nonluminous, nonpoisonous. *Metallic rhombohedral P.* is an allotropic form produced by heating phosphorus in a sealed tube with melted lead. Its specific gravity is 2.34. Medicinally, *P.* is used as an alterative in osteomalacia and in rickets, in sexual impotence, threatened cerebral degeneration, neuralgia, chronic alcoholism, morphinomania, furunculosis, etc. Dose $\frac{1}{12}$ to $\frac{1}{8}$ gr. (0.0065-0.0013). Ordinary *P.* is exceedingly poisonous; it causes a widespread fatty degeneration, most marked in the liver. Preparations: Elixir phosphori (U. S. P.). Dose m xv -[vj] (1.0-4.0). Oleum phosphoratum (U. S. P., B. P.) and Spiritus phosphori (U. S. P.). Dose m j - v (0.065-0.32). Pilulæ phosphori (U. S. P.) contain each gr. $\frac{1}{10}$. **Phosphotungstic Acid** (*fōs'-fo-tung'-stik*) $\text{H}_{11}\text{PW}_{10}\text{O}_{38} + \text{H}_2\text{O}$. A crystalline compound of phosphoric and tungstic acids, used as a test for alkaloids and peptones.

Phosphuret (*fos'-fu-ret*) [φωσφόρος, phosphorus]. A phosphid.

Phosphureted (*fos'-fu-ret-ed*). Synonym of *Phosphureted*.

Photo- (*fo'-to-*) [φῶς, light]. A prefix denoting relation to light.

Photobiotic (*fo-to-bi-ot'-ik*) [φῶς, light; βίος, life]. Living habitually in the light.

Photochemic (*fo-to-kem'-ik*) [φῶς, light; χημεία, chemistry]. Pertaining to the chemic action of light.

Photochemistry (*fo-to-kem'-is-tre*) [φῶς, light; χημεία, chemistry]. That branch of chemistry treating of the chemic action of light.

Photoelectricity (*fo-to-e-lek-tris'-it-e*) [φῶς, light; *electricity*]. Electricity produced under the influence of light.

Photogene (*fo'-to-jen*) [φῶς, light; γεννᾶν, to produce]. 1. A retinal impression; an after-image. 2. A liquid derived from bituminous shale.

Photogenic (*fo-to-jen'-ik*) [φῶς, light; γεννᾶν, to produce]. Light-producing.

Photography (*fo-to-graph'-ra-fe*) [φῶς, light; γράφειν, to write]. The art of producing an image of an object (*Photograph*) by throwing the rays of light reflected from it upon a surface coated with a film of a substance, such as a silver-salt, that is readily decomposed by light, subsequently treating the film with certain agents (*developers*) that bring out the image, and then dissolving the salt unacted upon by the light.

Photohemotachometer (*fo-to-hem-o-tach-om'-et-er*) [φῶς, light; αἷμα, blood; τάχος, swiftness; μέτρον, measure]. A hemotachometer in which the changes in level of the column of blood are photographed.

Photokinetic (*fo-to-kin-et'-ik*) [φῶς, light; κινητικός, causing movement]. Causing movement by means of light.

Photolyte (*fo'-to-lit*) [φῶς, light; λύνειν, to loosen]. A substance that is decomposed by the action of light.

Photomagnetism (*fo-to-mag'-net-izm*) [φῶς, light; *magnet*]. Magnetism produced by the action of light.

Photometer (*fo-tom'-et-er*) [φῶς, light; μέτρον, measure]. An instrument for measuring the intensity of light.

Photometry (*fo-tom'-et-re*) [φῶς, light; μέτρον, measure]. The measurement of the intensity of light.

Photomicrograph (*fo-to-mi'-kro-graf*) [φῶς, light; μικρός, small; γράφειν, to write]. A photograph of a small or microscopic object, usually made with the aid of a microscope, and of sufficient size for observation with the naked eye. Compare *Microphotograph*.

Photomicrography (*fo-to-mi'-krog'-ra-fe*) [φῶς, light; μικρός, small; γράφειν, to write]. The art of producing photomicrographs.

Photophobia (*fo-to-fo'-be-ah*) [φῶς, light; φόβος, fear]. Intolerance of light.

Photophone (*fo'-to-fon*) [φῶς, light; φωνή, sound]. An apparatus for the graphic representation of the character of sound-waves by means of flames.

Photopsia (*fo-top'-se-ah*) [φῶς, light; ὄψις, sight]. Subjective sensations of sparks or flashes of light occurring in certain morbid conditions of the optic nerve, the retina, or the brain.

Photoxylin, Photoxylon (*fo-toks'-il-in, fo-toks'-il-on*) [φῶς, light; ξύλον, wood]. A substance produced from wood-pulp by the action of sulphuric acid and potassium nitrate. It serves as a substitute for collodion in minor surgery, and as a medium for mounting microscopic specimens.

Photuria (*fo-tu'-re-ah*) [φῶς, light; *urina*, urine]. The passage of phosphorescent urine.

Phren (*fren*) [φρήν]. 1. Diaphragm. 2. Mind.

Phrenetic (*fren-et'-ik*) [φρήν, mind]. Maniacal; delirious.

Phrenic (*fren'-ik*) [φρήν, diaphragm, mind]. 1. Pertaining to the diaphragm, as P. nerve, P. artery. 2. Pertaining to the mind.

Phrenitis (*fren-i'-tis*) [φρήν, 1, mind; 2, diaphragm; *itis*, inflammation]. 1. Inflammation of the brain. 2. Inflammation of the diaphragm.

Phrenograph (*fren'-o-graf*) [φρήν, diaphragm; γράφειν, to write]. An instrument for registering the movements of the diaphragm.

Phrenology (*fren-ol'-o-je*) [φρήν, mind; λόγος, science]. The theory that the various faculties of the mind occupy distinct and separate areas in the brain-cortex, and that the predominance of certain faculties can be predicted from modifications of the parts of the skull overlying the areas where these faculties are located.

Phrenopathy (*fren-op'-ath-e*) [φρήν, mind; πάθος, disease]. Mental disease.

Phrenosin (*fren'-o-sin*) [φρήν, mind]. A nitrogenous body obtained from brain-tissue.

Phthalate (*thal'-at*). A salt of phthalic acid.

Phthalic Acid (*thal'-ik*) [from *naphthalene*], C₈H₄O₄. A crystalline substance derived from naphthalene.

Phtheiriasis, Phthiriasis (*thi-ri'-as-is*) [φθειρ, louse]. See *Pediculosis*.

Phthisic (*tiz'-ik*) [φθίσις, a wasting]. 1. Affected with phthisis. 2. A person affected with phthisis.

Phthisical (*tiz'-ik-al*) [φθίσις, a wasting]. Pertaining to or affected with phthisis.

Phthisis (*ti'-sis or te'-sis*) [φθίειν, to waste]. 1. A wasting away or consumption, as P. bulbi, shrinking of the eye ball. 2.

Any disease characterized by emaciation and loss of strength, especially pulmonary tuberculosis. **P.**, Fibroid. **1.** Interstitial pneumonia. **2.** Chronic tuberculosis of the lungs attended with the formation of fibrous tissue, which contracts, causes shrinking of the affected part, and sometimes bronchiectasis by traction on the bronchi. **P. florida**, an acute, rapidly fatal pulmonary tuberculosis; galloping consumption. **P.**, Laryngeal, tuberculosis of the larynx. **P.**, Pulmonary. **1.** Tuberculosis of the lung. **2.** Any one of a variety of interstitial pneumonias, such as Grinders' **P.**, Miners' **P.**, Stone-cutters' **P.**, etc. **P. ventriculi**, atrophy of the mucous membrane and thinning of the coats of the stomach.

Phylaxin (*fi-laks'-in*) [*φύλαξ*, a guardian]. A defensive proteid found in animals that have acquired an artificial immunity to a given infectious disease. The phyllaxins are of two varieties; one having the power to destroy pathogenic microorganisms, called *mycophyllaxin*; one that counteracts the poisons of the microorganisms, called *toxophyllaxin*.

Phyletic (*fi-let'-ik*) [*φύλον*, a tribe]. Pertaining to phylogeny.

Phylogenesis, Phylogeny (*fi-lo-gen'-es-is*, *fi-loj'-en-e*) [*φύλον*, a tribe; *γενᾶν*, to beget]. The evolution of a group or species of animals or plants from the simplest form; the evolution of the species, as distinguished from *ontogeny*, the evolution of the individual.

Phylogenetic (*fi-lo-gen-et'-ik*) [*φύλον*, a tribe; *γενᾶν*, to beget]. Pertaining to phylogeny.

Phyma (*fi'-mah*) [*φύμα*, a growth]. **1.** Formerly, any one of a variety of swellings of the skin. **2.** A localized plastic exudate larger than a tubercle; a circumscribed swelling of the skin.

Phymatosis (*fi-ma-to'-sis*) [*phyma*]. Any disease characterized by the formation of phymata.

Physalis (*fiz'-al-is*) [*φυσάλις*, bladder]. A large giant epithelial cell of giant-cell carcinoma.

Physic (*fiz'-ik*) [*φῆσις*, nature]. **1.** The science of medicine. **2.** A medicine, especially a cathartic. **3.** To administer medicines; also to purge.

Physical (*fiz'-ik-al*) [*φυσικός*, physical]. **1.** Pertaining to nature; also pertaining to the body or material things. **P. Diagnosis**, the investigation of disease by direct aid of the senses, sight, touch, and hearing. **P. Examination**, examination of the patient's body to determine the condition of the various organs and parts. **P. Signs**, the phenomena observed on inspection, palpation, percussion, auscultation, mensuration, or com-

binations of these methods. **2.** Pertaining to physics.

Physician (*fi-izh'-an*) [*φῆσις*, nature]. One who practises medicine.

Physicochemic (*fi-izh'-o-kem'-ik*) [*φῆσις*, nature; *χημεία*, chemistry]. Pertaining to both physics and chemistry.

Physics (*fiz'-iks*) [*φῆσις*, nature]. The science of nature, especially that treating of the properties of matter and of the forces governing it.

Physiognomy (*fi-e-og'-no-mē*) [*φῆσις*, nature; *γνώμη*, knowledge]. **1.** The science treating of the methods of determining character by a study of the face. **2.** The countenance.

Physiologic (*fi-e-o-loj'-ik*) [*φῆσις*, nature; *λόγος*, science]. **1.** Pertaining to physiology. **2.** Pertaining to natural or normal processes, as opposed to those that are pathologic. **P.**

Antidote, an antidote that neutralizes a poison by effects on the system that are antagonistic to those of the poison. **P. Unit**. See *Unit*.

Physiologist (*fi-e-ol'-o-jist*) [*φῆσις*, nature; *λόγος*, science]. One versed in physiology.

Physiology (*fi-e-ol'-o-jē*) [*φῆσις*, nature; *λόγος*, science]. The science that treats of the functions of organic beings. **P.**, Morbid, the study of diseased functions or of functions modified by disease.

Physique (*fi-izh')* [*Fr.*]. Physical structure or organization.

Physocoele (*fi'-so-sel*) [*φῆσις*, air; *κύστη*, tumor]. **1.** A swelling containing air or gas. **2.** Emphysema of the scrotum; a hernia filled with flatus.

Physometra (*fi'-so-mē'-trah*) [*φῆσις*, air; *μήτρα*, uterus]. A distention of the uterus with gas, produced by the decomposition of its contents.

Physostigma (*fi-so-stig'-mah*) [*φῆσις*, air; *στιγμα*, stigma]. Calabar bean; ordeal-nut. The seed of *Physostigma venenosum*, of the natural order Leguminosae, which is used by the natives of Africa as an ordeal-poison. It contains two alkaloids—eserin or physostigmin, and calabarin. It acts as a general depressant, producing motor paralysis and in poisonous doses causing death by paralysis of the respiration. It is a miotic, and in small doses stimulates the heart and intestinal peristalsis. In medicine it is employed as a motor depressant in tetanus and other spasms; as a stimulant in intestinal atony and dilatation, in asthma and emphysema. Preparations: Extractum physostigmatis (U. S. P., B. P.). Dose gr. $\frac{1}{16}$ – $\frac{1}{8}$ (0.008–0.016). Tinctura physostigmatis (U. S. P.). Dose m-v-xx (0.32–1.3).

Physostigmin (*fi-so-stig'-min*) [*physostigma*], $C_{15}H_{21}N_3O_2$. An alkaloid found in

the seed of *Physostigma venenosum*, or Calabar bean. It is also termed eserine. *Physostigminæ salicylas*, eserine salicylate, is used internally in doses of gr. $\frac{1}{16}$ (0.0008); but its chief use is for instillation into the eye as a miotic in conditions of mydriasis, and to lessen intraocular tension in glaucoma. It is used in these conditions in solution of the strength of from one to two grains to the fluid-ounce. *Physostigminæ sulphas*, eserine sulphate, is used in the same manner as the preceding.

Phytalbumose (*fi-tal'-bu-mōs*) [*φυτόν*, plant; *albumose*]. A vegetable albumose.

Phyto- (*fi'-to-*) [*φυτόν*, plant]. A prefix signifying relation to plants.

Phytogenesis (*fi-to-jen'-es-is*) [*φυτόν*, plant; *γένεσις*, origin]. The science of the origin and development of plants.

Phytogenous (*fi-toj'-en-us*) [*φυτόν*, plant; *γεννᾶν*, to produce]. Produced by plants.

Phytolacca (*fi-to-lak'-ah*) [*φυτόν*, plant; *lacca*, lac]. The poke, *P. decandra*, a plant of the natural order *Phytolaccaceæ*, the fruit (*Phytolaccæ fructus*, or poke-berry) and root of which (*Phytolaccæ radix*, or poke-root) are official in the U. S. P. Poke is emetocathartic and slightly narcotic. It has been used in rheumatism, and locally in granular conjunctivitis and parasitic skin-diseases. Dose of the powdered root gr. x-xxx (0.65-2.0). *Extractum phytolaccæ radicis fluidum* (U. S. P.). Dose $\mathfrak{m}\nu$ -xxx (0.32-2.0).

Phytoparasite (*fi-to-par'-as-it*) [*φυτόν*, a plant; *παράσιτος*, a parasite]. A vegetable parasite.

Phytopathology (*fi-to-path-ol'-o-je*) [*φυτόν*, plant; *πάθος*, disease; *λόγος*, science]. 1. The science of diseases of plants. 2. The science of diseases due to vegetable organisms.

Phytoplasm (*fi-to-plasm*) [*φυτόν*, plant; *πλάσμα*, plasma]. Vegetable protoplasm.

Phytosis (*fi-to'-sis*) [*φυτόν*, plant]. Any disease due to vegetable organisms.

Phytosterin (*fi-tos'-ter-in*) [*φυτόν*, plant; *στέαρ*, fat]. A fat-like substance, similar to cholesterol, present in plant-seeds and sprouts.

Pia, Pia Mater (*pi-a mah'-ter*) [L., kind or tender mother]. The vascular membrane enveloping the surface of the brain and spinal cord, and consisting of a plexus of blood-vessels held in a fine areolar tissue.

Pia-arachnitis (*pi-ah-ar-ak-ni'-tis*) [*pia*, pia; *αράχνη*, spider's web; *ιτις*, inflammation]. Inflammation of the pia-arachnoid; leptomeningitis.

Pia-arachnoid (*pi-ah-ar-ak'-noid*) [*pia*, pia; *αράχνη*, a spider's web; *ειδος*, like]. The pia and arachnoid considered as one structure.

Pial (*pi'-al*) [*pia*, pia]. Pertaining to the pia mater.

Pian (*pi'-an*). See *Frambesia*.

Piano-players' Cramp. A painful spasm of the muscles occurring in piano-players as the result of overuse of the muscles in playing; a form of occupation-neurosis.

Pica (*pi'-kah*) [L., magpie]. A craving for unnatural and strange articles of food; a symptom present in certain forms of insanity, hysteria, and chlorosis, and during pregnancy.

Picea (*pis'-e-ah*) [L., the pitch-pine]. A genus of coniferous trees. *P. alba*, the white spruce. *P. excelsa*, the common fir or pitch-pine. It yields resin and turpentine. *P. nigra*, the black spruce. *P. vulgaris*. Synonym of *P. excelsa*.

Piceous (*pis'-e-us*) [*pix*, pitch]. Resembling pitch.

Picolin (*pik'-o-lin*) [*picea*, the pitch-pine; *oleum*, oil], C_8H_7N . Methyl-pyridine, a liquid obtained by distillation from coal-tar.

Picramic Acid (*pik'-ram'-ik*) [*πικρός*, bitter; *amin*], $C_8H_5N_3O_6$. Picric acid in which one NO₂ radicle has been replaced by NH₂.

Picrate (*pik'-rat*) [*πικρός*, bitter]. A salt of picric acid.

Picric Acid (*pik'-rik*) [*πικρός*, bitter]. See *Acid*, *Picric*.

Picrocarmin (*pik-ro-kar'-min*) [*πικρός*, bitter; *carmin*]. A preparation for staining microscopic specimens. Its composition is as follows: carmin 1.0; ammonia 5.0; distilled water 50.0 parts. After solution, 50 parts of a saturated watery solution of picric acid are added, and the mixture allowed to stand in a wide-mouthed bottle until the ammonia has evaporated. It is then filtered.

Picrotoxin (*pik-ro-toks'-in*) [*πικρός*, bitter; *τοξικόν*, poison]. A bitter neutral principle prepared from *Anamirta paniculata* (*Cocculus indicus*). Picrotoxin stimulates the motor and inhibitory centers in the medulla, especially the respiratory and vagus centers; it causes epileptiform spasms by irritation of the motor centers of the cerebrum or cord. Its action is much like that of strychnine. It has been used in an ointment (gr. x to $\frac{3}{4}$) in tinea capitis and in pediculosis. It is useful for the night-sweats of phthisis and in the complex of symptoms known as vasomotor ataxia. Dose gr. $\frac{1}{16}$ to $\frac{1}{8}$ (0.001-0.003).

Piebald Skin. See *Leukoderma* and *Vitiligo*.

Piedra (*pe-a'-drah*) [S. Am.]. A disease of the hair marked by the formation of hard, pin-head-sized nodules on the shaft of the hair; it is thought to be due to a micrococcus.

Piezometer (*pi-e-zom'-et-er*) [*πιέζειν*, to press; *μέτρον*, measure]. An apparatus for measuring the degree of compression of gases.

Pigment (*pig'-ment*) [*pingere*, to paint]. A

dye-stuff; a coloring matter. Pigments may be in solution or in the form of granules or crystals. **P.**, **Hematogenous**, any pigment derived from the blood. Hematogenous pigments are hemoglobin, hematoïdin, hemosiderin, and the bile-pigments, which are indirectly derived from the blood-pigment. **P.**, **Metabolic**, a pigment formed by the metabolic action of cells. Melanin is the type of metabolic pigments.

Pigmentary (*pij'-men-ta-re*) [*pijere*, to paint]. Pertaining to or containing pigment; characterized by the formation of pigment.

Pigmentation (*pij-men-ta'-shun*) [*pijere*, to paint]. Deposition of or discoloration by pigment.

Pilary (*pi'-lar-e*) [*pilus*, hair]. Pertaining to the hair.

Pilastered (*pi-las'-terd*) [*pila*, pillar]. Flanged so as to have a fluted appearance; arranged in pilasters or columns. **P. Femur**, a condition of the femur in which the backward concavity of the shaft is exaggerated and the linea aspera prominent.

Piles (*pils*) [*pila*, ball]. See *Hemorrhoids*.

Pill [*pilula*, dim. of *pila*, ball]. A small, round mass containing one or more medicinal substances and used for internal administration. **P.**, **Blaud's**. See *Ferrum*. **P.**, **Blue**. See *Mercury*. **P.**, **Compound Cathartic**. See *Colocynthis*. **P.**, **Griffith's**. Synonym of *P.*, *Blaud's*. **P.**, **Lady Webster's**, pill of aloes and mastic. **P.-mass**, a cohesive mass used to hold together the ingredients of a pill.

Pillar (*pi'-lar*) [*pila*, a pillar]. A columnar structure acting as a support. **P.** of the **Abdominal Ring**, one of the columns on either side of the abdominal ring. **P.** of the **Fauces**, one of the folds of mucous membrane on either side of the fauces.

Pilocarpin (*pi-lo-kar'-pin*). See *Pilocarpus*.

Pilocarpus (*pi-lo-kar'-pus*) [*pilus*, hair; *καρπός*, fruit]. *Jaborandi*; a South American shrub of the natural order Rutaceæ, the leaves of which yield two alkaloids, pilocarpin, $C_{11}H_{16}N_2O_2$, and jaborin, $C_{22}H_{33}N_4O_4$, which resembles atropin in action. *Jaborandi* and the alkaloid pilocarpin taken internally produce salivation, perspiration, and contraction of the pupil. They are employed as diaphoretics in dropsy, Bright's disease, uremia, in rheumatism, and in the early stage of cold. Pilocarpin is used locally as a miotic. Dose of *jaborandi* gr. xx-lx (1.3-4.0). Extractum *jaborandi* (R. P.) gr. ij-x (0.13-0.65). Extractum pilocarpi fluidum (U. S. P.) m xl-lx (2.6-4.0). Infusum *jaborandi* (B. P.). Dose f 3-ij (32.0-64.0). Pilocarpin hydrochlorate (Pilocarpinæ hydrochloras U. S. P.). Dose gr. 1 (0.008). Pilocarpin nitrate (Pilocarpinæ nitras B. P.). Dose gr. 1 (0.008).

Pilomotor (*pi-lo-mo'-tor*) [*pilus*, hair; *movere*, to move]. Causing movement of the hair. **P. Nerves**, nerves causing contraction of the *erectores pilorum*. **P. Reflex**, the appearance of "goose-skin" when the skin is irritated.

Pilonidal (*pi-lo-ni'-dal*) [*pilus*, hair; *nidus*, nest]. Containing an accumulation of hairs in a cyst. **P. Fistula**, a fistula in the neighborhood of the rectum depending upon the presence of a tuft of hair in the tissues.

Pilose, **Pilous** (*pi'-lôs*, *pi'-lus*) [*pilosus*, hairy]. Hairy.

Pilosis (*pi-lo'-sis*) [*pilus*, hair]. The abnormal or excessive development of hair.

Pilula (*pi'-lu-lah*) [L.]. A pill.

Pilular (*pi'-lu-lar*) [*pilula*, dim. of *pila*, ball]. Of the nature of, or pertaining to pills.

Pilule (*pi'-ul*) [*pilula*, a small pill]. A small pill.

Pimenta (*pi-men'-tah*) [Sp., *pimiento*]. Allspice, the nearly ripe fruit of *P. officinalis*, a tree of the natural order Myrtaceæ. It has a fragrant aromatic odor, due to the presence of a volatile oil, *Oleum pimentæ*. **P.** is used as an aromatic carminative in flatulence and locally in chilblains. Dose gr. x-xi (0.65-2.6). *Oleum pimentæ*. Dose m ij-v. (0.13-0.32).

Pimpinella (*pin-pin-el'-ah*). A genus of umbelliferous plants. **P. anisum**, yields anise. **P. saxifraga**, is said to be diaphoretic, diuretic, and stomachic, and has been employed in asthma, dropsy, amenorrhœa, etc. Dose 3 ss (2.0).

Pimple (*pin'-pl*) [AS., *pipel*, a pimple]. A small pustule or papule.

Pincet, **Pincette** (*pin-set'*) [Fr.]. A small forceps.

Pine (*pin*) [*pinus*]. A genus of trees of the order Coniferae, yielding turpentine, pitch, tar, and other substances.

Pineal (*pi'-ne-al*) [*pinus*, a pine-cone]. Belonging to or shaped like a pine-cone.

P. **Body**, **P. Gland**, a small, reddish-gray, vascular body situated behind the third ventricle, which is embraced by its two peduncles; it is also called the conarium, from its conic shape. **P. Eye**, a rudimentary third, median, or unpaired eye of certain lizards, with which the pineal body of the mammalia is homologous.

Pinguecula, **Pinguicula** (*pin-gwek'-u-lah*, *pin-gwik'-u-lah*) [dim. of *pinguis*, fat]. A small, yellowish-white patch situated on the conjunctiva, between the cornea and the canthus of the eye; it is composed of connective tissue.

Piniform (*pi'-nif-orm*) [*pinus*, pine; *forma*, form]. Shaped like a pine-cone.

Pink-eye. A contagious mucopurulent conjunctivitis occurring especially in horses.

Pink-root. See *Spigelia*.

Pinna (*pin'-ah*) [L.]. The projecting part of the external ear; the auricle.

Pint (*pin*) [Sp., *pinta*, a spot, from L., *pingere*, to paint]. The eighth part of a gallon; octarius. Symbol O.

Pinta Disease [Sp., spot]. Spotted sickness; a tropical contagious disease characterized by scaly spots variable in color, shape, and size. The disease usually begins on the face and extremities, and is probably due to a parasite.

Pinus (*pi'-nus*). See *Pine*.

Pip. A contagious disease of fowls characterized by a secretion of thick mucus in the throat and mouth.

Piper (*pi'-per*) [L.]. See *Pepper*.

Pin-worm. See *Oxyuris*.

Piperazin (*pi-per-a'-zin*), $C_4H_{10}N_2$. Diethylendiamin, a crystalline substance produced by the action of ammonia on ethylene bromid or chlorid. It is readily soluble in water. In watery solutions it acts as an excellent solvent of uric acid. It is used internally in cases of gout, lithemia, diabetes, and as a solvent for uric acid calculi. Dose gr. xv a day; for hypodermic use it is best employed in a two per cent. solution.

Piperic (*pi-per'-ik*) [*piper*, pepper]. Pertaining to or containing pepper. **P. Acid**, $C_{12}H_{10}O_4$, a monobasic acid obtained by decomposing piperin.

Piperidin (*pi-per'-id-in*) [*piper*, pepper], $C_5H_{11}N$. A liquid base produced in the decomposition of piperin.

Piperin (*pi'-per-in*). See *Pepper*.

Pipet, Pipette (*pip-et'*) [Fr., dim. of *pipe*]. A glass tube open at both ends, but usually drawn out to a smaller size at one end. It is used for transferring small portions of a liquid from one vessel to another.

Pipsissewa (*pip-sis'-e-wah*). See *Chimaphila*.

Pirogoff's Operation. A method of amputation at the ankle, in which the greater part of the calcaneum is retained to give length and surface to the stump. See *Operations*, Table of.

Piscidia erythrina (*pis-id'-e-ah er-e-thri'-nah*) [*piscis*, fish; *cadere*, to kill]. Jamaica dogwood, a tree of the order Leguminosæ, the bark of which has been used for stupefying fish. It contains a neutral principle, piscidin. P. has been used as an anodyne in neuralgia, whooping-cough, and in insomnia. Dose of the fluid extract f ʒj (4.0).

Piscidin (*pis-id'-in*). See *Piscidia erythrina*.

Pisiform (*pis'-if-orm*) [*pisum*, a pea; *forma*, form]. Pea-shaped. **P. Bone**, a small bone on the inner and anterior aspect of the carpus.

Pit [AS., *pyt*, from *putus*, a well or pit]. 1. A depression, as the pit of the stomach, the armpit. 2. To indent by pressing; to become indented on pressure.

Pitch [ME., *picchen*, to throw]. The height of a sound; that quality which depends upon the relative rapidity of the vibrations that produce the sound.

Pitch [AS., *pic*, from *pix*, pitch]. A hard, but viscous, shining substance, breaking with a conchoidal fracture, obtained from various species of pine and from tar. **P.**, Burgundy, *Pix burgundica* (U. S. P., B. P.) is the prepared resinous exudation of *Abies excelsa*, and is used in the form of plaster as a counterirritant in chronic rheumatism. Preparations: *Emplastrum picis burgundicæ* (U. S. P.) and *Emplastrum picis* (B. P.); *Emplastrum picis cantharidatum* (U. S. P.); *Emplastrum calcfaciens* (B. P.), warming plaster. **P.**, Canada (*Pix canadensis*), is obtained from *Abies canadense*, and was formerly used for making plasters. **P.**, Liquid, *Pix liquida* (U. S. P., B. P.). See *Tar*.

Pith [AS., *piṭṭa*, pith]. 1. The soft cellular tissue found in the center of the stalks of plants. 2. To cut off all connection of the brain-centers of an animal with the periphery by piercing the brain and spinal cord.

Pituita (*pit-u'-it-ah*) [L.]. Phlegm; mucus.

Pituitary (*pit-u'-it-a-re*) [*pituita*, phlegm]. Secreting or containing mucus. **P. Body**, a small, reddish-gray, vascular body, weighing about ten grains, contained within the sella turcica of the skull. It consists of two portions, the large anterior or oral, and the small posterior or cerebral division. The anterior lobe is derived as a diverticulum from the primitive oral cavity; the posterior lobe descends as an outgrowth from the brain, communicating in fetal life with the third ventricle. The stalk of this outgrowth remains as the infundibulum. The pituitary body has attracted much attention on account of pathologic changes in its structure in certain obscure diseases, such as acromegaly, myxedema, and others. In some cases of the affection first named, it has been much enlarged. It is also called hypophysis cerebri. **P. Membrane**, the Schneiderian membrane.

Pituitous (*pit-u'-it-us*) [*pituita*, phlegm]. Containing or resembling mucus.

Pityriasis (*pit-ir-i'-as-is*) [*πύριον*, bran].

1. A term applied to various skin-affections characterized by fine, branny desquamation. 2. *Seborrhœa sicca*. **P. capillitii**, **P. capitis**, alopecia furfuracea. **P. circinata et marginata**, a disease characterized by an eruption of rose-colored spots on the trunk, the limbs, and in the axillæ, associated with slight fever and itching. **P. pilaris**. See *Kerato-*

sis pilaris. **P. rosea**. See *P. maculata et circumata*. **P. rubra**, dermatitis exfoliativa, a chronic inflammatory skin-disease, beginning in one or more localized patches, which coalesce and gradually invade the whole body. The skin is deep-red in color, and covered by whitish vesicles that constantly reform. The disease lasts months or years, and generally ends fatally. **P. versicolor**. See *Tinea versicolor*.

Pityroid (*pit'-ir-oid*) [*πίτυρον*, bran; *είδος*, like]. Branny.

Pix (*piks*) [L.]. Pitch. **P. liquida**. See *Tar*.

Placebo (*pla-se'-bo*) [L., I will please, from *placere*, to please]. A medicine given for the purpose of pleasing or humoring the patient, rather than for its therapeutic effect.

Placenta (*pla-sen'-tah*) [*πλάκους*, a cake]. The organ on the wall of the uterus, to which the embryo is attached by means of the umbilical cord and from which it receives its nourishment. It is developed about the third month of gestation, from the chorion of the embryo and the decidua serotina of the uterus. The villi of the chorion enlarge and are received into depressions of the decidua, and around them blood-sinuses form, into which, by diffusion, the waste-materials brought from the fetus by the umbilical arteries pass, and from which the blood receives oxygen and food-material, being returned to the fetus by the umbilical vein. At term the placenta weighs one pound, is one inch thick at its center, and seven inches in diameter. **P.**, **Adherent**, one that is abnormally adherent to the uterine wall after childbirth. **P.**, **Battledore**, one in which the insertion of the cord is at the margin of the placenta. **P. prævia**, a placenta that is fixed to that part of the uterine wall that becomes stretched as labor advances, so that it precedes the advance of the presenting part of the fetus. Being detached before the birth of the child, it generally causes grave hemorrhage. **P.**, **Retained**, one not expelled by the uterus after labor. **P. succenturiata**, an accessory placenta.

Placental (*pla-sen'-tal*) [*placenta*]. Pertaining to the placenta. **P. Murmur**, **P. Souffle**, a sound attributed to the circulation of blood in the placenta.

Placentation (*pla-sen'-tah-shun*) [*placenta*]. The formation and mode of attachment of the placenta.

Placentitis (*pla-sen-ti'-tis*) [*placenta*; *itis*, inflammation]. Inflammation of the placenta.

Plagioccephalic (*pla-je-o-sef'-al'-ik*) [*πλάγιος*, oblique, twisted; *κεφαλή*, head]. Having a skull exhibiting plagioccephaly.

Plagioccephaly (*pla-je-o-sef'-al'-e*) [*πλάγιος*, oblique; *κεφαλή*, head]. A malformation of

the head, produced by the closing of half of the coronal suture, giving an oblique growth to the cranial roof.

Plague (*plag*) [*πληγή*, a stroke]. A contagious disease at present endemic in Eastern Asia, but in former times occurring epidemically in Europe and Asia Minor. After a period of incubation of from three to eight days, the disease begins with fever, pain, and swelling of the lymphatic glands, chiefly the femoral, inguinal, axillary, and cervical. Headache, delirium, vomiting, and diarrhea may be present. In cases that are going on to recovery the temperature usually falls in about a week. The cause of the disease is believed to be a bacillus found by Kitasato in the blood, buboes, and internal organs of the victims of the plague.

Planta (*plan'-tah*) [L.]. The sole of the foot.

Plantar (*plan'-tar*) [*planta*, sole of the foot]. Pertaining to the sole of the foot. **P. Arch**. See *Arch*. **P. Fascia**, the dense triangular shaped aponeurosis occupying the middle and sides of the sole of the foot beneath the integument. **P. Reflex**. See *Reflexes*, *Table of*.

Plantaris (*plan-ta'-ris*) [*planta*, sole of the foot]. See *Muscles*, *Table of*.

Plantigrade (*plan'-te-grād*) [*planta*, sole; *gradī*, to walk]. Bringing the entire length of the sole of the foot to the ground in walking, as is seen in the bear.

Plaque (*plak*) [Fr.]. A patch. **P.**, **Blood-**. See *Blood-plaque*.

Plasma (*plaz'-mah*) [*πλάσμα*, a thing molded, from *πλάσσειν*, to mold]. 1. The fluid part of the blood and the lymph. See *Blood-plasma*. **P.-cells**, large, granular cells found in the connective tissue. **P.**, **Lymph-**, the fluid part of the lymph. **P.**, **Muscle-**. See *Muscle*. 2. Glycerite of starch.

Plasmatic (*plaz-mat'-ik*) [*plasma*]. Pertaining to plasma. **P. Layer**, the layer of plasma next to the wall of a capillary.

Plasmin (*plaz'-min*) [*plasma*]. A name given to the precipitate obtained from blood by treating it with a saturated solution of sodium sulphate, allowing the corpuscles to subside, then precipitating the plasma with sodium chlorid, and washing the precipitate with a saturated solution of sodium chlorid.

Plasmodium (*plaz-mo'-de-um*) [*πλάσμα*, a thing molded; *είδος*, form]. The mass of protoplasm formed by the fusion of two or more amebiform bodies. **P. malariae**, a protozoon parasite found in the blood of persons suffering from malaria. Many different forms have been observed, which by some are held to be stages in the life-history of the same organism, by others to be different species. The development and multipli-

cation of the organism take place within the red corpuscles. In intermittent fever the small intracorporeal body enlarges and gradually fills the entire cell, becoming pigmented at the same time. The pigment-granules are formed from the hemoglobin of the blood corpuscle. When the plasmodia have acquired a certain size, the pigment-granules begin to aggregate at the center of the body. At this time, in the quartan form, the organism becomes radially striate, or roset-shaped. The further step consists in the breaking up of the organisms into a number of small spheric bodies, the young plasmodia. The blood-corpuscle is destroyed, and the small, free elements enter other red cells. Peculiar crescentic bodies, intraglobular and extraglobular, have also been described. They may occur in acute forms, but are most constant in malarial cachexia. The significance of certain flagellate bodies has not been fully determined.

Plasmogen (*plas'-mo-jen*) [*πλάσμα*, a thing molded; *γεννάν*, to produce]. Formative protoplasm, germ-plasm, bioplasm.

Plasson (*plas'-son*) [*πλάσσειν*, to form or mold]. Primitive or undifferentiated protoplasm; the protoplasm of the cell in the non-nucleated or cytode stage.

Plaster (*plas'-ter*) [*emplastum*, from *ἐν*, in; *πλάσσειν*, to mold]. 1. An adhesive, semi-solid substance spread upon cloth or other flexible material for application to the surface of the body. *P.*, Adhesive, resin-plaster. *P.*, Court-, a plaster prepared by spreading a mixture of isinglass upon silk. *P.*, Diachylon-. See *Plumbum*. *P.*-mull, a plaster made by incorporating with mull or thin muslin a mixture of gutta-percha and some medicament dissolved in benzoin. It is used in skin-diseases. *P.*, Mustard-, one made by spreading upon muslin powdered mustard, or a mixture of mustard and flour reduced to the consistency of paste by the addition of water. *P.*, Spice-, a plaster composed of yellow wax, suet, turpentine, oil of nutmeg, olibanum, benzoin, oil of peppermint, and oil of cloves, and used to relieve abdominal pain in children. *P.*, Warming, a plaster of pitch and cantharides, the *Emplastum picis cum cantharide*. 2. Plaster of Paris, a mixture of calcium sulphate (gypsum) and water, having the property of becoming hard during drying. It is used for surrounding parts, such as joints, fractured limbs, etc., with a stiff casing, to prevent mobility.

Plastic (*plas'-tik*) [*πλάσσειν*, to mold]. 1. Formative; building up tissues; repairing defects, as *P.* surgery, *P.* operation. 2. Capable of being molded. *P.* Lymph, the inflammatory exudate that covers wounds or inflamed serous surfaces, and which becomes

organized by the development in it of blood-vessels and connective tissues.

Plasticity (*plas-tis'-it-ē*) [*πλάσσειν*, to mold]. 1. Plastic force. 2. The quality of being plastic.

Plastid (*plas'-tid*) [*πλάσσειν*, to mold]. An elementary organism; a cell or cytode.

Plastidule (*plas'-tid-ul*) [dim of *plastid*, from *πλάσσειν*, to mold]. A protoplasmic molecule; one of the physical units of which living matter is composed.

Plastin (*plas'-tin*) [*πλάσσειν*, to mold]. A phosphorized proteid, constituting the chief proteid of protoplasm.

Plate (*plat*) [*πλᾶτις*, broad]. A flattened part, especially a flattened process of bone, as the cribriform *P.* of the ethmoid bone, the orbital *P.* of the frontal bone, the tympanic *P.* of the temporal bone. *P.*, Approximation-, one of the plates of decalcified bone or other material that are used in enterectomy to bring the resected ends of intestine together. *P.*-culture, a method of obtaining pure cultures of bacteria by pouring the inoculated culture-medium upon sterile glass-plates and allowing it to solidify. *P.*, Dorsal, one of the two longitudinal ridges on the dorsal surface of the embryo which subsequently join to form the neural canal. *P.*, Frontal, in the fetus, a cartilaginous plate interposed between the lateral parts of the ethmoid cartilage and the lesser wings and anterior portion of the sphenoid bone. *P.*, Frontonasal, the middle of the facial plates, which subsequently forms the external nose. *P.*, Lateral Mesoblastic, the thick portion of the mesoblast situated one on each side of the notochord. Each plate splits into two portions, the outer divisions coalescing to form the body-wall, or somatopleure, the inner, to form the splanchnopleure, or visceral covering.

Platinic (*plat-in'-ik*) [*platinum*]. Containing platinum as a quadrivalent element.

Platinode (*plat'-in-ōd*) [*platinum*; *ὁδός*, way]. The collecting plate of an electric battery, so called because formerly often made of platinum.

Platinous (*plat'-in-us*) [*platinum*]. Containing platinum as a bivalent element.

Platinum (*plat'-in-um*) [*Sp.*, *platina*, dim. of *plata*, silver]. A silver-white metal occurring native or alloyed with other metals; atomic weight 195; sp. gr. 21.5; quantivalence 11 and 14; symbol Pt. It is fusible only at very high temperatures, and is insoluble in all acids except aqua regia. On account of these properties it is extensively used for chemic apparatus—crucibles, foils, wire, etc.; it is also employed as a reagent. *P.* occurs, aside from its ordinary metallic form, as a spongy mass (spongy *P.*) and as a fine

metallic powder (P., black), which is capable of condensing a great deal of oxygen, and hence acts as a powerful oxidizing agent. P. forms two sets of compounds, a platinum series, in which it acts as a dyad, and a platinic series, in which it acts as a tetrad. Platonic chlorid, PtCl₄, is used as a reagent to detect potassium and ammonium; also in syphilis, in doses of gr. $\frac{1}{4}$ – $\frac{1}{2}$ (0.008–0.03).

Platy- (*plat'-e*) [*πλάτης*, broad]. A prefix signifying broad.

Platycephalic, **Platycephalus** (*plat-is-ef-al'-ik*, *plat-is-ef'-al-us*) [*πλάτης*, wide; *κεφαλή*, head]. Having a broad skull with a vertical index of less than 70.

Platycnemía (*plat-e-kne'-me-ah*) [*πλάτης*, broad; *κνήμη*, leg]. The state of being platycnemic.

Platycnemic (*plat-e-kne'-mik*) [*πλάτης*, broad; *κνήμη*, leg]. Having a tibia which is exaggerated in breadth; broad-legged.

PlatyhERIC (*plat-e-hi-er'-ic*) [*πλάτης*, broad; *ιερός*, holy; sacrum]. Having a broad sacrum; having a sacral index of more than 100.

Platypellic (*plat-e-pel'-ik*) [*πλάτης*, wide; *πέλλα*, basin]. Having a broad pelvis.

Platypodia (*plat-e-pod'-de-ah*) [*πλάτης*, broad; *πούς*, foot]. Flat-footedness.

Platyrrhine (*plat'-ir-in*) [*πλάτης*, broad; *ῥίς*, nose]. Having a broad and flat nose; having a nasal index above 53.

Platysma myoides (*plat-is'-mah mi-oi'-des*) [*πλάτης*, broad]. See *Muscles*, Table of.

Playfair's Treatment. See *Rest-cure*.

Pledget (*piej'-et*) [origin obscure, perhaps dim. of *plug*]. A small flattened compress.

Pleochroic (*ple-o-kro'-ik*). See *Pleochromatic*.

Pleochroism (*ple-oh'-ro-izm*) [*πλέων*, more; *χρῶμα*, color]. The property possessed by some bodies, especially crystals, of presenting different colors when viewed in the direction of different axes.

Pleochromatic (*ple-o-kro-mat'-ik*) [*πλέων*, more; *χρῶμα*, color]. Pertaining to or exhibiting pleochroism.

Pleomorphic (*ple-o-mor'-fik*) [*πλέων*, more; *μορφή*, form]. Having more than one form.

Pleomorphism (*ple-o-mor'-fiz-m*) [*πλέων*, more; *μορφή*, form]. The state of being pleomorphic, *i. e.*, of existing in widely different forms.

Plesiomorphic (*ple-se-o-mor'-fus*) [*πλησιος*, near; *μορφή*, form]. Crystallizing in similar forms but differing in chemic composition.

Plessimeter (*ples-im'-et-er*). See *Pleximeter*.

Plessor (*ples'-or*). See *Plexor*.

Plethora (*pleth'-o-rah*) [*πληθώρα*, *πλήθος*, fulness]. A state characterized by an excess of blood in the vessels, and marked by

reddish color of the face, a full pulse, a feeling of fulness and tension in the head, drowsiness, and a tendency to nose-bleed.

Plethoric (*pleth'-o-rík*) [*πληθός*, fulness]. Pertaining to or characterized by plethora.

Plethysmograph (*pleth-is'-mo-graf*) [*πληθισμός*, increase; *γράφειν*, to write]. An instrument for ascertaining changes in the volume of an organ or part, dependent upon changes in the quantity of the blood.

Pleura (*plu'-rah*) [*πλευρά*, rib]. The serous membrane enveloping the lung (pulmonary P., visceral P.), and which, being reflected back, lines the inner surface of the thorax (costal P., parietal P.). P., **Diaphragmatic**, the reflection of the pleura upon the upper surface of the diaphragm.

Pleural (*plu'-ral*) [*πλευρά*, rib]. Pertaining to the pleura.

Pleuralgia (*plu-ral'-je-ah*) [*πλευρά*, rib; *ἄλγος*, pain]. Intercostal neuralgia.

Pleurapophysis (*plu-rap-off'-is-is*) [*πλευρά*, rib; *ἀπόφύσις*, process]. One of the lateral processes of a vertebra, having the morphologic valence of a rib.

Pleurisy (*plu'-ris-i*) [*πλευρά*, rib]. Pleuritis; inflammation of the pleura. It may be acute or chronic. Three chief varieties are usually described, depending upon the character of the exudate: (a) Fibrinous or plastic; (b) Serofibrinous; (c) Purulent. In fibrinous P., the pleura is covered with a layer of lymph of variable thickness, which, in the acute form, can be readily stripped off. Serofibrinous P., is characterized by the presence of a considerable quantity of fluid containing flocculi of lymph, and the deposit of some fibrin on the pleural surface. Purulent P., or empyema, is characterized by the presence of a purulent exudate. Acute pleurisy is marked by sharp and stabbing pain (stitch) in the side, increased by breathing and coughing; by fever, and by a friction-fremitus felt on palpation and a to-and-fro friction-sound heard on auscultation. In the serofibrinous variety a liquid effusion takes place, the signs of which are: bulging of the intercostal spaces and chest-wall, absence of vocal fremitus, displacement of the heart, movable dulness with a curved upper line, and a tympanitic percussion-note (Skodaic resonance) beneath the clavicle and above the level of the effusion. Chronic P. may be dry or serofibrinous. P., **Encysted**, pleurisy in which the effusion is circumscribed by adhesions. P., **Hemorrhagic**, a variety in which the exudate contains blood. P., **Mediastinal**, inflammation of the pleural layers about the mediastinum. P., **Metapneumonic**, pleurisy dependent upon a pneumonia.

Pleuritic (*plu-ris'-ik*) [*πλευρά*, rib; *ιτις*, in-

flammation]. Pertaining to, affected with, or of the nature of pleurisy.

Pleuritis (*plu-ri'-tis*) [*πλευρά*, rib; *ιτις*, inflammation]. See *Pleurisy*.

Pleuro- (*plu'-ro-*) [*πλευρά*, rib]. A prefix denoting connection with the pleura or with a side.

Pleurodynia (*plu-ro-din'-e-ah*) [*πλευρά*, rib; *δύνη*, pain]. A sharp pain in the intercostal muscles, of rheumatic origin.

Pleurogenic, Pleurogenous (*plu-ro-jen'-ik*, *plu-roj'-en-us*) [*πλευρά*, rib; *γεννᾶν*, to produce]. Originating in the pleura.

Pleuroperitoneal (*plu-ro-per-e-ton-e'-al*) [*πλευρά*, rib; *περιτόναιον*, peritoneum]. Pertaining to the pleura and the peritoneum.

Pleuropneumonia (*plu-ro-nu-mo'-ne-ah*) [*πλευρά*, rib; *πνεῦμον*, lung]. Combined inflammation of the pleura and the lung, especially a contagious variety occurring in cattle.

Pleurosoma (*plu-ro-so'-mah*) [*πλευρά*, rib; *σῶμα*, a body]. A variety of monsters of the species *celosoma*, in which there is a lateral evagination with atrophy or imperfect development of the upper extremity on the side of the evagination.

Pleurothotonos (*plu-ro-thot'-o-nos*). See *Pleurothotonos*.

Pleurothotonos (*plu-ro-thot'-o-nos*) [*πλευροθθεν*, from the side; *τείνειν*, to bend]. A form of tetanic spasm of the muscles in which the body is bent to one side.

Pleurotomy (*plu-rot'-o-me*) [*πλευρά*, rib; *τομή*, a cutting]. Incision into the pleura.

Plexiform (*pleks'-if-orm*) [*plexus*; *forma*, form]. Resembling a network or plexus.

Pleximeter (*pleks-im'-et-er*) [*πληξίς*, a stroke; *μέτρον*, measure]. A disc placed on the body to receive the stroke in mediate percussion.

Pleximetric (*pleks-e-met'-rik*) [*πληξίς*, a stroke; *μέτρον*, measure]. Pertaining to or performed with a pleximeter.

Plexor (*pleks'-or*) [*πληξίς*, stroke]. A hammer used for performing percussion.

Plexus (*pleks'-us*) [*plectere*, to knit]. A network, especially an aggregation of vessels or nerves forming an intricate network. **P.**, **Brachial**, a plexus formed in the neck by the union of the anterior branches of the lower four cervical and the greater part of the first dorsal nerves. Its branches are the rhomboid, subclavian, suprascapular, external anterior thoracic, musculocutaneous, subscapular, median, musculospiral, posterior thoracic, internal anterior thoracic, internal cutaneous, lesser internal cutaneous, and ulnar nerves. **P.**, **Cervical**, a plexus in the neck formed by the anterior branches of the upper four cervical nerves. Its branches are the *superficial*, to skin of the head and neck;

and the *deep*, the phrenic, communicans noni, two muscular, and two communicating branches. **P.**, **Choroid**. See *Choroid*. **P.**, **Pampiniform**, the spermatic plexus, a plexus of veins collecting the blood from the testicle.

Plica (*pli'-kah*) [L.]. A fold. **Plicæ palmaræ**, radiating folds in the mucous membrane of the cervix. **P. polonica**, Polish plait; a matted, entangled condition of the hair, due to want of cleanliness in certain diseases of the scalp. **P. semilunaris**, a conjunctival fold in the inner canthus of the eye, the rudiment of the *membrana nictitans* of birds.

Plicate (*pli'-kat*) [*plicare*, to fold]. Folded, plaited.

Plumbic (*plum'-bik*) [*plumbum*, lead]. Pertaining to or containing lead.

Plumbism (*plum'-bism*) [*plumbum*, lead]. Lead-poisoning.

Plumbum (*plum'-bum*) [L.]. Lead, a bluish-white metal occurring in nature chiefly as the sulphid, PbS , known as galena; atomic weight 207; sp. gr. 11.38; quantivalence II and IV; symbol Pb . The salts of lead are poisonous, producing, in sufficient doses, gastroenteritis; ingested in small quantities over a long period of time chronic lead-poisoning is produced. See *Lead-poisoning*. **Plumbi acetat**, lead acetate, $Pb(C_2H_3O_2)_2 \cdot 3H_2O$; sugar of lead. It is used as an astringent in diarrhea and dysentery, as a hemostatic, and as an astringent and sedative in gonorrhea, leukorrhea, conjunctivitis, etc. Dose gr. j-ij (0.065-0.20). From it are prepared: **Liquor plumbi subacetatis** (U. S. P.) (see *Plumbi subacetat*); **Unguentum plumbi acetatis** (B. P.); **Suppositoria plumbi composita** (B. P.); **Pilula plumbi cum opio** (B. P.). **Plumbi carbonas**, lead carbonate, white lead, $(PbCO_3)_2 \cdot Pb(OH)_2$, is used as a local sedative in ointments and in face-powders. The prolonged use of the latter has caused poisoning. From it is prepared **Unguentum plumbi carbonatis** (U. S. P.). **Plumbi chloridum**, lead chlorid, $PbCl_2$, is used like the carbonate. **Plumbi iodidum**, lead iodid, PbI_2 , is used as a local astringent and absorbent in **Unguentum plumbi iodidi** (U. S. P., B. P.) and **Emplastrum plumbi iodidi** (B. P.). **Plumbi nitras**, lead nitrate, $Pb(NO_3)_2$, is used locally as a sedative to excoriated surfaces, as sore nipples, chapped hands; in gonorrhea, leukorrhea; in onychia maligna, etc. *Ledoyen's* disinfecting fluid is a solution of lead nitrate of the strength of one dram to the ounce. **Plumbi oxidum**, lead oxid, litharge, PbO , is used for making lead plaster (**Emplastrum plumbi**), as an external application to irritated surfaces, and in the arts, for glazing pottery, and as an ingredient of fluid glass. Lead Plaster with olive oil

constitutes Unguentum diachylon (U. S. P.). **Plumbi oxidum rubrum**, red lead, minium, is used extensively in the arts as a paint and in the manufacture of glass. **Plumbi subacetat**, lead subacetate, a basic salt, is used in the form of *Liquor plumbi subacetatis* (U. S. P., B. P.), Goulard's extract, and *Liquor plumbi subacetatis dilutus* (U. S. P., B. P.), lead-water, as a sedative and astringent in inflammations and burns. It is also a constituent of *Ceratum plumbi subacetatis*. **Plumbi tannas**, lead tannate, is used as a sedative astringent.

Plummer's Pills. See *Antimonium*.

Plutomania (*plu-to-ma'-ne-ah*) [*πλοῦτος*, wealth; *μανία*, madness]. An insane belief that one is the possessor of great wealth.

Pneograph (*ne'-o-graf*) [*πνεῖν*, to breathe; *γράφειν*, to write]. An instrument for recording the force and character of the current of air during respiration.

Pneometer (*ne'-om'-et-er*). Synonym of *Spirometer*.

Pneoscope (*ne'-o-skōp*) [*πνεῖν*, to breathe; *σκοπεῖν*, to examine]. An instrument for measuring respiratory movements.

Pneumarthrosis (*nu-mar-thro'-sis*) [*πνεῦμα*, air; *ἄρθρον*, a joint]. A collection of air or gas in a joint.

Pneumathemia (*nu-ma-the'-me-ah*) [*πνεῦμα*, air; *αἷμα*, blood]. The presence of gas in the blood-vessels.

Pneumatic (*nu-mat'-ik*) [*πνεῦμα*, air]. 1. Pertaining to air or gas. 2. Pertaining to respiration. 3. Pertaining to compressed or rarefied air. **P. Speculum.** See *Siegle's Speculum*. **P. Trough**, a trough partly filled with water for facilitating the collection of gases.

Pneumatics (*nu-mat'-iks*) [*πνεῦμα*, air]. The branch of physics treating of the physical properties of air and gases.

Pneumato- (*nu-mat'-o-*) [*πνεῦμα*, air, breath]. A prefix denoting pertaining to air, gas, or breath.

Pneumatocele (*nu'-mat-o-sel*) [*πνεῦμα*, air; *κύλη*, tumor]. 1. A swelling containing air or gas. 2. A hernia of the lung. 3. A swelling of the scrotum produced by the presence of gas.

Pneumatogram (*nu'-mat-o-gram*) [*πνεῦμα*, breath; *γράμμα*, inscription]. A tracing showing the frequency, duration, and depth of the respiratory movements.

Pneumatology (*nu-mat-ol'-o-je*) [*πνεῦμα*, breath; *λόγος*, knowledge]. 1. The science of respiration. 2. The science of gases; also their use as therapeutic agents.

Pneumatometer (*nu-mat-om'-et-er*). See *Spirometer*.

Pneumatometry (*nu-mat-om'-et-re*) [*πνεῦμα*, air; *μετρον*, measure]. 1. The measurement

of the force of respiration. 2. The treatment of pulmonary and circulatory diseases by means of a pneumatic apparatus.

Pneumatosis (*nu-mat'-o'-sis*) [*πνεῦμα*, air]. The presence of gas or air in abnormal places, or in an excessive quantity where a little exists normally.

Pneumatotherapy (*nu-mat-o'-ther'-ap-e*) [*πνεῦμα*, air; *θεραπεία*, treatment]. The treatment of diseases by means of compressed or rarefied air.

Pneumatothorax (*nu-mat-o-tho'-raks*). Synonym of *Pneumothorax*.

Pneumaturia (*nu-mat-u'-re-ah*) [*πνεῦμα*, air; *ουρίνα*, urine]. The evacuation of urine containing free gas.

Pneumectomy (*nu-mek'-to-me*). See *Pneumonectomy*.

Pneumo- (*nu'-mo-*) [*πνεῦμα*, air, or *πνεῦμον*, lung]. A prefix denoting pertaining to the lung. See *Pneumono-*.

Pneumobacillus (*nu-mo-bas-il'-us*). See *Bacteria*, *Table of*.

Pneumocoele (*nu'-mo-sel*). Synonym of *Pneumatocele*.

Pneumocentesis (*nu-mo-sen-te'-sis*) [*πνεῦμον*, lung; *κέντησις*, puncture]. Paracentesis of the lung, especially for the purpose of evacuating a cavity.

Pneumococcus (*nu-mo-kok'-us*). See *Bacteria*, *Table of*.

Pneumokoniosis (*nu-mo-kon-e-o'-sis*). See *Pneumonokoniosis*.

Pneumogastric (*nu-mo-gas'-trik*) [*πνεῦμον*, lung; *γαστήρ*, stomach]. 1. Pertaining conjointly to the lungs and the stomach. 2. Pertaining to the pneumogastric or vagus nerve. **P. Nerve.** See *Nerves*, *Table of*.

Pneumograph (*nu'-mo-graf*) [*πνεῦμον*, lung; *γράφειν*, to write]. An instrument for recording the movements of the chest in respiration.

Pneumography (*nu-mog'-ra-fe*) [*πνεῦμον*, lung; *γράφειν*, to write]. A description of the lungs.

Pneumohemothorax (*nu-mo-hem-o-tho'-raks*) [*πνεῦμα*, air; *αἷμα*, blood; *θώραξ*, thorax]. A collection of air or gas, and blood, in the pleural cavity.

Pneumohydropericardium (*nu-mo-hi-dro-per-e-kar'-de-um*) [*πνεῦμα*, air; *ὕδωρ*, water; *περικάρδιον*]. An accumulation of air and fluid in the pericardial cavity.

Pneumohydrothorax (*nu-mo-hi-dro-tho'-raks*) [*πνεῦμα*, air; *ὕδωρ*, water; *θώραξ*, thorax]. A collection of air or gas, and fluid, in the pleural cavity.

Pneumokoniosis. See *Pneumonokoniosis*.

Pneumolith (*nu'-mo-lith*) [*πνεῦμον*, lung; *λίθος*, a stone]. A calculus of the lung.

Pneumometer (*nu-mom'-et-er*). Synonym of *Spirometer*.

Pneumometry (*nu-mom'-et-re*). Same as *Spirometry*.

Pneumonectasia, Pneumonectasis (*nu-mom-ek-ta'-se-ah, nu-mom-ek'-tas-is*) [*πνεύμων*, lung; *ἐκτασις*, distention]. Emphysema of the lung.

Pneumectomy (*nu-mom-ek'-to-me*) [*πνεύμων*, lung; *ἐκτομή*, excision]. Excision of a portion of a lung.

Pneumonia (*nu-mo'-ne-ah*) [*πνεύμων*, lung].

Inflammation of the lung, pneumonitis. Used without qualification, the term implies lobar pneumonia, *q. v.* **P.**, **Alcoholic**, the croupous pneumonia of drunkards, often associated with delirium, and very fatal. **P.**, **Apex-**, **P.**, **Apical**, croupous pneumonia of the apex of a lung. **P.**, **Aspiration-**, a bronchopneumonia due to the inspiration of food-particles or other irritant substances into the lung. **P.**, **Catarrhal**. Synonym of *Bronchopneumonia*. **P.**, **Central**, a croupous pneumonia beginning in the interior of a lobe of the lung. The physical signs are obscure until the inflammation reaches the surface. **P.**, **Croupous**. See *P.*, *Lobar*. **P.**, **De-glutition-**. Synonym of *P.*, *Aspiration-*. **P.**, **Desquamative**, a form characterized chiefly by an intense desquamation of the cells lining the air-vesicles, a proliferation of the connective-tissue cells of the septa between the vesicles, and the exudation of a scanty albuminous fluid. The exudate generally undergoes caseous degeneration. **P.**, **Embolic**, pneumonia due to embolism of the vessels of the lung. **P.**, **Fibrinous**. See *P.*, *Lobar*. **P.**, **Fibroid**, **P.**, **Fibrous**. Synonym of *P.*, *Interstitial*. **P.**, **Hypostatic**, a lobular *P.* occurring in the dependent portions of the lungs of persons debilitated by age or disease, and depending on the weakened circulation and respiration and the dorsal decubitus. **P.**, **Insular**. Synonym of *P.*, *Lobular*. **P.**, **Interstitial**, fibroid *P.*, cirrhosis of the lung; a chronic inflammation of the lung, characterized by an increase of the connective tissue. It may be due to the inhalation of excessive quantities of coal-dust or other irritant particles (pneumonokoniosis); it may be a termination of lobar pneumonia or bronchopneumonia; it may be due to tuberculosis, or to extension of a chronic inflammation from the pleura. **P.**, **Lobar**, croupous *P.*, lung-fever; an acute infectious disease characterized by an inflammation of one or more lobes of the lung, the affected parts becoming consolidated, owing to the exudation of cells and fibrin into the air-vesicles. The exciting cause is usually the diplococcus pneumoniae of Fränkel, but other microorganisms may produce it. The disease sets in with a chill, which is followed by a rapid rise of the temperature, hurried respiration,

cough, and the expectoration of a peculiar rust-colored sputum. The fever remains high until about the ninth day, when it falls by crisis. The disease generally involves the lower lobe of the right lung and is accompanied by a fibrinous pleurisy. In the early stage the lung is intensely congested (stage of congestion); soon there is an exudation into the vesicles causing solidification of the lobe (stage of consolidation). The lung in the beginning of this stage is dark-red and resembles liver-tissue (stage of red hepatization), later it becomes gray (stage of gray hepatization). In favorable cases the exudate is absorbed and expectorated (stage of resolution). **P.**, **Lobular**. Synonym of *Bronchopneumonia* and *P.*, *Catarrhal*. **P.**, **Massive**, lobar *P.* in which not only the air-cells, but the bronchi of an entire lobe, or even of a lung, are filled with the fibrinous exudate. **P.**, **Pleurogenic**, **P.**, **Pleurogenous**, *P.* secondary to disease of the pleura. **P.**, **Purulent**, one characterized by the formation of pus; it appears under three forms: suppuration of the minute bronchi and air-vesicles—purulent catarrh; true abscess of the lung; suppurative lymphangitis and perilymphangitis. **P.**, **Septic**, lobular *P.* due to the inspiration of septic material or to septic emboli. **P.**, **Syphilitic**, inflammation of the lung due to syphilis and manifesting itself as the white pneumonia of the fetus; as gum-mata of the lung; as interstitial pneumonia, taking its origin at the root of the lung and passing along the bronchi and vessels; and as acute syphilitic phthisis, analogous to acute pneumonic phthisis. **P.**, **Tubular**. Synonym of *Bronchopneumonia*. **P.**, **White**, a catarrhal form of pneumonia occurring in a syphilitic fetus and resulting in death. By an overgrowth of epithelium in the air-vesicles the cells die, and fatty degeneration follows, giving the lungs a white appearance, with the imprint of the ribs on their surface.

Pneumonic (*nu-mom'-ik*) [*πνεύμων*, lung]. Pertaining to the lungs or to pneumonia. **P.** **Phthisis**, tuberculosis affecting a whole lobe of the lung.

Pneumonitis (*nu-mom-i'-tis*) [*πνεύμων*, lung; *ιτις*, inflammation]. Pneumonia.

Pneumono- (*nu-mom'-o-*) [*πνεύμων*, lung]. A prefix denoting pertaining to the lungs.

Pneumonocele (*nu-mom'-o-sel*) [*πνεύμων*, lung; *κῆλη*, tumor, hernia]. Hernia of the lung.

Pneumonokoniosis (*nu-mom-o-kon-e-d'-sis*) [*πνεύμων*, lung; *κόκκιοι*, dust]. A general term applied to chronic induration or fibrous inflammation of the lungs due to the inhalation of dust. Various names are given to it according to the kind of dust causing the inflammation: *anthracosis*, that due to the in-

halation of coal-dust; *siderosis*, that due to inhalation of metallic dust; *chalicosis*, that due to the inhalation of mineral dust.

Pneumonometer (*nu-mon-om'et-er*). Synonym of *Spirometer*.

Pneumonomycosis (*nu-mon-o-mi-ko'-sis*) [*πνεύμων*, lung; *μύκης*, fungus]. Any disease of the lung due to fungi.

Pneumopericardium (*nu-mo-per-e-kar'-de-um*) [*πνεύμα*, air; *περί*, about; *καρδία*, heart]. The presence of air in the pericardial sac. It is due to traumatism, or to communication between the pericardium and the esophagus, stomach, or lungs, and is marked by tympany over the precordial region, and peculiar metallic heart-sounds.

Pneumopyopericardium (*nu-mo-pi-o-per-e-kar'-de-um*) [*πνεύμα*, air; *πίον*, pus; *περί*, about; *καρδία*, heart]. The presence of air or gas and pus in the pericardial sac.

Pneumopyothorax (*nu-mo-pi-o-tho'-raks*) [*πνεύμα*, air; *πίον*, pus; *θώραξ*, chest]. The presence of air and pus in the pleural cavity.

Pneumorrhagia (*nu-mor-a'-je-ah*). See *Hemoptysis*.

Pneumotherapy (*nu-mo-ther'-ap-e*) [*πνεύμα*, air, or *πνεύμων*, lung; *θεραπεία*, treatment]. 1. The treatment of diseases of the lung. 2. See *Pneumatotherapy*.

Pneumothorax (*nu-mo-tho'-raks*) [*πνεύμα*, air; *θώραξ*, chest]. The presence of air or gas in the pleural cavity. It is produced by perforating wounds of the chest, by the rupture of an abscess or tuberculous cavity of the lung, by the rupture of an emphysematous vesicle, or the evacuation of an empyema into the lung or through the chest-wall. It is marked by dyspnea, shock, pain, a tympanitic (sometimes a dull) percussion-note over the affected side, displacement of the heart, bell-tympany, and diminished respiratory murmur.

Pneumotomy (*nu-mot'-o-mo*) [*πνεύμων*, lung; *τομή*, a cutting]. Incision of the lung.

Pneumotoxin (*nu-mo-toks'-in*) [*πνεύμων*, lung; *τοξικόν*, a poison]. A poisonous albuminoid body produced by the pneumococcus, and believed to be the cause of many of the symptoms of lobar pneumonia. Antipneumotoxin is the name given to the antitoxin supposed to exist in the blood of persons convalescent from lobar pneumonia.

Pneumotyphus (*nu-mo-ti'-fus*) [*πνεύμων*, lung; *τίφος*, typhus]. 1. Typhoid fever beginning with pneumonia dependent upon the typhoid bacillus. 2. Pneumonia occurring in the course of typhoid fever.

Pock (*pok*) [A.S., *poc*, a pustule]. A pustule of an eruptive fever, especially of smallpox. P.-marked, marked with the cicatrices of the smallpox pustule.

Podagra (*pod-a'-grah*) [*πούς*, foot; *ἄγγρα*, seizure]. Gout, especially of the great toe or the joints of the foot.

Podalgia (*pod-al'-je-ah*) [*πούς*, foot; *ἄλγος*, pain]. Pain in the foot.

Podalic (*pod-al'-ik*) [*πούς*, foot]. Pertaining to the feet. P. Version, the operation of changing the position of the fetus in utero so as to bring the feet to the outlet.

Podelcoma (*pod-el-ko'-mah*). See *Fungus-foot*.

Podencephalus (*pod-en-sef'-al-us*) [*πούς*, foot; *ἐγκέφαλος*, brain]. A variety of monster of the species exencephalus, in which there is a protrusion of the cranial contents from the top of the head.

Pododynia (*pod-o-din'-e-ah*) [*πούς*, foot; *ὀδύνη*, pain]. Pain in the foot, especially a neuralgic pain in the heel unattended by swelling or redness.

Podophyllin (*pod-o-fil'-in*). See *Podophyllum*.

Podophyllotoxin (*pod-o-fil-o-toks'-in*). See *Podophyllum*.

Podophyllum (*pod-o-fil'-um*) [*πούς*, foot; *φύλλον*, leaf]. The May-apple or mandrake, *P. peltatum*, of the natural order Berberidæ. The rhizome and roots (P., U. S. P., B. P.) contain an active resin, podophyllin (Resina podophylli, U. S. P.), podophyllotoxin (C₂₂H₂₄O₈ + 2H₂O), having properties similar to those of the resin; picropodophyllin, a derivative of podophyllotoxin; and podophylloquercetin. P. is used in medicine as a laxative in chronic constipation and as a cathartic in hepatic congestion and bilious fever. Dose of the resin (Podophylli resina, B. P.), gr. 1/2 - 1/2 (0.008-0.03); of podophyllotoxin, gr. 1/2 - 1/2 (0.016-0.03). Extractum podophylli (U. S. P.), dose gr. j-v (0.065-0.32). Extractum podophylli fluidum (U. S. P.), ℥ij-xx (0.13-1.3). Tinctura podophylli (B. P.), ℥xx-45 (1.0-4.0).

Poikilocyte (*poi-kil-o-sit*) [*ποικίλος*, varied; *κύτος*, cell]. A large red blood-corpuscle of irregular shape. Poikilocytes are most abundant in the blood in pernicious anemia, but also occur in other forms of anemia.

Poikilocytosis (*poi-kil-o-si-to'-sis*) [*ποικίλος*, varied; *κύτος*, cell]. A condition of the blood characterized by the presence of poikilocytes.

Poikilothermic (*poi-kil-o-ther'-mik*) [*ποικίλος*, varied; *θερμῆ*, heat]. Varying in temperature according to the surroundings; cold-blooded.

Point [*punctum*, point, from *pungere*, to prick]. 1. The sharp end of an object, especially one used to pierce anything. 2. The limit at which anything occurs, as the melting-P., freezing-P. 3. A mark made by a sharp object; a minute spot or area. P.,

Craniometric. See *Craniometric Points*. **P.**, **Critical**, of gases, a temperature at or above which a gas cannot be liquefied by pressure alone; of liquids, that temperature at which a liquid, regardless of the pressure to which it is subjected, assumes a gaseous form. **P.**, **Dew-**, the temperature at which the atmospheric moisture is deposited as dew. **P. of Election**, in surgery, that point at which a certain operation is done by preference. **P.**, **Far**, the remotest point of distinct vision. **P.**, **Hystero-epileptogenous**, **P.**, **Hysterogenic**. See *Zone*. **P.**, **McBurney's**. See *McBurney*. **P.**, **Motor**. See *Motor Point*. **P.**, **Near**, the nearest point at which the eyes can accommodate to see distinctly. **P.**, **Principal**, one of the two points in the optic axis of a lens that are so related that lines drawn from these points to the corresponding points in the object and its image are parallel. **P.**, **Valleix's**

(*Pointe douloureux*), any one of the points of tenderness at the exit or in the course of a nerve the seat of neuralgia.

Point [*pungere*, to prick]. Of an abscess, to come to the surface.

Poison (*poi'-zon*) [*Fr.*, from *potio*, a draught]. A substance that when introduced into the body either destroys life or impairs seriously the functions of one or more of its organs.

P., **Irritant**, one that causes irritation at the point of entrance or at the point of elimination.

P., **Muscle-**. 1. A substance that impairs or destroys the proper functions of muscles. 2.

A poisonous albumin developed during muscular activity. **P.**, **Narcotic**, one affecting the cerebral centers, producing stupor. **P.**,

Ordeal, any one of the vegetable poisons, such as physostigma, used by savages in the trial of accused persons to determine their guilt or innocence.

TABLE OF POISONS.

NAME.	SYMPTOMS OF POISONING.	TREATMENT AND ANTIDOTES. (Antidotes in <i>Italics</i> .)
Acid, Carbolic (<i>Phenol</i>).	Immediate burning pain from mouth to stomach; giddiness, loss of consciousness, collapse; partial suppression of urine, which is smoky in color; characteristic odor; white, corrugated patches in mouth.	Stomach-pump; <i>magnesium sulphate</i> or <i>sodium sulphate</i> , atropin.
Acid, Chromic.	Dark yellow stains; abdominal pain; vomiting and purging; collapse.	Evacuate stomach; <i>chalk</i> , milk, or albumin; demulcent drinks.
Acid, Hydrochloric (<i>Muriatic</i>).	Pain throughout digestive tract; vomiting, feeble pulse, clammy skin, collapse; eschars externally; yellow stains on clothing, but none on skin.	<i>Alkalies</i> ; demulcent drinks; oil; stimulants (intravenous injection).
Acid, Hydrocyanic (<i>Prussic</i>).	Sudden unconsciousness, slow, labored respirations, slow pulse, staring eyes, purple face, general convulsions then relaxation and collapse; odor of peach-kernels. Death may be almost instantaneous.	Stomach-pump if possible; dilute ammonia; alternate cold and warm affusions; atropin and cardiac stimulants; artificial respiration.
Acid, Nitric.	Yellow stains on skin; otherwise similar to <i>Acid, Sulphuric</i> .	<i>Alkalies</i> ; <i>soap</i> ; demulcents; stimulants.
Acid, Oxalic.	Hot, acrid taste; burning, vomiting, collapse; sometimes general paralysis, numbness, and stupor.	<i>Lime</i> or <i>chalk</i> .
Acid, Salicylic.	Mydriasis; quick and deep respiration; delirium; dyspnea; lessened arterial pressure; deafness; olive-green urine.	
Acid, Sulphuric.	Black stains; pain throughout digestive tract, vomiting, often of tarry matter, feeble pulse, clammy skin, profuse and bloody salivation.	<i>Chalk</i> ; <i>magnesia</i> ; <i>soap</i> ; demulcent drinks.
<i>Aconitum napellus</i> (<i>Monkshood</i>). <i>Aconite</i> .	Sudden collapse, slow, feeble and irregular pulse and respirations, tingling in the mouth and extremities, giddiness, great muscular weakness, sometimes pained in the abdomen; pupils dilated, but may be contracted; marked anesthesia of skin; mind clear, convulsions at times.	<i>Tannic acid</i> solution for washing out stomach; <i>digitalis</i> , atropin, and stimulants; artificial respiration; warmth and friction; absolute quiet in recumbent position.

TABLE OF POISONS.—Continued.

NAME.	SYMPTOMS OF POISONING.	TREATMENT AND ANTIDOTES. (Antidotes in <i>Italics</i> .)
Alcohol.	Confusion of thought, giddiness, tottering gait, slight cyanosis, narcosis from which patient can be aroused; full pulse; deep, stertorous breathing; injection of eyes, dilatation of pupils, low temperature. Convulsions may occur.	Evacuate stomach; coffee; battery; amyl nitrite; hot and cold douches.
Ammonium and its compounds.	Intense gastroenteritis often with bloody vomiting and purging; lips and tongue swollen and covered with detached epithelium; violent dyspnea; characteristic odor.	<i>Vegetable acids</i> ; demulcents.
Antimony and its compounds.	Metallic taste, violent vomiting, becoming bloody; feeble pulse; pain and burning in the stomach; violent serous purging, becoming bloody; cramps in extremities, thirst, great debility; sometimes prostration, collapse, unconsciousness and convulsions without vomiting or purging.	<i>Tannic acid</i> ; demulcent drinks; opium; alcohol; external heat.
Antipyrin (<i>Phenazone</i>).	Headache, nausea, vomiting, a rash like that of measles, vertigo, drowsiness, deafness, confusion of ideas, cyanosis, collapse.	Recumbent position; warmth; strychnin; stimulants; oxygen; artificial respiration.
Apomorphin.	Violent vomiting, paralysis of motor and sensory nerves, delirium, depression of respiration and of heart.	Cardiac and respiratory stimulants.
Arsenic and its compounds.	Violent burning pain in the stomach, retching, thirst, purging of blood and mucus with flakes of epithelium, tenesmus, suppression of urine; sense of constriction in throat; pulse small and frequent.	<i>Hydrated sesquioxide of iron</i> ; <i>precipitated carbonate of iron</i> ; emetics; castor oil; demulcents.
Atropa belladonna (<i>Deadly Nightshade</i>). Atropin. Belladonna. Homatropin.	Heat and dryness of the mouth and throat; pupils widely dilated, scarlet rash, noisy delirium; quick pulse, at first corded, later feeble; rapid respirations, early strong, late shallow and feeble; retention of urine; sometimes convulsions, collapse, and paralysis.	Evacuate stomach; <i>tannic acid</i> ; stimulants; coffee; pilocarpin; artificial respiration; physostigmin may be of benefit; evacuation of bladder.
Caffein.	Burning pain in the throat, giddiness, faintness, nausea, numbness, abdominal pain, great thirst, dry tongue, tremor of extremities, diuresis, weak pulse, cold skin, collapse.	Emetics; stimulants; warmth; morphin and atropin.
Calabar Bean. See	<i>Physostigma</i> .	
Camphor.	Characteristic odor; languor, giddiness, disturbance of vision, delirium, convulsions, clammy skin, smarting in the urinary organs; pulse quick and weak; no pain, no vomiting, no purging.	Evacuate stomach; stimulants; warmth; hot and cold douches.
Cannabis indica (<i>Indian Hemp</i>).	Pleasurable intoxication, sense of prolongation of time, anesthesia with loss of strength, especially in legs; pupils dilated; rapid pulse; heavy sleep.	Evacuate stomach; stimulants.
Cantharis vesicatoria (<i>Spanish Fly</i>). Cantharides.	Burning in mouth and stomach; vomiting and purging soon becoming bloody; tenesmus, salivation, aching pains in back, strangury, priapism; unconsciousness only very late; convulsions at times.	Evacuate stomach; demulcent drinks; morphin; hot bath for the strangury; anesthetics may be necessary for the pain.
Carbolic Acid. See	<i>Acid, Carbolic</i> .	

TABLE OF POISONS.—*Continued.*

NAME.	SYMPTOMS OF POISONING.	TREATMENT AND ANTIDOTES. (Antidotes in <i>Italics</i> .)
Chloral Hydrate.	Deep sleep, loss of muscular power, lividity; reflexes diminished, pulse weak, respirations slowed, pupils contracted during sleep, but dilated on waking, temperature low.	Evacuate stomach; heat to the extremities; massage; coffee per rectum; strychnin; amyl nitrite; artificial respiration.
Chloroform.	Excitement and intoxication followed by anesthesia and unconsciousness, later profound narcosis. Pulse and respirations fail progressively or suddenly.	Draw tongue forward; artificial respiration; faradic current; hot and cold douches; amyl nitrite; ammonia injected into a vein; evacuation of the stomach if chloroform has been taken by mouth.
Coal-gas.	Headache, giddiness, loss of muscular power, unconsciousness, pupils dilated, breathing labored, coma; odor of the gas.	Fresh air; artificial respiration; ammonia; stimulants; oxygen; coffee; hot and cold douches.
Cocain.	Faintness, giddiness, nausea; pulse small, rapid, intermittent; dilated pupils, severe prostration, respiration slow and feeble.	Stimulants; amyl nitrite; artificial respiration.
Colchicum autumnale (Meadow-saffron).	Not unlike those of malignant cholera; gripping pain in the stomach, vomiting and continuous purging of seromucous material; intense thirst, muscular cramps, great prostration, collapse, dilated pupils, pain in the extremities.	Evacuate stomach; <i>tannic</i> or <i>gallic acid</i> ; demulcent drinks; stimulants; morphin.
Conium maculatum (Hemlock).	Weakness of the legs, gradual loss of all voluntary power, nausea, ptosis, dilatation of pupils, inability to speak or swallow.	Evacuate stomach; <i>tannic</i> or <i>gallic acid</i> ; stimulants; warmth; artificial respiration; atropin.
Croton tiglium (<i>Croton-oil</i>).	Intense pain in abdomen, vomiting, purging, watery stools, pinched face, small and thready pulse, moist skin, collapse.	Evacuate stomach; demulcent drinks; camphor; stimulants; morphin; poultices to abdomen.
Cyanogen and its compounds.	Similar to <i>Acid, Hydrocyanic, q. v.</i>	
Datura stramonium (Thorn-apple, Jamestown Weed).	Symptoms and treatment similar to those of <i>Atropin, q. v.</i>	
Digitalis purpurea (Foxglove).	Purging, with severe pain, violent vomiting, vertigo, feeble pulse, although heart's action is tumultuous, eyes prominent, pupils dilated, sclera blue; delirium and convulsions.	Evacuate stomach; <i>tannic</i> and <i>gallic acids</i> ; stimulants; aconite; recumbent position.
Erythroxylon coca.	See <i>Cocain</i> .	
Fly, Spanish. See <i>Cantharis</i> .		
Gelsemium sempervirens (Yellow Jessamine).	Symptoms appear in about twenty minutes; great muscular weakness; diplopia, ptosis, internal squint, widely dilated pupils, dimness of vision, labored respiration, weak pulse.	Evacuate stomach; atropin; stimulants; artificial respiration; hot and cold douches.
Hellebore, Green and Hemlock. See <i>Conium</i> .	White. See <i>Veratrum</i> .	
Iodin and its compounds.	Pain in throat and stomach; vomiting, purging, vomit yellow from iodine, or blue if starch present in stomach; giddiness, faintness, convulsive movements.	Evacuate stomach; <i>starch</i> ; amyl nitrite; morphin.
Iodoform.	Slight delirium, drowsiness, high temperature, rapid pulse. Symptoms resemble meningitis.	
Jaborandi. Pilocarpin.	Copious sweating, dizziness, salivation, vomiting, diarrhea, tearing pain in eyeballs, myopia, pupils much contracted.	Evacuate stomach; stimulants; atropin.

TABLE OF POISONS.—Continued.

NAME.	SYMPTOMS OF POISONING.	TREATMENT AND ANTIDOTES. (Antidotes in <i>Italics</i> .)
Lead acetate.	Sweet metallic taste, vomiting of white matter, great thirst, pain in abdomen, abdominal muscles usually rigid, constipation or diarrhea with black stools, cramps in the legs, paralysis of the extremities, convulsions; in the chronic forms, a blue line at margin of the gums.	Evacuate stomach; <i>dilute sulphuric acid</i> ; <i>Epsom</i> or <i>Glauber's salts</i> ; milk; morphia; potassium iodid to eliminate the poison.
<i>Lobelia inflata</i> (<i>Indian Tobacco</i>).	Severe vomiting, with intense depression and prostration, giddiness, tremors, convulsions, collapse.	Evacuate stomach; <i>tannic</i> or <i>gallic acid</i> ; stimulants; <i>strychnin</i> ; warmth; recumbent position.
Mercuric chlorid.	Acrid metallic taste, burning heat in throat and stomach, vomiting, diarrhea, with bloody stools, lips and tongue white and shriveled, pulse small and frequent, death in coma or convulsions; pain may be absent. Secondary symptoms: hectic fever, coppery taste, fetid breath, gums swollen, salivation.	Albumin in some form; raw white of egg or flour; evacuate stomach; opium; potassium iodid.
Morphin. See <i>Opium</i> .		
Nitric Acid. See <i>Acid</i> .	<i>Nitric</i> .	
Nitroglycerin.	Throbbing headache, pulsation over entire body, diastolic pulse, flushed face, mental confusion, anxiety, sudden collapse.	Recumbent position; cold to head; ergot; atropin.
<i>Nux vomica</i> . See	<i>Strychnos</i> .	
Opium. Morphin. Narcotin. Codein. Laudanum.	Preliminary mental excitement, acceleration of heart; soon weariness, sensation of weight in the limbs, sleepiness, diminished sensibility, pin-point pupils; pulse and respiration slow and strong; patient can be roused with difficulty; later this becomes impossible; reflexes abolished, respiration slow, irregular, and stertorous, pulse rapid and feeble.	1. Evacuate stomach by mustard or stomach-pump. 2. Arouse patient to maintain respiration by exercise, flagellation with wet towels, cold and hot douches alternately. 3. Stimulate by <i>atropin</i> , coffee, alcohol if pulse fails; external heat. Inhalations of oxygen; injection of dilute solution of <i>potassium permanganate</i> .
Oxalic Acid. See <i>Acid</i> .	<i>Oxalic</i> .	
Paris Green. See <i>Arsenic</i> .	<i>nit.</i>	
Phosphorus.	Vomiting and pain; vomit may be luminous in the dark; characteristic odor. After several days deep jaundice, coffee-colored vomit, hepatic tenderness, albuminuria, marked fall in temperature, coma, failure of pulse and respiration.	<i>Sulphate of copper</i> as an emetic, then as an antidote in small doses with opium; purgation.
<i>Physostigma venenosum</i> (<i>Calabar Bean</i>).	Giddiness, prostration, loss of power in the lower limbs, muscular twitching, contracted pupils, mind clear.	Evacuate stomach; <i>atropin</i> ; <i>strychnin</i> ; stimulants; artificial respiration.
Prussic Acid. See	<i>Acid, Hydrocyanic</i> .	
Santonin.	Disturbance of color-vision—objects first assume a bluish tinge, then yellow; tinnitus, dizziness, pain in the abdomen, failure of respiration, convulsions, stupor.	Evacuate stomach; stimulants; chloral.
Savin.	Pain, vomiting, bloody stools and tenesmus; disordered respirations, coma, convulsions, and collapse.	Evacuate stomach; castor-oil in large dose; morphia; poultices to the abdomen.
Silver-salts.	Pain, vomiting, and purging; vomit white and cheesy, rapidly turning black in the sunlight; vertigo, coma, convulsions, paralysis, and marked disturbance of respiration.	<i>Salt</i> and water; evacuate stomach; a large amount of milk.

TABLE OF POISONS.—*Continued.*

NAME.	SYMPTOMS OF POISONING.	TREATMENT AND ANTIDOTES. (Antidotes in <i>Italics</i> .)
St. Ignatius Bean. Strychnos ignatii. Strychnos nux vomica Nux vomica. Strychnin. Brucin.	Tetanic convulsions in paroxysms at varying intervals of from five minutes to half an hour; opisthotonos during paroxysm; eyeballs prominent, pupils dilated, respiration impeded, pulse feeble and rapid; anxiety.	Evacuate stomach; <i>tannic acid</i> followed by an emetic; catheterize; keep patient quiet; bromids and chloral; amyl nitrite or chloroform to control convulsions; artificial respiration if indicated.
Tobacco. Nicotin.	Nausea, vomiting, weakness, weak pulse, cold and clammy skin, collapse, pupils contracted, then dilated.	Evacuate stomach; <i>tannic acid</i> ; strychnin; stimulants; warmth; recumbent position.
Veratrum album (<i>White Hellebore</i>). Veratrum viride (<i>Green Hellebore</i>).	Pain and burning in alimentary tract; vomiting and diarrhea; slow, weak pulse; labored respiration; pupils usually dilated; there may be convulsions.	Evacuate stomach; ether hypodermatically; opium; stimulants; coffee; warmth; recumbent position.

Poke-root. See *Phytolacca*.

Polar (*po'-lar*) [*polus*, a pole]. Pertaining to or situated near a pole. **P. Bodies, P. Cells,** or **P. Globules**, two minute cells thrown off by the unfecundated ovum during maturation. **P. Method**, a method of applying electricity, in which the pole the distinctive effect of which is wanted is placed over the part to be treated, and the other pole over some indifferent part.

Polarimeter (*po-lar-im'-et-er*) [*polus*, a pole; *μέτρον*, a measure]. An instrument for determining the degree to which an optically active substance changes the plane of polarization to the right or to the left.

Polariscope (*po-lar'-is-kop*) [*polus*, pole; *σκοπεῖν*, to view]. An instrument for studying the polarization of light; a polarimeter.

Polarity (*po-lar'-it-ē*) [*polus*, pole]. The state of having poles or points of intensity presenting opposite qualities.

Polarization (*po-lar-i-sa'-shun*) [*polus*, pole]. 1. The act of polarizing, or the state of being polarized. 2. A condition produced in a ray of light by absorption, reflection, or refraction, by means of which the vibrations are restricted and take place in one plane only (plane P.), or in curves (circular or elliptic P.). The plane of P. is altered or rotated when the light is passed through a quartz-crystal or solutions of certain substances (rotatory P.). 3. The deposit of gas-bubbles (hydrogen) on the electronegative plate of a galvanic battery, whereby the flow of the current is impeded, and owing to the negative plate covered with hydrogen being more electropositive than the zinc-plate, the difference in potential between the two plates is reduced.

Polarize (*po'-lar-iz*) [*polus*, pole]. To endow with polarity; to place in a state of polarization.

Polarizer (*po'-lar-i-ser*) [*polus*, pole]. An object, such as a Nicol-prism, by means of which light is polarized.

Pole (*pōl*) [*πόλος*, *polus*, a pole]. 1. Either extremity of the axis of a body, as of the fetus, the crystalline lens, etc. 2. One of two points at which opposite physical qualities, *e. g.*, electricity or magnetism, are concentrated; specifically, the electrode of a galvanic battery, which is positive (positive P.) when connected with the electronegative plate of the battery (carbon, copper, platinum), or negative (negative P.) when connected with the electropositive plate (zinc). **P.-changer**, a switch or key for changing or reversing the direction of a current produced by an electric battery.

Poliencephalitis (*pol-e-en-sef-al-i'-tis*) [*πολιός*, gray; *ἐγκέφαλος*, brain; *ιτις*, inflammation]. Inflammation of the gray matter of the brain. **P. acuta**, an acute inflammation of the cerebral cortex, which when occurring in children, gives rise to infantile cerebral palsy. **P., Anterior Superior**, an inflammatory disease of the gray matter of the third ventricle, of the anterior portion of the fourth, and of that about the Sylvian aqueduct. It is characterized by ophthalmoplegia, chiefly external, and a peculiar somnolent state.

Poliomyelitis (*pol-e-o-mi-el-i'-tis*) [*πολιός*, gray; *μυελός*, marrow; *ιτις*, inflammation]. Inflammation of the gray matter of the spinal cord. **P., Acute Anterior**, infantile paralysis, an acute inflammation of the anterior horns of the gray matter of the spinal cord. It is most common in children, coming on during the period of the first dentition and producing a paralysis of certain muscle-groups or of an entire limb. The onset is sudden, and the paralysis is usually most extensive in the beginning, a certain amount of improvement taking place subsequently. The affected

muscles atrophy rapidly, the reflexes in them are lost, and reaction of degeneration develops. From contraction of antagonistic muscles deformities occur later in life. **P.**, Chronic Anterior. Synonym of *Progressive Muscular Atrophy*.

Poliomyelopathy (*pol-e-o-mi-el-ōp'-ath-e*) [*πολιός*, gray; *μυελός*, marrow; *πάθος*, disease]. Disease of the gray matter of the spinal cord and medulla oblongata.

Poliosis (*pol-e-ō'-sis*) [*πολιός*, hoary or gray]. A condition characterized by absence of pigment in the hair.

Politzerization (*pol-its-er-i-zā'-shun*) [after Adam Politzer, an Austrian otologist]. The inflation of the middle ear by means of a rubber-bag.

Politzer's Bag. A pear-shaped rubber bag with a conical tip, used for inflating the middle ear. The tip is introduced into the nostril, and the bag compressed while the other nostril is closed and the patient performs the act of swallowing. The latter opens the Eustachian tube and allows the air to enter.

Pollakiuria (*pol-ak-e-u'-re-ah*) [*πολλάκις*, often; *urina*, urine]. Abnormally frequent micturition.

Pollex (*pol'-eks*) [L.]. 1. The thumb. 2. The great toe.

Pollution (*pol-u'-shun*) [*polluere*, to defile]. 1. The act of defiling or rendering impure, as P. of drinking-water. 2. The production of the sexual orgasm by means other than sexual intercourse.

Poly- (*pol'-e-*) [*πολύς*, many]. A prefix denoting many.

Polyarthritis (*pol-e-ar-thri'-tis*) [*πολύς*, many; *άρθρον*, joint; *ιτις*, inflammation]. Inflammation of many joints. **P. rheumatica acuta**. See *Rheumatism, Acute Articular*.

Polyarticular (*pol-e-ar-tik'-u-lar*) [*πολύς*, many; *articulus*, joint]. Affecting many joints; the term multiarticular is preferable.

Polyatomic (*pol-e-at-ōm'-ik*) [*πολύς*, many; *atomic*]. 1. Containing several atoms. 2. Having several hydrogen-atoms replaceable by bases.

Polybasic (*pol-e-bā'-sik*) [*πολύς*, many; *βάσις*, base]. 1. Of acids, having several hydrogen-atoms replaceable by bases. 2. Formed from a polybasic acid by the replacement of more than one hydrogen atom by a base.

Polycholia (*pol-e-kō'-le-ah*) [*πολύς*, much; *χολή*, bile]. Excessive secretion of bile.

Polychromatic (*pol-e-kro-mat'-ik*) [*πολύς*, many; *χρώμα*, color]. Many-colored.

Polyclinic (*pol-e-klin'-ik*) [*πολύς*, many; *κλίνη*, a bed]. A hospital in which many diseases are treated.

Polycoria (*pol-e-kō'-re-ah*) [*πολύς*, many; *κόρη*, pupil]. The existence of more than one pupil in the iris.

Polycrotic (*pol-e-krot'-ik*) [*πολύς*, many; *κρότος*, pulse]. Of the pulse, presenting several waves for each cardiac systole.

Polycystic (*pol-e-sis'-tik*) [*πολύς*, many; *κίστις*, a sac]. Containing many cysts.

Polycythemia (*pol-e-si-thē'-me-ah*) [*πολύς*, many; *κύτος*, cell; *αἷμα*, blood]. A state of the blood characterized by an excess of red corpuscles.

Polydactylism (*pol-e-dak'-til-izm*) [*πολύς*, many; *δάκτυλος*, a finger]. The existence of supernumerary fingers or toes.

Polydipsia (*pol-e-dip'-se-ah*) [*πολύς*, much; *δίψα*, thirst]. Excessive thirst.

Polyemia (*pol-e-e'-me-ah*) [*πολύς*, much; *αἷμα*, blood]. Abnormal increase of the total mass of the blood; plethora. **P. hyperalbuminosa**, an excess of albumin in the blood-plasma. **P. polycythæmica**, an increase of the red corpuscles. **P. serosa**, a condition in which the amount of blood-serum is increased.

Polyesthesia (*pol-e-es-thē'-ze-ah*) [*πολύς*, many; *αἴσθησις*, sensation]. An abnormality of sensation in which a single touch is felt in two or more places at the same time.

Polygalactia (*pol-e-gal-ak'-te-ah*) [*πολύς*, much; *γάλα*, milk]. Excessive secretion of milk.

Polygalin (*pol-ig'-al-in*). See *Senega*.

Polyganglionic (*pol-e-gang-gle-on'-ik*) [*πολύς*, many; *γάγγλιον*, ganglion]. 1. Having several ganglia. 2. Affecting several lymphatic glands at once.

Polygnathus (*pol-ig'-na-thus*) [*πολύς*, many; *γνάθος*, jaw]. A form of monster in which the parasite is attached to the jaws of the host.

Polygyria (*pol-e-jir'-e-ah*) [*πολύς*, many; *γίρος*, gyre]. The existence of an excessive number of convolutions in the brain.

Polyhydramnios (*pol-e-hi-dram'-ne-ōs*) [*πολύς*, many; *ἵδωρ*, water; *ἀμνιον*, amnion]. An excessive production of liquor amnii.

Polyidrosis (*pol-e-id-ro'-sis*) [*πολύς*, much; *ἰδρωσις*, sweating]. Excessive sweating.

Polymastia (*pol-e-mas'-te-ah*) [*πολύς*, many; *μαστός*, breast]. The presence of more than two breasts or nipples. The supernumerary organs may be below the breast proper, in the axilla, or elsewhere on the body.

Polymelia (*pol-e-me'-le-ah*) [*πολύς*, many; *μέλος*, limb]. A malformation consisting in the presence of more than the normal number of limbs.

Polymelus (*pol-im'-el-us*) [*πολύς*, many; *μέλος*, a limb]. A monster having more than the normal number of limbs.

Polymeric (*pol-e-me'-ik*) [*πολύς*, many; *μέρος*, part]. Exhibiting polymerism.

Polymeride (*pol-im'-er-id*) [*πολύς*, many;

μέρος, a part]. In chemistry, a compound having the property of polymerism.

Polymerism (*pol-im'-er-izm*) [*πολύς*, many; *μέρος*, a part]. 1. The existence of more than a normal number of parts. 2. A variety of isomerism. See *Isomeric* and *Polymerization*.

Polymerization (*pol-e-mer-i-sa'-shun*) [*πολύς*, much; *μέρος*, a part]. The apparent fusion or union of two or more molecules of a compound, forming a more complex molecule, with a higher molecular weight and somewhat different physical and chemical properties.

Polymorphic, Polymorphous (*pol-e-mor'-fik*, *pol-e-mor'-fus*) [*πολύς*, many; *μορφή*, form]. Having or occurring in several forms; of a crystal, crystallizing in several forms.

Polymorphism (*pol-e-mor'-fizim*) [*πολύς*, many; *μορφή*, form]. The state of being polymorphous.

Polyneuritis (*pol-e-nu-ri'-tis*). See *Neuritis*, *Multiple*.

Poly-nuclear (*pol-e-nu'-kle-ar*) [*πολύς*, many; *nucleus*, nucleus]. See *Multinuclear*, the preferable term.

Polyopia, Polyopsia (*pol-e-o'-pe-ah*, *pol-e-op'-se-ah*) [*πολύς*, many; *ὄψις*, sight]. A condition in which more than one image of an object is formed upon the retina.

Polyorchis (*pol-e-or'-kis*) [*πολύς*, many; *ὄρχις*, a testicle]. One who has more than two testicles.

Polyotia (*pol-e-o'-she-ah*) [*πολύς*, many; *ὄτις*, ear]. A condition in which there is more than one auricle on a side.

Polyp (*pol'-ip*) [*πολύς*, many; *πούς*, foot]. A tumor having a distinct pedicle. See *Polypus*.

Polypapilloma tropicum (*pol-e-pap-il-o'-mah*) [*πολύς*, many; *papilla*, papilla; *δμα*, tumor]. Frambesia.

Polyparesis (*pol-e-par'-es-is*) [*πολύς*, much; *πάρεσις*, weakness]. General paresis.

Polyphagia (*pol-e-fa'-je-ah*) [*πολύς*, much; *φάγειν*, to eat]. Bulimia.

Polypharmacy (*pol-e-far'-mas-e*) [*πολύς*, many; *φάρμακον*, a drug]. The prescription of many drugs at one time; the excessive use of drugs.

Polyphrasia (*pol-e-fra'-se-ah*) [*πολύς*, much; *φράσις*, speech]. A morbid state characterized by excessive speaking; morbid loquacity; verberation.

Polyplastic (*pol-e-plas'-tik*) [*πολύς*, much; *πλάσσειν*, to mold]. 1. Of cells, having many substances in their composition. 2. Undergoing many modifications during development.

Polypnea (*pol-ip-ne'-ah*) [*πολύς*, many; *πνοία*, breathing]. Great rapidity of respiration; panting respiration.

Polypoid (*pol'-e-poid*) [*polypus*; *εἶδος*, like]. Resembling a polyp.

Polypus (*pol'-e-pus*) [*πολύς*, many; *πούς*, foot; *pl.*, polypi]. A tumor having a pedicle, found especially on mucous membranes, as in the nose, bladder, rectum, uterus, etc. **P.**, **Blood-**. Synonym of **P.**, **Placental**. **P.**, **carneous**. Synonym of **Sarcoma**. **P.**, **Fibrinous**, a polypoid mass on the uterine wall, resulting from the deposition of fibrin from retained blood. The mass may be attached to portions of an ovum or to thrombi at the placental site. **P.**, **Fibrous**, a polypus composed chiefly of fibrous tissue. **P.**, **Mucous**, a soft **P.** resulting either from a localized inflammatory hyperplasia of a mucous membrane or from the formation of a true myxoma. **P.**, **Placental**, a fibrinous polyp resulting from the deposit of fibrin upon a portion of retained placenta. **P.**, **Soft**. Synonym of **P.**, **Mucous**. **P.**, **Vascular**, a polypoid angioma.

Polyrrhea (*pol-e-re'-ah*) [*πολύς*, much; *ῥοία*, flow]. An excessive secretion of fluid.

Polysarcia (*pol-e-sar'-se-ah*) [*πολύς*, much; *σάρξ*, flesh]. Excessive corpulency; obesity.

Polysarcous (*pol-e-sar'-kus*) [*πολύς*, much; *σάρξ*, flesh]. Corpulent; exhibiting polysarcia.

Polyscelia (*pol-e-ske'-le-ah*) [*πολύς*, many; *σκέλος*, leg]. Excess in the number of legs.

Polyscelus (*pol-is'-kel-us*) [*πολύς*, many; *σκέλος*, leg]. A monster having supernumerary legs.

Polyscope (*pol'-is-kōp*) [*πολύς*, much; *σκοπεῖν*, to observe]. An instrument provided with an electric light which is introduced into a cavity in order to illuminate its internal surfaces.

Polysomia (*pol-e-so'-me-ah*) [*πολύς*, many; *σῶμα*, body]. A monster having more than a single body or trunk.

Polyspermism (*pol-e-sper'-mizm*) [*πολύς*, much; *σπέρμα*, seed]. An excessive quantity of seminal fluid.

Polytrichia, Polytrichosis (*pol-e-trik'-e-ah*, *pol-e-trik'-o'-sis*) [*πολύς*, many; *θρίξ*, hair]. Excessive development of hair.

Polyuria (*pol-e-u'-re-ah*) [*πολύς*, much; *urina*, urine]. The passage of an excessive quantity of urine. The causes of temporary **P.** are excessive ingestion of fluids, cold, suppression of perspiration, the use of diuretics; it occurs in the crisis of fevers, in hysteria, and nervous excitement. A permanent **P.** is met with in diabetes mellitus, diabetes insipidus, chronic interstitial nephritis, and in amyloid disease of the kidneys.

Pomade (*po-mad'*) [*ponium*, apple]. A perfumed ointment for applying to the scalp.

Pomatium (*po-ma'-tum*). See *Pomade*.

Pomegranate (*pum-gran'-et*). See *Granatum*.

Pompholyx (*pom'-fo-lik's*) [*πομφόλυξ*, bubble]. Cheiropompholyx; dysidrosis; a rare disease characterized by bullous eruptions on the palms of the hands and between the fingers. It occurs in depressed states of the nervous system, and is more common in women than in men.

Pons (*pom'-fus*) [*πομπός*, blister]. See *Pontile*.

Pomum (*po'-mum*) [L.]. Apple. **P. adam's**, Adam's apple, the prominence in the throat at the base of the neck caused by the projection of the thyroid cartilage.

Pons, bridge]. A process or bridge of gray matter connecting two parts of an organ. **P. cerebelli**. Same as *P. Varolii*. **P. posterior**, the posterior perforated space. **P. sphenoidal**, or simply **Pons**, a convex white structure situated at the base of the brain, serving to connect the various divisions of the brain with each other. It is placed in front of the medulla oblongata, behind the cerebrum, and beneath the cerebellum, and rests upon the sphenobasilar groove. In structure the pons consists chiefly of nerve-fibers, but contains also areas of gray matter, the *basal nuclei*. On section two portions, a superior and an inferior, are distinguishable. The latter contains transverse commissural fibers and is traversed by the pyramidal tracts. The superior or dorsal portion of the pons consists largely of gray matter which is continued up from below, especially the formation reticularis and the dorsal tracts of gray substance. In places the gray matter is grouped to form the nuclei of the fifth, sixth, seventh, and eighth nerves. In the lower part of the pons there is a small, folded gray mass, the superior olivary nucleus. It is enclosed by some transverse fibers known as the trapezium. The P. is connected with the medulla, with the cerebellum (by the middle peduncles), and with the cerebrum (by the crura cerebri).

Pontile, Pontine (*pon'-til*, *pon'-ten*) [*pons*, bridge]. Pertaining to the pons Varolii. **P. Hemiplegia**, a hemiplegia due to a lesion of the pons. When the lesion is situated low down in the pons, below the decussation of the fibers of the facial nerve, and above that of the pyramidal tracts, the hemiplegia is *alternate*, i.e., the arm and leg on one side, and the face on the other, are paralyzed.

Poplar (*pop'-lar*). See *Populus*.

Popliteal (*pop-lit'-e-al*) [*poples*, ham, hock, or knee]. Pertaining to or situated in the ham, as *P. artery*, *P. nerve*, *P. space*. **P. Aneurysm**, aneurysm of the popliteal artery. **P. Artery**. See *Arteries*, Table of.

Poppy (*pop'-e*) [AS., *popig*, poppy]. *Papaver somniferum*, a plant of the order *Papaveraceae*. **P. capsules** (*Papaveris capsulae*, B. P.) possess effects similar to those of opium, but far milder in degree. **Decoctum papaveris** (B. P.) is used chiefly as an anodyne application. **Extractum papaveris** (B. P.). Dose gr. v-x (0.32-0.65). **Syrupus papaveris** (B. P.). Dose [ʒss-ʒj] (2.0-4.0). **Red poppy** is used in making the *Syrupus rhæadus* of the B. P. Dose [ʒj] (4.0).

Populin (*pop'-u-lin*). See *Populus*.

Populus (*pop'-u-lus*) [L.]. Poplar, a genus of trees of the order *Salicinea*, several species of which contain salicin, $C_{10}H_{12}O_6$, and populin or benzoyl-salicin, $C_{20}H_{22}O_8$. Poplar-buds have been used in rheumatism and Bright's disease. The leaves and buds are also employed for anodyne ointments.

Porcupine-disease. See *Schthyosis*.

Pore (*por*) [*πόρος*, a pore]. A minute circular opening on a surface, as a P. of the skin, the opening of the duct of a sudoriparous gland.

Porencephalia, **Porencephalus** (*por-en-sef'-a'-le-ah*, *por-en-sef'-al-us*) [*πόρος*, cavity; *ἐνκεφαλος*, brain]. A condition characterized by the presence of depressions on the surface of the brain, due to a congenital arrest of development or to an acquired defect.

Porencephalous (*por-en-sef'-al-us*) [*πόρος*, cavity; *ἐνκεφαλος*, brain]. Affected with porencephalia.

Pornography (*por-nag'-ra-fe*) [*πόρνη*, a prostitute; *γράφειν*, to write]. 1. A treatise on prostitution. 2. Obscene writing.

Porocoele (*po'-ro-sel*) [*πίρος*, callus; *κύημα*, tumor]. A scrotal hernia in which the coverings are indurated and thickened.

Poroplastic (*por-o-plas'-tik*) [*πόρος*, pore; *πλάσσειν*, to mold]. Porous and plastic. **P. Felt**, a porous felt which is readily molded; it is used in the preparation of splints and jackets.

Porosity (*por-os'-i-t-e*) [*πόρος*, cavity]. The condition of being porous.

Porous (*po'-rus*) [*πόρος*, cavity, pore]. Having pores.

Porphyritization (*por-fir-i-za'-shun*) [*porphyry*, a kind of rock]. Pulverization, so-called because generally performed on a tablet of porphyry.

Porriigo (*por-i'-go*) [L.]. An old term applied to several diseases of the scalp. **P. decalvans**, alopecia areata. **P. favosa**, favus. **P. larvalis**, impetigo of the scalp conjoined with eczema.

Porro's Operation. See *Operations*, Table of.

Porta (*por'-tah*) [L., gate]. The hilus of an organ through which the vessels enter. **P. hepatis**, **P. jecoris**, the transverse fissure of the liver, through which the portal vein enters the organ.

Portal (*pôr'-tal*) [*porta*, gate]. Pertaining to the porta or hilus of an organ, especially to the porta hepatis, or to the vein entering at the porta hepatis (P. vein). P. Circulation, the passage of blood from the stomach, spleen, and intestines through the portal vein and the liver. P. Fissure. See *Fissure*. P. Vein, the large vein entering the liver at the transverse fissure, and bringing to it the blood from the digestive tract and the spleen.

Porte-caustique (*pört-kôs-ték'*). A holder for the stick of caustic.

Porter (*pôr'-ter*). See *Malt-liquors*.

Portio (*pört'-she-o*) [L.]. A part. P. dura, the hard part of the seventh pair of nerves of Willis, the facial nerve. P. inter duram et mollem, a small funiculus between the portio dura and the portio mollis of the seventh cranial nerve. P. intermedia Wrisbergii. Synonym of P. inter duram et mollem. P. mollis, the soft part of the seventh cranial nerve, the auditory nerve.

Port-wine Stains. See *Nevus*.

Porus (*pô'-rus*) [L.]. A pore. P. acusticus externus, the opening of the external auditory canal. P. acusticus internus, the opening of the internal auditory canal into the cranial cavity. P. opticus, the opening in the center of the lamina cribrosa of the sclerotic transmitting the central artery of the retina.

Position (*po-zish'-un*) [*ponere*, to place]. Place; location; attitude; posture. P., Dorsal, one in which the patient lies on the back. P., Edebohl's. See P., Simon's. P., English. See P., Left Lateral Recumbent. P., Knee-chest, P., Genupectoral; one in which the patient rests upon the knees and chest, the arms being crossed above the head. P., Knee-elbow, P., Genucubital; one in which the patient lies upon the knees and elbows, the head resting upon the hands. P., Left Lateral Recumbent, P., English, P., Obstetric; one in which the patient lies on the left side with the right thigh and knee drawn up. P., Lithotomy-, P., Dorsosacral; one in which the patient lies on the back with the legs flexed on the thighs, and the thighs flexed on the abdomen and abducted. P. of the Fetus, the relation of the presenting part of the fetus to the pelvis of the mother. See *Table of Fetal Positions*, under *Fetus*. P., Sim's, P., Semiprone; one in which the patient lies on the left side, with the right knee and thigh drawn up and the left arm placed along the back; the chest is inclined forward so that the patient rests upon it. P., Simon's, one in which the patient lies on the back, with the legs flexed on the thighs and the thighs on the abdomen, the hips being elevated and the thighs strongly abducted. P., Trendelenburg's, one in which

the patient lies on the back on a plane inclined about 45°, the feet and legs hanging over the end of the table.

Positive (*poz'-it-iv*) [*positivus*, from *ponere*, to place]. Real, existing, actual. In mathematics and physiology, denoting one of two quantities or conditions assumed as primary or fundamental; opposed to one assumed as negative; denoting a quantity greater than zero; to be added, additive. P. Electricity, the kind of electricity developed by rubbing glass with silk. P. Electrode, P. Pole; the electrode or pole connected with the negative plate of a battery. P., Element, P., Plate; that plate of a battery which is acted upon by the fluid, e. g., the zinc-plate in the zinc-carbon battery.

Posology (*pos-ol'-oj-e*) [*πῶσος*, how much; λόγος, science]. That branch of medical science dealing with the dosage of medicines.

Post- (*pöst-*) [L.]. A prefix denoting after.

Postanal (*pöst-a'-nal*) [*post*, after; *anus*]. Situated behind the anus.

Postaxial (*pöst-aks'-e-al*) [*post*, after; *axis*, axis]. Situated behind the axis; in the arm, behind the ulnar aspect; in the leg, behind the fibular aspect.

Postcentral (*pöst-sent'-tral*) [*post*, behind; *centrum*, center]. 1. Situated behind a center. 2. Situated behind the central fissure of the brain, as the P. convolution.

Postclavicular (*pöst-kla-vik'-u-lar*) [*post*, after; *clavicula*, clavicle]. Situated behind the clavicle.

Postconvulsive (*pöst-kon-vul'-siv*) [*post*, after; *convellere*, to tear away]. Coming on after a convulsion.

Postdiastolic (*pöst-di-as-tol'-ik*) [*post*, after; *διαστολή*, diastole]. Occurring after the diastole.

Postdicrotic (*pöst-di-krot'-ik*) [*post*, after; *dicrotic*]. Occurring after the dicrotic wave of the pulse, as the P. wave in the sphygmographic tracing.

Postdiphtheric (*pöst-dif-ther'-ik*) [*post*, after; *diphtheric*]. Occurring after an attack of diphtheria, as P. paralysis.

Postembryonic (*pöst-em-bre-on'-ik*) [*post*, after; *ἐμβρυον*, embryo]. Occurring after the embryonic stage.

Postepileptic (*pöst-ep-e-lep'-tik*) [*post*, after; *ἐπιληψία*, epilepsy]. Occurring after an epileptic attack.

Posterior (*pös-te'-re-or*) [comparative of *posterus*, after, behind]. Placed behind or to the back of a part.

Posteroexternal (*pös-ter-o-eks-tur'-nal*) [*posterus*, behind; *external*]. Occupying the outer side of a back part, as the P. column of the spinal cord.

Posterointernal (*pös-ter-o-in-tur'-nal*) [*post*, behind; *internal*]. Occupying the

inner side of a back part, as the P. column of the spinal cord.

Posterolateral (*pōs-te-ro-lat'-er-al*) [*posterus*, behind; *latus*, side]. Situated behind and at the side of a part.

Posteroparietal (*pōs-te-ro-par-i'-et-al*) [*posterus*, behind; *paries*, wall]. Situated at or opposite the posterior part of the parietal bone.

Posterosuperior (*pōs-te-ro-sū-pe'-re-or*) [*posterus*, hinder; *superior*, superior]. Situated behind and above a part.

Posterotemporal (*pōs-te-ro-tem'-por-al*) [*posterus*, behind; *tempus*, temple]. Situated at or opposite the posterior part of the temporal bone.

Postfebrile (*pōst-feb'-ril*) [*post*, after; *febris*, a fever]. Occurring after a fever.

Postglenoid (*pōst-gle'-noid*) [*post*, behind; *γλήνη*, socket; *ειδός*, like]. Situated behind the glenoid fossa, as the P. tubercle, also called simply P.

Posthemiplegic (*pōst-hem-e-ple'-jik*) [*post*, after; *hēmi*, half; *πληγή*, stroke]. Occurring after or following an attack of hemiplegia. P. Chorea, choreiform movements in paralyzed limbs.

Posthethomy (*pōs-thet'-o-me*) [*πρόθη*, penis; *τομή*, a cutting]. Circumcision.

Posthitis (*pōs-thi'-tis*) [*πρόθη*, prepuce; *itis*, inflammation]. Inflammation of the prepuce.

Postolith (*pōst'-tho-lith*) [*πρόθη*, prepuce; *λίθος*, stone]. A preputial calculus.

Posthumous (*pōst'-tu-mus*) [*post*, after; *humus*, the ground]. Occurring after death. P. Child, one born after the death of the father, or taken by the Cesarean operation from the body of its mother after death.

Posthyoid (*pōst'-hi'-oid*) [*post*, behind; *ὠψήθης*, hyoid]. Posterior to the hyoid bone.

Posthypnotic (*pōst-hip-not'-ik*) [*post*, after; *ἵπνως*, sleep]. Succeeding the hypnotic state; acting after the hypnotic state has passed off, as P. suggestion.

Posticus (*pōs-ti'-kus*) [L.]. Posterior.

Postmortem (*pōst-mor'-tem*) [*post*, after; *mors*, death]. 1. Occurring after death. 2. An examination of the body after death; an autopsy.

Postnasal (*pōst-na'-s al*) [*post*, behind; *nasus*, nose]. Situated behind the nose. P. Catarrh, catarrhal inflammation of the nasopharynx.

Postparalytic (*pōst-par-ah-lit'-ik*) [*post*, after; *παράλυσις*, paralysis]. Following an attack of paralysis.

Postpartum (*pōst-par'-tum*) [*post*, after; *partus*, birth]. Following childbirth; as P. hemorrhage.

Postpharyngeal (*pōst-far-in'-je-al*) [*post*, behind; *φάρυγξ*, pharynx]. Situated behind the pharynx.

Postpone (*pōst-pōn'*) [*post*, after; *ponere*, to place]. Of a paroxysm, to occur after the regular time.

Postpyramidal (*pōst-pe-ram'-id-al*) [*post*, behind; *pyramis*, pyramid]. Situated behind the pyramidal tract. P. Nucleus, the nucleus funiculi gracilis.

Postural (*pōst-tur-al*) [*ponere*, to place]. Pertaining to posture or position; performed by means of a special posture, as P. treatment.

Postuterine (*pōst-ut'-er-in*) [*post*, behind; *uterus*, uterus]. Situated behind the uterus.

Potable (*pō'-ta-bl*) [*potare*, to drink]. Drinkable; fit to drink.

Potash (*pōt'-ash*) [from Irish, *pot*, a pot, and *ash*]. 1. Potassium hydroxid, caustic potash. 2. Potassium carbonate.

Potassa (*pō-tas'-uh*) [L.]. Potash. P. cum calce, Vienna-paste, a caustic paste made by rubbing together equal parts of potassium hydroxid and quick-lime. P. sulphurata, a mixture of salts of potassium, consisting chiefly of the sulphid. It is used in chronic rheumatism, gout, and locally in diseases of the skin. Dose gr. ij-x (0.13-0.65).

Potassium (*pō-tas'-e-um*) [*potash*, from Irish, *pot*, a pot, and *ash*]. A silver-white, soft, very ductile metal, belonging to the alkali-group, and having a sp. gr. of 0.865; its valence is one, atomic weight 39, symbol K (from the Latin *kalium*). It forms compounds with nearly all nonmetals. The most important compounds are: P. acetate (Potassii acetas, U. S. P., B. P.), $K_2H_3O_2$, used in rheumatism, as a diuretic in dropsies, and in cutaneous diseases. Dose gr. xx-5j (1.3-4.0). P. bicarbonate (Potassii bicarbonas, U. S. P., B. P.), $KHCO_3$, used as an antacid. Dose gr. xx-5j (1.3-4.0). P. dichromate, P. bichromate (Potassii bichromas, U. S. P., B. P.), $K_2Cr_2O_7$, used as a caustic, and for the preparation of battery-fluids and of preserving fluids for tissues; it was formerly employed as an alternative. P. bitartrate (Potassii bitartaras, U. S. P., B. P.), cream of tartar, $KHC_4H_4O_6$, used as a mild cathartic and as a hydragogue diuretic in dropsies, and for making refrigerant drinks in febrile affections. It is an ingredient of Pulvis jalapæ compositus. Dose 5j-5ss (4.0-16.0). P. bromid. See Bromin. P. carbonate (Potassii carbonas, U. S. P., B. P.), used as an antacid in dyspepsia and as an antilithic. P. chlorate (Potassii chloras, U. S. P., B. P.), $KClO_3$, used locally and internally in various forms of stomatitis, in mercurial ptyalism, in pharyngitis. Dose gr. x-xx (0.65-1.3). In over-doses it is an irritant poison. From it are prepared Trochisci potassii chlo-

ratis. **P. citrate** (Potassii citras, U. S. P., B. P.), $K_2C_6H_5O_7 \cdot H_2O$, used as a diaphoretic in fevers, in bronchitis, and in rheumatism, the uric acid diathesis, etc. Dose gr. xx-xxx (1.3-2.0). Potassii citras effervescens (U. S. P.) is used like the preceding. Dose $\frac{3}{4}$ -ij (4.0-8.0). **P. cyanid** (Potassii cyanidum, U. S. P., B. P.), KCN, is similar in properties to hydrocyanic acid. Dose gr. $\frac{1}{4}$ (0.008). It is also used locally in neuralgia. **P. ferrocyanid** (Potassii ferrocyanidum, U. S. P., B. P.), $K_4Fe(CN)_6 \cdot 3H_2O$, yellow prussiate of potash, has been used in the night-sweats of phthisis. Dose gr. x-xv (0.65-1.0). It is extensively employed in the arts, as a reagent, and in pharmacy to prepare diluted hydrocyanic acid. **P. hydroxid**, KOH (Potassa, U. S. P., B. P.), caustic potash. This is a white solid, usually occurring in the form of pencils, and having powerful escharotic properties. From it are prepared, Liquor potasse and Potassa cum calce. The former is an antacid, diuretic, and antilithic. Dose mxx-xxx (0.65-2.0). **P. hypophosphite** (Potassii hypophosphis, U. S. P.), KH_2PO_2 . Dose gr. x-xxx (0.65-2.0). See *Hypophosphite*. **P. iodid** (Potassii iodidum, U. S. P., B. P.), KI, is used as an alterative in syphilis; in chronic poisoning by lead and mercury; as an absorbent of inflammatory exudates; in chronic rheumatism, etc. Dose gr. ij-x (0.13-0.65); in syphilis several drams may be administered daily. From it are prepared Unguentum iodi and Unguentum potassii iodidi. **P. nitrate** (Potassii nitras, U. S. P., B. P.), KNO_3 , saltpeter, is used as a refrigerant diuretic and diaphoretic; in asthma the inhalation of fumes produced by burning niter-paper (paper dipped in a solution of the nitrate and dried) is useful. Dose gr. x-xx (0.65-1.3). From it are prepared Argenti nitras dilutus and Charta potassii nitratis. **P. permanganate** (Potassii permanganas, U. S. P., B. P.), $KMnO_4$, is used as an antiseptic and deodorant, in amenorrhea, and as an antidote to opium-poisoning and poisoning by snake-bites. Dose as an emmenagogue gr. j-ij (0.065-0.13). It is also employed as a reagent. Condry's fluid is an aqueous solution of P. permanganate (2 parts in 100). **P. and sodium tartrate** (Potassii et sodii tartras, U. S. P., B. P.), $KNaC_4H_4O_6 \cdot 4H_2O$, Rochelle salt, is used as a saline cathartic; it is an ingredient of Seidlitz powder. Dose $\frac{3}{4}$ ss-j (16.0-32.0). **P. sulphate** (Potassii sulphas, U. S. P., B. P.), K_2SO_4 , is used as a laxative and purgative. Dose $\frac{3}{4}$ ss- $\frac{3}{4}$ iv (2.0-16.0). **P. tartrate** (Potassii tartras, B. P.), $K_2C_4H_4O_6 \cdot H_2O$, is used as a mild purgative in febrile affections. Dose $\frac{3}{4}$ j- $\frac{3}{4}$ j (4.0-32.0). Other compounds are: **P. bin-**

oxalate, salt of sorrel, a constituent of wood-sorrel. **P. bisulphate**, $KHSO_4$, has been used as a laxative and refrigerant. Dose $\frac{3}{4}$ j-ij (4.0-8.0). **P. cobaltinitrate** has been employed in asthma, nephritis, and valvular heart-disease. Dose gr. $\frac{1}{4}$ - $\frac{1}{2}$ (0.016-0.032). **P. ferricyanid**, $K_3Fe(CN)_6$, is used in the arts and as a reagent. **P. iodate** is used in stomatitis and pharyngitis as a substitute for P. chlorate. Dose gr. iv-vij (0.26-0.52). **P. nitrite**, KNO_2 , is used like the other nitrites. Dose gr. iij (0.2). **P. phosphate**, K_2HPO_4 , has been used as an alterative in scrofula and pulmonary tuberculosis. Dose gr. x-xxx (0.65-2.0). **P. silicate**, soluble glass, is used instead of plaster of Paris for making fixed dressings. **P. sulphocyanate**, KCNS, is a constituent of saliva.

Potato-culture. A culture of microorganisms on potato.

Potential (*po-ten'-shal*) [*potens*, able]. 1. Capable of acting or doing work, as P. energy. 2. In electricity, a state of tension or of P. energy capable of doing work. If two bodies of different P. are brought together, a current is established between them that is capable of doing work.

Potio, **Potion** (*po'-she-o*, *po'-shun*) [*potio*, a drink]. A drink or draught.

Potomania (*po-to-ma'-ne-ah*) [*πῶτος*, drink; *μανία*, madness]. Delirium tremens.

Pott's Disease [Percival Pott, an English surgeon]. Caries of the vertebrae, generally of tuberculous origin. The symptoms are stiffness of the spinal column, pain on motion, tenderness on pressure, undue prominence of one or more of the spines; in certain cases spasmodic pain in the abdomen; formation of abscess; occasionally, in late stages, paralysis. **P.'s Fracture.** See *Fracture*.

Pouch [*Fr.* *poché*, a pocket]. A sac or pocket. **P. of Douglas**, **P., Recto-uterine**, cul-de-sac of Douglas, a pouch formed by a fold of the peritoneum between the uterus and the rectum.

Poultice (*pol'-tis*) [*puls*, porridge]. A soft, semiliquid mass made of some cohesive substance mixed with water, and used for application to the skin, for the purpose of supplying heat and moisture or acting as a local stimulant. Synonym, *Cataplasma*.

Pound [*AS.* *pund*, pound]. A unit of measure of weight. The Troy P. contains 12 oz. or 5760 grains; the avoirdupois P. contains 16 oz. or 7000 grains. Symbol lb. See *Weights and Measures*. **P., Foot-**, the force necessary to raise one pound through the height of one foot.

Poupart's Ligament. See *Ligaments*, *Table of*.

Powder [Fr., *poudre*, from *pulvis*, powder].

1. A collection of fine particles. 2. In pharmacy, a collection of fine particles of one or more substances capable of passing through a sieve having meshes of a certain fineness; also a single dose of such a substance. **P.**, Aromatic. See *Cinnamon*. **P.**, Dover's, *Pulvis ipecacuanhæ et opii*. See under *Opium*. **P.**, James'. See *Antimony*. **P.**, Seidlitz-. See *Potassium and sodium tartrate*. **P.**, Tully's, a powder composed of camphor, prepared chalk, and licorice, and morphin sulphate, 1 part. It is used as a substitute for Dover's powder.

Pox [*pocks*, pl. of *pock*, a vesicle or pustule]. 1. A term applied to any disease possessing a vesicular or pustular eruption. 2. Vulgarly, syphilis. **P.**, Chicken-. See *Varicella*. **P.**, Cow-. See *Cow-pox*. **P.**, Small-. See *Varicella*.

Practice (*prak'-tis*) [*practicare*, to practice]. The practice of medicine; the application of the principles of medicine to the diagnosis and treatment of disease.

Practitioner (*prak-tish'-un-er*) [*practicare*, to practice]. One who practices medicine. (This is a badly-formed word.)

Præ- (*pre-*) [*l.*]. See *Pre-*.

Præputium (*pre-pu'-she-un*). See *Prepuce*.

Prævia (*pre'-ve-ah*) [sem. of *prævi*, from *præ*, before; *via*, a way]. Coming before, standing before. **P.**, Placenta. See *Placenta*.

Prairie-itch. A name applied to various forms of skin-diseases associated with pruritus, occurring in men engaged in work on prairies, among lumbermen, and others, and either due to scabies or pruritus hiemalis.

Prealbuminuric (*pre-al-bu-min-ur-ik*) [*præ*, before; *albumen*, albumin; *urina*, urine]. Occurring before the appearance of albuminuria.

Preataxic (*pre-at-aks'-ik*) [*præ*, before; *á priv.*, *τάξις*, order]. Occurring before ataxia.

Preaxial (*pre-aks'-e-al*) [*præ*, before; *axis*, axis]. Situated in front of the axis; in a limb, situated on the internal or anterior aspect—in the arm, on the radial, in the leg, on the tibial side.

Prebacillary (*pre-bas'-il-a-re*) [*præ*, before; *bacillus*, bacillus]. Occurring before the invasion of the system by bacilli.

Precancerous (*pre-kan'-ser-us*) [*præ*, before; *cancer*]. Occurring before the development of a carcinoma.

Precardiac (*pre-kar'-de-ak*) [*præ*, before; *kardia*, heart]. Anterior to the heart.

Precentral (*pre-sen'-tral*) [*præ*, before; *central*, center]. Situated in front of the central fissure of the brain. **P.** *Convolution*, a convolution in front of the central

fissure of the brain; the ascending frontal convolution.

Precipitant (*pre-sip'-it-ant*). See under *Precipitation*.

Precipitate (*pre-sip'-it-at*) [*precipitare*, from *præ*, before; *caput*, head]. The solid substance thrown down from a solution of a substance on the addition of a reagent which deprives it of its solubility. Abbreviation: *Ppt.* **P.**, Red, *Hydrargyrum oxidum rubrum*. See *Mercury*. **P.**, White, $N_2(Hg_2)Cl_2$, *Hydrargyrum ammoniatum*. See *Mercury*.

Precipitate (*pre-sip'-it-at*) [*precipitare*, from *præ*, before; *caput*, head]. To throw down in an insoluble form.

Precipitate (*pre-sip'-it-at*) [*precipitare*, from *præ*, before; *caput*, head]. Headlong; hasty. **P.** Labor. See *Labor*.

Precipitation (*pre-sip-it-a'-shun*) [*precipitare*]. The process of throwing down solids from the liquids which hold them in solution. Precipitates are crystalline, amorphous, curdy, flocculent, granular, or gelatinous, according to the form assumed. The agent causing precipitation is the *precipitant*, the solid thrown down, the *precipitate*.

Precordia (*pre-kor'-de-ah*) [*præ*, before; *cor*, the heart]. 1. A name applied by the ancients to the diaphragm, the thoracic viscera, or the epigastric region. 2. The area of the chest overlying the heart.

Precordial (*pre-kor'-de-al*) [*præ*, before; *cor*, heart]. Pertaining to the precordia.

Precordium (*pre-kor'-de-un*). Singular of *Precordia*.

Precuneus (*pre-ku-ne'-us*) [*præ*, before; *cuneus*, wedge]. The quadrate lobule of the parietal lobe situated in front of the cuneus of the occipital lobe.

Predialstolic (*pre-di-as-to'-lik*) [*præ*, before; *διαστολή*, diastole]. Occurring before the diastole.

Predicrotic (*pre-di-krot'-ik*) [*præ*, before; *dicrotic*]. Preceding the dicrotic wave or elevation of the sphygmographic tracing.

Predigest (*pre-di-jest'*) [*præ*, before; *digerere*, to digest]. To digest by artificial means before introducing into the body.

Predisposing (*pre-dis-po'-sing*) [*præ*, before; *disponere*, to dispose]. Rendering susceptible or liable to attack.

Predisposition (*pre-dis-po-zish'-un*) [*præ*, before; *disponere*, to dispose]. The state of having a susceptibility to disease.

Prefrontal (*pre-frun'-tal*) [*præ*, before; *frons*, forehead]. Situated in the anterior part of the frontal lobe of the brain.

Preglobulin (*pre-glob'-u-lin*) [*præ*, before; *globulin*]. An albuminous body found in cell-protoplasm, insoluble in water, soluble in a ten per cent. sodium chlorid solution and in dilute alkaline solutions.

ELY'S TABLE OF THE DURATION OF PREGNANCY.

EXPLANATION.—Find in the upper horizontal row the date of last menstruation; the figure beneath will show the expiration of 280 days or ten months of 28 days each.

January.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	November.
February.	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	1	2	3	4	5	6	7	
March.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	December.
April.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	January.
May.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	February.
June.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	March.
July.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	April.
August.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	May.
September.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	June.
October.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	July.
November.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	August.
December.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	September.
January.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	October.

PREGNANCY

519

PREMALIGNANT

Pregnancy (*preg'-nan-se*) [*pregnans*, from *pra*, before; *gignere*, to beget]. The state of being pregnant, or with child; the state of the woman from conception to the expulsion of the ovum. The normal duration of P. is 280 days, or ten lunar months, or nine calendar months. P., **Extrauterine**, ectopic gestation; the development of the ovum outside of the cavity of the uterus. Pregnancy may take place within the Fallopian tube (**Tubal P.**), in the ovary (**Ovarian P.**), in the part of the tube that traverses the uterine wall (**Interstitial P.**), or in the abdominal cavity (**Abdominal P.**). **P., False**, any condition in the abdomen that simulates pregnancy. P., **Molar**, pregnancy in which the ovum is converted into a mole. P., **Multiple**, that form in which the uterus contains two or more developing ova. P., **Mural**, a variety of extrauterine P in which the ovum develops in the wall of the uterus. P., **Signs of**, those clinical manifestations by which the existence of pregnancy may be demonstrated. There are three so-called absolute signs: Ballotement, fetal movements, and the fetal heart-sounds.

Pregnant (*preg'-nant*) [*pragnans*, from *pra*, before; *gignere*, to beget]. With child; gravid.

Prehemiplegic (*pre-hem-e-pleg'-ik*) [*pra*, before; *hmi*, half; *πληγῆ*, stroke]. Occurring before an attack of hemiplegia, as P. chorea.

Prehensile (*pre-hen'-sil*) [*prehendere*, to lay hold of]. Adapted for grasping.

Prehension (*pre-hen'-shun*) [*prehendere*, to lay hold of]. The act of grasping or seizing.

Prelum (*pre'-lum*) [L.]. Press. P. abdominale, the squeezing of the abdominal viscera between the diaphragm and the rigid abdominal wall, as in the processes of defecation, micturition, and parturition.

Premalignant (*pre-mal-ig'-nant*) [*pra*, before; *malign-*

- nant*]. Occurring before the development of malignancy.
- Premature** (*prel'-mat-ār*) [*præ*, before; *maturare*, to ripen]. Occurring before the proper time, as P. labor.
- Premaxilla** (*pre-maks-il'-ah*) [*præ*, before; *maxilla*, jaw]. The intermaxillary bone.
- Premenstrual** (*pre-men'-stru-al*) [*præ*, before; *menstrual*]. Preceding menstruation.
- Premolar** (*pre-mo'-lar*) [*præ*, before; *mola*, millstone]. 1. Situated in front of the molar teeth. 2. One of the two anterior permanent molars.
- Premonitory** (*pre-mon'-it-o-re*) [*præ*, before; *monere*, to warn]. Forewarning; foreboding, as P. symptoms, those which forebode the onset of an attack of a disease.
- Prenatal** (*pre-na'-tal*) [*præ*, before; *natus*, born]. Existing or occurring before birth.
- Preoccupation** (*pre-ok-u'-pa'-shun*) [*præ*, before; *occupare*, to occupy]. The condition of being so engrossed in one's own thoughts as not to answer or hear when accosted.
- Preparation** (*prep-ar-a'-shun*) [*præparare*, to make ready]. 1. The act of making ready. 2. Anything made ready, especially, in anatomy, any part of the body prepared or preserved for illustrative or other uses.
- Prepatellar** (*pre-pat-el'-ar*) [*præ*, before; *patella*, patella]. Situated in front of the patella, as P. bursa.
- Prepuce** (*pre'-piūs*) [*præputium*, prepuce]. The foreskin of the penis, a fold of skin lined by mucous membrane, and covering the glans penis.
- Preputial** (*pre-pu'-she-al*) [*præputium*, prepuce]. Pertaining to the prepuce.
- Presbycusis, Presbykousis** (*prez-be-koo'-sis*) [*πρεσβυς*, old; *ἀκοῦειν*, to hear]. The lessening of the acuteness of hearing that occurs in old age.
- Presbyope** (*prez'-be-ōp*) [*πρεσβυς*, old; *ὤψ*, eye]. One who is presbyopic.
- Presbyopia** (*prez-be-ō'-pe-ah*) [*πρεσβυς*, old; *ὤψ*, eye]. The condition of vision in the aged, due to diminished power of accommodation from impaired elasticity of the crystalline lens, whereby the near point of distinct vision is removed farther from the eye.
- Presbyopic** (*prez-be-ōp'-ik*) [*πρεσβυς*, old; *ὤψ*, eye]. Affected with presbyopia.
- Prescription** (*pre-skript'-shun*) [*præ*, before; *scribere*, to write]. A formula written by the physician to the apothecary, designating the substances to be administered. A P. consists of the heading, usually the symbol R (standing for the Latin word *recipe*, imperative of *recipere*, to take), the names and quantities of the ingredients, the directions to the apothecary, the directions to the patient, the date, and the signature.
- Present** (*pre-sent'*) [*presentare*, to place before]. Of a part of the fetus, to appear first at the os uteri.
- Presentation** (*pre-sen-ta'-shun*) [*presentare*, to place before]. In obstetrics, that part of the fetal body which presents itself to the examining finger at the os uteri.
- Presphenoid** (*pre-sfel'-noid*) [*præ*, before; *σφήν*, wedge; *ειδός*, like]. The anterior part of the body of the sphenoid bone.
- Pressor** (*pres'-or*) [*premere*, to press]. Stimulating. P. Nerve. See *Nerve*.
- Pressure-sense**. The sense by which knowledge is obtained of the amount of weight or pressure which is exerted upon a part of the body.
- Presterium** (*pre-ster'-num*) [*præ*, before; *sternum*, the breast-bone]. The manubrium or superior segment of the sternum.
- Presystolic** (*pre-sis-tol'-ik*) [*præ*, before; *συστολή*, systole]. Preceding the systole of the heart, as the P. murmur, P. thrill.
- Pretibial** (*pre-tib'-e-al*) [*præ*, before; *tibia*, tibia]. Situated in front of the tibia.
- Pretuberculous** (*pre-tu-ber'-ku-lus*) [*præ*, before; *tuberculum*, tubercle]. Preceding the development of tuberculosis.
- Preventive** (*pre-ven'-tiv*) [*prevenire*, to anticipate, to prevent]. Warding off. P. Medicine, the department of medicine dealing with the means and methods of preventing disease.
- Prevertebral** (*pre-ver'-te-bral*) [*præ*, before; *vertebra*, vertebra]. Situated in front of the vertebrae.
- Priapism** (*pri'-ap-izm*) [*πριαπισμός*]. Persistent erection of the penis, usually unaccompanied by sexual desire. It is caused by injuries to the spinal cord, or to the penis, and by vesical calculus.
- Prickle-cell**. A cell possessing delicate rod-shaped processes by which it is connected with neighboring cells. Such cells are found in the middle strata of the skin (**Prickle-layer**).
- Prickly Heat**. See *Miliaria*.
- Primæ viæ** (*pri'-me vi'-e*) [L., "the primary passages"]. The alimentary canal.
- Primary** (*pri'-ma-ry*) [*primus*, first]. First in time or in importance. P. Amputation, one that is done before the development of inflammation, usually within the first twenty-four hours. P. Bubo, a simple adenitis of an inguinal lymphatic gland, resulting from mechanical irritation. It is also known as *bubon d'emblée*. P. Dementia, a form of insanity occurring in young adults, and characterized by an extreme degree of apathy, the patient lying motionless, absolutely listless, without wants and seemingly without perception of his surroundings. P. Lesion, the original lesion which forms the starting-point

for secondary lesions. **P. Sore**, the initial sclerosis or chancre of syphilis.

Primigravida (*pri-me-grav'-id-ah*) [*primus*, first; *gravidus*, pregnant]. A woman pregnant for the first time.

Primipara (*pri-mip'-ar-ah*) [*primus*, first; *parere*, to bear]. A woman bearing or giving birth to her first child.

Primiparous (*pri-mip'-ar-us*) [*primus*, first; *parere*, to bear]. Bearing a child or being in labor for the first time.

Primitive (*prim'-it-iv*) [*primus*, first]. First-formed; original. **P. Groove**, the enlargement and deepening of the primitive streak. See **P. Streak**. **P. Streak**, **P. Trace**, a streak appearing at the end of the germinal disc, and indicating the first trace of the embryo.

Primordial (*pri-mor'-de-al*) [*primus*, first; *ordiri*, to arise]. Existing in the beginning, first-formed, primitive, original, of the simplest character. **P. Kidney**, the Wolfian body. **P. Ova**, cells lying among the germ-epithelium of the surface of the embryonic ovary.

Princeps (*prin'-seps*) [*primus*, first; *caput*, head]. First, original. **P. cervicis**, a branch of the occipital artery supplying the trapezius muscle. **P. pollicis**, a branch of the radial artery, going to the palmar surface of the thumb.

Principle (*prin'-si-pl*) [*principium*, from *princeps*]. A constituent of a compound having a definite composition and representing its essential or characteristic properties. **P.**, Proximate. See *Proximate*.

Prinos (*pri'-nos*) [*L.*]. Black alder, a shrub of the order Aquifoliaceæ. The bark and the berries of *P. verticillatus* have been employed as tonics and astringents in diarrhea, and locally, as an application to ulcers. Dose ʒ ss-j (2.0-4.3); of a decoction f ʒj-ij (32.0-64.0). Unof.

Prism (*prizm*) [*πρίσμα*, from *πρίξτιν*, to saw]. A solid whose bases or ends are similar plane figures and whose sides are parallelograms. In optics, a transparent solid with triangular ends and two converging sides. It breaks up white light into its component colors and bends the rays of light toward the side opposite the angle (the base of the **P.**), and is used to measure or correct imbalance of the ocular muscles. **P.-diopter**, **P.-dioptry**, a unit of prismatic refractive power: the refractive power of a prism that deflects a ray of light one centimeter on a tangent plane situated at a distance of one meter. **P.**, **Enamel-**, one of the prismatic columns of from four to six sides composing the enamel of teeth, closely packed together and generally vertical to the surface of the underlying dentine. **P.**, **Nicol-**. See *Nicol-prism*.

Prismatic (*pris-mat'-ik*) [*πρίσμα*, a prism]. Prism-shaped; produced by the action of a prism, as **P. colors**.

Prismometer (*pris-mop-tom'-et-er*) [*πρίσμα*, prism; *ὤψ*, eye; *μέτρον*, measure]. An instrument for estimating refractive defects of the eye by means of two prisms placed base to base.

Pro- [*L.*]. A prefix meaning before.

Probang (*pro'-bang*) [*probare*, to test]. A rod of whalebone or other flexible material used for making local applications to the esophagus or larynx, or removing foreign bodies. **P.**, **Ball-**, a probang having an ivory bulb attached to one end. **P.**, **Horsehair-**, **P.**, **Bristle-**, one having on the end a sheath of horsehair or bristles that can be made to spread like an umbrella as the instrument is drawn out. **P.**, **Sponge-**, one provided with a small sponge at one end.

Probe (*prōb*) [*probare*, to test]. A slender, flexible rod for exploring a channel. **P.**, **Anel's**, a delicate probe for exploring or dilating the lacrimal puncta and lacrimal canals. **P.**, **Blunt**, one with a blunt extremity. **P.**, **Bowman's**, one of a series of graduated probes used for dilating the nasal duct. **P.**, **Drum-**, one provided with a drum or reverberator to enable the ear to detect contact with foreign bodies. **P.**, **Electric**, one having two insulated wires, so that contact with a bullet or metal completes the circuit, and thus indicates the presence of such a foreign body. **P.**, **Eyed**, a probe having a slit at one end through which a tape or ligature can be passed. **P.**, **Lacrimal**, one used for dilating the lacrimal passages. **P.**, **Meerschm-**, a probe, the end of which is tipped with meerschm, which becomes streaked with the lead by contact with a bullet. **P.**, **Nélaton's**, one that is capped with unglazed porcelain upon which a leaden ball makes a metallic streak. **P.**, **Vertebrated**, one made of small links so jointed together that it is very flexible.

Process (*pros'-es*) [*processus*, from *procedere*, to go]. 1. A course of action; a group of phenomena, as the inflammatory **P.** 2. A prominence or outgrowth, as the spinous **P.** of a vertebra, the axis-cylinder **P.** of a nerve. **P.**, **Alveolar**. See *Alveolar*. **P.**, **Auditory**, the curved plate of bone surrounding the external auditory meatus for the greater part of its circumference, and serving for the attachment of the cartilage of the external ear. **P.**, **Basilar**. See *Basilar*. **P.**, **Ciliary**. See *Ciliary*. **P.**, **Conoid**. See *Conoid Tubercle*. **P.**, **Coracoid**. See *Coracoid*. **P.**, **Coronoid**. 1. A thin, flattened process projecting from the anterior portion of the upper border of the ramus of the lower jaw, and serving for the attachment of the temporal

muscle. 2. A triangular projection from the upper end of the ulna, forming the lower part of the greater sigmoid cavity. **P., Deiters'**, the axis-cylinder process of a nerve-cell. **P., Dendritic.** See *P., Protoplasmic.* **P., Ensiform**, the cartilaginous tip at the lower end of the sternum. **P., Ethmoid**, one of the projections from the superior border of the inferior turbinated bone. **P., External Angular**, the projection on the outer extremity of the supraorbital arch of the frontal bone. **P., Falciform: 1.** A process of the fascia lata of the thigh, forming the outer and upper margin of the saphenous opening. It is known also as the *Falciform P. of Burns*. 2. The falx cerebri. **P., Hamular: 1.** A hook-like process of bone on the lower extremity of the internal pterygoid plate, around which the tendon of the tensor palati turns. 2. Of the lacrimal bone, the hook-like termination of the lacrimal crest. **P., Internal Angular**, the inner extremity of the supraorbital arch of the frontal bone. **P., Jugular.** See *Jugular.* **P., Lacrimal**, a short process of the inferior turbinated bone that articulates with the lacrimal bone. **P., Long (of the Incus)**, a slender process that descends vertically from the body of the incus and articulates, by the lenticular process, with the head of the stapes. **P., Long (of the Malleus)**, a long delicate process that passes from the neck of the malleus outward to the Glaserian fissure, to which it is connected by cartilaginous and ligamentous fibers. **P., Malar**, a triangular eminence of the superior maxilla by which it articulates with the malar bone. **P., Mammillary**, one of the tubercles on the posterior part of the superior articular processes of the lumbar vertebrae. **P., Mastoid.** See *Mastoid.* **P., Maxillary**, a thin plate of bone descending from the ethmoid process of the inferior turbinated bone, and hooking over the lower edge of the orifice of the antrum. **P., Nasal (of the Superior Maxilla)**, a thick, triangular process of bone that projects upward, inward, and backward by the side of the nose, forming a part of its lateral wall. **P., Odontoid**, the tooth-like process of the axis which ascends and articulates with the atlas. **P., Olecranon**, the olecranon. **P., Olivary**, a small oval eminence situated behind the optic groove of the sphenoid bone. **P., Orbital (of the Palate Bone)**, a process directed upward and outward from the upper portion of the palate bone. **P., Orbital (of the Superior Maxilla)**, a process projecting from the orbital margin of the superior maxilla. **P., Palate**, a thick process projecting horizontally inward from the inner surface of the superior maxillary bone, and forming part of the floor of the nostril and the

roof of the mouth. **P., Postglenoid**, a small tubercle separating the glenoid fossa from the auditory process. **P., Protoplasmic. 1.** Any one of the processes of nerve-cells that are not continued as axis-cylinders. 2. A pseudopod of an ameboid cell. **P., Pterygoid (of the Palate Bone)**, a pyramidal process projecting from the posterior border of the palate bone and articulating with the sphenoid bone. **P., Pterygoid (of the Sphenoid Bone)**, one descending perpendicularly from the point of junction of the body with the greater wing of the sphenoid bone, and consisting of an external and internal plate. **P. of Rau.** Synonym of *P., Long (of the Malleus)*. **P., Short (of the Incus)**, a conic process projecting almost horizontally backward from the body of the incus and attached by ligamentous fibers to the margin of the opening leading into the mastoid cells. **P., Short (of the Malleus)**, a slight projection from the root of the manubrium of the malleus, lying in contact with the tympanic membrane. **P., Sphenoidal**, a thin plate of bone directed upward and inward from the vertical plate of the palate bone. **P., Spinous (of the Ilium)**, a prominent eminence on the anterior or posterior border of the ilium. The upper process on the anterior border is the *anterior superior spinous process*; below it is the *anterior inferior spinous process*. On the posterior border there are also two processes—a *posterior superior* and a *posterior inferior*. **P., Spinous (of the Sphenoid Bone)**, a rough prominence descending from the posterior part of the greater wing of the sphenoid bone. It receives the attachment of the internal lateral ligament of the jaw and the tensor palati muscle. **P., Spinous (of the Tibia)**, an eminence of bone on the upper surface of the tibia, between the two articular surfaces, and nearer to the posterior than the anterior border. **P., Spinous (of a Vertebra)**, the prominent backward projection from the middle of the posterior portion of the arch of a vertebra. **P., Styloid (of the Fibula)**, a pointed eminence projecting upward from the posterior portion of the head of the fibula. **P., Styloid (of the Radius)**, a projection from the external border of the lower extremity of the radius. **P., Styloid (of the Temporal Bone)**, a sharp spine about an inch in length descending downward, forward, and inward from the inferior surface of the petrous portion of the temporal bone. **P., Styloid (of the Ulna)**, a projection from the inner and posterior portion of the lower extremity of the ulna. **P., Superior Maxillary**, an eminence on the face of the embryo which gives rise to the superior maxilla and the malar bone. **P., Superior Vermiform**, the upper part of the median lobe of the cere-

bellum, connecting the two lateral hemispheres. **P.**, **Temporal**, the posterior angle of the malar bone by which it articulates with the zygomatic process of the temporal bone. **P.**, **Transverse**, a process projecting outward from the side of a vertebra, at the junction of the pedicle and the lamina. **P.**, **Unbranched**. Synonym of **P.**, **Axis-cylinder**. **P.**, **Unciform (of the Ethmoid Bone)**, a hook-like projection from the inferior portion of each lateral mass of the ethmoid bone. It articulates with the ethmoidal process of the inferior turbinated bone. **P.**, **Unciform (of the Hippocampal Gyrus)**, a hook-like projection from the anterior end of the gyrus hippocampi. **P.**, **Unciform (of the Unciform Bone)**, a hook-like projection from the palmar surface of the unciform bone. **P.**, **Vaginal (of Peritoneum)**, the process of peritoneum which the testicle in its descent carries in advance, and which in the scrotum forms the tunica vaginalis testis. **P.**, **Vaginal (of the Sphenoid Bone)**, a projection from the inferior surface of the body of the sphenoid bone, running horizontally inward from near the base of the pterygoid process. **P.**, **Vaginal (of the Temporal Bone)**, a sheath-like plate of bone which extends from the carotid canal to the mastoid process. It separates behind into two laminae that enclose the styloid process. **P.**, **Xiphoid**, the ensiform cartilage. **P.**, **Zygomatic (of the Malar Bone)**, a long, serrated process which articulates with the zygomatic process of the temporal bone. **P.**, **Zygomatic (of the Temporal Bone)**, a long projection from the lower part of the squamous portion of the temporal bone, articulating with the malar bone. **Processus (pro-ses'-us)** [L.]. A process. **P. ad cerebrum**, the superior cerebellar peduncle. **P. ad medullam**, the inferior peduncle of the cerebellum. **P. ad pontem**, the middle peduncle of the cerebellum. **P. ad testes**, the superior cerebellar peduncles. **P. clavatus**, a thickening on the posterior pyramid of the medulla near the apex of the fourth ventricle. **P. cochleariformis**, a thin plate of bone separating the canal for the Eustachian tube from that for the tensor tympani muscle. **P. e cerebello ad cerebrum**, the superior cerebellar peduncle. **P. e cerebello ad medullam**, the inferior cerebellar peduncle. **P. e cerebello ad pontem**, the middle cerebellar peduncle. **P. e cerebello ad testes**, the superior peduncles of the cerebellum. **P. gracilis**, a long delicate process passing from below the neck of the malleus to the Glaserian fissure. **P. hamatus**. Synonym of *Process, Unciform*.

Procidencia (pro-se-den'-she-ah) [*pro*, before; *cadere*, to fall]. Prolapse.

Procreation (pro-kre-a'-shun) [*procreare*, to bring forth]. The act of begetting offspring.

Proctalgia (prok-tal'-je-ah) [*πρωκτός*, anus; *ἄλγος*, pain]. Neuralgic pain in the rectum.

Proctectomy (prok-tek'-to-me) [*πρωκτός*, anus; *ἐκτομή*, excision]. Excision of the rectum.

Proctitis (prok-ti'-tis) [*πρωκτός*, anus; *ιτις*, inflammation]. Inflammation of the anus or rectum.

Procto- (prok'-to-) [*πρωκτός*, anus]. A prefix signifying anus or rectum.

Proctocoele (prok'-to-sel) [*πρωκτός*, anus; *κήλη*, hernia]. Prolapse of the rectum.

Proctodeum, or Proctodæum (prok-to-de'-um) [*πρωκτός*, the anus; *δαειν*, to divide]. An invagination of the ectoderm in the embryo that grows inward toward the cloaca until the ectoderm and endoderm come into contact; the membrane formed between the two finally breaks through, the gut then opening externally.

Proctodynia (prok-to-din'-e-ah) [*πρωκτός*, anus; *ὀδύνη*, pain]. Pain about the anus or in the rectum.

Proctoplasty (prok'-to-plas-te) [*πρωκτός*, anus; *πλάσσειν*, to form]. Plastic surgery of the anus.

Proctoptosis (prok-top-to'-sis) [*πρωκτός*, anus; *πτώσις*, a falling]. Prolapse of the rectum.

Proctorrhaphy (prok-tor'-a-fe) [*πρωκτός*, anus; *ῥαφή*, suture]. Suture of the rectum or anus.

Proctotomy (prok-to'-o-me) [*πρωκτός*, anus; *τομή*, a cutting]. Incision into the rectum, especially for stricture.

Procursive (pro-kur'-siv) [*pro*, forward; *currere*, to run]. Running forward, as **P.** epilepsy, a form in which the patient runs during the epileptic attack.

Prodromal (pro-dro'-mal) [*πρό*, before; *δρόμος*, a running]. Precursory; of the nature of a prodrome.

Prodrome (pro'-dröm) [*πρό*, before; *δρόμος*, a running]. A forerunner of a disease; a symptom indicating the approach of a disease.

Productive (pro-duk'-tiv) [*pro*, before; *ducere*, to lead]. Forming, especially forming new tissue, as a **P.** inflammation.

Proencephalus (pro-en-sef'-al-us) [*πρό*, before; *ἐγκέφαλος*, brain]. A monster characterized by a protrusion of the brain through a fissure in the frontal region.

Professional (pro-fesh'-un-al) [*profiteri*, to profess]. 1. Pertaining to a profession, especially to the medical profession; in keeping with medical ethics. 2. Produced by the practice of a profession, as **P.** neurosis (writer's cramp, telegrapher's cramp, etc.).

Profundus (*pro-fun'-dus*) [*profundus*, deep]. Deep-seated; applied to certain muscles, and in the feminine, *profunda*, to certain arteries. See *Arteries*, *Table of*.

Progenitor (*pro-jen'-it-or*) [*pro*, before; *gignere*, to beget]. Ancestor, or forefather.

Progeny (*pro-jen'-e*) [*pro*, before; *gignere*, to beget]. Offspring; descendants.

Proglottis (*pro-glot'-is*) [*pro*, before; *γλῶσσα*, tongue: *pl.*, *Proglottides*]. A mature segment of a tapeworm.

Prognathous (*prog'-na-thus*) [*pro*, before; *γνάθος*, jaw]. Having a projecting lower jaw.

Prognosis (*prog-no'-sis*) [*pro*, before; *γνώσις*, knowledge]. A judgment in advance concerning the duration, course, and termination of a disease.

Prognostic (*prog-nos'-tik*) [*pro*, before; *γνώσις*, knowledge]. Pertaining to prognosis.

Prognosticate (*prog-nos'-tik-āt*) [*pro*, before; *γνώσις*, knowledge]. To give a prognosis.

Progression (*pro-gresh'-un*) [*progredi*, to advance]. The act of advancing or moving forward. **P.**, **Backward**, a backward walking, a rare symptom of certain nervous lesions. **P.**, **Cross-legged**, walking with the legs almost crossing, a condition sometimes observed in bilateral hip-disease and in cerebral spastic palsy.

Progressive (*pro-gres'-iv*) [*progredi*, to go forward]. Gradually extending. **P.** **Muscular Atrophy**, chronic anterior poliomyelitis in which the large ganglion-cells of the anterior horns are gradually destroyed, which leads to atrophy of the muscles. **P.** **Ossifying Myositis**, a chronic inflammation of muscles, associated with a tendency to bony deposits in them.

Projectile (*pro-jek'-til*) [*pro*, before; *jacere*, to throw]. Throwing forward. **P.** **Vomiting**, a form of vomiting sometimes observed in diseases of the brain, in which the material is suddenly projected out of the mouth to some distance, generally without nausea.

Projection (*pro-jek'-shun*) [*pro*, before; *jacere*, to throw]. 1. The act of throwing forward. 2. A part extending beyond the level of the surrounding surface. 3. The referring of impressions made on the organs of sense to the position of the object producing them. **P.** **systems**, the name given to the pathways connecting the cerebral cortex with the periphery. The first **P.** system corresponds to the fibers passing through the corona radiata; the second, to the tracts proceeding downward to the gray matter from the third ventricle to the end of the spinal cord; the third, to the peripheral nerves.

Prolapse (*pro-laps'*) [*prolabi*, to slip down]. The falling forward or downward of a part.

P. of the Cord, premature expulsion of the umbilical cord during parturition. **P. of the Iris**, protrusion of the iris through a corneal wound.

Proliferate (*pro-lif'-er-āt*) [*proles*, offspring; *ferre*, to bear]. To multiply; to generate.

Proliferation (*pro-lif'-er-ā'-shun*) [*proles*, offspring; *ferre*, to bear]. The act of proliferating or multiplying, as **P.** of cells.

P., **Atrophic**, the multiplication of cells in atrophic tissues.

Proliferative, **Proliferous** (*pro-lif'-er-ā-tic*, *pro-lif'-er-us*) [*proles*, offspring; *ferre*, to bear]. Multiplying; characterized by the formation of new tissues or by cell-proliferation. **P.** **Cyst**, a cyst in which the lining epithelium proliferates and produces projections from the inner surface of the cyst.

Prolific (*pro-lif'-ik*) [*proles*, offspring; *facere*, to make]. Fruitful.

Proliferous (*pro-lif'-er-us*) [*proles*, offspring; *gerere*, to bear]. Germinating; producing offspring. **P.** **Disc**. See *Discus proligerus*.

Prominence (*prom'-in-ens*) [*pro*, before; *minere*, to jut]. 1. A projection on the surface of a part, especially on a bone. 2. The state of being prominent. **P.**, **Genital**, an accumulation of cells on the ventral aspect of the embryonic cloaca, from which the generative organs are developed.

Promontory (*prom'-on-to-re*) [*pro*, before; *mons*, mountain]. A projecting prominence.

P. of the Sacrum, the prominence formed by the angle between the upper extremity of the sacrum and the last lumbar vertebra.

Pronation (*pro-nā'-shun*) [*prona*, to bend forward]. 1. The condition of being prone; the act of placing in the prone position. 2. Of the hand, the turning of the palm downward.

Pronator (*pro-nā'-tor*) [*prona*, to bend forward]. That which pronates, a term applied to several muscles. See *Muscles*, *Table of*.

Prone (*prōn*) [*pronus*, prone]. Lying with the face downward; of the arm, having the palm directed downward; the opposite of supine.

Pronephron, **Pronephros** (*pro-nef'-ron*, *pro-nef'-rus*) [*pro*, before; *νεφρός*, a kidney]. The anterior of the three segments of the Wolffian body opening by means of the Müllerian duct into the cloaca. It is the first part of the urogenital system to be differentiated in the vertebrate embryo.

Pronucleus (*pro-nū'-kle-us*) [*pro*, before; *nucleus*, nucleus]. One of the two nuclear bodies of a newly fecundated ovum, the male **P.** and the female **P.**, the fusion of which results in the formation of the first embryonic nucleus.

Proof-spirit. See *Spirit*.

Prootic (*pro-o'-tik*) [*πρό*, before; *οὖς*, ear]. In front of the ear.

Propane (*pro'-pan*) [*propionic*] C_3H_8 . A hydrocarbon, the third member of the marsh-gas series, occurring in petroleum.

Prop-cells. Columnar or fusiform cells placed in the intervals of the rods and hair-cells of the organ of Corti. They are also known as supporting cells and cells of Deiter.

Propenyl (*pro'-pen-il*). See *Glyceryl*.

Propepsin (*pro-pep'-sin*) [*πρό*, before; *πέπτειν*, to cook]. The zymogen of pepsin, found in the cells of the gastric glands.

Propeptone (*pro-pep'-tōn*). See under *Pep-tone*.

Propeptonuria (*pro-pep-ton-u'-re-ah*) [*πρό*, before; *peptone*; *urina*, urine]. The appearance of propeptone in the urine. It is said to occur in fevers, diphtheria, osteomalacia, and during the administration of storax or phosphorus.

Properitoneal (*pro-per-e-ton-e'-al*) [*πρό*, before; *περιτόναιον*, peritoneum]. Situated in front of the peritoneum. **P. Hernia**, a hernia the sac of which extends in various directions within the abdominal walls.

Prophylactic (*pro-fil-ak'-tik*) [*προφυλάσσειν*, to keep guard before]. 1. Pertaining to prophylaxis. 2. A remedy or agent that prevents the development of disease.

Prophylaxis (*pro-fil-aks'-is*) [*προφυλάσσειν*, to keep guard before]. Prevention of disease; measures preventing the development or spread of disease.

Propionic Acid (*pro-pe-on'-ik*) [*πρώτος*, first; *πῖον*, fat], $C_3H_5O_2$, a monobasic acid of the fatty acid series, occurring in sweat, chyme, and occasionally in diabetic urine.

Proprius (*pro'-pre-us*) [L.]. Individual; special, as *Flexor proprius pollicis*, the special flexor of the thumb.

Proptosis (*prop-to'-sis*) [*πρό*, forward; *πτῶσις*, a falling]. A falling downward; prolapse.

Propulsion (*pro-pul'-shun*) [*pro*, before; *pellere*, to push]. 1. The act of pushing or driving forward. 2. A falling forward in walking, a condition seen in paralysis agitans. See *Festination*.

Propyl (*pro'-pil*) [*propionic*], C_3H_7 . The radicle of propane.

Propylamin (*pro-pil'-am-in*) [*propyl*, from *propionic*; *amin*]. A liquid basic compound having the formula C_3H_7N , and existing in two forms, a normal P., boiling at $47^\circ C.$, and isopropylamin, boiling at $31.5^\circ C.$ Normal P. has been obtained from cultures of bacteria of feces; isopropylamin has been found among the distillation-products of beet-root molasses. See *Plomains*, Table of.

Propylene (*pro'-pil-en*) [*propyl*, from *pro-*

pienic], C_3H_4 . A gaseous hydrocarbon belonging to the series of the olefins.

Pro re nata (*pro re na'-tah*) [L.]. A phrase signifying according to the circumstances of the case.

Proscoplex (*pro-sko'-leks*) [*πρό*, before; *σκῶληξ*, a worm; *pl.*, *Proscolices*]. The embryo of a cestode worm immediately after leaving the egg.

Prosector (*pro-sek'-tor*) [*pro*, for; *sector*, cutter]. An officer of a medical college who prepares subjects for anatomic dissection, or to illustrate didactic lectures.

Prosencephalon (*pros-en-sef'-al-on*) [*πρός*, before; *ἐγκέφαλος*, brain]. The forebrain; that part of the anterior cerebral vesicle from which are developed the hemispheres, the corpus callosum, the anterior commissure, the fornix, the septum lucidum, the anterior perforated space, the olfactory lobes, and the corpus striatum.

Prosopalgia (*pros-o-pal'-je-ah*) [*πρόσωπον*, face; *ἄλγος*, pain]. Face-ache; neuralgic pain in the distribution of the trigeminal nerve; tic douloureux.

Prostatagia (*pros-ta-tal'-je-ah*) [*προστάτης*, prostate; *ἄλγος*, pain]. Pain in the prostate gland.

Prostate, or Prostate Gland (*pros'-tat*) [*προστάτης*, prostate; *πρό*, before; *ιστάναι*, to stand]. The organ surrounding the neck of the bladder and beginning of the urethra (prostatic urethra). It consists of two lateral lobes and a middle lobe, and is composed of muscular and glandular tissue, the prostatic glands. The prostate often becomes enlarged in advanced life, and may then interfere with the emptying of the bladder.

Prostatectomy (*pros-ta-tek'-to-me*) [*προστάτης*, prostate; *ἐκτομή*, excision]. Excision of the prostate.

Prostatic (*pros-tat'-ik*) [*προστάτης*, prostate]. Relating to the prostate. **P. Calculus**, a stone lodged in the prostate gland. **P. Plexus**. 1. A collection of veins surrounding the neck and base of the bladder and the prostate gland. 2. A plexus of nerves derived from the pelvic plexus, and distributed to the prostate gland, seminal vesicles, and erectile tissue of the penis. **P. Urethra**, that portion of the urethra surrounded by the prostate gland.

Prostatitis (*pros-ta-ti'-tis*) [*προστάτης*, prostate; *ιτις*, inflammation]. Inflammation of the prostate gland.

Prostatorrhœa (*pros-ta-tor-e'-ah*) [*προστάτης*, prostate; *ῥοία*, flow]. A thin urethral discharge coming from the prostate gland in cases of prostatitis or masturbation.

Prostatotomy (*pros-ta-tot'-o-me*) [*προστάτης*, prostate; *τομή*, a cutting]. Incision into the prostate gland.

Prosthesis (*pros'-thes-is*) [*πρός*, to; *θεῖς*, a placing]. Replacement of a missing part by an artificial substitute.

Prosthetic (*pros-thet'-ik*) [*πρός*, to; *θεῖς*, a placing]. Pertaining to prosthesis.

Prosthetics (*pros-thet'-iks*) [*πρός*, to; *θεῖς*, a placing]. The branch of surgery that deals with prosthesis.

Prostrate (*pros'-trāt*) [*pro*, before; *sternere*, to spread]. Lying at full length.

Prostration (*pros-tra'-shun*) [*pro*, before; *sternere*, to spread]. 1. The condition of being prostrate. 2. Extreme exhaustion of nervous or muscular force. P., Nervous, general exhaustion from excessive expenditure of nervous energy.

Protagon (*pro'-tag-on*) [*πρωτός*, first; *ἀγεῖν*, to lead], $C_{10}H_{10}N_2O_{10}$. A crystalline glucosid found in nervous tissue, and yielding when boiled with baryta the decomposition-products of lecithin.

Proalbumose (*prot-al'-bu-mōs*). Same as *Protoalbumose*.

Protamin (*pro'-ta-min*) [*πρωτός*, first; *αμίν*], $C_{10}H_{10}N_2O_2$. An organic base found with nuclein in the spawn of salmon.

Protean (*pro'-te-an*) [*Proteus*, a Grecian sea-god who had the power of changing his shape]. Taking on many shapes; as a P. disease, P. eruption.

Protective (*pro-tek'-tiv*) [*pro*, before; *tegere*, to cover]. Covering so as to shield against harm; guarding against harm, as a P. dressing. P. Proteids. See *Proteids*, *Defensive*.

Proteid (*pro'-te-id*) [*πρωτός*, first]. Any one of the important and essential nitrogenous constituents of animal (animal P.) and vegetable (vegetable P.) tissues. They are colloid, noncrystallizable, and levorotary, are precipitated from solutions by alcohol and various metallic salts, and coagulated by heat and mineral acids. Proteids are divided by Landois into: (a) *native albumins*: serum-albumin, egg-albumin, metalbumin, and par-albumin; (b) *derived albumins, or albuminates*: acid-albumin, syntonin, alkali-albumin, and casein; (c) *globulins*: crystallin, vitellin, paraglobulin, or serum-globulin, fibrinogen, myosin, globin; (d) *fibrins*; (e) *coagulated proteids*; (f) *albumoses and peptones*; (g) *larducin*. P., Bacterial, one formed by the action of a microorganism. P., Bacterial Cellular, any proteid found in the bodies of bacteria. P., Defensive, one of the proteids existing in the blood and rendering the system immune to infectious diseases.

Protein (*pro'-te-in*) [*πρωτός*, first]. A substance formerly regarded as a proteid deprived of its sulphur, and looked upon as an alkali-albumin.

Proteolysis (*pro-te-ol'-is-is*) [*πρωτός*, first; *λύσις*, solution]. The change produced in proteids by ferments that convert them into diffusible bodies.

Proteolytic (*pro-te-ol'-ik*) [*πρωτός*, first; *λύσις*, solution]. Pertaining to, characterized by, or effecting, proteolysis.

Proteose (*pro'-te-ōs*) [*πρωτός*, first]. Any one of a group of bodies formed in gastric digestion, intermediate between the food-proteids and the peptones, called antipeptones, hemipeptone, etc.

Proteus (*pro'-te-us*) [L.]. A genus of bacteria. See *Bacteria*, *Table of*.

Prothesis (*proth'-es-is*). See *Prosthesis*.

Prothetic (*pro-thet'-ik*). See *Prosthetic*.

Protiodid (*pro-ti'-o-did*) [*πρωτός*, first; *ἰώδης*, like a violet]. A salt containing the least amount of iodine of the iodides of the same base. See *Proto*.

Proto- (*prō'-to-*) [*πρωτός*, first]. 1. A prefix signifying first. 2. In chemistry, a prefix signifying the lowest of a series of compounds of the same elements, as *protiodid*, *protoclorid*, *protozoid*.

Protoalbumose (*pro-to-al'-bu-mōs*). See *Albumose*.

Protoblast (*pro'-to-blast*) [*πρωτός*, first; *βλαστός*, germ]. A cell without a cell-wall.

Protocatechuic Acid (*pro-to-kat'-e-chu'-ik*) [*πρωτός*, first; *catechu*], $C_7H_6O_4$. Dioxycarboxylic acid, an acid sometimes occurring in the urine.

Protoclorid (*pro-to-kle'-rid*). See *Proto* (second definition).

Protogaster (*pro-to-gas'-ter*) [*πρωτός*, first; *γαστήρ*, stomach]. The primitive intestinal cavity of a gastrula.

Protoglobulose (*pro-to-glob'-u-lōs*) [*πρωτός*, first; *globulus*, a ball]. One of the primary products of the digestion of globulin.

Protoiodid (*pro-to-i'-o-did*). Synonym of *Protiodid*.

Protileukocyte (*pro-to-lu'-ko-sit*) [*πρωτός*, first; *λευκός*, white; *κύτος*, cell]. One of the minute lymphoid cells found in the red bone-marrow and also in the spleen.

Protomyosinose (*pro-to-mi-ō'-sin-ōs*) [*πρωτός*, first; *μῦς*, muscle]. A product of the primary digestion of myosin.

Protonephron (*pro-to-nef'-ron*) [*πρωτός*, first; *νεφρός*, kidney]. The pronephron, metanephron, and mesonephron taken together; the primitive kidney.

Protopathic (*pro-to-path'-ik*) [*πρωτός*, first; *πάθος*, disease]. Primary; relating to the first lesion; idiopathic; opposed to *deutero-pathic*.

Protoplasm (*pro'-to-plasm*) [*πρωτός*, first; *πλάσσειν*, to mold]. The viscid material constituting the essential substance of living cells, upon which all the vital functions of

nutrition, secretion, growth, reproduction, irritability, motility, depend. When highly magnified the protoplasm of most cells appears as a network (spongioplasm), containing a more fluid substance (hyaloplasm) in its meshes.

Protoplasmic (*pro-to-plas'-mik*) [*πρώτος*, first; *πλάσσειν*, to mold]. 1. Pertaining to protoplasm. 2. Composed of protoplasm. **P. Process.** See *Process*.

Protospasm (*pro'-to-spasm*) [*πρώτος*, first; *σπασμός*, spasm]. A spasm beginning in one part and extending to others.

Protosulphate (*pro-to-sul'-fāt*). See *Proto-* (2d def.).

Prototype (*pro'-to-tīp*) [*πρώτος*, first; *τύπος*, type]. An original type; a type after which others are copied.

Protoxid (*pro-toks'-id*). See *Proto-* (2d def.).

Protozoon (*pro-to-zo'-on*) [*πρώτος*, first; *ζῷον*, animal]. One of the lowest class of the animal kingdom, comprising organisms which consist of simple cells or colonies of cells, and which possess no nervous system and no circulatory organs.

Protyl (*pro'-til*) [*πρώτος*, first]. The supposed primitive universal element.

Proud Flesh. Exuberant granulation-tissue.

Provisional (*pro-vish'-un-āl*) [*pro*, before; *videre*, to see]. For temporary use, as *P. callus*.

Proximad (*proks'-im-ad*) [*proximus*, nearest; *ad*, toward]. Toward the proximal end.

Proximal (*proks'-im-āl*) [*proximus*, nearest]. Nearest to the body or the median line of the body, or some other point considered as the center of a system.

Proximate (*proks'-im-āl*) [*proximus*, nearest]. Nearest; immediate, as *P. cause*.

P. Principle, a substance existing under its own form in the animal solids or fluids, and that can be extracted by means not altering or destroying its chemic properties.

Prune - juice Expectorant. A peculiar bloody sputum, of a dark purple color, resembling prune-juice. It occurs in low forms of croupous pneumonia, in gangrene and carcinoma of the lung.

Prunum (*pru'-num*) [L.] Prune, the fruit of *Prunus domestica*, of the order Rosaceæ. Prunes are laxative.

Prunus (*pru'-nus*) [L.]. A genus of trees of the order Rosaceæ. *P. domestica*, is the source of the prune. *P. serotina*, yields wild-cherry bark (*Prunus Virginiana*, U. S. P.), which contains a volatile oil, hydrocyanic acid, tannic acid, a resin, and other substances. It is used as a tonic and sedative in gastric debility and general irritation of the system, and is a common ingredient of cough-mixtures. Prepara-

tions: *Extractum pruni virginianæ fluidum*. Dose f 3j (4.0). *Infusum pruni virginianæ*. Dose f 3ij-f 3iij (64.0-96.0). *Syrupus pruni virginianæ*. Dose f 3ss (16.0).

Pruriginous (*pru-rij'-in-us*) [*prurire*, to itch]. Pertaining to or resembling prurigo.

Prurigo (*pru-ri'-go*) [*prurire*, to itch]. 1. A chronic inflammatory disease of the skin, characterized by small pale papules and severe itching. The papules are deeply seated, and are most prominent on the extensor surfaces of the limbs. The disease begins in early life and is usually incurable. 2. *Pruritus*.

Pruritic (*pru-rit'-ik*) [*prurire*, to itch]. Pertaining to pruritus; itching.

Pruritus (*pru-ri'-tus*) [*prurire*, to itch]. Itching, a peculiar, uncomfortable sensation due to irritation of the peripheral sensory nerve, and leading to scratching. It may be a symptom of other diseases of the skin, or an independent neurosis. The latter receives its name from the part affected, as *P. ani*, *P. vaginæ*. *P. hiemalis*, a form affecting certain persons only in winter, especially in dry climates. *P. senilis*, the *P.* of the aged, at times due to degenerative changes in the skin.

Prussian Blue. The ferrocyanid of iron, used in the arts as a dye; it was formerly employed in medicine as a febrifuge, tonic, and alterative. Dose gr. iij-v (0.2-0.32).

Prussiate (*prus'-e-āt*). A salt of prussic, or hydrocyanic acid.

Prussic Acid. See *Acid, Hydrocyanic*.

Psalterium (*sal-te'-re-um*) [*ψαλτήριον*, a lute]. The third compartment of the stomach of ruminants. It is also called manyplies.

Psammocarcinoma (*sam-o-kar-sin-o'-mah*) [*ψάμμος*, sand; *carcinoma*]. A carcinoma containing calcareous deposits.

Psammoma (*sam-o'-mah*) [*ψάμμος*, sand; *ῥμα*, tumor]. A firm tumor found in the membranes of the brain, the choroid plexus, and in other parts, and characterized by the presence of peculiar mineral concretions. The tumor is generally a fibrosarcoma.

Psammous (*sam'-us*) [*ψάμμος*, sand]. Sandy; sabulous.

Pseud-, Pseudo-, (*sūd-, su'-dō-*) [*ψευδής*, false]. A prefix meaning false.

Pseudaconitin (*su-dak-on'-it-in*) [*ψευδής*, false; *aconite*], $C_{36}H_{48}NO_{12}$. An extremely poisonous alkaloid from *Aconitum ferox*.

Pseudacousma, Pseudacusic (*su-dak-ooz'-mah, su-dak-u'-sis*, [*ψευδής*, false; *ἀκούειν*, to hear]. A disturbance of hearing in which the person's own voice sounds strange or peculiar.

Pseudarthrosis (*su-dar-thro'-sis*) [*ψευδής*, false; *ἄρθρον*, a joint]. A false joint.

Pseudencephalus (*su-den-sef'-ul-us*) [*ψευ-*

δῆς, false; *ἐγκέφαλος*, brain]. A species of monster characterized by a partial development of the frontal, parietal, and occipital bones, while the brain is represented by a bunch of membranes, blood-vessels, connective, and possibly nervous tissue, at the base of the skull.

Pseudoesthesia (*su-des-the'-ze-ah*) [*ψευδής*, false; *αἴσθησις*, feeling]. An imaginary sensation for which there is no corresponding object; a sensation in parts of the body that have been removed by accident or surgical operation.

Pseudo- (*su'-do-*). See *Pseud-*.

Pseudoactinomycosis, Pseudactinomycosis (*su-do-ak-tin-o-mi-ko'-sis*, *su-dak-tin-o-mi-ko'-sis*) [*ψευδής*, false; *ἀκτίς*, ray; *μύκης*, fungus]. A form of pulmonary tuberculosis in which the sputum contains granular bodies resembling the grains of actinomycosis. They consist of a crystalline substance similar to leucin.

Pseudoangina (*su-do-an-ji'-nah*) [*ψευδής*, false; *ἀνγερῆς*, to strangle]. False angina; hysterical angina; an attack of cardiac pain somewhat resembling angina pectoris but less grave in character, longer in duration, and usually not associated with organic heart-disease. It occurs in neurotic women, and is generally brought on by emotional excitement.

Pseudarthrosis (*su-do-ar-thro'-sis*). See *Pseudarthrosis*.

Pseudobacterium (*su-do-bak-te'-re-um*) [*ψευδής*, false; *βακτήριον*, a little rod or staff]. Any object resembling a bacterium.

Pseudobulbar (*su-do-bul'-bar*) [*ψευδής*, false; *βολβός*, bulb]. Not really bulbar. **P. Paralysis**, symmetric disease of both cerebral hemispheres involving the centers or paths of the nerves of speech, and thus resembling disease of the medulla oblongata.

Pseudocoele (*su'-do-sel*) [*ψευδής*, false; *κοῖλος*, hollow]. The fifth ventricle of the brain.

Pseudocrisis (*su-do-kri'-sis*) [*ψευδής*, false; *κρίσις*, separating]. A false crisis; a sudden fall of temperature resembling the crisis of a disease, but subsequently followed by a rise of temperature and a continuation of the disease. **P.** is common in pneumonia.

Pseudocroup (*su'-do-kraṑ*) [*ψευδής*, false; *ἄσθμα*, *hρόpan*, to cry aloud]. False croup; laryngismus stridulus.

Pseudocyesis (*su-do-si-e'-sis*) [*ψευδής*, false; *κύησις*, pregnancy]. False pregnancy; the belief in the existence of pregnancy on the part of a woman when none exists.

Pseudodiphtheria (*su-do-dif-the'-re-ah*) [*ψευδής*, false; *diphtheria*]. An inflammation characterized by the presence of a false membrane not due to the Klebs-Löffler bacillus.

P. bacillus, a nonpathogenic bacillus resembling in form and growth the true diphtheria-bacillus. It is now considered an attenuated form of the true bacillus.

Pseudoephedrin (*su-do-ef'-ed-rin*), $C_{10}H_{17}NO$. An alkaloid found in *Ephedra vulgaris* and isomeric with ephedrin.

Pseudoerysipelas (*su-do-er-e-sip'-el-as*) [*ψευδής*, false; *erysipelas*]. Inflammation of the subcutaneous cellular tissue resembling erysipelas.

Pseudoglioma (*su-do-gli-o'-mah*) [*ψευδής*, false; *γλία*, *glia*; *όμα*, tumor]. A name given to inflammatory changes of the vitreous humor, due to iridochoroiditis, and resembling glioma of the retina.

Pseudogonococcus (*su-do-gon-o-ko'-kus*) [*ψευδής*, false; *gonococcus*]. A diplococcus found in the normal urethra, resembling the gonococcus.

Pseudogonorrhea (*su-do-gon-or-e'-ah*) [*ψευδής*, false; *gonorrhea*]. A simple nonspecific urethritis.

Pseudohermaphroditism (*su-do-her-me'-fro-dism*) [*ψευδής*, false; *hermaphroditism*]. A condition simulating hermaphroditism.

Pseudohydrophobia (*su-do-hi-dro-fo'-be-ah*) [*ψευδής*, false; *ὕδωρ*, water; *φόβος*, fear]. A condition resembling hydrophobia, at times produced by dread of the disease; lysophobia.

Pseudohypertrophic (*su-do-hi-per-tro'-fik*) [*ψευδής*, false; *ὑπέρ*, over; *τροφή*, nutrition]. Pertaining to or characterized by pseudohypertrophy. **P. Muscular Paralysis**. See *Paralysis, Pseudohypertrophic*.

Pseudohypertrophy (*su-do-hi-per'-tro-fe*) [*ψευδής*, false; *ὑπέρ*, above; *τροφή*, nutrition]. False hypertrophy; increase in the size of an organ or part on account of overgrowth of an unimportant tissue. It is accompanied by diminution in function.

Pseudoleukemia (*su-do-lu-ke'-me-ah*) [*ψευδής*, false; *leukemia*]. See *Hodgkin's Disease*. **P.**, Infantile, Von Jaksch's disease, a form of anemia occurring in young children, usually dependent on a rachitic diathesis, and not associated with much leukocytosis.

Pseudoleukocythemia (*su-do-lu-ko-si-the'-me-ah*) [*ψευδής*, false; *λευκός*, white; *κύτος*, cell; *αἷμα*, blood]. Synonym of *Pseudoleukemia*.

Pseudolipoma (*su-do-lip-o'-mah*) [*ψευδής*, false; *λίπος*, fat; *όμα*, tumor]. A localized edema resembling an accumulation of fat, occurring above the clavicle and about the knee, especially in cases of rheumatism.

Pseudomembrane (*su-do-mem'-brān*) [*ψευδής*, false; *membrana*, membrane]. A false membrane, such as is seen in diphtheria.

Pseudomembranous (*su-do-mem'-bra-nus*) [*ψευδής*, false; *membrana*, membrane].

Characterized by, or pertaining to false membranes. **P. Inflammation**, any inflammation characterized by the formation of a false membrane.

Pseudoneuroma (*su-do-nu-ro'-mah*) [*ψευδής*, false; *νεῦρον*, nerve; *δῆμα*, tumor]. A false neuroma. See *Neuroma*.

Pseudoparalysis (*su-do-par-al'-is-is*) [*ψευδής*, false; *παράλυσις*, paralysis]. Paralysis of motion, apparently but not really due to a lesion of the nervous system. **P. of Rickets**, the inability to walk in severe cases of rickets, due to distortion of the bones. **P.**, **Syphilitic**, an inflammatory condition of the epiphyses of the bones in syphilis causing a marked impairment of motion.

Pseudoparasite (*su-do-par'-a-sit*) [*ψευδής*, false; *παρά*, beside; *σίτος*, food]. Any object resembling a parasite.

Pseudopod, **Pseudopodium** (*su'-do-pod*, *su-do-po'-de-um*) [*ψευδής*, false; *πούς*, foot]. A protrusion of a portion of the substance of an amoeboid cell.

Pseudoscarlatina (*su-do-skar-la-te'-nah*) [*ψευδής*, false; *scarlatina*, scarlet fever]. A febrile disease associated with a rash like that of scarlet fever, occurring as a result of gonorrhea, or after puerperal infection.

Pseudosclerosis (*su-do-skle-ro'-sis*) [*ψευδής*, false; *σκληρός*, hard]. An affection similar in symptoms to multiple sclerosis of the nervous system, but without the anatomic lesions.

Pseudosmia (*su-dos'-me-ah*) [*ψευδής*, false; *ὀσμή*, smell]. Perversion of the sense of smell; an olfactory hallucination.

Pseudostoma (*su-dos'-to-mah*) [*ψευδής*, false; *στόμα*, mouth]. An apparent aperture between endothelial cells that have been stained with silver nitrate.

Pseudotabes (*su-do-ta'-bez*) [*ψευδής*, false; *tabes*, a wasting]. A disease simulating or resembling tabes dorsalis or tabes mesenterica.

Pseudotuberculosis (*su-do-tu-ber-ku-lo'-sis*) [*ψευδής*, false; *tuberculosis*]. A disease resembling tuberculosis, but not caused by the tubercle-bacillus.

Pseudotypoid (*su-do-ti'-phoid*) [*ψευδής*, false; *typhoid*]. Spurious typhoid, a disease simulating typhoid fever, but in which the true lesions of this disease as well as the typhoid bacilli are absent.

Pseudoxanthin (*su-do-xan'-thin*) [*ψευδής*, false; *ξανθος*, yellow]. 1. $C_4H_5N_5O$, a leukomycin isolated from fresh beef. 2. A body isomeric with xanthin, obtained by action of sulphuric acid upon uric acid.

Psilosis (*si-lo'-sis*) [*ψιλος*, bare]. 1. The removal of the hair from a part; depilation. 2. A certain form of tropical diarrhea.

Psoas (*so'-as*) [*ψῶα*, loin]. One of two mus-

cles, *P. magnus* and *P. parvus*. See *Muscles*, *Table of*. **P.-abscess**, an abscess, usually dependent upon tuberculous disease of the vertebra, making its way along the sheath of the psoas muscle and pointing at the front of the thigh, below Poupart's ligament, to the outer side of the spine of the pubis.

Psodymus (*sod'-im-us*) [*ψῶα*, loin; *δίδυμος*, double]. A monster with two heads and chests, and conjoined abdominal and pelvic cavities.

Psora (*so'-rah*) [*ψώρα*, from *ψάειν*, to scratch]. Scabies.

Psoriasis (*so-re-as'-ik*). See *Psoriatic*.

Psoriasis (*so-ri'-as-is*) [*ψώρα*, the itch]. A chronic inflammatory disease of the skin, characterized by the development of reddish patches covered with whitish scales. The disease affects especially the extensor surfaces of the body. **P. annularis**. Synonym of *P. circinata*. **P.**, **Buccal**, *P. buccalis*. Synonym of *Leukoplakia buccalis*. **P. circinata**, *P.* in which the central part of the lesions has disappeared, leaving ring-shaped patches. **P. diffusa**, a form in which there is coalescence of large contiguous lesions. **P. guttata**. See *P. punctata*. **P. gyrata**, *P.* with a serpentine arrangement of the patches. **P. punctata**, a form in which the lesions consist of minute red papules which rapidly become surmounted by pearly scales. **P. universalis**, a form in which the lesions are all over the body.

Psoriatic (*so-re-at'-ik*) [*ψώρα*, itch]. Pertaining to, or affected with psoriasis.

Psorophthalmia (*so-roff-thal'-me-ah*) [*ψώρα*, itch; *ὀφθαλμός*, eye]. Marginal blepharitis.

Psorosperm (*so'-ro-sperm*) [*ψώρα*, the itch; *σπέρμα*, seed]. A unicellular organism belonging to the Protozoa; a coccidium.

Psorospermial, **Psorospermic** (*so-ro-sperm'-e-al*, *so-ro-sperm'-ik*) [*ψώρα*, itch; *σπέρμα*, seed]. Pertaining to, or affected with psorosperms.

Psorospermiasis (*so-ro-sperm-i'-as-is*) [*ψώρα*, itch; *σπέρμα*, seed]. A state characterized by the presence of psorosperms.

Psorospermiosis (*so-ro-sperm-o'-sis*) [*ψώρα*, itch; *σπέρμα*, seed]. A diseased condition associated with the presence of psorosperms. **P.**, **Proliferative Follicular**. Synonym of *Keratosis follicularis* and *Darier's Disease*.

Psychiatric (*si-ke-at'-rik*) [*ψυχή*, mind; *ιατρική*, healing art]. Pertaining to psychiatry.

Psychiatry (*si-ki'-at-re*) [*ψυχή*, mind; *ιατρική*, healing art]. The science and treatment of the diseases of the mind.

Psychic, **Psychical** (*si'-kik*, *si'-hik-al*) [*ψυχή*, mind]. Pertaining to the mind. **P. Blindness**. See *Word-blindness*. **P. Deafness**. See *Deafness*. **P. Infection**,

mental infection; the development of a mental condition or disease through an influence acting upon the mind.

Psycho- (*si'-ko-*) [*ψυχή*, mind]. A prefix denoting connection with the mind.

Psychoauditory, Psychauditory (*si'-ko-au'-dit-o-re*, *si'-kro'-dit-o-re*) [*ψυχή*, mind; *αἰνέειν*, to hear]. Pertaining to the psychic perception of sound. **P. Area**, the cortical area concerned in the conscious perception of sound.

Psychocortical (*si'-ko-kor'-tik-al*) [*ψυχή*, mind; *cortex*, cortex]. Pertaining to that part of the cerebral cortex concerned in the conscious perception of sensations.

Psychology (*si'-kol'-o-je*) [*ψυχή*, mind; *λόγος*, science]. The science having for its object the investigation of the mind or consciousness.

Psychometry (*si'-kom'-et-re*) [*ψυχή*, mind; *μέτρον*, measure]. The measurement of the duration of psychic processes.

Psychomotor (*si'-ko-mo'-tor*) [*ψυχή*, mind; *κινεῖν*, to move]. Pertaining to voluntary movement, as the P. area, disposed chiefly along each side of the central fissure.

Psychoneurosis (*si'-ko-nu-ro'-sis*) [*ψυχή*, mind; *νεῦρον*, nerve]. Mental disease not dependent on any organic lesion.

Psychopathia (*si'-ko-pa'-the-ah*). **Psycho-**pathy. **P. Sexualis**, P. characterized by perversion of the sexual functions.

Psychopathy (*si'-kop'-ath-e*) [*ψυχή*, mind; *πάθος*, disease]. Any disease of the mind.

Psychophysical (*si'-ko-fiz'-ik-al*) [*ψυχή*, mind; *φυσικός*, physical]. Pertaining to psychophysics. **P. Law**. See *Law, Fechner's*.

Psychophysics (*si'-ko-fiz'-iks*) [*ψυχή*, mind; *φυσικός*, physical]. The study of mental processes by physical methods; the study of the relation of stimuli to the sensations which they produce, especially the determination of the differences of stimulus required to produce recognizable differences of sensation; experimental psychology.

Psychosensory (*si'-ko-sen'-so-re*) [*ψυχή*, mind; *sensus*, sense]. Pertaining to or concerned in the conscious perception of sensory impulses.

Psychosis (*si'-ko'-sis*) [*ψυχή*, mind]. A disease of the mind, especially one without demonstrable organic lesions.

Psychotherapy (*si'-ko-ther'-ap-e*) [*ψυχή*, mind; *θεραπεία*, treatment]. The treatment of disease by mental influence, or by suggestion.

Psychrometer (*si'-krom'-et-er*) [*ψυχρός*, cold; *μέτρον*, measure]. An instrument for determining the atmospheric moisture by estimating the amount of cold required to precipitate it.

Psychrophore (*si'-kro-fōr*) [*ψυχρός*, cold;

φορεῖν, to carry]. An instrument for applying cold to deeply-seated parts, as *e. g.*, a double-current catheter for applying cold to the posterior part of the urethra.

Psychrotherapy (*si'-kro-ther'-ap-e*) [*ψυχρός*, cold; *θεραπεία*, treatment]. The treatment of disease by the use of cold.

Psydracia (*si'-dra'-se-ah*) [*ψιθύραξ*, blister, pimple]. An old term for eczema.

Ptarmic (*tar'-mik*) [*πταρμός*, a sneezing]. 1. Pertaining to the act of sneezing; sternutatory. 2. A substance that produces sneezing.

Pterion (*te'-re-on*) [*πτερόν*, wing]. See *Craniometric Points*.

Pterygium (*ter'-ij'-e-um*) [dim. of *πτερυξ*, wing]. A triangular patch of mucous membrane growing on the conjunctiva, usually on the nasal side of the eye. The apex of the patch points toward the pupil, the fan-shaped base toward the canthus.

Pterygoid (*ter'-ig'-oid*) [*πτερυξ*, wing; *ειδός*, like]. 1. Wing-shaped, as the P. plate of the sphenoid bone. 2. Pertaining to the P. canal, P. plate, P. plexus, etc.

Pterygomaxillary (*ter'-ig'-o-maks'-il-o-re*) [*πτερυξ*, wing; *ειδός*, like; *maxilla*, maxilla]. Pertaining to the pterygoid process and the maxilla. **P. Fissure**, an elongated fissure formed by the divergence of the superior maxillary bone from the pterygoid process of the sphenoid bone. **P. Ligament**, a ligament extending from the apex of the internal pterygoid plate to the posterior end of the internal oblique line of the lower jaw.

Pterygopalatine (*ter'-ig'-o-pal'-at-in*) [*πτερυξ*, wing; *palatine*]. Situated between the pterygoid plate of the sphenoid bone and the palate bone, as the P. canal.

Ptilosis (*ti'-lo'-sis*) [*πτεῖλον*, feather]. Loss of the hair, especially loss of the eye-lashes.

Ptisan (*tis'-an*) [*πτεῖσιν*, peeled barley]. 1. Barley-water. 2. A decoction of barley used as a medicinal drink.

Ptomain (*to'-ma-in*) [*πτῶμα*, corpse]. A basic compound resembling the alkaloids, formed during the decomposition of animal or vegetable tissues; a putrefactive or animal alkaloid. See *Table*.

Ptomainemia (*to'-ma-in-e'-me-ah*) [*πτῶμα*, corpse; *αἷμα*, blood]. The presence of ptomaines in the blood.

Ptomatropin (*to-mat'-ro-pin*) [*πτῶμα*, corpse; *atropin*]. A ptomain resembling the atropin in its physiologic and chemic properties, and found in decomposing meat, in the organs of patients dead of typhoid fever, etc.

Ptoxis (*to'-sis*) [*πτέειν*, to fall]. Drooping of the upper eyelid, due to paralysis or atrophy of the levator palpebre superioris.

Ptyalagogue (*ti'-al'-a-gog*) [*πτεῖλον*, saliva; *ἀγωγός*, leading]. A sialagogue.

TABLE OF PTOMAINS.

NAME.	FORMULA.	SOURCE.	PHYSIOLOGIC ACTION.
Amidovalerianic acid.	$C_8H_{11}NO_2$	Putrefying fibrin and meat.	Nontoxic.
Amylamin.	$C_8H_{13}N$	Cod-liver oil, horn, putrid yeast.	Toxic.
Asellin.	$C_{28}H_{22}N_4$	Cod-liver oil.	Toxic.
Betain.	$C_5H_{13}NO_2$	Mussel, human urine.	Nontoxic.
Butylamin.	$C_4H_{11}N$	Cod-liver oil.	Toxic (?).
Cadaverin.	$C_6H_{14}N_2$	Putrefying animal tissues, cultures of comma-bacillus.	Slightly toxic.
Caproylamin.	See <i>Hexylamin</i> .		
Cholin.	$C_8H_{15}NO_2$	Decomposing animal tissues, proteus and comma-bacillus cultures.	Toxic.
Collidin (?).	$C_8H_{11}N$	Putrid mixture of pancreas and gelatin.	
Diethylamin.	$C_4H_{11}N$	Putrefying fish (pike).	Nontoxic.
Dihydrolutidin.	$C_7H_{11}N$	Cod-liver oil.	Toxic.
Dimethylamin.	C_2H_7N	Putrid gelatin, yeast, fish.	Nontoxic.
Eczemin.		Extracted from the urine in eczema. A white, crystalline substance, soluble in water, feebly alkaline in reaction.	Toxic.
Ethylamin.	C_2H_7N	Putrefying yeast, wheat-flour.	Nontoxic.
Ethylidenediamin (?).	$C_2H_6N_2$	Putrefying haddock.	Toxic.
Gadinin.	$C_7H_{13}NO_2$	Decomposing haddock.	Toxic.
Hexylamin.	$C_6H_{15}N$	Putrid yeast.	Toxic.
Hydrocollidin (?).	$C_8H_{13}N$	Putrefying mackerel, horse-flesh, and ox-flesh.	Toxic.
Hydrocoridin.	$C_{10}H_{17}N$	Agar-culture of bacterium album.	
Methylamin.	CH_5N	Putrefying fish.	Nontoxic.
Methyl-guanidin.	$C_2H_7N_2$	Putrefying horse-flesh.	Toxic.
Morrhucic acid.	$C_8H_{13}NO_2$	Cod-liver oil.	
Morrhuin.	$C_{19}N_{27}N_2$	Cod-liver oil.	Diuretic and diaphoretic.
Muscarin.	$C_8H_{15}NO_2$	Decomposing haddock.	Toxic.
Mydalein.		Putrefying human organs.	Toxic.
Mydatoxin.	$C_8H_{13}NO_2$	Putrefying human organs.	Toxic.
Mydin.	$C_8H_{11}NO$	Putrid human tissues, blood-serum cultures of bacillus of typhoid fever.	Nontoxic.
Mytilotoxin.	$C_8H_{13}NO_2$	Poisonous mussel (<i>Mytilus edulis</i>).	Toxic.
Neuridin.	$C_6H_{14}N_2$	Putrefying flesh.	Nontoxic.
Neurin.	$C_6H_{13}NO$	Putrefying flesh.	Toxic.
Parvolin (?).	$C_9H_{15}N$	Putrefying mackerel and horse-flesh.	
Peptotoxin.		Decomposing proteid substances.	Toxic.

TABLE OF PTOMAINS.—*Continued.*

NAME.	FORMULA.	SOURCE.	PHYSIOLOGIC ACTION.
Phillogosin.		Cultures of staphylococcus pyogenes aureus.	Powerful local irritant.
Propylamin.	C_3H_9N .	Gelatin-cultures of bacteria of human feces.	
Putrescin.	$C_4H_{13}N_2$.	Putrefying flesh, gelatin, etc.	Slightly toxic.
Pyocyanin.	$C_{14}H_{14}NO_2$.	Cultures of bacillus pyocyaneus.	Nontoxic.
Pyridin-base (?).	$C_{10}H_{11}N$.	Putrefying sea-polypa.	
Pyridin-base.	$C_8H_{11}N$.	Putrefying sea-polypa.	
Saprin.	$C_8H_{14}N_2$.	Putrefying human liver and spleen.	Nontoxic.
Spasmotoxin.		Cultures of tetanus-bacillus.	Toxic.
Susotoxin.	$C_{10}H_{20}N_2$ (?).	Cultures of hog-cholera bacillus.	Toxic.
Tetanin.	$C_{12}H_{20}N_2O_4$.	Cultures of tetanus-bacillus.	Toxic.
Tetanotoxin.	$C_4H_{11}N$ (?).	Cultures of tetanus-bacillus.	Toxic.
Triethylamin.	$C_6H_{15}N$.	Putrefying fish (haddock).	Nontoxic.
Trimethylamin.	C_3H_9N .	Herring-brine.	Nontoxic.
Trimethylenediamin.	$C_4H_{10}N_2$ (?).	Beef-broth cultures of comma-bacillus.	Toxic.
Typhotoxin.	$C_7H_{17}NO_2$.	Beef-broth cultures of bacillus of typhoid fever.	Toxic.
Tyretoxicon.		Poisonous cheese, ice-cream, milk, etc.	Toxic.
Unnamed.	$C_5H_{12}N_2O_4$.	Flesh, bones, etc.	Toxic.
Unnamed.	$C_4H_{11}NO_2$.	Cultures of tetanus-bacillus.	Nontoxic.
Unnamed.	$C_7H_{15}N_2$.	Sugar undergoing alcoholic fermentation.	Nontoxic.
Unnamed.	$C_7H_{17}NO_2$.	Putrefying horse-flesh.	Toxic.
Unnamed.	$C_7H_{12}N_2O_5$.	Flesh, bones, etc.	Toxic.
Unnamed.	$C_{10}H_{15}N$.	Putrid fibrin.	Toxic.
Unnamed.	$C_{12}H_{20}N_4$.	Fermenting cane-sugar.	
Unnamed.	$C_{14}H_{25}N_3O$.	Putrefying fibrin.	
Unnamed.	$C_{17}H_{30}N_4$.	Putrefying mackerel, horse-flesh, and ox-flesh.	
Unnamed.	$C_{20}H_{33}N$.		
Unnamed.	$C_9H_9NO_4$.	Urine in influenza.	Toxic, causing fever, and killing animals in eight hours.

Ptyalin (*ti'-al-in*) [$\pi\tau\acute{\iota}\alpha\lambda\omicron\nu$, saliva]. A diastatic ferment found in saliva, having the property of converting starch into dextrin and sugar. The starch first becomes converted into achroodextrin and erythrodextrin; these by hydration into maltose, and the latter, by further hydration, into dextrose.

Ptyalinogen (*ti-al-in'-o-jen*) [$\pi\tau\acute{\iota}\alpha\lambda\omicron\nu$, saliva;

γεννᾶν, to produce]. The hypothetic antecedent of ptyalin.

Ptyalism (*ti'-al-izm*) [$\pi\tau\acute{\iota}\alpha\lambda\omicron\nu$, saliva]. Salivation. P., Mercurial (or simply P.). See *Salivation*.

Ptyalize (*ti'-al-iz*) [$\pi\tau\acute{\iota}\alpha\lambda\omicron\nu$, saliva]. To produce ptyalism.

Puberty (*pu'-ber-te*) [*pubertas*, from *puber*,

adult]. 1. The period at which the generative organs become capable of exercising the function of reproduction, signalized in the boy by a change of voice and discharge of semen, in the girl by the appearance of the menses.

Pubes (*pu'-bes*) [L.]. 1. The pubic hair.

2. The hairy region covering the os pubis.

3. The os pubis, or pubic bone; that portion of the os innominatum forming the front of the pelvis.

Pubic (*pu'-bik*) [*pubes*, *pubes*]. Pertaining to the pubes.

Pubiotomy, Pubeotomy (*pu-be-ot'-o-me*) [*pubes*, *pubes*; *rough*, a cutting]. The operation of dividing the pubic bone to facilitate delivery in cases of pelvic malformation. See also *Symphysiotomy*.

Pubofemoral (*pu-bo-fem'-or-al*) [*pubes*, *pubes*; *femur*, *femur*]. Pertaining to the pubes and the femur.

Pubovesical (*pu-bo-ves'-ik-al*) [*pubes*, *pubes*; *vesica*, *bladder*]. Pertaining to the pubes and bladder.

Puccoon, Yellow. The *Hydrastis canadensis*.

Pudendal (*pu-den'-dal*) [*pudere*, to be ashamed]. Pertaining to the pudenda.

Pudendum (*pu-den'-dum*) [*pudere*, to be ashamed]. The external genital organs, especially those of the woman, generally used in the plural, *Pudenda*.

Pudic (*pu'-dik*) [*pudicus*, from *pudere*, to be ashamed]. Pertaining to the pudenda, as the P. artery.

Puerile (*pu'-er-il*) [*puer*, boy]. Boyish; childish; pertaining to childhood. P. Respiration, exaggerated breath-sounds with expiration prolonged and high-pitched, such as is heard in healthy children.

Puerpera (*pu-ur'-pe-rah*) [*puer*, child; *parere*, to bear]. A woman who is in labor, or has recently been delivered.

Puerperal (*pu-ur'-pe-ral*) [*puerpera*]. Pertaining to, caused by, or following childbirth, as P. convulsions, P. eclampsia. P. Fever, an acute, febrile disease of women in childbed, due to septic infection. P. Insanity, insanity occurring during the puerperium, usually within five or ten days after delivery. It may take the form of mania (P. mania), melancholia (P. melancholia), or dementia (P. dementia).

Puerperium (*pu-er-pe'-re-um*) [*puer*, child; *parere*, to bear]. The state of a woman in labor or of one who has just been delivered; the period from delivery to the time when the uterus has regained its normal size, which is about six weeks.

Puff-ball. See *Lycoperdon*.

Pugil, Pugillus (*pu'-jil, pu-jil'-us*) [L.]. A handful.

Pulex (*pu'-leks*) [L.]. The flea; an insect parasitic on the skin of man and animals. P. irritans, a species common in Europe and parasitic on the skin of man; its bite causes severe itching and localized swelling. P. penetrans, the chigoe, or jigger-flea, a species the female of which burrows under the skin of the feet to deposit its ova, producing a severe irritation that may proceed to serious inflammation.

Pullulation (*pul-u-la'-shun*) [*pullulare*, to put forth, to bud, to sprout]. The act of sprouting or budding, a mode of reproduction seen, e. g., in the yeast-plant.

Pulmometer (*pul-mom'-et-er*) [*pulmo*, a lung; *metron*, measure]. See *Spirometer*.

Pulmometry (*pul-mom'-et-re*) [*pulmo*, lung; *metron*, measure]. See *Spirometry*.

Pulmonary (*pul'-mon-a-re*) [*pulmo*, lung]. Pertaining to or affecting the lungs, as P. arteries, P. emphysema.

Pulmonic (*pul-mom'-ik*) [*pulmo*, lung]. 1. Pertaining to the lungs; pulmonary. 2. Pertaining to the pulmonary artery, as P. valves. 3. Produced at the P. valve, as P. murmur. P. Fever, croupous pneumonia.

Pulp [*pulpa*]. 1. The soft fleshy part of fruit. 2. The soft part in the interior of an organ, as the P. of the spleen, the P. of a tooth.

Pulpefaction (*pul-pe-fak'-shun*) [*pulpa*, pulp; *facere*, to make]. Conversion into a pulpy substance.

Pulpy (*pul'-pe*) [*pulpa*, pulp]. Resembling pulp; characterized by the formation of a substance resembling pulp.

Pulsatile (*pul'-sat-il*) [*pellere*, to strike]. Pulsating; throbbing.

Pulsatilla (*pul-sat-il'-ah*) [L.]. The herb of Anemone pulsatilla and of Anemone pratensis, of the order Ranunculaceæ, containing a crystalline principle, anemonin, C₁₅H₁₂O₆. P. is employed in amenorrhea, dysmenorrhea, and in inflammations of mucous membranes. Anemonin is used in bronchitis and asthma. Dose gr. 1/3 (0.022). Dose of P. in powder, gr. ij-ijj (0.13-0.20).

Pulsating (*pul'-sa-ting*) [*pellere*, to strike]. Exhibiting pulsation. P. Aorta, the pulsation of the abdominal aorta seen in nervous and anemic persons. P. Empyema, an accumulation of pus in the pleural cavity that transmits the pulsations of the heart.

Pulsation (*pul-sa'-shun*) [*pulsatio*, from *pellere*, to strike]. A beating or throbbing. P., Suprasternal, pulsation at the suprasternal notch. It may be due to aneurysm, dilated aortic arch, or the presence of an anomalous artery.

Pulse (*puls'*) [*pulsus*, the pulse]. The intermittent change in the shape of an artery due to an increase in the tension of its walls

following the contraction of the heart. The pulse is usually counted at the wrist (radial P.), but may be taken over any artery that is palpable, as the temporal, brachial, femoral, dorsalis pedis, etc. P., **Anacrotic**, one the sphygmographic tracing of which is characterized by notches in the ascending limb. P., **Angry**. Synonym of *P. Wiry*. P., **Capillary**, an intermittent filling and emptying of the capillaries of the skin. It is common in aortic regurgitation, and is seen under the finger-nail or on the forehead. P., **Catacrotic**, one with an elevation in the line of descent in the sphygmographic tracing. P., **Cordy**, a tense pulse. P., **Corrigan's**. See *Corrigan's Pulse*. P., **curve**, the tracing of the pulse, called a sphygmogram, made by the sphygmograph. P., **Dicrotic**, one in which the dicrotic wave or recoil wave is exaggerated. It is observed when the arterial tension is low, and gives to the finger the impression of two beats. P., **Entoptic**, the subjective illumination of a dark visual field with each heart-beat, a condition sometimes noted after violent exercise, and due to the mechanic irritation of the rods by the pulsating retinal arteries. P., **Full**, one in which the artery is filled with a large volume of blood and conveys a feeling of being distended. P., **Hard**, one characterized by high tension. P., **High Tension**, one due to increase of the peripheral resistance, together with a corresponding increase in the force of the ventricular systole. It is gradual in its impulse, long in duration, slow in subsiding, with difficulty compressible, and the artery between the beats feels like a firm round cord. P., **Hyperdicrotic**, P., **Hyperdicrotous**, a pulse of which the aortic notch falls below the base line, indicating very low tension, a symptom of great exhaustion. P., **Infrequent**, one the rhythm of which is slower than normal; *i. e.*, in which the heart-beats are fewer in a given time than normal. P., **Intermittent**, one in which one or more beats are dropped. P., **Irregular**, one in which the beats occur at irregular intervals, or in which the force, or both rhythm and force, vary. P., **Jerky**, a pulse in which the artery is suddenly and markedly distended, as in aortic regurgitation. P., **Jugular**, pulsation of the jugular veins in the neck. It is due to tricuspid regurgitation. P., **Locomotive**. Synonym of *Corrigan's Pulse*. P., **Low Tension**, one sudden in its onset, short, and quickly declining. It is easily obliterated by pressure. P., **Paradoxic**, one that is weaker during inspiration, a condition sometimes observed in adherent pericardium. P., **Quick**, one that strikes the finger rapidly, but leaves it also rapidly. P., **Slow**, one indicating a

lengthened systolic contraction of the heart and prolonged diastole,—often used to signify a pulse of slow rate. P., **Soft**, a pulse that is readily compressed. P., **Thready**, one that is scarcely perceptible, feeling like a thread under the finger. P., **Venous**, a pulse observed in a vein. P., **Water-hammer**. See *Corrigan's Pulse*. P., **Wiry**, a small, rapid, tense pulse, feeling like a cord under the finger. It is observed in acute peritonitis.

Pulsimeter (*pul'-sim'-et-er*) [*pulsus*, pulse; *μέτρον*, measure]. An instrument for determining the rate or force of the pulse.

Pulsus (*pul'-sus*) [*L.*]. Pulse. P. **alternans**, one in which there is a regular alternation of strong and weak beats. The weak beat may be imperceptible, in which case two heart-beats correspond to only one beat of the pulse. P. **bigeminus**, one in which the beats occur in pairs, so that a longer pause follows every two beats. P. **celer**, a quick, short pulse. P. **celer et altus**, a quick, full pulse, seen especially in aortic regurgitation. P. **paradoxicus**, P. **paradoxicus**. See *Pulse, Paradoxic*. P. **quadrigeminus**, P. **trigeminus**, a pulse in which a pause occurs after every fourth or third beat respectively.

Pultaceous (*pul'-ta'-shus*) [*puls*, pottage]. Having the consistence of pulp or pottage.

Pulverization (*pul'-ver-i-za'-shun*) [*pulvis*, powder]. The act of reducing a substance to powder.

Pulverulent (*pul'-ver'-u-lent*) [*pulvis*, powder]. Resembling, or of the nature of a powder.

Pulvinar (*pul'-vin-ar*) [*L.*, couch]. The posterior portion of the optic thalamus.

Pulvis [*L.*, *pl. pulveres*]. A powder. P. **antimonialis**. See *Antimony*. P. **aromaticus**, a mixture of cinnamon, ginger, nutmeg, and cardamom seeds; used as a carminative. Dose gr. x-xxx (0.65-2.0). P. **cretæ compositus**, consists of prepared chalk, powdered acacia and sugar, and is used as a mild astringent. Dose, gr. x-3j (0.65-4.0). P. **effervescens compositus**, Seidlitz-powder, a preparation consisting of two powders: the white paper contains 35 grains of tartaric acid, the blue paper, 40 grains of sodium bicarbonate and 2 drams of Rochelle salt. P. **glycyrrhizæ compositus**, consists of senna, licorice, fennel, washed sulphur, sugar; it is used as a laxative. Dose, ʒ ss-ij (2.0-8.0). P. **ipêcacuanhæ et opii**. See *Dover's Powder*. P. **jalapæ compositus**, consists of jalap, 35 parts; cream of tartar, 65 parts; it is used as a hydragogue cathartic. Dose ʒ ss-j (2.0-4.0). P. **rhei compositus**, consists of rhubarb, magnesia, and ginger; it is used as a mild laxative. Dose ʒ ss-j (2.0-4.0).

Pump. An apparatus either drawing up a liquid into its hollow chamber, or, after sucking up the liquid, forcibly ejecting it from one end. **P., Air-**, one used to exhaust the air from a chamber or to force more air into a chamber already filled with air. **P., Breast-**, a pump for removing milk from the breast. **P., Stomach-**, one for removing the contents of the stomach in cases of poisoning.

Pumpkin-seed. The seed of *Cucurbita pepo* of the order Cucurbitaceæ (Pepo of the U. S. P.). The seed is used against tapeworm. Dose $\frac{3}{4}$ iv (128.0).

Puncta (*punk'-tah*) [pl. of *punctum*, a point]. See *Punctum*. **P. dolorosa**, tender or painful points at the exit or in the course of nerves the seat of neuralgia; also called Valdeix's points. **P. lacrimalia**, the orifices of the lacrimal canaliculi in the eyelids near the inner canthus. **P. vasculosa**, minute red spots studding the cut surface of the white central mass of the brain. They are produced by the blood escaping from divided blood-vessels.

Punctate, Punctated (*punk'-tāt, punk'-tāt'*) [*punctum*, point]. Dotted; full of minute punctures.

Punctum (*punk'-tum*) [L.]. A point. **P. cæcum**. See *Blind Spot*. **P. proximum**, See *Near Point*. **P. remotum**. See *Far Point*.

Puncture (*punk'-chūr*) [*pungere*, to prick]. A hole made by a pointed instrument. **P., Lumbar**, puncture of the spinal canal for the withdrawal of cerebrospinal fluid in hydrocephalus, first suggested by Quincke.

Punctured (*punk'-tūrd*) [*pungere*, to prick]. Produced by a prick, as *P. wound*.

Pungent (*pun'-jent*) [*pungere*, to prick]. Acrid; penetrating; producing a pricking or painful sensation.

Pupil (*pu'-pil*) [*pupilla*, a little girl; the name is believed to be derived from the small images seen in the pupil]. The aperture in the iris of the eye for the passage of light. **P., Argyll Robertson**, a myotic pupil that responds on accommodative effort, but not to light,—a condition seen in locomotor ataxia. **P., Artificial**, an aperture made by iridectomy when the normal pupil is occluded.

Pupillary (*pu'-pil-a-re*) [*pupil*]. Pertaining to the pupil. **P. Membrane**, a membrane covering the eye of the fetus until the seventh month of gestation. **P. Membrane, Persistent**. See *Membrane*. **P. Reflex**. See *Reflexes, Table of*.

Pupillometer (*pu'-pil-om'-et-er*) [*pupilla*, pupil; *μέτρον*, a measure]. An instrument for measuring the pupil of the eye.

Pupilloscopy (*pu'-pil-os'-ko-pe*) [*pupilla*,

pupil; *σκοπεῖν*, to inspect]. Examination of the pupil; skiascopy.

Purgation (*pur'-ga'-shun*) [*purgare*, to cleanse]. The act of purging.

Purgative (*pur'-ga-tiv*) [*purgare*, to purge]. 1. Producing purgation. 2. A drug producing copious evacuations of the bowel.

Purging Nut. The seed of *Curcas purgans*, having cathartic properties.

Purge (*purj*) [*purgare*, to purge]. 1. To cause free evacuation of the bowel. 2. A drug that causes free evacuation of the bowel.

Puriform (*pu'-re-form*) [*pus*, pus; *forma*, form]. Resembling pus.

Purkinje (*poor'-kin-ye*) [I. E. *Purkinje*, a Bohemian physiologist, 1787-1869]. **P.'s Cells**, large ganglion-cells of the cerebellar cortex, disposed as a single row at the junction of the nuclear and the molecular layer, and presenting pyriform or flask-shaped bodies, 60-70 μ in their longest diameter. **P.'s Figures**, shadows of the retinal blood-vessels upon the retina. **P.'s Vesicle**, the nucleus of the human ovum; the germinal vesicle.

Purkinje-Sanson's Images. Three pairs of images of one object seen in an observed pupil: the first, erect, reflected from the anterior surface of the cornea; the second, erect, reflected from the anterior surface of the lens; the third, inverted, reflected from the posterior capsule of the lens.

Purple, Visual. See *Rhodopsin*.

Purpura (*pur'-pur-ah*) [L.]. A disease characterized by hemorrhages into the skin, taking the form of petechiæ, maculæ, or large patches. It may occur as an independent affection, or be symptomatic of other diseases. **P. fulminans**, a grave form of *P.*, developing in young children as a sequel to acute infectious diseases. It is of short duration, is marked by extensive extravasations, grave constitutional symptoms, and usually ends fatally. **P. hæmorrhagica**, morbus maculosus Werlhofii; land-scurvy. See *Werlhof's Disease*, under *Diseases, Table of*.

Purpuric (*pur'-pu'-rik*) [*purpura*, purple]. Pertaining to or resembling purpura.

Purpurin (*pur'-pu-rin*) [*purpura*, purple], $C_{16}H_8O_6$. 1. A dye present with alizarin in the madder-root, but also prepared artificially. 2. Uroerythrin, a red coloring matter sometimes present in urinary deposits.

Purring Thrill. A fine trembling vibration like the purring of a cat, perceived by palpation over the precordium. It may be due to aneurysm, or to valvular heart-lesion, especially mitral stenosis.

Purulence (*pu'-ru-lens*) [*pus*, pus]. The state of being purulent.

Purulent (*pu'-ru-lent*) [*pus*, pus]. Having the character of or containing pus; character-

ized by the formation of pus. **P. Catarrh**, an inflammation of a mucous membrane accompanied by the production of pus. **P. Edema**, a general infiltration of pus together with much fluid.

Puruloid (*pu'-ru-loid*) [*pus*, *pus*; *ειδος*, like]. Resembling pus; puriform.

Pus [*L.*]. A liquid substance consisting of cells and an albuminous fluid (liquor puris), formed in certain kinds of inflammation.

P., Blue, pus colored blue by the bacillus pyocyaneus. **P.-corpuscles**, the corpuscles found in pus. **P., Curdy**, pus containing cheesy-looking flakes. **P., Ichorous**, pus that is thin and acrid. **P., Laudable**, a whitish, inodorous pus, formerly thought to be essential to the healing of wounds. **P., Sanious**, pus mixed with blood. **P.-tube**.

See *Pyosalpinx*.
Pustula maligna. Anthrax.
Pustulant (*pus'-tu-lant*) [*pustula*, a pustule]. 1. Causing the formation of pustules. 2. An irritant substance giving rise to the formation of pustules.

Pustular (*pus'-tu-lar*) [*pustula*, pustule].

Characterized by the presence of pustules.

Pustulation (*pus-tu'-la'-shun*) [*pustula*, pustule]. The formation of pustules.

Pustule (*pus'-tūl*) [*pustula*, pustule]. A small circumscribed elevation of the skin containing pus. **P., Malignant**, anthrax.

Pustulocrustaceous (*pus-tu-lo-krus-ta'-shus*) [*pustula*, pustule; *crusta*, crust].

Characterized by the formation of pustules and crusts.

Putamen (*pu-ta'-men*) [*putamen*, husk]. 1.

The stone of a drupe, or shell of a nut. 2.

The outer darker part of the lenticular nucleus of the brain.

Putrefaction (*pu-tre-fak'-shun*) [*putridus*, putrid; *facere*, to make].

The decomposition of nitrogenous organic matter under the influence of microorganisms, accompan-

ied by the development of disagreeable odors, due to the evolution of ammonia, hydrogen sulphid, and other gases, and the production of aromatic bodies. In addition,

many other compounds are formed, among which ptomains are the most important. The

end-products are water, nitrogen, methane, and carbon dioxide.

Putrefactive (*pu-tre-fak'-tiv*) [*putridus*, putrid; *facere*, to make]. Pertaining to or causing putrefaction.

Putrescent (*pu-tres'-ent*) [*putrescere*, to become rotten]. Undergoing putrefaction.

Putrescin (*pu-tres'-in*) [*putrescere*, to become rotten], $C_4H_{12}N_2$. A poisonous ptomain, a clear, rather thin liquid, of a disagreeable odor, boiling at 156° – $157^{\circ}C$. See *Ptomains*, Table of.

Putrid (*pu'-brid*) [*putridus*, rotten]. Rot-

ten; characterized by putrefaction. **P. Fever**. Synonym of *Typhus Fever*.

Putrilage (*pu'-tril-āj*) [*putris*, rotten].

Putrescent material.

Pyæmia (*pi-ē'-me-ah*). See *Pyemia*.

Pyarthrosis (*pi-ar-thro'-sis*) [*πύον*, pus; *ἄρθρον*, joint]. Suppuration of a joint.

Pyelitis (*pi-el-ī'-tis*) [*πύελος*, a trough; *τις*, inflammation].

Inflammation of the pelvis of the kidney. It may be due to the irritation of calculi, to tuberculosis, or to acute

specific fevers, but most commonly is secondary to diseases of the bladder. It is

marked by pain and tenderness in the lumbar region, by a remittent fever, and by the pres-

ence in the urine of albumin, mucus, epithelial cells from the pelvis of the kidney, pus-

corpuscles in large amount, and frequently blood. The urine is generally acid. **P.,**

Calculus, that due to calculi.

Pyelonephritis (*pi-el-o-nef-rī'-tis*) [*πύελος*, trough; *νεφρός*, kidney; *τις*, inflammation].

Inflammation of the kidney and its pelvis.

Pyelonephrosis (*pi-el-o-nef-rō'-sis*). Syn-

onym of *Pyelonephritis*.

Pyemia (*pi-ē'-me-ah*) [*πύον*, pus; *αἷμα*, blood].

A disease due to the presence of pyogenic microorganisms in the blood and

the formation, wherever these organisms lodge, of embolic or metastatic abscesses.

The symptoms are intermittent or remittent fever, associated with sweats and chills, rapid

emaciation, slight jaundice, abscesses in dif-

ferent parts of the body, and often a purpuric eruption. The disease is generally fatal.

P., Arterial, a name given to pyemia pro-

duced by disorganization of a cardiac throm-

bos and the dissemination of emboli through

the arterial circulation.

Pyemic (*pi-em'-ik*, *pi-ē'-mik*) [*πύον*, pus; *αἷμα*, blood]. Pertaining to or affected with

pyemia.

Pygodidymus (*pi-go-did'-im-us*) [*πυγή*, buttock; *δίδυμος*, twins]. A double monster

united by the buttocks.

Pygomelus (*pi-gom'-el-us*) [*πυγή*, buttock; *μέλος*, member]. A monster with a parasite

attached to the hypogastric region or to the

buttock.

Pygopagus (*pi-gop'-ag-us*) [*πυγή*, buttock; *παγος*, joined]. A monster with conjoined

buttocks or backs.

Pyin (*pi'-in*) [*πύον*, pus]. An albuminous

substance of complex constitution occurring

in pus. It may be separated by adding

sodium chloride and filtering.

Pylephlebitis (*pi-le-fleb-ī'-tis*) [*πύλη*, gate; *φλέψ*, vein; *τις*, inflammation].

Inflammation of the portal vein. The condition

is usually secondary to disease of the intestines,

is generally suppurative in character, and

gives rise to the symptoms of pyemia.

Pylethrombosis (*pi-le-throm-bo'-sis*) [πύλη, gate; θρόμβος, clot]. Thrombosis of the portal vein.

Pylorectomy (*pi-lo-rek'-to-me*) [πυλωρός, a gate-keeper; εκτομή, excision]. Excision of the pylorus.

Pyloric (*pi-lo'-rik*) [πυλωρός, a gate-keeper]. Pertaining to the pylorus. **P. Glands**, glands situated in the region of the pylorus and secreting the gastric juice.

Pylorus (*pi-lo'-rus*) [πυλωρός, gate-keeper]. 1. The circular opening of the stomach into the duodenum. 2. The fold of mucous membrane and muscular tissue surrounding the aperture between the stomach and the duodenum.

Pyo- (*pi'-o*) [πύον, pus]. A prefix denoting pertaining to pus.

Pyocolpos (*pi-o-kol'-pos*) [πύον, pus; κόλπος, vagina]. An accumulation of pus within the vagina.

Pyoctanin (*pi-ok'-tan-in*). See *Pyoktanin*.

Pyocyanin (*pi-o-si'-an-in*) [πύον, pus; κύανος, blue], $C_{14}H_{14}NO_2$. A colored substance derived from blue pus, and from cultures of the bacillus pyocyaneus.

Pyocyte (*pi'-o-sit*) [πύον, pus; κύτος, cell]. A pus-corpusele.

Pyogenic (*pi-o-jen'-ik*) [πύον, pus; γεννάν, to beget]. Producing pus. **P. Membrane**, the thin, yellow layer of tissue forming the wall of an abscess. **P. Microorganisms**, the microorganisms producing pus. The ordinary **P. microorganisms** are the staphylococci and the streptococci. Under certain circumstances pus may be produced by the pneumococcus of Fränkel, the bacillus coli communis, the bacillus of typhoid fever, the gonococcus, and others.

Pyohæmia (*pi-o-he'-me-ah*). See *Pyemia*.

Pyoid (*pi'-oid*) [πύον, pus; εἶδος, like]. Resembling pus.

Pyoktanin (*pi-ok'-tan-in*) [πύον, pus; κτείνειν, to kill]. A name given to methyl-violet and methyl-blue on account of their germicidal properties. **P.** has been used in diphtheria, cystitis, gonorrhea, ulcers, and inflammations of the conjunctiva, and as an injection in carcinoma.

Pyometra (*pi-o-me'-trah*) [πύον, pus; μήτρα, womb]. A collection of pus in the uterus.

Pyonephrosis (*pi-o-nef-ro'-sis*) [πύον, pus; νεφρός, kidney]. An accumulation of pus in the pelvis of the kidney.

Pyopericarditis (*pi-o-per-e-kar-di'-tis*) [πύον, pus; pericarditis]. Suppurative pericarditis.

Pyophthalmia (*pi-off-thal'-me-ah*) [πύον, pus; ὀφθαλμός, eye]. Purulent ophthalmia.

Pyophylactic (*pi-o-fil-ak'-tik*) [πύον, pus; φυλάσσειν, to guard]. Protecting against pus. **P. Membrane**, the pyogenic membrane.

Pyopneumothorax (*pi-o-nu-mo-tho'-raks*) [πύον, pus; πνεῦμα, air; θώραξ, thorax]. An accumulation of air or gas and pus in the pleural cavity.

Pyorrhea (*pi-or-e'-ah*) [πύον, pus; ῥοία, a flow]. A purulent discharge.

Pyosalpinx (*pi-o-sal'-pinks*) [πύον, pus; σάλπιγξ, tube]. An accumulation of pus in the Fallopian tube.

Pyothorax (*pi-o-tho'-raks*) [πύον, pus; θώραξ, thorax]. An accumulation of pus in the pleural cavity; empyema.

Pyoxanthin, Pyoxanthose, (*pi-o-san'-thin, pi-o-san'-thōs*) [πύον, pus; ξανθός, yellow]. A yellow substance sometimes found in pus, and resulting from the oxidation of pyocyanin.

Pyramid (*pir'-am-id*) [πυραμῖς]. A solid having a polygonal base and triangular planes for its sides, which meet at a point called the apex or vertex. **P., Anterior**, one of the two pyramidal bundles of white matter on either side of the anterior median fissure of the medulla. **P. of the Cerebellum**, a conic projection forming the central portion of the inferior vermiform process. **P. of Ferrein**, one of the prolongations of the Malpighian pyramid into the cortex of the kidney; it is also known as medullary ray. **P., Malpighian**, one of the conic masses composing the medullary substance of the kidney. **P., Posterior**, one of the two narrow bundles of white matter placed on either side of the posterior median fissure of the medulla oblongata. They are continuous with the posterior median columns of the spinal cord.

Pyramidal (*pe-ram'-id-al*) [πυραμῖς, pyramid]. Shaped like a pyramid. **P. Tracts**. See *Tracts*.

Pyramidalis (*pe-ram-id-a'-lis*). Pyramidal, as **P. muscle**. See *Muscles, Table of*.

Pyrethrum (*pi-re'-thrum*). Pellitory; the root of *Anacyclus pyrethrum*, a plant of the order Compositæ. **P.** is used as a sialogogue and masticatory in headache, toothache, and neuralgic affections of the face. It is employed either in powder or in the form of the tincture, Tinctura pyrethri (U. S. P.).

Pyretic (*pi-ret'-ik*) [πυρετός, fever]. Pertaining to or affected with fever.

Pyretin (*pi'-re-tin*) [πυρετός, fever]. A name given to various empyreumatic substances found in pitch.

Pyretogenic, Pyretogenous (*pi-ret-oi-en'-ik, pi-ret-oi'-en-us*) [πυρετός, fever; γεννάν, to produce]. Causing or producing fever.

Pyretogenin (*pi-ret-oi'-en-in*) [πυρετός, fever; γεννάν, to produce]. A substance formed by microorganisms, and said to have the property of producing fever when inoculated into animals.

Pyretology (*pi ret-ol'-o-je*) [*πυρετός*, fever; *λόγος*, treatise]. The science on the nature of fevers.

Pyrexia (*pi-riks'-e-ah*) [*πύρεξις*, fever]. Elevation of temperature above the normal; fever.

Pyrexial (*pi-riks'-e-al*) [*πύρεξις*, fever]. Pertaining to pyrexia.

Pyridin (*pi-rid-in*) [*πύρ*, fire], C_5H_5N . A liquid base obtained as a distillation-product from tobacco, coal tar, and other organic matter, and forming the first of a long and important series of homologous bases. P. has been used in asthma by inhalation. P. **Tricarboxylic Acid**, $C_5H_2(COOH)_3N$, is antiseptic, antipyretic, antiperiodic, and antispasmodic. It has been used in malaria, asthma, and typhoid fever. Dose gr. ij-x (0.13-0.65).

Pyriform (*pi-'e-form*) [*pyrus*, pear; *forma*, a form]. Pear-shaped.

Pyriformis (*pi-'e-for'-mis*). Pyriform, as P. muscle. See *Muscles*, Table of.

Pyro- (*pi-'ro*) [*πύρ*, fire]. A prefix signifying fire or heat.

Pyroarsenic Acid (*pi-ro-ar-sen'-ik*) [*πύρ*, fire; *arsenium*, arsenic], $H_4As_2O_7$. A tetrabasic acid produced when arsenic is heated to $180^\circ C$.

Pyroboric Acid (*pi-ro-bor'-ik*) [*πύρ*, fire; *boron*], $H_2B_4O_{10}$. A dibasic acid produced by heating boric acid.

Pyrocatechin (*pi-ro-kat'-e-chin*) [*πύρ*, fire; *catechu*], $C_6H_4O_2$, $C_6H_4(OH)_2$. Catechol; a crystalline substance formed by the dry distillation of catechu and sometimes occurring in the urine. It has been used as an antipyretic. Dose gr. j-ij (0.005-0.13).

Pyrodim (*pi-ro'-din*) [*πύρ*, fire], $C_6H_5.C_2H_5.O.N_2H_2$. Hydrazetin, acetyl-phenyl-hydrazin; a crystalline, poisonous substance, used as a substitute for chrysarobin in psoriasis, and in other cutaneous affections; also as an antipyretic. Dose gr. $\frac{1}{2}$ - $\frac{1}{3}$ (0.03-0.2).

Pyrogallic Acid (*pi-ro-gal'-ik*). See *Acid*.

Pyrogenic (*pi-ro-jen'-ik*) [*πύρ*, fire; *γεννάν*, to produce]. Producing fever.

Pyroligneous (*pi-ro-lig'-ne-us*) [*πύρ*, fire; *lignum*, wood]. Pertaining to the destructive distillation of wood. P. Acid, wood-vinegar. See *Acid*, *Pyroligneous*.

Pyromania (*pi-ro-ma'-ne-ah*) [*πύρ*, fire; *μανία*, madness]. A monomania for incendiarism.

Pyrometer (*pi-rom'-et-er*) [*πύρ*, fire; *μέτρον*, measure]. An instrument for measuring the intensity of heat of too high a degree to be estimated by the ordinary thermometer.

Pyrophosphoric Acid (*pi-ro-fos-for'-ik*).

See *Acids*, Table of.

Pyrosis (*pi-ro'-sis*) [*πύρ*, fire]. An affection of the stomach characterized by a burning sensation, accompanied by eructations of an acrid, irritating fluid; heartburn.

Pyrosoma bigemina (*pi-ro-so-mah bi-jem'-in-ah*) [*pyrus*, pear; *σώμα*, body]. The parasite supposed to cause Texas fever in cattle.

Pyrotoxina bacterica (*pi-ro-toks'-in-ah bak-ter'-ik-ah*) [*πύρ*, fire; *τοξικόν*, poison]. A pyrogenic substance believed to be produced by many forms of bacteria.

Pyroxylin (*pi-roks'-il-in*) [*πύρ*, fire; *ξύλον*, wood]. Gun-cotton; cotton-fiber treated with a mixture of nitric and sulphuric acids, by which the cellulose is changed into various nitrocompounds. Soluble gun-cotton, Pyroxylinum of the U. S. P., is used in the preparation of collodium. The explosive gun-cotton is the hexanitrate of cellulose.

Pyrrol (*pir'-ol*) [*πύρ*, fire; *oleum*, oil], $C_4H_7(NH)$. A liquid base obtained in the distillation of Dippel's oil and other organic substances. P. tetriodid. See *Iodol*.

Pythogenic (*pi-tho-jen'-ik*) [*πιθαί*, to rot; *γεννάν*, to produce]. Producing or arising from decomposition. P. Fever. Synonym of *Typhoid Fever*.

Pyuria (*pi-u'-re-ah*) [*πύρ*, pus; *urina*, urine]. The passage of urine containing pus.

Q

Q. S. Abbreviation of *quantum sufficit*—as much as suffices.

Quack (*kwak*). One who practices quackery; a pretender of medical skill; a vender of nostrums; a medical charlatan.

Quackery (*kwak'-er-ee*). The practice of medicine by a quack.

Quadrangular (*kwo-d-rang'-gu-lar*) [*quad-*

rangulum, a four-cornered figure]. Having four angles, as the Q. lobe of the cerebellum.

Quadrant (*kwo-d'-rant*) [*quadratus*, squared].

1. The fourth part of a circle, subtending an angle of 90 degrees. 2. One of the four regions into which the abdomen may be divided for purposes of physical diagnosis.

Quadrante (*kwo-d'-rāt*) [*quadratus*, square].

- Square; four-sided. **Q. Lobule.** See *Pre-cuneus*.
- Quadratus** (*kwod-ra'-tus*) [L.]. Squared; having four sides. **Q. Muscle.** See *Muscles, Table of*.
- Quadri-** (*kwod'-re-*) [L.]. A prefix denoting four, or four times.
- Quadribasic** (*kwod-re-ba'-sik*) [*quadri*, four; *basis*, base]. In chemistry, applied to an acid having four replaceable hydrogen-atoms.
- Quadriceps** (*kwod'-re-seps*) [*quadri*, four; *caput*, head]. Four-headed, as a Q. muscle. See *Muscles, Table of*.
- Quadrigeminal** (*kwod-re-jem'-in-al*) [*quadrigeninus*, fourfold]. Fourfold; consisting of four parts, as the Q. bodies. See *Corpora quadrigemina*.
- Quadiurate** (*kwod-re-u'-rat*) [*quadri*, four; *urate*]. A term applied to the hyperacid urate of human urine, and the urine of birds and reptiles. The quadriurates have the general formula of $MH(C_5H_3N_4O_8) \cdot H_2C_3H_2 \cdot N_4O_8$.
- Quadrivalent** (*kwod-riv'-al-ent*) [*quadri*, four; *valere*, to be worth]. In chemistry, having a combining power equivalent to that of four hydrogen-atoms.
- Quadruplet** (*kwod-ru'-plet*) [*quadruplare*, to make fourfold]. Any one of four children born at one birth.
- Quaker-button.** A popular name for *nux vomica*.
- Qualitative** (*kwol'-e-ta-tiv*) [*qualitas*, quality]. Pertaining to quality. **Q. Analysis.** See *Analysis*.
- Quantitative** (*kwon'-tit-a-tiv*) [*quantus*, how much; *valere*, to be worth]. Pertaining to quantity. **Q. Analysis.** See *Analysis*.
- Quantivalence** (*kwon-tiv'-al-ens*) [*quantus*, how much; *valere*, to be worth]. The combining power of an element or radicle expressed in terms of the number of atoms of hydrogen with which it will unite. Univalent or monad atoms, as chlorine, are saturated with one atom. Bivalent or dyad atoms require two; trivalent or triad, as boron, take three; quadrivalent or tetrad, quinquivalent or pentad, sexvalent or hexad, require two, three, four, five, and six atoms of hydrogen respectively.
- Quarantine** (*kwor'-an-ten*) [It., *quaranta*, forty]. The time (formerly forty days) during which vessels or travellers from ports infected with contagious or epidemic diseases are required by law to remain outside the port of their destination, as a safeguard against the spreading of such diseases. 2. Also, the place of detention. 3. The act of detaining vessels or travellers from suspected ports or places for purposes of inspection or disinfection. **Q., Land-**, the isolation of a person or district on land for similar purposes.
- Quart** (*kwort*) [*quartus*, fourth]. The fourth part of a gallon.
- Quartan** (*kwor'-tan*) [*quartus*, fourth]. 1. Recurring on the fourth day. 2. A form of intermittent fever, the paroxysms of which occur every fourth day. **Q., Double**, a variety characterized by milder and severer paroxysms, each occurring every fourth day.
- Quarter-evil** (*kwor'-ter-e-vil*). An infectious disease of cattle prevalent during the summer-months, and characterized by the appearance of irregular emphysematous swellings of the subcutaneous tissue and muscles, especially over the quarters, hence the name. It is also called symptomatic anthrax, or black-leg.
- Quartipara** (*kwor-tip'-ar-ah*) [*quartus*, fourth; *parere*, to bring forth]. A woman in her fourth pregnancy. See *Multipara*.
- Quartiparous** (*kwor-tip'-ar-us*) [*quartus*, fourth; *parere*, to bring forth]. Pregnant four times.
- Quassation** (*kwass-a'-shun*) [*quassatio*, a shaking or shattering]. The reduction of barks, roots, and other drugs to morsels, in preparation for further pharmaceutic treatment; cassation. *
- Quassia** (*kwosh'-e-ah*) [after *Quassi*, a negro slave who first used it]. The wood of several trees of the order Simarubæ. **Q.** of the U. S. P. and B. P. is the wood of *Picræna excelsa*, and contains the bitter principle, quassin, $C_{22}H_{44}O_{10}$. **Q.** is a simple bitter, and is used in dyspepsia and constipation; in the form of an enema it is employed against seat-worms. Preparations: Extractum quassiæ (U. S. P., B. P.), dose gr. j-ij (0.065-0.2). Extractum quassiæ fluidum (U. S. P.), dose fʒ ss-j (2.0-4.0). Infusum quassiæ (B. P.), dose fʒ ij (64.0). Tinctura quassiæ (U. S. P.) dose, fʒ j (4.0). Quassin, dose gr. ʒ (0.006).
- Quassin** (*kwos'-in*). See *Quassia*.
- Quaternary** (*kwat-ter'-na-re*) [*quaterni*, four each]. 1. Consisting of four elements. 2. Fourth in order.
- Quebrachin** (*ke-brah'-kin*). See *Quebracho*.
- Quebracho** (*ke-brah'-ko*) [from Pg. *quebracho*, ax-breaker]. The *Aspidosperma quebracho-blanco*, of the order Apocynaceæ. It contains the following alkaloids:—Aspidospermin, $C_{22}H_{40}N_2O_8$; aspidospermatin, $C_{22}H_{40}N_2O_8$; aspidosamin, C_{77} ; quebrachin, $C_{21}H_{38}N_2O_8$, and quebrachamin. **Q.** is used in emphysema, bronchitis, and in asthma. Extractum adispermatis fluidum (U. S. P.), dose m xv-fʒ j (1.0-4.0). Dose of aspidospermin gr. ʒ-½ (0.016-0.032).
- Queen-root.** See *Stillingia*.
- Quercin** (*kwor'-sin*) [*quercus*, oak], $C_6H_4(OH)_4$. A bitter crystallizable carbohydrate extracted from acorns and oak-bark.

Quercitannic Acid (*kwer-se-tan'-ik*) [*quercus*, oak; *tannin*, tannin], $C_{11}H_{18}O_9$. A variety of tannic acid found in oak-bark.

Quercite (*kwer'-it*) [*quercus*, oak], $C_6H_7(OH)_5$. A sweet principle found in acorns.

Quercitrin (*kwer'-sit-rin*) [*quercus*, oak; *citrus*, lemon], $C_{30}H_{50}O_{30}$. A glucosid found in the bark of *Quercus tinctoria*.

Quercus (*kwer'-kus*) [L.]. The oak. See *Oak*.

Quicklime [AS., *cwic*, alive; *lime*]. Calcic oxid.

Quicksilver [AS., *cwic*, alive; *silver*]. Mercury.

Quickening (*kwi'-en-ing*) [AS., *cwic*, alive]. The first feeling on the part of the pregnant woman of fetal movements, occurring between the fourth and fifth month of pregnancy.

Quill-suture. See *Suture*.

Quillaja (*kwi'-d'-yah*) [Chilean, *quillean*, to wash]. A genus of trees of the order Rosaceæ. The bark of *Q. saponaria* [Quillaja, U. S. P.], soap-bark, contains saponin, and produces a froth when agitated in water. *Q.* is used in pulmonary affections and as a sternutatory, and in the arts as a substitute for soap. *Tinctura quillajæ* (U. S. P.), dose [3j] (4.0).

Quince (*quins*). See *Cydonium*.

Quincke's Capillary Pulse. See *Pulse, Capillary*.

Quincke's Operation. See *Puncture, Lumbar*.

Quinidin (*kwin'-id-in*) [Peruvian, *kina*, bark], $C_{20}H_{21}N_3O_9$. An alkaloid of cinchona-bark isomeric with quinin, which it resembles in action, differing only in being less powerful. **Q. sulphate** (Quinidinæ sulphas U. S. P.) is used as an antiperiodic, in doses of gr. xx-lx (1.3-4.0).

Quinic (*kwin'-ik*) [Peruvian, *kina*, bark], Pertaining to quinin. **Q. Acid**, $C_7H_{12}O_6$, an acid occurring in cinchona-bark, in the ivy, oak, elm, ash, coffee-plant, etc.

Quinin (*kwi'-nin* and *kwin'-en*) are common pronunciations, but as the Fr. *qu* is merely copied from the Sp. *qu* (which had a *k* sound) and as, moreover, the Peruvian word, from which our word was originally derived, had a *k* sound (without the *u* element), it follows that, at least, from an etymologic standpoint, our pronunciation should be *kin-en'* [Peruvian, *kina*, bark], $C_{20}H_{21}N_3O_9 \cdot 3H_2O$. A bitter amorphous or crystalline alkaloid obtained from the bark of various species of cinchona. It is soluble in 900 parts of water, readily soluble in alcohol, ether, and in chloroform, and gives a beautiful emerald-green color when it or its salts are treated with a solution of chlorin and then with ammonia. *Q.* acts as a stimulant to the nervous system, causing in large doses cere-

bral congestion and lessening of the reflexes; it is a slight respiratory stimulant and a depressant to the circulation; it lessens the ameboid movement of the white corpuscles, and during fever is strongly antipyretic; it also possesses antiseptic properties. In large doses it causes ringing in the ears, a feeling of fullness in the head, dizziness, slight deafness, and at times disturbances of vision; occasionally also a rise of temperature (*Q.-fever*). *Q.* is used as an antiperiodic in malaria, in which disease it has a specific action; it is also employed as an antipyretic in other febrile affections, as a tonic in convalescence, as a stimulant to the uterus during parturition, in whooping-cough, coryza, and in hay-fever. Doses of *Q. sulphate* (Quininæ sulphas, U. S. P., B. P.), the salt most commonly employed—in malaria gr. v-xxiv (0.32-1.6) before the paroxysms; and gr. ij-iv (0.13-0.26) as a prophylactic; as a tonic gr. j-ij (0.065-0.13); in whooping-cough, gr. ias (0.10) for each year of the child's age, or locally in solution of gr. i-ij (0.065-0.13) to the ounce (32.0) by the atomizer. Other salts, given in about the same doses as the sulphate are: **Q. bisulphate** (Quininæ bisulphas, U. S. P.); **Q. hydrobromate** (Quininæ hydrobromas, U. S. P.); **Q. hydrochlorate** (Quininæ hydrochloras, U. S. P.); **Quinin and urea hydrochlorate**, employed for hypodermic use chiefly. **Q. sulphovinate** has been used hypodermically; **Q. tannate** has been employed in whooping-cough; **Q. valerianate** is used in doses of gr. j-ij (0.065-0.13) in nervous debility and hemicrania. From *Q.* are prepared *Ferri et quininæ citras* and *Ferri et quininæ citras solubilis*. See *Ferrum*. From *Q. sulphate*, *Syrupus ferri, quininæ et strychninæ phosphatum* is prepared. **Quininism, Quinism** (*kwin'-in-izm*, *kwin'-izm*) [Peruvian, *kina*, bark]. Cinchonism. **Quinoidin** (*kwin'-oi'-din*). See *Chinoidinum*. **Q., Animal-**, a basic substance obtained from animal tissues and having the property of fluorescence like quinin. **Quinolin** (*kwin'-o-lin*) [*quinin*], C_8H_7N . A liquid alkaloid obtained in the destructive distillation of quinin, or cinchonin, with potassium hydroxid. It is antipyretic and antiseptic. **Q. hydrochlorate**, **Q. salicylate**, and **Q. tartrate**, have been used. Dose of the last gr. xv (1.0).

Quinone (*kwin'-on*) [*quinin*], $C_6H_4O_2$. A yellow crystalline substance obtained by heating quinic acid with manganese dioxide and sulphuric acid. It is the lowest member of a series of bodies known as quinones.

Quinotannic Acid (*kwin-o-tan'-ik*) [*quinin*; *tannin*]. A form of tannic acid found in cinchona-bark.

Quinovin (*kwín'-o-vín*) [Peruvian, *kina*, bark], $C_{20}H_{30}O_{11}$. Kinovin, a bitter glucosid found in cinchona-bark.

Quinivalent (*kwín-kwín'-al-ent*) [*quinque*, five; *valere*, to be worth]. Having a valence of five; capable of combining with five atoms of hydrogen or their equivalent.

Quinsy (*kwín'-ze*) [*κίων*, dog; *ἀγχειν*, to choke]. Acute inflammation of the tonsils, usually tending to suppuration.

Quintan (*kwín'-tan*) [*quintus*, fifth]. An intermittent fever, the paroxysms of which recur every four days, *i. e.*, on the 5th, 9th, 13th, etc.

Quintipara (*kwín-típ'-a-rah*). See *Multi-para*.

Quintuplet (*kwín-tu'-plet*) [*quintuplex*, five-fold]. One of five children born at one time.

Quitch-grass (*kwitch'-gras*). See *Triticum repens*.

Quiz (*kwis*) [*quæere*, to ask (?)]. A recitation, conducted by questions and answers, in which the student familiarizes himself with his studies.

Quizzer (*kwiz'-er*) [*quæere*, to ask (?)]. One who conducts a quiz.

Quotidian (*kwot-id'-e-an*) [*quot*, as many as; *dies*, day]. 1. Recurring every day. 2. An intermittent fever, the paroxysms of which recur daily. **Q.**, Double, a fever having two paroxysms a day, usually differing in character.

R

R. The abbreviation of *Recipe*, take; of right, of Réaumur, and of resistance (electric).

Rabic (*rab'ik*) [*rabies*, rage]. Pertaining to rabies, as *R. virus*.

Rabid (*rab'id*) [*rabidus*, mad]. Affected with rabies, or hydrophobia; pertaining to rabies, as *R. virus*.

Rabies (*ra'-be-zs*) [L.]. Lyssa, or hydrophobia. The latter term is generally applied to the human disease consequent upon the bite of a rabid dog or other animal. **Rabies** is an acute infectious disease of animals dependent upon a specific virus, and communicable to man by inoculation. All animals are liable to the disease, but it occurs most frequently in the wolf, the cat, and the dog, and is chiefly propagated by the latter, which is specially susceptible. The nature of the poison is as yet unknown; it has a special affinity for the nervous system, and is found in the secretions, particularly in the saliva. See *Hydrophobia*.

Racemose (*ras'-e-mōs*) [*racemus*, a bunch of grapes]. Resembling a bunch of grapes, as *R. gland*. **R. Aneurysm**, aneurysm by anastomosis.

Rachialgia (*ra-ke-al'-je-ah*) [*ράχις*, spine; *ἄλγος*, pain]. Neuralgic pain in the spinal column.

Rachidial (*ra-kid'-e-al*) [*ράχις*, the spine]. Pertaining to the spine.

Rachidian (*ra-kid'-e-an*) [*ράχις*, spine]. Spinal; vertebral.

Rachilysis (*ra-kil'-is-is*) [*ράχις*, spine; *λύσις*, a loosening]. A method of correcting lateral curvature of the spine by appropriate traction and pressure.

Rachiodynia (*ra-ke-o-din'-e-ah*) [*ράχις*, spine; *δύσπη*, pain]. Spasmodic pain in the spinal column.

Rachiotomy (*ra-ke-ot'-o-me*) [*ράχις*, spine; *τέμνειν*, to cut]. 1. The operation of cutting into or through the vertebral column. 2. The operation of cutting through the spine of the fetus to facilitate delivery.

Rachischisis (*ra-kis'-kis-is*). See *Spina bifida*.

Rachitic (*ra-kit'-ik*) [*ράχις*, spine; *ιτις*, inflammation]. Affected with, resembling, or produced by rachitis; rickety. **R. Rosary**, the row of nodules appearing on the ribs, at their junction with the cartilages, in rachitis.

Rachitis (*ra-ki'-tis*) [*ράχις*, spine; *ιτις*, inflammation]. Rickets, a constitutional disease of infancy, characterized by impaired nutrition and changes in the bones, the symptoms being a diffuse soreness of the body, slight fever, and profuse sweating about the head and neck, and changes in the osseous system, consisting in a thickening of the epiphyseal cartilages and periosteum and a softening of the bones. Through the action of the muscles on the soft bones various deformities are produced, while the perosteal hyperplasia leads to nodular hyperostoses, especially about the head, giving the latter a square appearance (*caput quadratum*). Dentition and closure of the fontanels are delayed. Nervous symptoms are often present, as feverishness, laryngismus stridulus, and convulsions. The liver and spleen are usually enlarged. The etiology is obscure—it has been ascribed to deficiency in the earthy salts, to defect in the osteoblasts, and to

microspinal infection. *R. adularum*, osteomielitis; mollities osium.

Radseyyge (*rad-de-sei'-ge*) [*Norwegian*]. A disease also known as Scandinavian syphilis, or Norwegian leprosy, and characterized by ulceration and other cutaneous lesions. It is probable that under the name of *R.* are included syphilitic and leprosy lesions.

Radial (*rad-de-al*) [*radius*, a spoke; *ad*, toward]. Toward the radial side.

Radial (*rad-de-al*) [*radius*, a spoke]. 1.

Radiating; diverging from a common center.

2. Pertaining to or in relation with the radius or bone of the forearm, as the *R.* artery.

Radiant (*rad-de-ant*) [*radius*, a spoke].

Radiating; diverging from a common center.

Radiate (*rad-de-it*) [*radius*, a spoke]. Diverging from a central point.

Radiation (*ra-de-ah'-shun*) [*radiare*, to radiate].

1. The act of radiating or diverging from a central point, as *R.* of light; divergence from a center; having the appearance of rays. 2. In cerebral anatomy, certain groups of fibers that diverge after leaving their place of origin. *R.*, Optic, a large strand of fibers continuous with those of the corpus radiata, derived mainly from the pulvinar, the external and internal geniculate bodies, and the optic tract, and radiating into the occipital lobes. *R.*, Striothalamic, a system of fibers connecting the corpus striatum with the optic thalamus and the subthalamic region. *R.*, Thalamic, certain tracts of fibers from the optic thalami, that radiate into the hemispheres.

Radical (*rad-ik-al*) [*radix*, a root]. 1. Belonging to the root; going to the root, or attacking the cause of a disease, as a radical operation. 2. See *Radicle* (2d def.).

Radicle (*rad-ik-l*) [*dim. of radix*]. 1. A little root, as *R.* of a nerve, one of the ultimate fibrils of which a nerve is composed; *R.* of a vein, one of the minute vessels uniting to form a vein. 2. An atom or group of atoms constituting the essential part of the molecule of a compound; a group of atoms that remains intact in the ordinary decomposition of the molecule; especially a group of atoms that acts in combination as a simple element but is incapable of existence in the free state, as NH_4 , ammonium, C_6H_5 , phenyl. *R.*, Alcohol, one that forms an alcohol when combined with the elements of water.

Radicular (*rad-ik'-u-lar*). Pertaining to a root or to a radicle; specifically, pertaining to the roots of the spinal nerves.

Radio- (*ra'-de-o-*) [*radius*, a spoke]. A prefix denoting connection with the radius.

Radiobicipital (*ra-de-o-bi-sip'-it-al*) [*radius*, radius; *biceps*, two-headed]. Pertaining to the radius and the biceps.

Radiocarpal (*ra-de-o-kar'-pal*) [*radius*, radius; *carpus*, carpus]. Pertaining to the radius and the carpus.

Radiodigital (*ra-de-o-dij'-it-al*) [*radius*, radius; *digitus*, a digit]. Pertaining to radius and the finger.

Radiography (*ra-de-og'-ro-fi*). See *Sinigraphy*.

Radiohumeral (*ra-de-o-har'-mer-al*) [*radius*, radius; *humerus*, the humerus]. Pertaining to the radius and the humerus.

Radiopalmar (*ra-de-o-pal'-mar*) [*radius*, radius; *palmus*, palm]. Pertaining to the radius and the palm.

Radioulnar (*ra-de-o-ul'-nar*) [*radius*, radius; *ulna*, ulna]. Pertaining to the radius and ulna.

Radius (*rad-de-us*) [*radius*, a spoke of wheel]. 1. A ray. 2. The outer of two bones of the forearm.

Radix (*rad-dir*) [*L.*]. A root.

Radzyge (*rad-si'-ge*). See *Radseyyge*.

Rag-picker's disease, "Haderstrækkel." An acute febrile disease occurring in women engaged in sorting rags in paper-factories. It is supposed to be due to the inhalation of anthrax-bacilli or spores, and is characterized by an exudation into the pulmonary tissue bronchial glands, and pleural cavity.

Railway-spine. A term given by Erichsen to a varied group of spinal symptoms consequent on slight injuries or concussions received in railway accidents. The condition is classed with the traumatic neuroses, and a form of neurasthenia. It is frequently cause for litigation. See *Erichsen's Diseases*.

Rainey's Corpuscles or Tubes. Protoplasts occurring in the muscles of the jaw and other animals.

Raisins (*ra'-sins*). Dried grapes.

Rale (*rah-l*) [*Fr., râler*, to rattle]. A ventitious sound heard over the chest during respiration and indicating some local disturbance. Rales are either *dry* or *moist* (produced by the bubbling of air through liquid). They are also classified according to their place of production, as *laryngeal*, *tracheal*, *bronchial*, *vesicular*, *cardiac*, *pleural*, *pericardial*. A table of rales is appended.

Ramification (*ram-e-fik-a'-shun*) [*ramus*, branch; *facere*, to make]. 1. The act of branching. 2. A branch.

Ramify (*ram-e-fi*) [*ramus*, a branch; *face*, to make]. To form branches; to branch.

Ramus (*ra'-mus*) [*L.*]. 1. A branch, especially of a vein, artery, or nerve. 2. A slender process of bone projecting like a branch twig from a large bone, as the *R.* of the lower jaw; ascending *R.* of the ischium. 3. A branch of the sympathetic ganglia connecting it with the sympathetic ganglia.

TABLE OF RALES.

VARIETY.	WHEN HEARD.	HOW AND WHERE PRODUCED.	CHARACTER.	CONDITION IN WHICH HEARD.
Amphoric.	Inspiration and expiration.	By movement of air in a tense-walled cavity containing air and communicating with a bronchus.	Large, musical, and tinkling.	In tuberculous and abscess-cavities.
Bubbling, large.	Inspiration and expiration.	By passage of air through frothy mucus in the trachea and larger bronchi.	Larger than the medium bubbling; moist.	Bronchitis and pulmonary engorgement.
Bubbling, medium.	Inspiration and expiration.	By passage of air through mucus in the larger tubes.	Larger than the small bubbling; moist.	In capillary bronchitis, especially in children.
Bubbling, small.	Inspiration and expiration.	By passage of air through mucus in the bronchioles.	Small, moist; like the bursting of small bubbles.	In capillary bronchitis, especially in children.
Cavernous.	Inspiration and expiration.	By passage of air through a small cavity with flaccid walls, that collapse with expiration.	Hollow and metallic.	In the third stage of pulmonary tuberculosis.
Clicking.	Inspiration only.	By passage of air through softening material in smaller bronchi.	Small, sticky.	In pulmonary tuberculosis, early stage.
Consonating.	Inspiration and expiration.	When bronchial tubes are surrounded by consolidated tissue.	Bright, clear, ringing.	Tuberculous pneumonia.
Crackling, dry.	In inspiration.	By the breaking down of lung-tissue.	Sharp, short, and clicking.	In the second or softening stage of pulmonary tuberculosis and in pulmonary gangrene.
Crackling, large.	Inspiration and expiration.	By fluid in very small cavities.	Larger than the medium crackling; dry.	In pulmonary tuberculosis and pneumonia, after formation of small cavities.
Crackling, medium.	Chiefly in inspiration.	By fluid in the finer bronchi.	Larger than the small crackling; dry.	Softening of tuberculous deposit or pneumonic exudation.
Crackling, small.	Chiefly in inspiration.	By fluid in the finer bronchi.	Small, dry; like the breaking of small shells.	Softening of tuberculous deposit or pneumonic exudation.
Crepitant.	End of inspiration.	By passage of air into vesicles collapsed, or containing fibrinous exudation. Usually at the base of the lungs.	Small; like rubbing the hair between the fingers.	Pneumonia, early stage; edema of lungs; hypostatic pneumonia; localized in pulmonary tuberculosis.
Dry.	Inspiration and expiration.	By narrowing of the bronchial tubes from thickening of the mucous lining, from spasmodic contraction of the muscular coat, viscid mucus within, or pressure from without.	Large and sonorous, or small and hissing, or whistling.	In bronchitis, asthma, and localized in beginning pulmonary tuberculosis.

TABLE OF RALES.—*Continued.*

VARIETY.	WHEN HEARD.	HOW AND WHERE PRODUCED.	CHARACTER.	CONDITION IN WHICH HEARD.
Extrathoracic.		In the trachea or larynx.		
Friction.	Inspiration and expiration; most distinct at the end of inspiration.	By the rubbing together of serous surfaces, roughened by inflammation or deprived of their natural secretion.	Grazing, rubbing, grating, creaking, or crackling.	In pleurisy and pericarditis.
Gurgling.	Inspiration and expiration.	By the passage of air through fluid in cavities, on coughing.	Larger than the large bubbling; moist; like the bursting of large bubbles.	Pulmonary tuberculosis after formation of cavities.
Cuttural.		In the throat.		
Moist.		By the passage of air through bronchi containing fluid.		
Mucous (of Laennec).	Inspiration and expiration.	By viscid bubbles bursting in the bronchial tubes.	A modification of the subcrepitant.	Pulmonary emphysema.
Rale redux, rale de retour.	Inspiration and expiration.	By the passage of air through fluid in a bronchial tube.	Crackling, unequal.	In pneumonia, in the stage of resolution.
Sibilant.	Inspiration and expiration.	By narrowing of the smaller bronchi from viscid mucus adhering to the walls, from thickening of the lining membrane or spasmodic contraction.	High-pitched and even hissing or piping.	In bronchitis, asthma, and localized in beginning pulmonary tuberculosis.
Sonorous.	Inspiration and expiration.	By lessened caliber of the larger bronchi, from spasm, tumefaction of mucous lining or external pressure.	Low-pitched and snoring.	Most frequent in bronchitis and spasmodic asthma.
Subcrepitant.	Inspiration and expiration.	By passage of air through mucus in the capillary bronchial tubes.	Small, moist.	Capillary bronchitis.

Rancid (*ran'-sid*) [*rancidus*, sour]. Having a rank or musty smell or taste; a term applied to fats and oils that have undergone decomposition with the development of volatile principles.

Rancidity (*ran-sid'-it-e*) [*rancidus*, rancid]. The state of being rancid.

Range (*ranj*) [Fr. *ranger*, to dispose]. Scope, extent. **R.** of Accommodation. See *Accommodation*.

Ranine (*ra'-nen*) [*rana*, a frog]. 1. Pertaining to a frog. 2. Pertaining to a ranula or to the region in which a ranula occurs, as R. artery.

Ranula (*ran'-u-lah*) [*rana*, frog]. A cystic tumor beneath the tongue, due to the occlusion of the duct of the sublingual or submax-

illary gland, or of a mucous gland of the floor of the mouth; frog-tongue.

Ranvier's Nodes (*ron(g)'-ve-az*) [*Ranvier*, a French histologist]. See *Node*.

Rape (*rap*) [*rapere*, to seize]. Sexual intercourse with a woman against her will or without her consent.

Raphania (*ra-fa'-ne-ah*) [*ραφανίς*, radish]. A nervous affection attended with spasmodic disorder of the joints and the limbs. It has been attributed to a poisonous principle in the seeds of the wild radish, which become mixed with grain. The affection is allied to ergotism and pellagra.

Raphe (*raf'-a*) [*ράφη*, a seam]. A seam or ridge, especially one indicating the line of junction of two symmetric halves.

Rarefaction (*rar-e-fak'-shun*) [*rarus*, rare; *facere*, to make]. The act of rarefying or of decreasing the density of a substance, especially the air. **R. of Bone**, the process of rendering bone more porous.

Rarefy (*ra'-re-fi*) [*rarus*, rare; *facere*, to make]. To make less dense or more porous.

Rarefying Osteitis. See *Osteoporosis*.

Rash [OF., *rasche*, from *radere*, to scrape]. A superficial eruption of the skin or mucous membrane. **R., Caterpillar-**, a localized eruption attributed to the irritant action of the hairs of certain caterpillars. **R., Drug-**, one produced by drugs. **R., Medicinal**. See *R., Drug-*. **R., Mulberry-**, an eruption resembling an exanthem of measles, sometimes occurring in typhus. **R., Nettle-**. See *Urticaria*. **R., Rose-**. See *Roseola*. **R., Scarlet**. See *Scarlet Fever*. **R., Tooth-**, any rash attributed to dentition.

Raspatory (*ras'-pa-to-re*) [*raspatorium*, from *radere*, to scrape]. A rasp or file for trimming the rough surfaces of bones or for removing the periosteum.

Raspberry (*raz'-ber-e*). The fruit of *Rubus idaeus*, a plant of the order Rosaceae. A syrup (*Syrupus rubi idaei*, U. S. P.) is used as a vehicle and as a drink in fevers.

Rasura (*ra-su'-rah*) [L.]. 1. The process of rasping, shaving, or scraping. 2. Scrapings; filings.

Ratanhia, Ratany (*rat-an'-e-ah, rat'-an-e*) [Peruv., *ratana*]. See *Krameria*.

Rathke's Pouch (*rat'-káz*) [M. H. Rathke, a German anatomist]. A diverticulum from the oral cavity into the cranial cavity, forming the anterior lobe of the pituitary body.

Rational (*ra'-shon-al*) [*ratio*, reason]. Based upon reason; reasonable. In therapeutics, opposed to empiric. **R. Formula**, in chemistry, a structural formula. See *Formula*. **R. Symptoms**, the symptoms elicited by questioning the patient, as opposed to those ascertained by physical examination.

Ratsbane (*rats'-ban*). 1. Arsenous trioxid. 2. A name given to any rat-poison containing arsenic.

Rattle (*rat'-l*) [AS., *hratelan*, to rattle]. A rale. **R., Death-**, a gurgling sound observed in dying persons, due to the passage of the air through mucus in the trachea.

Rau, Process of. The long process at the junction of the handle with the neck of the malleus.

Rauber's Layer. A superficial stratum of flat cells occurring in the center of the embryonal spot at an early stage in the development of the blastodermic membranes.

Rauschbrand (*raush'-brond*). The German name for black-leg.

Ray (*ra*) [*radius*, ray]. 1. A beam of light or heat; one of the component elements of light or heat. 2. One of a number of lines diverging from a common center. **R., Actinic**, a solar ray that produces chemic changes. **R., Cathode, R., Röntgen**. See *X-rays*. **R., Chemic**. See *R., Actinic*. **R.-fungus**. See *Actinomyces*. **R., Medullary, of the Kidney**, any one of the bundles of tubules that are the continuation into the cortex of the Malpighian pyramids.

Raynaud's Disease (*ra-nôz*). A trophoneurosis characterized by three grades of intensity: (a) Local syncope, observed most frequently in the extremities, and producing the condition known as dead fingers or dead toes. (b) Local asphyxia, which usually follows local syncope, but may develop independently. The fingers, toes, and ears are the parts usually affected. In the most extreme degree the parts are swollen, stiff, and livid, and the capillary circulation is almost stagnant. (c) Local or symmetric gangrene. Small areas of necrosis appear on the pads of the fingers and of the toes, also at the edges of the ears and tip of the nose. Occasionally symmetric patches are seen on the limbs or trunk, and in severe cases terminate in extensive gangrene. Some cases are attended by hemoglobinuria. The pathology of the disease is obscure.

Re-. A Latin prefix signifying back or again.

Reaction (*re-ak'-shun*) [*re*, again; *agere*, to act]. 1. Counteraction; opposite action; interaction. 2. The response of an organ or part to a stimulus. 3. In chemistry, (a) interaction of two or more substances when brought in contact; (b) the response to a certain test, as *Acid R.*, *Alkaline R.*, responding to the test for acid and alkali respectively. **R., Amphigenous, R., Amphoteric**. See *Amphoteric*. **R. of Degeneration**, the R. obtained when an electric stimulus is applied to a muscle deprived of its trophic nerve-influence. It is characterized by the following conditions: Diminution or abolition of the excitability of the muscles for the faradic current, with a temporary increase in excitability for the galvanic current. In the nerves there is a diminution or abolition of both faradic and galvanic excitability. The reactions of the muscle to the galvanic current (the true R. of degeneration) may be formulated as follows:—

An ClC < Ca ClC	} Muscle normal.
An OC > Ca OC	
An ClC = Ca ClC	
An OC = Ca OC	
An ClC > Ca ClC	
An OC < Ca OC	} Muscle in first stage of degeneration.
An ClC > Ca ClC	
An OC < Ca OC	} Muscle in more advanced stage of degeneration.
An ClC > Ca ClC	

AnClC = anodal closing contraction; AnOC = anodal opening contraction; CaClC = cathodal closing contraction; CaOC = cathodal opening contraction.

R., Electric, a response in a muscle or a part produced by electric stimulation. **R. of Exhaustion**, a variety of **R.** to electric excitation seen in states of exhaustion, in which a certain reaction produced by a given current-strength cannot be reproduced without an increase of current-strength. **R.**, Hemiopic Pupillary, of Wernicke, a reaction obtained in some cases of hemianopia in which a pencil of light thrown on the blind side of the retina gives rise to no movement in the iris, but thrown upon the normal side produces contraction of the iris. It indicates that the lesion producing the hemianopia is situated at or anterior to the geniculate bodies. **R.**, Myotonic, a **R.** seen in Thomsen's disease, in which there is quantitative increase in the faradic excitability. **R.**, Neutral, a **R.** indicating the absence of both acid and alkaline properties. **R.-period**, **R.-time**, the interval between the application of a stimulus and the beginning of the corresponding motor act.

Reagent (*re-aj-jent*) [*re*, again; *agere*, to act]. A substance used to produce a chemic reaction.

Real Focus. See *Focus*. **R. Image**. See *Image*.

Realgar (*re-al'-gar*) [Arab., dust of the mine; *rahj*, dust; *al*, the; *ghâr*, mine]. Arsenous disulphid, As_2S_3 .

Reamputation (*re-am-pu-ta'-shun*) [*re*, again; *amputation*]. An amputation upon a member on which the operation has been performed before.

Réaumur's Thermometer. See *Thermometer*.

Receiver (*re-se'-ver*) [*recipere*, to receive]. 1. The vessel receiving the products of distillation. 2. In an air-pump, the jar in which the vacuum is produced.

Receptaculum (*re-sep-tak'-u-lum*) [*recipere*, to receive]. A receptacle. **R. chyli**, the sac-like beginning of the thoracic duct opposite the last dorsal vertebra.

Recessus (*re-ses'-us*) [L.]. A depression or recess. **R. hemiellipticus**. See *Fovea hemielliptica*. **R. hemisphaericus**. See *Fovea hemisphaerica*. **R. pharyngeus**, a pouch-like process of the mucosa of the pharynx situated below the opening of the Eustachian tube.

Recipe (*res'-ip-e*). 1. The imperative of *recipere*, used as the heading of a physician's prescription and signifying take. Symbol **R.** 2. Also the formula itself.

Reciprocal Proportions, Law of. See *Law, Dalton's*.

Reclination (*rek-lin-a'-shun*) [*reclinare*, to recline]. An old operation for cataract, called also "couching," in which the lens was pushed back into the vitreous chamber.

Recrement (*rek'-re-ment*) [*recrementum*, from

re, again; *crescere*, to grow]. A secretion that is reabsorbed after fulfilling its function. **Recrementitious** (*rek-re-men-tish'-us*) [*recrement*]. Pertaining to or of the nature of a recrement.

Recrudescence (*re-kru-des'-ens*) [*re*, again; *crudescere*, to become raw]. An increase in the symptoms of a disease after a remission or a short intermission.

Rectal (*rek'-tal*). Pertaining to the rectum or performed through the rectum. **R. Alimentation**. See *Alimentation*. **R. Crises**, attacks of rectal pain and tenesmus occurring in locomotor ataxia.

Rectification (*rek-tif-ik-a'-shun*) [*rectus*, straight; *facere*, to make]. 1. A straightening, as **R.** of a crooked limb. 2. The redistillation of weak spirit in order to strengthen it.

Rectify (*rek'-tif-i*) [*rectus*, straight; *facere*, to make]. To make right or straight; to refine. **Rectified Spirit**, alcohol containing 84 per cent. of spirit.

Rectitis (*rek-ti'-tis*). See *Proctitis*.

Recto- (*rek'-to-*). A prefix meaning pertaining to the rectum.

Rectocele (*rek'-to-sel*) [*rectum*, rectum; *celia*, tumor]. Prolapse of the rectum into the vagina.

Rectotomy (*rek-tot'-o-me*) [*rectum*, rectum; *tomein*, to cut]. The operation of cutting into the rectum.

Rectourethral (*rek-to-u-re'-thral*) [*rectum*, rectum; *urethra*]. Pertaining to the rectum and the urethra.

Rectouterine (*rek-to-u'-ter-in*) [*rectum*, rectum; *uterus*]. Pertaining to the rectum and the uterus.

Rectovaginal (*rek-to-vaj'-in-al*) [*rectum*, rectum; *vagina*, vagina]. Pertaining to the rectum and the vagina. **R. Fistula**, an opening between the vagina and the rectum.

Rectovesical (*rek-to-ves'-ik-al*) [*rectum*, rectum; *vesica*, the bladder]. Pertaining to the rectum and the bladder.

Rectum (*rek'-tum*) [*rectus*, straight]. The lower part of the large intestine, extending from the sigmoid flexure to the anus. It begins opposite the left sacroiliac synchondrosis, passes obliquely downward to the middle of the sacrum, and thence descends in the median line to terminate at the anus.

Rectus (*rek'-tus*) [L.]. Straight; applied to anything having a straight course, as an artery (arteriæ rectæ of the kidney) or a muscle. **R. Muscle**. See *Muscles, Table of*.

Recurrans (*re-kur'-rens*) [*re*, back; *currere*, to run]. Relapsing fever.

Recurrent (*re-kur'-ent*) [*re*, back; *currere*, to run]. 1. Returning. 2. In anatomy, turning back in its course, as **R. laryngeal nerve**. **R. Fever**, relapsing fever.

Red [AS., *redd*, red]. The least refrangible

of the spectral colors; of a color resembling that of the blood. **R. Bark.** See *Cinchona*. **R.-blindness.** See *Blindness, Color-*. **R. Gum,** a red papular eruption of infants; also called strophulus. **R. Lead,** red lead oxid, formerly used in plasters. **R. Nucleus.** See *Nucleus*. **R. Pepper.** See *Capsicum*. **R. Precipitate.** See *Mercury*. **R. Softening,** a form of acute softening of the brain or spinal cord, characterized by a red punctiform appearance due to the presence of blood.

Reduce (*re-dūs'*) [*re*, back; *ducere*, to lead]. 1. To restore a part to its normal relations, as to **R.** a hernia or fracture. 2. In chemistry, to bring back to the metallic form; to deprive of oxygen.

Reduced (*re-dūs'd'*) [*re*, back; *ducere*, to lead]. 1. Restored to the proper place. 2. In chemistry, brought back into the metallic form as **R.** iron. 3. Diminished in size. **R. Eye.** See *Eye*.

Reducible (*re-dū'sibl'*) [*re*, back; *ducere*, to lead]. Capable of being reduced.

Reducin (*re-dū'sin*). A leukomatin, $C_{12}H_{16}N_4O_8$, found in urine.

Reduction (*re-dūk'-shun*) [*re*, back; *ducere*, to lead]. The act of reducing. **R. en masse,** the **R.** of a strangulated hernia still surrounded by its sac, thus failing to relieve the strangulation.

Reduplicated (*re-dū'plik-a-ted*) [*re*, again; *duplicare*, to double]. Doubled, as **R.** heart-sounds. See *Reduplication*.

Reduplication (*re-du'plik-a'-shun*) [*re*, again; *duplicare*, to double]. A doubling. **R. of the Heart-sounds,** a doubling of either the first or second sound of the heart.

Redux (*re'-duks*) [*L.*]. Returning. **R., Crepitus,** the small mucous rales heard in the early stage of the resolution of lobar pneumonia.

Reel, Cerebellar. The peculiar staggering gait in diseases of the cerebellum, particularly in tumor.

Refine (*re-fin'*) [*re*, again; *finire*, to finish]. To purify; to separate a substance from foreign matter.

Reflection (*re-flek'-shun*) [*re*, back; *flectere*, to bend]. A bending or turning back; specifically, the turning back of a ray of light from a surface upon which it impinges without penetrating.

Reflector (*re-flek'-tor*) [*re*, back; *flectere*, to bend]. A polished surface by which light is reflected.

Reflex (*re'-fleks*) [*re*, back; *flectere*, to bend]. 1. Anything reflected or thrown back. 2. A reflex act. **R. Act,** an act following immediately upon a stimulus without the intervention of the will. **R. Arc,** the mechanism necessary for a **R.** action; it consists of an afferent or sensory nerve; a nerve-center to change this sensory impulse into a motor one; and an efferent or motor nerve to carry a motor impulse to the muscle or group of muscles.

TABLE OF REFLEXES.

NAME.	HOW OBTAINED.	EFFECT PRODUCED.	SIGNIFICANCE.
Abdominal.	Sharp, sudden stroking of abdominal wall from margin of ribs downward.	Contraction of muscles about umbilicus.	Shows integrity of spinal cord from eighth to twelfth dorsal nerve.
Ankle-clonus.	By sudden complete flexion of foot, by pressing hand against sole.	Clonic contractions of tendo Achillis, dependent upon alternate contraction and relaxation of anterior tibial and calf-muscles.	
Biceps.	Tapping tendon of biceps.	Contraction of biceps muscle.	Normal; increased by causes increasing knee-jerk.
Bulbocavernous.	See <i>Virile</i> .		
Chin.	See <i>Jaw-jerk</i> .		
Ciliospinal.	By irritation of the skin of the neck.	Pupillary dilatation.	
Corneal.	Same as <i>Eyelid-closure</i> .		
Cremasteric.	Stimulation of skin on front and inner aspect of thigh.	Retraction of testicle on same side.	Shows integrity of cord between the first and second pairs of lumbar nerves.

TABLE OF REFLEXES.—*Continued.*

NAME.	HOW OBTAINED.	EFFECT PRODUCED.	SIGNIFICANCE.
Crossed.	Stimulation of one side of body.	Reflex on opposite side of body.	
Deep.	Reflexes developed by percussion of tendons or bones.		
Dorsal.	Same as <i>Erector spinæ</i> .		
Epigastric.	Stimulation of skin in fifth or sixth intercostal space near axilla.	Dimpling in the epigastrium, due to contraction of the highest fibers of the rectus abdominis muscle.	Shows integrity of cord from fourth to seventh dorsal nerves.
Erector spinæ.	Stimulation of skin along border of erector spinæ muscle.	Local contraction of these muscles.	Integrity of dorsal region of cord.
Eyelid-closure.	Irritation of conjunctiva.	Closure of lid.	
Femoral.	Irritation of skin on upper anterior aspect of thigh.	Plantar flexion of first three toes and of the foot, and extension of the knee-joint.	Disease of spinal cord, e.g., in some cases of transverse myelitis.
Front-tap.	See <i>Tendo Achillis</i> .		
Gluteal.	Firm sudden stroking of skin over buttock.	Contraction of glutei.	Shows integrity of cord at fourth and fifth lumbar nerves.
Interscapular.	See <i>Scapular</i> .		
Iris-contraction.	See <i>Pupillary</i> .		
Jaw-jerk, or jaw-clonus.	Downward stroke with a hammer on the lower jaw hanging passively or gently supported by the hand.	Clonic movements of inferior maxilla.	Rarely present in health; increased in sclerosis of lateral columns of cord.
Knee-jerk.	By striking patellar tendon after rendering it tense by flexing the knee at right angle.	Contraction of quadriceps muscle, the foot being jerked forward.	Normal in health. <i>Absent</i> in locomotor ataxia, destructive lesions of lower part of cord, multiple neuritis, affections of the anterior gray cornua, infantile paralysis, meningitis, diphtheric paralysis, atrophic palsy, pseudohypertrophic muscular paralysis, diabetes, etc. <i>Increased</i> in diseases of the pyramidal tracts, in spinal irritability, tumors of brain, cerebrospinal sclerosis, lateral sclerosis, after epileptic seizures or unilateral convulsions.
Laryngeal.	Irritation of fauces, larynx, etc.	Cough.	
Lumbar.	Same as <i>Erector spinæ</i> .		
Nasal.	Irritation of Schneiderian membrane.	Sneezing.	
Obliquus.	Irritation of skin below Poupart's ligament.	Contraction of fibers of external oblique in females; corresponds to cremasteric in males, although it can also be caused in males.	
Palatal.	Irritation of palate.	Swallowing.	
Palmar.	Tickling of palm.	Contraction of digital flexors.	Shows that cervical region of cord is normal.
Patellar.	Same as <i>Knee-jerk</i> .		
Patellar, Paradoxical.	Percussing patellar tendon with the patient in the dorsal decubitus.	Contraction of the adductor, but not of the quadriceps muscle. If the patient be in the sitting posture the normal reflex is elicited.	Spinal concussion.
Penis.	See <i>Virile</i> .		
Periosteal.	Tapping the bones of the forearm or leg.	Sharp contractions of the muscles.	Indicates disease of the lateral columns of spinal cord.

TABLE OF REFLEXES.—*Continued.*

NAME.	HOW OBTAINED.	EFFECT PRODUCED.	SIGNIFICANCE.
Peroneal.	Stroke on peroneal muscles when tense or when the foot is turned inward.	Reflex movements.	
Pharyngeal.	Irritation of pharynx.	Swallowing.	
Plantar.	Stroking sole of foot.	Contraction of toes.	
Platysma.	Pinching the platysma myoides muscle.	Dilatation of pupil.	
Pupillary.	Exposure of retina to light.	Contraction of iris.	Absent in basal meningitis, etc.
Pupillary, paradoxical.	Stimulation of retina by light.	Dilatation of pupil.	
Scapular.	Irritation of interscapular region.	Contraction of scapular muscles.	Shows integrity of cord between upper two or three dorsal and lower two or three cervical nerves.
Skin.	See <i>Platysma</i> .		
Sole.	Same as <i>Plantar</i> .		
Spinal.	Those reflex actions emanating from centers in the spinal cord.		
Superficial.	Such as are developed from irritation of the skin.		
Tendo Achillis, or front-tap contraction.	By striking muscles on anterior part of leg, while in extension, the foot being extended by the hand upon the sole.	Reflex contraction of gastrocnemius.	Considered as a delicate test of heightened spinal irritability.
Toe.	Strong flexion of great toe.	Involuntary flexion of foot, then flexion of leg, and, lastly, flexion of the thigh on the pelvis.	Met with in cases in which the knee-jerk and other tendon-reflexes are strongly developed.
Virile.	Sharp percussion of back of penis, the sheath having been made tense.	Retraction of bulbocavernous portion.	Occurs in health.
Wrist-clonus.	By pressing hand backward, causing extreme extension.	A series of jerking movements of the hand.	In the late rigidity of hemiplegia.

Refract (*re-frakt'*) [*re*, back; *frangere*, to break]. 1. To bend back. 2. To change direction by refraction. 3. To estimate the degree of ametropia present in an eye.

Refracta dosi (*re-frakt'-tah do'-si*) [L.]. In broken doses.

Refraction (*re-frak'-shun*) [*re*, back; *frangere*, to break]. 1. The act of refracting or bending back. 2. The deviation of a ray of light from a straight line in passing obliquely from one transparent medium to another of different density. 3. The state of refractive power, especially of the eye; the ametropia or emmetropia present. **R., Angle of**, the angle formed by a refracted ray of light with the perpendicular at the point of refraction.

R., Double, the power possessed by certain substances, as Iceland-spar, of dividing a ray of light and thus producing a double image of an object. **R., Dynamic**, the static refraction of the eye, plus that secured by the action of the accommodative apparatus. **R., Index of**, the refractive power of any substance as compared with air. It is the quotient of the angle of incidence divided by the angle of refraction of a ray passing through the substance. **R., Static**, that of the eye when accommodation is at rest.

Refractive (*re-frak'-tiv*) [*re*, back; *frangere*, to break]. Refracting; capable of refracting or bending back; pertaining to refraction.

Refractivity (*re-frak-tiv'-it-e*) [*re*, back;

frangere, to break]. Power of refraction; ability to refract.

Refractometer (*re-frak-tom'et-er*) [*re*, back; *frangere*, to break; *metron*, measure]. 1.

An instrument for measuring the refraction of the eye. 2. An instrument for the determination of the refractive indices of liquids.

Refractory (*re-frak'tor-e*) [*re*, back; *frangere*, to break]. 1. Resisting treatment. 2. Resisting the action of heat; slow to melt.

Refracture (*re-frak'tur*) [*re*, back; *frangere*, to break]. The breaking again of fractured bones that have joined by faulty or improper union.

Refrangibility (*re-fran-jib-il'-it-e*) [*re*, back; *frangere*, to break]. Capability of undergoing refraction.

Refresh (*re-fresh'*) [*re*, again; *friscus*, new]. In surgery, to give to an old lesion the character of a fresh wound.

Refrigerant (*re-frij'-er-ant*) [*refrigerans*, from *re*, again; *frigus*, cold]. 1. Cooling; lessening fever. 2. A medicine or agent having cooling properties, or lowering body-temperature.

Refrigeration (*re-frij'er-a'-shun*) [*re*, again; *frigus*, cold]. The act of lowering the temperature of a body, by conducting away its heat to a surrounding cooler substance.

Refusion (*re-fu'-shun*) [*refusio*, an overflowing]. The act of withdrawing blood from the vessels, exposing it to the oxygen of the air, and passing it back again.

Regeneration (*re-jen'er-a'-shun*) [*re*, again; *generare*, to beget]. The repair of lost or diseased structures.

Regimen (*rej'im-en*) [*regere*, to rule]. The regulated use of food, and the sanitary arrangement of surroundings to suit existing conditions of health or disease.

Region (*re'-jun*) [*regio*, a region]. One of the divisions of the body, possessing either natural or arbitrary boundaries. **Rs. of the Abdomen.** See *Abdomen*. **R., Epigastric.** See *Abdomen*. **R., Hypochondriac.** See *Abdomen*. **R., Hypogastric.** See *Abdomen*. **R., Iliac.** See *Abdomen*. **R., Infraaxillary,** the space between the anterior and posterior axillary lines. **R., Infraclavicular,** the space just below the clavicle. **R., Infra-mammary,** the space between a line drawn along the upper border of the xiphoid cartilage and the margin of the false ribs. **R., Infrascapular,** the region on either side of the vertebral column below a horizontal line drawn through the inferior angle of each scapula. **R., Infraspinous,** that included between the spine of the scapula and a line passing through the angle of the scapula. **R., Inguinal,** the region of the groin. **R., Interscapular,** the space between the scapulae. **R., Ischiorectal,** the region be-

tween the ischium and the rectum. **R., Lumbar.** See *Abdomen*. **R., Mammary,** the space on the anterior surface of the chest between the third and the sixth ribs. **R., Precordial,** the surface of the chest covering the heart. **R., Supraclavicular,** the space above the clavicle. **R., Supraspinous,** the region corresponding to the supraspinous fossa of the scapula. **R., Umbilical.** See *Abdomen*.

Regional (*re'-jun-al*) [*region*]. Pertaining to a region. **R. Anatomy,** the branch of anatomy that treats of the relations of the structures in a region of the body to each other and to the body-surface.

Regressive (*re-gres'-iv*) [*re*, back; *gradi*, to go]. Going back, returning, subsiding.

Regular (*reg'-u-lar*) [*regere*, to rule]. According to rule or custom. **R. School of Medicine,** the great mass of the profession, whose practice is based on the results of experience and experimental research without adherence to any exclusive theory of therapeutics. **R. Physician,** one belonging to the regular school.

Regurgitation (*re-gur-jit-a'-shun*) [*re*, again; *gurgitare*, to engulf]. 1. A back-flow of blood through a heart-valve that is defective. 2. The return of food from the stomach to the mouth soon after eating, without the ordinary efforts of vomiting.

Reichert's Cartilage. The cartilaginous structure that appears in the embryo on the hyoid arch, eventually becoming the styloid process, the stylohyoid ligament, and the lesser cornu of the hyoid bone.

Reichmann's Disease. A chronic disease of the stomach characterized by permanent gastric hypersecretion, associated with marked dilatation of the stomach, with thickening of its walls, and hypertrophy of the glands. It is accompanied by violent attacks of pain with vomiting, and may be followed by the formation of a round ulcer on the wall of the stomach.

Reid's Base Line. A line drawn from the infraorbital ridge through the middle of the external auditory meatus, and used for localizing the fissures of the brain on the skull.

Reil, Island of. See *Island of Reil*.

Reimplantation (*re-im-plan-tat'-shun*) [*re*, again; *plantare*, to plant]. In dental surgery, the replacing of a drawn tooth into its socket.

Reinfection (*re-in-fek't-shun*) [*re*, again; *inficere*, to infect]. Infection a second time with the same kind of virus.

Reinoculation (*re-in-ok-u-lat'-shun*) [*re*, again; *inoculare*, to inoculate]. Inoculation a second time with the same kind of virus.

Reinsch's Test. A test for arsenic. The suspected fluid is strongly acidulated with

hydrochloric acid and boiled, some slips of bright copper being added; a grayish coating on the copper may be shown to be arsenic by heating in a glass tube held obliquely, when, if it is arsenic, a crystalline coating will be sublimated on the glass above the copper.

Reinversion (*re-in-ver'-shun*) [*re*, again; *invertere*, to turn upside down]. The act of reducing an inverted uterus by the application of pressure to the fundus.

Reissner's Membrane. The membrane extending from the lamina spiralis to the outer wall of the cochlea, separating the scala media from the scala vestibuli.

Relapse (*re-laps'*) [*re*, again; *labi*, to fall]. A return of an attack of a disease shortly after the beginning of convalescence.

Relapsing Fever. An acute infectious disease due to the spirochæta Obermaieri. After a period of incubation of from five to seven days, the disease sets in with chill, fever, and pains in the back and limbs. The spleen enlarges, sweats and delirium occur, and the symptoms continue for five or six days, then suddenly cease by crisis. After a variable interval, usually in about a week, a second paroxysm occurs, which may be followed by a third and fourth. The disease prevails where conditions of overcrowding and defective food-supply obtain; hence the name, famine-fever.

Relaxant (*re-laks'-ant*) [*re*, again; *laxare*, to loosen]. 1. Loosening; causing relaxation. 2. An agent that diminishes tension.

Relaxation (*re-laks-a'-shun*) [*re*, again; *laxare*, to loosen]. A diminution of tension in a part; a diminution in functional activity, as, e. g., R. of the skin.

Remak's Fibers (*rem'-aks*) [*Remak*, a German physiologist]. See *Fiber*. R.'s Ganglion, a ganglion of nerve-cells in the sinus venosus of the frog's heart.

Remedial (*re-me'-de-al*) [*re*, again; *mederi*, to heal]. Having the nature of a remedy; relieving; curative.

Remedy (*rem'-e-d-e*) [*re*, again; *mederi*, to heal]. Anything used in the treatment of disease.

Remission (*re-mish'-un*) [*re*, back; *mittere*, to send]. 1. Abatement or subsidence of the symptoms of a disease. 2. The period of diminution of the symptoms of a disease.

Remittent (*re-mit'-ent*) [*re*, back; *mittere*, to send]. Characterized by remissions. R. Fever, a malarial fever characterized by periods of remission without complete apyrexia.

Ren [*I.*]. Kidney.

Renal (*re'-nal*) [*ren*, kidney]. Pertaining to the kidney. R. Calculus, a concretion in the kidney. R. Inadequacy, the condition in which the amount of urinary solids, and often the quantity of urine itself, is con-

siderably diminished. It is probably due to an exhausted condition of the epithelial cells of the kidney.

Reniform (*ren'-e-form*) [*ren*, kidney; *forma*, form]. Kidney-shaped.

Rennet (*ren'-et*) [*AS.*, *rinnan*, to run]. The prepared inner membrane of the fourth stomach of the calf, or an infusion of this membrane. It contains a milk-curdling ferment that decomposes casein.

Rennin (*ren'-in*) [*AS.*, *rinnan*, to run]. The milk-curdling ferment of the gastric juice; also called rennet.

Repellent (*re-pel'-ent*) [*re*, back; *pellere*, to push]. 1. Driving back. 2. Causing resolution of morbid processes.

Repercolation (*re-per-ko-la'-shun*) [*re*, again; *percolare*, to percolate]. Repeated percolation; the passage of a percolate for a second time, or oftener, through the percolator.

Repercussion (*re-per-kush'-un*) [*re*, again; *percutere*, to percuss]. 1. Ballottement. 2. A driving in or dispersion of a tumor or eruption.

Repletion (*re-ple'-shun*) [*re*, again; *plere*, to fill]. The condition of being full.

Reposition (*re-po-sish'-un*) [*re*, back; *ponere*, to place]. The act of returning into place, as R. of the uterus.

Repositor (*re-pos'-it-or*) [*re*, back; *ponere*, to place]. An instrument for replacing parts that have become displaced, especially for replacing a prolapsed umbilical cord; an instrument used in the replacement of a displaced uterus.

Reproduction (*re-pro-duk'-shun*) [*re*, again; *producere*, to produce]. The act of producing again; the procreation of one's kind; the producing of something like that lost.

Reproductive (*re-pro-duk'-tiv*) [*re*, again; *producere*, to produce]. Pertaining to reproduction, as the R. organs.

Repulsion (*re-pul'-shun*) [*re*, back; *pellere*, to drive, to push]. 1. The act of repelling or driving back or apart. 2. The influence tending to drive two bodies apart; the opposite of attraction.

Resection (*re-sek'-shun*) [*re*, again; *secare*, to cut]. The operation of cutting out. R. of a Joint, the cutting away of the ends of the bones forming a joint, or a portion of bone, nerve, or other structure.

Reserve Air. See *Respiration*.

Reservoir of Pecquet (*pek'-a'*). See *Receptaculum chyli*.

Residual Air (*re-sid'-u-al*). See *Respiration*.

Residue (*res'-id-u*) [*re*, again; *sedere*, to sit]. That remaining after a part has been removed.

Resilience (*re-sil'-e-ens*) [*re*, back; *salire*, to leap]. The quality of being elastic or resilient.

Resilient (*re-sil'-e-ent*) [*re*, back; *salire*, to leap]. Rebounding; elastic. **R. Stricture**, one that contracts again immediately after being dilated.

Resin (*res'-in*) [*resina*]. One of a class of vegetable substances exuding from various plants, and characterized by being soluble in alcohol, in ether, and in the volatile oils, and insoluble in water; they are readily fusible, and inflammable. They are obtained in pharmacy by treating the substances containing them with alcohol, and then precipitating the alcoholic solution with water. The chief resins are Resina (U. S. P., B. P.), from turpentine; Resina copaibæ (U. S. P.), dose gr. x-xx (0.65-1.3); Resina jalapæ (U. S. P., B. P.), dose gr. ij-v (0.13-0.32); Resina podophylli, dose gr. $\frac{1}{8}$ - $\frac{1}{2}$ (0.008-0.032); Resina scammonii (U. S. P., B. P.), dose gr. iv-viii (0.26-0.52). **R. - plaster**, emplastrum resinæ; adhesive plaster.

Resinous (*res'-in-us*) [*resina*, a resin]. Having the nature of a resin.

Resistance (*re-sis'-tans*) [*resistere*, to withstand]. In electricity, the opposition offered by a conductor to the passage of the current. **R. - coil**, a coil of wire for increasing the resistance in a circuit. **R., Essential**, or **R., Internal**, the resistance to conduction within the battery itself. **R., Extraordinary**, or **R., External**, the resistance to conduction outside of the battery.

Resolution (*rez-a-lu'-shun*) [*resolvere*, to resolve]. The return of a part to the normal state after a pathologic process.

Resolve (*re-solv'*) [*resolvere*, to dissolve]. 1. To return to the normal state after some pathologic process. 2. To separate anything into its component parts.

Resolvent (*re-solv'-vent*) [*resolvere*, to dissolve]. 1. Causing solution or dissipation of tissue. 2. An agent causing resolution.

Resolving Power. The capability of a lens of making clear the finest details of an object.

Resonance (*rez'-o-nans*) [*re*, again; *sonare*, to sound]. 1. The sound obtained on striking a hollow object, especially the note obtained on percussing the chest or abdomen. 2. The sound of the voice as transmitted to the ear applied to the chest. **R., Cracked-pot**, a sound elicited by percussing over a pulmonary cavity communicating with a bronchus. **R., Skodaic**, the increased percussion-resonance over the upper part of a lung when the lower part is compressed by a pleural effusion. **R., Tympanitic**, a hollow sound elicited on percussion over the intestines and over large pulmonary cavities with thin yielding walls. **R., Vesicular**, the normal pulmonary R. **R., Vesiculotympanitic**, an admixture of vesicular and tympanitic R. **R., Vocal**, the sound heard on auscultation of the chest

during ordinary speech. **R., Whispering**, the sound heard on auscultation of the chest during the act of whispering.

Resonator (*rez'-o-na-tor*) [*re*, again; *sonare*, to sound]. An instrument used to intensify sounds.

Resorcin (*re-sor'-sin*) [*resina*, resin; *Orcus*, Pluto], $C_6H_4(OH)_2$. A crystalline substance isomeric with pyrocatechin and hydroquinone, and usually prepared by fusing sodium benzene disulphonate with sodium hydroxid. It is an antipyretic and antiseptic, but is chiefly used in ointments for chronic skin-diseases. Dose gr. ij-iv (0.13-0.26).

Resorption (*re-sorp'-shun*) [*re*, again; *sorbere*, to absorb]. The absorption of morbid deposits, as of the products of inflammation.

Respirable (*res'-pir-a-bl*) [*re*, again; *spirare*, to breathe]. Capable of being inspired and expired; capable of furnishing the gaseous interchange in the lung necessary for life.

Respiration (*res-pir-a'-shun*) [*re*, again; *spirare*, to breathe]. 1. The interchange

between the gases of living organisms and the gases of the medium in which they live, through any channel, as cutaneous R. 2. The act of breathing with the lungs; the taking into and the expelling from the lungs of air. It consists of two acts—inspiration, or the taking in of the atmospheric air, and expiration, the expelling of the modified air. Expired air contains less oxygen and more carbon dioxide than inspired air. The volume of air taken into the lungs and given out during an ordinary R. (*tidal air*) is 500 c.c.; the volume that can be inspired in addition by a forcible inspiration (*complemental air*) is 1500 c.c.; that which remains in the chest after a normal expiration (*reserve* or *supplemental air*) is 1500 c.c.; the amount remaining in the chest after the most complete expiration (*residual air*) is from 1200-1600 c.c. The volume of air that can be forcibly expelled after the most forcible inspiration is termed *total* or *respiratory capacity* and is equal to the tidal air, complemental air, and reserve air, or about 3500 c.c. See *Breath* and *Breathing*. **R., Abdominal**, a type of R. caused by the contraction of the diaphragm and the elasticity of the abdominal walls and viscera. It is more common in men than in women. **R., Artificial**, the artificial production of the normal respiratory movements. See *Artificial Respiration*. **R., Costal**, a type of R. in which the chest movement predominates over the diaphragmatic movement, seen especially in women. **R., Metamorphosing**, a form of R. in which inspiration is at first harsh, but toward the end becomes blowing and accompanied by adventitious sounds. It is produced by the entrance of air into a cavity through a narrow opening.

Respirator (*res-pir-a-tor*) [*re*, again; *spirare*, to breathe]. An appliance by which the inspired air, in passing through it, is warmed, purified, or medicated.

Respiratory (*res'-pir-a-to-re*) [*re*, again; *spirare*, to breathe]. Pertaining to respiration. **R. Bundle**, the ascending root of the glossopharyngeal nerve, probably arising in the posterior horns of the cord. **R. Center**. See *Center*. **R. Murmur**, the sound produced by the air entering and escaping from the lungs during respiration. **R. Nerve**, one of two nerves supplying important muscles of respiration. The external is the posterior thoracic nerve; the internal, the phrenic nerve. **R. Quotient**, the ratio of the amount of oxygen taken in by the lungs to the carbon dioxide given off in the same period of time. **R. Tract**, all the air-passages and air-cells concerned in respiration.

Restiform (*res'-te-form*) [*restis*, a rope; *forma*, form]. Corded or cord-like. **R. Body**, a part of the medulla oblongata, which as the inferior cerebellar peduncle connects the medulla with the cerebellum. It contains fibers from the lateral column of the spinal cord (the lateral cerebellar tract), from the posterior column, and from the inferior olivary nucleus.

Restitution (*res-tit-u'-shun*) [*re*, again; *statuere*, to set up]. The act of restoring.

Resuscitation (*re-sus-it-a'-shun*) [*re*, again; *suscitare*, to raise up]. The bringing back to life of one apparently dead.

Retch [AS., *hræc*, a cough]. To strain at vomiting.

Re-te (*re'-te*) [L.]. A net or net-like structure. **R. Malpighii**. See *R. mucosum*. **R. mirabile**, a cluster of fine vessels produced by the splitting of an artery into numerous branches. If the branches do not reunite, it is called unipolar *R. mirabile*; if they reunite, bipolar *R. mirabile*. **R. mirabile duplex**, a *R. mirabile* consisting both of veins and arteries. **R. mucosum**, the deeper layers of the epidermis. **R. testis**, the network of seminal tubules in the corpus Highmori of the testicle.

Retention (*re-ten'-shun*) [*re*, back; *tenere*, to hold]. The act of retaining or holding back. **R.-cyst**. See *Cyst*. **R. of Urine**, the holding of the urine in the bladder on account of some hindrance to urination.

Reticular (*ret-ik'-u-lar*) [*reticulum*, dim. of *rete*, net]. Resembling a net; formed by a network. **R. Formation**. See *Formatio*.

R. Lamina, the membrane covering the organ of Corti. **R. Layer of the Skin**, the deep layer of the skin, consisting of interlacing bands of white and yellow fibrous tissue.

Reticulated (*ret-ik'-u-la-ted*). See *Reticular*.

Reticulum (*re-tik'-u-lum*) [*reticulum*, dim. of *rete*, net]. A network.

Retina (*ret'-in-ah*) [*rete*, a net]. The delicate membrane of the eye representing the terminal expansion of the optic nerve, and extending from the point of entrance of the nerve forward to its termination in the ora serrata. It consists of the following layers, named from behind forward: (*a*) the pigment-layer; (*b*) the neuroepithelial layer, comprising the layer of rods and cones (Jacob's membrane, bacillary layer), the outer limiting membrane, and the outer nuclear layer; (*c*) the cerebral layer, comprising the outer reticular layer (outer granular layer), the inner nuclear layer, the inner reticular layer (inner granular layer), the ganglion-cell layer, the nerve-fiber layer. These layers are cemented together by a supporting framework of connective tissue, the fibers of Müller, or radiating fibers.

Retinaculum (*ret-in-ak'-u-lum*) [*re*, back; *tenere*, to hold]. A band or membrane holding back an organ or part. **R. ligamenti arcuati**, the short external lateral ligament of the knee-joint. **R. Morgagni**, **R. of the Ileocecal Valve**, the ridge formed by the coming together of the valve-segments at each end of the opening between the cecum and the ileum. **R. peroneorum inferius**, a fibrous band running over the peroneal tendons as they pass through the grooves on the outer side of the calcaneum. **R. peroneorum superius**, the external annular ligament of the ankle-joint. **R. tendinum**, the annular ligament of the wrist or ankle.

Retinal (*ret'-in-al*) [*retina*]. Pertaining to or affecting the retina.

Retinitis (*ret-in-ī'-tis*) [*retina*, retina; *itis*, inflammation]. Inflammation of the retina. **R., Albuminuric**, **R. due to nephritis**. **R., Diabetic**, **R. occurring in diabetes**. **R., Hemorrhagic**, **R. associated with hemorrhages**. **R., Leukemic**, a form occurring in leukemia and characterized by pallor of the retinal vessels and optic disc and hemorrhages.

Retinochoroiditis (*ret-in-o-ko-roi-di'-tis*) [*retina*, retina; *χόροιν*, the chorion; *ιδος*, like; *itis*, inflammation]. Inflammation of the retina and choroid.

Retinol (*ret'-in-ol*) [*resin*, a resin], $C_{27}H_{46}$. A liquid hydrocarbon obtained in the destructive distillation of resin. It is used as a solvent and has also been employed in gonorrhea.

Retinoscopy (*ret-in-ō'-ko-pe*) [*retina*, retina; *σκοπεῖν*, to observe]. A method of determining the refraction of the eye by observation of the movements of the retinal images and shadows through the ophthalmoscopic mirror; skiascopy.

Retort (*re-tort'*) [*re*, back; *torquere*, to twist]. A vessel employed in distillation, consisting of an expanded globular portion and a long neck, and containing the liquid to be distilled.

Retractile (*re-trak'-til'*) [*re*, back; *trahere*, to draw]. Capable of being drawn back.

Retractility (*re-trak'-til'-it-ē*) [*re*, back; *trahere*, to draw]. The power of retracting or drawing back.

Retraction (*re-trak'-shun*) [*re*, back; *trahere*, to draw]. The act of retracting or drawing back; a R. of the muscles after amputation.

Retractor (*re-trak'-tor*) [*retrahere*, to draw back]. An instrument for drawing back the lips of a wound so as to give a better view of the deeper parts.

Retrahens aurem (*re-trah'-hens aw-rem*). Drawing back the ear. See *Muscles*, Table of.

Retro- (*re'-tro-*) [*retro*, back]. A prefix meaning backward or behind.

Retrobulbar (*re-tro-bul'-bar*) [*retro*, back; *bulbus*, bulb]. Situated or occurring behind the eyeball. R. Neuritis, inflammation in the orbital part of the optic nerve.

Retrocedent (*re-tro-sed'-dent*) [*retro*, back; *cedere*, to go]. Going back; disappearing from the surface. R. Gout, a form of gout in which the joint inflammation suddenly disappears and is replaced by affections of the internal organs.

Retrocession (*re-tro-sesh'-un*) [*retro*, back; *cedere*, to go]. The act of going back.

Retroclulsion (*re-tro-klu'-shun*) [*retro*, back; *claudere*, to shut]. A form of acupressure in which the pin is passed first above the artery into the tissues on the other side, then below the artery into the tissues upon the side first entered.

Retrocollic (*re-tro-koll'-ik*) [*retro*, back; *collic*, the neck]. Pertaining to the muscles at the back of the neck. R. Spasm, spasm of the muscles at the back of the neck, causing retraction of the head.

Retroflexion (*re-tro-flek'-shun*) [*retro*, back; *flectere*, to bend]. The state of being bent backward. R. of the Uterus, a condition in which the uterus is bent backward upon itself, producing a sharp angle in its axis.

Retrograde (*ret'-ro-grād*, or *re'-tro-grād*) [*retro*, backward; *gradī*, to go]. Going backward; undoing. R. Embolism, embolism in which the embolus has gone against the normal direction of the blood-stream.

Retrography (*re-trog'-ra-fē*) [*retro*, back; *γράφειν*, to write]. Backward writing; mirror-writing.

Retroinsular (*re-tro-in'-su-lar*) [*retro*, behind; *insula*, island]. Situated behind the island of Reil, as the R. convolutions.

Retronasal (*re-tro-nal'-sal*) [*retro*, behind; *nusus*, nose]. Situated behind the nose or nasal cavities.

Retroocular (*re-tro-ok'-u-lar*) [*retro*, behind; *oculus*, the eye]. See *Retrobulbar*.

Retroperitoneal (*re-tro-per-i-ton-e'-al*) [*retro*, behind; *περίτονιον*, peritoneum]. Situated behind the peritoneum.

Retropharyngeal (*re-tro-far-in'-je-al*) [*retro*, behind; *φάρυγξ*, the pharynx]. Situated behind the pharynx, as R. abscess.

Retropulsion (*re-tro-pul'-shun*) [*retro*, back; *pellere*, to drive]. 1. A driving or turning back, as of the fetal head. 2. A running backward; a form of walking sometimes seen in paralysis agitans.

Retrosternal (*re-tro-ster'-nal*) [*retro*, back; *sternum*]. Situated behind the sternum.

Retrotarsal (*re-tro-tar'-sal*) [*retro*, back; *ταρσός*, tarsus]. Situated behind the tarsus, as, e. g., the R. fold of the conjunctiva.

Retrouterine (*re-tro-ū'-ter-in*) [*retro*, behind; *uterus*, uterus]. Behind the uterus. R. Hematocoele, a blood-tumor behind the uterus in the pouch of Douglas.

Retrovaccination (*re-tro-vak'-sin-a'-shun*) [*retro*, back; *vaccination*]. Vaccination with virus from a cow that had been inoculated with the virus of smallpox from a human subject.

Retroversion (*re-tro-ver'-shun*) [*retro*, backward; *vertere*, a turning]. A turning back. R. of Uterus, a condition in which the uterus is tilted backward without curvature of its axis.

Retroverted (*re-tro-ver'-ted*) [*retro*, backward; *vertere*, to turn]. Tilted or turned backward, as a R. uterus.

Reuss's Test. A test for atropin, consisting in the development of an odor of blossoms when the substance is treated with sulphuric acid and an oxidizing agent.

Revulsant (*re-vul'-sant*) [*re*, back; *vellere*, to pluck]. 1. Causing revulsion. 2. See *Revulsive*.

Revulsion (*re-vul'-shun*) [*re*, back; *vellere*, to pluck]. A plucking or driving backward; specifically, the diverting of disease from one part to another by the sudden withdrawal of the blood from the part.

Revulsive (*re-vul'-siv*) [*re*, back; *vellere*, to pluck]. 1. Causing revulsion. 2. An agent that causes revulsion.

Rhabdomyoma (*rah-do-mi-o'-mah*) [*ράβδος*, a rod; *μῦς*, a muscle; *δύα*, a tumor]. A form of myoma characterized by the presence of striated muscular fibers.

Rhachi-. See *Rachi-*.

Rhagades (*rag'-ad-ē*) [*ράγας*, fissure]. Linear cracks or fissures, especially in the skin.

Rhamnus (*ram'-nus*) [*ῥάμνος*, buckthorn].

The *Rhamnus purshiana*, a small tree of the order Rhamnaceæ. Its bark (*R. purshiana*, U. S. P., *Rhamni purshiani cortex*, B. P.) is commonly known as Cascara sagrada or chittam-bark, and is used as a laxative in habitual constipation. *Extractum rhamni purshianæ fluidum* (U. S. P.), *Extractum cascariæ sagradæ liquidum* (B. P.). Dose $\text{m}\frac{\text{xv}}{\text{xlv}}$ (1.0–3.0).

Rhaphania (*raf'-a'-ne-ah*). See *Raphania*.

Rhaphe (*ra'-fe*). See *Raphe*.

Rhatany (*rat'-an-e*). See *Krameria*.

Rheo- (*re'-o*) [*ῥέος*, current]. A prefix denoting pertaining to a current.

Rheocord (*re'-o-kord*) [*ῥέος*, current; *χορδή*, a cord]. See *Rheostat*.

Rheometer (*re-om'-et-er*) [*ῥέος*, current; *μέτρον*, a measure]. 1. A galvanometer. 2. An apparatus for measuring the velocity of the blood-current.

Rheophore (*re'-o-för*) [*ῥέος*, current; *φέρειν*, to carry]. An electrode.

Rheoscope (*re'-o-sköp*) [*ῥέος*, current; *σκοπεῖν*, to see]. An instrument for demonstrating the existence of an electric current; a galvanoscope.

Rheostat (*re'-o-stat*) [*ῥέος*, current; *στάναι*, to stand]. An instrument introduced into an electric current and offering a known resistance, for the purpose of regulating the strength of the current.

Rheotome (*re'-o-töm*) [*ῥέος*, current; *τέμνειν*, to cut]. An instrument for breaking and making a galvanic circuit; an interrupter.

Rheotrope (*re'-o-tröp*) [*ῥέος*, current; *τρέπεν*, to turn]. An apparatus for reversing the direction of an electric current.

Rheum (*re'-um*). See *Rhubarb*.

Rheum (*rüm*) [*ῥεύμα*, from *ῥέειν*, to flow]. Any watery or catarrhal discharge. **R.**, Salt-, eczema.

Rheumatic (*ru-mat'-ik*) [*ῥεύμα*, flux]. Pertaining to, of the nature of, or affected with rheumatism. **R. Diathesis**, the condition of body tending to the development of rheumatism. **R. Fever**, acute articular rheumatism. **R. Gout**. Synonym of *Rheumatoid Arthritis*.

Rheumatism (*ru-ma-tizm*) [*ῥεύμα*, flux]. A constitutional disease characterized by pain in the joints and muscles, tending to recur, and associated with exposure to cold and wet. **R.**, **Acute Articular**, is characterized by fever, by swelling of various joints, beginning usually in one and rapidly spreading to others, by acid sweats, and by a marked tendency to involve the endocardium, less frequently the pericardium, pleura, and peritoneum. The iris and conjunctiva may also become affected. **R.**, **Chronic**, a chronic form in which the symptoms are milder and in which the disease attacks either the muscles (muscular **R.**) or

the joints. **R.**, **Gonorrheal**, joint-inflammation occurring in association with gonorrheal urethritis. It generally involves but one joint; if several are affected, it is usually the smaller joints. The course is chronic.

Rheumatismal (*ru-ma-tiz'-mal*) [*ῥεύμα*, flux]. Pertaining to rheumatism.

Rheumatoid (*ru'-ma-toid*) [*ῥεύμα*, flux; *εἶδος*, resemblance]. Resembling rheumatism. **R. Arthritis**, a disease of the joints characterized by chronic inflammatory and degenerative changes involving the structure of the various articulations and resulting in rigidity and deformity. The mildest type is known as Heberden's nodosities, in which small nodules appear at the sides of the distal phalanges of the fingers; in severer forms nearly every articulation may become ankylosed.

Rheumic (*ru'-mik*) [*ῥεύμα*, flux]. Pertaining to rheum. **R. Diathesis**, one that gives rise to cutaneous eruptions.

Rhigolene (*rig'-o-len*) [*ῥίγος*, cold]. A very volatile liquid obtained from petroleum by distillation, and used as a local anesthetic. Its rapid evaporation freezes and benumbs the part upon which it is sprayed.

Rhin- (*rin-*), **Rhino-** (*ri'-no-*) [*ῥίς*, nose]. A prefix signifying pertaining to the nose.

Rhinencephalon (*ri-nen-sef'-al-on*) [*ῥίς*, nose; *ἐγκέφαλος*, brain]. The olfactory lobe of the brain.

Rhineurynter (*ri-nu-rin'-ter*) [*ῥίς*, nose; *εὐρίπτειν*, to dilate]. A distensible bag or sac which is inflated after insertion into the nostril.

Rhinion (*rin'-e-on*) [*ῥίς*, nose]. The lower point of the suture between the nasal bones. See *Cranimetric Points*.

Rhinitis (*ri-ni'-tis*) [*ῥίς*, the nose; *ιτις*, inflammation]. Inflammation of the nasal mucous membrane. **R.**, **Acute**, coryza; cold in the head. **R.**, **Chronic**, a form usually due to repeated attacks of acute **R.**, and producing in the early stages hypertrophy of the mucous membrane (hypertrophic **R.**) and in the later stages atrophy (atrophic **R.**), and the presence of dark, offensively-smelling crusts. **R.**, **Fibrinous**, a rare form of rhinitis characterized by the development of a false membrane in the nose. **R.**, **Pseudomembranous**. See *R.*, *Fibrinous*. **R.**, **Syphilitic**, a chronic form due to syphilis, and usually attended by ulceration and caries of the bone and an offensive discharge (ozena). **R.**, **Tuberculous**, that due to the tubercle-bacillus; it is usually associated with ulceration and caries of the bones.

Rhinocephalus (*ri-no-sef'-al-us*) [*ῥίς*, nose; *κεφαλή*, head]. A monster in which the nose resembles a tube and the eyes are fused below the nose.

Rhinolalia (*ri-no-la'-le-ah*) [*ῥίς*, the nose; *λαλία*, speech]. Imperfect articulation due to undue closure (*R. clausa*), or undue patulousness (*R. aperta*), of the posterior nares.

Rhinologic (*ri-no-loj'-ik*) [*ῥίς*, nose; *λόγος*, science]. Pertaining to rhinology.

Rhinologist (*ri-nol'-o-jist*) [*ῥίς*, nose; *λόγος*, science]. A specialist in the treatment of diseases of the nose.

Rhinology (*ri-nol'-o-je*) [*ῥίς*, nose; *λόγος*, science]. The science of the anatomy, functions, and diseases of the nose.

Rhinometer (*ri-nom'-et-er*) [*ῥίς*, nose; *μέτρον*, measure]. An instrument for measuring the nose.

Rhinophyma (*ri-no-fī'-mah*) [*ῥίς*, nose; *φύμα*, tumor]. A form of acne rosacea of the nose characterized by a marked hypertrophy of the blood-vessels and the connective tissue, producing a lobulated appearance of the nose.

Rhinoplasty (*ri'-no-plas-te*) [*ῥίς*, nose; *πλάσσειν*, to mold]. A plastic operation upon the nose. **R.**, **Indian**. See *Operation, Indian*, in *Operations, Table of*. **R.**, **Tagliacotian**. See *Operation, Tagliacotian*, in *Operations, Table of*.

Rhinopolypus (*ri-no-pol'-ip-us*) [*ῥίς*, nose; *πολύπος*]. Polypus of the nose.

Rhinorrhagia (*ri-nor'-a'-je-ah*) [*ῥίς*, nose; *ῤῥγνῖναι*, to burst forth]. Hemorrhage from the nose.

Rhinoscleroma (*ri-no-skle-ro'-mah*) [*ῥίς*, nose; *σκληρός*, hard; *ῥμα*, tumor]. A new growth of almost stony hardness, affecting the anterior nares and adjacent parts. The disease commences in the mucous membrane of the anterior nares and adjoining skin, the lesions consisting of flat, isolated, or coalescent nodules. The cause is thought to be a bacillus.

Rhinoscope (*ri'-no-skōp*) [*ῥίς*, nose; *σκοπεῖν*, to observe]. An instrument for examination of the cavities of the nose.

Rhinoscopy (*ri-nos'-kō-pe*) [*ῥίς*, nose; *σκοπεῖν*, to observe]. Examination of the nasal fossæ by means of the rhinoscope; that of the anterior nares is termed anterior *R.*; that of the posterior nares, posterior *R.*

Rhizome (*ri'-zōm*) [*ῥίζα*, root]. A subterranean stem, having roots at its nodes and a bud or shoot at its apex.

Rhodophan (*ro'-dō-fan*) [*ῥόδον*, rose; *φανής*, appearing]. A red pigment found in the retinal cones.

Rhodopsin (*ro-dop'-sin*) [*ῥόδον*, rose; *ὤψ*, eye]. The visual purple contained in the retinal rods.

Rhœados Petala (*re'-a-dos pet'-a-lah*), **Rhœas** (*re'-as*). See under *Poppy*.

Rhomboid (*rom'-boid*) [*ῥόμβος*, a rhomb; *εἶδος*, resemblance]. Having a shape simi-

lar to that of a rhomb, a quadrilateral figure with opposite sides equal and parallel, and oblique angles. **R. Fossa**, the fourth ventricle of the brain. **R. Ligament**. See *Ligaments, Table of*.

Rhonical (*rong'-ke-al*) [*ῥογχός*, a snoring]. Relating to or produced by a rhonchus, as *R. fremitus*.

Rhonchus (*rong'-kus*) [*ῥογχός*, snore]. A rattling sound produced in the throat or bronchial tubes during respiration. See *Rule*.

Rhotacism (*ro'-tas-izm*) [*ῥο*, the letter *r*]. The use of the *r* sound in place of other speech-sounds; the too-strong utterance of the letter *r*.

Rhubarb (*ru'-barb*) [*L., Rheum*]. A genus of plants of the order Polygonaceæ. The bark of *Rheum officinale*, or *Rheum palmatum* (*Rheum*, *U. S. P.*, *Rhei radix*, *B. P.*), contains chrysophanic acid, tannic acid (rheotannic acid), and several coloring principles, and is used as a laxative, stomachic, and astringent. Its chief uses are in dyspepsia with constipation, in the diarrhea of children, and in the beginning of bilious fevers. Dose gr. v-xxx (0.32-2.0). Preparations and doses: Extractum rhei (*U. S. P.*, *B. P.*), gr. x-xv (0.65-1.0). Extractum rhei fluidum (*U. S. P.*), mxx-xxx (0.65-2.0). Infusum rhei (*B. P.*), f 3j-ij (32.0-64.0). Mistura rhei et sodæ (*U. S. P.*), f 3ss-j (2.0-4.0). Pilulæ rhei (*U. S. P.*), each of which contains gr. ij (0.20) of rhubarb. Pilulæ rhei compositæ (*U. S. P.*), Pilulæ rhei composita (*B. P.*), contain rhubarb and aloes. Dose 2-4 pills. Pulvis rhei compositus (*U. S. P.*, *B. P.*), Gregory's powder, 3ss-j (2.0-4.0). Syrupus rhei (*U. S. P.*, *B. P.*), f 3j (4.0). Syrupus rhei aromaticus (*U. S. P.*), f 3j (4.0). Both of the foregoing are used chiefly for children, in the doses given. Tinctura rhei (*U. S. P.*, *B. P.*), f 3j-ij (4.0-8.0). Tinctura rhei aromatica (*U. S. P.*), f 3ss-j (2.0-4.0). Tinctura rhei dulcis (*U. S. P.*), f 3j-ij (4.0-12.0). Vinum rhei (*B. P.*), f 3j-iv (4.0-16.0).

Rhus (*rus*) [*ῥοῖς*, sumac, gen., *rhôis*]. A genus of shrubs or small trees of the order Anacardiæ. The berries of *R. glabra*, or sumach (*R. glabra*, *U. S. P.*), are used as an astringent in inflammations of the mouth and throat, in the form of a decoction or the official Extractum rhois glabræ fluidum. **R. toxicodendron**, the poison-ivy, is a powerful irritant and produces in susceptible persons a violent dermatitis with vesicles and intense itching (ivy-poisoning). The active agent seems to be an acid called toxicodendric acid. In overdoses taken internally it acts as a narcotic poison. It has been employed in chronic rheumatism and in incontinence of

urine. **R. venenata**, swamp-sumach, is also poisonous.

Rhytidosis (*rit-id-o'-sis*) [*ῥυτίς*, a wrinkle]. A wrinkling.

Rib. One of the 24 long, flat, curved bones forming the wall of the thorax. **R.**, **False**, one of the five lower ribs not attached to the sternum directly. **R.**, **Floating**, one of the last two ribs which have one end free. **R.**, **True**, one of the seven upper ribs that are attached to the sternum.

Rice (*ris*). The *Oryza sativa*, a plant of the Gramineæ; also its seed. **R.** is used as a food, as a demulcent, and, in the form of rice-water, as a drink in fevers.

Rice-water Evacuations. The name given to the bowel-discharges in cholera.

Richter's Hernia. See *Diseases, Table of*. **Ricin** (*ris'-in*) [*ricinus*, castor-oil]. A poisonous proteid found in the castor-oil bean.

Ricinin (*ris'-in-in*) [*ricinus*, castor-oil]. A poisonous crystalline substance obtained from castor-oil.

Ricinus (*ris'-in-us*) [*L.*, a tick, from the resemblance of the seed to that insect]. The *R. communis*, or castor-oil plant, a plant or tree of the order Euphorbiaceæ. The fixed oil expressed from its seeds is the Oleum ricini or castor-oil of the pharmacopeia, and is used as a cathartic in constipation, colic, and irritative diarrheas. Dose $\frac{1}{3}$ ss (16.0).

Rickets (*rik'-ets*) [*ῥάχις*, spine]. See *Rachitis*.

Rider's Bone. An osseous formation in the adductor muscles of the leg, from long-continued pressure of the leg against the saddle.

Rigidity (*rij-id'-it-e*) [*rigidus*, stiff]. Stiffness; immobility. **R.**, **Cadaveric**, rigor mortis. **R.**, **Postmortem**-, rigor mortis.

Rigor (*ri'-gor*) [*rigor*, from *rigere*, to be cold]. Chill. **R. mortis**, the muscular rigidity that occurs a short time after death, due to chemic changes resulting in coagulation of the muscle-plasma and the development of an acid reaction.

Rima (*ri'-mah*). A chink. **R. glottidis**, the cleft between the true vocal bands; the glottis.

Rimula (*rim'-u-lah*) [dim. of *rima*, a chink]. A small cleft or fissure, especially of the spinal cord or the cerebellum.

Ring-worm. *Tinea trichophytina*, a contagious disease of the skin due to a vegetable parasite, the trichophyton. See *Tinea*.

Rinne's Test (*rin'-nes*). A test to determine the condition of the various parts of the ear, performed by applying a vibrating tuning-fork first over the mastoid process, leaving it there until the patient seems no longer to hear the sound, and then as quickly as possible bringing it immediately in front of the external

meatus, avoiding all contact with the head or ear. If the patient is then able to hear the sound of the tuning-fork once more, it indicates that the conduction through the air is better than through the bone.

Riolan's Muscle (*ri'-o-lanz*). The ciliary portion of the orbicularis palpebrarum.

Risorius (*ri-so'-re-us*) [*ridere*, to laugh]. Laughing. **R. Muscle**. See *Muscles, Table of*.

Risus (*ri'-sus*) [*L.*]. A grin or laugh. **R. sardonicus**, the sardonic grin, a peculiar grinning distortion of the face produced by spasm of the muscles about the mouth, seen in tetanus.

Ritter's, or Ritter-Valli's Law. The law that if a nerve is separated from its center, the irritability is at first increased, the primary increase as well as the subsequent decrease of irritability traveling from the central to the peripheral end.

Rivini, or Rivinus, Ducts of. The excretory ducts of the sublingual gland.

Rivolta's Disease. Synonym of *Actinomycosis*.

Rob. A confection made of fruit-juice, especially of that of the mulberry.

Roberts' Test. For glucose in the urine. See *Fermentation-test*. Every degree of specific gravity lost corresponds to 0.23 per cent. of glucose.

Robertson's Pupil. See *Pupil, Argyll Robertson's*.

Roborant (*ro'-bo-rant*) [*robur*, an oak]. Tonic, strengthening.

Roccella (*rok-sel'-ah*). A genus of plants of the Roccellæ. **R. tinctoria**, the litmus-plant.

Rochelle Salt (*ro-shel'*). Potassium and sodium tartrate.

Rock-oil. See *Petroleum*.

Rodent Ulcer. Jacob's Ulcer; *noli me tangere*; a superficial ulceration of the skin, occurring usually in the eyelid or the side of the nose, and frequently persisting for years; it is a form of epithelioma.

Rods and Cones, Rod-and-Cone Layer. See *Retina*.

Rolandic (*ro-lan'-dik*). Described by Rolando, as, *e. g.*, the *R. fissure*.

Roller-bandage. A bandage made into a cylindrical roll.

Romberg's Sign or Symptom. The increased incoordination of movement in locomotor ataxia caused by placing the feet in juxtaposition and closing the eyes. It is also called the Brach-Romberg symptom.

Rongeur-forceps (*rông-shur*). A strong pair of forceps for breaking off pieces of bone, especially in enlarging a trephine-opening.

Röntgen Rays. See *X-rays*.

Röntography (*roen-tog'-ra-fe*). See *Skiography*.

Root [*Icel.*, *rot*, *root*]. 1. The descending axis of a plant. 2. The part of an organ embedded in the tissues, as, *e.g.*, the R. of a tooth. **R. of a nerve**, one of two bundles of nerve-fibers, the anterior and posterior roots, joining to form a nerve trunk. **R.-sheath**, the epithelium of the hair-follicle. **R.-zone**, a name given to the column of Burdach of the spinal cord.

Rosa (*ro'-zah*) [*L.*]. See *Rose*.

Rosacea (*ro-sa'-se-ah*). See *Acne rosacea*.

Rosanilin (*ro-san'-il-in*) [*rosa*, *rose*; *anilin*], $C_{20}H_{19}N_3O$. A colorless, crystalline derivative of anilin. R. hydrochlorid or R. acetate is the red dye fuchsin.

Rosary, Rachitic. See *Rachitis*.

Rose (*rös*) [*ῥόδον*, from Arab., *ward*, a rose]. A genus of plants of the order Rosaceae. The petals of the **Pale Rose** (*Rosa centifolia*, U. S. P., *Rosæ centifoliæ petala*, B. P.) are used in the preparation of **R.-water** (*Aqua rosæ fortior*, U. S. P., and *Aqua rosæ*, U. S. P., B. P.) and cold cream (*Unguentum aque rosæ*, U. S. P.). The petals of the **Red Rose** (*Rosa gallica*, U. S. P., *Rosæ gallicæ petala*, B. P.) are slightly astringent and tonic, but are chiefly employed as a vehicle. From them are prepared *Confectio rosæ* (U. S. P.); *Confectio rosæ gallicæ* (B. P.); *Extractum rosæ fluidum* (U. S. P.), used as an astringent and vehicle, dose [ʒ]-ij (4.0-8.0); *Pilulæ aloes et mastiches*, Lady Webster's dinner pill, used as a laxative. The fruit of the **Dog-Rose** (*Rosæ caninæ fructus*, B. P.) is used in Europe as a vehicle. From *Rosa damascena* is prepared *Oleum rosæ* (U. S. P.), attar or otto of rose, employed as a perfume and flavoring agent. **R.-cold**, hay-fever, so-called because believed to be due to exhalations from the rose.

Rosemary (*rös'-ma-re*) [*rosmarinus*, marine dew; from *ros*, dew; *marinus*, marine]. The *Rosmarinus officinalis*, a plant of the order Labiatae, yielding a volatile oil (*Oleum rosmarini*, U. S. P., B. P.), used as a stimulant and in rubefacient liniments. Dose mʒij-vj (0.20-0.39). From it is prepared spirit of R. (*Spiritus rosmarini*, B. P.), used as a perfume and in liniments.

Rosenmüller's Body or Organ. The parovarium.

Roseola (*ro-sē'-o-lah*) [*roseus*, rosy]. 1. Rose-rash, a name given to any rose-colored eruption. 2. Synonym of *Rubella*. **R. cholericæ**, an eruption sometimes appearing in cholera. **R.**, Syphilitic, an eruption of rose-colored spots appearing early in secondary syphilis. **R. typhosa**, the eruption of typhoid or typhus fever. **R. vaccina**, a general rose-colored eruption sometimes occurring during vaccinia.

Rosin (*ros'-in*) [*resina*]. Colophony.

Rosmarinus (*rös'-ma'-ri'-nus*). See *Rosemary*.

Rosolic Acid (*ro-zol'-ik*), $C_{20}H_{16}O_3$. A substance used as a test for acids and alkalies. Acids decolorize it; with alkalies it gives a red color.

Rostellum (*ros-tel'-um*) [dim. of *rostrum*, a beak]. A little beak, especially the hook-bearing portion of the head of certain worms.

Rostrum (*ros'-trum*) [*L.*]. A beak; a projection or ridge, as the R. of the sphenoid bone or of the corpus callosum.

Rot [*AS.*, *rotian*, to rot]. Decay. **R.**, **Potato-**. See *Mildew*.

Rotary (*ro'-ta-re*) [*rotare*, to turn, from *rota*, wheel]. Producing or characterized by rotation.

Rotation (*ro-ta'-shun*) [*rotare*, to turn, from *rota*, wheel]. The act of turning about an axis passing through the center of a body, as, *e.g.*, R. of the eye, R. of the arm.

Rotator (*ro-ta'-tor*) [*rotare*, to turn]. Anything, especially a muscle, that produces rotation.

Rötheln (*rä'-teln*). See *Rubella*.

Rottlera (*rot-le'-rah*). See *Kamala*.

Rotula (*rot'-u-lah*) [dim. of *rota*, wheel]. The patella.

Rouleau (*roo'-lo*) [*Fr.*; pl. *rouleaux*]. A roll, especially a roll of red blood-corpuscles resembling a roll of coins.

Round Ligament. 1. One of the ligaments of the uterus, passing through the inguinal canal. 2. One of the ligaments of the liver, lying in the longitudinal fissure. 3. One of the ligaments of the hip-joint—*ligamentum teres*.

Round Worm. The *Ascaris*.

Rubber (*rub'-er*). Caoutchouc, india-rubber. **R.-dam**, a sheet of rubber used to confine the flow of secretions or of discharges from a wound.

Rubefacient (*ru-be-fa'-se-ent*) [*ruber*, red; *facere*, to make]. 1. Causing redness of the skin. 2. An agent that causes redness of the skin.

Rubella (*ru-bel'-ah*) [dim. of *rubeola*]. Epidemic roseola; German measles; French measles; Rötheln (G.). An acute contagious eruptive disease, of short duration and mild character. After a period of incubation varying from one to three weeks, the disease sets in abruptly with pains in the limbs, sore throat, and slight fever. The eruption appears at the end of the first day, and consists of red papules, and disappears usually without desquamation in about three days. The disease is associated with enlargement of the superficial cervical and posterior auricular glands. See *Exanthemata*, *Table of*.

Rubeola (*ru-bel'-o-lah*) [*rubeus*, red]. See *Measles*.

Rubia (*ru'-be-ah*). The *R. tinctorum* or dyers' madder, containing the coloring principles alizarin ($C_{14}H_8O_4$) and purpurin ($C_{14}H_8O_5$). *R.* is used as a dye.

Rubigo (*ru-bi'-go*) [L.]. Rust.

Rubus (*ru'-bus*) [L.]. Blackberry. A genus of plants of the order Rosaceæ. The bark of the root of *R. canadensis*, *R. villosus*, and *R. trivialis* is the *R.* of the U. S. P., and is used as an astringent tonic in diarrhea. Dose gr. xx-xxx (1.3-2.0). Extractum rubi fluidum (U. S. P.). Dose f 3 ss-j (2.0-4.0). Syrupus rubi. Dose f 3 j-ij (4.0-8.0). The fruit of *R. idæus* (*R. idæus*, U. S. P.), the raspberry, is used to prepare Syrupus rubi idæi, which is used as a tonic.

Ructus (*ruk'-tus*) [L.]. A belching of gas from the stomach.

Rue (*ru*) [*ῥύη*, *ruë*]. The *Ruta graveolens*, a plant of the order Rutaceæ, yielding an oil (*Oleum rutæ*, B. P.), which is a local irritant and has been employed in amenorrhea and menorrhagia. Dose m ij-v (0.13-0.32).

Ruga (*ru'-gah*) [L.]. A fold or ridge.

Rugose (*ru'-gôs*) [*ruga*, a fold]. Characterized by folds.

Rugosity (*ru-gos'-it-e*) [*ruga*, fold]. A condition of being in folds.

Rugous (*ru'-gus*). See *Rugose*.

Rum [Malay, *brain*, rum]. A spirit obtained from the molasses of the sugar-cane by fermentation and distillation.

Rumen (*ru'-men*) [L.]. The first stomach of ruminants, also called the paunch, from

which the food is returned to the mouth for remastication.

Rumex (*ru'-meks*) [L.]. Yellow dock, a genus of plants of the order Polygonaceæ. The root of *R. crispus* (*Rumex*, U. S. P.) is astringent and tonic, and has been employed externally and internally in various diseases of the skin. Extractum rumicis fluidum (U. S. P.). Dose f 3 j (4.0).

Rumination (*ru-min-a'-shun*) [*ruminare*, to chew cud]. See *Merycismus*.

Rump [Icel., *rumpe*, rump]. The region near the end of the backbone; the buttocks.

Rupia (*ru'-pe-ah*) [*ῥύπος*, filth]. A form of eruption occurring especially in tertiary syphilis, and characterized by the formation of large, dirty-brown, stratified, conic crusts.

Rupial (*ru'-pe-al*) [*ῥύπος*, filth]. Resembling or characterized by rupia.

Rupture (*rup'-tūr*) [*rumperé*, to break]. 1. A forcible tearing of a part, as, *e. g.*, *R.* of the uterus, *R.* of the bladder. 2. Hernia.

Rust [AS., *rust*, rust]. 1. The oxid and hydroxid of iron formed on the surface of iron exposed to the air. 2. A disease common on cereals, causing rust-like masses to break out on the tissues of the plant.

Ruta (*ru'-tah*) [L.]. See *Rue*.

Rutting (*rut'-ing*) [ME., *rut*, rut]. The period of sexual excitement in female animals coincident with the rupture of ovisacs.

Ruysch, Membrane of. See *Membrane*.

Rye (*ri*) [AS., *ryga*, rye]. The *Secale cereale* and its grain. The grain is used in the manufacture of bread. See *Graminaceæ*.

S

S. i. The symbol of sulphur. 2. An abbreviation for *sinister*, left; in prescriptions, for *signa*, sign.

S. romanum. The sigmoid flexure of the colon.

Sabadilla (*sab-ad-il'-ah*). Cevadilla, the *Schoenocaulon officinale* (*Asagrea officinalis*), a plant of the order Melanthaceæ, containing the alkaloids veratrin, sabadin, and sabadinin. *S.* is an emetocathartic, and was formerly used as a teniacide and to destroy vermin in the hair. Its chief value is as a source of veratrin. It is official in the B. P.

Sabbatia (*sab-a'-she-ah*) [after Liberatus *Sabbati*, an Italian botanist]. A genus of the order Gentianaceæ. *S. angularis* is the American centaury.

Sabina (*sa-bi'-nah*) [L.]. See *Savine*.

Sabulous (*sab'-u-lus*) [*sabulum*, sand]. Gritty; sandy.

Saccate, or **Saccated** (*sak'-at*, *sak'-a-ted*) [*saccus*, a sac]. Sac-shaped; contained in a sac; encysted.

Saccharated (*sak'-ar-a-ted*) [*σάκχαρον*, sugar]. Containing sugar.

Saccharification (*sak-ar-if-ik-a'-shun*) [*σάκχαρον*, sugar; *facere*, to make]. The act of converting into sugar.

Saccharimeter (*sak-ar-im'-et-er*) [*σάκχαρον*, sugar; *μέτρον*, measure]. An apparatus for determining the amount of sugar in solutions, either in the form of a hydrometer, which indicates the strength in sugar by the specific gravity of the solution; or of a polarimeter, which indicates the strength in

sugar by the number of degrees of rotation of the plane of polarization.

Saccharin (*sak'-ar-in*) [*σακχαρον*, sugar], $C_6H_5SO_2N$. A crystalline substance nearly 280 times sweeter than cane-sugar, and used as a substitute for the latter in diabetes. It is also employed as an antiseptic.

Saccharine (*sak'-ar-in*) [*σακχαρον*, sugar]. Containing sugar; sugary; as sweet as sugar.

Saccharometer (*sak'-ar-om'-et-er*) [*σακχαρομετρον*, sugar; *μετρον*, measure]. See *Saccharimeter*.

Saccharomyces (*sak'-ar-o-mi'-set*) [*σακχαρομυκης*, sugar; *μυκης*, fungus]. A genus of unicellular vegetable organisms of which the yeast-plant is a common example.

Saccharose (*sak'-ar-ös*) [*σακχαροσ*, sugar], $C_{12}H_{22}O_{11}$. 1. A crystalline carbohydrate, cane-sugar, occurring in the juice of many plants, chiefly in sugar-cane, in some varieties of maple, and in beet-roots; it melts at $160^{\circ}C$; at $190-200^{\circ}C$ it changes into a brown noncrystallizable mass called caramel, used in coloring liquids. It is not directly fermentable, and does not reduce alkaline copper-solutions. 2. Any one of a group of carbohydrates isomeric with cane-sugar.

Saccharum (*sak'-ar-um*) [*σακχαρον*, sugar]. Sugar. The S. of the pharmacopeia (Saccharum, U. S. P., S. purificatum, B. P.) is cane-sugar. See *Saccharose*, first definition. S. lactis, milk-sugar.

Sacciform (*sak'-sif-orm*) [*saccus*, a pouch; *forma*, form]. Resembling a sac.

Saccular (*sak'-u-lar*) [*saccus*, sac]. Sac-shaped, as, e. g., a S. aneurysm.

Sacculated (*sak'-u-la-ted*) [*saccus*, sac]. Divided into small sacs.

Sacculation (*sak-u-la'-shun*) [*saccus*, sac]. 1. The state of being sacculated. 2. The formation of small sacs.

Saccule (*sak'-ül*) [*sacculus*, dim. of *sac*, a sac]. 1. A small sac. 2. See *Sacculus*, second definition.

Sacculus (*sak'-u-lus*) [dim. *saccus*, sac]. 1. A small sac. 2. The smaller of two vestibular sacs of the membranous labyrinth of the ear. S. laryngis, the laryngeal pouch between the superior vocal bands and the inner surface of the thyroid cartilage.

Saccus (*sak'-us*) [L.]. A sac. S. endolymphaticus. A small sac contained in the aqueduct of the vestibule and serving to establish a communication between the endolymph and the subdural space.

Sachse's Test. A test for sugar in the urine, consisting in the reduction of the test-solution, a solution of mercuric iodid, potassium iodid, and potassium hydrate. It is generally employed as a quantitative test.

Sacrad (*sa'-krad*) [*sacrum*, the sacrum; *ad*, to]. Toward the sacrum.

Sacral (*sa'-kral*) [*sacrum*, sacrum]. Pertaining to the sacrum.

Sacra media (*sa'-krah me'-de-ah*) [*arteria* understood]. The artery running down the middle of the anterior surface of the sacrum and representing the termination of the aorta.

Sacro- (*sa'-kro-*) [*sacrum*, sacrum]. A prefix denoting relation to the sacrum.

Sacroanterior (*sa-kro-an-te'-re-or*) [*sacrum*, sacrum; *anterior*]. Of the fetus, having the sacrum directed anteriorly.

Sacrococcygeal (*sa-kro-kok-sij'-e-al*) [*sacrum*, sacrum; *κόκκυξ*, coccyx]. Pertaining to the sacrum and the coccyx.

Sacrocoxitis (*sa-kro-koks-i'-tis*). See *Sacroiliac Disease*.

Sacroiliac (*sa-kro-il'-e-ak*) [*sacrum*, sacrum; *ilium*, ilium]. Pertaining to the sacrum and the ilium. S. Disease, an inflammation, usually tuberculous, of the sacroiliac joint, characterized by pain, tenderness, and swelling, and elongation of the limb.

Sacrolumbalis (*sa-kro-lum'-ba'-lis*). See *Muscles*, Table of.

Sacrolumbar (*sa-kro-lum'-bar*) [*sacrum*, sacrum; *lumbus*, loin]. Pertaining to the sacrum and the loins. S. Angle, the angle formed by the articulation of the sacrum and the last lumbar vertebra.

Sacroposterior (*sa-kro-pöst-e'-re-or*) [*sacrum*, sacrum; *posterior*]. Of the fetus, having the sacrum directed backward.

Sacrosciatic (*sa-kro-si-at'-ik*) [*sacrum*, sacrum; *sciatic*]. Pertaining to the sacrum and the ischium, as the S. notch, S. ligaments.

Sacrouterine (*sa-kro-u'-ter-in*) [*sacrum*, sacrum; *uterus*]. Pertaining to the sacrum and the uterus.

Sacrovertebral (*sa-kro-ver'-te-bral*) [*sacrum*; *vertebra*, vertebra]. Pertaining to the sacrum and the vertebrae.

Sacrum (*sa'-krum*) [*sacer*, sacred; *os*, bone, understood]. A curved triangular bone composed of five united vertebrae, situated between the last lumbar vertebra above, the coccyx below, and the two ossa innominata on either side, and forming the posterior boundary of the pelvis.

Saddle-joint. An articulation in which each surface is concave in one direction and convex in the other.

Saddle-nose. A nose of which the bridge is sunken in.

Saemisch's Ulcer (*sa'-mish*). A serpiginous ulcer of the cornea. See *Diseases*, Table of.

Sæptum (*sæp'-tum*). See *Septum*.

Saffron (*saf'-ron*) [Arab., *zafarân*, saffron]. The *Crocus sativus*, a plant of the order Iridaceæ. Its stigmas (Crocus, U. S. P., B. P.) contain a glucosid, coloring matter (crocin),

and a bitter principle. S. is used as a coloring and flavoring agent, and in the form of a tea to bring out the eruption of the exanthematous diseases.

Safranin (*saf'-ra-nin*) [Fr., *safran*, saffron], $C_{18}H_{18}N_4$. A coal-tar dye used in microscopy, especially in studying karyokinesis. It is a powerful cardiac and respiratory poison.

Safrene (*saf'-ren*) [Fr., *safran*, saffron], $C_{10}H_{18}$. A hydrocarbon obtained from sassafras.

Safrol (*saf'-rol*) [Fr., *safran*, saffron; *oleum*, oil], $C_{10}H_{10}O_2$. The stearoptene of the oil of sassafras, used in headache, neuralgia, and subacute rheumatism. Dose m_x -xx (0.65-1.3).

Sagapenum (*sag-a-pe'-num*). A fetid gum-resin, believed to be the concrete juice of *Ferula persica*. Its properties resemble those of asafetida and galbanum.

Sage (*saj*) [*salvia*]. The *Salvia officinalis*, a plant of the order Labiatae. Its leaves (*Salvia*, U. S. P.) contain several terpenes, an oil, salviol, $C_{10}H_{18}O$, and camphor. S. is tonic, astringent, and aromatic, and is used in dyspepsia. It was formerly employed in colliquative sweats. Dose gr. xx-xxx (1.3-2.0).

Sagittal (*saj'-it-al*) [*sagitta*, an arrow]. 1. Arrow-like, as the S. suture of the skull. 2. Pertaining to the anteroposterior median plane of the body. S. Nucleus, the nucleus of the oculomotor nerve.

Sago (*sa'-go*) [Malay, *sagu*, sago]. The starch obtained from the sago-palms, used as a food and demulcent. S.-spleen, a spleen presenting on section the appearance of sago-grains, as a result of amyloid degeneration of the Malpighian bodies.

Saigon Cinnamon. A variety of cinnamon obtained from Saigon, the capital of French Cochinchina.

Saint Anthony's Fire. Erysipelas; anthrax.

Saint Gothard's Disease. Ankylostomiasis.

Saint Hubert's Disease. Hydrophobia.

Saint Ignatius' Bean. See *Ignatia*.

Saint Vitus' Dance. Chorea.

Sal [*sal*, salt]. 1. Salt. 2. Any substance resembling salt. S. aeratus, sodium bicarbonate. S. alembroth. See *Alembroth*. S. ammoniac, ammonium chlorid. S. communis, common salt. S. de duobus, potassium sulphate. S. Glauberi, sodium sulphate. S. kissingense, a salt obtained from the mineral springs of Kissingen, in Bavaria. S. seignetti, potassium and sodium tartrate. S. sodæ, crystallized sodium carbonate. S. volatiliss, ammonium carbonate.

Salaam Convulsion (*sa-lahm'*) [Arab., *salm*, saluting]. A clonic spasm of the muscles of the trunk, producing a bowing movement; it is usually due to hysteria.

Salamandarin (*sal-a-man'-dar-in*). See *Leukomains*, Table of.

Salep (*sal'-ep*) [Arab., *sahleb*, salep]. The dried tubers of various species of the genus *Orchis* and the genus *Eulophia*. S. is used as a food, like sago and tapioca.

Saleratus (*sal-er-a'-tus*). See *Sal aeratus*.

Salicin (*sal'-is-in*) [*salix*, willow], $C_{13}H_{18}O_7$. A crystalline glucosid found in the bark and leaves of the willow. S. (*Salicinum*, U. S. P., B. P.) is used as a substitute for salicylic acid, in doses of gr. xx-xxx (1.3-2.0).

Salicylage (*sal'-is-il-aj*) [*salix*, willow]. The addition of salicylic acid to foods for their preservation.

Salicylamid (*sal-is-il'-am-id*) [*salix*, willow; *amid*], $C_6H_4(OH)CONH_2$. A tasteless compound produced by treating methyl salicylate with an alcoholic solution of ammonia, and used as a substitute for salicylic acid.

Salicylate (*sal-is-il-at*) [*salix*, willow]. A salt of salicylic acid. The salicylates of lithium, methyl, and sodium, which are official, and those of ammonium and strontium, which are unofficial, are used in rheumatism, in doses of gr. x-xv (0.65-1.0). Bismuth S. is employed as an intestinal antiseptic; naphthol S. is betol; phenyl S. is salol.

Salicylated (*sal-is-il-a-ted*) [*salix*, willow]. Impregnated with salicylic acid.

Salicylic Acid (*sal-is-il'-ik*). See *Acid*, *Salicylic*.

Salicyluric Acid (*sal-is-il-u'-rik*) [*salix*, willow; *uric*]. A compound of glycol and salicylic acid found in the urine after the administration of salicylic acid.

Salifiable (*sal-if-i'-a-bl*) [*sal*, salt; *feri*, to become]. Forming a salt by union with an acid.

Saligenin (*sal-ij'-en-in*) [*salix*, willow; *γεννᾶν*, to produce], $C_7H_8O_4$. Orthooxybenzylalcohol, a substance obtained from salicin by boiling with dilute hydrochloric or sulphuric acid.

Salimeter (*sal-im'-et-er*) [*sal*, salt; *μέτρον*, measure]. An hydrometer for ascertaining the strength of saline solutions.

Saline (*sa'-len* or *sa'-lin*) [*sal*, salt]. 1. Salty; containing salt or substances resembling salt. S. Solution, a 0.6 per cent. solution of sodium chlorid; normal S. solution. 2. A salt of an alkali or alkaline earth.

Salipyrin (*sa-lip'-ir-in*, *sal-e-pi'-rin*) [*salix*, willow; *πῦρ*, fire], $C_{11}H_{13}N_3O_3C_7H_5O_2$. Salicylate of antipyrin, consisting of 57.7 parts of salicylic acid and 42.3 parts of antipyrin; it is soluble in water, and is used in rheumatism, neuralgia, and as an antipyretic. Dose gr. xv-xxx (1.0-2.0).

Saliretin (*sal-e-ret'-in*) [*salix*, willow; *πῦρ*,

an, resin], $C_{14}H_{14}O_3$. An amorphous resinous body, produced by treating saligenin with acids.

Saliva (*sa-li'-vuh*) [L.]. The mixed secretion of the parotid, submaxillary, sublingual, and mucous glands of the mouth. It is opalescent, tasteless, alkaline, and has a specific gravity of from 1004-1009, and contains serum-albumin, globulin, mucin, urea, an amylolytic ferment called ptyalin, and a proteolytic and a lipolytic ferment; also, salts, among which is potassium sulphocyanid, derived especially from the parotid gland. Among formed elements are epithelial cells, salivary corpuscles, and bacteria. The functions of saliva are to moisten the food and lubricate the bolus, to dissolve certain substances, to facilitate tasting, to aid in deglutition and articulation, and to digest starches, which it converts into maltose, dextrin, and glucose.

Salivary (*sal'-iv-a-ry*) [*saliva*, saliva]. Pertaining to or producing saliva; formed from saliva. **S. Calculus**, a calcareous concretion found in the salivary ducts.

Salivate (*sal'-iv-at*) [*salivare*, to spit out]. To cause an excessive discharge of saliva.

Salivation (*sal'-iv-a'-shun*) [*saliva*, saliva]. An excessive secretion of saliva; a condition produced by mercury, pilocarpin, and by nervous disturbances.

Salix (*sa'-liks*) [L.]. See *Willow*.

Salol (*sal'-ol*) [*salix*, willow], $C_{13}H_{10}O_3$. Phenyl salicylate; a white crystalline substance, used as an intestinal and urinary antiseptic, and as a substitute for salicylic acid. It is decomposed in the intestines into salicylic acid and phenol. Dose gr. v-xv (0.32-1.0).

Salophen (*sal'-o-fen*) [*salix*, willow], $C_8H_5OH.CO_2.C_6H_4.NH(C_2H_5O) = C_{13}H_{13}NO_4$. Acetylparamidophenyl salicylate, a crystalline substance containing fifty per cent. of salicylic acid, and used as a substitute for the latter, and as an intestinal antiseptic. Dose gr. xv (1.0).

Salpingectomy (*sal-pin-jek'-to-me*) [*σάλπιγξ*, tube; *ἐκτομή*, excision]. Excision of the Fallopian tube.

Salpingitic (*sal-ping-jit'-ik*) [*σάλπιγξ*, tube; *ιτις*, inflammation]. Pertaining to or affected with salpingitis.

Salpingitis (*sal-ping-jit'-tis*) [*σάλπιγξ*, tube; *ιτις*, inflammation]. 1. Inflammation of the Fallopian tube. 2. Inflammation of the Eustachian tube.

Salpingo- (*sal-ping'-go-*) [*σάλπιγξ*, tube]. A prefix denoting relation to the Fallopian or the Eustachian tube.

Salpingo-oophorectomy (*sal-ping-go-o-o-for-ek'-to-me*) [*σάλπιγξ*, tube; *ὄον*, egg; *φορέω*, bearing; *ἐκτομή*, excision]. Excision of the Fallopian tube and the ovary.

Salpingo-oophoritis (*sal-ping-go-o-o-for-itis*) [*σάλπιγξ*, tube; *ὄον*, egg; *φορέω*, bearing; *ιτις*, inflammation]. Inflammation of the Fallopian tube and the ovary.

Salpingostaphylinus (*sal-ping-go-staf-il-i'-nus*) [*σάλπιγξ*, tube; *στονύλη*, uvula]. The abductor muscle of the Eustachian tube. **S. internus**. Synonym of *Levator palati*. See *Muscles*, Table of.

Salpingostomy (*sal-ping-gos'-to-me*) [*σάλπιγξ*, tube; *στόμα*, mouth]. The operation of making an artificial fistula between a Fallopian tube and the body-surface.

Salpingotomy (*sal-ping-got'-o-me*) [*σάλπιγξ*, tube; *τομή*, a cutting]. The operation of cutting into a Fallopian tube.

Salt [*sal*, salt]. 1. Sodium chlorid. 2. Any compound of a base and an acid. **S., Acid**, a salt formed from a dibasic or polybasic acid in which only a part of the replaceable hydrogen-atoms has been replaced by the base. **S., Basic**, a S. containing an excess of the basic element, and formed by the union of a normal salt with a basic oxid or hydroxid. **S., Bay-**, sodium chlorid; also the sea-salt obtained by the evaporation of sea-water by solar heat. **S., Common**, sodium chlorid. **S., Diuretic**, potassium acetate. **S., Double**, one in which the hydrogen-atoms of an acid are replaced by two metals. **S., Epsom-**, magnesium sulphate. **S.-frog**, a frog from whose vascular system all blood has been artificially removed and replaced by normal salt-solution; it is also known as Cohnheim's frog. **S., Glauber's**, sodium sulphate. **S., Halogen**, **S., Haloid**, any salt of the halogen-elements, bromin, chlorin, fluorin, and iodin. **S., Monsel's**, subsulphate of iron, used chiefly in solution as a styptic. **S., Neutral**, one formed by the replacement of all the hydrogen-atoms of an acid by a base or a radicle. **S., Normal**. Synonym of *S., Neutral*. **S.-rheum**, chronic eczema. **S., Rochelle-**, sodium and potassium tartrate. **S., Rock-**, native sodium chlorid, occurring in crystalline masses. **S., Sea-**, the sodium chlorid obtained by the evaporation of sea-water. **S., Smelling**, any pungent, irritant salt which when inhaled usually acts reflexly as a respiratory or circulatory stimulant. Ammonium carbonate is generally used. **S.-solution**, a solution of sodium chlorid in distilled water. One containing from 0.6 to 0.75 per cent. of sodium chlorid is known as a *normal* or *physiologic salt-solution*, and is used in physiologic experiments on living tissues. In medicine, it has been employed to restore to the system the fluids lost by severe hemorrhage or profuse diarrheal discharges. The solution is introduced into the subcutaneous tissues or into a vein; sometimes also into the rectum.

Salts. A saline cathartic, especially magnesium sulphate, sodium sulphate, or Rochelle-salt.

Saltatoric, Saltatory (*sal-tat-or'-ik*, *sal'-tat-o-re*) [*saltare*, to dance]. Dancing or leaping. **S. Spasm**, a clonic spasm that causes the patient to leap or jump when he attempts to stand.

Saltpeter (*sawlt-pe'-ter*) [*sal*, salt; *πῆτρα*, rock]. Potassium nitrate. **S.**, Chile-, sodium nitrate.

Salubrious (*sa-lu'-bre-us*) [*salus*, health]. Healthful.

Salufer (*sal'-u-fer*) [*salus*, health; *ferre*, to bring]. Sodium silicofluorid.

Salve (*sahv*) [*AS.*, *sealf*, salve]. Ointment. **S.**, Deshler's, compound resin cerate.

Salvia (*sal'-ve-ah*) [*L.*]. See *Sage*.

Salzer's Operation. Excision of the whole of the third division of the fifth nerve.

Sambucus (*sam-bu'-kus*) [*L.*]. Elder; a shrub or tree of the order Caprifoliaceæ. The flowers of **S. canadensis** (*Sambucus*, U. S. P., B. P.) and the berries are sudorific; the latter have been used as an alterative in rheumatism and syphilis. The inner bark has been employed in epilepsy, dropsy, and various chronic diseases. *Aqua sambuci* (B. P.) is used as a vehicle.

Sanative (*san'-a-tiv*) [*sanare*, to heal]. Promoting health; healing.

Sanatorium (*san-at-o'-re-um*) [*sanare*, to heal]. An establishment for the treatment of the sick; especially a private hospital. See *Sanitarium*.

Sanatory (*san'-at-o-re*). See *Sanative*.

Sand [*AS.*, *sand*, sand]. An aggregation of fine grains of silicic oxid. **S.-bath.** 1. A vessel containing dry sand in which a substance requiring a slowly rising or uniform temperature may be heated. 2. See *Bath*.

Sandalwood. 1. Red saunders (*Santalum rubrum*, U. S. P., *Pterocarpus ligni*, B. P.), the wood of *Pterocarpus santalinus*, of the order Leguminosæ. It is used as a coloring agent. 2. The wood of *Santalum album*, of the order Santalaceæ, containing a volatile oil, *Oleum santali* (U. S. P., B. P.), used in bronchitis and gonorrhea. Dose $\text{m}\text{xv}-\text{xx}$ (1.0-1.3).

Sandarac (*san'-dar-ak*). A white, transparent resin produced by *Callitris quadrivalvis*, a tree of North Africa. It is now little used except as a varnish and incense.

Sane (*sān*) [*sanus*, whole]. Of sound mind.

Sanguification (*sang-gwif-ik-a'-shun*) [*sanguis*, blood; *facere*, to make]. 1. The formation of blood. 2. Conversion into blood, as, *e. g.*, the **S.** of substances absorbed from the intestinal tract.

Sanguinaria (*sang-gwin-a'-re-ah*) [*sanguis*, blood]. Blood-root, a genus of plants of the order Papaveraceæ. The rhizome of **S.**

canadensis (*Sanguinaria*, U. S. P.) contains several alkaloids, of which sanguinarin, $\text{C}_{20}\text{H}_{15}\text{NO}_4$, is the most important; it is emetic and narcotic; in large doses an irritant and convulsant. It is used chiefly as an expectorant in bronchitis. Dose, as an expectorant, gr. j-v (0.065-0.32); as an emetic, gr. v-x (0.32-0.65). *Extractum sanguinarie fluidum* (U. S. P.), dose $\text{m}\text{j}-\text{x}$ (0.065-0.65); *Tinctura sanguinarie* (U. S. P.), dose $\text{m}\text{xxx}-\text{lx}$ (2.0-4.0).

Sanguine (*sang'-gwin*) [*sanguis*, blood]. 1. Resembling blood; bloody. 2. Hopeful; active, as **S.** temperament.

Sanguineous (*sang'-gwin'-e-us*) [*sanguis*, blood]. 1. Pertaining to the blood; containing blood. 2. Sanguine. **S. Cyst**, a cyst containing blood-stained fluid.

Sanguinolent (*sang-gwin'-o-lent*) [*sanguis*, blood]. Tinged with blood.

Sanguis (*sang'-gwis*) [*L.*]. Blood.

Sanguisuga (*sang'-gwi-su'-gah*) [*sanguis*, blood; *sugere*, to suck]. See *Leech*.

Sanies (*sa'-ne-tz*) [*L.*]. A thin, fetid, greenish, seropurulent fluid discharged from an ulcer, wound, or fistula.

Sanious (*sa'-ne-us*) [*sanies*]. Pertaining to or resembling sanies, as **S.** pus.

Sanitarium (*san'-it-a'-re-um*) [*sanitas*, health]. A place where the conditions are such as especially to promote health; a resort for convalescents.

Sanitary (*san'-it-a-re*) [*sanitas*, health]. Pertaining to health. **S. Science**, the science that includes a consideration of all that can be done for the prevention of disease and the promotion of the public health.

Sanitation (*san-it-a'-shun*) [*sanitas*, health]. The act of securing a healthful condition; the application of sanitary measures.

Sanity (*san'-it-e*) [*sanitas*, from *sanus*, sound]. Soundness of mind.

Santalum (*san'-tal-um*). See *Sandal-wood* (2d def.).

Santonica (*san-ton'-ik-ah*) [*σάντονικον*, wormwood]. Levant wormseed, the unexpanded flower-heads of *Artemisia pauciflora* (U. S. P.), or *A. maritima*, var., *Stechmanniana* (B. P.), of the order Compositæ, the essential constituent of which is santonin, $\text{C}_{15}\text{H}_{18}\text{O}_3$ (*Santoninum*, U. S. P., B. P.). *Santonin* is a neutral crystalline principle, producing, in overdoses, xanthopsia, giddiness, stupor, at times convulsions, and death from failure of respiration. The urine is colored yellow. **S.** and santonin are used as vermicides against the lumbricoid worm. Dose of **S.** gr. x-xxx (0.65-2.0); of *Santonin*, gr. j-ij (0.065-0.13); *Trochisci santonini* (U. S. P.) contain each about one-half grain (0.033) santonin; *Trochisci santonini* (B. P.) contain each one grain santonin.

Sodium santoninate was formerly used as a substitute for santonin, but has produced poisoning.

Santonin (*san'-to-nin*). See *Santonica*.

Santorini's Cartilage (*sahn-to-re'-nēz*) [*Santorini*, an Italian anatomist]. See *Cartilage*.

Saphenous (*sa-fe'-nus*) [*σαφηνής*, manifest].

Apparent; superficial; manifest; applied to two veins of the lower limb, the internal or long S. vein and external or short S. vein, situated just beneath the surface; also applied to the nerves accompanying these veins. **S. Opening**, an opening in the fascia lata at the upper part of the thigh through which the long S. vein and nerve pass.

Sapid (*sa'-pid*) [*sapere*, to taste]. Capable of being tasted.

Sapo (*sa'-po*) [L.]. See *Soap*.

Saponaceous (*sap-o-na'-se-us*) [*sapo*, soap]. Having the nature of soap.

Saponification (*sa-pon-if-ik-a'-shun*) [*sapo*, soap; *facere*, to make]. The act of converting into soap; the process of treating a neutral fat with an alkali, which combines with the fatty acid, forming a soap.

Saponify (*sa-pon'-e-fi*) [*sapo*, soap; *facere*, to make]. To convert into soap; to convert a neutral fat by the action of an alkali into free glycerol and a salt of the alkali, the latter forming a soap.

Saponin (*sap'-o-nin*) [*sapo*, soap], $C_{27}H_{54}O_{18}$. A glucosid contained in the roots of soapwort and other plants, and in aqueous solution forming a strong lather.

Sapotoxin (*sap-o-tox'-in*) [*sapo*, soap; *ροξικόν*, poison], $C_{17}H_{30}O_{10}$. A poisonous glucosid obtained from saponin.

Sapphism (*saf'-izm*) [from *Σαφώ*, *Sappho*, a Greek poetess]. Tribadism.

Sapremia (*sap-re'-me-ah*) [*σαπρός*, putrid; *αἷμα*, blood]. A febrile condition due to introduction into the blood of the ptomaines produced in putrefaction.

Saprin (*sap'-rin*) [*σαπρός*, putrid]. A non-poisonous ptomaine formed in the putrefaction of animal tissues.

Saprogenic, Saprogenous (*sap-ro-jen'-ik*, *sap-roj'-en-us*) [*σαπρός*, putrid; *γεννᾶν*, to beget]. 1. Causing putrefaction. 2. Produced by putrefaction.

Saprophyte (*sap-ro-fi't*) [*σαπρός*, putrid; *φυτόν*, a plant]. A vegetable organism living on dead organic matter.

Saprophytic (*sap-ro-fi't-ik*) [*σαπρός*, putrid; *φυτόν*, plant]. Growing in dead organic matter, as, e. g., *S. bacteria*.

Sarcin (*sar'-sin*) [*σάρξ*, flesh]. See *Hypoxanthin*.

Sarcina (*sar-si'-nah*) [L., a bundle]. A genus of Schizomycetes consisting of cocci dividing in three directions, thus producing cubic masses. See *Bacteria*, Table of.

Sarco- (*sar'-ko-*) [*σάρξ*, flesh]. A prefix denoting composed of or pertaining to flesh.

Sarcocele (*sar'-ko-sel*) [*σάρξ*, flesh; *κύστη*, a tumor]. Any fleshy swelling of the testicle.

Sarcocystis (*sar'-ko-sis'-tis*) [*σάρξ*, flesh; *κύστις*, cyst]. A group of the sporozoa. **S. Miescheri**, a parasite found in pork and beef.

Sarcode (*sar'-kōd*) [*σάρξ*, flesh]. Proto-plasm.

Sarcolactic Acid (*sar-ko-lak'-tik*). See *Acid*.

Sarcolemma (*sar-ko-lem'-ah*) [*σάρξ*, flesh; *λίμμα*, husk]. The delicate membrane enveloping a muscle fiber.

Sarcoma (*sar'-ko'-mah*) [*σάρξ*, flesh; *μαῖα*, tumor]. A tumor made up of embryonal connective tissue. It is characterized by a great preponderance of cells and very little homogeneous or fibrillar intercellular substance. **S., Alveolar**, one in which groups of sarcoma-cells are contained in alveolar spaces. **S., Angiolithic**. Synonym of *Pseudomoma*. **Chondrosarcoma**, one containing cartilaginous tissue. **S., Giant-celled**, one containing giant-cells as a prominent feature. **Lymphosarcoma**, a round-celled sarcoma in which the cells are held in a reticulum.

S., Melanotic, a sarcoma, usually spindle-celled, in which the cells contain melanin. **S., Myeloid**. See *S., Giant-celled*. **Myxosarcoma**, one which in part has undergone myxomatous degeneration. **S., Round-celled**, one made up of round cells. There are two varieties, the small round-celled and the large round-celled sarcoma. **S., Spindle-celled**, one made up of spindle-cells. It is also known as recurrent fibroid.

Sarcomatosis (*sar-ko-mat'-o'-sis*) [*sarcoma*]. The formation of multiple sarcomatous growths in various parts of the body. **Sarcomatous** (*sar-ko'-mat-us*) [*σάρξ*, flesh; *μαῖα*, tumor]. Of the nature of or resembling sarcoma.

Sarcomere (*sar'-ko-mēr*) [*σάρξ*, flesh; *μερὸς*, a part]. One of the segments into which a muscle-fibril appears to be divided by transverse septa.

Sarcoplasm (*sar'-ko-plasm*) [*σάρξ*, flesh; *πλάσσειν*, to mold]. The hyaline or finely granular interfibrillar material of muscle-tissue.

Sarcoplast (*sar'-ko-plast*) [*σάρξ*, flesh; *πλάσσειν*, to mold]. A cell lying between muscular fibrillae and capable of developing into a muscular fiber.

Sarcoptes (*sar'-kop'-tes*) [*σάρξ*, flesh; *κόπτειν*, to cut]. A genus of mites, including *S. hominis*, the itch mite.

Sarcosepsis (*sar'-ko-sep'-sis*) [*σάρξ*, flesh; *σεπσις*]. The presence of bacteria directly in the tissues.

Sarcosin (*sar'-ko-sin*) [*σάρξ*, flesh], $C_3H_5NO_4$. Methylglycocoll, a crystalline sub-

stance produced when creatin and caffein are heated with baryta.

Sarcosporidia (*sar-ko-spor-id'-e-ah*) [*σάρξ*, flesh; *σπορός*, seed]. A variety of psorosperms found in the muscles of cattle, sheep, swine, and other mammals.

Sarcostyle (*sar'-ko-stil*) [*σάρξ*, flesh; *στυλος*, a pillar]. One of the fine longitudinal fibrillæ of which a striated muscle-fiber is composed and into which it can be split up.

Sarcous (*sar'-kus*) [*σάρξ*, flesh]. Pertaining to muscle. **S. Element**, one of the dark prisms of the ultimate fibrillæ of striped muscle-fibers.

Sardonic Grin (*sar-don'-ik*) [*Σαρδία*, Sardinia, because resembling the grimace produced by eating a certain Sardinian plant]. See *Risus sardonius*.

Sarkin (*sar'-kin*). See *Hypoxanthin*.

Sarsaparilla (*sar-sap-ar-il'-ah*) [Sp.]. The *Smilax officinalis* and other species of *Smilax*, of the order Liliaceæ. The root (*Sarsaparilla*, U. S. P., *Sarsæ radix*, B. P.) contains a crystalline glucosid, parillin, $C_{26}H_{70}O_{18}$. S. has been employed as an alternative in syphilis, rheumatism, and scrofulous affections. Preparations and doses: Decoctum sarsæ (B. P.), f3 iv-vj (128.0-192.0). Decoctum sarsaparillæ compositum (U. S. P.), Decoctum sarsæ compositum (B. P.), f3 iv-vj (128.0-192.0); Extractum sarsaparillæ fluidum (U. S. P.), mxxx-lx 2.0-4.0; Extractum sarsæ liquidum (B. P.), f3 j-iv (8.0-16.0); Extractum sarsaparillæ fluidum compositum (U. S. P.), mxxx-lx (2.0-4.0); Syrupus sarsaparillæ compositus (U. S. P.), f3 j-iv (4.0-16.0).

Sartian Disease (*sar'-shun*). An endemic affection of the tropics, characterized by red indurated spots that finally ulcerate.

Sartorius (*sar-to'-re-us*) [*sartor*, tailor]. The tailor's muscle, so-called from being concerned in crossing the one leg over the other. See *Muscles*, Table of.

Sassafras (*sas'-a-fras*) [Sp., from *saxifraga*, from *saxum*, rock; *frangere*, to break]. The *S. variifolium*, a tree of the order Laurineæ. The root-bark (*Sassafras*, U. S. P., *Sassafras radix*, B. P.) is employed as an aromatic stimulant. The pith (*Sassafras medulla*, U. S. P.) yields a mucilage (*Mucilago sassafras medullæ*, U. S. P.) that is used as an application to inflamed eyes, and as a demulcent drink in inflammation of the mucous membranes and kidneys.

Sassy Bark (*sas-e*). See *Casca*.

Saturated (*sat'-u-ra-ted*) [*saturare*, to fill]. 1. Of a liquid, containing in solution all of a substance that it can dissolve. 2. Of a chemic compound, having all the affinities of its component atoms satisfied, a term especially applied to the hydrocarbons.

Saturation (*sat-u-ra'-shun*) [*saturare*, to fill]. 1. A state in which a liquid holds in solution all of a substance that it can dissolve; the state of being or becoming saturated. 2. Of a chemic compound, a state in which the affinities of all its atoms are satisfied.

Satureja (*sa-tur-e'-ya*). A plant of the order Labiatæ. **S. hortensis** resembles thyme, and is used as a culinary herb.

Saturnine (*sat'-ur-nin*) [*Saturnus*, a Roman deity; the alchemists' name for lead]. Pertaining to or produced by lead.

Saturnism (*sat'-ur-nism*) [*Saturnus*, a Roman deity; the alchemists' term for lead]. Lead-poisoning; plumbism.

Satyriasis (*sat'-ir-i'-as-is*) [*σάτυρος*, a satyr]. Excessive venereal desire in the man.

Saunders (*sawm'-ders*). See *Santalum*.

Sauriderma (*saw-re-der'-mah*) [*σαύρα*, lizard; *δέρμα*, skin]. Ichthyosis.

Sauriosis (*saw-re-o'-sis*) [*σαύρα*, lizard]. Ichthyosis.

Sausage-poisoning. A state of gastroenteritis produced by the ingestion of decomposed sausage. It is also known as botulism or allantiasis.

Savill's Disease. An epidemic skin-disease characterized by the appearance of a papular rash, followed by a branny desquamation, and by marked constitutional symptoms. A fatal result may follow.

Savine (*sav'-in*). *Juniperus sabina*, a shrub of the order Conifere. The tops (*Sabina*, U. S. P., *Sabinæ cacumina*, B. P.) contain a volatile oil (*Oleum sabinæ*) and possess marked irritant properties. S. is employed in amenorrhea, chronic rheumatism, gout, and as a local application to warts, ulcers, and parasitic affections of the skin. Preparations and doses: Extractum sabinæ fluidum (U. S. P.), mlij-vijj (0.20-0.52); Tinctura sabinæ (B. P.), mxx-f3j (1.3-4.0); Unguentum sabinæ (B. P.). Dose of the oil, mlij-v (0.13-0.32).

Saw [AS., *saga*, saw]. An instrument having a thin blade with sharp teeth on one edge, and used for dividing bones and other hard substances. **S.**, Butcher's, one in which the blade can be fixed at any angle. **S.**, Chain-, one in which the teeth are set in links movable upon each other, the saw being moved by pulling alternately upon one and the other handle. **S.**, Crown-. See *Trephine*. **S.**, Hey's, a serrated disc affixed to a handle, and used for enlarging an opening in a bone.

Sayre's Apparatus or Jacket. A jacket of plaster-of-Paris molded to support the spine in diseases of the vertebral column.

Scab [*scabere*, to scratch]. 1. The crust formed by the desiccation of the secretion of an ulcer. 2. Scabies.

Scabies (*skā'-be-ē*) [*scabere*, to scratch]. Itch; a disease of the skin caused by an animal parasite, the *Sarcoptes scabiei*, or itch-mite. The insect forms burrows or cuniculi beneath the skin, and causes irritation, with vesicles, papules, or pustules, which are frequently modified by scratching.

Scabrities (*skā-brīt'-ē-ēs*) [*scaber*, rough]. Roughness; scabbiness.

Scala (*skā'-lah*) [*l.*]. A staircase or ladder.

S. media, the space between the membrane of Reissner and the basilar membrane, containing the essential peripheral organs of hearing. **S. tympani**, the canal lying below the osseous lamina and the basilar membrane of the internal ear. **S. vestibuli**, the canal bounded by the osseous lamina and the membrane of Reissner. See *Ear*.

Scald (*skawld*) [*excaldere*, to wash in hot water]. The burn caused by hot liquids or vapors.

Scald (*skawld*) [*Icel.*, *skalli*, a bare head]. A disease of the skin accompanied by the formation of scabs.

Scale (*skāl*) [*AS.*, *scale*, a husk; a scale]. The dry semiopaque lamina of horny epidermis, shed from the skin in health and in various diseases.

Scale [*scala*, ladder]. Anything bearing marks placed at regular intervals and used as a standard in measuring, as barometric S.

Scalene (*skā'-lēn*) [*σκαληνός*, uneven]. Having unequal sides. **S. Muscle**. See *Scalenus, Muscles, Table of*.

Scall (*skawld*). See *Scald*, 2d definition.

Scalp [*AS.*, *scale*, a shell or husk]. The hairy integument covering the cranium.

Scalpel (*skāl-pel'*) [*scalpere*, to cut]. A small knife having a convex edge.

Scalprum (*skāl'-prum*) [*scalpere*, to scrape]. A toothed raspatory used in trephining and in removing carious bone.

Scaly (*skā'-lē*) [*AS.*, *scale*, scale]. 1. Resembling scales; characterized by scales, as S. desquamation. 2. Covered with or having scales.

Scanning, or **Scanning Speech** [*scandere*, to climb]. A peculiar, slow, and measured form of speech, occurring in various nervous affections, especially in multiple sclerosis.

Scaphocephalic, **Scaphocephalous** (*skaf-o-sef'al-ik*, *skaf-o-sef'-al-us*) [*σκάφη*, a skiff; *κεφαλή*, head]. Having a boat-shaped head, from early ossification of the sagittal suture, which projects like the keel of a boat.

Scaphocephaly (*skaf-o-sef'-al-ē*) [*σκάφη*, boat; *κεφαλή*, head]. The condition of having a skull characterized by a projecting keel-like sagittal suture, due to its premature ossification.

Scaphoid (*skāf'-oid*) [*σκάφη*, boat; *εἶδος*, like]. Boat-shaped. **S. Abdomen**, the

sunken abdomen, seen in meningitis and in great emaciation. **S. Bone**, a name given to a boat-shaped bone of the tarsus and of the carpus. **S. Fossa**. 1. A depression in the base of the internal pterygoid plate of the sphenoid bone. 2. A depression between the helix and antihelix of the auricle.

Scapula (*skap'-u-lah*) [*L.*]. The shoulder-blade, the large, flat, triangular bone forming the back of the shoulder.

Scapular (*skap'-u-lar*) [*scapula*, the shoulder-blade]. Pertaining to the shoulder-blade.

S. Line, a vertical line drawn on the back through the inferior angle of the scapula. **S. Point**, a tender point developed in neuralgia of the brachial plexus and situated at the inferior angle of the scapula. **S. Reflex**. See *Reflexes, Table of*. **S. Region**, the region of the back corresponding to the position of the scapula, the spine of which divides it into a supraspinous and an infraspinous region.

Scapulary (*skap'-u-lar-ē*) [*scapula*, scapula]. A bifurcated bandage, the two ends of which pass over the shoulders, while the single end passes down the back, all three being fastened to a body-bandage.

Scapulo- (*skap'-u-lo-*) [*scapula*, scapula]. A prefix denoting relation to the scapula.

Scapuloclavicular (*skap'-u-lo-kla-vik'-u-lar*) [*scapula*, scapula; *clavus*, key]. Pertaining to the scapula and the clavicle.

Scapulohumeral (*skap'-u-lo-hu'-mee-al*) [*scapula*, scapula; *humerus*, humerus]. Pertaining to the scapula and the humerus.

Scar (*skar*) [*L.*, *eschara*, scar, from *εσχάρα*, a fire-place]. See *Cicatrix*.

Scarfskin (*skarft'-skin*) [*AS.*, *scarfe*, a fragment; *skin*]. The epidermis or cuticle.

Scarification (*skar-if-ik-a'-shun*) [*scarifi-care*, to scarify]. The operation of making numerous small, superficial incisions.

Scarificator (*skar-if-ik-a'-tor*) [*scarificare*, to scarify]. An instrument used in scarification, consisting of a number of small lancets operated by a spring.

Scarify (*skar'-if-i*) [*scarificare*, to scarify]. To make a number of small, superficial incisions.

Scarlatina (*skar-lat-e'-nah*) [*Pers.*, *sagānā*, scarlet cloth]. See *Scarlet Fever*. **S. anginosa**, scarlet fever with marked inflammation of the throat. **S. maligna**. See *Scarlet Fever*.

Scarlatinal, **Scarlatinous** (*skar-lat-i'-nah*, *skar-lat-i'-nus*, *skar-lat'-in-us*) [*scarlatinus*, scarlet]. Pertaining to or caused by scarlet fever. **S. Nephritis**, the acute catarrhal nephritis arising in the course of or during the convalescence from scarlet fever.

Scarlet Fever. An acute, contagious, febrile disease, having a period of incubation varying from several hours to a week, setting in

with vomiting or a chill, which is followed by high fever, rapid pulse, sore throat, and the appearance at the end of the first or the second day of the disease of a punctiform, scarlet-red eruption. The tongue, at first heavily coated and red at the tip and edges, soon shows prominence of the papillæ, which are red and swollen (strawberry-tongue). The eruption, at the appearance of which all the symptoms become intensified, gradually fades after five or six days, and is followed by a scaly desquamation. A peculiarity of scarlet fever is the tendency to involve the kidneys. **Malignant Scarlet Fever** is characterized by an abrupt onset, high fever, convulsions, coma, and death, usually before the appearance of the eruption.

Scarpa's Fascia [*Scarpa*, an Italian anatomist]. The deep layer of the superficial abdominal fascia.

Scarpa's Foramen. See *Foramina*, Table of.

Scarpa's Liquor [*Scarpa*, an Italian anatomist]. The endolymph.

Scarpa's Triangle. See *Triangle*.

Scatol (*skā'-tol*). See *Skatol*.

Scatula (*skat'-u-lah*) [L.]. An oblong, flat box for powders or pills.

Scavenger (*skav'-en-jer*) [AS., *scedwian*, to show]. One who cleans; a remover of waste and filth. **S.-cells**, wandering cells that take up debris. S.-cells are common in the nervous system.

Schacher's Ganglion. The ophthalmic ganglion.

Schede's Method. A method of treating caries of bone. The diseased tissue is scraped away and the cavity allowed to fill with a blood-clot. The latter is kept moist and aseptic by a covering of gauze and protective.

Scheiner's Experiment. An experiment illustrating refraction and accommodation of the eye. The person looks through two pinholes made in a card and placed at a less distance than the diameter of the pupil. If the eye is emmetropic, or if accurately focused, the two sets of rays, passing through the pinholes, unite and form a single image. In a myopic or a hyperopic eye the object appears double.

Schema (*skē'-mah*) [*σχῆμα*, form]. 1. A simple design to illustrate a complex mechanism. 2. An outline of a subject.

Schematic (*skē-mat'-ik*) [*σχῆμα*, form]. Pertaining to or of the nature of a schema.

S. Eye, one showing the proportions of a normal or typical eye.

Scherlievo (*skār-le-a'-vo*) [Ital.]. A form of ulcerative syphilis prevalent in the Austrian seaports during the last century.

Scheurlen's Bacillus (*shoir'-lēns*). A bacil-

lus at one time thought to be the cause of carcinoma.

Schindylesis (*skin-dil-e'-sis*) [*σχινδύλησις*, a cleavage]. A form of articulation in which a plate of one bone is received into a fissure of another bone.

Schisto- (*skis'-to-*) [*σχιστός*, cleft]. A prefix meaning split or fissured.

Schistocephalus (*skis-to-sef'-al-us*) [*σχιστός*, cleft; *κεφαλή*, head]. 1. Having a fissured skull. 2. A monster with a fissured skull.

Schistocyte (*skis'-to-sit*) [*σχιστός*, cleft; *κύτος*, cell]. A blood-corpuscle in the stage of dividing.

Schistoglossia (*skis-to-glos'-e-ah*) [*σχιστός*, cleft; *γλῶσσα*, tongue]. Cleft tongue.

Schistoprosopus (*skis-to-pros-o'-pus*) [*σχιστός*, cleft; *πρόσωπον*, face]. 1. Having a cleft or fissured face. 2. A monster having a fissure of the face.

Schistorrhachis (*skis-tor'-a-kis*) [*σχιστός*, cleft; *ράχης*, spine]. Spina bifida.

Schistosomus (*skis-to-so'-mus*) [*σχιστός*, cleft; *σῶμα*, a body]. A variety of monster in which there is a lateral or median eventration extending the whole length of the abdomen, the lower extremities being absent or rudimentary.

Schistothorax (*skis-to-tho'-raks*) [*σχιστός*, cleft; *θώραξ*, chest]. Fissure of the thorax.

Schizogenesis (*skiz-o-jen'-es-is*) [*σχίζειν*, to cleave; *γένεσις*, production]. Reproduction by fission.

Schizomycetes (*skiz-o-mi-se'-tes*) [*σχίζειν*, to cleave; *μύκης*, a fungus]. The cleft fungi or bacteria, so-called because multiplying by fission.

Schlemm, Canal of. A channel within the sclera close to the corneal juncture, whereby the aqueous humor finds its way into the general circulation.

Schlemm's Ligament. One of two ligaments connected with the shoulder-joints.

Schneiderian Membrane (*shni-de'-re-an*) [C. V. *Schneider*, a German anatomist]. The nasal mucous membrane.

Schönlein's Disease. See *Diseases*, Table of.

Schott's Method. A method of treating heart-disease by resisted exercise and special forms of baths.

Schrager's Lines (*shrah'-gers*). A coarse striation, concentric with the outline of the pulp-cavity, produced by the parallel curvings of dentinal tubules.

Schreiner's Base (*shri'-nurz*). See *Spermin*.

Schwann, Sheath of. The neurilemma of a nerve-fiber.

Schwann, White Substance of. The myelin of a medullated nerve-fiber.

Schweinerotlauf (*shwi'-na-rüt'-lowf*)

[Ger.]. Rouget du porc [Fr.]. Hog-erysipelas, an infectious disease of hogs, characterized by fever and an eruption of reddish or brownish spots, and due to a special bacillus. See *Bacillus erysipelatos suis*, *Bacteria*, *Table of*.

Schweinfurth Green (*shwīn'-foort*). Synonym of *Paris-green*.

Sciatic (*si-at'-ik*) [*ischion*, ischium]. 1. Pertaining to the ischium, as, e. g., the S. notch. 2. Pertaining to the sciatic nerve, as S. neuralgia.

Sciatica (*si-at'-ik-ah*) [*ischiatricus*, from *ischion*, ischium]. A disease characterized by neuralgic pain along the course of the sciatic nerve. It usually follows exposure to cold and wet, and is dependent upon inflammation of the nerve. In addition to pain there are numbness and tingling, tenderness along the course of the nerve, and eventually wasting of the muscles.

Scilla (*sīl'-ah*) [L.]. See *Squill*.

Scirrroid (*skīr'-oid*) [*σκίρρῶς*, hard; *είδος*, like]. Resembling a scirrhous.

Scirrhoma (*skīr-o'-mah*) [*σκίρρῶς*, hard; *δύμα*, tumor]. See *Scirrhous*.

Scirrhosarca (*skīr-o-sar'-kah*) [*σκίρρῶς*, hard; *σάρξ*, flesh]. Hardening of the flesh, especially of new-born infants; sclerema of infants.

Scirrhous (*skīr'-us*) [*σκίρρῶς*, hard]. Hard.

Scirrhus (*skīr'-us*) [*σκίρρῶς*, hard]. A hard carcinoma.

Scissors (*siz'-ors*) [*scindere*, to cut]. An instrument consisting of two blades held together by a rivet, and crossing each other so that in closing they cut the object placed between them.

Scissura (*siz-n'-rah*) [*scindere*, to cut]. A fissure.

Sclera (*skle'-rah*) [*σκληρός*, hard]. The sclerotic coat of the eye; the firm, fibrous, outer membrane of the eyeball, continuous with the sheath of the optic nerve behind and with the cornea in front.

Scleral (*skle'-ral*) [*σκληρός*, hard]. Pertaining to the sclera.

Scleratitis (*skle-rat'-it-is*). Same as *Scleritis*.

Sclerectasia (*skle-rek-ta'-se-ah*) [*σκληρός*, hard; *εκτασις*, extension]. Localized bulging of the sclera.

Sclerectomy (*skle-rek'-to-me*) [*σκληρός*, hard; *ἐκτομή*, excision]. Excision of a portion of the sclera.

Sclerema (*skle-re'-mah*) [*σκληρός*, hard]. A hardening. **S. adultorum**. See *Scleroderma*. **S. neonatorum**, a disease of the new-born, characterized by a hardening of the subcutaneous tissue, especially of the legs and feet, and probably dependent on a coagulation of the fat.

Scleriosis (*skle-ri'-as-is*) [*σκληρίσις*, hard]. Scleroderma.

Scleritis (*skle-ri'-tis*) [*σκληρίτις*, hard; *ιτις*, inflammation]. Inflammation of the sclerotic coat of the eye. It may exist alone (simple S. or episcleritis) or may be combined with inflammation of the cornea, iris, or choroid.

Sclero- (*skle'-ro-*) [*σκληρός*, hard]. 1. A prefix meaning hard. 2. A prefix denoting connection with the sclera.

Sclerоchoroiditis (*skle-ro-ko-roid-i'-tis*) [*σκληρόσις*, hard; *choroid*; *ιτις*, inflammation]. Inflammation of the choroid and the sclerotic coat of the eye.

Sclerоcorneal (*skle-ro-kor'-ne-al*) [*σκληρόσις*, hard; *corneus*, horny]. Pertaining conjointly to the sclerotic coat and the cornea of the eye.

Sclerodactylia, **Sclerodactyly** (*skle-ro-dak-tīl'-e-ah*, *skle-ro-dak'-tīl-e*) [*σκληρόσις*, hard; *δάκτυλος*, finger]. A disease of the fingers (or toes) allied to scleroderma. It is usually symmetric, occurs chiefly in women, and leads to marked deformity.

Scleroderma (*skle-ro-der'-mah*) [*σκληρόσις*, hard; *δέρμα*, skin]. A disease characterized by a progressive induration of the skin, occurring either in circumscribed patches (see *Morphea*) or diffusely. The skin becomes hard, pigmented, and firmly attached to the underlying tissues; destructive changes may also occur, and joints may become immobile from adhesions of the skin. The cause of S. is not known.

Sclerogenous (*skle-roj'-en-us*) [*σκληρόσις*, hard; *γεννάν*, to beget]. Producing a hard substance.

Scleroma (*skle-ro'-mah*) [*σκληρόσις*, hard]. Abnormal hardness or induration of a part.

S. adultorum. Synonym of *Scleroderma*.

Scleromucin (*skle-ro-mu'-sin*) [*σκληρόσις*, hard; *mucus*, mucus]. A gummy substance obtained from ergot, and considered one of its active principles.

Scleronyxis (*skle-ron-ik'-sis*) [*σκληρόσις*, hard; *νύξις*, a pricking]. Puncture of the sclera.

Sclerosarcoma (*skle-ro-sar-ko'-mah*) [*σκληρόσις*, hard; *σάρξ*, flesh; *δύμα*, tumor]. A hard, fleshy tumor of the gums.

Sclerosed (*skle'-rōsd*) [*σκληρόσις*, hard]. Affected with sclerosis; hardened.

Sclérose en plaques (*skla-ras' on(g)-plahk*). Synonym of *Sclerosis*, *Multiple*.

Sclerosis (*skle-ro'-sis*) [*σκληρόσις*, hard]. Hardening, especially a hardening of a part from an overgrowth of fibrous tissue; applied particularly to hardening of the nervous system from atrophy or degeneration of the nerve-elements and hyperplasia of the interstitial tissue; also to a chronic inflammation of the arteries characterized by thickening of their

coats. **S.**, Diffuse, one extending through a large part of the brain and cord. **S.**, Disseminated, a form in which numerous sclerotic patches are scattered through the brain and cord. **S.**, Insular. See *S.*, Multiple. **S.**, Lateral. See *Lateral Sclerosis*. **S.**, Multiple. See *Charcot's Disease*, in *Diseases*, Table of.

Scleroskeleton (*skle-ro-skell'-et-on*) [*σκληρόσ, hard; σκελετόν, a dry body*]. The part of the skeleton including the bones occurring in tendons and ligaments.

Sclerostenosis (*skle-ro-sten-o'-sis*) [*σκληρόσ, hard; στένωσις, constriction*]. 1. Sclerosis with stenosis. 2. *Scleroderma*.

Sclerotic (*skle-rot'-ik*) [*σκληρόσ, hard*]. Hard, indurated. **S.** Coat. See *Sclera*.

Sclerotica (*skle-rot'-ik-ah*) [*σκληρόσ, hard*]. See *Sclera*.

Scleroticochoroiditis (*skle-rot'-ik-o-ko-roid-i'-tis*). See *Sclerochoroiditis*.

Scleroticonyx (*skle-rot'-ik-on-ik'-sis*). See *Scleronyxis*.

Scleritis (*skle-ro-ti'-tis*) [*σκληρόσ, hard, sclera; ιτις, inflammation*]. See *Scleritis*.

Sclerotium (*skle-ro'-she-um*) [*σκληρόσ, hard*]. A thick mass of hyphae constituting a resting-stage in the development of some fungi, as the ergot.

Sclerotome (*skle'-ro-tōm*) [*σκληρόσ, hard; τέμνειν, to cut*]. 1. A knife used in sclerotomy. 2. A hard tissue separating successive myotomes in certain of the lower vertebrates.

Sclerotomy (*skle-rot'-o-me*) [*σκληρόσ, hard; τέμνειν, to cut*]. The operation of incising the sclera. **S.**, Anterior, the making of an incision through the sclera anterior to the ciliary body, and entering the anterior chamber, as is done in glaucoma. **S.**, Posterior, **S.** by an incision through the sclera behind the ciliary body, and entering the vitreous chamber.

Scolex (*sko'-leks*) [*σκόληξ, a worm*]. The head of a tapeworm, giving rise to the chain of proglottides.

Scoliorachitic (*sko-le-o-ra-kil'-ik*) [*σκολιός, curved; ραχίς, spine; ιτις, inflammation*]. Pertaining to or produced by scoliosis and rickets.

Scoliosis (*sko-le-o'-sis*) [*σκολιός, curved*]. A morbid lateral curvature of the spine.

Scoliotic (*sko-le-ot'-ik*) [*σκολιός, curved*]. Pertaining to or marked by scoliosis.

Scoop [*AS., skopa, a scoop*]. An instrument resembling a spoon, for the extraction of bodies from cavities, as an ear-S., lithotomy-S.

Scoparin (*sko'-par-in*). See *Scoparius*.

Scoparius (*sko-pa'-re-us*) [*scopa, a broom*]. The *Cytisus scoparius*, a shrub of the order Leguminosae. The tops (*S.*, *U. S. P.*, *Scoparii cacumina*, *B. P.*) contain the alkaloid

spartein, $C_{15}H_{24}N_2$, and a neutral principle, scoparin, $C_{11}H_{12}O_{10}$. **S.** is diuretic and cathartic, these actions probably depending upon scoparin. *Extractum scoparii fluidum* (*U. S. P.*). Dose mxx-xl (1.3-2.6). For properties of spartein see *Sparteina*.

Scopolamine, **Scopolein** (*sko-po'-lam-in, sko-po'-le-in*). See *Scopolia*.

Scopolia (*sko-po'-le-ah*) [after *Scopoli*, an Italian naturalist]. A genus of the Solanaceae, resembling the genera *Atropa* and *Hyoscyamus*. The rhizome of *S. japonica* and *S. carniolica* contains the alkaloid scopolamine or scopolein, $C_{17}H_{21}NO_4$, used as a mydriatic. **Scorbutic** (*skor-bu'-tik*) [*scorbutus, scurvy*]. Pertaining to, affected with, or caused by scorbutus.

Scorbutus (*skor-bu'-tus*) [*L.*]. See *Scurvy*.

Scotodinia (*sko-to-din'-e-ah*) [*σκότος, darkness; δίνοσ, a whirl*]. Vertigo associated with the appearance of black spots before the eyes.

Scotograph (*sko'-to-graf*) [*σκότος, darkness; γράφειν, to write*]. 1. An instrument for aiding the blind to write. 2. A name given to the picture produced by means of the so-called X-rays. See *X-rays*.

Scotoma (*sko-to'-mah*) [*σκότος, darkness*]. An area in the visual field, rays of light from which are not at all (**Absolute S.**) or imperfectly (**Relative S.**) perceived. **S.**, Central, one limited to the region of the visual field corresponding to the macula lutea. **S.**, Color-, color-blindness limited to a part of the visual field, and which may exist without interruption of the field for white light. **S.**, Flittering, a **S.** with serrated margins extending peripherally and producing a large defect in the visual field. It is also called fortification-spectrum. **S.**, Negative, a defect due to the destruction of the retinal center, and which is not noticeable to the patient. **S.**, Positive, a **S.** perceptible to the patient as a dark spot before his eyes. **S.**, Relative, a scotoma within which perception of light is only partially impaired. **S.**, Ring-, **S.**, Annular, a zone of scotoma surrounding the center of the visual field. **S.** scintillans, **S.**, Scintillating. See *S.*, Flittering.

Screatus (*skre-a'-tus*) [*L.*]. 1. A hawking. 2. A neurosis characterized by paroxysms of hawking.

Scrivener's Palsy. See *Writer's cramp*.

Scrobiculus (*skro-bik'-u-lus*) [*L.*]. A small pit. **S. cordis**, the depression at the epigastrium; the pit of the stomach.

Scrofula (*skrof'-u-lah*) [*scrofa, a sow*]. A term formerly applied to a peculiar condition characterized by enlargement of the lymphatic glands and necrosis of the bones; it is at present considered a form of tuberculosis.

Scrofulide (*skrof'-u-lid*) [*scrofa*, sow]. See *Scrofuloderm*.

Scrofuloderm (*skrof'-u-lo-derm*) [*scrofa*, sow; *δέρμα*, the skin]. A disease of the skin due to scrofula, and generally characterized by superficial irregular ulcers with undermined edges. The cause is the tubercle bacillus.

Scrofulosis (*skrof-u-lo'-sis*) [*scrofa*, sow]. The state characterized by the presence of scrofula; a scrofulous diathesis.

Scrofulous (*skrof'-u-lus*) [*scrofa*, sow]. Having the nature of, affected with, or produced by scrofula.

Scrotal (*skro'-tal*) [*scrotum*, scrotum]. Pertaining to, or contained in the scrotum, as *S. hernia*.

Scrotum (*skro'-tum*) [L.]. The pouch containing the testicles, consisting of skin, dartos, spermatic fascia, cremasteric fascia, infundibuliform fascia, and parietal tunica vaginalis.

Scruple (*skru'-pl*) [*scrupulus*, dim. of *scrupus*, a sharp stone]. In apothecaries' weight, 20 grains; represented by the sign \mathfrak{D} .

Scurf (*skerf*) [AS., *scurf*, scurf]. A bran-like desquamation of the epidermis, especially from the scalp; dandruff.

Scurvy (*sker'-ve*) [AS., *scurf*, scurf]. A disease observed among persons who have been deprived of proper food for a length of time; it is characterized by spongy gums, extravasations of blood beneath the skin, hemorrhages from the mucous membranes, fetor of the breath, and painful contractions of the muscles. It is most common among sailors living on salt-meats. S., Land-. See *Purpura hemorrhagica*.

Scurvy-grass. The *Cochlearia officinalis*, a plant of the order Crucifere, the properties of which reside in a volatile oil resembling oil of mustard. S. is used in scurvy and in chronic rheumatism.

Scute (*sküt*) [*scutum*, a shield]. See *Scutum*.

Scutellaria (*sku-tel'-a'-re-ah*) [*scutellum*, a little shield]. A genus of the Labiatae. *S. lateriflora*, skullcap, is the S. of the U. S. P., and is employed in neuralgia, chorea, delirium tremens, and other nervous affections. *Extractum scutellariae fluidum* (U. S. P.). Dose f3j-ij (4.0-8.0). Dose of *Scutellarin*, an impure precipitate from an alcoholic tincture, gr. iij-iv (0.20-0.26).

Scutulum (*sku'-tu-lum*) [dim. of *scutum*, a shield]. Any one of the thin plates of the eruption of favus.

Scutum (*sku'-tum*) [*scutum*, a shield]. A shield-like plate of bone. *S. tympanicum*, the semilunar plate of bone separating the attic of the tympanum from the outer mastoid cells.

Scybala (*sib'-al-ah*). Plural of *Scybalum*, q. v.

Scybalous (*sib'-al-us*) [*σύνβαλον*, fecal matter]. Of the nature of a scybalum.

Scybalum (*sib'-al-um*) [*σύνβαλον*, fecal matter]. A mass of abnormally hard fecal matter.

Seamstress's Cramp. A painful cramp affecting the fingers of seamstresses; an occupation-neurosis analogous to writer's cramp.

Searcher (*serch'-er*) [*circare*, to go around]. A sound used for the detection of stones in the bladder.

Sea-sickness. A condition occurring in persons aboard ships, produced by the rolling of the ship, and characterized by vertigo, nausea, retching, and prostration. A similar state may be induced by riding in cars, elevators, etc.

Sea-tangle. See *Laminaria*.

Seat-worm. See *Oxyuris*.

Sebaceous (*se-ba'-shus*) [*sebum*, suct, fat].

Pertaining to sebum; secreting sebum, as the S. Glands or Follicles, compound acellular glands associated with the hair-follicles, and secreting a semifluid substance, the sebum, composed of oil-droplets and broken-down epithelial cells. *S. Cyst*, a cystic tumor formed by occlusion of the duct of a S. gland, with retention of the secretion, dilatation and thickening of the wall of the gland. It contains a grayish-white, cheesy material.

Sebadilla (*seb-ad-il'-ah*). See *Sabadilla*.

Sebiparous (*seb-ip'-ar-us*) [*sebum*, fat; *παρρε*, to produce]. Secreting sebum.

Sebolith (*seb'-o-lith*) [*sebum*, fat; *λίθος*, stone]. A concretion in a sebaceous gland.

Seborrhea (*seb-or'-e'-ah*) [*sebum*, suct; *ῥοία*, a flow]. A functional disease of the sebaceous glands, characterized by an excessive secretion of sebum, which collects upon the skin in the form of an oily coating or of crusts or scales. *S. congestiva*, *Lupus erythematosus*. *S. capillitii*, *S. capitis*, S. of the scalp. *S. corporis*, S. of the trunk. *S. faciei*, S. of the face. *S. nigra*, *S. nigricans*, seborrhea with the formation of dark-colored crusts, the coloration being usually from dirt. *S. oleosa*, a form characterized by an excessive oiliness of the skin, especially about the forehead and nose. *S. sicca*, the commonest form of S., characterized by greasy, brownish-gray scales.

Sebum (*se'-bum*) [L.]. The secretion of the sebaceous glands. See *Sebaceous Glands*.

Secale (*se-ka'-le*) [L.]. Rye. *S. cornutum*. See *Ergot*.

Secondary (*sek'-un-da-re*) [*secundarius*, from *secundus*, second]. 1. Second in the order of time or development, as the S. lesions of syphilis. 2. Second in relation; subordinate; produced by a cause considered primary. *S. Amputation*, an amputation done after

the subsidence of inflammatory symptoms. **S. Cataract.** See *Cataract*. **S. Coil,** the coil of wire in which the induced current is generated. **S. Degeneration,** of nerve fibers, a degeneration following injury or disease of the trophic centers. **S. Hemorrhage.** See *Hemorrhage*.

Second Intention. See *Healing*.

Second Nerve. The optic nerve.

Secrete (*se-kre't*) [*secernere*, to separate]. To separate; specifically, to separate from the blood, or form out of materials furnished by the blood a certain substance termed a secretion.

Secretion (*se-kre'-shun*) [*secernere*, to secrete]. 1. The act of secreting or forming from materials furnished by the blood a certain substance which is either eliminated from the body or is used in carrying on special functions. 2. The substance secreted. **S., External,** a S. thrown out upon the external or internal surface of the body. **S., Internal,** a S. that is not thrown out upon a surface, but is absorbed into the blood.

Secretory (*se'-kre-to-re*) [*secernere*, to separate]. Pertaining to secretion; performing secretion.

Sectio (*sek'-she-o*) [L.]. See *Section*. **S. abdominis.** See *Celiotomy*. **S. agrippina,** Cesarean section. **S. alta,** suprapubic cystotomy. **S. cadaveris,** an autopsy. **S. cæsarea,** Cesarean section. **S. franconiana,** suprapubic cystotomy. **S. lateralis,** lateral lithotomy. **S. mediana,** median lithotomy.

Section (*sek'-shun*) [*secare*, to cut]. 1. The act of cutting or dividing. 2. A cut; a cut surface. **S., Abdominal.** See *Celiotomy*. **S., Cesarean.** See *Cesarean Operation*. **S., Frontal,** a S. dividing the body into dorsal and ventral parts. **S., Sagittal,** a S. parallel with the sagittal suture, and hence with the median plane of the body, and serving to divide the body into equal parts.

Secundines (*sek'-un-denz*) [*secundus*, second]. The placenta, part of the umbilicus, and the membranes discharged from the uterus after the birth of the child.

Secundipara (*se-kun-dip'-ar-ah*) [*secundus*, second; *parere*, to bring forth]. See *Multipara*.

Sedation (*se-da'-shun*) [*sedare*, to soothe]. 1. A state of lessened functional activity. 2. The production of a state of lessened functional activity.

Sedative (*sed'-at-iv*) [*sedare*, to soothe]. 1. Quieting or lessening functional activity. 2. An agent lessening functional activity.

Sediment (*sed'-im-ent*) [*sedimentum*, from *sedere*, to sit]. The material settling to the bottom of a liquid.

Sedimentation (*sed-im-en-ta'-shun*) [*sedimentum*, sediment, from *sedere*, to sit]. The

process of producing the deposition of a sediment, especially the rapid deposition by means of a centrifugal machine.

Segment (*seg'-ment*) [*segmentum*, from *secare*, to cut]. A small piece cut from the periphery of anything; a part bounded by a natural or imaginary line.

Segmental (*seg-men'-tal*) [*segmentum*, a segment, from *secare*, to cut]. 1. Pertaining to a segment; made up of segments. 2. Undergoing or resulting from segmentation. **S. Duct,** the duct of the pronephros. **S. Organs,** a tubular structure found in the embryos of amniotic animals, and comprising the pronephros, the mesonephros, and the metanephros.

Segmentation (*seg-men-ta'-shun*) [*segmentum*, a segment, from *secare*, to cut]. The process of dividing into two equal parts, as, *e.g.*, the S. of the ovum. **S.-cavity,** the central space in the ovum produced by S. **S.-cells.** See *S.-sphere*. **S.-nucleus.** See *Nucleus*. **S.-sphere,** one of the cells of an ovum formed by S.

Seidlitz-powder (*sid'-litz*). Pulvis effervescens compositus. See *Potassium*.

Selection (*se-lek'-shun*) [*seligere*, to choose]. The act of choosing. **S., Natural,** the selective action of external conditions, whereby characters favorable to the species of animal or plant are preserved. **S., Sexual,** the selection produced by preferences of the one sex for a member of the other sex in some way specially endowed.

Self-abuse, Self-pollution. See *Masturbation*.

Self-limited. Limited by reason of inherent qualities; applied to diseases that run a definite limited course independent of treatment.

Sella (*sel'-ah*) [L.]. A saddle. **S. turcica** [Turkish saddle], the pituitary fossa of the sphenoid bone, lodging the pituitary body.

Selters, Seltzer Water (*sel'-ters, sellz'-er*). An effervescent mineral water obtained at Selters in Prussia.

Semeiography (*sem-i-og'-ra-fe*) [*σημειον*, sign; *γράφειν*, to write]. Symptomatology.

Semeiology (*sem-i-ol'-o-je*) [*σημειον*, sign; *λόγος*, discourse]. Symptomatology.

Semeiotic (*sem-i-ol'-ik*) [*σημειον*, sign]. Pertaining to symptoms.

Semeiotics (*sem-i-ol'-iks*) [*σημειον*, sign]. Symptomatology.

Semelincident (*sem-el-in'-sid-ent*) [*semel*, once; *incidere*, to happen]. Happening only once in the same person; as a S. disease.

Semen (*se'-men*) [*serere*, to sow]. 1. A seed. 2. The fecundating fluid of the male, chiefly secreted by the testicles, composed of liquor seminis, seminal granules, oil-globules, and spermatozoa.

Semi- (*sem'-e-*) [L.]. A prefix denoting half. **Semicircular** (*sem-e-sir'-ku-lar*) [*semi*, half; *circulus*, a circle]. Having the form of a half-circle. **S. Canals.** See *Ear*.

Semilunar (*sem-e-lu'-nar*) [*semi*, half; *luna*, moon]. Resembling a half-moon in shape, as, e. g., the S. bone of the carpus, the S. cartilage of the knee, the S. ganglion of the abdominal sympathetic nerve or of the trifacial nerve (Gasserian ganglion), the S. valves of the heart. **S. Space of Traube**, the tympanic area at the lower part of the left chest corresponding to the stomach.

Semimembranous (*sem-e-mem'-bra-nus*) [*semi*, half; *membranosus*, like a membrane]. Partly membranous, as, e. g., the S. muscle (**Semimembranosus**). See *Muscles*, *Table of*.

Seminal (*sem'-in-al*) [*semen*, seed]. Pertaining to the semen. **S. Cyst**, a cyst of the spermatic cord or testicle containing semen. **S. Vesicles.** See *Vesicles*.

Seminiferous (*sem-in-iff'-er-us*) [*semen*, seed; *ferre*, to carry]. Producing semen, as the S. tubules of the testicle.

Seminormal (*sem-e-nor'-mal*) [*semi*, half; *norma*, rule]. Half-normal. **S. Solution**, one containing in solution half the quantity of the substance contained in the normal solution.

Semis (*se'-mis*) [L.]. Half; abbreviated in prescription to ss., which is placed after the sign indicating the measure.

Semispinalis (*sem-e-spi-na'-lis*). See *Muscles*, *Table of*.

Semisulcus (*sem-e-sul'-kus*) [*semi*, half; *sulcus*, a sulcus]. A half-sulcus which uniting with another sulcus forms a complete sulcus.

Semitendinous (*sem-e-ten'-din-us*) [*semi*, half; *tendo*, a tendon]. Partly tendinous, as, e. g., a S. muscle (**Semitendinosus**). See *Muscles*, *Table of*.

Senega (*sen'-e-ga*) [L.]. The *Polygala senega*, a plant of the order *Polygalæ*. Its root (S., U. S. P., *Senegæ radix*, B. P.) contains a bitter principle, senegin or polygalic acid (or polygalin), which is probably identical with saponin. S. is used as a stimulant, expectorant, and diuretic; in large doses it is emetocathartic. It is chiefly employed in bronchitis and laryngitis, as a diuretic in dropsy, and in amenorrhea. Dose gr. x-xx (0.65-1.3). *Extractum senegæ fluidum* (U. S. P.). Dose mxx-xx (0.65-1.3). *Infusum senegæ* (B. P.). Dose f 3 j (32.0). *Syrupus senegæ* (U. S. P.). Dose f 3 j-ij (4.0-8.0). *Tinctura senegæ* (B. P.). Dose f 3 j (4.0). Polygalic acid is employed in doses of from gr. 1/4-j (0.016-0.065).

Senegin (*sen'-e-gin*). See *Senega*.

Senile (*se'-nil*) [*senilis*, resembling the state

produced by old age; from *senex*, old]. Pertaining to or caused by old age.

Senility (*sen-il'-it-e*) [*senilis*, from *senex*, old]. The state of being senile; the weakness characteristic of old age.

Senn's Bone-plates [after Nicholas Senn, an American surgeon]. Plates of decalcified bone used in intestinal anastomosis.

Senn's Test. The introduction of hydrogen-gas into the bowel through the rectum, for the detection and localization of an abnormal opening.

Senna (*sen'-ah*) [Arab., *senā*]. The leaflets of various species of *Cassia*, a genus of the order *Leguminosæ*. S. of the U. S. P. is derived from *Cassia acutifolia*. S. of the B. P. is of two varieties—**Alexandrian S.**, from *Cassia acutifolia*, and **East India or Tinnevely S.**, from *Cassia angustifolia*. S. contains cathartic acid, a glucosid representing the purgative properties of S., the bitter principles, sennapierin and sennacrol, and a coloring matter, chrysophan. S. is used as a purgative, generally in combination with an aromatic to prevent griping. Dose 3 ss-ij (2.0-8.0). Preparations and doses: *Confectio sennæ* (U. S. P., B. P.). Dose 3 ij (8.0). *Extractum sennæ fluidum* (U. S. P.). Dose f 3 j-iv (4.0-16.0). *Infusum sennæ* (B. P.). Dose f 3 iv (128.0). *Infusum sennæ compositum* (U. S. P.), black draught, contains S., manna, and magnesium sulphate. Dose f 3 iv (128.0). *Pulvis glycyrrhizæ compositus* (U. S. P., B. P.), compound liquorice powder. Dose gr. xxx-lx (2.0-4.0). *Syrupus sennæ* (U. S. P., B. P.). Dose f 3 j-iv (4.0-16.0). *Tinctura sennæ* (B. P.). Dose f 3 j-iv (4.0-16.0).

Sennacrol (*sen'-ak-rol*) [Arab., *senā*, senna; *acris*, sharp]. See *Senna*.

Sennapierin (*sen-ap-ik'-rin*) [Arab., *senā*, senna; *πικρός*, bitter]. See *Senna*.

Sensation (*sen-sa'-shun*) [*sentatio*, from *sentire*, to feel]. A feeling or impression produced by the stimulation of an afferent nerve.

Sense (*sens*) [*sensus*, from *sentire*, to feel]. 1. Any one of the faculties by which stimuli from the external world or from within the body are received and transformed into sensations. The faculties receiving impulses from the external world are the senses of sight, hearing, touch, smell, and taste, which are the special senses, and the muscular and temperature-sense. Those receiving impulses from the internal organs (visceral senses) are the hunger-S., thirst-S., and others. 2. A sensation.

Sensibility (*sen-sib-il'-it-e*) [*sensibilitas*, from *sentire*, to feel]. The ability to receive and feel impressions; of a nerve or end-organ, to receive and transmit impulses.

Sensible (*sen'-sibl*) [*sensibilis*, from *sentire*, to feel.] Perceptible by the senses, as, *e. g.*, S. perspiration; capable of receiving an impression through the senses.

Sensitive (*sen'-sit-iv*) [*sensitivus*]. 1. Capable of feeling; capable of transmitting sensation. 2. Reacting to a stimulus.

Sensorial (*sen-so'-re-al*) [*sensorium*, the organ of sensation]. Pertaining to the sensorium.

Sensorimotor (*sen-so-re-mo'-tor*) [*sensus*, feeling; *motor*]. Both sensory and motor; concerned with the perception of sensory impulses and with motor impulses. **S. Centers**, centers that are concerned both with the perception of sensation and with motor impulses.

Sensorium (*sen-so'-re-um*) [L.]. A center for sensations, especially the part of the brain concerned in receiving and combining the impressions conveyed to the individual sensory centers.

Sensory (*sen'-so-re*) [*sentire*, to feel]. Pertaining to or conveying sensation. **S.**

Aphasia. See *Aphasia*. **S. Aura**, an aura affecting the special senses. **S. Crossway**, the posterior third of the posterior limb of the internal capsule, where the afferent fibers conveying sensory impulses cross to the opposite side. **S. Epilepsy**, various disturbances of sensation occurring in paroxysms that replace the epileptic convulsion. **S. Nerve**, one that conveys sensations from the periphery to the centers.

Sentient (*sen'-she-ent*) [*sentire*, to feel]. Capable of feeling.

Separator (*sep'-ar-a-tor*) [*separare*, to separate]. 1. Anything that separates, especially an instrument for separating the teeth. 2. An instrument for detaching the pericranium or periosteum.

Sepsin (*sep'-sin*) [*σῆψιν*, to make rotten]. A poisonous ptomain obtained from decomposed yeast and blood. See *Ptomain*, *Table of*.

Sepsis (*sep'-sis*) [*σῆψις*]. A state of poisoning produced by the absorption of putrefactive substances. **S.**, **Puerperal**, sepsis occurring after childbirth, from absorption of putrefactive products from the parturient canal.

Septan (*sep'-tan*) [*septem*, seven]. Recurring every seventh day, as, *e. g.*, S. fever.

Septemia (*sep-te'-me-ah*). See *Septicemia*.

Septentrionalin (*sep-ten-tre-o-nal'-in*). An alkaloid obtained from *Aconitum lycoctonum*. It is a sensory paralyzant, and has been suggested as a local and general anesthetic.

Septic (*sep'-tik*) [*σηπτικός*, putrid]. Pertaining to or produced by putrefaction. **S. Intoxication**, a form of poisoning resulting from the absorption of products of putrefaction.

Septicemia (*sep-te-se'-me-ah*) [*σηπτικός*, putrid; *αἷμα*, blood]. Blood-poisoning; a form of poisoning resulting from the presence in the blood of the products of pyogenic or putrefactive microorganisms. **S.**, **Mouse-**, a form of S. occurring in mice and produced by the bacillus murisepticus. It is usually fatal in from 40 to 60 hours, the animal becoming early apathetic. **S.**, **Rabbit-**, a form of S. occurring in rabbits and due to a special bacillus, the bacillus septicemiac hamorrhagicae, or bacillus of chicken-cholera. **S.**, **Sputum-**, a form of S. produced by inoculation with microorganisms found in sputum, especially the pneumococcus.

Septicemic (*sep-te-sem'-ik*, *sep-te-se'-mik*) [*σηπτικός*, putrid; *αἷμα*, blood]. Pertaining to or affected with septicemia.

Septicin (*sep'-tis-in*) [*σηπτικός*, putrid]. A ptomain obtained from decaying flesh.

Septicopyemia (*sep-tik-o-pi-e'-me-ah*) [*σηπτικός*, putrid; *πύον*, pus; *αἷμα*, blood]. Combined septicemia and pyemia. **S.**, **Primary**, that in which the general infection is produced by the same bacteria as those causing the primary lesion, and **S.**, **Secondary**, that in which the general infection is due to other bacteria than those causing the primary lesion.

Septimipara (*sep-tim-ip'-ar-ah*) [*septimus*, seventh; *parere*, to bear]. See *Multipara*.

Septivalent (*sep-tiv'-al-ent*) [*septem*, seven; *valere*, to be worth]. Having an atomicity of seven.

Septometer (*sep-tom'-et-er*) [*septum*, septum, 1st def.; *σηπτός*, putrid, 2d def.; *μέτρον*, measure]. 1. An instrument for determining the thickness of the nasal septum. 2. An apparatus for determining organic impurities in the air.

Septum (*sep'-tum*) [*sepire*, to hem in]. A partition; a division-wall. **S. atriorum**, **S. auricularum**, the S. between the right and left auricles of the heart. **S. crurale**, the layer of areolar tissue closing the femoral ring. **S. lucidum**, a thin translucent septum forming the internal boundary of the lateral ventricles of the brain and enclosing between its two lamina the fifth ventricle. **S.**, **Nasal**, the S. between the two nasal cavities. **S.**, **Pectiniform**, that between the corpora cavernosa of the penis. **S.**, **Recto-vaginal**, the tissue forming the partition between the rectum and the vagina. **S. ventriculorum**, the S. between the two ventricles of the heart.

Septuplet (*sep'-tu-plet*) [*septem*, seven]. One of seven offspring born from a single gestation.

Sequela (*se-kwe'-lah*) [*sequi*, to follow]. A diseased or abnormal condition following an attack of a disease, and directly or indirectly dependent upon it.

Sequestration (*se-kwes-tra'-shun*) [*sequestrare*, to separate]. 1. The formation of a sequestrum. 2. The isolation of persons suffering from disease for purposes of treatment or of protecting others.

Sequestrectomy (*se-kwes-trek'-to-me*). See *Sequestrotomy*.

Sequestrotomy (*se-kwes-trot'-o-me*) [*sequestrum*, sequestrum; *tomē*, a cutting]. The operation of removing a sequestrum.

Sequestrum (*se-kwes'-trum*) [*sequestrare*, to separate]. A dead piece of bone that has become separated from the living bone.

Seralbumin (*ser-al-bu'-min*) [*serum*, serum; *albumin*]. Serum-albumin, the albumin found in the blood.

Serial (*se'-re-al*) [*series*, a succession]. Following in regular order; occurring in rows. **S. Sections**, microscopic sections made in consecutive order and arranged in the same manner.

Sericeps (*ser'-is-eps*) [*onpukē*, silken; *forceps*, forceps]. A device made of loops of ribbon, used in place of the forceps in making traction upon the fetal head.

Sero- (*se'-ro-*) [*serum*, serum]. A prefix denoting serous.

Serocystic (*se-ro-sis'-tik*) [*serum*, serum; *kyōstis*, bladder]. Composed of cysts filled with a serous fluid.

Serofibrinous (*se-ro-fi'-brin-us*) [*serum*, serum; *fibrin*]. Composed of serum and fibrin, as, *e. g.*, a S. exudate; characterized by the production of a S. exudate, as, *e. g.*, a S. inflammation.

Serolin (*se'-ro-lin*) [*serum*, serum; *oleum*, oil]. A neutral fatty constituent of blood, occurring in small amount; its nature is undetermined.

Seropurulent (*se-ro-pu'-ru-lent*) [*serum*, serum; *pus*, pus]. Composed of serum and pus, as, *e. g.*, a S. exudate.

Seropus (*se'-ro-pus*) [*serum*, serum; *pus*, pus]. A fluid consisting of serum and pus.

Serosa (*se-ro'-sah*) [*serosus*, serous; *membrana*, understood]. A serous membrane.

Seroserous (*se-ro-se'-rus*) [*serum*, serum]. Pertaining jointly to two serous surfaces.

Serosynovitis (*se-ro-si-no-vi'-tis*) [*serum*, serum; *synovitis*]. A synovitis accompanied by an increase of the synovial fluid.

Serotherapy (*se-ro-ther'-ap-e*) [*serum*, serum; *θεραπεῖα*, therapy]. The treatment of disease by means of the blood-serum of animals or individuals that are immune to or convalescent from an infectious disease.

Serotina (*ser-o-ti'-nah*) [*serotinus*, late]. See *Decidua serotina*.

Serous (*se'-rus*) [*serum*, serum]. 1. Pertaining to, characterized by, or resembling serum. 2. Producing serum, as a S. gland; containing serum, as a S. cyst. **S. Effu-**

sion, an effusion of serum. **S. Exudate**, an exudate consisting largely of serum. **S. Inflammation**, an inflammation characterized by the formation of a S. exudate. **S. Membrane**. See *Membrane*.

Serpens (*ser'-pens*) [L.]. Creeping.

Serpentaria (*ser-pen-ta'-re-ah*) [L.]. Virginia snake-root, the root of several species of *Aristolochia*, of the order of *Aristolochiaceae*. The rhizoma and rootlets of *Aristolochia serpentaria* and *Aristolochia reticulata* constitute the S. of the U. S. P. (*Serpentariae rhizoma*, B. P.). S. contains a volatile oil, a bitter principle, and a nitrogenous principle called aristolochin. It is a stimulant, tonic, diaphoretic, and diuretic, and is used in the eruptive fevers to bring out the eruption, in intermittent fever, and in dyspepsia. **Extractum serpentariae fluidum** (U. S. P.). Dose mxx-xxx (1.3-2.0). **Infusum serpentariae** (B. P.). Dose fʒj (32.0). **Tinctura serpentariae** (U. S. P., B. P.). Dose fʒj (4.0). **Tinctura cinchonae composita** (U. S. P., B. P.). Dose fʒj (4.0). **Serpentine** (*ser'-pen-tin*) [*serpens*, serpent]. Sinuous; snake-like.

Serpiginous (*ser-pi'-in-us*) [*serpiginosus*, from *serpere*, to creep]. Creeping. **S. Ulcer**, one that extends in one direction while healing in another.

Serpigo (*ser-pi'-go*) [L.]. Ringworm.

Serrate, **Serrated** (*ser'-at*, *ser'-a-ted*) [*serra*, a saw]. Provided with sharp projections like the teeth of a saw.

Serration (*ser'-a-shun*) [*serra*, a saw]. The state or condition of being serrate.

Serratus (*ser'-a-tus*) [L.]. Serrated; applied to muscles arising or inserted by a series of processes resembling the teeth of a saw. See *Muscles*, Table of.

Serre-fine (*sar-fen'*) [Fr.]. A small spring-forceps for seizing and compressing bleeding vessels.

Serre-nœud (*sar-noe*) [Fr.]. An instrument used for drawing tight a ligature thrown around a part, as around the pedicle of a tumor.

Serrulate (*ser'-u-lat*) [*serrula*, dim. of *serra*, a saw]. Minutely notched or serrated.

Serum (*se'-rum*) [L.]. 1. The clear, yellowish fluid separating from the blood after the coagulation of the fibrin. 2. Any clear fluid resembling the S. of the blood. **S.-albumin**, the albumin found in the blood-serum and other animal fluids. **S.-globulin**. See *Paraglobulin*. **S. lactis**, whey. **S.-unit**. See *Unit*.

Sesame (*ses'-am-e*). See *Sesamum*.

Sesamoid (*ses'-am-oid*) [*σῆσαμον*, sesame; *εἶδος*, like]. Resembling a sesame-seed. **S. Bone**, a small bone developed in a tendon subjected to much pressure.

Sesamum (*ses'-am-um*) [*σάμινον*]. A genus of plants of the order Pedaliaceæ. **S.** indicum and **S.** orientale yield a bland, sweetish oil, sesame-oil, teel-oil, benne-oil (*Oleum sesami*, U. S. P.), employed like olive-oil.

Sesqui- (*ses'-kwe-*) [L.]. A prefix denoting one and one-half.

Sesquioxid (*ses'-kwe-oks'-id*) [*sesqui-*, one and one-half; *ὀξύς*, acid]. A compound of oxygen and another element, containing three parts of oxygen to two of the other element.

Sesquisalt (*ses'-kwe-savult*) [*sesqui*, one and one-half; *salt*]. A salt containing one and one-half times as much of the acid as of the radicle or base.

Sessile (*ses'-il*) [*sessilis*, from *sedere*, to sit]. Attached by a broad base; not pedunculated; as, *e. g.*, a S. tumor.

Seton (*se'-ton*) [*seta*, a bristle]. 1. A thread or skein of threads drawn through a fold of the skin, so as to produce a fistulous tract; it is used as a counterirritant. 2. The tract thus produced.

Setschenow's Inhibitory Center (*setch'-en-ofs.*). A cerebral center for the inhibition of reflex movements, situated in the corpora quadrigemina and the medulla oblongata.

Seven-day Fever. Relapsing fever.

Sevum (*se'-vum*) [L.]. Suet.

Sewer-gas. The mixture of gases emanating from sewers.

Sewing Spasm. See *Seamstress's Cramp*.

Sextan (*seks'-tan*) [*sex*, six]. Occurring every sixth day, as, *e. g.*, a S. fever.

Sextipara (*seks'-tip'-ar-ah*) [*sex*, six; *parere*, to bear]. See *Multipara*.

Sextuplet (*seks'-tūp'-let*) [*sex*, six]. One of six offspring of a single gestation.

Sexual (*seks'-u-al*) [*sexus*, sex]. Pertaining to or characteristic of sex, as the S. organs.

Sexvalent (*seks'-val-ent*) [*sex*, six; *valere*, to be worth]. Having an atomicity of six as compared with that of hydrogen.

Shadowgram. See *X-rays*.

Shadow-test. See *Skiascopy*.

Shakes (*shaks*). See *Ague*.

Shaking Palsy. See *Paralysis agitans*.

Sharpey's Perforating Fibers. Transverse or perpendicular fibers transfixing and joining the lamellæ of bone.

Shaven-beard Appearance. A peculiar appearance of the agminated glands of the intestine in typhoid fever, resembling that of a recently-shaven beard.

Sheath (*shēth*) [AS., *scodh*, sheath]. A covering. **S.**, Primitive, **S.** of Schwann. See *Neurilemma*.

Sheep-pox. A contagious pustular disease of sheep, similar to cow-pox.

Sherry-wine (*sher'-e*). See *Vinum xericum*.

Shin [AS., *scina*, shin]. The sharp anterior margin of the tibia. **S.-bone**, the tibia.

Shingles (*shing'-gles*). Herpes zoster.

Ship-fever. Typhus fever.

Shiver (*shiv'-er*) [ME., *chiveren*, to shiver]. A slight tremor or shaking of the body due to cold, etc.

Shock [Fr., *shoc*, shock]. 1. A sudden grave depression of the system produced by operations, accidents, or strong emotion. It is due to a profound influence on the nervous system. If not fatal it is followed by a stage of reaction. 2. The agent causing a general or local depression, as, *e. g.*, an electric S.

Shoddy Fever. A diseased condition caused by the inhalation of the dust in shoddy-factories; it is characterized by feverishness, headache, nausea, dryness of the mouth, dyspnea, cough, and expectoration.

Shoemaker's Spasm. An occupation-neurosis, analogous to writer's cramp, occurring in shoemakers.

Short Circuit. One in which an electric current encounters an abnormally small resistance.

Short-sight. Myopia.

Shoulder (*shōl'-der*) [AS., *sculder*, shoulder].

The region where the arm joins the trunk, formed by the meeting of the clavicle and the scapula, and the overlying soft parts. **S.-blade**, the scapula. **S.-girdle**. See *Girdle*.

Show (*sho*) [AS., *scawian*, to look]. 1. A bloody discharge from the birth-canal prior to labor. 2. The first appearance of a menstrual flow.

Shower-bath. See *Bath*.

Shrapnell's Membrane. See *Membrane*.

Si. A symbol for silicon.

Sialagogue (*si-al'-a-gog*) [*σίαλον*, spittle; *ἀγωγός*, leading]. 1. Producing a flow of saliva. 2. A drug producing a flow of saliva.

Sialorrhea (*si-al-or'-e-a*) [*σίαλον*, spittle; *ῥοία*, a flow]. Salivation. **S.**, **Pancreatic**, a flow of pancreatic juice.

Sibbens (*sib'-ens*) [Gael., *subhan*, raspberries]. A disease formerly endemic in the Scotch highlands, and by some identified with syphilis, by others with yaws.

Sibilant (*sib'-il-ant*) [*sibilare*, to hiss]. Hissing or whistling, as, *e. g.*, a S. rale.

Sibilus (*sib'-il-us*) [*sibilare*, to hiss]. A sibilant rale.

Sick [AS., *seōc*, sick]. Ill; not well. **S.-headache**, migraine.

Sickness (*sik'-nes*) [AS., *seōc*, sick]. 1. The state of being unwell. 2. Nausea. **S.**, **African Sleeping**. See *African Lethargy*.

S., **Falling**, epilepsy. **S.**, **Green**, chlorosis. **S.**, **Monthly**, the menstrual epoch.

S., **Mountain-**, a sensation of nausea, with impeded respiration and irregular heart's

action, due to the rarefied air of high altitudes.

Siderosis (*sid-er-o'-sis*) [*σιδηρος*, iron]. A pigmentation by a deposit of particles of iron; specifically, a chronic interstitial pneumonia caused by the inhalation of particles of iron.

Siegle's Otoloscope or Speculum. An instrument consisting of a glass-covered box with a conical projection and a rubber tube attached laterally. When the conical projection is inserted firmly into the external auditory canal, and the air is compressed or rarefied, the movements of the drum-membrane may be observed. It is also used for the purpose of rendering the articulations of the ossicles mobile.

Sieve (*siv*) [AS., *sife*, sieve]. An apparatus with a reticulated bottom, used for the separation of fine from coarse particles. The gauge of the sieve is usually expressed in the number of meshes per square inch.

Sigaultian Operation, Sigault's Operation [J. R. *Sigault*, a French obstetrician]. Symphysiotomy.

Sigh (*si*) [AS., *scan*, to sigh]. A prolonged and deep inspiration followed by a shorter expiration; suspirium.

Sight (*sit*) [AS., *sikt*, sight]. The act of seeing; the special sense concerned in seeing.

S., Day-, hemeralopia. S., Far, S., Long, hyperopia. **S., Night-, nyctalopia. S., Old,** presbyopia. **S., Short,** myopia.

Sigmatism (*sig'-mat-izm*) [*sigma*, the Greek letter *s*]. 1. Defective utterance of the sound of *s*. 2. The too frequent use of the *s* sound in speech.

Sigmoid (*sig'-moid*) [*σιγμα*, the letter *s*; *eidōs*, likeness]. Shaped like the letter S.

S. Cavities, two depressions on the head of the ulna; the *greater* is for articulation with the humerus; the *lesser*, on the outer side of the coronoid process, is for articulation with the radius. **S. Flexure**, an S-shaped bend in the colon between the descending portion and the rectum, usually occupying the left iliac fossa. 2. Pertaining to the S. flexure of the colon, as the S. artery, the S. mesocolon.

Sigmoidostomy (*sig-moi-dos'-to-me*) [*σιγμα*, the letter *s*; *στόμα*, mouth]. The formation of an artificial anus in the sigmoid flexure of the colon.

Sign (*sin*) [*signum*, a mark]. A mark or evidence; in a restricted sense, a physical S. **S., Objective, S., Physical**, one apparent to the observer. **S., Subjective**, one only recognized by the patient. A table of *Eponymic Signs and Symptoms* is appended.

TABLE OF EPONYMIC SIGNS AND SYMPTOMS OF DISEASES.

NAME.	DESCRIPTION.	HOW ELICITED.	DISEASE IN WHICH PRESENT.	SIGNIFICANCE.
Abadie's sign.	Spasm of the levator palpebræ superioris muscle.	Inspection.	Exophthalmic goiter.	
Allis's sign.	Relaxation of the fascia between the crest of the ilium and the trochanter major.	Inspection.	Fracture of the neck of the femur.	
Argyll Robertson pupil.	A pupil that acts in accommodation but not to light.	Usual tests for accommodation and light.	Locomotor ataxia.	
Bacelli's sign (<i>aphonic pectoriloquy</i> .)	Reverberation of the whispered voice, heard through the chest-wall.	Whispered voice of patient auscultated through the chest.	Pleural effusion.	
Baruch's sign.	The resistance of the rectal temperature to a bath of 75° for fifteen minutes, with friction.	Immersion in bath of 75° F.	Typhoid fever.	
Bernhardt's symptom.	Paresthetic and painful sensations on the outer and anterior aspect of the thigh, in the distribution of the external cutaneous nerve.	Condition is subjective.		In some cases dependent on displacement of external cutaneous nerve.

TABLE OF EPONYMIC SIGNS AND SYMPTOMS OF DISEASES.—*Continued.*

NAME.	DESCRIPTION.	HOW ELICITED.	DISEASE IN WHICH PRESENT.	SIGNIFICANCE.
Biermer's change of sound.	See <i>Gerhardt's change of sound</i> (in this table).	Paretic dementia.		
Biernacki's symptom.	Analgesia of ulnar nerve.			
Bouillaud, Tintement métallique of.	A peculiar clink sometimes heard at the right of the apex-beat of the heart.	Auscultation.	Cardiac hypertrophy.	
Brach-Romberg symptom.	See <i>Romberg's symptom</i> .			
Brown-Séquard's paralysis.	Hemiparaplegia with hemianesthesia of opposite side.		Lesion of lateral half of spinal cord.	
Burton's sign.	Blue line at junction of teeth with gums.	Visual examination.	Chronic lead-poisoning.	
Cheyne-Stokes' respiration.	A succession of respirations becoming progressively shorter and more shallow, then an intermission of varying duration, followed by progressive increase in depth and length of respirations.		In various affections in which the brain is implicated. Cerebraledema. Uremia.	
Chvostek's sign.	Sudden spasm of one side of the face.	A slight tap upon the side of the face.	Postoperative tetany.	
Clark's (Alonzo) sign.	Obliteration of the hepatic dulness due to tympanitic distention of the abdomen.	Percussion.	Appendicitis and other peritoneal inflammations.	The presence of the inflated bowel, or of gas in front of the liver.
Corrigan's line.	A purple line at the junction of the teeth with the gums.	Visual examination.	Chronic copper-poisoning.	
Corrigan's pulse.	A forcible pulse-wave, which quickly recedes.	By finger or sphygmograph.	Aortic insufficiency.	
Dalrymple's sign.	Abnormal widening of the palpebral aperture.	Visual examination.	Exophthalmic goiter.	
Davidsohn's sign.	Reflection of light through the pupil in transillumination.	Electric light in the mouth.		Health.
Drummond's whiff.	A whiff heard at the open mouth, during respiration proceeding from the glottis.	Auscultation.	Aortic aneurysm.	
Dietl's crises.	Sharp paroxysmal pains occurring in case of movable kidney.			Probably dependent on acute hydronephrosis from twisting of ureter.
Duroziez's murmur.	A double murmur occasionally heard in the femoral artery.	Auscultation.	Aortic incompetence.	
Farre's tubercles.	Superficial masses felt on the surface of the liver.	Palpation.	Carcinoma of the liver.	
Fillipovitch's sign.	A saffron-like coloration of the prominent parts of palms of hands and soles of feet.	Inspection.	Typhoid fever.	

TABLE OF EPONYMIC SIGNS AND SYMPTOMS OF DISEASES.—*Continued.*

NAME.	DESCRIPTION.	HOW ELICITED.	DISEASE IN WHICH PRESENT.	SIGNIFICANCE.
Fisher's brain-murmur.	A systolic murmur over the anterior fontanel or in the temporal region of infants.	Auscultation.	Rickets and other conditions.	
Flint's murmur.	A second murmur frequently heard at the apex; it has a rumbling quality and may be presystolic, and is probably produced at the mitral orifice.	Auscultation.	Aortic incompetence.	
Foerster's shifting type.	Variations in the field of vision.	Perimetric limits differ according as they are determined by moving the disc from the center outward or from without to the center.	Anesthesia of the retina, traumatic neuroses, etc.	
Friedreich's respiratory change of sound.	The pitch of the percussion-note becomes increased at the height of a deep inspiration.	On percussion.	Pulmonary tuberculosis and other conditions.	
Friedreich's sign.	Diastolic collapse of the cervical veins.	Palpation. Inspection.	Adherent pericardium.	
Garel's sign.	Absence of luminous perception on the affected side of the walls and sinuses about the mouth— <i>antrum</i> of Highmore.	Electric transillumination.	Disease of the <i>antrum</i> .	
Gerhardt's change of sound.	A change of percussion-note according to the patient's position, whether upright, lying on the back, or on the side. Due to changes in the form of the air-space and fluid-contents of the thoracic cavity.	Percussion in the different positions.	Pneumothorax, pulmonary tuberculosis.	
Gerhardt's sign.	Absence of movement of larynx in dyspnea due to aneurysm of aorta.	Inspection.	In dyspnea from other causes, the movements of larynx are extensive.	
Glasgow's sign.	Systolic sound in brachial artery.	Auscultation.	Latent aneurysm of aorta.	
von Graefe's sign.	Failure of the upper lid to follow the eyeball in glancing downward.	By having the patient alternately rotate the eyes up and down.	Exophthalmic goiter.	
Grancher's sign.	The expiratory murmur equals in pitch that of the inspiratory.	Auscultation.	Pulmonary condensation.	Obstruction to expired air.
Gubler's tumor.	A prominence seen on the dorsum of the carpus.	By flexing carpus.	Wrist-drop of chronic lead poisoning.	Probably some effusion into the synovial sacs.
Guyon's sign.	Renal ballotement.	Palpation.	Floating kidney.	

TABLE OF EPONYMIC SIGNS AND SYMPTOMS OF DISEASES.—Continued.

NAME.	DESCRIPTION.	HOW ELICITED.	DISEASE IN WHICH PRESENT.	SIGNIFICANCE.
Heberden's nodosities.	Hard nodules, usually on the distal joints of the fingers.	Palpation.	Rheumatoid arthritis, usually in advanced life.	
Hegar's sign.	A softening of the lower uterine segment.	By forefinger in the rectum and the thumb in the vagina, with pressure from above.	Pregnancy.	
Hick's (Braxton) sign.	Intermittent uterine contraction. Begins to be apparent at end of third month. May also be produced by any tumor that distends uterus.	Palpation.	Pregnancy.	
Hippocratic facies.	An anxious face, with pinched features and sunken eyes.	By involuntary contraction of the facial muscles.	In peritonitis and fatal diseases.	
Hippocratic fingers.	Clubbing of the fingertips, with incurvation of the nails.	Inspection.	Pulmonary tuberculosis and other wasting diseases.	
Hippocratic succussion.	Splashing sound.	By shaking body of patient.	Pyopneumothorax; hydropneumothorax.	Air and fluid in the pleural cavity.
Hutchinson's patch.	Dull-red coloration of the cornea.	Ciliary injection.	Interstitial keratitis.	Syphilis.
Hutchinson's teeth.	Upper central permanent incisor teeth are peg-shaped, and notched on the cutting edge.	Inspection.	Inherited syphilis.	
Hutchinson's trio of symptoms.	Notched teeth, interstitial keratitis, and otitis.		Inherited syphilis.	
Jaccoud's sign.	Prominence of the aorta in the suprasternal notch.	Inspection.	Leukemia (and pseudoleukemia).	
Jacquemin's sign.	Violet color of the mucous membrane of vagina; appears about the fourth week of gestation.	Inspection.	Pregnancy.	Venous congestion.
Jadelot's lines.	Various lines on the face of infants.	Inspection.	Various diseased conditions.	See <i>Jadelot's Lines</i>
Jorissen's sign.	Pulse does not become accelerated on changing from the horizontal to the erect position.	Change of position as described.	Pregnancy.	
Josseraud's sign.	A loud metallic second sound over the pulmonic area.	Auscultation.	Acute pericarditis.	
Keen's sign.	Increased diameter through the leg at the malleoli.	Measurement.	Fracture of fibula (Pott's).	
Küster's sign.	Presence of a cystic tumor in the median line anterior to the uterus.	Palpation and inspection.	Ovarian dermoids.	.

TABLE OF EPONYMIC SIGNS AND SYMPTOMS OF DISEASES.—*Continued.*

NAME.	DESCRIPTION.	HOW ELICITED.	DISEASE IN WHICH PRESENT.	SIGNIFICANCE.
Laennec's perles.	Rounded gelatinous masses.	In sputum.	Bronchial asthma.	Mucous molds of the smaller bronchial tubes.
Laennec's rale.	A modified subcrepitant rale.	Auscultation.	Pulmonary emphysema.	Mucus in the bronchioles.
Litten's sign.	See <i>Diaphragm-phenomenon</i> .			
Mannkopf's symptom.	Increase in the frequency of the pulse.	By pressure on peripheral points.	Pain.	Not present in simulated pain.
Oliver's sign.	Tracheal tugging.	By grasping the larynx between the thumb and finger and pressing upward.	Aneurysm of the aorta.	
Palmoplantar sign.	See <i>Filipovitch's sign</i> .			
Parkinson's facies.	Face expressionless, "wooden;" movements of the lips slow; eyebrows elevated. The whole expression is immobile and mask-like.	Inspection.	Paralysis agitans.	
Parkinson's mask.	See <i>Parkinson's facies</i> .			
Parrot's nodes.	Osteophytes of the skull.	Palpation.	Of syphilitic origin.	Inherited syphilis.
Parrot's sign.	Dilatation of the pupil.	By pinching the skin of the neck.	Meningitis.	
Paul's sign.	A feeble apex-beat, with a forcible impulse over the body of the heart.	Palpation.	Pericarditis.	Pericardial adhesions.
Porter's sign.	See <i>Oliver's sign</i> .			
Pott's boss.	Projecting spinous process.	Palpation.	Pott's disease.	Vertebral caries.
Quincke's pulse or symptom.	Blanching of the fingernails at each diastole of the heart.	Inspection.	Aortic insufficiency.	Very marked regurgitation.
Raynaud's phenomena.	A white and cold condition of the fingers, alternating with burning heat and redness.		Raynaud's disease, <i>q. v.</i>	Vasomotor disturbance.
Ritter's tetanus.	Tetanus of a muscle.	By suddenly breaking the circuit while a strong constant current is passing through a nerve.		Health.
Ritter-Rollet phenomenon.	Flexion of the foot by gentle electric stimulation. Extension of the foot by energetic stimulation.	Voltaic or faradic stimulation.		
Romberg's symptom.	Swaying of the body.	By standing patient with feet close together and with eyes shut.	Locomotor ataxia.	

TABLE OF EPONYMIC SIGNS AND SYMPTOMS OF DISEASES.—*Continued.*

NAME.	DESCRIPTION.	HOW ELICITED.	DISEASE IN WHICH PRESENT.	SIGNIFICANCE.
Rosenbach's sign.	Abolition of abdominal reflex.		In inflammatory intestinal diseases.	
Rosenthal's hyperacid vomiting.	The vomiting of very acid material.		Exaggerated secretion of HCl in the gastric juice.	Rosbach's disease.
Seguin's signal symptom.	The contraction of one muscle or group of muscles, preceding the epileptic attack.	Involuntary.	Epilepsy.	
Skeer's symptom.	A small circle that forms in the iris near the pupil in both eyes simultaneously.	Inspection.	Tuberculous meningitis.	
Skoda's consonating rales.	Bronchial rales heard through consolidated pulmonary tissue.	Auscultation.	Pneumonia.	Mucus in bronchial tubes surrounded by consolidated structure.
Skoda's resonance, sign, or tympany.	A tympanitic note on percussion.	By percussing the chest above a large pleural effusion or above the line of consolidation in pneumonia.	Heard when a pleural effusion extends up to the 4th rib or above.	Vicarious action of the portion of lung not involved (apex or upper lobe).
Stairs-sign.	Difficulty in descending stairs.		Early symptoms of locomotor ataxia.	
Stellwag's symptom.	Apparent widening of the palpebral aperture.	Retraction of upper eyelid.	Exophthalmic goiter.	
Stokes's sign.	Violent throbbing in the abdomen to the right of the umbilicus.	Palpation.	Acute enteritis.	
Tache cérébrale.	The appearance of a red line.	By drawing the finger-nail over the skin.	Tuberculous meningitis, acute fevers, and other conditions.	
Tarnier's sign.	The effacement of the angle between the upper and lower uterine segments.	Digital examination.	Pregnancy.	Inevitable abortion.
Trousseau's phenomenon or sign.	Muscular spasm, which continues as long as pressure is applied.	Pressure on the large arteries or on the nerve-trunk.	Tetany.	Heightened neuromuscular irritability.
Trousseau's spots.	See <i>Tache cérébrale</i> .			
Valleix's points.	Tender spots along the course of a nerve.	Pressure.	Neuralgia.	
Vigouroux's symptom.	Diminished electric resistance of skin.	Electric stimulation.	Exophthalmic goiter.	
Weber's symptom.	Paralysis of the oculomotor nerve of one side and hemiplegia of the opposite side.			
Wernicke's symptom.	See <i>Reaction, Hemiplegic Pupillary</i> .			

TABLE OF EPONYMIC SIGNS AND SYMPTOMS OF DISEASES.—*Continued.*

NAME.	DESCRIPTION.	HOW ELICITED.	DISEASE IN WHICH PRESENT.	SIGNIFICANCE.
Westphal's foot-phenomenon.	Ankle-clonus.	By suddenly flexing foot on leg.	Lateral sclerosis and various other conditions.	
Westphal's paradoxical contraction.	The tonic contraction of a muscle caused by the passive approximation of its extremities.	In anterior tibial, by rapid dorsal flexion of the foot.	Paralysis agitans (Westphal), and sclerosis of the posterolateral columns of the cord.	
Westphal's sign.	Abolition of the knee-jerk.	Percussion over the quadriceps-tendon.	Locomotor ataxia, destructive lesions of lower part of cord, peripheral neuritis, etc.	
Westphal-Erb symptom.	Same as <i>Westphal's sign</i> .			
Williams' tracheal tone.	A dull tympanitic resonance, becoming higher pitched on opening the mouth.	Percussion.	Pleural effusion.	Arises from the vibration of air in a large bronchus surrounded by compressed lung.
Wintrich's sign.	Change in the pitch when the mouth is opened and closed.	Percussion.	Tuberculosis of lung.	Pulmonary cavity.

Signa (*sig'-nah*) [*signare*, to make a sign]. Mark. In prescription-writing, a term placed before the physician's directions to the patient concerning the medicine prescribed; abbreviated to S. or Sig.

Signature (*sig'-nat-ür*) [*signare*, to make a sign]. 1. The part of the prescription that is to be placed on the label. 2. A distinguishing character. **Doctrine of Signatures**, a theory that the medicinal uses of plants or other objects can be determined from the signatures or peculiar characters.

Sikimin (*sik'-im-in*) [*sikkim*, a region of the Himalaya]. A poisonous principle derived from *Illicium religiosum*.

Silent Region of the Spinal Cord. The gray matter of the cord.

Silica (*sil'-ik-ah*) [*silex*, flint]. Silicon dioxide, SiO_2 , occurring in nature in the form of quartz, flint, and other minerals.

Silicate (*sil'-ik-ät*) [*silex*, flint]. A salt of silicic acid.

Silicic Acid (*sil'-is'-ik*) [*silex*, flint], H_2SiO_4 . A tetrabasic acid, forming the silicates. See *Sodium silicate*.

Silicon (*sil'-ik-on*) [*silex*, flint]. A nonmetallic element occurring widely distributed in nature as silica, SiO_2 , and in the form of silicates. Atomic weight 28.3; symbol Si; valence four. It resembles carbon in its chemic behavior.

Silicosis (*sil'-ik-o'-sis*) [*silex*, flint]. A deposit of particles of silica in the tissues; specifically, a chronic fibroid condition of the lung or the bronchial lymphatic glands, produced by the inhalation of particles of silica.

Silk-worm Gut. The thread drawn from the silk-worm killed when ready to spin the cocoon.

Silver (*sil'-ver*). See *Argentum*.

Silver-fork Deformity. A peculiar deformity of the wrist and hand in Colles' fracture, resembling a fork.

Simaruba (*sim-ar-ü'-bah*). A genus of trees of the order Simarubaceæ. The bark of the root of *S. officinalis* has been used as a simple bitter.

Simple (*sim'-pl*) [*simplex*, single]. Not complex; consisting of but one substance, or containing only one active substance; not compound.

Simples (*sim'-plz*) [*simplex*, simple]. A term for herbs having a medicinal value.

Sims' Position. See *Postures*, Table of.

Sims' Speculum [J. Marion Sims, an American gynecologist]. See *Speculum*.

Simulation (*sim-u-la'-shun*) [*simulare*, to feign]. A feigning or counterfeiting.

Sinalbin (*sin-al'-bin*). See *Mustard*.

Sinapin (*sin'-ap-in*) [*σίναπι*, mustard], $\text{C}_{16}\text{H}_{23}\text{NO}_5$. A substance occurring as a sulphocyanate in white mustard.

Sinapis (*sin-a'-pis*) [*σίβημι*, mustard]. See *Mustard*.

Sinapism (*sin'-ap-izm*) [*σίβημι*, mustard]. A mustard-plaster.

Sinapized (*sin'-ap-īzd*) [*σίβημι*, mustard]. Containing mustard.

Sincalin (*sing'-ka-lin*). A base found in mustard and identified with cholin.

Sincipital (*sin - sip' - it - al*) [*sinciput*]. Pertaining to the sinciput.

Sinciput (*sin'-sip-ut*) [*semi*, half; *caput*, head]. The superior and anterior part of the head.

Sinew (*sin'-u*) [AS., *sinu*, sinew]. See *Tendon*.

Singultus (*sing-gul'-tus*). See *Hiccup*.

Sinigrin (*sin'-ig-rin*). See *Mustard*.

Sinistrad (*sin'-is-trad*) [*sinister*, left; *ad*, toward]. Toward the left.

Sinistral (*sin'-is-tral*) [*sinister*, left]. On the left side.

Sinistrin (*sin'-is-trin*) [*sinister*, left]. A substance resembling dextrin, found in squills.

Sinus (*si'-nus*) [*sinus*, a gulf or hollow]. 1.

A hollow or cavity; a recess or pocket. 2.

A large channel containing blood, especially

one containing venous blood. 3. A suppurating tract. S., Air-, a cavity within bones

containing air, especially one communicating

with the nasal passages. S., Aortic. See

S. of *Valsalva*. S., Cavernous, a large

venous S. extending from the sphenoid fis-

sure to the apex of the petrous portion of

the temporal bone, communicating behind

with the inferior and superior petrosal

sinuses and receiving the ophthalmic vein

in front. S., Circular, a venous S. sur-

rounding the pituitary body, and communicat-

ing on each side with the cavernous S.

S., Coronary (of the heart), a large venous

S. in the transverse groove between the left

auricle and left ventricle of the heart. S.,

Frontal, one of the two irregular cavities

in the frontal bone containing air and com-

municating with the nose by the infundib-

ulum. S., Inferior Longitudinal, a venous

S. which extends along the posterior half of

the lower border of the falx cerebri and ter-

minates in the straight S. S., Inferior

Petrosal, a large venous S. arising from the

cavernous S., running along the lower margin

of the petrous portion of the temporal bone,

and joining the lateral S. to form the internal

jugular vein. S. of Kidney, the prolonga-

tion inward of the hilum of the kidney. S.

of the Larynx, the ventricle of the larynx.

S., Lateral, a venous S. which begins at the

torcular Herophili and runs horizontally on the

inner surface of the occipital bone to the base

of the petrous portion of the temporal bone,

where it unites with the inferior petrosal S.

to form the internal jugular vein. S. of Morgagni, the interval between the upper border of the superior constrictor muscle and the basilar process of the occipital bone. S., Occipital, a small venous S. in the attached margin of the falx cerebelli, opening into the torcular Herophili. S. pocularis. See *Uterus masculinus*. S., Prostatic. See *Uterus masculinus*. S., Rhomboid, S. rhomboideus, the fourth ventricle of the brain. S., Sphenoid, the air-space in the body of the sphenoid bone, communicating with the nasal cavity. S., Straight, a venous S. running from the inferior longitudinal S. along the junction of the falx cerebri and tentorium to the lateral S. S., Superior Longitudinal, a venous S. which runs along the upper edge of the falx cerebri, beginning in front at the crista galli and terminating at the torcular Herophili. S., Superior Petrosal, a venous S. running in a groove in the petrous portion of the temporal bone extending from the posterior part of the cavernous S. to the lateral S. S., Terminal, S. terminalis, a vein that encircles the vascular area of the blastoderm, and empties either by one trunk, the anterior vitelline vein, into the left vitelline vein, or by two trunks into both vitelline veins. S., Transverse, a S. uniting the inferior petrosal sinuses. S., Urogenital, the canal or duct into which, in the embryo, the Wolffian ducts and the bladder empty, and which opens into the cloaca. S. of Valsalva, one of the pouch-like dilatations of the aorta or pulmonary artery opposite the segments of the semilunar valves.

Sinus-phlebitis (*si-nus-fle-bi'-tis*) [*sinus*, a gulf; *phlebitis*]. Inflammation of one of the sinuses of the cranial cavity.

Sinus-thrombosis (*si-nus-throm-bo'-sis*) [*sinus*, a gulf; *thrombosis*]. Thrombosis of the sinuses of the dura mater of the brain. It is usually septic in character and is apt to lead to pyemia. The most frequent cause is disease of the middle ear.

Siphon (*si'-fon*) [*σίφων*, a tube]. A tube bent at an angle, one arm of which is longer than the other, for the purpose of removing liquids from a cavity or vessel.

Siphonoma (*si-fon-o'-mah*) [*σίφων*, siphon; *μα*, tumor]. A tumor composed of fine tubes; also known as *Henle's tubular tumor*.

Sirenomelus (*si-ren-om'-el-us*) [*σειρήν*, mermaid; *μέλος*, limb]. A form of monster in which the lower extremities are intimately fused, the feet being absent.

Sirup (*sir'-up*). See *Syrup*.

Sitiophobia (*sit-e-o-fo'-be-ah*). See *Sitophobia*.

Sitomania (*si-to-ma'-ne-ah*) [*σιτος*, food; *μανία*, madness]. 1. A periodic craving for food; periodic bulimia. 2. Sitophobia.

Sitophobia (*si-to-fo'-be-ah*) [*σίτος*, food; *φόβος*, fear]. Morbid aversion to food.

Situs (*si'-tus*) [*situs*, site]. A position. **S. viscerum inversus**, an anomaly in which the viscera of the body are changed from the normal to the opposite side of the body.

Sitzbath (*sitz'-bath*) [Ger., *Sitz*, a seat; *bath*]. A hip-bath; a bath taken in a sitting posture.

Skatol (*skat'-ol*) [*σκατός*, gen. of *σκήω*, dung], $C_8H_9N_3$. A nitrogenous compound produced by the decomposition of proteids in the intestinal canal.

Skeletal (*skel'-et-al*) [*skeleton*]. Pertaining to or connected with the skeleton or supporting structure of a body.

Skeletization (*skel-et-i-zā'-shun*) [*σκελετών*, skeleton]. The process of converting into a skeleton; gradual wasting of the soft parts, leaving only the skeleton.

Skeletogenous (*skel-et-ōj'-en-us*) [*σκελετόν*, skeleton; *γεννᾶν*, to produce]. Producing a skeleton or skeletal tissues.

Skeletography (*skel-et-ōg'-ra-fe*) [*σκελετόν*, skeleton; *γράφειν*, to write]. A description of the skeleton.

Skeletology (*skel-et-ōl'-o-jē*) [*σκελετόν*, skeleton; *λόγος*, science]. The branch of anatomy treating of the skeleton.

Skeleton (*skel'-et-on*) [*σκελετόν*, a dried body, from *σκέλλειν*, to dry up]. A supporting structure, especially the bony framework (osseous **S.**) supporting and protecting the soft parts of an organism. **S., Cartilaginous**, the cartilaginous structure from which the bony **S.** is formed through ossification.

Skiagraphy (*ski-ag'-ra-fe*). See *Skiography*.

Skiascopy (*ski-as'-ko-pe*) [*σκία*, shadow; *σκοπεῖν*, to see]. See *Retinoscopy*.

Skin [ME., *skin*, skin]. The protective covering of the body composed of the epidermis, scarf-skin, or cuticle, and the corium, or true **S.** The epidermis consists of a deep layer, the stratum Malpighii, and three superficial layers, the stratum granulosum, the stratum lucidum, and the stratum corneum. The corium, derma, or true **S.**, consists of a papillary and reticular layer (stratum papillare and stratum reticulare), the former projecting upward in the form of papillæ. The true **S.** is made up of elastic tissue, white fibrous tissue, and nonstriped muscular tissue (the arrectores pili). The subcutaneous tissue consists of fibroelastic and adipose tissue. The appendages of the **S.** are the nails, hairs, and sweat- and sebaceous glands, which are derivatives of the epithelial layer of the **S.** In the skin are also placed terminal nerve-organs subserving the sense of touch. **S.-bound Disease**. See *Scleroderma*. **S., Glossy**, a peculiar shiny, glazed skin seen in conditions in which the

trophic nerve-supply to the **S.** is cut off, as after injury to a nerve. **S., Goose-**. See *Goose-skin*. **S.-grafting**, the application of pieces of the outer layers of healthy skin to a granulating surface for the purpose of hastening its cicatrization.

Skiography (*ski-ōg'-ra-fe*) [*σκία*, shadow; *γράφειν*, to write]. Photography by the Röntgen or X-rays. Skotography, Skiagraphy, Radiography, Electrophotography, Electroskiography, Röntgenography, and the New Photography are some of the names that have been proposed to designate the method.

Skoda's Sign, S.'s Resonance, or **S.'s Tympany** [*Skoda*, an Austrian physician]. The tympanic percussion-note above the level of a pleural effusion.

Skodaic Resonance. See *Skoda's Sign*.

Skotography (*sko-tōg'-ra-fe*). See *Skiography*.

Skull (*skul*) [Icel., *skáli*, a bowl]. The bony framework of the head, consisting of the cranium and the face. The cranium is made up of the occipital, frontal, sphenoid, and ethmoid bones, and the two parietal and two temporal bones. The face is composed of two nasal, two superior maxillary, two lacrimal, two malar, two palate, and two inferior turbinated bone, and the vomer and inferior maxillary bone. **S.-cap.** 1. The top of the skull. 2. See *Scutellarium*.

Skunk-cabbage. The *Dracopis foetidum*, the rhizome of which is stimulant, antispasmodic, and narcotic, and has been used in asthma, rheumatism, hysteria, and dropsy.

Sleep. The periodic state of rest in which voluntary consciousness and activity cease.

S.-epilepsy. See *Narcolepsy*. **S., Hypnotic**, **S., Magnetic**, **S., Mesmeric**. See *Hypnotism*. **S.-walking**. See *Sonambulism*.

Sleeping Sickness. See *African Lethargy*.

Slender Column. See *Funiculus gracilis*.

S. Lobe of Cerebellum, a small lobe in the inferior surface of the cerebellum.

Slide (*slid*). A small, rectangular plate of glass upon which objects intended for examination with the microscope are placed.

Sling. A swinging bandage for supporting an arm or other part.

Slough (*sluf*) [ME., *slouk*, the skin of a snake]. A mass of soft tissues destroyed by gangrene.

Sloughing (*sluf'-ing*) [ME., *slouk*, the skin of a snake]. Pertaining to or characterized by sloughs.

Small-pox. See *Varicella*.

Smec-cell. See *Batteries*, *Table of*.

Smegma (*smeg'-mah*) [*σμήγμα*, a cleansing substance]. Sebum. **S. preputii**, or simply **S.**, the substance secreted by the sebaceous glands of the prepuce.

Smell. 1. The perception of odor. 2. Odor. **Smelling-salts.** A name applied to various preparations of ammonium carbonate flavored with aromatic substances.

Smilax (*smi'-laks*). See *Sarsaparilla*.

Smith's Cramp. An occupation-neurosis occurring in smiths and characterized by painful cramps in the arm or hand.

Sn. Symbol for tin (L., *stannum*).

Snake-root. See *Cimicifuga*, *Senega*, and *Serpentaria*.

Snare [AS., *snear*, a cord]. A loop of wire or other material used in removing projecting growths.

Sneeze (*snez*) [AS., *fnéosan*, to sneeze]. A sudden, noisy, spasmodic expiration through the nose.

Snellen's Types. See *Test-types*.

Snore, Snoring [ME., *snoren*, to snore].

1. To breathe through the nose in such manner as to cause a vibration of the soft palate, thereby producing a rough, audible sound. 2. The sound so produced.

Snow-blindness. See *Blindness*.

Snuffles. Coryza, especially of infants, which is frequently due to inherited syphilis.

Soap [*sapo*]. A chemic compound made by the union of certain fatty acids with an alkali or other metal. According to the alkali used, the S. formed is a potash-S., soda-S., ammonia-S., lead-S., lime-S., etc. **S.-bark.** See *Quillaja*. **S., Castile, S.** made from olive-oil. **S., Hard.** See *Soda-S.* **Potash-S.** (soft S.) is made from linseed-oil and potash (Sapo mollis, U. S. P.) or from olive-oil and potash (Sapo mollis, B. P.). From it is prepared Linimentum saponis mollis (U. S. P. **Soda-S.** is made from soda and olive oil (Sapo, U. S. P., Sapo durus, B. P.). From it are prepared Emplastrum saponis (U. S. P., B. P.), used as a local sedative; Linimentum saponis (U. S. P., B. P.), liquid opodeldoc, used as a sedative liniment in rheumatic affections and sprains. **Soda-S.** also enters into the composition of various pills. **S., Soft.** See *Potash-S.*

Sob. A convulsive inspiration due to contraction of the diaphragm and spasmodic closure of the glottis.

Socaloin (*so-kul'-o-in*) [*Socotra*, an island in the Indian Ocean; *álon*, aloes]. See *Aloes*.

Socia parotidis (*so'-se-ah par-ot'-id-is*). A small separate lobe of the parotid gland.

Socket (*sok'-et*) [ME., *soket*]. The concavity into which a movable part is inserted.

Soda (*so'-dah*) [Ital., from L., *solidus*, solid]. 1. Sodium oxid, Na_2O . 2. Sodium carbonate or sodium bicarbonate. **S., Baking,** sodium bicarbonate. **S., Caustic,** sodium hydroxid. **S.-soap.** See *Soap*. **S., Washing,** sodium carbonate. **S.-water,** water impregnated with carbon dioxide.

Sodic (*so'-dik*) [*soda*, soda]. Derived from or containing soda.

Sodium (*so'-de-um*) [*soda*]. A metallic element of the alkaline group of metals, melting at 95.6°C ., and having a specific gravity of 0.97, an atomic weight of 23, and a valence of one. Symbol Na, from the Latin natrium. S. occurs widely distributed in nature, and forms an important constituent of animal tissues. It has a strong affinity for oxygen and other nonmetallic elements. It is also a constituent of many medicinal preparations, the most important of which are the following: **S. acetate** (Sodii Acetas, U. S. P.), $\text{NaC}_2\text{H}_3\text{O}_2 \cdot 3\text{H}_2\text{O}$, is diuretic. Dose gr. xx- \mathfrak{z} ij (1.3-8.0). **S. arsenate** (Sodii arsenas, U. S. P., B. P.), $\text{N}_2\text{HAsO}_4 \cdot 7\text{H}_2\text{O}$, is used like the other preparations of arsenic. Dose gr. $\frac{1}{2}$ - $\frac{1}{4}$ (0.005-0.02). From it is prepared Liquor sodii arsenatis. **S. aurochlorid**, chlorid of gold and sodium. See *Aurum*. **S. benzoate** (Sodii benzoas, U. S. P., B. P.), $\text{NaC}_7\text{H}_5\text{O}_2$, is used in gout, rheumatism, lithemia, influenza, etc. Dose \mathfrak{z} j-ij (4.0-8.0). **S. bicarbonate** (Sodii bicarbonas, U. S. P., B. P.), NaHCO_3 , is used as an antacid in dyspepsia, gout, rheumatism, lithemia, and diabetes. Dose gr. x- \mathfrak{z} j (0.65-4.0). **S. bisulphite** (Sodii bisulphis, U. S. P.), NaHSO_3 , is used to check gastric fermentation, and as a parasiticide in skin-diseases. Dose gr. viij-xxx (0.52-2.0). **S. borate**, or borax (Sodii boras, U. S. P., B. P.), $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$, is used in dysmenorrhea, in uric acid diathesis, in stomatitis, and as an antiseptic. In overdoses it is a depressant poison. **S. bromid** (Sodii bromidum, U. S. P., B. P.), NaBr, is used like the other bromids. Dose \mathfrak{z} ss- \mathfrak{z} ij (2.0-8.0). **S. carbonate** (Sodii carbonas, U. S. P., B. P.), $\text{Na}_2\text{CO}_3 \cdot 10\text{H}_2\text{O}$, is used as an antacid, and locally in diseases of the skin and in superficial burns. Dose gr. x- \mathfrak{z} ss (0.65-2.0). Dried S. carbonate (Sodii carbonas exsiccatus, U. S. P., B. P.) is used like the carbonate. Dose v-xv (0.32-1.0). **S. chlorate** (Sodii chloras, U. S. P.), NaClO_3 , has medical properties similar to those of potassium chlorate, but is more soluble. Dose gr. v-xv (0.32-1.0). **S. chlorid** (Sodii chloridum, U. S. P., B. P.), NaCl, common salt, is a constituent of animal fluids and tissues and of food. In medicine it is used as a stomachic; in hemoptysis; as an application to sprains and bruises; as a tonic and stimulant in the form of salt-water baths; as a cathartic; and in the form of a 0.6-0.75 per cent. solution to replace the loss of fluids from hemorrhage or profuse diarrhea, being used as an intravenous, subcutaneous, or rectal injection. **S. citrate** is used as a purgative. Dose \mathfrak{z} x-xiv (4.0-56.0). **S.**

citrotartrate (Sodii citrotraras effervescens, B. P.) is refrigerant and laxative. Dose \mathfrak{z} j-ij (4.0-8.0). S. copaivate, $\text{NaC}_{10}\text{H}_{12}\text{O}_6$, is used like copaiba. S. dithiosalicylate is used like salicylic acid in rheumatism. Dose gr. iij (0.2). S. ethylate, NaOC_2H_5 , is used as a caustic. S. hydrate, S. hydrazid, soda, caustic soda (Soda, U. S. P., Soda caustica, B. P.), NaOH , is an extremely corrosive substance, occurring in the form of white pencils; it is used as a caustic and as an antacid, like the bicarbonate. From it is prepared Liquor sodæ (U. S. P., B. P.). Dose \mathfrak{m} v-xxx (0.32-2.0). S. hypochlorite, NaOCl , is a constituent of Liquor sodæ chloratæ, U. S. P. (Liquor sodæ chlorinatæ, B. P.), Labarraque's solution. S. hypophosphite (Sodii hypophosphis, U. S. P., B. P.), $\text{NaH}_2\text{PO}_2 \cdot \text{H}_2\text{O}$, is used like the other hypophosphites in pulmonary tuberculosis, acrofula, rickets, etc. Dose gr. x-xxx (0.65-2.0). S. hyposulphite (Sodii hyposulphis, U. S. P.), $\text{Na}_2\text{S}_2\text{O}_5 \cdot 5\text{H}_2\text{O}$, is used to check fermentation, and locally in parasitic diseases of the skin and mouth. Dose gr. x-xx (0.65-1.3). S. iodid (Sodii iodidum, U. S. P., B. P.), NaI , is used like potassium iodid. S. nitrate (Sodii nitras, U. S. P., B. P.), NaNO_3 , has been used in dysentery. Dose \mathfrak{z} ss- \mathfrak{z} j (16.0-32.0). S. nitrite (Sodii nitris, U. S. P., B. P.), NaNO_2 , is used like the other nitrites, but its effects are more slowly produced and more permanent. Dose gr. j-ij (0.065-0.2). See *Nitrite*. S. nitroprussid, $\text{Na}_2\text{Fe}(\text{CN})_5\text{NO}$, is used as a reagent. S. orthophosphate, S. phosphate (Sodii phosphas, U. S. P., B. P.), $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$, is a purgative and cholagogue, and is used in jaundice, gall-stones, diarrhea of children, and rickets. Dose gr. iij-x (0.2-0.65) for children; gr. xx-xl (1.3-2.6) for adults; as a purgative, \mathfrak{z} j-ij (32.0-64.0). Sodii phosphas effervescens (B. P.) is used like the phosphate. Dose \mathfrak{z} ij-iv (8.0-16.0). S. and Potassium tartrate, Rochelle salt. See *Potassium*. S. pyrophosphate (Sodii pyrophosphas, U. S. P.), $\text{Na}_4\text{P}_2\text{O}_7 \cdot 10\text{H}_2\text{O}$, is used for preparing ferric pyrophosphate. S. salicylate (Sodii salicylas, U. S. P., B. P.), $\text{NaC}_7\text{H}_5\text{O}_3$, has the properties and uses of salicylic acid. Dose gr. xv-xxx (1.0-2.0). S. santoninate. See *Santonin*. S. silicate, Na_2SiO_3 or Na_3SiO_3 , is used in preparing Liquor sodii silicatis, which is employed as a surgical dressing. S. silicofluorid, Na_2SiF_6 , is said to be antiseptic. Its solution is known as *salufer*. S. soziodol is used as a parasiticide in skin-diseases and as an antiseptic. S. sulphate, Glauber's salt (Sodii sulphas, U. S. P., B. P.), $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$, is a hydragogue cathartic and diuretic. Dose \mathfrak{z} ss-

\mathfrak{z} j (16.0-32.0). Sodii sulphas effervescens (B. P.) is used for the same purposes. Dose \mathfrak{z} ij-iv (8.0-16.0). S. sulphite (Sodii sulphis, U. S. P., B. P.), $\text{Na}_2\text{SO}_3 \cdot 7\text{H}_2\text{O}$, is used like S. hyposulphite. S. sulphocarbolate (Sodii sulphocarbolas, U. S. P., B. P.), $\text{NaSO}_3\text{C}_6\text{H}_4(\text{OH}) \cdot 2\text{H}_2\text{O}$, is used locally as an antiseptic and internally as an antiseptic in intestinal fermentation. Dose gr. x-xxx (0.65-2.0). S. tartrate, $\text{Na}_2\text{C}_4\text{H}_4\text{O}_6 \cdot 2\text{H}_2\text{O}$, is used as an antacid and as a refrigerant in fevers. See *S. citrotraras*. S. valerianate (Sodii valerianas, B. P.) is used as a nervous stimulant. Dose gr. j-v (0.065-0.32).

Sodom (*sod'-om-e*) [*Sodom*, a city of ancient Palestine]. Sexual connection by the anus. **Sommering's Yellow Spot**. The macula lutea.

Soft. Yielding readily to pressure; not hard. **S. Palate**. See *Palate*. **S. Water**, one containing but little mineral matter and forming free lather with soap.

Softening (*sof'-en-ing*). The act of becoming soft.

Solanin (*sol'-lan-in*) [*solanum*, the nightshade]. An alkaloid found in various species of solanum.

Solanum (*so-la'-num*) [L.]. A genus of the Solanaceæ including the tomato, potato, bitter-sweet, and black nightshade. *S. carolinense*, horse-nettle, has been used in epilepsy.

Solar Plexus (*sol'-lar*) [*solaris*, from *sol*, the sun, so called because of the radiating nerves]. A plexus consisting of a network of nerves and ganglia (Solar Ganglia) and situated behind the stomach and in front of the aorta and crura of the diaphragm. It receives the great splanchnic nerves and filaments from the right pneumogastric nerve, and supplies branches to all the abdominal viscera.

Sole (*söl*) [AS., *sole*]. The plantar surface of the foot. **S.-reflex**. See *Reflexes*, *Table of*.

Soleus (*so-le'-us*). A flat muscle of the calf. See *Muscles*, *Table of*.

Solferino (*sol-fer-e'-no*) [an Italian city, the scene of a battle]. A synonym of *Fuchsin*.

Solid (*sol'-id*) [*solidus*]. 1. Firm; dense; not fluid or gaseous. 2. Not hollow. 3. A firm body; a body the molecules of which are in a condition of strong mutual attraction.

Solidism (*sol'-id-izm*) [*solidus*, solid]. The theory that diseases depend upon alterations in the solids of the body.

Solitary (*sol'-it-a-re*) [*solitarius*, solitary]. Single; existing separately; not collected together. **S. Bundle**, a strand of nerve-fibers in the medulla. **S. Follicles**, **S. Glands**, minute lymphatic nodules in the mucous membrane of the intestines.

Solubility (*sol-u-bil'-it-e*) [*solubilis*, from

solvere, to dissolve]. The state of being soluble.

Soluble (*sol'-u-bl*) [*solubilis*]. Capable of being dissolved.

Solution (*so-lu'-shun*) [*solutio*, from *solvere*, to loosen]. 1. A separation or break, as S. of continuity. 2. The process of dissolving a solid, or of being dissolved. 3. A liquid in which a substance has been dissolved. S., Donovan's. See *Arsenic*. S., Fowler's. See *Arsenic*. S., Labarraque's. See *Sodium*. S., Monsel's. See *Ferrum*. S., Normal. See *Normal*. S., Normal Saline. See *Saline*. S., Saturated. See *Saturated*.

Solutol (*sol-u'-tol*) [*solvere*, to dissolve]. An alkaline solution of cresol in sodium cresylate, used as a disinfectant.

Solvent (*sol'-vent*) [*solvere*, to dissolve]. 1. Capable of dissolving. 2. A liquid capable of dissolving.

Solveol (*sol'-ve-ol*). A neutral solution of cresol in sodium cresylate, used as a disinfectant.

Solvine (*sol'-vin*) [*solvere*, to dissolve]. One of a series of liquids obtained from certain oils by the action of concentrated sulphuric acid; it is a powerful solvent, and also possesses the property of dissolving the red corpuscles.

Somacule (*so'-mak-ul*) [dim. of *σῶμα*, the body]. A physiologic unit corresponding to, but greatly more complex than, the chemic molecule.

Somatic (*so-mat'-ik*) [*σωματικός*, from *σῶμα*, body]. 1. Pertaining to the body. 2. Pertaining to the framework of the body and not to the viscera.

Somatology (*so-mat-ol'-o-je*) [*σῶμα*, body; *λόγος*, science]. The study of anatomy and physiology, or of organized bodies; biology apart from psychology.

Somatome (*so'-mat-ōm*) [*σῶμα*, body; *τομή*, a cutting]. A transverse segment of an organized body; a somite.

Somatopleure (*so-mat'-o-plūr*) [*σῶμα*, the body; *πλευρά*, the side]. The body-wall.

Somite (*so'-mit*) [*σῶμα*, the body]. 1. A segment of the body of an embryo. 2. One of a series of segments of the mesoblast on each side of the dorsal ridge of the embryo; a protovertebra; a protovertebral or mesoblastic S.

Somnal (*som'-nal*) [*somnus*, sleep]. A crystalline substance, a compound of chloral hydrate and urethane. It is diuretic and hypnotic. Dose gr. xxx (2.0).

Somnambulism (*som-nam'-bu-lizm*) [*somnus*, sleep; *ambulare*, to walk]. 1. The condition of half-sleep, in which the senses are but partially suspended; also sleep-walking, a condition in which the individual walks during sleep. 2. The type of hypnotic sleep

in which the subject is possessed of all his senses, often having the appearance of one awake, but whose will and consciousness are under the control of the hypnotizer.

Somnifacient (*som-ne-fa'-shent*) [*somnus*, sleep; *facere*, to make]. 1. Producing sleep. 2. A medicine producing sleep.

Somniloquence, Somniloquism, Somniloquy (*som-nil'-o-kwens*, *som-nil'-o-kwism*, *sam-nil'-o-kwe*) [*somnus*, sleep; *loqui*, to talk]. The act of talking during sleep.

Somnolentia (*som-no-len'-she-ah*) [L.]. Sleep-drunkenness, a condition of incomplete sleep in which a part of the faculties are abnormally excited, while the others are in repose.

Sonometer (*so-nom'-et-er*) [*sonus*, a sound; *μέτρον*, a measure]. 1. An instrument for determining the pitch of sounds and their relation to the musical scale. 2. An instrument for testing hearing.

Sonorous (*so-no'-rus*) [*sonus*, sound]. Capable of producing a musical sound; of rales, low-pitched.

Soot-cancer. Epithelioma of the scrotum; so-called from its frequency in chimney-sweeps.

Sophistication (*so-fis-tik-a'-shun*) [*σοφιστικός*, deceitful]. The adulteration or imitation of a substance.

Sopor (*so'-por*) [L.]. Sleep, especially the profound sleep symptomatic of a morbid condition.

Soporiferous (*so-por-iff'-er-us*). See *Soporific*.

Soporific (*so-por-iff'-ik*) [*sopor*, sleep; *facere*, to make]. 1. Producing sleep. 2. A remedy producing sleep.

Sorbefacient (*sor-be-fa'-shent*) [*sorbere*, to suck; *facere*, to make]. Promoting absorption.

Sordes (*sor'-dēs*) [*sordere*, to be foul]. Filth, dirt, especially the crusts that accumulate on the teeth and lips in continued fevers.

Sore. 1. Painful; tender. 2. An ulcer or wound.

Souffle (*soo'-fl*) [Fr.]. A blowing sound. S., Fetal, an inconstant murmur heard over the uterus during pregnancy, and supposed to be due to the compression of the umbilical cord. S., Funic or Funicular, a hissing sound, synchronous with the fetal heart-sounds, heard over the abdomen of a pregnant woman, and supposed to be produced in the umbilical cord. S., Splenic, a sound said to be audible over the spleen in cases of malaria and leukemia. S., Placental, S., Uterine, a sound heard in the latter months of pregnancy, and caused by the entrance of blood into the dilated arteries of the uterus.

Sound [1. *sonus*, sound; 2. Fr., *sonder*, to probe]. 1. The sensation produced by

stimulation of the auditory nerve by aerial vibrations. 2. An instrument for introduction into a channel or cavity, for determining the presence of constriction, foreign bodies, or other morbid conditions, and for the purpose of treatment.

Southern-wood. See *Artemisia*.

Southey's Drainage-tubes. Small tubes employed for draining away the fluid from limbs that are the seat of extensive anasarca.

Sozin (*so'-zin*) [*σώζω*, to save, keep]. A defensive proteid occurring naturally in the animal body. One capable of destroying microorganisms is termed a *mycosozin*, one antagonizing bacterial poisons, *toxosozin*.

Soziodol, Soziodolic Acid (*so-zí'-o-dol*, *so-sí-o-dol'-ik*) [*σώζω*, to save; *ιώδης*, like a violet], $C_8H_7I_3(SO_3H)OH$. A crystalline, odorless powder used as an antiseptic, disinfectant, and parasiticide, chiefly in the form of its salts, of which the following have been employed: Sodium S., potassium S., zinc S., and mercury S.

Sozolic Acid (*so-so'-lik*). See *Aseptol*.

Space-sense. A sense by which we judge of the relation of objects in space; it is a part of the sense of sight.

Spanemia (*span'-e'-me-ah*) [*σπάνις*, scarcity; *αἷμα*, blood]. Anemia.

Spanish Fly. See *Cantharis*. **S. Windlass.** See *Windlass*.

Sparadrap (*spar'-a-drap*) [*sparadrapum*]. A plaster, spread upon cotton, linen, silk, leather, or paper.

Spargosis (*spar-go'-sis*) [*σπαργάω*, to be distended]. 1. Enlargement of the breasts from accumulation of milk. 2. Elephantiasis.

Sparteín (*spar'-te-in*). An alkaloid found in *Scoparius*, the sulphate of which (*Sparteínæ* sulphas, U. S. P.) is used as a cardiac stimulant and diuretic. Dose gr. $\frac{1}{6}$ – $\frac{1}{2}$ (0.01–0.032).

Spasm (*spazm*) [*σπασμός*, spasm]. A sudden muscular contraction. **S., Clonic,** a S. broken by relaxations of the muscles. **S., Habit,** a S. acquired from habit. **S., Handicraft.** See *Occupation-neurosis*. **S., Tetanic, S., Tonic,** a S. that persists without relaxation for some time.

Spasmo- (*spaz'-mo-*) [*σπασμός*, spasm]. A prefix denoting pertaining to spasm.

Spasmodic (*spaz'-mod'-ik*) [*σπασμός*, spasm]. Pertaining to or characterized by spasm. **S. Spinal Paralysis.** See *Lateral Sclerosis*.

Spasmophilia (*spaz-mo-fil'-e-ah*) [*σπασμός*, spasm; *φιλέω*, to love]. A morbid tendency to convulsions.

Spastic (*spas'-tik*) [*σπαστικός*, spastic]. Pertaining to or characterized by spasm; produced by spasm. **S. Diplegia.** See *Paraplegia*, *Infantile Spasmodic*. **S. Paralysis.** See *Paralysis*.

Spasticity (*spas-tis'-it-e*) [*σπαστικότητα*, from *σπάειν*, to draw]. The state of being spastic.

Spatula (*spat'-u-lah*) [L., dim. of *spatha*, a ladle]. A flexible blunt blade used for spreading ointments.

Spay [Gael., *spath*]. To remove the ovaries.

Spearmint (*spe'-mint*). See *Mentha viridis*.

Species (*spe'-shes*) [L.]. 1. A subdivision of a genus of animals or plants the individuals of which are either identical in character or differ only in unimportant and inconstant details. 2. A powder made from the leaves or other parts of various plants, and used to prepare a medicinal infusion.

Specific (*spe-sif'-ik*) [*species*, species; *facere*, to make]. 1. Of or pertaining to a species, or to that which distinguishes a thing or makes it of the species of which it is. 2. A medicine which has a distinct curative influence on an individual disease. 3. Produced by a single microorganism, as a S. disease; in a restricted sense, syphilitic. **S. Gravity.** See *Gravity*. **S. Heat.** See *Heat*.

Specificity (*spe-sif-is'-it-e*) [*species*, species; *facere*, to make]. The quality of being specific.

Specillum (*spe-sil'-um*) [L.]. A probe.

Specimen (*spe'-im-en*) [L.]. An example; a sample.

Spectacles (*spek'-tah-ls*) [*spectaculum*, a show, from *spectare*, to view]. A pair of lenses set in a frame, and used for the correction of optic defects of the eye.

Spectral (*spek'-tral*) [*spectrum*, image]. Pertaining to a spectrum.

Spectrometer (*spek-trom'-et-er*) [*spectrum*; *μέτρον*, measure]. An instrument for determining the deviation of a ray of light produced by a prism or diffraction-grating, or for ascertaining the wave-length of a ray of light.

Spectrometry (*spek-trom'-et-re*) [*spectrum*, image; *μέτρον*, measure]. The use of the spectrometer.

Spectroscope (*spek'-tro-sköp*) [*spectrum*, an image; *σκοπεῖν*, to see]. An instrument for the production and examination of the spectrum.

Spectroscopic (*spek-tro-sköp'-ik*) [*spectrum*, image; *σκοπεῖν*, to view]. Pertaining to the spectroscope.

Spectrum (*spek'-trum*) [L., an image]. 1. The band of rainbow-colors produced by decomposing light by means of a prism or a diffraction-grating. 2. An after-image, or ocular spectrum. **S.-analysis**, determination of the nature of bodies by the character of their spectra.

Speculum (*spek'-u-lum*) [L.]. 1. A mirror. 2. An instrument for dilating the opening of a cavity of the body in order that the interior may be more easily visible, as vaginal S., rectal

S., nasal S., etc. S., Sims', a vaginal S. invented by J. Marion Sims.

Speech [AS., *sprecan*, to speak]. 1. The faculty of expressing thought by spoken words; the act of speaking. 2. The words spoken. S.-center, the cerebral center for speech. See *Center*.

Sperm, Sperma (*spur'-mah*) [σπέρμα, seed]. The semen. S.-cell, a spermatoblast.

Spermaceti (*sper-mas-e'-le*) [σπέρμα, seed; κίτος, whale], $C_{18}H_{31}O_2$, $C_{18}H_{33}$. A white, semitransparent substance (Cetaceum, U. S. P., B. P.), consisting of a mixture of various fats of which cetyl palmitate, $C_{18}H_{33}(C_{18}H_{31}O_2)$, is the most important. It is obtained from the head of the sperm-whale, and is used internally as an emollient and as an ingredient of various ointments. Ceratum cetacei (U. S. P.) and Unguentum cetacei (B. P.) are prepared from it.

Spermatic (*spur'-mat'-ik*) [σπέρμα, seed]. 1. Pertaining to the semen. 2. Conveying the semen, as, *e. g.*, the S. cord. 3. Pertaining to the S. cord, as, *e. g.*, the S. fascia.

Spermatin (*spur'-mat-in*) [σπέρμα, seed]. An odorless, mucin-like substance found in semen.

Spermato- (*spur'-mat'-o-*) [σπέρμα, seed]. A prefix meaning pertaining to the semen.

Spermatoblast (*spur'-mat-o-blast*) [σπέρμα, seed; βλαστός, sprout]. A cell resulting from the division of the spermatogenic cell and developing into a spermatozoon.

Spermatocoele (*spur'-mat-o-sel*) [σπέρμα, seed; κύλη, tumor]. A spermatic cyst or encysted hydrocele containing spermatozoa.

Spermatocystitis (*spur'-mat-o-sis-ti'-tis*) [σπέρμα, seed; κύστις, cyst; ιτις, inflammation]. Inflammation of the seminal vesicles.

Spermatocyte (*spur'-mat-o-sit*) [σπέρμα, seed; κύτος, cell]. The germinal cell from which the spermatozoon develops.

Spermatogenesis, Spermatogeny (*spur'-mat-o-jen'-es-is*, *spur'-mat-og'-en-e*) [σπέρμα, seed; γεννάν, to produce]. The formation of spermatozoa.

Spermatogenic (*spur'-mat-o-jen'-ik*) [σπέρμα, seed; γεννάν, to beget]. Producing spermatozoa, as, *e. g.*, the S. cells of the testicle.

Spermatogonium (*spur'-mat-o-go'-ne-um*) [σπέρμα, seed; γονή, generation]. A formative seminal cell or mass of spermatoblasts.

Spermatophore (*spur'-mat-o-för*) [σπέρμα, seed; φέρειν, to bear]. 1. The part of the spermospore that is not converted into a spermatoblast. 2. A semitransparent capsule surrounding a group of spermatozoa.

Spermatorrhea (*spur'-mat-or-e'-ah*) [σπέρμα, seed; ροία, a flow]. Involuntary discharge of semen without sexual excitement.

Spermospore (*spur'-mat-o-spör*) [σπέρμα,

seed; σπόρος, a seed]. A primitive cell giving rise by division to spermatoblasts.

Spermatozoid (*spur'-mat-o-so'-id*). See *Spermatozoon*.

Spermatozoon (*spur'-mat-o-so'-on*) [σπέρμα, semen; ζών, animal]. The male element capable of fecundating the ovum. It consists of an oval head, and a long, mobile cilium or tail. It is the essential element of the semen.

Spermaturia (*spur'-mat-u'-re-ah*) [σπέρμα, seed; urina, urine]. The presence of semen in the urine.

Spermin (*spur'-min*) [σπέρμα, seed], C_2H_8N . A nonpoisonous base obtained from sputum, human semen, the organs of leukemic patients, etc. It has been used in neurasthenia, senile debility, diabetes mellitus, and pulmonary tuberculosis. S. phosphate constitutes the Charcot-Leyden crystals.

Spermoblast (*spur'-mo-blast*). See *Spermatoblast*.

Spermoplasm (*spur'-mo-plasm*) [σπέρμα, seed; πλάσσειν, to mold]. The protoplasm of the spermatoblast.

Spermosphere (*spur'-mo-sfēr*) [σπέρμα, seed; σφαίρα, sphere]. A mass of spermatoblasts.

Spermospore (*spur'-mo-spör*). See *Spermatozoon*.

Sphacelate, Sphacelated (*sfas'-el-at*, *sfas'-el-a-ted*) [σφάκελος, dead]. Necrosed; gangrenous; mortified.

Sphacelation (*sfas'-el-a'-shun*) [σφάκελος, dead]. The formation of a sphacelus; moist gangrene.

Sphaceloderma (*sfas'-el-o-der'-mah*) [σφάκελος, dead; δέρμα, skin]. Gangrene of the skin, especially symmetric gangrene, or Raynaud's disease.

Sphacelus (*sfas'-el-us*) [σφάκελος, gangrene]. A slough.

Sphaerobacteria (*sfe-ro-bak-te'-re-ah*). See *Spherobacteria*.

Spheno- (*sfe'-no-*) [σφήν, wedge]. A prefix denoting pertaining to the sphenoid bone.

Sphenobasilar (*sfe-no-bas'-il-ar*) [σφήν, wedge; βάσις, base]. Pertaining conjointly to the sphenoid bone and the basilar portion of the occipital bone.

Sphenoccipital (*sfe-nok-sip'-it-al*) [σφήν, wedge; occiput, occiput]. Pertaining to the sphenoid and the occipital bone.

Sphenocephalus (*sfe-no-sef'-al-us*) [σφήν, wedge; κεφαλή, head]. A variety of monster in which the two eyes are well separated, the ears united under the head, the jaws and mouth distinct, and the sphenoid bone altered in shape, so that it is analogous in form to what is found normally in birds.

Sphenoethmoid (*sfe-no-eth'-moid*) [σφήν, wedge; ἔθμός, sieve; εἶδος, like]. Relating to both the sphenoid and ethmoid bones.

Sphenoid (*sfe'-noid*) [*σφήν*, wedge; *ῥίδος*, like]. Wedge-shaped, as the S. bone, or S.

Sphenoido- (*sfe-noid'-do-*) [*σφήν*, wedge; *ειδος*, like]. Pertaining to the sphenoid bone.

Sphenoidofrontal (*sfe-noid-do-frun'-tal*) [*σφήν*, wedge; *frons*, forehead]. Pertaining to the sphenoid and frontal bones.

Sphenoidoparietal (*sfe-noid-do-par-i'-et-al*) [*σφήν*, wedge; *paries*, wall]. Pertaining to the sphenoid and parietal bones.

Sphenomaxillary (*sfe-no-mak'-il-a-re*) [*σφήν*, wedge; *maxilla*, maxilla]. Pertaining to the sphenoid and maxillary bones, as, e. g., the S. fossa.

Sphenoccipital (*sfe-no-ok-sip'-it-al*). See *Sphenoccipital*.

Sphenoorbital (*sfe-no-or'-bit-al*) [*σφήν*, wedge; *orbita*, orbit]. Pertaining to the sphenoid bone and the orbit.

Sphenopalatine (*sfe-no-pal'-at-in*) [*σφήν*, wedge; *palatum*, palate]. Pertaining to the sphenoid bone and the palate, as, e. g., the S. foramen.

Sphenoparietal (*sfe-no-par-i'-et-al*) [*σφήν*, wedge; *paries*, wall]. Pertaining to the sphenoid and parietal bones.

Sphenopetrosal (*sfe-no-pe-tros'-sal*) [*σφήν*, wedge; *πέτρα*, rock]. Pertaining to the sphenoid bone and the petrous portion of the temporal bone.

Sphenotic (*sfe-not'-ik*) [*σφήν*, wedge; *ὅς*, ear]. A part of the sphenoid bone, existing as a distinct bone in the fetus, and forming the parts adjacent to the carotid groove.

Sphenotribe (*sfe-no-trīb*) [*σφήν*, wedge; *τρίβειν*, to rub]. An instrument for crushing the basal portion of the fetal skull.

Spheric, Spherical (*sfer'-ik*, *sfer'-ik-al*). Having the shape of or pertaining to a sphere.

S. Aberration. See *Aberration*.

Spherobacteria (*sfe-ro-bak-te'-re-ah*) [*σφαῖρα*, sphere; *βακτηριον*, a rod]. The micrococci.

Spherometer (*sfe-rom'-et-er*) [*σφαῖρα*, sphere; *μέτρον*, a measure]. An instrument for determining the degree of curvature of a sphere or part of a sphere, especially of optic lenses, or of the tools used for grinding them.

Sphincter (*sfinck'-ter*) [*σφίγγειν*, to bind]. A muscle surrounding and closing an orifice, as, e. g., the anal S., the pyloric S., etc.

Sphincterotomy (*sfinck-ter-ot'-o-me*) [*sphincter*; *τομή*, a cutting]. The operation of incising a sphincter.

Sphingoin (*sphing'-go-in*) [*σφίγγειν*, to bind]. A leukomatin derived from cerebral tissue.

Sphygmie (*sfig'-mik*) [*σφυγμός*, pulse]. Pertaining to the pulse.

Sphygmo- (*sfig'-mo-*) [*σφυγμός*, pulse]. A prefix signifying pertaining to the pulse.

Sphygmochronograph (*sfig-mo-kro'-no-graf*) [*σφυγμός*, pulse; *χρόνος*, time; *γράφειν*, to write]. A form of sphygmograph.

Sphygmogram (*sfig'-mo-gram*) [*σφυγμός*, pulse; *γράφειν*, to write]. The tracing made by the sphygmograph.

Sphygmograph (*sfig'-mo-graf*) [*σφυγμός*, pulse; *γράφειν*, to write]. An instrument for recording graphically the features of the pulse and the variations in blood pressure.

Sphygmographic (*sfig-mo-graf'-ik*) [*σφυγμός*, pulse; *γράφειν*, to write]. Pertaining to the sphygmograph.

Sphygmology (*sfig-mol'-o-je*) [*σφυγμός*, pulse; *λόγος*, treatise]. The branch of medicine dealing with the characters of the pulse.

Sphygmomanometer (*sfig-mo-man-om'-et-er*) [*σφυγμός*, pulse; *μάνος*, rare; *μέτρον*, measure]. An instrument for measuring the tension of the blood-current.

Sphygmometer (*sfig-mom'-et-er*). See *Sphygmograph*.

Sphygmophone (*sfig'-mo-fon*) [*σφυγμός*, pulse; *φωνή*, sound]. A sphygmograph in which the vibrations of the pulse produce a sound.

Sphygmoscope (*sfig'-mo-skop*) [*σφυγμός*, pulse; *σκοπεῖν*, to examine]. An instrument for showing the movements of the heart or the pulsations of a blood-vessel.

Sphygm systole (*sfig-mo-sist'-to-le*) [*σφυγμός*, pulse; *συστολή*, contraction]. That part of the sphygmogram produced under the influence of the cardiac systole upon the pulse.

Sphyrotomy (*sfi-rot'-o-me*) [*σφαῖρα*, malleus; *τομή*, a cutting]. An operation consisting in the removal of the malleus or its handle, together with a portion of the membrana tympani.

Spica (*spi'-kah*) [L.]. A spike. **S.-bandage.** See *Bandage*.

Spicule (*spik'-ul*) [dim. of *spica*, a spike]. A minute, sharp-pointed body, as a S. of bone.

Spider-cells. The cells of the neuroglia. See *Deiters' Cells*.

Spigelia (*spi-je'-le-ah*) [after Adrian van der Spiegel, a Belgian physician]. Pink-root, a genus of plants of the order Loganiaceæ. The rhizome and rootlets of *S. marilandica* (S., U. S. P.) contain a volatile alkaloid, spigelin, and are used as an anthelmintic against the round worm. Dose gr. x-xx (0.65-1.3) for a child; ʒj-ij (4.0-8.0) for an adult. Dose of Extractum spigeliæ fluidum (U. S. P.) ʒx-xx (0.65-1.3) for a child; ʒj-ij (4.0-8.0) for an adult.

Spigelian Lobe (*spi-je'-le-an*) [after Adrian van der Spiegel, a Belgian physician]. A small lobe of the liver projecting from the back part of the inferior surface of the right lobe.

Spikenard (*spik'-nard*). A name given to the rhizome of various species of *Valeriana*.

Spiloma (*spi-lo'-mah*). See *Nævus vascularis*.

Spiloplaxia (*spi-lo-plaks'-e-ah*) [σπίλος, spot; πλάξ, a broad surface]. A synonym of *Leprosy*.

Spina (*spi'-nah*) [L.]. Spine. **S. bifida**, a protrusion of the spinal membranes through a congenital cleft of the lower part of the vertebral column. **S. cervina**. See *Buck-thorn*.

Spinal (*spi'-nal*) [*spina*, spine]. 1. Pertaining to the spine. 2. Pertaining to the S. cord. **S. Accessory Nerve**. See *Nerves*, *Table of*. **S. Canal**. See *Canal*, *Vertebral*. **S. Column**, the vertebral column, composed of vertebrae, intervertebral cartilages, and ligaments. **S. Cord**, the neural structure occupying the vertebral canal and extending from the atlas to the first lumbar vertebra, and terminating in the filum terminale. It is covered by the **S. membranes** (the pia mater, arachnoid, and dura mater) and is divided into symmetric halves by the anterior and posterior median fissures. These halves are joined together by the anterior white commissure and the gray commissure. In the middle of the latter is the central canal, a continuation of the ventricular cavities of the brain. Each half of the S. cord consists of an internal mass of gray matter and an outer covering of white matter. The former is subdivided into the anterior and posterior horns, which are made up of ganglion-cells, nerve-fibers, and delicate fibrillae, and a modified neuroglia, the substantia gelatinosa. The white matter is divided by the two gray horns into three columns: the anterior, lateral, and posterior. These are again subdivided into distinct physiologic tracts. Thus the anterior column includes the direct pyramidal tract (Türk's column) and the anterior ground-bundle, or anterior radicular zone, which is continuous with the adjacent part of the lateral column. In the latter the following tracts are distinguished: the crossed pyramidal, direct cerebellar, anterolateral (Gowers' tract), and mixed lateral tract. The posterior column contains the postero-median tract (Goll's column) and the posterolateral or posteroexternal tract (Burdach's column). The spinal cord is the conductor of impulses from and to the brain, as well as a center for reflex acts. **S. Epilepsy**. See *Epilepsy*. **S. Irritation**, a form of neurasthenia characterized by pain in the back, tenderness along the spines of the vertebrae, fatigue on slight exertion, and occasionally numbness and tingling in the limbs. **S. Nerves**, the 31 pairs of nerves arising from the S. cord, and grouped into 8 *Cervical*, 12 *Dorsal*, 5 *Lumbar*, 5 *Sacral*, 1 *Coccygeal*. Each arises by two roots, a dorsal (gangliated) root and a ventral (anterior, nongangliated) root. On the dorsal root is the

spinal ganglion. Beyond the ganglion the two roots unite to form, in the spinal canal, the mixed trunk of a spinal nerve. The anterior roots supply efferent fibers to all the voluntary muscles of the trunk and extremities, to the smooth muscular fibers of the bladder, ureter, uterus, etc., vasomotor, inhibitory, secretory, and trophic fibers. The posterior roots carry afferent impulses.

Spinalis (*spi-na'-lis*) [*spina*, spine]. 1. Spinal. 2. A muscle attached to the spinous processes of the vertebrae. See *Muscles*, *Table of*.

Spinant (*spi'-nant*) [*spina*, spine]. A drug or other agent increasing the reflex excitability of the spinal cord; strychnin is a spinant.

Spindle-tree. See *Euonymus*.

Spine (*spin*) [*spina*, a thorn]. 1. A sharp process of bone. 2. The back-bone or spinal column.

Spinobulbar (*spi-no-bul'-bar*) [*spina*, spine; *bulbus*, the medulla oblongata]. Pertaining to the spinal cord and the medulla oblongata.

Spinoneural (*spi-no-nu'-ral*) [*spina*, spine; νεύρον, nerve]. Pertaining to the spinal cord and the peripheral nerves.

Spinous (*spi'-nus*) [*spina*, spine]. Resembling or pertaining to a spine, as, *e. g.*, the S. process of the tibia or of a vertebra.

Spintherism (*spin'-ther-izm*) [σπινθηρίζειν, to emit sparks]. The sensation of sparks dancing before the eyes.

Spiral (*spi'-ral*) [*spira*, a spire]. 1. Winding like the threads of a screw, as, *e. g.*, a S. bandage. 2. A curve having a S. course.

Spirem (*spi'-rem*) [*spira*, a coil, a twist]. The close skein, or mother skein, of chromatin-fibrils in a cell undergoing mitotic division.

Spirillum (*spi-ril'-um*) [*spirillum*, dim. of *spira*, a coil]. A genus of bacteria having a spiral shape. See *Bacteria*, *Table of*. **S.-fever**. See *Relapsing Fever*.

Spirit (*spir'-it*) [*spiritus*, breath, from *spirare*, to breathe]. 1. The soul. 2. An alcoholic solution of a volatile substance. See *Spiritus*. 3. Alcohol. **S., Corn-**, whisky obtained by the distillation of corn. **S., Potato-**, whisky obtained by the distillation of potatoes. **S., Proof-**, diluted alcohol (Alcohol dilutum, U. S. P.; Spiritus tenuior, B. P.) containing about 41 per cent. by weight of absolute ethyl alcohol. **S., Rectified**, (Spiritus rectificatus, B. P.), containing 16 per cent. of water.

Spirituos (*spir'-it-u-us*) [*spiritus*, spirit]. Alcoholic; pertaining to alcoholic liquors.

Spiritus (*spir'-it-us*) [L.]. See *Spirit*. **S. frumentii**, whisky, a spirit obtained by the distillation of fermented grain. **S. chloroformi** is used as a carminative. Dose ℥ x-lx (0.65-4.0). **S. juniperi**, gin or whisky

with which juniper-berries and hops have been distilled. **S. odoratus**, cologne-water. **S. vini gallici**, brandy; a liquor obtained by the distillation of wine. For other varieties of Spiritus, see the different drugs.

Spirobacteria (*spi-ro-bak-te'-re-ah*) [*spira*, a spiral; *βακτηριον*, bacterium]. Spiral bacteria.

Spirocheta, Spirochæta (*spi-ro-ke'-tah*) [*spira*, a coil; *χούτη*, a bristle]. A genus of bacteria characterized by flexible, spiral filaments. See *Bacteria, Table of*.

Spirograph (*spi'-ro-graf*) [*spirare*, to breathe; *γράφειν*, to write]. An instrument for registering the movements of respiration.

Spirometer (*spi-rom'-et-er*) [*spirare*, to breathe; *μέτρον*, a measure]. An instrument for measuring the quantity of air taken in and given out in forcible respiration.

Spirometry (*spi-rom'-et-re*) [*spirare*, to breathe; *μέτρον*, measure]. The determination by means of the spirometer of the amount of air taken in and given out during respiration.

Spirophore (*spi'-ro-for*) [*spirare*, to breathe; *φέρειν*, to carry]. An instrument for performing artificial respiration.

Spirulina (*spi-ro-liv'-nah*) [*spirula*, from *spira*, a coil]. A spiral microorganism of spindle shape.

Spissated (*spis'-a-ted*). See *Inspissated*.

Spissitude (*spis'-it-ud*) [*spissare*, to thicken]. The state of being inspissated.

Spittle (*spit'-l*). See *Saliva*.

Splanchnic (*splangk'-nik*) [*σπλάγχνα*, viscera]. Pertaining to or supplying the viscera. **S. Nerves**, three nerves, the great, lesser, and least, or renal S., derived from the sympathetic system.

Splanchno- (*splangk'-no-*) [*σπλάγχνα*, viscera]. A prefix denoting pertaining to the viscera.

Splanchnography (*splangk-nog'-ra-fe*). See *Splanchnology*.

Splanchnology (*splangk-nol'-o-je*) [*σπλάγχνα*, viscera; *λόγος*, treatise]. The branch of anatomy treating of the viscera.

Splanchnopleure (*splangk'-no-plur*) [*σπλάγχνα*, viscera; *πλευρά*, the side]. The visceral layer of mesoderm forming the covering of the digestive tube.

Splanchnoscopy (*splangk-nos'-ko-pe*) [*σπλάγχνα*, viscera; *σκοπεῖν*, to examine]. Visual examination of the viscera.

Splanchnoskeleton (*splangk-no-skel'-et-on*) [*σπλάγχνα*, viscera; *σκελετόν*, skeleton]. That portion of the skeleton related to the viscera.

Splanchnotomy (*splangk-nol'-o-me*) [*σπλάγχνα*, viscera; *τέμνειν*, to cut]. Dissection of the viscera.

Splay-foot. See *Talipes*.

Spleen (*splen*) [*σπλήν*, spleen]. One of the abdominal viscera, situated just below the diaphragm on the left side, and connected with the hematopoietic system. It is covered by a fibroelastic capsule from which trabeculae radiate into the organ. In the spaces formed by these are found collections of lymphoid tissue (the Malpighian corpuscles) and the splenic pulp. The Malpighian corpuscles surround the small branches of the splenic artery. The splenic pulp consists of a delicate reticulum containing large connective-tissue cells, lymphoid cells, and red corpuscles. The spleen receives a large amount of blood, which in passing from the termination of the splenic artery to the beginning of the splenic vein is probably not held within walls, but comes in direct contact with the lymphoid tissue. The S. normally weighs about 200 grams. **S., Accessory**, a detached portion of splenic tissue in the neighborhood of the spleen. **S., Floating**. See *S., Wandering*. **S. Sago-**, one of which the Malpighian follicles are the seat of amyloid change. **S., Wandering**, one that is movable, owing to relaxation of its attachments.

Splen- (*splen-*) [*σπλήν*, spleen]. A prefix denoting pertaining to the spleen.

Splenadenoma (*splen-ad-en-o'-mah*) [*σπλήν*, spleen; *άδην*, gland; *όμα*, tumor]. Hyperplasia of the lymphoid tissue of the spleen.

Splenalgia (*splen-al'-je-ah*) [*σπλήν*, spleen; *άλγος*, pain]. Neuralgic pain in the spleen.

Splenculus (*splen'-ku-lus*) [*σπλήν*, spleen]. An accessory spleen.

Splendid Line of Haller. See *Lines, Table of*.

Splenectomy (*splen-ek'-to-me*) [*σπλήν*, spleen; *εκτομή*, excision]. Excision of the spleen.

Splenemia (*splen-e'-me-ah*) [*σπλήν*, spleen; *αἷμα*, blood]. Splenic leukemia.

Splenic (*splen'-ik*) [*σπλήν*, spleen]. Pertaining to or affecting the spleen. **S. Apoplexy**, **S. Fever**. See *Anthrax*.

Splenification (*splen-if-ik-a'-shun*) [*σπλήν*, spleen; *ποιεῖν*, to make]. See *Splenization*.

Splenified (*splen-if-id'*) [*σπλήν*, spleen; *ποιεῖν*, to make]. Of a tissue, resembling the tissue of the spleen, as, e. g., S. bone-marrow.

Splenitis (*splen-i'-tis*) [*σπλήν*, spleen; *τις*, inflammation]. Inflammation of the spleen.

Splenium (*sple'-ne-um*) [*σπληνιον*, bandage]. 1. A bandage. 2. The rounded posterior extremity of the corpus callosum.

Splenius (*sple'-ne-us*) [*σπληνιον*, bandage]. Shaped like a splenium, as, e. g., the S. muscle or simply splenius. See *Muscles, Table of*.

Splenization (*splen-i-a'-shun*) [*σπλήν*, spleen]. The change in an organ, especially the lung, produced by congestion, whereby it comes to resemble the tissue of the spleen.

Spleno- (*splen'-o-*). Same as *Splen-*.

Splenocele (*splen'-o-sel*) [*σπλήν*, spleen; *κήλη*, tumor]. Hernia of the spleen.

Splenic (*splen-o-kol'-ik*) [*σπλήν*, spleen; *κόλον*, colon]. Pertaining to the spleen and the colon.

Splenomalacia (*splen-o-mal'-a'-se-ah*) [*σπλήν*, spleen; *μαλακία*, softness]. Softening of the spleen.

Splenomegalia, Splenomegaly (*splen-o-meg-a'-le-ah*, *splen-o-meg'-al-e*) [*σπλήν*, spleen; *μέγας*, large]. Enlargement of the spleen, especially simple enlargement of the spleen without leukemia; by some it is considered merely as Hodgkin's disease of splenic type.

Splenopathia (*splen-o-path'-e-ah*). Splenopathy. *S. leukocythæmica*, splenic leukocythemia.

Splenopathy (*splen-op'-ath-e*) [*σπλήν*, spleen; *πάθος*, suffering]. Any disease of the spleen.

Splenopexis (*splen-o-peks'-is*) [*σπλήν*, spleen; *πέζις*, a fixing in]. Fixation of a wandering spleen to the abdominal wall by means of sutures.

Splenophrenic (*splen-o-fren'-ik*) [*σπλήν*, spleen; *φρήν*, diaphragm]. Pertaining to the spleen and the diaphragm.

Splenopneumonia (*splen-o-nu-mo'-ne-ah*) [*σπλήν*, spleen; *πνεύμων*, lung]. Pneumonia with splenization of the lung.

Splenotomy (*splen-ot'-o-me*) [*σπλήν*, spleen; *τέμνειν*, to cut]. 1. The operation of incising the spleen. 2. Dissection of the spleen.

Splint [Swedish, *splinte*, a kind of spike]. A piece of wood, metal, or other material for keeping the ends of a fractured bone or other movable parts in a state of rest.

Spodogenous (*spo-doj'-en-us*) [*σποδός*, ashes; *γεννάν*, to produce]. Pertaining to or produced by waste-material, as, *e. g.*, *S.* enlargement of the spleen, a swelling of the spleen produced by the accumulation of the detritus of red corpuscles.

Spondyl-, Spondylo- (*spon'-dil-, spon'-dil-o-*) [*σπόνδυλος*, vertebra]. A prefix denoting pertaining to a vertebra.

Spondylarthritis (*spon-dil-ar-thri'-tis*) [*σπόνδυλος*, vertebra; *άρθρον*, joint; *ιτις*, inflammation]. Inflammation of a vertebral articulation.

Spondylitis (*spon-dil-i'-tis*) [*σπόνδυλος*, a vertebra; *ιτις*, inflammation]. Inflammation of a vertebra, especially Tuberculous *S.*, or Pott's disease.

Spondylizema (*spon-dil-i-se'-mah*) [*σπόνδυλος*, vertebra; *ἰζημα*, a subsiding]. The settling of a vertebra into the place of a subjacent one that has been destroyed.

Spondylolisthesis (*spon-dil-o-lis-the'-sis*) [*σπόνδυλος*, vertebra; *όλισθησις*, a slipping]. Deformity of the spinal column produced by the gliding forward of the lumbar vertebrae in such a manner that they overhang the brim

and obstruct the inlet of the pelvis; especially the separation of the last lumbar vertebra from, and its slipping forward on, the sacrum.

Spondylolisthetic (*spon-dil-o-lis-the'-ik*) [*σπόνδυλος*, vertebra; *όλισθησις*, a slipping]. Pertaining to or caused by spondylolisthesis.

Spondylotomy (*spon-dil-ot'-o-me*) [*σπόνδυλος*, vertebra; *τομή*, a cutting]. See *Rachiotomy*.

Sponge (*spunj*) [*σπόγγος*]. A marine animal of the class Porifera, having a porous, horny skeleton; also the skeleton itself, used as an absorbent. *S.-bath*, the application of water to the surface of the body by means of a *S. S.-graft*. See *Graft*. *S.-tent*. See *Tent*.

Spongiiform (*spun'-je-form*) [*σπόγγος*, sponge; *forma*, a form]. Resembling a sponge.

Spongin (*spun'-jin*) [*σπόγγος*, sponge]. The horny substance forming the skeletal fibers of the sponge.

Spongioblast (*spun'-je-o-blast*) [*σπόγγος*, a sponge; *βλαστός*, a germ]. A variety of cell derived from the ectoderm of the embryonic neural tube, and forming later the neuroglia.

Spongiopilin (*spun-je-o-pi'-lin*) [*σπόγγος*, sponge; *πίλος*, felt]. Felted or woven cloth, into which tufts of sponge are incorporated and one side of which is coated with rubber; it is used as a poultice.

Spongioplasm (*spun'-je-o-plasm*) [*σπόγγος*, sponge; *πλάσσειν*, to mold]. The fine, elastic, protoplasmic threads forming the reticulum of cells.

Spongy (*spun'-je*) [*σπόγγος*, sponge]. Having the texture of sponge. *S. Portion of the Urethra*, that contained in the corpus spongiosum of the urethra.

Spontaneous (*spon-ta'-ne-us*) [*spons*, will]. Voluntary; occurring without extraneous impulse; automatic.

Spoon. An instrument consisting of an oval or circular bowl fixed to a handle; it is used in surgery to scrape away dead tissue, granulations, etc.

Sporadic (*spor-ad'-ik*) [*σποραδικός*, scattered]. Scattered; occurring in an isolated manner. *S. Cholera*, cholera morbus.

Spore (*spor*) [*σπόρος*, seed]. A seed; a reproductive body of lower vegetable organisms, as of bacteria.

Sporiferous (*spor-if'-er-us*) [*σπόρος*, seed; *ferre*, to bear]. Spore-bearing.

Sporoblast (*spor'-o-blast*) [*σπόρος*, seed; *βλαστός*, germ]. One of the four round bodies produced by the process of endogenous cell-formation in a coccidium.

Sporocyst (*spor'-o-sist*) [*σπόρος*, seed; *κίστις*, a bag]. The mother-cell of a spore.

Sporogenesis (*spor-o-jen'-es-is*) [*σπόρος*,

seed; *γενεαίς*, generation]. The development of spores; reproduction by spores.

Sporogony (*spor-og'ō-nē*). See *Sporogenesis*.

Sporophore (*spor'ō-for*) [*σπόρος*, seed; *φέρειν*, to bear]. That portion of a fungus bearing the spores.

Sporozoa (*spor-ō-zō'ah*) [*σπόρος*, seed; *ζῶον*, an animal]. A class of parasitic Protozoa including the Gregarinidia, parasitic in various worms and arthropods; the Coccidia, or oviform psorosperms, parasitic in the hepatic and intestinal epithelium of various mammals, including man; the Saracsporidia, or tubuliform psorosperms, parasitic in the muscles of various animals; the Myxosporidia, the psorosperms of fishes; the Microsporidia, the psorosperms of articulates.

Sporulation (*spor-u-lā'shun*) [*σπόρος*, spore]. The production of spores. **S.**, **Arthrogenous**, the change of bacteria into resistant forms, which are capable of germinating again under favorable conditions.

Spot. See *Macula*. **S.**, **Blind**. See *Blind Spot*.

Spotted Fever. 1. Cerebrospinal fever. 2. Typhus.

Sprain (*sprān*) [OF., *espreindre*, from L., *exprimere*, to press out]. A wrenching of a joint producing a stretching or laceration of the ligaments. **S.-fracture**, an injury in which a tendon is torn from its attachment together with a shell of bone.

Spray (*sprā*). A liquid blown into minute particles by a strong current of air.

Spruce (*sprūs*). See *Picea*.

Sprue (*sprā*). Synonym of *Thrush*.

Spunk (*spunk*). See *Agaricus*.

Spurge (*spurj*). See *Eupatorium*.

Spurious (*spu're-us*) [*spurius*, false]. False.

Spurred Rye. See *Ērgot*.

Sputum (*spu'tum*) [*spuere*, to spit; *pl.*, *Spula*]. The secretion ejected from the mouth in spitting. It consists of saliva and mucus from the nasal fossa and the fauces. In diseased conditions of the air-passages or lungs it may be purulent, mucopurulent, fibrinous, or bloody. **S. crudum**, the scanty, viscid expectoration of the early stages of acute bronchitis. **S.**, **Nummular**, a sputum characterized by round, coin-like masses; it is seen in pulmonary tuberculosis. **S.**, **Rusty**, the dark-colored sputum of lobar pneumonia, the color of which is due to the admixture of blood.

Sputum-septicemia (*spu-tum-sep-tē-sē'mē-ah*). See *Septicemia*.

Squama (*skwa'mah*) [L.]. A scale or scale-like mass, as, e. g., the S. of the temporal bone.

Squamoparietal (*skwa-mo-par'ē-et'al*) [*squama*, scale; *paries*, wall]. Pertaining to the squamous portion of the temporal bone.

Squamosal (*skwa-mō'al*) [*squama*, scale]. The squamous portion of the temporal bone.

Squamosphenoid (*skwa-mo-sfe'noid*) [*squama*, scale; *sphenoid*]. Pertaining to the squamous portion of the temporal bone and to the sphenoid bone.

Squamotemporal (*skwa-mo-tem'pō-rāl*) [*squama*, scale; *temporal*]. Pertaining to the squamous portion of the temporal bone.

Squamous (*skwa'mus*) [*squamosus*, scaly]. 1. Of the shape of a scale, as, e. g., the S. portion of the temporal bone. 2. Scaly. **S.**, **Suture**, the suture between the squamous portion of the temporal bone and the frontal and parietal bones.

Squamozygomatic (*skwa-mō-zī-gō-mat'ik*) [*squama*, scale; *zygomatic*]. Pertaining to the squamous and zygomatic portions of the temporal bone.

Square Lobe. 1. The lobus quadratus of the liver. 2. A lobe on the upper surface of the cerebellar hemisphere.

Squarrous (*skwar'us*) [*ισχάρα*, a scab]. Scurfy.

Squill (*skwil*). The bulb of *Urginea maritima* (U. S. P.), or *Urginea scilla* (B. P.), of the order Liliaceae. Squill (Scilla, U. S. P., B. P.) contains several bitter principles, *scillitin*, *scillipicrin*, *scillitoxin*, and *scillin*; a carbohydrate, *sinistrin*, and other substances. S. is expectorant, diuretic, and emetocathartic, and is used in dropsy and in croup. Dose gr. j-ij (0.065-0.13). Preparations and doses: Acetum scillæ (U. S. P., B. P.), m_x -xxx (0.65-2.0); Extractum scillæ fluidum (U. S. P.), m ij-ijj (0.13-0.20); Oxy-mel scillæ (B. P.), f $\frac{3}{4}$ j-ij (4.0-8.0); Pilula scillæ composita (B. P.), gr. v-x (0.32-0.65); Syrupus scillæ (U. S. P., B. P.), f $\frac{3}{4}$ ss-j (2.0-4.0); Syrupus scillæ compositus, hivesyrup, m xx-xxx (1.3-2.0); Tinctura scillæ, (U. S. P., B. P.), m v-xxx (0.32-2.0).

Squint (*skwint*) [origin obscure]. See *Strabismus*.

Squirting Cucumber. See *Elaterium*.

Sr. Symbol for strontium.

Ss. Abbreviation for *semis*, one-half.

Stabile (*sta'bīl*) [*stabilis*, from *stare*, to stand]. Not moving; fixed. **S. Current**, an electric current produced by holding the electrodes in a fixed position.

Staccato Speech. See *Scanning Speech*.

Stadium (*sta'de-um*). [L.]. Stage. **S. acmes**, the height of a disease. **S. augmenti**, the period in which there is increase in the intensity of the disease. **S. caloris**, the period during which there is fever; the hot stage. **S. decrementi**, defervescence of a febrile disease; the period in which there is a decrease in the severity of the disease.

Staff. An instrument for passing into the bladder through the urethra and used as a guide in operations for stricture.

Stage (*stāj*) [*stare*, to stand]. 1. A definite period of a disease characterized by certain symptoms; a condition in the course of a disease. 2. The horizontal plate projecting from the pillar of a microscope, for supporting the slide or object. **S.**, **Algid**, a condition characterized by subnormal temperature, feeble, flickering pulse, various nervous symptoms, etc. It occurs in cholera and other diseases marked by exhausting intestinal discharges. **S.**, **Amphibolic**, the S. of a disease intervening between its height and its decline. **S.**, **Asphyxial**, the preliminary S. of Asiatic cholera, marked by extreme thirst, muscular cramps, etc., due to loss of water from the blood. **S.**, **Cold**, the rigor or chill of an attack of a malarial paroxysm. **S.**, **Expulsive** (*of labor*), the stage which begins when dilatation of the cervix uteri is complete and during which the child is expelled from the uterus. See *Labor*. **S.**, **First** (*of labor*), that stage in which the molding of the fetal head and the dilatation of the cervix are affected. **S.**, **Hot**, the febrile S. of a malarial paroxysm. **S.** of **Invasion**, the period in the course of a disease in which the system comes under the influence of the morbid agent. **S.** of **Latency**, the incubation-period of an infectious disease, or that period intervening between the entrance of the virus and the manifestations of the symptoms to which it gives rise. **S.**, **Placental** (*of labor*), the period occupied by the expulsion of the placenta and fetal membranes. **S.**, **Preeruptive**, the period of an eruptive fever following infection and prior to the appearance of the eruption. **S.**, **Second** (*of labor*). See *S.*, *Expulsive*. **S.**, **Sweating**, the third or terminal S. of a malarial paroxysm, during which there is sweating. **S.**, **Third** (*of labor*). See *S.*, *Placental*.

Stain (*stān*) [from *distain*, from *dis*, priv.; *tingere*, to color]. 1. A discoloration. 2. A pigment employed in microscopy to color the tissues or to produce certain reactions. The common microscopic stains are hematoxylin, carmin, osmic acid, and the anilin dyes.

Stammer (*stām'-er*) [AS., *stamur*, stammering]. To speak interruptedly or with hesitation.

Standard (*stan'-dard*) [*extendere*, to spread out]. Something used for comparison. **S.**-**solution**, a solution containing a definite quantity of a reagent.

Stannic (*stan'-ik*) [*stannum*, tin]. Containing tin as a tetrad element.

Stannous (*stan'-us*) [*stannum*, tin]. Containing tin as a dyad element.

Stannum (*stan'-um*) [L.]. See *Tin*.

Stapedectomy (*sta-pe-dek'-to-me*) [*stapes*; *ektomē*, excision]. Excision of the stapes.

Stapediovestibular (*sta-pe-de-o-ves-tib'-u-lar*) [*stapes*; *vestibulum*, vestibule]. Relating to the stapes and the vestibule.

Stapedius (*sta-pe'-de-us*) [*stapes*]. See *Muscles*, *Table of*.

Stapes (*sta'-pes*) [L., a stirrup]. The stirrup-shaped bone of the middle ear, articulating with the incus and the fenestra ovalis.

Staphisagria (*staf-is-ag'-re-ah*) [*σταφίς*, dried grape; *ἀγριος*, wild]. The Delphinium staphisagria, of the order Ranunculaceæ. The seeds (S., U. S. P., Staphisagria semina, B. P.) contain the alkaloids delphinin, $C_{27}H_{37}NO_3$, delphinoidin, $C_{42}H_{68}N_2O_7$, delphinin, $C_{37}H_{46}N_2O_4$, and staphisagrin, $C_{23}H_{33}NO_3$. S. has been used locally as an application in rheumatism, and as an ointment to destroy lice and itch-mites.

Staphylinopharyngeus (*staf-il-i-no-far-in'-je-us*) [*σταφυλή*, uvula; *φάρυγξ*, pharynx]. The palatopharyngeus. See *Muscles*, *Table of*.

Staphylinus (*staf-il-i'-nus*) [*σταφυλή*, uvula]. 1. Palatal. 2. See *S. medius*. **S. externus**, the tensor palati. **S. internus**, the levator palati. **S. medius**, the azygos uvulæ muscle. See *Muscles*, *Table of*.

Staphylitis (*staf-il-i'-tis*) [*σταφυλή*, uvula; *ιτις*, inflammation]. Inflammation of the uvula.

Staphylo- (*staf'-il-o-*) [*σταφυλή*, uvula]. A prefix denoting pertaining to the uvula.

Staphylococcemia (*staf-il-o-kok-se'-me-ah*) [*σταφυλή*, grape; *κόκκος*, a berry; *αἷμα*, blood]. The presence of staphylococci in the blood.

Staphylococcus (*staf-il-o-kok'-us*) [*σταφυλή*, grape; *κόκκος*, a berry]. See *Bacteria*, *Table of*.

Staphyloma (*staf-il-o'-mah*) [*σταφυλή*, grape; *δμα*, tumor]. A bulging of the cornea or sclera of the eye. **S.**, **Anterior**. See *Keratoglobus*. **S. cornæ**, a bulging of the cornea, due to a thinning of the membrane, with or without previous ulceration. **S.**, **Posterior**, **S. posticum**, a backward bulging of the sclerotic coat at the posterior pole of the eye.

Staphyloplasty (*staf'-il-o-plas-te*) [*σταφυλή*, uvula; *πλάσσειν*, to mold]. A plastic operation for the closure of cleft palate.

Staphylorrhaphy (*staf-il-or'-a-fe*) [*σταφυλή*, uvula; *ράφή*, suture]. Suture of a cleft palate.

Staphylotomy (*staf-il-o'-o-me*) [*σταφυλή*, uvula; *τομή*, a cutting]. 1. The operation of incising or removing the uvula. 2. The operation of incising a staphyloma.

Star-anise. See *Illicium*.

Starch (*starch*) [AS., *starc*, stiff; L., *Amylum*]. A carbohydrate widely distributed in the vegetable kingdom, occurring in peculiar concentrically marked granules or grains. When heated with water the granules swell

up, burst, partially dissolve, and form S. paste. The soluble portion is called granulose, the insoluble is cellulose. Iodin produces a characteristic blue coloration with S. The most important varieties of S. are: potato-S., leguminous S., wheat S. (*Amylum*, U. S. P., B. P.), sago-S., rice-S. S., **Animal**. See *Glycogen*. S., **Corn-**, a highly nutritive S. obtained from Indian corn. S.-**enema**, an enema consisting of starch-water. S., **Iodized**, starch that has been acted upon by iodine. S.-**water**, a mixture of wheat-S. and water, used chiefly as an emollient enema.

Star-grass. See *Aletris*.

Stars of Verheyn (*fär-hin'*). The star-shaped figures formed by the stellate veins of the kidney, beneath the capsule.

Starvation (*star-vat'-shun*) [AS., *steorfan*, to die]. Deprivation of food; the state produced by deprivation of food.

Stasis (*sta'-sis*) [*στάσις*, from *istānai*, to stand]. A standstill of the current of any of the fluids of the body, especially of the blood.

Stas-Otto Method. A method of extracting alkaloids or ptomaines from tissues. It depends upon the facts that the salts of the alkaloids are soluble in water and in alcohol, and generally insoluble in ether, while the free alkaloids are soluble in ether, and may be removed from alkaline fluids by agitation with ether. The method consists in treating the mass with 90 per cent. alcohol and tartaric or oxalic acid; digesting the whole for some time at about 70° C. and filtering; the filtrate is evaporated at a temperature not exceeding 35° C., and the residue taken up with absolute alcohol, filtered, and evaporated again at a low temperature. The residue is dissolved in water, alkalinized, and agitated with ether. After separation, the ether is removed and allowed to evaporate. The residue may be further purified by redissolving in water and again extracting with ether.

Static (*stat'-ik*) [*στατικός*, from *istānai*, to stand]. At rest; pertaining to matter or force at rest or in equilibrium. S. **Breeze**, a method of administering static electricity, consisting in the withdrawal of a static charge from a patient by means of a pointed electrode. S. **Electricity**. See *Electricity*.

Statics (*stat'-iks*) [*στατική*, from *istānai*, to stand]. The science relating to forces in a condition of equilibrium.

Station (*sta'-shun*) [*statio*, from *stare*, to stand]. Standing position or attitude.

Stationary (*sta'-shun-a-ree*) [*statio*, from *stare*, to stand]. Standing still; not moving. S. **Air**, the amount of air which is constantly in the lungs during normal respiration.

Statometer (*stat-om'-et-er*) [*στασίς*, standing; *μέτρον*, measure]. An instrument for measuring the degree of exophthalmus.

Status (*sta'-tus*) [L.]. A state. S. **arthriticus**, the nervous manifestations preceding an attack of gout. S. **epilepticus**, a condition in which epileptic attacks occur in rapid succession, the patient not regaining consciousness during the interval. S. **præsens**, the state of a patient at the time of examination. S. **typhosus**. See *Typhoid State*.

Stauroplegia (*stau-ro-ple'-je-ah*) [*σταυρός*, crossed; *πληγή*, a stroke]. Crossed hemiplegia.

Stavesacre (*stär-a'-ker*). See *Staphisagria*.

Steapsin (*ste-ap'-sin*) [*στέας*, fat]. A ferment assisting in the saponification of the fats and found in the pancreatic juice.

Stearate (*ste'-ar-at*) [*στέας*, fat]. A salt of stearic acid. Glycerol S. is called stearin, *q. v.*

Stearic Acid (*ste-ar'-ik*) [*στέας*, fat]. See *Acid*, *Stearic*, and *Stearin*.

Stearin (*ste'-ar-in*) [*στέας*, fat], $C_{18}H_{35}O_2$ ($C_{17}H_{33}O_2$). A compound of stearic acid and glyceryl occurring in the harder animal fats, especially in tallow. It crystallizes in white, pearly scales.

Stearopten (*ste-ar-op'-ten*) [*στέας*, fat; *πτερός*, winged, volatile]. The crystalline substance occurring naturally in solution in a volatile oil.

Stearrhea (*ste-ar-e'-ah*) [*στέας*, fat; *ῥοία*, a flow]. See *Schorrhæa*.

Steatite (*ste'-at-it*). See *Talcum*.

Steato- (*ste'-at-o-*) [*στέας*, fat]. A prefix meaning fatty.

Steatocele (*ste-at-o-sel*) [*στέας*, fat; *κύημα*, tumor]. A swelling formed by the collection of fatty matter in the scrotum.

Steatoma (*ste-at-o-mah*) [*στέας*, fat; *δύα*, tumor]. 1. A sebaceous cyst. 2. A lipoma.

Steatopygia (*ste-at-o-pij'-e-ah*) [*στέας*, fat; *πτεγή*, buttock]. Enormous fatness of the buttock, common among the women of some African tribes.

Steatopygous (*ste-at-op'-ig-us*) [*στέας*, fat; *πτεγή*, buttock]. Characterized by excessive development of the buttocks.

Steatorrhea (*ste-at-or-e'-ah*) [*στέας*, fat; *ῥοία*, flow]. See *Schorrhæa*.

Steatosis (*ste-at-o'-sis*) [*στέας*, fat]. 1. Fatty degeneration. 2. An abnormal accumulation of fat.

Steatozoon (*ste-at-o-zo'-on*) [*στέας*, fat; *ζῷον*, an animal]. The parasite, *Demodex folliculorum*, contained in comedones.

Steel. A form of iron holding an intermediate position between cast iron and wrought iron and containing from 0.5 to 1.5 per cent. of carbon. S., **Tincture of**, *tinctura ferri chloridi*; see under *Ferrum*. S., **Wine of**. Synonym of *Vinum ferri*, under *Ferrum*.

Stellate (*stel'-at*) [*stellā*, a star]. Star-shaped.

Stellulæ Verheynei. See *Stars of Verheyen*.
Stellwag's Symptom. See *Signs and Symptoms, Table of*.

Steno- (*sten'-o-*) [*στενός*, narrow]. A prefix meaning narrow or constricted.

Stenocardia (*sten-o-kar'-de-ah*) [*στενός*, narrow; *καρδία*, heart]. Angina pectoris.

Stenocephalous (*sten-o-sef'-al-us*) [*στενός*, narrow; *κεφαλή*, head]. Having a head narrow in one or more of its diameters.

Stenocephaly (*sten-o-sef'-al-e*) [*στενός*, narrow; *κεφαλή*, head]. Narrowing of the head in one or more of its diameters.

Stenochoria (*sten-o-ko'-re-ah*) [*στενός*, narrow; *χώρας*, space]. Narrowing; stenosis.

Stenopeic (*sten-o-pe'-ik*) [*στενός*, narrow; *ὄπαιος*, pierced]. Pertaining to or having a narrow slit; applied to lenses that allow the passage of rays only through a narrow slit.

Steno's Duct. See under *Duct*.

Stenosis (*sten-o'-sis*) [*στενός*, narrow]. Constriction or narrowing, especially of a channel or aperture, as, *e. g.*, aortic S., mitral S., etc.

Stenotic (*sten-o'-ic*) [*στενός*, narrow; *ὄπαιος*, pierced]. Pertaining to or having a narrow slit; applied to lenses that allow the passage of rays only through a narrow slit.

Stensen's Duct. See *Duct*.

Stensen's Experiment. Compression of the abdominal aorta of an animal, so as to cut off the blood-supply to the lumbar region of the spinal cord. It leads to rapid paralysis of the posterior portion of the body.

Stensen's Foramen. See *Foramina, Table of*.

Stephanion (*stef'-an'-e-on*) [*στέφανος*, a wreath or crown]. See *Cranimetric Points*.

Steppage-gait (*step'-aj-gat*). The peculiar high-stepping gait seen in tabes dorsalis and certain forms of multiple neuritis.

Stercobilin (*ster-ko-bi'-lin*) [*stercus*, dung; *bilis*, bile]. A coloring-matter found in feces, and identical with hydrobilirubin.

Stercoraceous (*ster-ko-ra'-shus*) [*stercus*, dung]. Fecal; having the nature of feces; containing feces, as, *e. g.*, S. vomiting.

Stercoral (*ster'-ko-ral*). See *Stercoraceous*.

Stercoremia (*ster-ko-re'-me-ah*) [*stercus*, dung; *αἷμα*, blood]. A condition of the blood resulting from arrest of intestinal excretion and the absorption of toxic matters from the feces.

Stercorin (*ster'-ko-rin*) [*stercus*, dung]. An extractive from the feces resembling cholesterol.

Stercus (*ster'-kus*) [*I.*, dung]. Feces.

Stere (*ster*) [*στερεός*, solid]. A measure of 1000 liters; a kiloliter.

Stereometry (*ste-re-om'-et-re*) [*στερεός*, solid; *μέτρον*, measure]. The measurement of the capacity of a hollow body.

Stereophantoscope (*ste-re-o-fan'-to-skop*)

[*στερεός*, solid; *φανός*, visible; *σκόπειν*, to see]. See *Stereophoroscope*.

Stereophoroscope (*ste-re-o-for'-o-skop*) [*στερεός*, solid; *φέρειν*, to carry; *σκοπεῖν*, to see]. A stereoscopic zoetrope, an instrument for producing a series of images apparently in motion.

Stereoplasm (*ste'-re-o-plasm*) [*στερεός*, solid; *πλάσσειν*, to mold]. The solid part of the protoplasm of cells.

Stereoscope (*ste'-re-o-skop*) [*στερεός*, solid; *σκοπεῖν*, to see]. An instrument by which two similar pictures of the same object are made to overlap so that the reflected images are seen as one, thereby giving the appearance of solidity and relief.

Stereoscopic (*ste-re-o-skop'-ik*) [*στερεός*, solid; *σκοπεῖν*, to view]. Pertaining to stereoscopy. **S.** Vision, binocular vision. See *Stereoscopy*.

Sterile (*ster'-il*) [*sterilis*, barren]. 1. Not fertile; not capable of reproducing. 2. Free from microorganisms or spores.

Sterility (*ster-il'-it-e*) [*sterilis*, sterile]. The condition of being sterile.

Sterilization (*ster-il-i-za'-shun*) [*sterilis*, barren]. The act of rendering anything sterile; the destruction of microorganisms, particularly by means of heat. **S.**, Intermittent, a method of sterilization in which an interval of time is allowed to elapse between the several heatings, giving an opportunity for any spores present to develop into adult microorganisms, in which form they readily succumb to the action of heat.

Sterilized (*ster'-il-izd*) [*sterilis*, sterile]. Rendered sterile.

Sterilizer (*ster'-il-i-zer*) [*sterilis*, sterile]. An apparatus for destroying the microorganisms attached to an object, especially by means of heat.

Sternal (*stur'-nal*) [*sternum*; *στέρνον*, sternum]. Pertaining to the sternum.

Sterno- (*stur'-no-*) [*sternum*; *στέρνον*, sternum]. A prefix denoting connection with the sternum.

Sternoclavicular (*stur-no-kla-vik'-u-lar*) [*sternum*, sternum; *clavicula*]. Pertaining to the sternum and the clavicle.

Sternocleidomastoid (*stur-no-cli-do-mas'-toid*) [*στέρνον*, sternum; *κλείς*, key; *mas-toid*]. Pertaining to the sternum, the clavicle, and the mastoid process, as, *e. g.*, the S. muscle. See *Muscles, Table of*.

Sternocostal (*stur-no-kos'-tal*) [*sternum*, sternum; *costa*, a rib]. Pertaining to the sternum and the ribs.

Sternohyoid (*stur-no-hi'-oid*) [*στέρνον*, sternum; *hyoid*]. Pertaining to the sternum and the hyoid bone, as, *e. g.*, the S. muscle. See *Muscles, Table of*.

Sternomastoid (*stur-no-mas'-toid*) [*στέρνον*,

sternum; mastoid]. Pertaining to the sternum and the mastoid process.

Sternopagus (*stur-nop'-ag-us*) [*στέρον*, sternum; *πάγος*, fastened]. A double monster the parts of which are united by the sternum.

Sternothyroid (*stur-no-thi'-roid*) [*στέρον*, sternum; *thyroid*]. Pertaining to the sternum and the thyroid cartilage, as, e. g., the S. muscle. See *Muscles*, Table of.

Sternotracheal (*stur-no-tra'-ke-al*) [*στέρον*, sternum; *trachea*]. Pertaining to the sternum and the trachea.

Sternum (*ster'-num*) [L.; *στέρον*, breast-bone]. The flat, narrow bone in the median line in the front of the chest, composed of three portions, the manubrium, the gladiolus, and the ensiform or xiphoid appendix.

Sternutation (*stur-nu-ta'-shun*) [*sternutatio*, a sneezing]. The act of sneezing.

Sternutatory (*stur-nu'-tat-o-ry*) [*sternutare*, to sneeze]. 1. Producing sneezing. 2. An agent that causes sneezing.

Stertor (*stur'-tor*) [L., a snoring]. Sonorous breathing, or snoring; the rasping, rattling sound produced when the larynx and the air-passages are obstructed by mucus.

Stertorous (*stur'-to-rus*) [*stertere*, to snore]. Characterized by stertor, as, e. g., S. breathing.

Stetho- (*steth'-o*) [*στήθος*, chest]. A prefix denoting pertaining to the chest.

Stethograph (*steth'-o-graf*) [*στήθος*, chest; *γράφειν*, to write]. An instrument recording the respiratory movements of the chest.

Stethometer (*steth-om'-et-er*) [*στήθος*, chest; *μέτρον*, measure]. An instrument for measuring the degree of expansion of the chest.

Stethophonometer (*steth-o-fo-nom'-et-er*) [*στήθος*, chest; *φωνή*, sound; *μέτρον*, measure]. An instrument for measuring the phenomena elicited by auscultation.

Stethoscope (*steth'-o-skop*) [*στήθος*, breast; *σκοπεῖν*, to view]. An instrument for ascertaining the condition of the organs of circulation and respiration by the sounds made by these organs. It consists of a hollow tube, one end being placed over the locality to be examined, the other at the ear of the examiner. The binaural S. consists of a Y-shaped tube, the flexible branches being applied each to an ear of the listener.

Stethoscopic (*steth-o-skop'-ik*) [*στήθος*, chest; *σκοπεῖν*, to view]. Pertaining to or detected by means of the stethoscope.

Stethoscopy (*steth-as'-ko-pe*) [*στήθος*, chest; *σκοπεῖν*, to view]. Examination with the aid of the stethoscope.

Sthenic (*sthen'-ik*) [*σθένος*, strength]. Strong, active. **S. Fever**, a form of fever marked by high temperature, quick and tense pulse, and highly colored urine.

Stibium (*stib'-e-um*). See *Antimonium*.

Sticking Plaster. Resin-plaster.

Stiff-neck Fever. Epidemic cerebrospinal meningitis.

Stigma (*stig'-mah*) [*στίγμα*, point]. A small spot or mark, especially a spot of hemorrhage in the palm or sole corresponding to the nail-marks of Christ, occurring in hysteric persons. 2. Any one of the marks or signs characteristic of a condition; generally used in the plural, as *hysteric stigmata*. 3. That part of a pistil which receives the pollen.

Stigmata maydis. See *Zen mays*.

Stigmatic (*stig-mat'-ik*) [*στίγμα*, spot]. Pertaining to a stigma.

Stigmatization (*stig-mat-i-sa'-shun*) [*στίγμα*, point]. The formation of stigmata.

Stilet, **Stilette** (*sti-let'*) [Fr., dim. of *stilus*, a point]. 1. A small, sharp-pointed instrument enclosed in a cannula. 2. A wire passed into a flexible catheter.

Still-born. Born lifeless.

Still-birth. The birth of a dead child.

Stillicidium (*stil-is-id'-e-um*) [*stillar*, a drop; *cadere*, to fall down]. The flow of a liquid drop by drop. **S. lacrimarum**, overflow of tears from obstruction of the canaliculus or nasal duct; epiphora. **S. narium**, coryza.

Stilling's Canal. See *Canal of Stilling*.

Stillingia (*stil-in'-je-ah*) [after Benjamin Stillingfleet, an English botanist]. A genus of plants of the order Euphorbiaceae. The root of *S. sylvatica*, queen's root or queen's delight (S., U. S. P.), is used as an alternative in syphilis, scrofula, diseases of the skin, etc. Dose of Extractum stillingiae fluidum (U. S. P.) fʒ ss-j (2.0-4.0).

Stimulant (*stim'-u-lant*) [*stimulus*, a goad].

1. Stimulating. 2. An agent that causes stimulation. **S., Cardiac**, one that increases the heart's action. **S., Cerebral**, one that exalts the action of the cerebrum. **S., Diffusive**, one that has a prompt but transient effect. **S., Hepatic**, one that excites the activity of the liver.

Stimulate (*stim'-u-lat*) [*stimulare*, to stimulate]. To quicken; to stir up; to excite; to increase functional activity.

Stimulation (*stim-u-la'-shun*) [*stimulus*, a goad]. 1. The act of stimulating. 2. The effect of a stimulant.

Stimuline (*stim'-u-len*) [*stimulus*, a goad]. A substance capable of stimulating the phagocytic action of cells.

Stimulus (*stim'-u-lus*) [L.]. A goad; an impulse; anything capable of causing stimulation. **S., Adequate**. See *S., Homologous*. **S., Chemic**, one due to or produced by chemic means. **S., Heterologous**, one acting upon the nervous elements of the sensory apparatus along their entire course. **S., Homologous**, one acting only upon the end-organ. **S., Mechanic**, one acting by mechanic means, as, e. g., pinching or striking.

Stirrup, Stirrup-bone (*stir' - up*). The stapes.

Stitch. 1. A sudden, sharp, lancinating pain. 2. See *Suture*.

Stoma (*sto' - mah*) [*στόμα*, mouth]. 1. A mouth. 2. A pore, as that between endothelial cells, establishing direct communication between adjacent lymph-channels.

Stomach (*stum' - ak*) [*στόμαχος*]. The most dilated part of the alimentary canal, situated below the diaphragm in the left hypochondriac, the epigastric, and part of the right hypochondriac regions. It is connected at one end (cardiac end) with the esophagus, at the other (pyloric end) with the duodenum. Its wall consists of four coats—the serous, muscular, submucous, and mucous. The mucous coat contains the gastric glands (cardiac and pyloric glands), which secrete the gastric juice and mucus. **S.-cough**, a reflex cough excited by irritation of the stomach. **S.-pump**, a pump for withdrawing the contents of the stomach. **S.-tooth**, a lower canine tooth, especially one of the first dentition. **S.-tube**, a flexible tube for irrigation or evacuation of the stomach.

Stomachal (*stum' - ak - al*) [*στόμαχος*, the stomach]. Pertaining to the stomach.

Stomachic (*stum - ak' - ik*) [*στόμαχος*, stomach]. 1. Pertaining to the stomach. 2. Stimulating the secretory activity of the stomach. 3. An agent having this property.

Stomatitis (*stom - at - itis*) [*στόμα*, mouth; *ιτις*, inflammation]. Inflammation of the mouth. **S. aphthosa**, **S., Aphthous**. See *Aphthæ*. **S., Catarrhal**, a simple form characterized by swelling of the mucous membrane, pain, and salivation. **S., Gangrenous**. See *Cancrum oris*. **S., Mercurial**, that arising from poisoning by mercury. **S., Mycotic**. See *Thrush*. **S., Scorbutic**, that due to scurvy. **S., Ulcerative**, a form characterized by the formation of small ulcers on the cheeks, lips, and tongue, with copious salivation, pain, fetid breath, slight fever, and at times great prostration.

Stomato- (*stum' - at - o*) [*στόμα*, mouth]. A prefix meaning pertaining to the mouth.

Stomatomycosis (*stom - at - o - mi - ko' - sis*) [*στόμα*, mouth; *μύκης*, fungus]. A disease of the mouth due to fungi, especially the *oidium albicans*.

Stomatonecrosis, Stomatonomia (*stom - at - o - ne - kro' - sis*, *stom - at - on' - o - mah*). See *Cancrum oris*.

Stomatopathy (*stom - at - op' - ath - e*) [*στόμα*, mouth; *πάθος*, disease]. Any disease of the mouth.

Stomatoplasty (*stom' - at - o - plas - te*) [*στόμα*, mouth; *πλάσσειν*, to form]. A plastic operation upon the mouth.

Stomatoscope (*stum' - at - o - skop*) [*στόμα*,

mouth; *σκοπεῖν*, to inspect]. An instrument for inspecting the cavity of the mouth.

Stomocephalus (*stom - o - sef' - al - us*) [*στόμα*, mouth; *κεφαλή*, head]. A variety of monster in which there is the same deformity as in rhinocephalus or in cyclocephalus, associated with a defect of the maxillary bones, so that the skin hangs in folds around the mouth.

Stomodæum (*stom - o - de' - um*) [*στόμα*, mouth; *δαίειν*, to divide]. The primitive oral cavity of the embryo, formed by a depression of the ectoderm and afterward forming the mouth and upper part of the pharynx.

Stone. A hardened mass of mineral matter. See *Calculus*. **S., Blue**, copper-sulphate crystals. **S., Gall**-, a biliary calculus.

Stool. The evacuation of the bowels.

Stop-needle. A lance-pointed needle used in the operation of discission, having an enlargement or shoulder upon the shank to prevent too deep penetration.

Storax (*sto' - raks*). See *Syrax*.

Strabismal, Strabismic (*stra - bis' - mal*, *strabis' - mik*) [*στραβισμός*, from *στραβός*, crooked]. Relating to or affected with strabismus.

Strabismus (*strab - is' - mus*) [*στραβισμός*, from *στραβός*, crooked]. Squint; that abnormality of the eyes in which the visual axes do not meet at the desired objective point, in consequence of incoordinate action of the external ocular muscles. **S., Alternating**, one in which either eye fixes alternately. **S., Concomitant**, one in which the squinting eye has full range of movement. **S., Convergent**, one in which the squinting eye is turned to the nasal side. **S., Divergent**, one in which the squinting eye is turned to the temporal side. **S., External**. See *S., Divergent*. **S., Internal**. See *S., Convergent*. **S., Paralytic**, that due to paralysis of one or more muscles. **S., Spastic**, that due to a spastic contraction of an ocular muscle.

Strain (*strān*) [OF., *estraindre*, from L., *stringere*, to draw tight]. 1. Excessive stretching; overuse of a part. 2. The condition produced in a part by overuse, as, *e. g.*, eye-strain. 3. To overexert; to use to excess; to make violent efforts.

Strait (*strāt*) [Fr., *étroit*, from *strictus*, drawn tight]. A narrow or constricted passage, as, *e. g.*, the inferior or superior S. of the pelvis. **S.-jacket**, a strong jacket placed on the insane or delirious to prevent injury to themselves or to others.

Stramonium (*stra - mo' - ne - um*). The *Datura stramonium*, Jamestown or Jimson weed, a plant of the order Solanaceæ. The leaves (*Stramonii folia*, U. S. P., B. P.) and seeds (*Stramonii semen*, U. S. P., *Stramonii semina*, B. P.) contain two alkaloids, daturin, identical with atropin, and hyoscyamin. The ac-

tion of *S.* resembles that of belladonna. It is used in asthma, dysmenorrhea, neuralgia, rheumatism, and pains of syphilitic origin. In asthma the leaves may be smoked in a tobacco-pipe. Locally *S.* is employed as an ointment or cataplasm in irritable ulcers and inflamed surfaces. Dose of the seeds gr. j (0.065); of the leaves gr. ij-ijj (0.13-0.20). Extractum stramonii seminis, U. S. P. (Extractum stramonii, B. P.). Dose gr. $\frac{1}{6}$ - $\frac{1}{4}$ (0.010-0.016). Extractum stramonii seminis fluidum, U. S. P. Dose \mathfrak{m} j- \mathfrak{m} ij (0.065-0.13). Tinctura stramonii seminis, U. S. P. Dose \mathfrak{m} v-xxx (0.32-2.0).

Strangling. See *Strangulation*.

Strangulated (*strang'-gu-la-ted*) [*strangulare*, to strangle]. 1. Choked. 2. Compressed so that the circulation is arrested, as, *e. g.*, *S.* hernia.

Strangulation (*strang'-gu-la'-shun*) [*strangulare*, to choke]. 1. The act of choking. 2. Constriction of a part producing arrest of the circulation, as, *e. g.*, *S.* of a hernia.

Strangury (*strang'-gu-re*) [*σπάγγε*, a drop; *οὐρον*, urine]. Painful urination, the urine being voided drop by drop.

Strap. 1. A long band, as of adhesive plaster. 2. To compress a part by means of bands, especially bands of adhesive plaster.

Stratification (*strat'-e-fik-a'-shun*) [*stratum*, a layer; *facere*, to make]. Arrangement in layers.

Stratified (*strat'-e-fid*) [*stratum*, a layer; *facere*, to make]. Arranged in layers.

Stratiform (*strat'-e-form*) [*stratum*, stratum; *forma*, form]. Formed into a layer. *S.* Fibrocartilage, fibrocartilage lining bony grooves through which the tendons of muscles pass.

Stratum (*stra'-tum*) [*L.*, from *sternere*, to strew]. A layer. *S.* corneum, *S.* granulosum, *S.* lucidum, *S.* Malpighii. See *Skin*.

Strawberry-tongue. The characteristic tongue of scarlet fever, in which the vessels of the fungiform papillae become turgid, causing the papillae to stand out as red points, in marked contrast with the thick coating of fur on the filiform papillae.

Streak (*strek*). A furrow, line, or stripe. *S.* Medullary. See *Medullary Groove*. *S.* Primitive, an opaque band extending some distance forward from the posterior margin of the area pellucida and forming the first noticeable sign of the development of the blastoderm. *S.* Meningitic. See *Tache cérébrale*.

Strengthening Plaster. Emplastrum roburans, or iron-plaster.

Strephotome (*stref'-o-tōm*) [*σπρίφειν*, to twist; *τέμνειν*, to cut]. An instrument shaped like a corkscrew, used to secure union in the operation for the radical cure of hernia.

Streptobacteria (*strep'-to-bak'-te'-re-ah*) [*σπείρτης*, twisted; *βακτηριον*, bacterium]. Short, rod-shaped bacteria forming chains.

Streptococcemia (*strep'-to-kak'-se'-ne-ah*) [*streptococcus*; *αἷμα*, blood]. A state of the blood characterized by the presence of streptococci.

Streptococcus (*strep'-to-kak'-kus*) [*σπείρτης*, twisted; *κόκκος*, a kernel]. See *Bacteria*, Table of. *S.-angina*, angina due to streptococci.

Streptothrix (*strep'-to-thriks*) [*σπείρτης*, twisted; *θρίξ*, the hair]. See *Cladethrix*, under *Bacteria*, Table of.

Stretcher (*strech'-er*). A cot or litter for carrying the sick.

Stria (*stri'-ah*) [*L.*]. A streak or white line. **Acoustic or Auditory Striæ**, transverse white lines on the lower part of the floor of the fourth ventricle, which unite with the auditory nerve-roots. **Striæ gravidarum**, the atrophic striæ observed upon the abdomen in pregnant women. **Striæ longitudinales**, long, slightly elevated lines on the upper surface of the corpus callosum.

Striate, Striated (*stri'-at*, *stri'-a-ted*) [*stria*, a furrow]. Striped. *S.* Body, the corpus striatum. *S.* Muscle. See *Afuscle*.

Striation (*stri'-at'-shun*) [*stria*, furrow]. 1. The state of being striated. 2. A striated structure.

Stricture (*strikt'-tūr*) [*strictura*, from *stringere*, to draw tight]. A narrowing of a canal from external pressure, or as a result of inflammation or other changes in its walls. *S.* Cicatricial, a *S.* due to cicatricial tissue. *S.* Functional. See *S.*, *Spasmodic*. *S.* Impermeable, or *S.* Impassable, one not permitting the passage of an instrument. *S.* Irritable, one in which the passage of an instrument causes great pain. *S.* Organic, one due to structural changes in or about a canal. *S.* Spasmodic, one due to muscular spasm.

Stricturotome (*strikt'-tu-ro-tōm*) [*strictura*, stricture; *τέμνειν*, to cut]. An instrument for dividing a stricture.

Stricturotomy (*strikt'-tu-rol'-o-mē*) [*strictura*, stricture; *τέμνειν*, to cut]. The operation of incising a stricture.

Stridor (*stri'-dor*) [*stridere*, to make a creaking sound]. A peculiar, harsh, vibrating sound produced during expiration.

Stridulous (*strid'-u-lus*) [*stridere*, to make a creaking sound]. Characterized by stridor. *S.* Laryngismus. See *Laryngismus*.

Strobilus (*strob'-il-us*) [*strobilus*, a pine-cone]. The tape-worm.

Stroboscope (*strob'-o-skop*). See *Zootrope*. **Stroboscopic** (*strob'-o-skop'-ik*) [*στροβίλος*, a twisting; *σκοπεῖν*, to view]. Pertaining to the stroboscope.

Stroke (*strôk*). A sudden attack; the word is used colloquially for apoplexy and paralysis.

Stroma (*stro'-mah*) [*στρώμα*, a bed]. The tissue forming the framework for the essential part of an organ.

Strongylus (*stron'-jil-us*) [*στρογγύλος*, round]. A genus of nematode worms found in the lower animals, and occasionally in man.

Strontium (*stron'-she-um*) [after *Strontian*, in Scotland]. A metallic element belonging to the group of alkaline earths. It has a specific gravity of 2.5, an atomic weight of 87.3, and a valence of two. Symbol Sr. It combines with oxygen to form S. oxid, or strontia, SrO. S. bromid, SrBr₂, 6H₂O (Strontii bromidum, U. S. P.), has been used in epilepsy, diabetes, gastrectasis, rheumatoid arthritis, and lithemia. Dose gr. xv-xxx (1.0-2.0). S. iodid, SrI₂, 6H₂O (Strontii iodidum, U. S. P.), is used like the other oxids. S. lactate, Sr(C₂H₃O₃)₂·3H₂O (Strontii lactas, U. S. P.), is used in nephritis, albuminuria, rheumatism, and gout. Dose gr. x-xxx (0.65-2.0).

Strophanthus (*strof'-an'-thus*) [*στροφῶς*, a twisted band; *ἄνθος*, flower]. A genus of plants of the order Apocynaceæ, some of the species of which are used for the preparation of arrow poison in Africa. The seeds of S. hispidus (S., U. S. P., B. P.) contain a crystalline glucosid, strophanthin, and an alkaloid, inein. S. is a muscle-poison, but in small doses is a cardiac and perhaps a vascular stimulant. It is used in the same cases as digitalis. Dose of Tinctura strophanthi (U. S. P., B. P.) $\text{m}\bar{\text{v}}$ -xv (0.32-1.0); of strophanthin gr. $\frac{3}{16}$ - $\frac{1}{8}$ (0.0002-0.0003).

Strophulus (*strof'-u-lus*) [*στροφῶς*, a twisted band]. Red gum, tooth-rash; a form of miliaria occurring in infants. S. pruriginosus, an eruption occurring in children, and characterized by disseminated, intensely itching papules.

Structural (*struk'-tu-ral*) [*structura*, structure]. Pertaining to or affecting the structure.

Struma (*stru'-mah*) [L.]. 1. Scrofula. 2. Goiter.

Strumiprivus (*stru'-me-pri'-vus*) [*struma*; *privus*, deprived of]. Deprived of the thyroid gland; due to the removal of the thyroid gland, as cachexia strumipriva.

Strumous (*stru'-mus*) [*struma*]. Scrofulous.

Strychnin (*strih'-nin*) [*στρογχινος*, strychnin], C₁₁H₁₂N₂O₂. One of the alkaloids of *Nux vomica*. S. sulphate, (C₁₁H₁₂N₂O₂)₂·H₂SO₄·5H₂O (Strychnine sulphas, U. S. P.). Dose of the alkaloid or salt gr. $\frac{1}{8}$ (0.003). For properties and uses see *Nux vomica*.

Strychninism (*strih'-nin-ism*) [*στρογχινος*, strychnin]. The state of being under the influence of strychnin.

Strychnos (*strih'-nos*). A genus of plants of the order Loganiaceæ, several species of which yield strychnin and brucin.

Stump [Icel., *stump*, stump]. The portion of a limb or other part left attached to the body after an amputation.

Stun [AS., *stunian*, to make a din]. To render temporarily insensible, as by a blow.

Stupe (*stüp*) [*slupa*, tow]. A cloth used for applying heat or counterirritation; especially a cloth wrung out of hot water and sprinkled with a counterirritant, as, e. g., turpentine-S.

Stupefacient, Stupefactive (*stu-pe-fa'-shent*, *stu-pe fak'-tiv*) [*stupere*, to be stunned; *facere*, to make]. Narcotic.

Stupor (*stu'-por*) [L.]. A state of partial unconsciousness from which the individual can be roused.

Stuttering (*stut'-er-ing*) [Icel., *stauta*, to stutter]. A hesitation in speech due to an inability to enunciate the syllables without repeated efforts.

Stye (*sti*). See *Hordeolum*.

Style, Stylet (*stil*, *sti-let'*) [*στυλος*, pillar].

1. A probe. 2. A wire inserted into a catheter or cannula, in order to stiffen the instrument or to perforate the tissues. See *Stilet*.

Stylo- (*sti'-lo-*) [*στυλος*, pillar]. A prefix denoting pertaining to the styloid process of the temporal bone.

Styloglossus (*sti-lo-glos'-us*). See *Muscles, Table of*.

Stylohyoid (*sti-lo-hi'-oid*) [*στυλος*, pillar; *hyoid*].

1. Pertaining to the styloid process of the temporal bone and the hyoid bone, as, e. g., the S. muscle. See *Muscles, Table of*. 2. Pertaining to the S. muscle.

Styloid (*sti'-loid*) [*στυλος*, pillar; *ειδος*, like]. Resembling a stylius.

Stylomastoid (*sti-lo-mas'-toid*) [*στυλος*, pillar; *mastoid*]. Pertaining to the styloid and mastoid processes.

Stylomaxillary (*sti-lo-maks'-il-a-re*) [*στυλος*, pillar; *maxilla*, maxilla]. Pertaining to the styloid process and the maxilla.

Stylopharyngeus (*sti-lo-far-in-je'-us*). See *Muscles, Table of*.

Stylus (*sti'-lus*) [L., a stake]. A pointed instrument for making applications.

Styptic (*stip'-tik*) [*στυπτικός*, astringent]. 1. Checking hemorrhage by contracting the blood-vessels. 2. An agent that checks hemorrhage by causing contraction of the blood-vessels.

Styracin (*sti'-ra-sin*). See *Styrax*.

Styrax (*sti'-raks*) [*στυραξ*, storax]. Storax; a balsam obtained from the inner bark of *Liquidambar orientalis*, or oriental sweetgum. It contains a volatile oil, styrol, several resins, an amorphous substance called storesin, cinnamic acid, and styracin (the cinamate of cinnamyl). It is stimulant, ex-

pectorant, and antiseptic, acting like benzoïn and tolu, and is used in bronchial affections and catarrh of the urinary passages. Externally it is an antiseptic and parasiticide. It is a constituent of Friar's balsam. Dose gr. v-xx (0.32-1.3).

Styrone (*sti'-ron*) [*στίραξ*, storax]. Cinnamic alcohol, $C_9H_{10}O$.

Styryl Alcohol (*sti'-ril*) [*στίραξ*, storax]. See *Styrone*.

Sub- [L.]. A prefix denoting under or beneath; in chemistry, a prefix denoting (1) the lower of two compounds of the same elements; (2) denoting a basic salt.

Subacetate (*sub-as'-et-at*) [*sub*, under; *acetum*, vinegar]. A basic acetate.

Subacid (*sub-as'-id*) [*sub*, under; *acidum*, acid]. Moderately acid.

Subacromial (*sub-ak-ro'-me-al*) [*sub*, under; *acromial*]. Below the acromion.

Subacute (*sub-ak-ūt'*) [*sub*, under; *acutus*, sharp]. Moderately acute.

Subanconeus (*sub-an-ko-ne'-us*). See *Muscles*, *Table of*.

Subaponeurotic (*sub-ap-on-u-ro't'-ik*) [*sub*, under; *aponeurosis*]. Beneath an aponeurosis.

Subarachnoid (*sub-ar-ak'-noid*) [*sub*, under; *arachnoid*]. Beneath the arachnoid membrane, as, e. g., the S. space.

Subarcuate (*sub-ar'-ku-at*) [*sub*, under; *arcus*, an arc]. Slightly arcuate.

Subastragalar, **Subastragaloid** (*sub-as-trag'-al-ar*, *sub-as-trag'-al-oid*) [*sub*, under; *astragalus*]. Below the astragalus.

Subcalcarine (*sub-kal-ka-rēn*) [*sub*, under; *calcarine*]. Situated beneath the calcarine fissure, as the S. convolution.

Subcapsular (*sub-kap'-u-lar*) [*sub*, under; *capsula*, capsule]. Beneath a capsule.

Subcarbonate (*sub-ka'-bon-at*) [*sub*, under; *carbo*, charcoal]. A basic carbonate.

Subclavian (*sub-kla'-ve-an*) [*sub*, under; *clavis*, key]. Lying under the clavicle, as, e. g., the S. artery.

Subclavicular (*sub-kla'-rik'-u-lar*) [*sub*, under; *clavis*, key]. Beneath the clavicle.

Subclavius (*sub-kla'-vus*). See *Muscles*, *Table of*.

Subconjunctival (*sub-kon-junk-ti'-val*) [*sub*, under; *conjunctiva*]. Situated beneath the conjunctiva.

Subconsciousness (*sub-kon'-shus-ness*) [*sub*, under; *conscious*, knowing]. Imperfect consciousness; that state in which mental processes take place without the mind being distinctly conscious of its own activity.

Subcoracoid (*sub-ko'-ak-oid*) [*sub*, under; *κόραξ*, a crow; *ὅσος*, like]. Situated below the coracoid process.

Subcortical (*sub-ko'-tik-al*) [*sub*, under; *cortex*, cortex]. Beneath the cortex.

Subcranial (*sub-kra'-ne-al*) [*sub*, under; *κράνιον*, cranium]. Situated beneath the cranium.

Subcrepitant (*sub-krep'-it-ant*) [*sub*, under; *crepitare*, to make a crackling noise]. Almost crepitant, as, e. g., S. rale. See *Rale*.

Subcrureus (*sub-kru'-re-us*). See *Muscles*, *Table of*.

Subculture (*sub-kul'-tūr*) [*sub*, under; *cultura*, culture]. In bacteriology, a secondary culture made from a primary culture.

Subcutaneous (*sub-ku-ta'-ne-us*) [*sub*, under; *cutaneus*, from *cutis*, the skin]. Beneath the skin; hypodermic.

Subcuticular (*sub-ku-tik'-u-lar*) [*sub*, under; *cutis*, skin]. Beneath the epidermis, as, e. g., a S. stitch.

Subdiaphragmatic (*sub-di-a-frag-mat'-ik*) [*sub*, under; *διάφραγμα*, diaphragm]. Under the diaphragm.

Subdural (*sub-du'-ral*) [*sub*, under; *dura*, dura]. Beneath the dura.

Subencephalon (*sub-en-sef'-al-on*) [*sub*, under; *ἐγκεφαλον*, brain]. The medulla oblongata, pons, and corpora quadrigemina taken together.

Subendocardial (*sub-en-do-kar'-de-al*) [*sub*, under; *ἐνδον*, within; *καρδία*, heart]. Beneath the endocardium.

Subepidermal, **Subepidermatic**, **Subepidermic** (*sub-ep-e-der'-mal*, *sub-ep-e-der-mat'-ik*, *sub-ep-e-der'-mik*) [*sub*, under; *ἐπι*, upon; *δέρμα*, skin]. Situated beneath the epidermis.

Subepithelial (*sub-ep-e-the'-le-al*) [*sub*, under; *ἐπι*, upon; *θήλη*, nipple]. Situated under an epithelial surface.

Suberin (*su'-ber-in*) [*suber*, cork]. The impure cellulose forming the cellular tissue of cork.

Subfascial (*sub-fash'-e-al*) [*sub*, under; *fascia*, fascia]. Beneath the fascia.

Subfebrile (*sub-feb'-ril*) [*sub*, under; *febris*, fever]. Slightly febrile.

Subflavous (*sub-fla'-vus*) [*sub*, under; *flavus*, yellow]. Somewhat yellow.

Subgallate (*sub-gal'-lat*). A basic gallate. S. of Bismuth, $BiC_7H_7O_7$. Dermatol, an odorless, yellow powder used as an antiseptic in diseases of the gastrointestinal tract and locally as a substitute for iodoform. Dose gr. v-x (0.32-0.65).

Subglenoid (*sub-gle'-noid*) [*sub*, under; *γλήνη*]. Beneath the glenoid fossa, as, e. g., S. dislocation of the humerus.

Subhyoid (*sub-hi'-oid*) [*sub*, under; *ὑοειδής*, hyoid]. Beneath the hyoid bone.

Subiculum (*sub-ik'-u-lum*) [*subex*, a layer]. The uncinate gyrus.

Subinflammation (*sub-in-flam-a'-shun*) [*sub*, under; *inflammatio*, inflammation]. A slight degree of inflammation.

Subintrant (*sub-in'-trant*) [*subintrare*, to enter secretly]. Entering secretly; applied to malarial fevers in which a new paroxysm begins before the termination of the preceding one.

Subinvolution (*sub-in-vo-lu'-shun*) [*sub*, under; *involutio*, a rolling up]. Imperfect involution. **S. of the Uterus**, the imperfect contraction of the uterus after delivery.

Subjacent (*sub-ja'-sent*) [*sub*, under; *jacere*, to lie]. Lying beneath.

Subject (*sub'-jekt*) [*sub*, under; *jacere*, to throw]. 1. An individual that serves for purposes of experiment or study, or that is under observation or treatment. 2. A cadaver.

Subjective (*sub-jek'-tiv*) [*sub*, under; *jacere*, to throw]. 1. Pertaining to the individual himself. 2. Of symptoms, experienced by the patient himself, and not amenable to physical exploration.

Sublatio (*sub-la'-she-o*) [L.]. Removal. **S. retinæ**, detachment of the retina.

Sublimate (*sub'-lim-at*) [*sublimare*, to lift up high]. A substance obtained by sublimation. **S.**, Corrosive, mercuric chlorid. See *Mercury*.

Sublimation (*sub-lim-a'-shun*) [*sublimare*, to lift up high]. The vaporization and condensation of a volatile solid.

Sublime (*sub-lim*) [*sublimare*, to lift up high]. 1. To subject to sublimation. 2. To undergo sublimation.

Sublingual (*sub-ling'-gwal*) [*sub*, beneath; *lingua*, tongue]. 1. Lying beneath the tongue. 2. Pertaining to the parts lying beneath the tongue.

Sublinguitis (*sub-ling'-gwi'-tis*) [*sub*, under; *lingua*, tongue; *itis*, inflammation]. Inflammation of the sublingual gland.

Sublobular (*sub-lob'-u-lar*) [*sub*, under; *lobulus*, a lobule]. Situated beneath a lobule. **S. Veins**, the radicles of the hepatic veins, situated at the base of a cluster of lobules.

Subluxation (*sub-luks-a'-shun*) [*sub*, under; *luxatio*, luxation]. Incomplete luxation.

Submammary (*sub-mam'-a-re*) [*sub*, under; *mamma*, breast]. Situated beneath the breast.

Submarginal (*sub-mar'-jin-al*) [*sub*, under; *margo*, margin]. Situated near the border or margin.

Submaxillary (*sub-maks'-il-a-re*) [*sub*, under; *maxilla*, maxilla]. 1. Lying beneath the lower maxilla, as, *e. g.*, the **S. gland**. 2. Pertaining to the **S. gland**.

Submental (*sub-men'-tal*) [*sub*, under; *mentum*, chin]. Situated under the chin.

Submerge (*sub-merj'*) [*sub*, under; *mergere*, to dip]. To place under the surface of a liquid.

Submersion (*sub-mer'-shun*) [*sub*, under; *mergere*, to dip]. The act of submerging;

the condition of being under the surface of a liquid.

Submucous (*sub-mu'-kus*) [*sub*, under; *mucosus*, mucous]. Situated beneath a mucous membrane.

Subnasal (*sub-na'-zal*) [*sub*, under; *nasus*, nose]. Situated below the nose. **S. Point**. See *Craniometric Points*.

Subnitrate (*sub-ni'-trat*) [*sub*, under; *nitrum*, niter]. A basic nitrate.

Subnormal (*sub-nor'-mal*) [*sub*, under; *norma*, rule]. Below normal.

Subnucleus (*sub-nu'-kle-us*) [*sub*, under; *nucleus*, nucleus]. Any one of the smaller groups of cells into which a large nerve-nucleus is divided by the passage through it of nerve-bundles.

Suboccipital (*sub-ok-sip'-it-al*) [*sub*, under; *occiput*]. Situated beneath the occiput.

Suboxid (*sub-oks'-id*) [*sub*, under; *ὀξύς*, acid]. One of two oxids containing the less oxygen.

Subparietal (*sub-par-i'-et-al*) [*sub*, under; *paries*, wall]. Situated beneath the parietal bone, convolution, or fissure.

Subpatellar (*sub-pat-el'-ar*) [*sub*, under; *patella*, knee-cap]. Situated beneath the patella.

Subpeduncular (*sub-pe-dung'-ku-lar*) [*sub*, under; *pedunculus*, peduncle]. Situated beneath a peduncle.

Subpericardial (*sub-per-e-kar'-de-al*) [*sub*, under; *περί*, around; *καρδία*, heart]. Situated beneath the pericardium.

Subperiosteal (*sub-per-e-os'-te-al*) [*sub*, under; *περί*, around; *ὀστίον*, bone]. Beneath the periosteum.

Subperitoneal (*sub-per-e-ton-e'-al*) [*sub*, under; *περιτόναιον*, peritoneum]. Beneath the peritoneum.

Subphrenic (*sub-fren'-ik*). Synonym of *Subdiaphragmatic*.

Subpleural (*sub-plu'-ral*) [*sub*, under; *πλευρά*, side]. Beneath the pleura.

Subpontine (*sub-pon'-tin*) [*sub*, under; *pons*, pons]. Beneath the pons.

Subpreputial (*sub-pre-pu'-she-al*) [*sub*, under; *preputium*, prepuce]. Beneath the prepuce.

Subpubic (*sub-pu'-bik*) [*sub*, under; *pubes*, pubis]. Situated beneath the pubic arch or symphysis.

Subpulmonary (*sub-pul'-mon-a-re*) [*sub*, under; *pulmo*, the lung]. On the ventral side of the lungs.

Subretinal (*sub-ret'-in-al*) [*sub*, under; *retina*, retina]. Beneath the retina.

Subsalt. A basic salt.

Subscapular (*sub-skap'-u-lar*) [*sub*, under; *scapula*, the shoulder-blade]. 1. Beneath the scapula, *e. g.*, as the **S. muscle**, or **Subscapularis**. 2. Pertaining to the **S. muscle**.

Subscapularis (*sub-skap-u-lar'is*). See *Mus. sc.*, *Table of*.

Subscription (*sub-skrip'-shun*) [*sub*, beneath; *scribere*, to write]. That part of a prescription containing the directions to the pharmacist, indicating how the ingredients are to be mixed and prepared.

Subserous (*sub-se'-rus*) [*sub*, under; *serous*, serous]. Beneath a serous membrane.

Subspinous (*sub-spi'-nus*) [*sub*, under; *spina*, spine]. 1. Beneath a spine. 2. Beneath the spinal column.

Substage (*sub-'s:tj*). The parts beneath the stage of a microscope, including the diaphragm, condenser, illuminator, and other accessories.

Substance of Rolando. See *Substantia gelatinosa*.

Substantia (*sub-stan'-she-ah*) [L.]. Substance. **S. cinerea**, the gray matter of the nervous system. **S. ferruginea**. Synonym of *Locus caeruleus*. **S. fusca**. See *Locus niger*. **S. gelatinosa**, a peculiar gelatinous tissue found in the spinal cord near the tip of the posterior horn. **S. grisea**, the gray matter of the spinal cord. **S. nigra**, the locus niger. **S. propria**, the essential tissue of a part; especially the middle layer (lamina propria) of the tympanic membrane.

Substernal (*sub-stur'-nal*) [*sub*, under; *sternum*]. Beneath the sternum.

Substitution (*sub-stit'-u'-shun*) [*sub*, under; *statuere*, to place]. The replacement of one thing by another. In chemistry, the replacing of one or more elements or radicles in a compound by other elements or radicles.

Subsulphate (*sub-sul'-fat*) [*sub*, under; *sulphur*, sulphur]. A basic sulphate.

Subsultus (*sub-sul'-tus*) [*sub*, under; *saltire*, to leap]. A jerking or twitching. **S. tendinum**, involuntary twitching of the muscles, especially of the hands and feet, seen in low fevers.

Subtarsal (*sub-tar'-sal*) [*sub*, beneath; *tarsus*]. Beneath the tarsus.

Subthalamic (*sub-thal-am'-ik*) [*sub*, under; *thalamus*, thalamus]. Beneath the optic thalamus.

Subtrochanteric (*sub tro kan-ter'-ik*) [*sub*, under; *τροχαντήρ*, trochanter]. Below the trochanter.

Subtrochlear (*sub trok'-le-ar*) [*sub*, under; *τροχία*, pulley]. Beneath the trochlea.

Subungual (*sub-ung'-gw'al*) [*sub*, under; *unguis*, nail]. Beneath the nail.

Suburethral (*sub-u-re'-thral*) [*sub*, beneath; *urethra*]. Beneath the urethra.

Subvaginal (*sub-vaj'-in-al*) [*sub*, beneath; *vagina*, sheath]. Beneath a sheath. **S. Space**, the space beneath the sheath of dura mater surrounding the optic nerve.

Subvertebral (*sub-ver'-te-bral*) [*sub*, under; *vertebra*, vertebra]. Beneath a vertebra.

Subzonal (*sub-zo'-nal*) [*sub*, under; *ζώνη*, zone]. Beneath the zona pellucida.

Subzygomatic (*sub-zi-go-mat'-ik*) [*sub*, under; *ζύγμα*, zygoma]. Below the zygoma.

Succedaneum (*suk-se-da'-ne-um*) [*sub*, under; *cedere*, to go]. A substitute. **S. Caput**. See *Caput*.

Succenturiate (*suk-sen-tu'-re-at*) [*succenturiare*, to receive as a substitute]. Accessory.

Succinate (*suk'-sin-at*) [*succinum*, amber]. A salt of succinic acid.

Succinic Acid (*suk-sin'-ik*). See *Acid*.

Succinum (*suk-si'-num*) [L.]. Amber, a fossil resin found in the alluvial deposits of Central Europe, and thought to be derived from an extinct species of pine. It contains a volatile oil, *Oleum succini*, used in hysteria, whooping-cough, amenorrhea, and locally as a rubefacient in chronic rheumatism, whooping-cough, and infantile convulsions. Dose ℥v-xv (0.32-1.0).

Succuba (*suk'-u-bah*) [*sub*, under; *cumbere*, to lie]. A female demon formerly believed to consort with men in their sleep. Cf. *In-cubus*.

Succulent (*suk'-u-lent*) [*succus*, a juice]. Juicy.

Succus (*suk'-us*) [L.]. 1. A vegetable juice. 2. An animal secretion. **S. entericus**, the intestinal juice, secreted by the glands of the intestinal mucous membrane. It is thin, opalescent, alkaline, and has a specific gravity of 1011. Its chief function is probably to act as a diluent. It contains an amylolytic and a proteolytic ferment. **S. gastricus**, the gastric juice.

Succussion (*suk-ush'-un*) [*sub*, under; *quater*, to shake]. A shaking, especially of the individual from side to side, for the purpose of determining the presence of fluid in a cavity or hollow organ of the body. **S.-sound**, **S.-splash**, the peculiar splashing sound heard when the patient is shaken in hydropneumothorax or pyopneumothorax, or in cases of dilated stomach containing fluid.

Sucrose (*su'-kros*). See *Saccharum*.

Suction (*suk'-shun*) [*suctio*, a sucking]. The act of sucking.

Sudamen, **Sudamina** (*su da'-men*, *su-dam'-in-ah*) [*sudor*, sweat]. An eruption of translucent whitish vesicles, due to a non-inflammatory disturbance of the sweat glands, consisting in a collection of sweat in the ducts of the sweat-glands or beneath the epidermis, and occurring in fevers and profuse sweating.

Sudation (*su da'-shun*) [*sudare*, to sweat]. The act of sweating.

Sudatorium (*su-dat'-re-um*) [*sudor*, sweat].

1. A hot air-bath. 2. A room for the administration of a hot air-bath.

Sudor (*su'-dor*) [L.]. Sweat. **S. anglicus**. See *Miliaria*. **S. cruentus**. Synonym of *Hematidrosis*. **S. nocturnus**, night-sweat. **S. sanguinosus**. See *Hematidrosis*. **S. urinosus**. See *Uridrosis*.

Sudoral (*su'-dor-al*) [*sudor*, sweat]. Pertaining to or characterized by sweating.

Sudoriferous (*su-dor-ij'-er-us*) [*sudor*, sweat; *ferre*, to bear]. Producing sweat.

Sudorific (*su-dor-ij'-ik*) [*sudor*, sweat; *facere*, to make]. 1. Inducing sweating. 2. An agent inducing sweating.

Sudoriparous (*su-dor-ip'-ar-us*) [*sudor*, sweat; *parere*, to beget]. Secreting sweat.

Suet (*su'-et*) [*sebum*, suet]. The internal fat of the abdomen of the sheep or cattle. Mutton-S. (Sevum, U. S. P., Sevum præparatum, B. P.) consists of stearin, palmitin, and olein, and is used as an emollient and in the preparation of ointments.

Suffocation (*suf-o-ka'-shun*). Interference with the entrance of air into the lungs by means other than external pressure on the trachea.

Suffocative Catarrh. Capillary bronchitis.

Suffumigation (*suf-u-mig-a'-shun*) [*suffumigatio*]. 1. The act of fumigating from underneath. 2. A substance used for fumigation.

Suffusion (*suf-u'-zhun*) [*sub*, under; *fundere*, to pour]. A spreading of a liquid over a surface; an extensive superficial extravasation of blood.

Sugar (*shooq'-ar*) [ME., *suger*, sugar]. The generic name of a class of sweet carbohydrates. See *Saccharum*. Chemically, sugars are divided as follows: Cane-sugar, $C_{12}H_{22}O_{11}$; glucose (grape-sugar or starch-sugar), $C_6H_{12}O_6$; lactose, sugar of milk; and inosite, a variety found in certain muscular tissues and in the juice of asparagus. **S.**, Beet-, saccharose obtained from species of Beta, especially the common beet, Beta vulgaris. **S.**, Diabetic, glucose. **S.**, Fruit-, levulose. **S.**, Grape-, glucose in the solid state. **S.**, Invert. See *Invert*. **S.**, Maple-, saccharose obtained from the sugar-maple. **Sugar-house Eczema**, an eczema sometimes observed in laborers employed in sugar refineries.

Suggestible (*sug-jes'-tib-l*) [*suggerere*, to suggest, from *sub*, under; *gerere*, to bring]. Amenable to suggestion.

Suggestion (*sug-jes'-chun*) [*suggerere*, to suggest, from *sub*, under; *gerere*, to bring]. 1. The artificial production of a certain psychic state in which the individual experiences such sensations as are suggested to him or ceases to experience those which he is instructed not to feel. 2. The thing suggested.

S., Posthypnotic, the command to do certain acts given the subject while in the hypnotic stage, and causing him to execute these acts after his return to his normal condition. **S.-therapy**, treatment of disordered states by means of suggestion.

Suggillation, Sugillation (*suj-il-a'-shun*) [*sugillare*, to beat black and blue]. An ecchymosis or bruise.

Sulcate (*sul'-kat*) [*sulcus*, a furrow]. Furrowed; grooved.

Sulcus (*sul'-kus*) [L.]. A furrow or groove; applied especially to the fissures of the brain. For subheadings see *Fissures*. **S.**, Intraparietal, that dividing the superior from the inferior parietal lobule. **S.**, Precentral, one situated in front of the fissure of Rolando and running nearly parallel with it. **S. spiralis**, the grooved extremity of the lamina spiralis of the cochlea. **S.**, Vertical. Same as *S.*, Precentral.

Sulfonal (*sul'-fo-nal*). See *Sulphonal*.

Sulph- (*sulf-*). See *Sulpho-*.

Sulphaminol (*sul-fam'-in-ol*) [*sulphur*, sulphur; *amin*], $C_{17}H_{19}SNO_3$. Thioxydiphenylamin, a powder obtained by the action of sulphur on the salts of methoxydiphenylamin. It is used as an antiseptic in diseases of the antrum and frontal sinuses and in laryngeal tuberculosis.

Sulphanilic Acid (*sul-fan-il'-ik*) [*sulphur*, sulphur]. See *Acids*, Table of.

Sulphate (*sul'-fat*) [*sulphur*, sulphur]. A salt of sulphuric acid.

Sulphid (*sul'-fid*) [*sulphur*]. A compound of sulphur with an element or basic radicle.

Sulphinid (*sul'-fin-id*). Saccharin.

Sulphite (*sul'-fit*) [*sulphur*]. A salt of sulphurous acid.

Sulpho- (*sul'-fo-*) [*sulphur*, sulphur]. A prefix denoting containing sulphur.

Sulphocarbolic Acid (*sul'-fo-kar-bol'-ik*). See *Acids*, Table of.

Sulphonal (*sul'-fo-nal*) [*sulphur*, sulphur], $C_8H_8S_2O_4$. Diethylsulphon-dimethylmethane, a crystalline substance soluble in 15 parts of boiling water and about 450 parts of cold water. It is used as an hypnotic in insomnia from functional causes. Dose gr. x-xl (0.65-2.6).

Sulphophenol (*sul'-fo-fe'-nol*). See *Sulphocarbolic Acid*.

Sulphosalicylic Acid. See *Salicylsulphonic Acid*.

Sulphovinic Acid (*sul'-fo-vil'-nic*), $C_4H_8H_3SO_3$, ethylsulphuric acid, a monobasic acid formed by the action of sulphuric acid on alcohol.

Sulphur (*sul'-fur*) [L., probably from Skt., *śulvāri*, sulphur]. A nonmetallic element found native in volcanic regions (volcanic S.), and occurring combined with several metals,

especially iron and copper, in the form of sulphids, called iron and copper pyrites. S. can exist in various allotropic forms. The ordinary S. is a yellow, brittle solid, having a specific gravity of 2, and an atomic weight of 31.98. Symbol S. Its valence is two, four, and six. S. combines with oxygen to form sulphurous oxid (S. dioxide), SO_2 , and sulphuric acid (S. trioxid), SO_3 , which by uniting with water form corresponding acids—sulphurous acid, H_2SO_3 , and sulphuric acid, H_2SO_4 . Other acids are also formed: hyposulphurous acid, H_2SO_2 , thiosulphuric acid, $\text{H}_2\text{S}_2\text{O}_3$, and a series of acids termed thionic acids, viz., $\text{H}_2\text{S}_2\text{O}_6$, $\text{H}_2\text{S}_3\text{O}_8$, $\text{H}_2\text{S}_4\text{O}_{10}$, and $\text{H}_2\text{S}_5\text{O}_{12}$. With hydrogen S. forms the offensively-smelling gas, hydrogen sulphid (hydrosulphuric acid or sulphureted hydrogen), H_2S . With metals and other bases it forms sulphids. S. is laxative and diaphoretic. It has been used in hemorrhoids, chronic rheumatism, gout, and locally in diphtheria and in various diseases of the skin, especially acne and scabies. S. is used in the form of S. lotum (U. S. P.), washed sulphur, dose $\mathfrak{z}\text{ss}$ – $\mathfrak{z}\text{ss}$ (2.0–16.0); S. precipitatum (U. S. P., B. P.), dose $\mathfrak{z}\text{j}$ – ij (4.0–12.0); S. sublimatum (U. S. P., B. P.), sublimed S. S. iodid (Sulphuris iodidum, U. S. P., B. P.), S_2I_2 , employed in various skin-diseases. From it is prepared Unguentum iodidi (B. P.). Unguentum sulphuris (U. S. P., B. P.) is prepared from washed S. Sulphurous oxid, SO_2 , is employed as a disinfectant by fumigation.

Sulphurated (*su'l-fu-ra-ted*) [*sulphur*, sulphur]. Combined with sulphur. S. Potassa. See *Potassa sulphurata*.

Sulphuret (*su'l-fu-ret*) [*sulphur*]. A sulphid.

Sulphureted (*su'l-fu-ret-ed*) [*sulphur*, sulphur]. Combined with sulphur. S. Hydrogen. See *Sulphur*.

Sulphuric (*su'l-fu-rik*) [*sulphur*, sulphur]. Combined with sulphur; derived from sulphur trioxid, SO_3 . S. Acid. See *Acid*, *Sulphuric*, and *Sulphur*.

Sulphurous (*su'l-fu-rus* or *su'l-fu-rus*) [*sulphur*, sulphur]. 1. Of the nature of sulphur. 2. Combined with sulphur; derived from sulphur dioxide, SO_2 . S. Acid. See *Acid*, *Sulphurous*, and *Sulphur*.

Sulphydrate (*sul-fu-hi-drat*) [*sulphur*, sulphur; *hydr*, water]. A compound of a base with the univalent radicle, Sulphydryl, SH.

Sulphydic Acid. Used improperly as a synonym of sulphureted hydrogen.

Sumbul (*sum'-bul*). The *Ferula sumbul* of the order Umbelliferae. The root (Sumbul, U. S. P., Sumbul radix, B. P.) contains angelic acid, $\text{C}_8\text{H}_8\text{O}_2$, and a little valerianic acid, $\text{C}_8\text{H}_{10}\text{O}_2$. S. is used as a nervine in neurasthenia, hysteria, and in anemia,

chronic bronchitis, etc. Dose $\mathfrak{z}\text{ss}$ – ij (2 8.0). Tinctura sumbul (U. S. P., B. P. $\mathfrak{m}\text{xx}$ – $\mathfrak{f}\mathfrak{z}\text{j}$ (1.3–4.0).

Summer-complaint. See *Cholera infantum*.

Summer-rash. Lichen tropicus.

Sun-burn. Superficial inflammation of skin caused by exposure to the sun. \mathfrak{f} stroke, insolation.

Super- (*su'-per-*) [L., above or upon]. prefix denoting above, upon, or excessive.

Superalimentation (*su-per-al-im-en-t-shun*) [*super*, over; *alimentation*]. Overfeeding.

Superciliary (*su-per-sil'-e-a-re*) [*super*, above; *cilium*, eyelash]. Pertaining to the eyebrow. S. Entropion, incurvation of the eyebrow against the conjunctiva.

Supercilium (*su-per-sil'-e-um*) [L.]. The eyebrow.

Superextension (*su-per-eks-ten'-shun*) [*super*, over; *extendere*, to extend]. Excessive extension.

Superfecundation (*su-per-fe-kun-da'-shun*) [*super*, over; *fecundus*, fertile]. The fertilization of more than one ovum of the same ovulation resulting from separate acts of coitus.

Superfetation (*su-per-fe-ta'-shun*) [*super*, over; *fetus*]. A fertilization of an ovum when there is another from a previous ovulation in the uterus.

Superficial (*su-per-fish'-al*) [*super*, over; *facies*, face]. Confined to or pertaining to the surface.

Superficialis (*su-per-fish-e-a'-lis*) [L.].

Superficial. 2. A superficial part, as, e. g., volæ, a superficial branch of the radial artery. **Superimpregnation** (*su-per-im-preg-na-shun*) [*super*, over; *impregnation*]. See *Superfetation*.

Superior (*su-pe'-re-or*) [comparative of *superus*, high]. Higher; denoting the upper of two parts.

Supernumerary (*su-per-nu'-mer-a-re*) [*super*, over; *numerus*, a number]. Existing more than the usual number.

Supersalt (*su'-per-sawlt*). An acid salt.

Supersaturate (*su-per-sat'-u-rāt*) [*super*, over; *saturare*, to saturate]. To saturate excess; to add more of a substance than liquid can dissolve.

Supination (*su-pin-a'-shun*) [*supinus*, on the back]. 1. The turning of the palm of the hand upward. 2. The condition of being supine.

Supinator (*su'-pin-a-tor*). See *Muscle Table of*.

Supplemental (*sup-le-men'-tal*) [*sub*, *ple*, to fill]. Additional. S. Air. See *Respiration*.

Suppository (*sup'-er-i-t-o-ry*) [*suppositum*, from *sub*, under; *ponere*, to place].

solid medicated compound designed to be introduced into the rectum, urethra, or vagina. Its consistency is such that while retaining its shape at ordinary temperatures, it readily melts at the temperature of the body. The basis of most suppositories is oil of theobroma. For urethral suppositories a mixture of gelatin and glycerol is used.

Suppression (*sup-resh'-un*) [*sub*, under; *primere*, to press]. A sudden cessation of secretion, as, *e. g.*, S. of the urine or menses.

Suppurant (*sup'-u-rant*) [*suppuration*]. 1. Promoting suppuration. 2. An agent promoting suppuration.

Suppuration (*sup-u-ra'-shun*) [*sub*, beneath; *pus*]. The formation of pus.

Suppurative (*sup'-u-ra-tiv*) [*suppuration*]. 1. Producing pus. 2. An agent that favors suppuration.

Supra- (*su'-prah-*) [L., above]. A prefix signifying upon or above.

Supraacromial (*su-prah-ak-ro'-me-al*) [*supra*, above; *acromion*]. Situated above the acromion.

Supraauricular (*su-prah-aw-rik'-u-lar*) [*supra*, above; *auricle*]. Above the external ear. S. Point. See *Cranio-metric Points*.

Suprachoroid (*su-prah-ko'-roid*) [*supra*, above; *choroid*]. Above the choroid or the choroid plexus.

Supraclavicular (*su-prah-kla-vik'-u-lar*) [*supra*, above; *clavis*, key]. Above the clavicle.

Supracondylar, Supracondylar (*su-prah-kon'-dil-ar*, *su-prah-kon'-dil-oid*) [*supra*, above; *condyle*]. Above the condyles.

Supracostal (*su-prah-kos'-tal*) [*supra*, above; *costa*, rib]. Above the ribs.

Supracotyloid (*su-prah-ko'-il-oid*) [*supra*, above; *cotyloid*]. Above the cotyloid cavity.

Supraglenoid (*su-prah-gle'-noid*) [*supra*, above; *glenoid*]. Above the glenoid cavity.

Supraglottic (*su-prah-glot'-ik*) [*supra*, above; *γλωττις*, glottis]. Above the glottis.

Suprahyoid (*su-prah-hi'-oid*) [*supra*, above; *hyoid*]. Above the hyoid bone.

Supramalleolar (*su-prah-mal'-o-lar*) [*supra*, above; *malleolus*, malleolus]. Above the malleoli.

Supramammary (*su-prah-mam'-a-re*) [*supra*, above; *mamma*, breast]. Above the mammae.

Supramarginal (*su-prah-mar'-jin-al*) [*supra*, above; *margo*, margin]. Above an edge or margin, as, *e. g.*, the S. convolution of the brain.

Supramastoid (*su-prah-mas'-toid*) [*supra*, above; *μαστός*, nipple; *εἶδος*, like]. Above the mastoid process.

Supramaxillary (*su-prah-maks'-il-a-re*) [*supra*, above; *maxilla*, maxilla]. Pertaining to the superior maxilla.

Supranuclear (*su-prah-nu'-kle-ar*) [*supra*, above; *nucleus*]. Above a nucleus.

Supraoccipital (*su-prah-ok-sip'-it-al*) [*supra*, above; *occiput*]. 1. Above the occipital bone. 2. The upper part of the occipital bone.

Supraorbital (*su-prah-or'-bit-al*) [*supra*, above; *orbita*, orbit]. 1. Above the orbit, as, *e. g.*, the S. nerve. 2. Pertaining to the S. nerve.

Suprapatellar (*su-prah-pat-el'-ar*) [*supra*, above; *patella*, patella]. Above the patella.

Suprapelvic (*su-prah-pel'-vik*) [*supra*, above; *pelvis*]. Above the pelvis.

Suprapineal (*su-prah-pi'-ne-al*) [*supra*, above; *pineal*]. Above the pineal gland.

Suprapontine (*su-prah-pon'-tin*) [*supra*, above; *pons*, bridge]. Above or in the superior part of the pons.

Suprapubic (*su-prah-pu'-bik*) [*supra*, above; *pubis*, pubis]. Above the pubes.

Suprarenal (*su-prah-re'-nal*) [*supra*, above; *ren*, the kidney]. 1. Above the kidney, as, *e. g.*, the S. capsule. 2. Pertaining to the suprarenal capsule. S. Body, S. Capsule, a small triangular organ situated above the kidney, and consisting of an external or cortical, and an internal or medullary portion. The cortex consists of polygonal cells disposed in three layers, the *zona glomerulosa*, *zona fasciculata*, and *zona reticularis*. Fibrous septa, derived from the capsule, extend into the organ and separate the groups of cells. The medulla contains cords and networks of polygonal cells, and in its center ganglion cells and nonmedullated nerve-fibers. The function of the S. body is not definitely known—it is believed to bear some relation to pigment-production.

Suprascapular (*su-prah-skap'-u-lar*) [*supra*, above; *scapula*, the shoulder-blade]. Above or in the upper part of the scapula.

Supraspinal (*su-prah-spi'-nal*) [*supra*, above; *spina*, spine]. Above a spine.

Supraspinales (*su-prah-spi-na'-lez*). See *Muscles, Table of*.

Supraspinatus (*su-pra-spi-na'-tus*) [*supra*, above; *spina*, spine]. Above the spine, as the S. muscle. See *Muscles, Table of*.

Supraspinous (*su-prah-spi'-nus*) [*supra*, above; *spina*, spine]. Above the spinous process of the scapula or of a vertebra. S. Fossa, the triangular depression above the spine of the scapula.

Suprasternal (*su-prah-stur'-nal*) [*supra*, above; *sternum*, sternum]. Above the sternum.

Supratemporal (*su-prah-tem'-po-ral*) [*supra*, over; *tempus*, time]. Above the temporal region.

Supratrochlear (*su-prah-trok'-le-ar*) [*supra*, above; *trochlea*]. Above the trochlea or pulley of the superior oblique muscle.

Supravaginal (*su-prah-vaj' in-al*) [*supra*, over; *vagina*, vagina]. 1. Above a sheath; on the outside of a sheath. 2. Above the vagina.

Sura (*su'-rah*) [*L.*]. The calf of the leg.

Sural (*su'-ral*) [*sura*, calf]. Pertaining to the calf of the leg.

Surcingle (*sur'-sin-gl*) [*super*, over; *cingulum*, a belt]. The tail of the corpus striatum.

Surdity (*sur'-dit-e*) [*surditas*]. Deafness.

Surdmutitas (*sur-do-mut'-lit-as*) [*surdus*, deaf; *mutus*, mute]. Deaf-mutism.

Surgeon (*sur'-jun*) [*OF.*, *chirurgien*, from *χειρ*, hand; *εργον*, to work]. One who practices surgery.

Surgery (*sur'-jer-e*) [*OF.*, *chirurgie*, from *χειρ*, hand; *εργον*, to work]. The branch of medicine dealing with diseases requiring operative procedure. **S.**, **Antiseptic**, the application of antiseptic methods in the treatment of wounds. **S.**, **Major**, that in which the operations are important and involve risks to life. **S.**, **Minor**, that part of **S.** including procedures not involving danger to life, as, *e. g.*, bandaging, the application of splints, dressings, sutures, counterirritation, cauterization, and blood-letting. **S.**, **Plastic**, repair of defects by transference of tissue.

Surgical (*sur'-jik-al*) [*surgery*]. 1. Pertaining to surgery. 2. Produced by **S.** operations. **S.** **Kidney**, suppuration of the kidney due to disease of the genitourinary tract. **S.** **Neck** of the humerus, the constricted part of the shaft below the tuberosities, so called because it is a common seat of fracture.

Sursumduction (*sur-sum-duk'-shun*) [*sursum*, up; *ducere*, to lead]. The power of the two eyes of fusing two images when one eye has a prism placed vertically before it.

Susotoxin (*su-so-tok'-sin*) [*sus*, pig; *τοξικον*, poison], $C_{10}H_{16}N_2$. A ptomain found in cultures of the bacillus of hog-cholera.

Suspension (*sus-pen'-shun*) [*sub*, under; *pendere*, to hang]. Hanging; a mode of treatment of tabes dorsalis and other nervous diseases, in which the patient hangs by the neck, chin, and shoulders.

Suspensory (*sus-pen'-so-re*) [*sub*, under; *pendere*, to hang]. Serving for suspension, as, *e. g.*, **S.** **ligament**, **S.** **bandage**.

Sustentaculum (*sus-ten-tak'-u-lum*) [*sustentare*, to support]. A support. **S.** **tali**, a process of the os calcis supporting the astragalus.

Susurrus (*su-su'-rus*) [*L.*]. A soft murmur.

Sutural (*su'-tu-ral*) [*sutura*, suture]. Pertaining to a suture.

Suture (*su'-tur*) [*sutura*, from *suere*, to sew]. 1. A line of joining or closure, as, *e. g.*, a cranial **S.** The following are the most important cranial sutures: **S.**, **Basilar**, the junction between the basilar surface of the occipital bone and the posterior surface of the body of the sphenoid. **S.**, **Coronal**, the

union of the frontal with the parietal bones transversely across the vertex of the skull. **S.**, **Ethmofrontal**, the union between the frontal and ethmoid bones. **S.**, **Ethmolacrimal**, the union between the lacrimal and ethmoid bones. **S.**, **Ethmosphenoid**, the union between the sphenoid and ethmoid bones. **S.**, **Frontal**, a **S.** which at birth joins the two frontal bones from the vertex to the root of the nose, but which afterward becomes obliterated. **S.**, **Frontomalar**, the union between the malar and frontal bones. **S.**, **Frontomaxillary**, the union between the superior maxillary and frontal bones. **S.**, **Frontonasal**, the union between the nasal and frontal bones. **S.**, **Frontoparietal**. See **S.**, **Coronal**. **S.**, **Frontosphenoid**, the union between the ala of the sphenoid bone and the frontal bone. **S.**, **Frontotemporal**, the union between the frontal and temporal bones. **S.**, **Intermaxillary**, the union between the superior maxillary bones. **S.**, **Internasal**, the union between the nasal bones. **S.**, **Interparietal**. See **S.**, **Sagittal**. **S.**, **Jugal**. See **S.**, **Sagittal**. **S.**, **Lambdoid**, the union between the two superior borders of the occipital bone and the parietal bones. **S.**, **Longitudinal**. See **S.**, **Sagittal**. **S.**, **Mastooccipital**. See **S.**, **Occipitomastoid**. **S.**, **Mastoparietal**. See **S.**, **Parietomastoid**. **S.**, **Maxillolacrimial**, the union between the lacrimal and superior maxillary bones. **S.**, **Mediofrontal**. See **S.**, **Frontal**. **S.**, **Metopic**. See **S.**, **Frontal**. **S.**, **Nasomaxillary**, the union between the superior maxillary and nasal bones. **S.**, **Occipital**. See **S.**, **Lambdoid**. **S.**, **Occipitomastoid**, the union between the mastoid portion of the temporal bone and the occipital bone. **S.**, **Occipitoparietal**. See **S.**, **Lambdoid**. **S.**, **Palatine**, the union between the palate bones. **S.**, **Parietomastoid**, the union between the mastoid portion of the temporal bone and the parietal bone. **S.**, **Petrooccipital**, the union between the occipital bone and the petrous portion of the temporal. **S.**, **Petrosphenoid**, the union between the great wing of the sphenoid bone and the petrous portion of the temporal. **S.**, **Sagittal**, the union between the superior borders of the parietal bones. **S.**, **Sphenomalar**, the union between the malar bone and the great wing of the sphenoid. **S.**, **Sphenoparietal**, the union between the ala magna of the sphenoid bone and the parietal bone. **S.**, **Sphenotemporal**, the union between the temporal and the sphenoid bone. **S.**, **Squamoparietal**. **S.**, **Squamosal**, the union between the squamous portion of the temporal bone and the parietal bone. **S.**, **Squamosphenoid**, the union between the great wing of the sphenoid and the squamous

portion of the temporal bone. 2. A stitch or series of stitches used in closing the lips of a wound. **S., Buried**, one completely covered by and not involving the skin. **S., Catgut-**, one in which the material employed is catgut. **S., Circular**, one that is applied to the entire circumference of a divided part, as the intestine. **S., Cobblers'**, one made by arming a needle with two threads. **S., Continuous, or Glovers'**, one in which the thread passes across the wound continually in the same direction, and is tied only at the beginning and end. **S., Czerny's**, for intestinal wounds, one in which the needle is passed from the serous surface through the wound, down to, but not including, the mucous membrane, and through the wound on the opposite side, and out on the serous surface. **S., Czerny-Lembert**, the application of Lembert sutures after the Czerny sutures are in place. **S., Dry**, one carried through adhesive plaster strips applied to the lips of the wound. **S., Hare-lip**, or **S., Twisted**, one in which the edges of the wound are transfixed with pins and approximation secured by twisting or wrapping the ends of the pins with thread. **S., Interrupted**, one of a series of sutures passed through the margins of the wound, and each of which is tied separately. **S., Lembert's**, an intestinal S. for wounds in which the needle is passed transversely to the wound through the peritoneal and muscular coats, and out again on one side of the wound, and then carried across the wound and made to penetrate the two outer coats as before. **S., Pin-**. See **S., Hare-lip**. **S., Quill-**, **S., Quilled**, one in which a doubled thread is passed and tied over quills or pieces of a soft catheter. **S., Relaxation-**, one introduced some distance from the wound-margin, carried through its depths, and made to emerge at some distance on the opposite side, to relieve the tension of the wound-sutures proper. **S., Shotted**, one in which both ends of the S. are passed through a perforated shot, which is then tightly compressed. **S., Subcuticular**, a buried, continuous S., in which the needle is passed horizontally into the true skin back and forth until the wound is closed. **Swallow's Nest**. The nidus hirundinis, a deep fossa of the cerebellum between the commissure of the flocculus and the uvula. **Swallowing**. Deglutition. **Swamp-sassafras**. The *Magnolia glauca*, the bark of which is aromatic and diaphoretic. **Sweat** (*svet*) [*AS., sveti*, sweat]. The secretion of the sudoriferous glands, consisting of a transparent, colorless, aqueous fluid, holding in solution neutral fats, volatile fatty acids, cholesterin, traces of albumin and urea, free lactic acid, sodium lactate, sodium chlorid, potassium chlorid, and traces of alkaline

phosphates. **S.-gland**, one of the small glands secreting the sweat, situated in the true skin and subcutaneous areolar tissue, consisting of a convoluted tube from which the excretory duct passes outward through the skin. In its passage through the epidermis the duct is more or less spiral. **Sweating Sickness**. See *Miliaria*. **Swedish Movements**. Certain systematic gymnastic exercises intended to exercise and develop the human body, and affect function, nutrition, etc. **Sweet**. Having a taste like that of sugar or honey. **S.-bread**, the thymus gland. **Swine-fever**. Hog-cholera. **Swine-plague**. An infectious disease of swine, due to the bacillus of swine-plague. **Swoon**. Syncope. **Sycosiform** (*si-ko'-se-form*) [*sycosis*; *forma*, form]. Resembling sycosis. **Sycosis** (*si-ko'-sis*) [*σῖκος*, a fig]. An inflammatory disease affecting the hair-follicles, particularly of the beard, and characterized by papules, pustules, and tubercles, perforated by hairs, together with infiltration of the skin and crusting. **S. parasitaria**, **S., Parasitic**, barber's itch, a disease of the hair-follicles, usually affecting the region covered by the beard, and due to the presence of the trichophyton fungus. **Sydenham's Chorea**. See *Chorea*. **Syllabic Utterance**. Scanning speech. **Sylvester's Method**. See *Artificial Respiration*. **Sylvian** (*sil'-ve-an*). Described by the anatomist Sylvius (1478-1555). **S. Artery**, the middle cerebral artery, lying in the fissure of Sylvius. **S. Aqueduct**. See *Aqueduct*. **S. Fissure**. See *Fissure*. **Sym-** (*sim-*). The same as *Syn-*. **Symbiosis** (*sim-be-d'-sis*) [*σῖν*, with; *βίος*, life]. The intimate association of two different living organisms, dependent on each other. **Symblepharon** (*sim-blef'-ar-on*) [*σῖν*, together; *βλέφαρον*, the eyelid]. Adhesion of the eyelids to the eyeball, known as **Anterior S.**, when the edge of the lid is adherent; **Posterior S.**, when the adhesion is at the conjunctival fold; and **Total S.**, when the entire lid is adherent. **Syme's Operation**. See *Operations, Table of*. **Symmelus** or **Symelus** (*sim'-el-us*) [*σῖν*, together; *μῆλος*, limb]. A species of monster characterized by imperfect development of the pelvis and lower extremities, with more or less intimate fusion of the latter. **Symmetric, Symmetrical** (*sim-et'-rik*, *sim-et'-rik-al*) [*σῖν*, together; *μέτρον*, a measure]. Pertaining to or exhibiting symmetry. **S. Gangrene**. See *Sphaceloderma*. **Sympathetic** (*sim-path-et'-ik*) [*σῖν*, with; *πάθος*, suffering]. 1. Pertaining to or pro-

duced by sympathy. 2. Conveying sympathy or S. impulses, as, *e. g.*, the S. System, a system of ganglia (S. ganglia) forming a chain from the cranium to the end of the spinal column, connected together by nerve-fibers, and supplying the viscera and blood-vessels. At intervals the S. nerves and ganglia form plexuses (S. plexuses). See *Plexus*, and *Ganglia*, *Table of*. S. Irritation, irritation of an organ arising from irritation of another related organ, as, *e. g.*, S. irritation of one eye from irritation of the other. S. Ophthalmia, inflammation of one eye arising subsequent to inflammation of the other eye.

Sympathic (*sim-path'-ik*). See *Sympathetic*.

Sympathy (*sim'-path-e*) [*σίμπαθία*, together; *πάθος*, suffering]. The mutual relation between parts more or less distant, whereby a change in the one has an effect upon the other.

Symphyseal (*sim-fiz'-e-al*) [*symphysis*]. Pertaining to a symphysis.

Symphysiotomy (*sim-fiz-e-ot'-o-me*) [*σύνφυσις*, symphysis; *τομή*, a cutting]. The operation of dividing the symphysis pubis, for the purpose of increasing the diameters of the pelvic canal and facilitating labor.

Symphysis (*sim'-fiz-i*) [*σίμψις*, together; *φύειν*, to grow]. The line of junction of two bones, as, *e. g.*, S. pubis.

Symplocarpus foetidus (*sim-plo-kar'-pus fel'id-us*). *Dracontium foetidum*, or skunk-cabbage, the rhizome of which is stimulant, antispasmodic, and narcotic, and has been used in asthma, chronic rheumatism, chorea, hysteria, etc. Dose gr. x-xx (0.65-1.3).

Symptom (*simf'-tom*) [*σύντομα*, a falling]. The change in a patient occurring during disease and serving to point out its nature and location. See *Signs and Symptoms*, *Table of*. S., Constitutional, S., General, one produced by the effect of the disease on the whole body. S., Local, one produced by localization of the disease in a special part. S., Objective, one observed by the physician. S., Subjective, one observed only by the patient.

Symptomatic (*simf'-tom at'-ik*) [*symptom*]. 1. Pertaining to or of the nature of a symptom. 2. Affecting symptoms, as, *e. g.*, S. treatment. S. Anthrax. See *Black-leg*.

Symptomatology (*simf'-tom-at-el'-o-je*) [*symptom*; *λόγος*, treatise]. The science of symptoms; the symptoms of disease taken together as a whole.

Sympus (*sim'-pus*) [*σίμψυς*, together; *πούς*, foot]. A monster in which there is a coalescence of the lower limbs.

Syn- (*sin-*) [*σύν*]. A prefix signifying with or together.

Synadelphus (*sin-ad-el'-fus*) [*σύν*, with; *ἀδελφός*, brother]. A monster having eight limbs with but one head and trunk.

Synæsthesia (*sin-es-the'-ze-ah*). See *Synæsthesia*.

Synalgia (*sin-al'-je-ah*) [*σύν*, together; *ἄλγος*, pain]. Pain felt in a distant part from an injury or stimulation of another part.

Synanthema (*sin-an'-them-ah*) [*συνανθεῖν*, to blossom together]. A group of efflorescences on the skin.

Synaptase (*sin-apt'-tās*). See *Emulsin*.

Synarthrodia (*sin-ar-throd'-de-ah*). See *Synarthrosis*.

Synarthrodial (*sin-ar-throd'-de-al*) [*σύν*, together; *ἄρθρον*, a joint]. Pertaining to or of the nature of a synarthrosis.

Synarthrosis (*sin-ar-throd'-sis*) [*σύν*, together; *ἄρθρον*, joint]. A form of articulation in which the bones are immovably bound together without any intervening synovial cavity. The forms are *Sutura*, in which processes are interlocked; *Schindylesis*, in which a thin plate of one bone is inserted into a cleft of another; and *Gomphosis*, in which a conic process is held by a socket.

Syncephalus (*sin-sef'-al-us*) [*σύν*, together; *κεφαλή*, head]. A monster with two heads fused into one.

Synchondrosis (*sin-kon-dro'-sis*) [*σύν*, together; *χόνδρος*, a cartilage]. A joint in which the surfaces are connected by a growth of cartilage.

Synchondrotomy (*sin-kon-drot'-o-me*) [*σύν*, together; *χόνδρος*, cartilage; *τέμνειν*, to cut]. A division of the cartilage uniting bones, especially of that of the symphysis pubis.

Synchronous (*sin'-kro-nus*) [*σύν*, together; *χρόνος*, time]. Occurring at the same time.

Synchysis scintillans (*sin'-kis-is sin'-til-lanz*) [*σύνχυσις*, a mixing together]. The presence of bright, shining particles in the vitreous humor of the eye.

Synclonus (*sin'-klo-nus*) [*σύν*, with; *κλόνος*, clonus]. 1. Clonic movements occurring simultaneously in several muscles. 2. A disease thus characterized, as, *e. g.*, chorea.

Syncopal (*sin'-ko-pal*) [*σύν*, together; *κόπτειν*, to strike or cut]. Pertaining to or characterized by syncope.

Syncope (*sin'-ko-pe*) [*σύν*, together; *κόπτειν*, to strike or cut]. Swooning or fainting, a partial or complete temporary suspension of the functions of respiration and circulation from cerebral anemia.

Syncytial (*sin-sit'-e-al*) [*σύν*, together; *κύτος*, a hollow]. Pertaining to a syncytium.

Syncytium (*sin-sit'-e-um*) [*σύν*, together; *κύτος*, a cell]. A mass of protoplasm with numerous nuclei.

Syndactylia, **Syndactylism**, **Syndactyly** (*sin-dak-til'-e-ah*, *sin-dak'-til-i-zm*, *sin-dak'-til-e*) [*σύν*, together; *δάκτυλος*, finger]. Adhesion between fingers or toes; webbed fingers, webbed toes.

Syndectomy (*sin-dek'-to-me*) [*σύν*, together; *δεῖν*, to bind; *ἐκτομή*, a cutting out]. See *Peritomy*.

Syndesmitis (*sin-dez-mi'-tis*) [*σύνδεσμος*, a ligament; *ιτις*, inflammation]. 1. Inflammation of a ligament. 2. Conjunctivitis.

Syndesmography (*sin-dez-mog'-ra-fe*) [*σύνδεσμος*, ligament; *γράφειν*, to write]. The branch of anatomy treating of ligaments.

Syndesmology (*sin-dez-mol'-o-je*) [*σύνδεσμος*, ligament; *λόγος*, treatise]. See *Syndesmography*.

Syndesmosis (*sin-dez-mo'-sis*) [*σύνδεσμος*, ligament]. A form of articulation in which the bones are connected by ligaments.

Syndesmotomy (*sin-dez-mol'-o-me*) [*σύνδεσμος*, ligament; *τέμνειν*, to cut]. 1. Dissection of the ligaments. 2. The division of a ligament.

Syndrome (*sin'-drōm*) [*σύν*, together; *δρόμος*, a running]. The aggregate symptoms of a disease; a complex of symptoms. **S. of Weber**, paralysis of the oculomotor nerve of one side and of the extremities, the face, and the hypoglossal nerve of the other side. It is usually due to a lesion limited to the inferior and inner part of one cerebral peduncle.

Synechia (*sin-e-ki'-ah* or *si-ne'-ke-ah*) [*σύν*, together; *ἔχειν*, to hold]. A morbid union of parts; especially, adhesion of the iris to a neighboring part of the eye, termed **Anterior S.**, when the adhesion is to the cornea; **Posterior S.**, when to the lens; and **Total S.**, when the entire iris is adherent.

Synechotomy (*sin-ek'-o'-o-me*) [*synechia*; *τέμνειν*, to cut]. The division of a synechia.

Synergic (*sin-ur'-jik*) [*σύν*, together; *ἔργον*, work]. Pertaining to synergy.

Synergist (*sin'-ur-jist*) [*σύν*, together; *ἔργον*, work]. An agent cooperating with another.

Synergy (*sin'-ur-je*) [*σύν*, together; *ἔργον*, work]. The cooperative action of two or more agents (synergists) or organs.

Synesthesia (*sin-es-the'-ze-ah*) [*σύν*, with; *αἰσθησις*, sensation]. A sensation felt in one part of the body as the result of an irritation of a distant organ or part.

Syngenesia (*sin-jen'-es-is*) [*σύν*, together; *γένεσις*, generation]. 1. The theory that the embryo is the product of the union of the male and female elements; also the theory that the embryo contains within itself the germs of all future generations developed from it. 2. Reproduction by union of male and female elements.

Syngnoscism (*sin-jig'-no-sizm*) [*σύν*, together; *γινώσκειν*, to know]. Hypnotism, so termed from the agreeing of one mind with another.

Synizesis (*sin-iz-e'-sis*) [*σύν*, together; *ἵκειν*, to sit]. Closure. **S. pupillæ**, closure of the pupil.

Synkinesis (*sin-kim-e'-sis*) [*σύν*, together; *κίνησις*, movement]. Involuntary movement taking place in one part of the body synchronously with or in consequence of a voluntary or reflex movement in another part.

Synneurosis (*sin-nu-ro'-sis*). See *Syndesmosis*.

Synocha (*sin'-o-kah*). See *Synochus*.

Synochus (*sin'-o-kus*) [*σύν*, together; *ἔχειν*, to hold on]. Any continued fever.

Synophthalmus (*sin-off-thal'-mus*). See *Cyclops*.

Synorchism (*sin-or'-kizm*) [*σύν*, together; *ὄρχις*, testicle]. Partial or complete fusion of the two testicles.

Synostosis (*sin-os-to'-sis*) [*σύν*, together; *ὀστέον*, bone]. A union of normally separate bones by osseous material.

Synotus (*sin-o'-tus*) [*σύν*, together; *ὅς*, ear]. A monster characterized by fused ears.

Synovia (*sin-o'-ve-ah*) [*σύν*, with; *ὥν*, an egg]. The clear, alkaline, lubricating fluid secreted within synovial membranes.

Synovial (*sin-o'-ve-al*) [*synovialis*, from; *σύν*, together; *ὥν*, egg]. Pertaining to the synovia. **S. Membrane**. See *Membrane*.

Synovitis (*sin-o-vi'-tis*) [*synovia*; *ιτις*, inflammation]. Inflammation of a synovial membrane.

Synthesis (*sin'-thes-is*) [*σύν*, with; *τίθημι*, to place]. In chemistry, the artificial formation of a compound by combining its constituents.

Synthetic (*sin-thet'-ik*) [*σύν*, with; *τίθημι*, to place]. Pertaining to or produced by synthesis.

Syntonin (*sin'-to-nin*) [*σύντονος*, contracted]. An acid-albumin obtained by the action of dilute hydrochloric acid upon the myosin of muscle.

Syphilide (*sif'-il-id*) [*syphilis*]. Any disease of the skin due to syphilis. Syphilides may be erythematous, macular, acneiform, lenticular, squamous, vesicular, pustular, bullous, tubercular, rupial, etc. **S.**, **Secondary**, any S. occurring during the secondary stage of syphilis. **S.**, **Tertiary**, any S. occurring during the tertiary stage of syphilis.

Syphilis (*sif'-il-is*) [*origin obscure*]. A chronic infectious disease, characterized by a variety of structural lesions of which the chancre, the mucous patch, and the gumma are the most distinctive. A bacillus has been found in the lesions by Lustgarten and others, but whether it is the real cause or not has not been definitely determined. The disease is generally acquired in sexual congress, hence its earliest manifestations appear upon the genital organs, but any abraded surface of the body, if brought in contact with the syphilitic poison, may give entrance to the infection. The earliest lesion of ac-

quired syphilis is the *chancre*, *initial sclerosis*, or *primary sore*, which appears after a period of incubation varying from two to three weeks. It is usually a reddish-brown papule with an ulcerated central spot, and has a slight serous or purulent discharge. Taken between the fingers, it is found to have a peculiar cartilaginous hardness. Microscopically it consists of an accumulation of round cells, epithelioid cells, with, perhaps, a giant cell here and there. The blood-vessels present a hyperplasia of the intima, to which in part the induration of the chancre is due. Very soon after the appearance of the chancre the nearest lymphatic glands become enlarged and indurated—the *indolent buboes* of syphilis. The *mucous patch*, *condyloma latum*, *moist papule*, or *mucous tubercle* is located upon mucous membranes, at mucocutaneous junctions, or where two skin-surfaces are in habitual contact, and is a flat, scarcely elevated patch, generally covered by a whitish pellicle. The *gumma* or *gummy tumor* is a rounded nodule, varying in size from the dimensions of a pea to those of a small apple. Its favorite seats are the periosteum of flat bones, the membranes of the brain, the liver, spleen, and testicle. It is usually soft and contains in its interior a gelatinous “gummy” material. Another important though not distinctive lesion produced by syphilis is a diffuse sclerosis of the blood-vessels, especially of the parenchymatous organs. The clinical course of syphilis is generally divided into three stages: the *primary* (**Primary S.**), characterized by the presence of the chancre and the indolent bubo; the *secondary* (**Secondary S.**), by the mucous patch, cutaneous eruptions, sore throat, and general enlargement of the lymphatic glands; the *tertiary* (**Tertiary S.**), by the gumma and by severe skin lesions. Between the appearance of the chancre and the secondary manifestations a period of six weeks usually elapses. The tertiary phenomena follow the secondary after a stage of quiescence of variable length. S. also bears an important, but as yet obscure, relation to certain diseases of the nervous system, such as locomotor ataxy and parietic dementia. S. can be transmitted from parent to offspring (**Hereditary S.**, **Congenital S.**). S., **Extragenital**, S. in which the first lesion is situated elsewhere than on the genital organs. S. **insontium**, S. of the innocent, *i. e.*, S. acquired in an innocent manner, or non-venereal S. S., **Marital**, S. acquired in lawful wedlock. S., **Nonvenereal**. Synonym of S. *insontium*. S. **œconomica**, a form of syphilis insontium, in which the disease is acquired through eating and drinking or household utensils, or by inci-

denal contact with syphilitic persons. S. **technica**, S. acquired in following one's occupation, as by physicians, midwives, nurses. S., **Venereal**, S. acquired in illegitimate sexual intercourse. S., **Visceral**, S. of the viscera—the lesions are either inflammatory or gummatous.

Syphilitic (*sif-il-it'-ic*) [*syphilis*]. Pertaining to or affected with syphilis.

Syphilization (*sif-il-i-zā'-shun*) [*syphilis*].

1. Inoculation with syphilis, especially inoculation for the purpose of conferring immunity to future attacks. 2. The state produced by inoculation with syphilis.

Syphiloderm (*sif-il-o'-derm*) [*syphilis*; δέρμα, skin]. See *Syphilide*.

Syphilographer (*sif-il-og'-ra-fer*) [*syphilis*; γράφειν, to write]. One who writes on syphilis.

Syphilography (*sif-il-og'-ra-fe*) [*syphilis*; γράφειν, to write]. A treatise on syphilis.

Syphiloid (*sif-il-o'-oid*) [*syphilis*; εἶδος, like].

1. Resembling syphilis. 2. A disease resembling syphilis.

Syphiloma (*sif-il-o'-mah*) [*syphilis*; δμα, tumor]. A syphilitic gumma.

Syphilophobia (*sif-il-o'-fo'-be-ah*) [*syphilis*, syphilis; φόβος, dread].

1. A condition in which the patient imagines himself to be infected with syphilis. 2. A morbid dread of syphilitic infection.

Syriac Ulcer. Synonym of *Diphtheria*.

Syringe (*sir'-inj*) [*σῆριξ*, a pipe]. An apparatus for injecting a liquid into a cavity.

Syringitis (*sir-in-jit'-tis*) [*σῆριξ*, tube; *τις*, inflammation]. Inflammation of the Eustachian tube.

Syringomyelia (*sir-ing-go-mi-el'-le-ah*) [*σῆριξ*, tube; *μυελός*, marrow]. A condition characterized by the presence of cavities in the substance of the spinal cord. It is believed to be the result of faulty development either affecting the central canal, so that the latter remains in an embryonal condition, and sends a diverticulum into the posterior portion of the cord, which may become constricted off, or groups of embryonal neuroglia-cells remain in the neighborhood of the central canal or in the posterior portion of the spinal cord, and in later life take on active development and form a gliomatous tumor. The degeneration of the latter then gives rise to cavities in the cord. S. occurs usually between the ages of 20 and 30, and is characterized by progressive atrophy of the muscles, especially of the upper extremity, by loss of the pain-sense and temperature-sense, with preservation of the tactile and muscular sense (the so called “dissociation symptom”), and by vasomotor and trophic changes in the skin, joints, etc.

Syringomyelocele (*sir-ing-go-mi-el'-le-ah*)

[σῦριγξ, tube; μνελός, marrow; κοιλία, cavity]. A form of spina bifida in which the protruding mass consists of membranes and nerve-substance, and the cavity of which communicates with the central canal of the spinal cord.

Syringotome (*sir-ing'-go-tōm*) [σῦριγξ, tube; τέμνειν, to cut]. An instrument for incising a fistula.

Syringotomy (*sir-ing-got'-o-me*) [σῦριγξ, tube; τομή, a cutting]. The operation of cutting a fistula, especially a fistula in ano.

Syrup (*sir'-up*) [*syropus*, syrup]. 1. A concentrated solution of sugar in water (*Syrupus*, U. S. P., B. P.). 2. A preparation composed of a solution of a medicinal substance in syrup.

Syrupy (*sir'-up-e*) [*siropus*, syrup]. Resembling a syrup.

Syssarcosis (*sis-ar-kō'-sis*) [σύν, together; σάρξ, flesh]. The union of bone by the interposition of muscular tissue.

Syssomus (*sis-so'-nus*) [σύν, together; σῶμα,

body]. A double monster joined by the trunks.

System (*sis'-tem*) [σίστημα, from σίν, together; στήναι, to stand]. 1. A methodic arrangement. 2. A combination of parts into a whole, as the digestive S., the nervous S. 3. The body as a whole. **S.-disease**, **S.-lesion**, a disease of the cerebro-spinal axis affecting a tract of nerve-fibers or nerve-cells having common anatomic relations and physiologic properties.

Systematic (*sis-tem-at'-ik*) [*system*]. Pertaining to or affecting a system.

Systemic (*sis-tem'-ik*) [*system*]. 1. Of or pertaining to a system. 2. Pertaining to the whole organism.

Systole (*sis'-to-le*) [σύν, together; στέλλειν, to place]. The contraction of the heart.

Systolic (*sis-tol'-ik*) [*systole*]. Pertaining to the systole; occurring during systole.

Syzygium (*sis-ij'-e-um*) [συζύγιος, yoked]. A genus of East Indian trees. **S. jambolanum** is used in diabetes.

T

T. 1. An abbreviation for tension. 2. An abbreviation for temperature.

Tabacosis (*tab-ak'-ō'-sis*) [*tabacum*, tobacco]. A state of poisoning produced by the excessive use of tobacco.

Tabacum (*tab-ak'-um*). See *Tobacco*.

Tabatière anatomique (*tahb-ah'-e-ar ahn-ah'-ōm'-ēk*) [Fr., anatomic snuff box]. The depression at the base of the thumb between the tendons of the extensor primi and extensor secundi internodii pollicis.

Tabby-cat Striation. Peculiar markings occurring on muscles that have undergone extreme fatty degeneration, especially seen in the heart-muscle.

Tabella (*ta-bell'-ah*) [L.]. A troche.

Tabes (*ta'-bēs*) [L.]. A wasting or consumption. **T. dorsalis**, locomotor ataxy, a disease dependent upon sclerosis of the posterior columns of the spinal cord. The symptoms are lightning-pains, unsteadiness and incoordination of voluntary movements, extending to the upper extremities; disorders of vision, among others the Argyll Robertson pupil; cutaneous anesthesia; girdle-sense; abolition of the patellar reflex; diminution of sexual desire; disturbance of the sphincters. **T., Hereditary**. See *Friedreich's Ataxia*. **T. mesenterica**, tuberculous disease of the mesenteric glands in children,

with progressive wasting. **T., Spasmodic**, lateral sclerosis of the spinal cord.

Tabetic (*tab-et'-ik*). See *Tabic*.

Tabic (*tab'-ik*) [*tabes*, wasting]. 1. Affected with tabes; of or pertaining to tabes. 2. Pertaining to or affected with tabes dorsalis.

Tabid (*tab'-id*). See *Tabic*, 1st definition.

Table (*ta'-bl*) [*tabula*]. 1. A flat-topped piece of furniture, as, e. g., an operation T., examining T. 2. A flat plate, especially one of bone, as, e. g., a T. of the skull.

Tablespoon. A large spoon, equivalent to about 15 c.c., or 4 fluidounces.

Tablet (*tab'-let*) [*tabula*, a table]. A lozenge; a troche.

Tabloid (*tab'-loid*) [*tabula*, table; εἶδος, like]. A preparation resembling a flat troche, or lozenge.

Tache (*tahsh*) [Fr.]. A spot. **Taches blanches**, certain white spots described by Hanot as occurring on the liver, especially on its convex surface, in infectious diseases. Microscopically they present a leukocytic infiltration and bacteria. **T. bleuâtre**, a spot of a delicate blue tint, sometimes observed on the skin of typhoid fever patients. **T. cérébrale**, **T. méningéale**, the red line made when the finger-nail is drawn over the skin; due to vasomotor paresis and occurring especially in meningeal irritation.

Tachycardia (*tak - e - kar' - de - ah*) [*ráxh*, quick; *cardia*, heart]. Excessive rapidity of the heart's action. **T.**, Essential, T. occurring in paroxysms, and due to functional disturbance of the cardiac nerves.

Tactile (*tak' - til*) [*tactus*, touch]. Pertaining to the sense of touch. **T. Cells**, cells representing special sensory nerve-endings, found in the deeper layers of the epidermis, or the adjacent stratum of corium. **T. Corpuscles**, special sensory nerve-endings exhibiting more complexity of structure than the T. cells.

Tactus (*tak' - tus*) [*tangere*, to touch]. Touch. **T. eruditus**, **T. expertus**, special sensitiveness of touch acquired by long experience.

Tædium vitæ (*tæ' - de - um vi' - te*) [L.]. Weariness of life.

Tænia (*tæ' - ne - ah*) [L., a band]. 1. A band or band-like structure. **T. fornicis**, one of the peduncles of the pineal gland. **T. hippocampi**, the corpus fimbriatum of the hippocampus major. **T. semicircularis**, a narrow band on the floor of the lateral ventricle, between the caudate nucleus and the optic thalamus. **T. violacea**, a bluish, longitudinal band, on the floor of the fourth ventricle. 2. See *Tapeworm*.

Tagetes (*ta - je' - tes*). A genus of plants of the order Compositæ. **T. erecta** and **T. patula**, French marigold, African marigold, are used as substitutes for *calendula*.

Tagliacotian Operation (*tah - le - ah - kô' - she - an*) [after *Tagliacoti*, an Italian surgeon]. See *Operations*, *Table of*.

Tagma (*tag' - mah*) [*τάγμα*, that which has been arranged, from *τάσσειν*, to arrange]. An aggregate of molecules.

Tail (*tal*). 1. The caudal extremity of an animal. 2. Anything resembling a tail.

Tailor's Spasm. An occupation-neurosis occurring in tailors, and characterized by spasm of the muscles of the arm and head.

Talc, **Talcum** (*talk*) [Arab., *talg*, talc], $4\text{MgO} \cdot 5\text{SiO}_2 \cdot \text{H}_2\text{O}$. A silicate of magnesium, a white, greasy powder, used as a dusting powder. It is also termed *steatite* or *soapstone*; the latter, however, contains also aluminum.

Talipes (*tal' - ip - ez*) [*talus*, ankle; *pes*, foot]. Club-foot, a deformity depending upon contraction of one or more muscles or tendons about the foot, either congenital or acquired. **T. calcaneus**, T. in which the patient walks upon the heel alone. **T. equinus**, T. in which the heel is elevated and the weight thrown upon the anterior portion of the foot. **T. planus**, flat-foot, splay-foot. **T. valgus**, T. in which the foot is everted. **T. varus**, a variety the reverse of the last, in which the foot is bent inward. Combinations of these occur, called *T. equinovarus*, *T. equinovalgus*, *T. calcaneovarus*, *T. calcaneovalgus*, etc.

Tallow (*tal' - o*) [O. Dutch, *taighe*, tallow]. The fat extracted from suet, the solid fat of cattle and sheep and other ruminants.

Talo- (*tal' - lo*) [*talus*, ankle]. A prefix denoting pertaining to the ankle or to the astragalus.

Talus (*tal' - hus*) [L.]. 1. The astragalus. 2. The ankle.

Tamarind (*tam' - ar - ind*) [Arab., *tamer*, a ripe date; Hind., India]. The *Tamarindus indica*, a tree of the order Leguminosæ. Its fruit (*Tamarindus*, U. S. P., B. P.) is laxative and refrigerant. Dose $\frac{5}{3}$ - $\frac{3}{3}$ (4.0 - 32.0). **Tamar indien**. An aromatic confection of senna.

Tambour (*tam' - hoor*) [Fr.]. A drum; a drum-like instrument used in physiologic experiments, and consisting of a metal cylinder over which is stretched an elastic membrane, and from which or to which passes a tube for transmitting a current of air. It is connected with another apparatus upon which changes in pressure in the T. are recorded.

Tampon (*tam' - pon*) [Fr.]. 1. A plug of cotton, sponge, or other material, inserted into the vagina, nose, or other cavity. 2. To plug with a tampon.

Tamponade (*tam - pon - ad'*) [Fr.]. The act of plugging with a tampon.

Tanacetum (*tan - as - e' - tum*). See *Tansy*.

Tanghinia (*tan - gin' - e - ah*). The *T. venenifera*, the ordeal-bean of Madagascar, a cardiac and respiratory poison. Its active principle is tanghinin.

Tangle. See *Laminaria*.

Tannate (*tan' - at*) [*tannin*]. A salt of tannic acid.

Tannic Acid. See *Acid*, *Tannic*.

Tannigen (*tan' - ij - en*) [*tannin*; γεννῶν, to produce]. A grayish-yellow powder derived from tannic acid, and used as an intestinal astringent.

Tannin (*tan' - in*). See *Acid*, *Tannic*.

Tanret's Test. A test for albumin, consisting in the development of a precipitate when an albuminous solution is brought in contact with the reagent, composed of mercuric chlorid, potassium iodid, acetic acid, and distilled water.

Tansy (*tan' - ze*) [O. Fr., *tanaisie*, from Low L., *tanacetum*, from *ἀθανασία*, immortality]. The *Tanacetum vulgare*, a plant of the order Compositæ. The leaves and tops (*Tanacetum*, U. S. P.) contain a bitter principle, *tanacetin*, $\text{C}_{11}\text{H}_{16}\text{O}_4$, tannic acid, and an essential oil (*Oleum tanacetii*). T. is an aromatic bitter and irritant narcotic, and has been used in malaria, in hysteria, and as an emmenagogue and anthelmintic. In overdoses it produces abdominal pain, vomiting, epileptiform convulsions, and death from

failure of respiration. Dose gr. xxx-3j (2.0-4.0); of the oil m.j-iv (0.065-0.26).

Tap. 1. A sudden slight blow. 2. To empty of fluid, as, *e. g.*, to tap a hydrocele.

Tapetum (*ta-pe'-tum*) [*τάπης*, a mat, or rug]. 1. The layer forming the roof of the posterior and middle cornua of the lateral ventricles of the brain; it is composed of fibers from the corpus callosum. 2. The brilliant, greenish layer of the eyes of nocturnal animals, which are by it visible in the dark. It is also known as the tapetum lucidum.

Tapeworm. One of the Cestoda, a class of worms parasitic in man and the lower animals. The adult worm (*strobilus*) consists of a head (*scolex*) and numerous segments (*proglottides*), which are capable of leading for some time a separate existence, are hermaphroditic, and contain numerous ova. If the ova are swallowed by the proper host, they develop into embryos (*proscolices*), which are transformed into the *cysticerci*, containing the *scolices*. If the meat of animals containing living scolices is eaten, the latter develop into the mature tapeworm, or strobilus. **T., Beef-** (*Tænia mediocanellata* or *saginata*), also termed the *unarmed T.*, the cysticercus of which occurs in beef. **T., Dog-** (*Tænia echinococcus*), also called hydatid *T.* The mature parasite lives in the intestine of the dog, the scolices occur in the internal organs of man and give rise to the echinococcus or hydatid cysts. **T., Fish-, T., Broad, T., Swiss** (*Bothriocephalus latus*), the cysticercus of which occurs in fish. **T., Pork-** (*Tænia solium*), also known as the *armed T.*, from the presence of several hooklets on the head, is derived from pork which contains the cysticerci. Other tapeworms occasionally found in man are: *Tænia cucumerina* or elliptica, most frequent in the dog and cat; *Tænia nana* has been found in man in Italy; *Tænia leptocephala*, common in the mouse, has also been observed in man.

Tapinocephalic (*tap-in-o-sef'-al'-ik*) [*ταπεινός*, low; *κεφαλή*, head]. Affected with tapinocephaly.

Tapinocephaly (*tap-in-o-sef'-al'-e*) [*ταπεινός*, low; *κεφαλή*, head]. Flatness of the top of the cranium.

Tapioca (*tap-e-o'-kah*) [Sp.]. A variety of starch obtained from the cassava or manioc plant, *Jatropha manihot*. It is used as a food.

Tapotement (*tap-et-mom*(g)) [Fr.]. In massage, the operation of percussing or tapping.

Tapping (*tap'-ing*). See *Paracentesis*.

Tar (*tahr*) [AS., *teoru*, tar]. An empyreumatic liquid resin obtained by the destructive distillation of the wood of various species of *Pinus*, of the order *Coniferae*. **Tar** (*Pix liquida*, U. S. P., B. P.) contains a great

variety of compounds, among which are pyroligneous acid, toluene, xylene, pseudocumene, cresol, phenol, guaiacol, creosol, paraffin, naphthalene, pyrocatechin, etc. It is employed in chronic bronchitis and in diseases of the urinary tract; externally in tinea capitis, psoriasis, chronic eczema, and other affections of the skin. Preparations: *Syrupus picis liquidæ* (U. S. P.), dose f3j-ij (4.0-8.0); *Unguentum picis liquidæ* (U. S. P., B. P.).

Tarantism (*tar'-an-tism*). A choreic affection, ascribed to the bite of a tarantula, and supposed to be cured by dancing.

Taraxacum (*tar-aks'-ak-um*). Dandelion, the *T. officinale* (*T. dens-leonis*), a plant of the order *Compositæ*. Its root (*T.*, U. S. P., *Taraxaci radix*, B. P.) contains two crystalline principles, *taraxacin* and *taraxacerin*, and is used in chronic congestion of the liver and spleen. Preparations and doses: *Decoctum taraxaci* (B. P.), f3ij (64.0); *Extractum taraxaci* (U. S. P., B. P.), gr. x (0.65); *Extractum taraxaci fluidum* (U. S. P.), f3j (4.0); *Succus taraxaci* (B. P.), f3ij-iv (8.0-16.0).

Tardieu's Spots. Ecchymotic spots found beneath the pleura and the pericardium after death from strangling. They have also been observed in death from asphyxia due to other causes.

Tarsal (*tar'-sal*) [*ταρσός*, instep]. 1. Pertaining to the tarsus of the foot. 2. Pertaining to the tarsus of the eye.

Tarsalgia (*tar-sal'-je-ah*) [*ταρσός*, tarsus; *ἀλγος*, a pain]. Pain, especially one of neuralgic character, in the tarsus.

Tarsectomy (*tar-sek'-to-me*) [*ταρσός*, tarsus; *ἐκτομή*, excision]. Excision of tarsal bones.

Tarsitis (*tar-si'-tis*) [*ταρσός*, tarsus; *ιτις*, inflammation]. Inflammation of the tarsus.

Tarso- (*tar'-so-*) [*ταρσός*, tarsus]. A prefix denoting pertaining to the tarsus.

Tarsometatarsal (*tar-so-met'-ah-tar'-sal*) [*ταρσός*, tarsus; *metatarsus*]. Relating to the tarsus and metatarsus.

Tarsophalangeal (*tar-so-fa-lan'-je-al*) [*ταρσός*, tarsus; *φάλαξ*, phalanx]. Pertaining to the tarsus and phalanges.

Tarsoplasty (*tar'-so-plas-te*) [*ταρσός*, tarsus; *πλάσσειν*, to form]. Plastic surgery of the eyelid.

Tarsorrhaphy (*tar-sor'-a-fe*) [*ταρσός*, tarsus; *ῥαφή*, suture]. The operation of sewing the eyelids together for a part or the whole of their extent.

Tarsotomy (*tar-sot'-o-me*) [*ταρσός*, tarsus; *τομή*, a cutting]. 1. The operation of cutting into the tarsus. 2. Tarsectomy.

Tarsus (*tar'-sus*) [*ταρσός*, tarsus]. 1. The instep, consisting of the os calcis, astragalus, cuboid, scaphoid, internal, middle, and ex-

ternal cuneiform bones. 2. The cartilage of the eyelid, called the tarsal cartilage, a dense connective tissue forming the support of the lid.

Tartar (*tar'-tar*) [Low L., *tartarum*, from Arab., *durd*, dregs]. 1. A hard mineral deposited on the inside of wine-casks, and consisting mainly of acid potassium tartrate (Cream of T.). T. emetic, antimony and potassium tartrate. See *Antimony*. 2. A hard incrustation on the teeth, consisting of mineral and organic matter.

Tartarated (*tar'-tar-a-ted*) [*tartar*]. Containing tartar. T. Antimony, tartar emetic. See *Antimony*.

Tartaric Acid. See *Acid, Tartaric*.

Tartarized (*tar'-tar-ized*). See *Tartarated*.

Tartarus (*tar'-tar-us*) [L.]. Tartar.

Tartrate (*tar'-trat*). A salt of tartaric acid.

Tartrated (*tar'-tra-ted*). Containing tartar; combined with tartaric acid.

Tashkend Ulcer (*tash-kend'*). See *Sartian Disease*.

Taste (*tast*). 1. The sensation produced by stimulation of special organs in the tongue (T. organs) by soluble bodies. 2. The faculty by which these sensations are appreciated. T., After-, a secondary T. perceived after the immediate T. has ceased. T.-bud, an oval, flask-shaped body, embedded in the epithelium of the tongue, and serving the sense of taste.

Tattooing (*tat-too'-ing*) [Tahitian]. The production of permanent colors in the skin by the introduction of foreign substances, such as carbon, India ink, etc., a common practice among sailors. T. of the Cornea, a method of hiding leukomatous spots.

Taurin (*tau'-rin*) [*taurus*, bull], $C_4H_7NSO_3$. Amidoethylsulphonic acid, a crystalline decomposition product of bile.

Taurocholic Acid (*tau-ro-kol'-ik*). See *Acid*.

Taxis (*taks'-is*) [*τάξις*, from *τάσσειν*, to arrange]. An arranging; a manipulation, especially manipulation for the reduction of hernia.

T.-bandage. See *Bandage*.

Te. Symbol for Tellurium.

Tea (*te*) [Chinese]. 1. The dried leaves of *Thea chinensis*, of the order *Ericaceae*, used for preparing a beverage, also called tea. 2. Any vegetable infusion used as a beverage.

Teale's Amputation. See *Operations, Table of*.

Tears. The secretion of the lacrimal gland.

Tease (*te:z*). To tear a tissue into its component parts with needles.

Teaspoon. A small spoon holding about 4 c.c., or one fluidram.

Teat (*teet*). Nipple.

Tectocephaly (*tek-to-seff'-al-e*) [*tectum*, a cover; *κεφαλή*, head]. The state of having a roof-shaped skull.

Tectorial (*tek-to'-re-al*) [*tectorium*, a cover].

Serving as a roof or covering. T. Membrane. See *Membrane of Corti*.

Teel-oil. See *Sesame-oil*.

Teething [AS., *tōð*, tooth]. The eruption of the first teeth in an infant; dentition.

Tegmen (*teg'-men*) [*tegere*, to cover]. A cover. T. tympani, the roof of the tympanic cavity.

Tegmental (*teg-men'-tal*) [*tegere*, to cover]. Pertaining to the tegmentum. T. Nucleus, the red nucleus. See *Nucleus*.

Tegmentum (*teg-men'-tum*) [*tegere*, to cover]. A covering; specifically, the dorsal portion of the crus cerebri and pons Varolii.

Tegument (*teg'-u-ment*) [*tegere*, to cover]. The integument.

Teichmann's Crystals. Hemin-crystals.

Teichopsia (*ti-kop'-se-ah*) [*τείχος*, wall; *ὄψις*, vision]. A temporary amblyopia, with subjective visual images like fortification-angles; it is probably due to vasomotor disturbances of the visual center.

Tela (*te'-lah*) [L.]. A web or tissue. T. choroidea, the membranous roof of the third and fourth ventricles of the brain. T. vasculosa, the choroid plexus.

Telangiectasis (*tel-an-je-ek'-ta-sis*) [*τέλος*, end; *ἀγγείον*, vessel; *ἐκτασις*, a stretching]. Dilatation of groups of capillaries or smaller blood-vessels.

Telangiectatic (*tel-an-je-ek'-tal'-ik*) [*τέλος*, end; *ἀγγείον*, vessel; *ἐκτασις*, a stretching]. Pertaining to or characterized by telangiectasis.

Telangioma (*tel-an-je-ek'-mah*) [*τέλος*, end; *ἀγγείον*, vessel; *ὄμα*, tumor]. A tumor composed of dilated capillaries.

Telephony (*tel-eg'-on-e*) [*τέλος*, end; *φωνή*, semen]. The influence of the paternal seed upon the offspring of a woman or animal by a later husband or male.

Telegrapher's Cramp. See *Occupation-disease*.

Telepathy (*te-lep'-ath-e*) [*τήρη*, far; *πάθος*, disease]. The action, real or supposed, of one mind upon another when the two persons are separated by a considerable distance; thought-transference.

Telluric (*tel lu'-rik*) [*tellus*, earth]. Derived from the earth.

Tellurium (*tel lu'-re-um*) [*tellus*, the earth]. A nonmetallic element of bluish-white color, having a specific gravity of 6.24, a quantitative of two, four, or six, an atomic weight of 128. Symbol Te.

Telolecithal (*tel-o-les'-ith-al*) [*τέλος*, end; *τεκθός*, yolk]. Of an ovum, having a relatively large mass of food-yolk placed eccentrically.

Telolemma (*tel-o-lem'-ah*) [*τέλος*, end; *λέμμα*, husk]. The membrane covering the eminence of Doyère, or the point of entrance of a motor nerve into a muscular fiber.

Temperament (*tem'-per-am-ent*) [*temperamentum*]. A term applied to mental disposition and physical constitution of an individual, as, *e. g.*, the bilious, lymphatic, nervous, and sanguine temperaments.

Temperature (*tem'-per-a-tūr*) [*temperatura*]. The degree of intensity of heat of a body, especially as measured by a scale termed a thermometer. **T.**, **Absolute**, that reckoned from the absolute zero of temperature, estimated at -273° C. **T.**, **Normal**, the temperature of the body in a state of health, *i. e.*, 98.6° F. **T.-sense**, the sense by which differences in temperature are appreciated, consisting of a sense for cold (*cryesthesia*) and a heat-sense (*thermoesthesia*). These are represented on the surface by different nerve-endings, the so-called cold and hot points.

Temple (*tem'-pl*) [*tempus*, time]. The portion of the head behind the eye and above the ear.

Temporal (*tem'-po-ral*) [*tempus*, time (temple)]. Pertaining to the temple, as, *e. g.*, the T. bone, T. artery.

Temporo- (*tem'-po-ro-*) [*tempus*, time]. A prefix denoting pertaining to the temple.

Temporoauricular (*tem-po-ro-aw-rik'-u-lar*) [*tempus*, time; *auricular*]. Pertaining to the temporal and auricular regions of the head.

Temporooccipital (*tem-po-ro-ok-sip'-it-al*) [*tempus*, time; *occiput*, occiput]. Pertaining to the temple and the occiput.

Temporofacial (*tem-po-ro-fa'-shal*) [*tempus*, time; *facies*, face]. Pertaining to the temple and the face.

Temporomalar (*tem-po-ro-ma'-lar*) [*tempus*, time; *mala*, cheek]. Pertaining to the temporal and malar bones.

Temporomastoid (*tem-po-ro-mas'-toid*) [*tempus*, time; *mastoid*]. Pertaining to the temporal and mastoid regions of the skull.

Temporomaxillary (*tem-po-ro-maks-il'-a-re*) [*tempus*, time; *maxilla*, maxilla]. Pertaining to the temporal region and the upper jaw.

Temporoparietal (*tem-po-ro-par-i'-et-al*) [*tempus*, time; *paries*, wall]. 1. Pertaining to the temporal and parietal bones. 2. Pertaining to the temporal and parietal lobes of the brain.

Temporosphenoidal (*tem-po-ro-sfe-noid'-al*) [*tempus*, time; *sphenoid*]. Pertaining to, or in relation with, the temporal and sphenoid bones.

Tenacious (*te-na'-shus*) [*tenax*, tough]. Tough; cohesive.

Tenacity (*te-nas'-it-e*) [*tenacitas*]. Toughness.

Tenaculum (*ten-ak'-u-lum*) [*tenere*, to hold]. A hook-shaped instrument for seizing and holding parts.

Tendinitis (*ten-din-i'-tis*). See *Tenontitis*.

Tendinous (*ten'-din-us*) [*tendon*]. Pertaining to or having the nature of tendon.

Tendo (*ten'-do*) [*L.*]. A tendon. **T. Achilles**, the common tendon of the gastrocnemius and soleus muscles inserted into the heel.

Tendon [*tendere*, to stretch]. A band of dense fibrous tissue forming the termination of a muscle and attaching the latter to a bone. **T.**, **Central**, the aponeurosis in the center of the diaphragm. **T.-reflex**, a reflex produced by stimulating the tendon of a muscle.

Tendosynovitis (*ten-do-si-no-vi'-tis*). See *Tenosynovitis*.

Tenesmus (*te-nex'-mus*) [*τείνειν*, to stretch]. A straining, especially the painful straining to empty the bowels or bladder without the evacuation of feces or urine.

Tenia (*te'-ne-ah*). See *Tapeworm*.

Teniicide (*te'-ne-as-ld*) [*tenia*, tapeworm; *cadere*, to kill]. 1. Destructive of tapeworms. 2. An agent that destroys tapeworms.

Teniafuge (*te'-ne-af-ūj*) [*tenia*, tapeworm; *fugare*, to drive]. 1. Expelling tapeworms. 2. An agent that expels tapeworms.

Tennis-arm, **T.-elbow**. A strain of the elbow, said to be frequent in tennis-players.

Teno- (*ten'-o-*) [*τένων*, tendon]. A prefix meaning pertaining to a tendon.

Tenon, **Capsule of**. A fibroelastic membrane surrounding the eyeball. It is covered by a continuous layer of endothelial plates, and corresponds to a synovial sac.

Tenonitis (*ten-on-i'-tis*) [*Tenon*, an anatomist; *itis*, inflammation]. Inflammation of Tenon's capsule.

Tenontitis (*ten-on-ti'-tis*) [*τένων*, tendon; *itis*, inflammation]. Inflammation of a tendon.

Tenontography (*ten-on-tog'-ra-fe*) [*τένων*, tendon; *γράφειν*, to write]. The descriptive anatomy of the tendons.

Tenontology (*ten-on-tol'-o-je*) [*τένων*, tendon; *λόγος*, treatise]. See *Tenontography*.

Tenorrhaphy (*ten-or'-a-fe*) [*τένων*, tendon; *ῥαφή*, suture]. The uniting of a divided tendon by sutures.

Tenosuture (*ten-o-su'-tūr*). Same as *Tenorrhaphy*.

Tenosynovitis (*ten-o-sin-o-vi'-tis*) [*τένων*, tendon; *synovia*, synovia; *itis*, inflammation]. Inflammation of a tendon and of its sheath.

Tenotome (*ten'-o-tōm*) [*τένων*, tendon; *τέμνειν*, to cut]. A knife for performing tenotomy.

Tenotomy (*ten-ot'-o-me*) [*τένον, tendon; τήνναι, to cut*]. The operation of cutting a tendon.

Tension (*ten'-shun*) [*tendere, to stretch*]. 1. The act of stretching; the state of being stretched. 2. In electricity, the power of overcoming resistance. **T.**, Intraocular, the pressure of the ocular contents upon the sclerotic coat. It may be estimated by means of an instrument called tonometer, or by palpation with the fingers, and is recorded by symbols as follows: **T**_n = normal tension; **T** + 1, **T** + 2, **T** + 3 indicate various degrees of increased **T.**, and **T** - 1, **T** - 2, **T** - 3 corresponding degrees of decreased **T.** **T. of Gases**, the tendency of a gas to expand on account of the mutual repulsion of its molecules.

Tensor (*ten'-sor*) [*tendere, to stretch*]. A stretcher; a muscle that serves to make a part tense. See *Muscles, Table of*.

Tent [*tenta*]. An instrument made of compressed sponge, **T.**, **Sponge-**, laminaria, **T.**, Laminaria, tupelo, **T.**, Tupelo-, or other material that increases in volume by the absorption of water; it is used chiefly for dilating the cervix of the uterus.

Tenth Cranial Nerve. The pneumogastric or vagus nerve.

Tentorium (*ten-to'-re-um*) [*tentorium, a tent*]. The partition between the cerebrum and the cerebellum formed by an extension of the dura mater.

Tephromyelitis (*tef-ro-mi-el-i'-tis*) [*τεφρός, ashen; myelitis*]. See *Poliomyelitis*.

Tephrosis (*tef-ro'-sis*) [*τεφρός, ashen*]. Incineration.

Tepid bath. See *Bath*.

Teratic (*ter-at'-ik*) [*τίρας, a monster*]. Monstrous.

Teratism (*ter-at-i-m*) [*τίρας, a monster*]. An anomaly of conformation, congenital or acquired.

Teratogeny (*ter-at-aj'-en-e*) [*τίρας, a monster; γένναι, to beget*]. The formation or bringing forth of monsters.

Teratoid (*ter-at-oid*) [*τίρας, monster; είδος, like*]. Resembling a monster. **T.** Tumor, a complex tumor due to the growth of tissue embryologically misplaced.

Teratology (*ter-at-ol'-o-je*) [*τίρας, monster; λόγος, science*]. The science of malformations and monstrosities.

Teratoma (*ter-at-ol'-mah*) [*τίρας, monster; άμα, tumor*]. A tumor containing teeth, hair, and other material not found in the part wherein it grows, and resulting from an embryonic misplacement of tissue or from the inclosure of parts of a rudimentary fetus.

Terchlorid (*ter-klo'-rid*) [*ter, three; χλωρός, green*]. Synonym of *Trichlorid*.

Terebene (*ter'-eb-en*) [*τέρι βινθος, terebinth-*

tree], C₁₀H₁₆. A hydrocarbon obtained the oxidation of oil of turpentine by means of sulphuric acid. It is soluble in alcohol and is used in bronchitis, dyspepsia, diseases of the genitourinary tract. **D** η_v-x (0.32-0.65).

Terebinthina (*ter-eb-in'-thin-ah*) [*τερίβιν, terebinth-tree*]. See *Turpentine*.

Terebinthinate (*ter-eb-in'-thin-at*) [*τερίβινθος, terebinth-tree*]. 1. Containing turpentine. 2. A member or derivative of the terepentine-group.

Terebration (*ter-eb-ra'-shun*) [*terebrare, bore*]. The operation of boring.

Teres (*te'-res*) [*terere, to rub*]. 1. Round, as, *e. g.*, the ligamentum **T.** 2. A muscle having a cylindrical shape, as, *e. g.*, **T. major**, **T. minor**. See *Muscles, Table of*.

Ter in die [*L.*]. Three times daily.

Term [*τέρμα, a limit*]. A limit; the time during which anything lasts.

Terminal (*ter'-min-al*) [*terminalis, from τέρμα, limit*]. Pertaining to the end; placed at or forming the end.

Ternary (*ter'-na-re*) [*ter, three times*]. A chemical compound, made up of three elements or radicals.

Teroxid (*ter-ok'-id*) [*ter, three; όξύς, acid*]. A trioxid.

Terpene (*ter'-pen*) [a modified form of *terebene*]. One of a number of hydrocarbons having the formula C₁₀H₁₆, and containing many volatile oils.

Terpin (*ter'-pin*). See *Terebene*.

Terra (*ter'-ah*) [*L.*]. Earth. **T. alba**, white clay. **T. japonica**, catechu. **T. ponderosa**, baryta or barium sulphate.

Terrain-cure (*ter-an-gy-kür*) [*Fr.*]. A method of treatment consisting in mountain climbing, dietetics, etc., for plethora, corpulence, neurasthenia, chlorosis, incipient pulmonary tuberculosis, etc.

Tersulphate (*ter-sul'-fat*) [*ter, thrice; sulphate*]. A salt in which the base is united with three sulphuric acid radicals.

Tertian (*ter'-she-an*) [*tertius, third*]. Recurring every other day, as, *e. g.*, **T. fever**, a form of intermittent fever.

Tertiary (*ter'-she-a-re*) [*tertius, third*]. Third in order. **T.** Syphilis. See *Syphilis*.

Tessellated (*tes'-el-a-ted*) [*tessellatus, from tessera, a square*]. Formed into little squares checked. **T.** Epithelium, flattened epithelial cells joined at their edges.

Test [*testum, crucible*]. 1. A trial. In chemistry, a characteristic reaction which distinguishes one body from others. 2. The reagent for producing a special reaction. **T. meal**, one given for the purpose of studying the secretory power of the stomach. **T. paper**, paper impregnated with a chemical reagent, and used for detecting the presence of

certain substances or conditions which cause a change in the color of the paper. T.-tube, a cylinder of thin glass closed at one end, used in various chemic procedures. T.-types, letters or figures of different sizes to test acuteness of vision. Those most commonly employed are Snellen's test-types, a series of letters which at proper distances subtend an angle of five minutes.

Testa (*tes'-tah*) [L.]. A shell. T. ovi, egg-shell. T. præparata, crushed and powdered oyster-shell.

Testibrachium (*tes-te-bra'-ke-um*) [*testis*, *testis*; *brachium*, arm]. The process connecting the cerebellum with the testis of the brain; the superior peduncle of the cerebellum.

Testicle (*tes'-tik-l*) [*testiculus*, dim. of *testis*]. See *Testis*, 1st definition. T., Irritable, neuralgia of the testicle. T., Undescended, the condition in which a testicle remains either in the pelvis or the inguinal canal.

Testicular (*tes-tik'-u-lar*) [*testiculus*, a testicle]. Pertaining to the testicle. T. Juice, a juice extracted from the testicles of animals, and used in the treatment of various nervous conditions.

Testis (*tes'-is*) [L.]. 1. One of the two glandular bodies in the scrotum that secrete the semen. T., Descent of, the passage of the testes through the inguinal canal into the scrotum. T. muliebria, the ovary. 2. One of the posterior pair of tubercles of the corpora quadrigemina.

Tetanic (*tel'-an'-ik*) [*tétanos*, tetanus]. 1. Pertaining to or resembling tetanus. 2. Producing tetanus.

Tetaniform (*tel'-an'-e-form*) [*tétanos*, tetanus; *forma*, form]. Resembling tetanus.

Tetanilla (*tel'-an-il'-ah*) [*tétanos*, tetanus]. Tetany.

Tetanic (*tel'-an-in*) [*tétanos*, tetanus], $C_{12}H_{20}N_2O_4$. A ptomain obtained from cultures of the bacillus of tetanus in beef-broth, and from the tissues of patients suffering from tetanus. It produces symptoms of tetanus in the lower animals. See *Ptomains*, Table of.

Tetanization (*tel'-an-i-za'-shun*) [*tétanos*, tetanus]. The production of tetanus, or of tetanic spasms.

Tetanoid (*tel'-an-oid*) [*tétanos*, tetanus; *εἶδος*, resemblance]. Resembling tetanus.

Tetanomotor (*tel'-an-o-mo'-tor*) [*tétanos*, tetanus; *motor*, a mover]. An instrument for stimulating a nerve mechanically and producing tetanus of the supplied muscle.

Tetanotoxin (*tel'-an-o-tox'-in*) [*tétanos*, tetanus; *τοξικόν*, poison], $C_8H_{11}N$. A poisonous ptomain obtained from cultures of the bacillus of tetanus. It produces first tremor, then paralysis and violent convulsions. See *Ptomains*, Table of.

Tetanus (*tel'-an-us*) [*tétanos*, from *τείνειν*, to stretch]. 1. An infectious disease characterized by tonic spasm of the voluntary muscles, an intense exaggeration of reflex activity, and peculiar convulsions. It is due to a bacillus, the bacillus of T., and has a period of incubation of from ten to twelve or more days. The first symptom is usually trismus or lock-jaw, which is soon followed by spasms of the other muscles. The poison may enter through a wound (traumatic T.); at times no point of entrance is discoverable (idiopathic T.). Puerperal T. follows labor. T. neonatorum is due to infection of the umbilicus or the circumcision-wound.

Tetany (*tel'-an-e*) [*tetanus*]. A disease characterized by intermittent, bilateral, painful, tonic spasms of the muscles, especially of the upper extremities. It is most common in young adults but may occur in others. The cause appears to be a toxic agent. It occurs in connection with typhoid fever, gastrointestinal inflammation, in rickets, dilatation of the stomach, and after extirpation of the thyroid gland.

Tetra- (*tel'-rah-*) [*τέτρα*, four]. A prefix meaning four.

Tetrabasic (*tel'-rah-ba-sik*) [*τέτρα*, four; *βάσις*, base]. Having four atoms of replaceable hydrogen.

Tetrabrachius (*tel'-rah-bra'-ke-us*) [*τέτρα*, four; *βραχίον*, arm]. A monster having four arms.

Tetrachirus (*tel'-rah-ki'-rus*) [*τέτρα*, four; *χείρ*, hand]. A monster with four hands.

Tetracoccus (*tel'-trah-kok'-us*) [*τέτρα*, four; *κόκκος*, berry]. A micrococcus occurring in groups of four.

Tetrachlorid (*tel'-rah-klo'-rid*) [*τέτρα*, four; *chlorid*]. A binary compound consisting of an element or radicle and four chlorine atoms.

Tetracid (*tel'-ras'-id*) [*τέτρα*, four; *acid*]. Having four atoms of hydrogen that are replaceable by acid radicles.

Tetrad (*tel'-rad*) [*τέτρα*, four]. An element having an atomicity of four.

Tetraiodopyrrol (*tel'-trah-i-o-do-pir'-rol*). See *Iodol*.

Tetramazia (*tel'-rah-ma'-ze-ah*) [*τέτρα*, four; *μαζός*, breast]. The presence of four breasts or mammary glands.

Tetramethylenediamin (*tel'-rah-meth-il-en-di-am'-in*), $C_4H_{10}(NH_2)_2$. Putrescin.

Tetramethyl-putrescin (*tel'-rah-meth-il-pu-tres'-in*), $C_8H_{20}N_4$. A crystalline base derived from putrescin, having very poisonous properties.

Tetranopsia (*tel'-ran-op'-se-ah*) [*τέτρα*, four; *ὄψις*, vision]. A contraction of the field of vision limited to one quadrant.

Tetrapus (*tel'-rah-pus*) [*τέτρα*, four; *πούς*,

foot]. 1. Having four feet. 2. A monster having four feet.

Tetrascelus (*tet-ras'-el-us*) [*τέτρα*, four; *σκελος*, leg]. A monster having four legs.

Tetraster (*tet-ras'-ter*) [*τέτρα*, four; *αστηρ*, star]. A karyokinetic figure characterized by an arrangement of four stars, due to a fourfold division of the nucleus.

Tetratonic (*tet-rat-on'-ik*) [*τέτρα*, four; *ἀτομος*, indivisible]. 1. Containing four atoms. 2. Having four atoms of replaceable hydrogen.

Tetravalent (*tet-rav'-al-ent*). See *Quadrivalent*.

Tetronal (*tet'-ron-al*) [*τέτρα*, four; $C_3H_8 \cdot S_2O_4$]. Diethylsulphondiethylmethane, an hypnotic resembling sulphonal. Dose gr. x-xx (0.65-1.3).

Tetroxid (*tet-roks'-id*) [*τέτρα*, four; *oxid*]. A binary compound composed of a base and four atoms of oxygen.

Tetter (*tet'-er*) [AS., *teter*]. A name for various skin-eruptions, particularly herpes, eczema, and psoriasis.

Texas Fever. An infectious disease of cattle, characterized by high fever, hemoglobinuria, and enlargement of the spleen. The disease is probably due to a specific organism.

Textural (*teks'-tu-ral*) [*textura*, texture]. Pertaining to the tissues.

Thalamencephalon (*thal-am-en-sef'-al-on*) [*θάλαμος*, couch; *ἐνκέφαλον*, the brain]. The posterior portion of the anterior brain-vesicle; the interbrain.

Thalamic (*thal-am'-ik*) [*thalamus*]. Pertaining to the optic thalamus.

Thalamocortical (*thal-am-o-kor'-tik-al*) [*θάλαμος*, couch; *cortex*]. Pertaining to the optic thalamus and the cortex of the brain.

Thalamolenticular (*thal-am-o-len-tik'-u-lar*) [*θάλαμος*, chamber; *lenticular*]. Pertaining to the optic thalamus and the lenticular nucleus.

Thalamus (*thal'-am-us*) [*θάλαμος*, couch]. A mass of gray matter at the base of the brain, developed from the wall of the vesicle of the third ventricle, and forming part of the wall of the latter cavity. The posterior part is called the pulvinar. The T. receives fibers from all parts of the cortex, and is also connected with the tegmentum and with fibers of the optic tract. It is also termed **T. opticus**, or **optic T.**

Thalassotherapy (*thal-as-o-ther'-a-p'e*) [*θάλασσα*, sea; *θεραπεία*, treatment]. Treatment of disease by sea-voyages, sea-bathing, etc.

Thallin (*thal'-in*) [*θαλλος*, a green shoot]. $C_{10}H_{11}ON$. A liquid basic substance, the tetrahydroparamethyloxychinolin. The sulphate and tartrate are used as germicides and antipyretics, and have been employed in typhoid fever and the fever of phthisis. Dose

gr. j-iv (0.065-0.26). In gonorrhea the sulphate is employed as an injection in the strength of a 1.5 per cent. solution.

Thallium (*thal'-e-um*) [*θαλλός*, a green shoot]. A metallic element having an atomic weight of 203.7, a specific gravity of 11.8; symbol, Tl. The salts are poisonous. See *Elements*, *Table of*.

Thallophyte (*thal'-o-fyt*) [*θαλλός*, a green shoot; *φυτόν*, a plant]. One of a class of very low cryptogams.

Thanato- (*than'-at-o-*) [*θάνατος*, death]. A prefix denoting pertaining to death.

Thanatognomonic (*than-at-og-no-mon'-ik*) [*θάνατος*, death; *γνώμων*, sign]. Indicative of death.

Thanatoid (*than'-at-oid*) [*θάνατος*, death; *ειδος*, like]. Resembling death.

Thanatometer (*than-at-om'-et-er*) [*θάνατος*, death; *μέτρον*, measure]. A thermometer introduced into a body-cavity to determine if the depression of temperature is so great as to be a sign of death.

Thanatophobia (*than-at-o-f'o-be-ah*) [*θάνατος*, death; *φόβος*, dread]. A morbid fear of death.

Thea (*he'-ah*). Tea, the dried leaves of *Thea chinensis*, a shrub of the order Ternstroemiaceæ, containing the alkaloid Thein, $C_8H_{10}N_2O_2$, identical with caffeine. T. is astringent and gently stimulant to the nervous system; its infusion is used as a beverage.

Thebain (*the'-ba-in*) [*Θηβαι*, Thebes], $C_{15}H_{21}NO_3$. An alkaloid found in opium, analogous to strychnin in its physiologic effects. **Thebesius**, **Foramina of**. See *Foramina*, *Table of*.

Thebolactic Acid (*theb-o-lak'-tik*). A variety of lactic acid occurring in milk.

Theca (*the'-kah*) [*θήκη*, a sheath]. A sheath, especially one of a tendon. **T. cerebri**, the cranium. **T. cordis**, the pericardium. **T. vertebralis**, the membranes of the spinal cord.

Thecal (*the'-kal*) [*θήκη*, a sheath]. Pertaining to a sheath.

Thecitis (*the-si'-tis*) [*θήκη*, a sheath; *τις*, inflammation]. Inflammation of the sheath of a tendon.

Thein (*the'-in*). See *Thea*.

Thelyblast (*thel'-e-blast*) [*θήλυς*, female; *βλαστος*, germ]. The female element of the bisexual nucleus; the ovum after the polar globules have been extruded.

Thenad (*the'-nad*) [*θεν*, palm; *ad*, toward]. Toward the thenar eminence.

Thenal (*the'-nal*) [*θεν*, the palm]. Pertaining to the palm, or the thenar eminence.

Thenar (*the'-nar*) [*θεν*, palm]. 1. The palm of the hand. 2. The fleshy prominence of the palm corresponding to the base of the thumb, also called **T. Eminence**.

Theobroma (*the-o-bro'-mah*) [*θεός*, a god; *βρώμα*, food]. A genus of trees of the Sterculiaceae. The seeds of *T. cacao* yield a fixed oil (*Oleum theobromatis*, U. S. P.), and contain the alkaloid theobromin, $C_7H_8N_4O_2$, which is closely related to caffeine and xanthin. The seeds are used in the preparation of chocolate and cocoa; the oil (butter of cacao) is employed as an ingredient of cosmetic ointments and for making pills and suppositories. Theobromin acts similarly to caffeine. The salicylate of sodium and theobromin, called *diuretin*, is an active diuretic. Dose gr. xv (I.O.).

Theobromin (*the-o-bro'-min*). See *Theobroma*.

Theomania (*the-o-ma'-ne-ah*) [*θεός*, a god; *μανία*, madness]. 1. Religious mania. 2. Insanity in which the patient believes himself to be a divine being.

Theophyllin (*the-o-phil'-in*) [*thea*, tea; *φύλλον*, leaf]. $C_7H_8N_4O_2$. An alkaloid occurring in tea and isomeric with theobromin and with paraxanthin.

Therapeutic (*ther-ap-u'-tik*) [*θεραπευτική*, treatment]. Pertaining to therapeutics; curative. **T. Test**, a method of diagnosis by administering certain remedies known to influence a given disease, *e. g.*, quinin in malaria, potassium iodid and mercury in syphilis.

Therapeutics (*ther-ap-u'-tik-s*) [*θεραπευτική*, *θεραπεία*, treatment]. The branch of medical science dealing with the treatment of disease.

Therapeutist (*ther-ap-u'-tist*) [*θεραπεία*, treatment]. One skilled in therapeutics.

Therapy (*ther'-ap-e*) [*θεραπεία*, treatment]. See *Therapeutics*.

Theriaca (*the-ri'-ak-ah*) [*θηριακή*, from *θηρίον*, a wild beast, because believed to be good against the poison of animals]. Treacle; molasses. **T. Andromachi**, Venice treacle, a compound containing nearly seventy ingredients, and used as an antidote against poisons.

Therm [*θέρμη*, heat]. The amount of heat required to raise the temperature of one gram of water from 0° C. to 1° C.; it is equal to 1000 calories.

Thermæsthesia (*thur-mes-the'-se-ah*). See *Thermesthesia*.

Thermal (*thur'-mal*) [*θέρμη*, heat]. 1. Pertaining to heat. 2. Hot, as, *e. g.*, T. spring.

Thermanesthesia (*thur-man-es-the'-se-ah*). See *Thurmoanesthesia*.

Thermesthesia (*thur-mes-the'-se-ah*) [*θέρμη*, heat; *αἴσθησις*, sensation]. 1. The heat-sense. 2. Sensitiveness to heat.

Thermic (*thur'-mik*) [*θέρμη*, heat]. Pertaining to heat. **T. Fever**, sunstroke; heat-fever.

Thermo- (*thur'-mo-*) [*θέρμη*, heat]. A prefix meaning heat.

Thermoanesthesia (*thur-mo-an-es-the'-se-ah*) [*θέρμη*, heat; *anesthesia*]. Loss of the perception of thermal impressions, a condition sometimes present in syringomyelia.

Thermocautery (*thur-mo-kaw'-ter-e*). See *Cautery*.

Thermochroic (*thur-mo-kro'-ik*) [*θέρμη*, heat; *χρῶμα*, color]. Transmitting some thermal rays and absorbing others.

Thermoelectricity (*thur-mo-e-lek-tris'-it-e*) [*θέρμη*, heat; *electricity*]. Electricity generated by heat.

Thermogenesis (*thur-mo-jen'-es-is*) [*θέρμη*, heat; *γεννᾶν*, to produce]. The production of heat.

Thermogenetic, Thermogenic, Thermogenous (*thur-mo-jen-et'-ik*, *thur-mo-jen'-ik*, *thur-moj'-en-us*) [*θέρμη*, heat; *γεννᾶν*, to produce]. Pertaining to thermogenesis; producing heat.

Thermograph (*thur'-mo-graf*) [*θέρμη*, heat; *γράφειν*, to write]. A device for registering variations of temperature automatically.

Thermohyperesthesia (*thur-mo-hi-per-es-the'-se-ah*) [*θέρμη*, heat; *ὑπερ*, above; *αἴσθησις*, sensation]. Abnormal sensitiveness to the application of hot bodies.

Thermoinhibitory (*thur-mo-in-hib'-it-o-re*) [*θέρμη*, heat; *inhibitory*]. Inhibiting the production of heat.

Thermolysis (*thur-mol'-is-is*) [*θέρμη*, heat; *λύσις*, a loosening]. 1. Dissipation of animal heat. 2. Chemic decomposition by means of heat.

Thermolytic (*thur-mo-lit'-ik*) [*θέρμη*, heat; *λίσις*, a loosening]. Pertaining to thermolysis.

Thermometer (*thur-mom'-et-er*) [*θέρμη*, heat; *μέτρον*, measure]. An instrument for measuring the intensity of heat, consisting of a substance capable of expanding and contracting, and a graduated scale by means of which variations in the volume of the substance can be determined. In the ordinary T. the expansive substance is mercury (mercurial T.), expanding into a vacuous capillary tube, the degree of heat being measured by the length of the column of mercury. T., Centigrade, T., Celsius-, one in which the freezing-point is at 0° and the boiling-point at 100°. T., Clinical, a self-registering thermometer for ascertaining the bodily temperature. T., Fahrenheit-, one in which the interval between the freezing-point and the boiling-point is divided into 180 equal parts, each called a degree, the zero-point being 32 degrees or divisions below the freezing-point of water. T., Reaumur-, one in which the freezing-point of water is 0° and the boiling-point 80°. T., Self-registering, one that by means of an index shows the highest (maximum-

T.) or lowest (minimum-T.) temperature to which it has been exposed. **T., Surface-**, one for registering the surface-temperature of any portion of the body.

Thermometric (*thur-mo-met'-rik*) [*θερμη, heat; μετρον, measure*]. Pertaining to a thermometer.

Thermometry (*thur-mom'-et-re*) [*θερμη, heat; μετρον, measure*]. The measuring of temperature by means of the thermometer.

Thermopile (*thur'-mo-pl'*) [*θερμη, heat; pile*]. A contrivance consisting of a series of connected metallic plates, in which, under the influence of heat, a current of electricity is produced, which acts upon a registering index. By means of it very minute amounts of heat can be measured.

Thermopneypnea (*thur-mo-pol-ip-ne'-ah*) [*θερμη, heat; πολλος, many; πναιν, to breathe*]. Rapid respiration due to high temperature.

Thermostat (*thur'-mo-stat*) [*θερμη, heat; στατος, standing*]. A device for automatically regulating and maintaining a constant temperature.

Thermosystaltic (*thur-mo-sis-tal'-tik*) [*θερμη, heat; συστειλλειν, to contract*]. Contracting under the influence of heat; pertaining to muscular contraction due to heat.

Thermotactic, Thermotaxic (*thur-mo-tak'-tik, thur-mo-taks'-ik*) [*θερμη, heat; τάσσειν, to arrange*]. Regulating the heat of the body, as, e. g., a T. center.

Thermotaxis (*thur-mo-taks'-is*) [*θερμη, heat; τάξις, from τάσσειν, to arrange*]. The regulation of the bodily heat.

Thermotherapy (*thur-mo-ther'-ap-e*) [*θερμη, heat; θεραπεια, cure*]. Treatment of disease by heat.

Thiersch's Method of Skin-grafting. A method in which long, broad strips of skin are removed from the arm or leg and placed on a wound which has previously been deprived of its granulations by means of a sharp curet.

Thigh (*thi*) [AS., *thēōh*, thigh]. The part of the lower limb extending from pelvis to knee.

Thilandin (*thi'-lan-in*) [*θειον, sulphur; lano-lin*]. A brownish-yellow substance derived from and resembling lanolin. It contains 3 per cent. of sulphur, and is used in the treatment of eczema and other diseases of the skin.

Thio- (*thi'-o-*) [*θειον, sulphur*]. A prefix denoting containing sulphur.

Thioacid (*thi-o-as'-id*). One of a group of acids produced by the substitution of sulphur for the oxygen in an oxygen-acid.

Thioalcohol (*thi-o-al'-ko-hol*). See *Mer-captan*.

Thiocyanic Acid (*thi-o-si-an'-ik*), CNHS. Sulphocyanic acid, a monobasic acid forming the thiocyanates or sulphocyanates. Potassium thiocyanate, CNKS, occurs in saliva.

Thioether (*thi-o-e'-ther*). See *Alkyl-mul-phids*.

Thiol (*thi'-ol*) [*θειον, sulphur*]. A substance prepared from gas-oil by heating with sulphur, and occurring in a dry and a liquid form. It has been used as an application to ulcers and in diseases of the skin.

Thionin (*thi'-o-nin*) [*θειον, sulphur*], $C_{12}H_2N_2S$. A sulphur-compound of the aromatic group, used as a stain in microscopy. Its solutions are of a dark blue color.

Thiophene (*thi'-o-fen*) [*θειον, sulphur*], C_4H_4S . A hydrocarbon of the aromatic series; a colorless, oily liquid, miscible with water. **T. sodium sulphonate**, $C_4H_4S.NaSO_3$, is a white powder; it is used in prurigo. The iodid, $C_4H_4I_2S$, has been used as a substitute for iodoform.

Thioresorcin (*thi-o-res-or'-sin*) [*θειον, sulphur; resorcin*], $C_6H_4(SO)_2$. A compound of sulphur and resorcin used as a powder or ointment as a substitute for iodoform.

Thiosinamin (*thi-o-sin'-am-in*) [*θειον, sulphur; sinapis, mustard*], $C_4H_4N_2S$. A crystalline substance prepared from mustard-oil and ammonia. It is used in lupus, glandular enlargements, and night-sweats.

Thiosulphuric Acid (*thi-o-sul-fu'-rik*) [*θειον, sulphur; sulphur*], $H_2S_2O_3$. An acid derived from sulphuric acid by the substitution of sulphur for part of the oxygen.

Thiourea (*thi-o-u-re'-ah*) [*θειον, sulphur; urea*], $CS(NH_2)_2$. Sulphocarbamid; a derivative of urea in which sulphur replaces the oxygen of the latter.

Third Cranial Nerve. The oculomotor nerve. See *Nerves, Table of*.

Thirst (*thurst*) [AS., *thurst*]. A desire for drink.

Thiry's Fistula (*thir'-rez*). A fistula for obtaining the intestinal juice. A piece of intestine about four inches long is separated from the bowel, without dividing the mesentery and its blood-vessels. One end of the tube is closed, the other is stitched to the abdominal wound. The two ends of intestine from which the piece was cut out are then united by sutures. From the excised piece a pure intestinal juice is obtained.

Thlipsencephalus (*thlip-sen-seft'-al-us*) [*θλιψις, pressure; ἐγκέφαλον, brain*]. A monster in which there is extensive exposure of the base of the skull from nondevelopment of the occipital bone and even of the upper vertebrae.

Thomsen's Disease [after J. *Thomsen*, a German physician]. Myotonia congenita, a disease commonly congenital and occurring in families, and characterized by tonic spasm or rigidity of the muscles, coming on when they are first put in action after a period of rest. As the muscles are used the stiffness

gradually wears off. See *Diseases, Table of*.

Thomsonianism (*tom-so'-ne-an-ism*). A system of medicine introduced by Samuel Thomson (1769-1843), of Massachusetts. It insisted on the use of vegetable remedies only.

Thoracentesis (*tho-ras-en-te'-sis*) [θώραξ, chest; κέντησις, a piercing]. Puncture of the thorax for the removal of fluid.

Thoracic (*tho-ras'-ik*) [θώραξ, thorax]. Pertaining to or situated in the chest or thorax, as, e. g., the T. aorta. T. Duct. See *Duct. T. Index*. See *Index*.

Thoraco- (*tho'-rak-o-*) [θώραξ, thorax]. A prefix denoting pertaining to the thorax.

Thoracocentesis (*tho-rak-o-sen-te'-sis*). See *Thoracentesis*.

Thoracodidymus (*tho-rak-o-did'-im-us*) [θώραξ, chest; διδυμος, double]. A double monster joined by the thoraces.

Thoracodynia (*tho-rak-o-din'-e-ah*) [θώραξ, thorax; ὀδύνη, pain]. Pain in the chest.

Thoracometer (*tho-rak-om'-et-er*) [θώραξ, thorax; μέτρον, a measure]. A stethometer.

Thoracopagus (*tho-rak-op'-ag-us*) [θώραξ, thorax; πηγνύναι, to join]. A double monster with portions of the thoraces coalescent.

Thoracoplasty (*tho'-rak-o-plas-te*) [θώραξ, thorax; πλάσσειν, to form]. Plastic operation upon the thorax.

Thoracoscopy (*tho-rak-os'-ko-pe*) [θώραξ, thorax; σκοπεῖν, to examine]. Examination of the chest, especially by the stethoscope.

Thoracotomy (*tho-rak-ot'-o-me*) [θώραξ, thorax; τέμνειν, to cut]. Incision of the thorax or chest-wall.

Thoradelphus (*tho-rad-el'-fus*) [θώραξ, thorax; ἀδελφός, brother]. A monster consisting of two individuals united above the umbilicus, with one head, four lower, and two upper extremities.

Thorax (*tho'-raks*) [θώραξ]. The chest; the framework of bones and soft tissues, bounded by the diaphragm below, the ribs and sternum in front, the ribs and dorsal portion of the vertebral column behind, and above by the structures in the lower part of the neck, and containing the heart enclosed in the pericardium, the lungs invested by the pleura, and the mediastinal structures. T., *Regions of*. See *Regions*.

Thorn-apple. See *Stramonium*.

Thornwaldt's Disease. See *Diseases, Table of*.

Thorough-wort. See *Eupatorium*.

Thought-transference. See *Telepathy*.

Thread-worm. See *Oxyuris*.

Three-day Fever. *Synonym of Dengue*.

Threpeology (*threp-sol'-o-je*) [θρεψις, nutrition; λόγος, treatise]. The science of nutrition.

Threshold (*thresh'-old*). The lower limit of stimulus capable of producing an impression upon consciousness.

Thrill. A fine vibration felt by the hand. A thrill may be felt on palpation over an aneurysm, over a heart the seat of valvular disease, and over hydatid cysts. T., *Purring*, a thrill resembling that felt when the hand is placed on the back of a cat.

Throat (*thrōt*) [AS., *throate*, throat]. 1. The anterior part of the neck. 2. The pharynx and larynx; the fauces. T., *Sore*, pharyngitis.

Throb. A pulsation or beating.

Throbbing (*throb'-ing*). A rhythmic beating. T. *Aorta*, exaggerated pulsation of the abdominal aorta perceptible to the patient.

Thrombo- (*throm'-bo-*) [θρόμβος, thrombus]. A prefix denoting pertaining to a thrombus.

Thromboarteritis (*throm-bo-ar-ter-i'-tis*) [θρόμβος, thrombus; ἀρτηρία, artery; ιτις, inflammation]. Inflammation of an artery associated with thrombosis.

Thromboid (*throm'-boid*) [θρόμβος, thrombus; εἶδος, like]. Resembling a thrombus.

Thrombophlebitis (*throm-bo-pleb-i'-tis*) [θρόμβος, thrombus; φλέψ, vein; ιτις, inflammation]. Inflammation of a vein associated with thrombosis.

Thrombosis (*throm-bo'-sis*) [θρόμβος, thrombus]. The formation of a thrombus.

Thrombotic (*throm-bot'-ik*) [θρόμβος, thrombus]. Pertaining to or produced by thrombosis.

Thrombus (*throm'-bus*) [θρόμβος, clot]. A clot of blood formed within the heart or blood-vessels, due usually to a slowing of the circulation or to alteration of the blood or vessel-walls.

Through-Illumination. Transillumination.

Through-drainage. A method of drainage in which a perforated tube is carried through the cavity to be drained, so that the latter can be flushed through and through by the injection of fluid into one end of the tube.

Thrush. A form of stomatitis due to a specific fungus, the *Oidium albicans*, or *Saccharomyces albicans*, and characterized by the presence of diffuse, white patches. It occurs especially in weakly children, but may affect adults depressed by wasting diseases.

Thuja (*thu'-yah, thu'-jah*). A genus of trees of the order *Coniferae*. T. *occidentalis*, arbor vitæ, has been used in intermittent fever, rheumatism, and scurvy, and as an emmenagogue.

Thumb (*thum*). The digit on the extreme radial side of the hand, differing from the other digits in having but two phalanges, and in that its metacarpal bone is separately movable.

Thus [*L., thus*]. True frankincense, or olibanum.

Thymacetin (*thi-mat'-et-in*) [*thymon*, thyme; *acetum*, vinegar], $C_8H_8(ClH_2)(C_2H_5)(OC_2H_5) \cdot NH(C_2H_5O)$. A derivative of thymol used as an antineuralgic. Dose gr. iij-xv (0.2-1.0).

Thyme (*tim*) [*thymon*, thyme]. The genus *Thymus*, of the order Labiate. *Thymus vulgaris* yields a volatile oil (*oleum thymi*, U. S. P.), in which are found cymene, thymene, and thymol.

Thymic (*thi'-mic*, also for first definition, *ti'-mik*) [*thymos*, thyme]. 1. Pertaining to or contained in thyme. 2. Pertaining to the thymus gland. **T. Asthma**, a form of laryngismus stridulus, consisting in a temporary suspension of respiration, attributed to enlargement of the thymus.

Thymol (*ti'-mol*, *thi'-mol*) [*thymon*, thyme; *oleum*, oil], $C_{10}H_{14}O$. A phenol derived from the volatile oils of *Thymus vulgaris*, *Monarda punctata*, and *Carum ajowan*. It is a crystalline solid, melting at $44^\circ C$, very slightly soluble in water, and is used as a local antiseptic and deodorant in ulcers, leukorrhea, and stomatitis, as an intestinal antiseptic, and as an anthelmintic. Dose gr. i-ij (0.065-0.13).

Thymus (*thi'-mus*) [*thymos*, thymus]. The T. gland, an organ situated in the anterior superior mediastinum. It continues to develop until the second year of life, afterward remains stationary until about the fourteenth, and then undergoes fatty metamorphosis and atrophy. The T. consists of lobules largely composed of lymphadenoid tissue in which minute concentric bodies, the corpuscles of Hassall, are found. The latter are remnants of epithelial structures. 2. A genus of labiate plants. See *Thyme*.

Thyreo- (*thi'-re-o-*). See *Thyro-*.

Thyro- (*thi'-ro*) [*thyreos*, a shield, and hence, from similarity of shape and function, the thyroid gland or cartilage]. A prefix signifying relationship to the thyroid gland.

Thyroarytenoid (*thi-ro-ar-it'-en-oid*) [*thyreos*, shield; *arytenoid*]. Pertaining to the thyroid and arytenoid cartilages, as, e. g., the T. ligaments; T. muscle (**Thyroarytenoides**).

Thyrocele (*thi'-ro-sel*) [*thyreos*, thyroid; *kelos*, a tumor]. A tumor affecting the thyroid gland; goiter.

Thyroepiglottic (*thi-ro-ep-e-glot'-ik*) [*thyreos*, shield; *epiglottis*]. Pertaining to the thyroid cartilage and the epiglottis, as, e. g., the T. muscle (**Thyroepiglottideus**).

Thyrohyoid (*thi-ro-hi'-oid*) [*thyreos*, shield; *hyoid*]. Pertaining to the thyroid cartilage and hyoid bone, as, e. g., the T. membrane.

Thyroid (*thi'-roid*) [*thyreos*, shield; *id*, like]. 1. Shield-shaped. 2. Pertaining to the T. gland. 3. Pertaining to the T. cartilage. 4. Pertaining to the T. foramen.

Cartilage, the largest of the laryngeal cartilages, united at an angle in front, called *pomum Adami*. **T. Foramen**. See *Foramen*, *Obturator*.

T. Gland, one of the so-called ductless glands, lying in front of trachea, and consisting of two lateral lobes connected centrally by an isthmus. The gland is composed of follicles lined by epithelium, producing a peculiar colloid material. The function of the organ is not definitely known, but is supposed to be the production of some substance necessary to the body's internal secretion that may counteract poison produced in the system. Hypertrophy of gland (goiter) is sometimes associated with peculiar disease known as exophthalmic goiter; absence of the gland leads to cretinism or myxedema. An extract prepared from the T. gland of animals (**T. Extract**) and other preparations of the gland are used medicinally. See *Organotherapy*.

Thyroidectomy (*thi-roi-dek'-to-me*) [*thyreos*, shield; *ektomē*, excision]. Excision of thyroid gland.

Thyroidin (*thi-roi'-din*) [*thyreos*, shield; *idin*, like]. An extract of the thyroid gland.

Thyroiditis (*thi-roi-di'-tis*) [*thyreos*, shield; *itis*, inflammation]. Inflammation of thyroid gland.

Thyroidotomy (*thi-roi-dot'-o-me*) [*thyreos*, shield; *tomē*, a cutting]. Incision of thyroid gland.

Thyrotomy (*thi-roi'-o-me*) [*thyreos*, shield; *tomē*, a cutting]. Incision or splitting of thyroid cartilage.

Tibia (*tib'-e-ah*) [*L., shin*]. The large bone of the leg, commonly called the shin-bone, articulating with the femur, fibula, and astragalus.

Tibial (*tib'-e-al*) [*tibia*, tibia]. Pertaining to or in relation with the tibia, as, e. g., T. muscle (*Tibialis*), T. artery, T. nerve.

Tibio- (*tib'-e-o-*) [*tibia*, tibia]. A prefix meaning pertaining to the tibia.

Tibiofemoral (*tib-e-o-fem'-or-al*) [*tibia*, tibia; *femur*, femur]. Pertaining to the tibia and the femur.

Tibiofibular (*tib-e-o-fib'-u-lar*) [*tibia*, tibia; *fibula*, fibula]. Pertaining to the tibia and the fibula.

Tibiotarsal (*tib-e-o-tar'-sal*) [*tibia*, tibia; *taraxos*, tarsus]. Pertaining to the tibia and the tarsus.

Tic (*tik*) [*Fr.*]. A twitching, especially of the facial muscles. **T. convulsif**, **T. convulsive**, spasm of the facial muscles.

douloureux, neuralgia of the trifacial nerve.

Tick (*tik*). A name applied to several species of *Acarus*.

Tickle (*tik'-l*). To touch so as to cause a peculiar sensation (tickling, or titillation), usually associated with laughing and reflex muscular movements.

T. i. d. An abbreviation for *ter in die*, three times a day.

Tidal Air. See *Respiration*. **T. Wave.** See *Wave*.

Tigilium (*tig'-le-um*). The *Croton tiglium* of the order Euphorbiaceæ. It contains a fixed oil, croton oil (*Oleum tiglii*, U. S. P., *Oleum crotonis*, B. P.), which is a powerful local irritant, and is used, locally, as a counter-irritant in gout, rheumatism, neuralgia, glandular swellings, etc. It is an active purgative, especially useful in cases when a prompt effect is desired, as in mania, coma, etc. Dose 1 to 2 drops (0.065-0.13).

Tilia (*til'-e-ah*) [L.]. A genus of exogenous trees—the linden, or basswood.

Timbre (*tam'-ber*) [Fr.]. The peculiar quality of a tone, other than pitch and intensity, that makes it distinctive. It depends upon the overtones of the vibrating body.

Tin. A silvery-white, metallic, malleable element, having a specific gravity of 7.3, an atomic weight of 118, an atomicity of two or four. Symbol Sn, from the Latin stannum.

T. Chlorid, stannous chlorid, $\text{SnCl}_2 + 2\text{H}_2\text{O}$, is used as a reagent.

Tincture (*tingk'-tūr*) [*tinctura*, from *tingere*, to tinge]. 1. A solution of the medicinal principles of a substance in a fluid other than water or glycerol. 2. Specifically, an alcoholic solution of a medicinal substance. **T., Ammoniated**, one made with ammoniated alcohol. **T., Ethereal**, one made with ether.

Tinea (*tin'-e-ah*) [L., a moth, a worm]. Ringworm; a generic term applied to a class of skin-diseases caused by parasitic fungi, formerly applied to many spreading cutaneous diseases. **T. amiantacea**, **T. asbestina**, seborrhea. **T. circinata**. See *T. trichophytina*. **T. cruris**. See *T. trichophytina*. **T. decalvans**, alopecia areata. **T. favosa**, favus. **T. furfuracea**, seborrhea sicca. **T. imbricata**, Tokelau ringworm, Bowditch Island ringworm, a disease occurring in the East Indies, and characterized by the formation of concentric scaly patches and intense itching. **T. kerion**, a markedly inflammatory form of ringworm of the scalp (*T. tonsurans*), giving rise to the formation of an edematous, boggy swelling discharging a mucoid secretion. **T. lupinosa**, favus. **T. sycosis**. See *T. trichophytina*. **T. tarai**, blepharitis ulcerosa. **T. tonsurans**. See *T. trichophytina*. **T. trichophytina**, ringworm, a spreading, contagious disease of the skin due

to a vegetable fungus, the *Trichophyton*. On the nonhairy parts of the body (*T. trichophytina corporis*, *T. circinata*) it presents itself by spreading, scaly patches, tending to clear in the center. On the thighs and scrotum (*T. trichophytina cruris*) it is apt to assume the appearance of eczema, hence it is also called *Excema marginatum*. The nails may be affected (*T. trichophytina unguium*), becoming grayish, opaque, and brittle. On the scalp (*T. trichophytina capitis*, *T. tonsurans*) it forms rounded, grayish, slightly elevated, scaly patches, with brittleness and loss of the hair. Ringworm of the bearded region (*T. trichophytina barbae*, *T. sycosis*, parasitic sycosis, barber's itch) forms at first rounded, scaly patches, which soon become nodular and lumpy, and tend to break down. **T. versicolor**, a disease of the skin due to a vegetable parasite, the *Microsporon furfur*. It is characterized by brownish-yellow macules that coalesce to form extensive areas of eruption. There is usually slight itching.

Tingible (*tinj'-ibl*) [*tingere*, to tinge]. Capable of being stained; stainable.

Tinkling (*tingk'-ling*). A chinking sound, heard over a pneumothorax or a large pulmonary cavity; also called metallic T.

Tinnitus (*tin-i'-tus* or *tin'-it-us*) [*tinnire*, to tinkle]. A subjective ringing, roaring, or hissing sound heard in the ears, also called **T. aurium**.

Tisane (*te-zahn'*). See *Plisan*.

Tissue (*tish'-u*) [Fr., *tissu*, from L., *texere*, to weave]. An aggregation of similar cells and fibers, forming a distinct structure, and entering as such into the formation of an organ or organism. **T., Adipose**, fatty tissue, a form of connective T. consisting of fat-cells lodged in the meshes of areolar T. **T., Areolar**, a form of connective T. consisting of cells and delicate, elastic fibers interlacing in every direction. **T., Cancellous**, the spongy T. of bones. **T., Cartilaginous**. See *Cartilage*. **T., Connective**, a general term for all those tissues of the body that support the essential elements or parenchyma. The most important varieties are adipose T., areolar T., osseous T., cartilaginous T., elastic T., fibrous T., lymphoid T. **T., Epithelial**. See *Epithelium*. **T., Erectile**, a spongy T. that becomes expanded and hard when filled with blood. **T., Lymphoid**, a form of connective T. in which reticular meshes contain lymphoid cells. **T., Mucous**, a connective T. such as is present in the umbilical cord of the fetus. **T., Muscular**. See *Muscle*. **T., Nervous**. See *Nerve*. **T., Osseous**. See *Bone*.

Titillation (*tit-il-a'-shun*) [*titillare*, to tickle]. The act of tickling; the sensation produced by tickling.

Titration (*tit-ra'-shun*) [Fr., *titre*, standard of fineness]. Volumetric analysis by the aid of standard solutions.

Titubation (*tit-u-ba'-shun*) [*titubare*, to stagger]. A staggering gait seen especially in diseases of the cerebellum.

Tl. Symbol of thallium.

Tobacco (*to-bak'-o*) [Sp., *tabaco*, tobacco]. The *Nicotiana tabacum*, a plant of the order Solanaceæ, the dried leaves of which (*Tabacum*, U. S. P., *Tabaci folia*, B. P.) contain a liquid alkaloid, nicotine, $C_{10}H_{14}N_2$, which is also present in the seeds and root. Nicotine is one of the most active poisons known. Tobacco-smoke contains a series of bases, among which are pyridin, picolin, lutidin, collidin, parvolin, and others. T. is used as a sedative in nearly all parts of the world, being smoked, chewed, or used as snuff. Its physiologic action is that of a nauseant, antispasmodic, and depressant; it is also a local irritant. In medicine it has been employed as a relaxant in intestinal obstruction, being given in the form of an enema, but it is now rarely used except in asthma and locally in hemorrhoids. **T.-amblyopia**, amblyopia produced by the prolonged and excessive use of T. **T.-heart**, an irritable state of the heart, characterized by irregular action and palpitation, produced by excessive indulgence in T. **T.**, Indian. See *Lobelia*.

Tocodynamometer (*tok-o-di-nam-on'-et-er*) [*rókos*, birth; *dynamometer*]. An instrument for measuring the force of the expulsive efforts of the uterus in childbirth.

Tocology (*tok-ol'-o-je*) [*rókos*, childbirth; *lógos*, science]. The science of obstetrics.

Toe [AS., *tā*]. A digit of the foot. **T.-reflex**. See *Reflexes*, *Table of*.

Toilet (*toi'-let*) [OF., *toilette*, a cloth]. In surgery, the cleansing, washing, and dressing of an operative wound.

Tokelau Ringworm. See *Tinea imbricata*.

Tolerance (*tol'-er-ans*) [*tolerare*, to bear]. The ability of enduring the influence of a drug or poison, particularly when acquired by a continued use of the substance.

Tollen's Test. A test for glucose, consisting in the reduction of an ammoniacal solution of silver nitrate to metallic silver.

Tolu (*to-lu'*) [Santiago de *Tolu* in New Granada, where it was first obtained]. Balsam of tolu (*Balsamum toltanum*, U. S. P., B. P.), a balsam obtained from *Toluifera balsamum*, of the order Leguminosæ. It is used as a stimulant expectorant. Dose gr. x-xxx (0.65-2.0). Preparations and doses: Syrupus toltanus, U. S. P., B. P., fʒj (4.0); Tinctura toltana, U. S. P., B. P., fʒj-ij (4.0-8.0).

Toluene (*to-lu'-u-en*) [*tolu*], C_7H_8 . Methyl-

benzene; a hydrocarbon obtained from coal-tar, and also produced in the dry distillation of tolu-balsam and many resins.

Toluidin (*tol-u'-id-in*) [*tolu*], $C_7H_7.NH_2$. A homologue of anilin, prepared from toluene.

Toluol (*tol'-u-ol*). Same as *Toluene*.

Tomentum (*to-men'-tum*) [L.]. A lock of wool. **T. cerebri**, the network of small blood-vessels of the pia penetrating the cortex of the brain.

Tome's Fibers. See *Fibers of Tome*.

Tomomania (*tom-o-ma'-ne-ah*) [*rough*, a cutting; *mania*, madness]. An excessive desire to perform operations.

Tone (*tōn*) [*róvos*, from *reivew*, to stretch]. 1. A distinct sound. 2. The normal state of tension of a part or of the body.

Tonga (*tong'-gah*). A mixture of various barks, probably of *Premna taitensis* and *Raphidophora vitiensis*, brought from the Feejee Islands, and used in neuralgia. Dose of a fluid extract fʒss (2.0).

Tongue (*tung*). The movable muscular organ attached to the floor of the mouth, and concerned in tasting, masticating, swallowing, and speaking. It consists of a number of muscles, and is covered by mucous membrane from which project numerous papillæ, and in which are placed the terminal organs of taste. **T., Black**, a condition in which the dorsal surface of the tongue is covered with a black coating. **T.-depressor**, a spatula for pushing down the tongue during the examination of the mouth and throat. **T., Fern-leaf Pattern**, a name given to a T. presenting a well-marked central furrow (midrib) with lateral branches. **T., Filmy**, one with whitish, symmetric patches on both sides. **T., Furred**, a coated T., the papillæ of which are prominent, giving the mucous membrane the appearance of a whitish fur. **T., Geographic**, one with localized thickening of the epithelium, giving to the surface the appearance of a geographic chart. **T., Hairy**, one with a hyperplasia of the papillæ, giving rise to hair-like projections. **T., Strawberry**, a hyperemic T., the fungiform papillæ of which are very prominent; it is seen especially in scarlet fever. **T.-swallowing**, a condition in which there is an abnormal mobility of the organ, so that it falls backward, giving rise to danger of suffocation. **T.-tie**, a congenital shortening of the frenum of the T., interfering with its mobility; ankyloglossia. **T., Wooden**, on the seat of actinomycosis.

Tonic (*ton'-ik*) [*tonos*, tone]. 1. Pertaining to tone; producing normal tone or tension. 2. An agent producing normal tone of an organ or part. **T., Cardiac**, one strengthening the heart-muscle. **T., Intestinal**, one strengthening the tone of the in-

testines. **T.**, Nervine, one increasing the tone of the nervous system. **T.**, Stomachic, one increasing the tone of the stomach. **T.**, Vascular, one increasing the tone or tension of the blood-vessels. 3. Characterized by continuous tension or contraction, as, *e. g.*, a **T.** spasm, **T.** convulsion.

Tonicity (*to-nis'-it-e*) [*tróvos*, tone]. The condition of normal tone or tension of organs; a state of tone.

Tonka Bean (*tong'-kah ben*). The seed of *Dipteryx odorata*, a tree of South America; it contains coumarin, and is used as a flavoring agent.

Tonometer (*ton-om'-et-er*) [*tróvos*, tone, tension; *μέτρον*, measure]. An instrument to measure tension, as, *e. g.*, that of the eyeball.

Tonsil (*ton'-sil*) [*tonsilla*]. 1. A small, almond-shaped body, situated on each side of the fauces, between the anterior and posterior pillars of the soft palate. It consists of an aggregation of from ten to eighteen lymph-follicles, and is covered by mucous membrane, which dips into certain depressions called crypts. 2. A small lobe of the cerebellar hemisphere, situated on the inferior mesial aspect. **T.**, Faucial. See *Tonsil*, 1st def. **T.**, Lingual, an accumulation of lymphadenoid tissue at the base of the tongue. **T.**, Pharyngeal, a mass of lymphadenoid tissue in the pharynx, between the Eustachian tubes.

Tonsillar (*ton'-sil'-ar*) [*tonsilla*, tonsil]. 1. Pertaining to the tonsil, as, *e. g.*, the **T.** artery. 2. Affecting the tonsil, as, *e. g.*, **T.** abscess.

Tonsillitis (*ton-sil'-it-is*) [*tonsilla*, tonsil; *itis*, inflammation]. Inflammation of the tonsil. **T.**, Follicular, a form in which the follicles are especially involved and project as whitish points from the surface of the tonsil. **T.**, Herpetic, a form characterized by an eruption of herpetic vesicles, which soon rupture, leaving small, circular ulcers that coalesce and become covered with a fibrinous exudation. The disease has an acute onset, a continuous fever, and a critical decline, affects those subject to herpes elsewhere, and tends to recur. **T.**, Lacunar. Same as **T.**, Follicular. **T.**, Mycotic, a form of **T.** due to fungi. **T.**, Pustular, one characterized by the formation of pustules, as in small-pox. **T.**, Suppurative. Synonym of *Quinsy*.

Tonsillotome (*ton'-sil-o-tóm*) [*tonsilla*, tonsil; *τομή*, a cutting]. An instrument for removing or cutting off the tonsil.

Tonsillotomy (*ton-sil'-o'-o-me*) [*tonsilla*, tonsil; *τομή*, a cutting]. The operation of cutting out the tonsil.

Tonus (*to'-nus*) [*tróvos*, tone]. See *Tone*.

Tooth (*tooth*) [*AS.*, *toð*, tooth]. One of the small, bone-like organs occupying the alve-

olar processes of the upper and lower jaws, and serving for tearing and comminuting the food. The teeth begin to appear in the human being about the seventh month; by the end of the third year, the eruption of the so-called temporary, deciduous, or milk-teeth, numbering 20, is completed. The permanent teeth begin to replace the deciduous teeth about the seventh year. In the adult, the permanent teeth number 32, or 16 in each jaw, and are divided as follows: 2 incisors, 1 canine, 2 bicuspids, and 3 molars in each lateral half of the jaw. Each tooth is composed of a crown, the exposed part, a constricted part, called the neck, and a part within the alveolus, called the fang or root. In structure a tooth consists of an outer hard substance, the enamel, encasing the crown; the dentine, within the enamel; and the pulp, a soft vascular tissue filling the pulp-cavity. The dentine of the root is surrounded by the cement or crusta petrosa. **T.**, Eye-, the upper canine tooth. Teeth, Hutchinson's. See *Hutchinson's Teeth*. **T.**, Stomach-, the lower canine tooth. **T.**, Wisdom, the third molar tooth.

Tophaceous (*to-fa'-shus*) [*tróvos*, stone]. Of the nature of tophi; sandy, or gritty.

Tophus (*to'-fus*) [*tróvos*, stone]. 1. The hard, stone-like deposits occurring in gout, especially about the knuckles and the cartilages of the ear, and consisting of sodium urate. 2. The tartar of the teeth.

Topic, Topical (*top'-ik*, *top'-ik-al*) [*tróvos*, place]. Local.

Topoalgia (*top-o-al'-je-ah*) [*tróvos*, place; *ἄλγος*, pain]. Localized pain, common in neurasthenia, and often appearing suddenly after emotional disturbances.

Topographic (*top-o-graf'-ik*) [*tróvos*, place; *γράφειν*, to write]. Pertaining to a locality. **T.** Anatomy, the study of the regions occupied by a part, or in which anything occurs.

Topography (*top-og'-ra-fe*) [*tróvos*, place; *γράφειν*, to write]. A study of the regions of the body or its parts, as, *e. g.*, cerebral **T.**

Torcular Herophili (*tor'-ku-lar her-off'-il-i*) [*L.*, the wine-press of Herophilus]. The expanded extremity of the superior longitudinal sinus, placed in a depression on the inner surface of the occipital bone. It receives the blood from the occipital sinus, and from it is derived the lateral sinus of the side to which it is deflected.

Tormen. See *Tormina*.

Tormina (*tor'-min-ah*) [*pl.* of *tormen*, a racking pain]. Gripping pains in the bowels.

Tornwaldt's Disease. See *Thornwaldt's Disease*.

Torpid (*tor'-pid*) [*torpere*, to be numb]. Affected with torpor.

Torpidity (*tor-pid'-it-e*). See *Torpor*.

Torpor (*tor'-por*) [*L.*]. Sluggishness, inactivity. **T. retinæ**, dulled perceptive power of the retina.

Torrefaction, Torrefication (*tor'-e-fak'-shun, tor'-e-fik'-a'-shun*) [*torrefacere*, to dry by heat]. Roasting; drying by means of high heat.

Torricellian Vacuum (*to-re-tshel'-e-an*) [*Torricelli*, an Italian physicist]. The vacuum above the mercury-column of a barometer.

Torsion (*tor'-shun*) [*torquere*, to twist]. A twisting.

Torticollis (*tor-te-coll'-is*) [*tortus*, twisted; *collum*, neck]. Wry-neck, a contraction of one or more of the cervical muscles, usually of one side, resulting in an abnormal position of the head. **T., Rheumatic**, stiff-neck, a form due to rheumatism of the sternocleidomastoid or other muscle of the neck. **T., Spasmodic**, spasmodic contraction of the muscles of the neck of one side, especially the sternocleidomastoid, causing a drawing of the head toward the opposite side.

Torula (*tor'-u-lah*) [*torulus*, a small tuft].

1. A genus of fungi reproducing by budding, many species of which are alcoholic ferments. **T. cerevisiæ**. See *Saccharomyces cerevisiæ*. 2. A chain of spheric bacteria.

Torus (*tor'-us*) [*L.*]. 1. A surface having a regular curvature, with two principal meridians of dissimilar curvature at right angles to each other. 2. An elevation or prominence. 3. The tuber cinereum of the brain. **T. frontalis**, a protuberance in the region of the frontal sinuses, at the root of the nose. **T. palatinus**, a protuberance on the surface of the hard palate, marking the point of junction of the intermaxillary and palatomaxillary sutures.

Touch (*tuch*) [*Fr., toucher*]. 1. The tactile sense. 2. The act of judging by the tactile sense; palpation, especially palpation of a part with the finger, as rectal **T.**, vaginal **T.** **T.-corpuscle**, a tactile corpuscle.

Tour de Maitre (*toor-d'-mâtr*) [*Fr.*, the master's turn]. A method of passing a catheter into the bladder in which it is introduced into the urethra with the convexity upward, the shaft lying obliquely across the left thigh of the patient, and as the point enters the bulb, the handle is swept around toward the abdomen, when the beak passes into the membranous urethra, and is carried into the bladder by depressing the shaft between the patient's thighs. A sound may be introduced into the uterus in an analogous manner, by entering the instrument with the convexity upward, and then sweeping the shaft around.

Tourette's Disease [Gilles de la Tourette, a French physician]. A convulsive form of

tic, characterized by motor incoordination with echolalia and coprolalia.

Tourniquet (*toor'-nik-et*) [*Fr.*, from *tourner*, to turn]. An instrument for controlling the circulation by means of compression, usually consisting of two metallic plates united by a thumb-screw, and a strap provided with a pad. The strap is fastened about the part, the pad being made to lie over the artery to be occluded. The screw is placed diametrically opposite the pad, and the strap tightened by separating the metallic plates of the screw.

T., Dupuytren's, one for compressing the abdominal aorta, consisting of a semicircle of metal with a pad at one extremity. **T., Esmarch's**, one consisting of a stout, elastic rubber band applied above the proximal turn of an elastic bandage passing around the part to be rendered bloodless. **T., Field's**, one consisting of a strap and buckle with a pad to be placed over the artery. **T., Horse-shoe**, one shaped like a horse-shoe, to compress only two points, and thus permit venous return. **T., Provisional**, one applied loosely, so that it may be tightened at once upon the recurrence of hemorrhage. **T., Signorini's**. See *T., Horse-shoe*.

Tow. The coarse part of flax or hemp, used as an absorbent.

Tox-, Toxic-, Toxo- (*toks-, toks'-ik-o-, toks'-o-*) [*τοξικόν*, poison]. A prefix signifying poisonous, or caused by a poison.

Toxalbumin (*toks-al-bu'-min*) [*τοξικόν*, poison; *albumin*]. A poisonous proteid. Toxalbumins have been obtained from cultures of bacteria and from certain plants.

Toxemia (*toks-e'-me-ah*) [*τοξικόν*, poison; *aima*, blood]. Blood-poisoning, a condition of the blood in which it contains poisonous products, either those produced by the body-cells or those due to the growth of microorganisms.

Toxemic (*toks-em'-ik, toks-e'-mik*) [*τοξικόν*, poison; *aima*, blood]. Pertaining to, affected with, or caused by toxemia.

Toxic (*toks'-ik*) [*τοξικόν*, poison]. Poisonous; produced by a poison.

Toxicant (*toks'-ik-ant*) [*τοξικόν*, poison]. 1. Poisonous or toxic. 2. A poisonous agent.

Toxicity (*toks-is'-it-e*) [*τοξικόν*, poisonous]. The quality of being toxic.

Toxicodendron (*toks-ik-o-den'-dron*). See *Rhus*.

Toxicoderma (*toks-ik-o-der'-mah*) [*τοξικόν*, poison; *derma*, skin]. Disease of the skin due to poison.

Toxicogenic (*toks'-ik-o-jen'-ik*) [*τοξικόν*, poison; *γεννάν*, to produce]. Producing poisons.

Toxicohemia (*toks'-ik-o-he'-me-ah*). See *Toxemia*.

Toxicology (*toks-ik-ol'-o-jé*) [*τοξικόν*, poison;

λόγος, science]. The science of the nature and effects of poisons, their detection, and the treatment of their effects.

Toxicomania (*toks-ik-o-ma'-ne-ah*) [τοξικόν, poison; μανία, madness]. 1. Morbid desire to consume poison. 2. Toxiphobia.

Toxicosis (*toks-ik-o'-sis*) [τοξικόν, poison]. A state of poisoning.

Toxiferous (*toks-if'-er-us*) [τοξικόν, poison; *ferre*, to bear]. Producing or conveying poison.

Toxin (*toks'-in*) [τοξικόν, poison]. 1. Any poisonous substance, albuminous in nature, produced by the action of bacteria on organic matter. 2. Any poisonous substance, either albuminous or basic, produced by the action of bacteria on organic matter.

Toxiphobia (*toks-e-fó'-be-ah*) [τοξικόν, poison; φόβος, fear]. Morbid dread of being poisoned.

Toxophylaxin (*tox-o-fil-aks'-in*) [τοξικόν, poison; φύλαξ, a protector]. See *Phylaxin*.

Toxozoin (*toks-o-zó'-zin*) [τοξικόν, poison; σώζω, to save]. See *Sozin*.

Toynbee's Experiment [*Toynbee*, an English otologist]. The rarefaction of the air of the middle ear by swallowing, with coincident closure of the mouth and nares.

Trabecula (*tra-bek'-u-lah*) [L., a small beam; *pl.*, *Trabecula*]. Any one of the fibrous bands extending from the capsule into the interior of an organ.

Trabs cerebri (*trabs ser'-eb-ri*) [L.]. The corpus callosum.

Trace (*trás*). A mark. T., Primitive. See *Primitive Trace*.

Tracheal (*tra'-ke-al*) [τραχεία, trachea]. Pertaining to, or produced in the trachea.

T. Tugging, the downward tugging movement of the larynx, sometimes observed in aneurysm of the aortic arch.

Trachealis Muscle (*trak-e-a'-lis*) [τραχεία, trachea]. The intrinsic transverse muscle-fibers found in the trachea.

Tracheitis (*tra-ke-i'-tis*) [τραχεία, trachea; *ιτις*, inflammation]. Inflammation of the trachea.

Trachelo- (*trak'-el-o-*) [τράχηλος, neck]. A prefix denoting pertaining to the neck.

Trachelomastoid (*trak'-el-o-mas'-toid*) [τράχηλος, neck; *mastoid*]. Pertaining to the neck and the mastoid process.

Tracheloplasty (*trak'-el-o-plas-te*) [τράχηλος, neck; *πλάσσειν*, to mold]. Plastic operation on the neck of the uterus.

Trachelorrhaphy (*trak-el-or'-a-fe*) [τράχηλος, neck; *ρῑφή*, suture]. Repair of a laceration of the cervix uteri.

Trachelotomy (*trak-el-ot'-o-me*) [τράχηλος, neck; *τομή*, a cutting]. Incision into the cervix uteri.

Tracheobronchitis (*tra-ke-o-brong'-ki'-tis*)

[τραχεία, trachea; βρόγχος, bronchus; *ιτις*, inflammation]. Inflammation of the trachea and bronchi.

Tracheocele (*tra'-ke-o-sel*) [τραχεία, trachea; *κήλη*, tumor]. 1. Protrusion of the mucous membrane of the trachea. 2. Goiter.

Tracheoesophageal (*tra-ke-o-e-sof-aj'-e-al*) [τραχεία, the trachea; οισοφάγος, esophagus]. Pertaining to the trachea and the esophagus.

Tracheolaryngeal (*tra-ke-o-lar-in'-je-al*) [τραχεία, trachea; *λάρυγξ*, larynx]. Pertaining to the trachea and the larynx.

Tracheoscopy (*tra-ke-os'-ko-pe*) [τραχεία, trachea; *σκοπεῖν*, to inspect]. Inspection of the interior of the trachea by means of a laryngoscopic mirror and reflected light.

Tracheostenosis (*tra-ke-o-sten-o'-sis*) [τραχεία, trachea; *στένωσις*, stenosis]. Stenosis of the trachea.

Tracheotomy (*tra-ke-ot'-o-me*) [τραχεία, trachea; *τομή*, a cutting]. The operation of cutting into the trachea through the cricothyroid membrane, or through the cricoid cartilage and the upper part of the trachea. T., Inferior, one performed below the isthmus of the thyroid gland. T., Superior, one performed above the isthmus of the thyroid gland. T.-tube, a metal tube placed in the opening made in tracheotomy, and through which breathing is carried on.

Trachoma (*tra-ko'-mah*) [τραχίς, rough]. Granular conjunctivitis, granular lids; a contagious disease of the eyelids characterized by small, sago-like elevations on the conjunctiva, and later by cicatricial contraction and deformity of the lids. The friction of the elevations (*T. granulations*) against the cornea often produces ulcer or pannus.

Trachomatous (*tra-ko'-mat-us*) [τραχίς, rough]. Affected with or pertaining to trachoma.

Tract [*tractus*]. A distinct, more or less defined region having considerable length, especially a collection of fibers, in the central nervous system.

Traction (*trak'-shun*) [*trahere*, to draw]. The act of drawing or pulling. T.-aneurysm, an aneurysm due to traction on the aorta by an incompletely atrophied ductus Botalli. T., Axis-, traction in the axis or direction of a channel, as of the pelvis, through which a body is to be drawn. Axis-T. Forceps, an obstetric forceps for performing axis-traction in the delivery of the fetus. T.-diverticulum, a circumscribed sacculatation of the esophagus from the traction of adhesions. T., Elastic, traction by an elastic force.

Tractor (*trak'-tor*) [*trahere*, to draw]. 1. An instrument for making traction. 2. See *Perkinsism*.

Tractoration (*trak-tor-a'-shun*) [*trahere*, to

draw]. Treatment by metallic tractors. See *Perkinsism*.

Tractus (*trak'-tus*). See *Tract*.

Tragacanth (*trag'-a-kanth*) [*τραγάκανθα*: *τραγος*, goat; *ἀκανθα*, thorn]. A gummy exudation from various species of *Astragalus*, of the order Leguminosæ, constituting the Tragacantha of the U. S. P. and B. P. It resembles gum-arabic, and is used as a demulcent, and is added to water to suspend insoluble powders and for making troches. Preparations: Mucilago tragacanthæ, U. S. P., B. P.; Pulvis tragacanthæ compositus, B. P., dose gr. xxx-5j (2.0-4.0); Glycerinum tragacanthæ, B. P.

Tragicus (*traj'-ik-us*). See *Muscles, Table of*.

Tragus (*tra'-gus*) [*τράγος*, goat]. The small prominence of cartilage projecting over the meatus of the external ear.

Trance (*trans*) [*transire*, to pass over]. A form of abnormal sleep from which the patient ordinarily cannot be aroused, and which is dependent upon hysteria (hysterical T.) or hypnotism.

Trans- [*trans*, across]. A prefix denoting through or across.

Transsection (*trans-sek'-shun*) [*trans*, across; *secare*, to cut]. A section made across the long axis of a part.

Transfer (*trans'-fer*) [*trans*, across; *ferre*, to carry]. The change of anesthesia or hyperesthesia from one part of the body to another, a phenomenon present in some cases of hysteria.

Transference (*trans-fer'-ens*) [*trans*, across; *ferre*, to carry]. 1. See *Transfer*. 2. See *Thought-transfer*.

Transfix (*trans-fik'-s*) [*trans*, across; *figere*, to fix]. To pierce through and through.

Transfixion (*trans-fik'-shun*) [*trans*, across; *figere*, to fix]. The act of piercing through and through.

Transforation (*trans-for'-a'-shun*) [*trans*, across; *forare*, to pierce]. The act of perforating, as, e. g., T. of the fetal skull.

Transformation (*trans-for-ma'-shun*) [*trans*, across; *formare*, to form]. A change of form or constitution; degeneration.

Transfusion (*trans-fu'-shun*) [*trans*, across; *fundere*, to pour]. The pouring of liquid from one vessel to another; especially the introduction into a vessel of the body of blood or other liquid. T., Arterial, T. of blood into an artery. T., Direct or Immediate, the T. of blood from one person to another without exposure of the blood to the air. T., Indirect or Mediate, the introduction of blood that was first drawn into a vessel. T., Venous, T. into a vein.

Transillumination (*trans-il-lu-min-a'-shun*) [*trans*, through; *illuminare*, to illuminate].

Illumination of the walls of a cavity by a light passed through them.

Translation (*trans-la'-shun*) [*trans*, across; *ferre*, to bear]. A change of location.

Translucent (*trans-lu'-sent*) [*trans*, across; *lucere*, to shine]. Permitting a partial transmission of light; somewhat transparent.

Transmigration (*trans-mi-gra'-shun*) [*trans*, across; *migrare*, to wander]. A wandering across or through; as, e. g., T. of the ovum; T. of the white corpuscles. T., External, the passage of an ovum from one ovary to the opposite oviduct without traversing the uterus. T., Internal, the passage of the ovum through its proper oviduct into the uterus and across to the opposite oviduct.

Transmission (*trans-mish'-un*) [*trans*, across; *mittere*, to send]. The communication or transfer of anything, especially disease, from one person or place to another.

Transparent (*trans-pa'-rent*) [*trans*, across; *parere*, to appear]. Having the property of permitting the passage of light-rays without material obstruction, so that objects beyond the body can be seen.

Transpiration (*trans-pir-a'-shun*) [*trans*, across; *spirare*, to breathe]. 1. The act of exhaling fluid or gas through the skin. 2. The material exhaled.

Transplantation (*trans-plan-ta'-shun*) [*trans*, across; *plantare*, to plant]. The operation of transplanting or of applying to a part of the body tissues taken from another body or from another part of the same body. See *Graft*.

Transposition (*trans-po-zish'-un*) [*trans*, across; *ponere*, to place]. A change of position. T. of the Viscera, a change in the position of the viscera whereby they are placed on the side opposite to that normally occupied.

Transudate (*trans-u-dāt*) [*trans*, through; *sudare*, to perspire]. A liquid or other substance produced by transudation.

Transudation (*trans-u-da'-shun*) [*trans*, through; *sudare*, to perspire]. 1. The passing of fluid through a membrane, especially of blood serum through the vessel-walls. 2. Transudate.

Transversalis (*trans-ver-sa'-lis*) [*trans*, across; *vertere*, to turn]. Transverse; an artery (T. colli) or a muscle (T. abdominis) running transversely. See *Arteries, Table of*, and *Muscles, Table of*. T. Fascia, the fascia on the inner surface of the T. muscle, between the latter and the peritoneum.

Transverse (*trans-vers')* [*trans*, across; *vertere*, to turn]. Cross-wise; at right angles to the longitudinal axis of the body. T. Presentation, a presentation of the fetus at right angles to the longitudinal axis of the uterus.

Transversus (*trans-ver'-sus*) [L.]. Transverse, as T. muscle. See *Muscles*, Table of.

Trapeziometacarpal (*tra-pe-ze-o-met-ah-kar'-pal*) [*trapezium*; *metacarpal*]. Pertaining to the trapezium and the metacarpus.

Trapezium (*tra-pe'-ze-um*) [τράπεζα, table]. 1. The first bone of the second row of carpal bones. 2. A tract of transverse fibers situated in the lower part of the pons, enclosing the superior olivary nucleus, and connected with the accessory auditory nucleus.

Trapezium (*tra-pe'-ze-us*) [τράπεζα, table]. See *Muscles*, Table of.

Trapezoid (*trap'-es-oid*) [τράπεζα, a table; εἶδος, like]. A geometric four-sided figure having two parallel and two diverging sides.

T. Bone, or simply **T.**, the second bone of the second row of the carpus. **T. Ligament**. See *Ligaments*, Table of.

Trapp's Formula. A formula employed in the estimation of the quantity of solid matters in urine. The last two figures of the specific gravity are multiplied by 2 (Trapp's coefficient) or by 2.33 (Häser's coefficient).

Traube-Rosenstein Theory (*traw-be-ro'-sen-shtn*). A theory that puerperal eclampsia is due to an anemia brought about by an edema of the brain, the result of some poison circulating in the blood.

Traube's Semilunar Space (*traw'-bes*). See *Semilunar*. **T. Curves**, **T.-Hering's Curves**, curves in the tracing of the blood-pressure occurring independent and after the cessation of the respiration.

Trauma (*traw'-mah*) [τραῦμα, a wound]. A wound; an injury.

Traumatic (*traw-mat'-ik*) [τραῦμα, a wound]. Pertaining to or caused by a wound or injury.

Traumaticin (*traw-mat'-is-in*) [τραῦμα, trauma]. A solution of gutta-percha in chloroform, used for closing superficial wounds.

Traumatism (*traw'-mat-izm*) [τραῦμα, wound]. 1. The condition produced by trauma. 2. Improperly, trauma.

Treacle (*tre'-akl*). The uncrystallized residue remaining after the refining of cane-sugar.

Treatment (*trét'-ment*) [tractare, to treat]. The means employed in effecting the cure of disease; the management of disease or of diseased patients.

Trehalose (*tre-hal'-os*), $C_{12}H_{22}O_{11}$. A carbohydrate resembling sugar, derived from ergot and from Trehala manna.

Trematode (*trem'-at-ód*) [τρήμα, hole; εἶδος, like]. A member of the Trematoda, a class of worms, some of which are parasitic in man and the lower animals.

Tremor (*trem'-or*) [tremere, to shake]. A trembling of the voluntary muscles. **T.**, **Fibrillary**, tremor caused by consecutive con-

tractions of separate muscle-fibrillæ. **T.**, **Intention-**, one appearing on voluntary movement. **T.**, **Purring**. Synonym of *Thrill*, *Purring*.

Tremulous (*trem'-u-lus*) [tremere, to shake]. Trembling, quivering, as, e. g., T. iris.

Trendelenburg Posture. See *Postures*, Table of.

Trepine (*tre-fín'*) [τρίπανον, from τρυπᾶν, to bore]. An instrument for cutting out a circular piece of bone, usually from the skull.

Trephining (*tre-fí'-ning*) [trephine]. The operation of cutting out a piece of bone with a trephine.

Trepidation (*trep-id-a'-shun*) [trepidare, to tremble]. 1. Trembling. 2. A peculiar oscillatory movement at times seen in the muscles after hemiplegia.

Tri- [τρεῖς, or tres, three]. A prefix denoting three.

Triacid (*tri-as'-id*) [tres, three; acidum, acid]. Of an alcohol, containing three atoms of hydrogen replaceable by a base.

Triad (*tri'-ad*) [tres, three]. See *Quantivalence*.

Trial-glasses, or **T.-lenses**. A graduated set of concave, convex, cylindric, and prismatic lenses used in testing vision.

Triallylamin (*tri-al-il-am'-in*). A volatile base having the formula $(C_3H_5)_3N$.

Triamin (*tri-am'-in*). A compound derived from three molecules of ammonia in which the hydrogen has in part or wholly been replaced by bases.

Triangle (*tri'-ang-gl*) [tres, three; angulus, angle]. A figure having three sides and three angles. **T. of Bryant**. See *T., Iliofemoral*. **T.**, **Carotid**, **Inferior** (*T. of necessity*), located in the neck; it is bounded in front by the median line of the neck, behind by the anterior margin of the sternomastoid, and above by the anterior belly of the omohyoid. Its floor is formed by the longus colli below and the scalenus anticus above. The common carotid artery, internal jugular vein, vagus nerve, superficialis colli nerve, a branch of the communicans noni, the inferior thyroid artery, the recurrent laryngeal nerve, the sympathetic nerve, the trachea, thyroid gland, and larynx are the important structures within it. **T.**, **Carotid**, **Superior** (*T. of election*), a T. located in the neck; it is bounded behind by the sternomastoid, in front by the anterior belly of the omohyoid, and above by the posterior belly of the digastric. Its floor is formed by the thyrohyoid, hyoglossus, and inferior and middle constrictors of the pharynx. The important structures contained within it are the common carotid artery and its bifurcation into the external and internal carotids, the superior thyroid artery, the lingual artery, the facial artery,

the occipital and ascending laryngeal arteries, the internal jugular vein and the veins corresponding to the arteries mentioned, the descendens noni, hypoglossal, pneumogastric, sympathetic, spinal accessory, superior laryngeal, and external laryngeal nerves. **T., Cephalic**, a T. on the anteroposterior plane of the skull, bounded by lines joining the occiput with the forehead and with the chin and a line joining the latter two. **T., Digastric**. See *T., Submaxillary*. **T. of Elbow**, a T. lying in front of the elbow with the base directed upward toward the humerus, and bounded externally by the supinator longus and internally by the pronator radii teres. Its floor is formed by the brachialis anticus and supinator brevis. Its contents are the brachial artery and veins, the radial and ulnar arteries, the median and musculospiral nerves, and the tendon of the biceps. **T. of Election**. See *T., Carotid, Superior*. **T., Facial**, a T. formed by lines uniting the basion with the alveolar and nasal points and a line joining the latter two. **T., Frontal**, a T. bounded by the maximum frontal diameter and lines joining its extremities and the glabella. **T. of Hesselbach**, a T. at the lower part of the abdomen, bounded below by the deep epigastric artery, on the outer side by Poupart's ligament, and on the inner side by the margin of the rectus muscle. **T., Iliofemoral**, a T. located at the hip. Its hypothenuse is formed by Nélaton's line, a second side by the continuation outward of a line drawn through the two superior iliac spines, and the third by a line drawn at right angles to this from the summit of the greater trochanter. **T., Inferior Occipital**, a T. having the bimas-toid diameter for its base and theinion for its apex. **T., Infraclavicular**, a T. situated below the clavicle; it is bounded above by the clavicle, below and to the inner side by the upper border of the great pectoral muscle, and to the outer side by the anterior border of the deltoid. It contains the axillary artery. **T., Inguinal**. Same as *T., Scarpa's*. **T. of Lesser**, a T. located in the neck. Its boundaries are as follows: at its upper border the hypoglossal nerve; the two sides are formed by the anterior and posterior bellies of the digastric muscle. It is covered by the skin, superficial and deep fasciæ, and apex of the submaxillary gland. The floor is formed by the hyoglossus muscle. It contains the ranine vein and hypoglossal nerve. **T., Lumbocostoabdominal**, a T. bounded anteriorly by the external oblique, superiorly by the lower border of the serratus posticus inferior and the point of the 12th rib, posteriorly by the outer edge of the erector spinæ, and inferiorly by the internal oblique. **T.,**

Mylohyoid, the space bounded by the mylohyoid and the two bellies of the digastric. **T. of Necessity**. Same as *T., Carotid, Inferior*. **T. of the Neck, Anterior**, a triangle bounded anteriorly by a line extending from the chin to the sternum, posteriorly by the anterior margin of the sternomastoid, the base being formed by the lower border of the body of the inferior maxilla and a continuation of this line to the mastoid process of the temporal bone. It is subdivided into three smaller triangles by the digastric muscle above and the anterior belly of the omohyoid below. These are named from below upward the *inferior carotid*, the *superior carotid*, and the *submaxillary*. **T. of the Neck, Posterior**, a T. bounded anteriorly by the sternomastoid muscle, posteriorly by the anterior margin of the trapezius; the base is formed by the upper border of the clavicle; the apex corresponds to the occiput. It is divided by the posterior belly of the omohyoid muscle into two triangles, the *occipital* or *upper*, and the *subclavian* or *lower*. **T., Occipital**, a T. with the following boundaries: anteriorly, the sternomastoid muscle; posteriorly, the trapezius, and below, the omohyoid muscle. Its important contents are the spinal accessory nerve, the ascending and descending branches of the cervical plexus, and the transversalis colli artery and vein. **T. of Petit**, a T. bounded below by the crest of the ilium and laterally by the external oblique and latissimus dorsi. **T., Scarpa's**, a T. located in the proximal third of the thigh. The boundaries are externally the sartorius muscle, internally the adductor longus muscle, and above Poupart's ligament. Its important contents are the femoral artery and vein, the anterior crural nerve, and the crural branch of the genitocrural nerve. **T., Subclavian**, a T. bounded above by the posterior belly of the omohyoid muscle, below by the upper border of the clavicle. Its base is formed by the sternomastoid muscle. It contains the subclavian artery and occasionally the vein, the brachial plexus of nerves, the suprascapular vessels, the transversalis colli artery and vein, and the external jugular vein. **T., Submaxillary**, a T. formed above by the lower border of the body of the inferior maxilla and a continuation of this line to the mastoid process of the temporal bone, below by the posterior belly of the digastric and the stylohyoid muscle, and anteriorly by the middle line of the neck. It contains the submaxillary gland, the facial artery and vein, the submental artery, the mylohyoid artery and nerve, and the stylomaxillary ligament, behind which is the external carotid artery. **T., Suboccipital**, a T. in the posterior part of the neck,

formed by the rectus capitis posticus major, and superior and inferior oblique muscles, and containing the vertebral artery. **T.**, Suprameatal, of McEwen, a space bounded by the lower posterior edge of the zygomatic root and the superior posterior edge of the external auditory canal.

Triangular (*tri-ang'-gu-lar*) [*triangle*]. Having three sides or angles, as, *e.g.*, the T. ligament.

Triangularis (*tri-ang'-gu-lar-ris*). A triangular muscle. See *Muscles*, Table of.

Triatomic (*tri-at-om'-ik*) [*tres*, three; *atom*]. 1. Consisting of three atoms. 2. Having three atoms of replaceable hydrogen.

Tribadism (*trib'-ad-izm*) [*τριβειν*, to rub]. Sexual intercourse between women produced by friction of the genitals.

Tribasic (*tri-ba'-sik*) [*tres*, three; *basis*, a base]. Having three hydrogen-atoms replaceable by bases.

Tribrommethane (*tri-bröm-meth'-än*) [*tres*, three; *bromin*; *methane*]. Bromoform.

Tribromid (*tri-bröm'-id*) [*τριεις*, three; *βρωμος*, stench]. A compound of bromin containing three atoms of bromin to one of the base.

Tricalcic (*tri-kal'-sik*) [*tres*, three; *calcium*]. Containing three atoms of calcium.

Triceps (*tri'-seps*) [*tres*, three; *caput*, head]. Three-headed; a muscle having three heads of origin. See *Muscles*, Table of.

Trichauxia (*trik-avuk'-is*) [*θριξ*, hair; *αυξησις*, increase]. Hypertrichosis.

Trichiasis (*trik-i'-as-is*) [*θριξ*, a hair]. A state of abnormal position of the eyelashes, so that they produce irritation by friction upon the globe. **T. of the Anus**, an incurvation of the hairs about the anus, so that they irritate the mucous membrane.

Trichina (*trik-i'-nah*) [*θριξ*, a hair]. A genus of nematode worms, of which one species, the *T. spiralis*, is parasitic in the hog and at times in man. See *Trichinosis*.

Trichinosis (*trik-in-o'-sis*) [*trichina*]. A disease produced by the ingestion of pork containing the *Trichina spiralis*. It is characterized by nausea, vertigo, fever, diarrhea, prostration, stiffness and painful swelling of the muscles, edema of the face, and in some cases perspiration, insomnia, and delirium.

Trichinous (*trik'-in-us*) [*trichina*]. Infested with or containing trichinae.

Trichloracetic Acid (*tri-klör-as-e'-tik*). See *Acid*, *Trichloracetic*.

Trichlorid (*tri-klör'-id*) [*τριεις*, three; *chlorid*]. A compound containing chlorin in the proportion of three atoms to one of the base.

Trichlorphenol (*tri-klör-fu'-ne'*) [*τριεις*, three; *χλωρος*, green; *φαινωμι*, to show].

(OH). A derivative of phenol used as a disinfectant.

Tricho- (*trik'-o-*) [*θριξ*, hair]. A prefix denoting pertaining to a hair.

Trichocephalus (*trik-o-sef'-al-us*) [*θριξ*, hair; *κεφαλή*, head]. A genus of nematode worms, the thread-worms. *T. dispar* is parasitic in the intestines, especially the large intestine.

Trichoglossia (*trik-o-glos'-e-ah*) [*θριξ*, hair; *γλωσσα*, tongue]. Hairy tongue, a thickening of the papillae, producing an appearance as if the tongue were covered with hair.

Trichoid (*trik'-oid*) [*θριξ*, hair; *ειδος*, like]. Resembling hair.

Trichology (*trik-ol'-o-je*) [*θριξ*, hair; *λογος*, science]. The science of the hair.

Trichomatose (*trik-o'-mat-ös*) [*θριξ*, hair]. Matted together.

Trichomatosis (*trik-o-mat-o'-sis*) [*θριξ*, hair]. An affection of the hair characterized by a matted condition due to fungoid growths. See *Plica Polonica*.

Trichomonas (*trik-om'-o-nas*) [*θριξ*, hair; *μονάς*, a monad]. A genus of infusorians.

T. vaginalis, a species occasionally found in the vagina.

Trichomycosis (*trik-o-mi-ko'-sis*) [*θριξ*, hair; *μύκης*, a mushroom]. A disease of the hair produced by a vegetable parasite.

Trichonosis, **Trichonosus** (*trik-on-o'-sis*, *trik-on-o'-sus*) [*θριξ*, hair; *νόσος*, disease]. Any disease of the hair.

Trichopathy (*trik-op'-ath-e*) [*θριξ*, hair; *πάθος*, disease]. Any disease of the hair.

Trichophyton (*tri-koff'-it-on*) [*θριξ*, hair; *φυτόν*, a plant]. A fungus parasitic upon the hair, and causing tinea trichophytina, or ringworm.

Trichorrhexis (*trik-or-eks'-is*) [*θριξ*, hair; *ρῆξις*, a breaking]. Brittleness of the hair.

T. nodosa, an atrophic condition of the hair, affecting more often the male beard, and characterized by irregular thickenings resembling nodes on the hair-shaft, the hairs often breaking with a "green-stick fracture" immediately through a node.

Trichosis (*trik-o'-sis*). See *Trichonosis*.

Trichotillomania (*trik-o-til-o-ma'-ne-ah*) [*θριξ*, hair; *τίλλειν*, to pluck out; *μανία*, mania]. An uncontrollable impulse to pull out one's hair.

Trichroic (*tri-kro'-ik*) [*τριεις*, three; *χρῶμα*, color]. Possessing trichroism.

Trichroism (*tri'-kro-izm*) [*τριεις*, three; *χρῶμα*, color]. The property of exhibiting three different colors when viewed under three different aspects.

Tricipital (*tri-sip'-it-al*) [*triceps*, three-headed]. 1. Three-headed. 2. To the triceps.

Tricornis (*tri-kor'-nis*) [*θ*, horn]. Having three horns.

Tricrotic (*tri-krot'-ik*) [*τρεις*, three; *κρότος*, stroke]. Having three waves corresponding to one pulse-beat.

Tricrotism (*tri'-krot-izm*) [*τρεις*, three; *κρότος*, stroke]. The quality of being tricrotic.

Tricuspid (*tri-kus'-pid*) [*tres*, three; *cuspis*, point]. 1. Having three cusps, as, *e. g.*, the T. valve. 2. Affecting or produced at the T. valve.

Triethylamin (*tri-eth-il-am'-in*) [*τρεις*, three; *ethyl*; *amin*], $C_6H_{15}N$. A ptomain obtained from putrid haddock. See *Ptomains*, Table of.

Trifacial Nerve (*tri-fa'-shal*) [*tres*, three; *facies*, face]. The fifth cranial nerve, so-called because it divides into three main branches that supply the face.

Trifolium (*tri-fol'-le-um*) [*tres*, three; *folium*, leaf]. Clover.

Trigeminal (*tri-jem'-in-al*) [*tres*, three; *geminus*, twin-born]. 1. Triple; dividing into three parts, as, *e. g.*, the T. nerve. 2. Pertaining to the T. nerve. See *Trifacial*.

Trigeminus (*tri-jem'-in-us*) [*tres*, three; *gemini*, double]. The trifacial nerve.

Trigger-finger. A condition in which flexion or extension of a finger is at first impeded, but finally accomplished with a jerk.

Trigone (*tri'-gon*) [*τρεις*, three; *γωνία*, angle]. Triangle. T. of the bladder,

Trigonum vesicæ, or simply T., a smooth, triangular space on the inside of the bladder, immediately behind the orifice of the urethra.

T., Olfactory, the gray root of origin of the olfactory tract.

Trihydrate (*tri-hi'-drat*) [*τρεις*, three; *hydrate*]. A compound containing the hydroxyl radicle in the proportion of three to one atom of the base.

Trihydric (*tri-hi'-driks*) [*τρεις*, three; *hydric*]. Containing three atoms of hydrogen replaceable by bases.

Trihydroxid (*tri-hi-drok'-id*). See *Trihydrate*.

Triiodid (*tri-i'-o-did*) [*τρεις*, three; *iodid*]. A compound containing iodine in the proportion of three atoms to one of the base.

Trimethylamin (*tri-meth-il-am'-in*) [*tres*, three; *methyl*; *amin*]. A colorless liquid ptomain obtained from herring-brine and various animal and vegetable substances. See *Ptomains*, Table of.

Trimethylenediamin (*tri-meth-il-en-di-am'-in*) [*tres*, three; *methylene*; *diamin*]. A ptomain obtained from cultures of the comma bacillus on beef broth. It causes convulsions and muscle-tremor. See *Ptomains*, Table of.

Trimethyl-xanthin (*tri-meth-il-zan'-thin*). See *Caffein*.

Trinitrate (*tri-ni'-tāt*) [*tres*, three; *nitrate*]. A nitrate containing three nitric-acid radicles.

Trinitrin (*tri-ni'-trin*). See *Nitroglycerin*.

Trinitrocellulose (*tri-ni-tro-sef'-u-lōs*). See *Pyroxylin*.

Trinitrophenol (*tri-ni-tro-fe'-nol*). See *Picric Acid*.

Triocephalus (*tri-o-sef'-al-us*) [*τρεις*, three; *κεφαλή*, head]. A monster characterized by an absence of the ocular, nasal, and buccal apparatus, the head being merely a small spheroidal mass.

Triolein (*tri-o'-le-in*) [*tres*, three; *oleum*, oil]. See *Olein*.

Trional (*tri'-on-al*) [*τρεις*, three]. $C_4H_5CH_2-C-(SO_2C_2H_5)_2$. Diethylsulphonemethylmethane, an hypnotic. Dose gr. xv (1.0).

Triorchid (*tri-or'-kid*) [*τρεις*, three; *ὄρχις*, a testicle]. 1. Having three testicles. 2. An individual having three testicles.

Trioxid (*tri-oks'-id*) [*τρεις*, three; *oxid*]. A compound containing oxygen in the proportion of three atoms to one of the base.

Tripalmitin (*tri-pal'-mit-in*) [*τρεις*, three; *palmitin*]. See *Palmitin*.

Tripier's Amputation. One differing from Chopart's only in that the portion of the os calcis below the sustentaculum tali is removed.

Triple (*trip'-l*) [*tripplus*, triple]. Three-fold. T. Phosphate, ammoniomagnesian phosphate, a phosphate occurring in urine and in phosphatic calculi.

Triplet (*trip'-let*) [*tripplus*, triple]. 1. One of three children born at one birth. 2. In optics, a system consisting of three lenses.

Triplex (*trip'-lex*) [L.]. Triple. T. Pills, Pilulæ triplices, pills containing three principal ingredients.

Triploblastic (*trip-lo-blas'-tik*) [*τριπλόος*, threefold; *βλαστός*, germ]. Possessing three blastodermic membranes.

Triplopia (*trip-lo'-pe-ah*) [*τριπλόος*, triple; *ὤψ*, eye]. A disturbance of vision in which three images of a single object are seen.

Triquetrous (*tri-kwet'-rus*) [*triquetrus*, three-cornered]. Three-cornered, as, *e. g.*, the T. bone (os triquetrum), a Wormian bone. See *Triquetrum*.

Triquetrum (*tri-kwet'-rum*) [*triquetrum*, three-cornered]. 1. Any one of the Wormian bones. 2. The cuneiform bone of the carpus.

Triradial, Triradiate (*tri-ra'-de-al, tri-ra'-de-at*) [*tres*, three; *radius*, ray]. Radiating in three directions.

Trismus (*tri'-mus*) [*τρισμός*, from *τρι*, to gnash]. Lockjaw, a tonic spasm of the muscles of mastication. T. nascentium, T. neonatorum, a form of T. occurring in newborn infants, and supposed to be due to septic infection of the umbilical stump.

Trisplanchnic (*tri-splan'-gik'-nik*) [*τριπλάνχνης*, three; *σπλάνχνην*, viscus]. Distributed to the viscera of the three largest cavities of the

body, as the T. nerve (the sympathetic nerve).

Tristearin (*tri-ste'-ar-in*) [*τρεῖς*, three; *στέαρ*, fat]. $C_5H_5(C_{18}H_{35}O_2)_3$. See *Stearin*.

Trisubstituted (*tri-sub'-stit-u-ted*) [*tres*, three; *substituere*, to substitute]. Having three atoms or radicles substituted by other atoms or radicles.

Trisulphid (*tri-sul'-fid*) [*tres*, three; *sulphur*]. A compound containing sulphur in the proportion of three atoms to one of the base.

Triticeoglossus (*trit-is-e-o-glos'-us*) [*tritium*, wheat; *γλῶσσα*, tongue]. An anomalous muscle having its origin from the arytenoid cartilage and its insertion in the side of the tongue.

Triticeous (*trit-ish'-us*) [*tritium*, wheat]. Having the shape of a grain of wheat. **T. Nodule**, corpus triticeum, a small cartilaginous nodule in the thyrohyoid ligament.

Triticum (*trit'-ik-um*) [L.]. A genus of the Gramineæ. T. of the U. S. P. is the rhizome of *Agropyrum repens*, and is used in cystitis and irritable bladder. Dose of Extractum tritici fluidum, U. S. P., f3 iij-vj (12 o-24 o). T. sativum, T. vulgare, is wheat.

Tritoxid (*tri-toks'-id*) [*τρίτος*, third; *ὀξύς*, acid]. Same as *Trioxid*.

Triturate (*trit'-u-rat*) [*triturare*, from *terere*, to rub]. 1. To reduce to fine powder. 2. A finely divided powder. In the U. S. P., a medicinal substance rubbed up with milk-sugar. **T.**, **Tablet-**, a T. compressed into tablet-form.

Trituration (*trit-u-ra'-shun*) [*triturare*, from *terere*, to rub]. The process of reducing a solid substance to a powder by rubbing.

Trivalent (*triv'-al-ent*) [*tres*, three; *valere*, to be worth]. Combining with or equivalent to three atoms of hydrogen.

Trivalve (*tri'-valu*) [*tres*, three; *valva*, door]. Having three valves.

Trocar (*tro'-kar*) [Fr., *trois-quarts*, from its triangular point]. An instrument for piercing the wall of a cavity in paracentesis.

Trochanter (*tro-kun'-ter*) [*τροχαντήρ*, from *τροχός*, a wheel or pulley]. One of two processes on the upper extremity of the femur below the neck. The greater T. is situated on the outer, and the lesser T. on the inner side of the bone.

Trochanteric (*tro-kun-ter'-ik*) [*τροχαντήρ*, a trochanter]. Pertaining to a trochanter.

Troche (*tro'-ke*) [*τροχός*, wheel]. A lozenge.

Trochiscus (*tro-kis'-kus*). See *Troche*.

Trochlea (*trok'-le-ah*) [*τροχιλία*, pulley]. A part having the nature of a pulley. T. of the Humerus, a surface acting as a pulley at the lower extremity of the humerus. T.

of the Superior Oblique, T. of the Orbit, the ligamentous ring or pulley, attached to the upper margin of the orbit, which transmits the tendon of the superior oblique muscle of the eye.

Trochlear (*trok'-le-ar*) [*τροχιλία*, a pulley].

1. Pertaining to or of the nature of a pulley. 2. Pertaining to the T. muscle. 3. Pertaining to the T. nerve.

Trochlearis (*trok-le-a'-ris*) [*τροχιλία*, a pulley]. Pulley-shaped, as, e. g., the T. muscle or simply T., the superior oblique muscle of the eye.

Trochoides (*tro-koi'-des*) [*τροχός*, wheel]. A pivot-joint or pulley-joint, such as the atlanto-axial joint.

Trommer's Test. A test for the presence of glucose, consisting in the reduction of an alkaline copper-solution. When the latter is boiled with the suspected fluid rendered alkaline by the addition of sodium hydroxid, the copper is precipitated in the form of a bright, orange-red, basic oxid.

Tropacocain (*tro-pa-ko'-ka-in*) [*atropin*; *cocain*]. An alkaloid obtained from a small-leaved coca-plant of Java.

Tropæolin, **Tropeolin** (*tro-pe'-o-lin*). One of a group of orange anilin dyes, so called from the resemblance of their colors to those of the flowers of *Tropæolum*, the garden-nasturtium. Its solutions are turned brown by free acids and are used as a test for such acids.

Tropein (*tro'-pe-in*). A salt of tropin and an organic acid.

Trophic (*trof'-ik*) [*τροφή*, nourishment]. Pertaining to nutrition. **T. Center**, a collection of ganglion-cells regulating the nutrition of a nerve and through that of the part supplied.

Trophoneurosis (*trof-o-nu-ro'-sis*) [*τροφή*, nourishment; *νεῦρον*, nerve]. Any disease of a part due to disturbance of the nerves or nerve-centers with which it is connected.

Trophoneurotic (*trof-o-nu-ro'-ik*) [*τροφή*, nourishment; *νεῦρον*, nerve]. Pertaining to or caused by a trophoneurosis.

Tropic Acid (*trof'-ik*) [from *atropin*], $C_9H_{10}O_3$. An acid produced by treating atropin with baryta-water, alkalies, or acids.

Tropidin (*trof'-id-in*) [*atropin*], $C_8H_{13}N$. A substance resulting from the decomposition of atropin in the presence of hydrochloric and glacial acetic acids; it is an oily fluid having an odor like coniin.

Tropin (*trof'-in*) [*atropin*], $C_8H_{15}NO$. A crystalline base obtained in the decomposition of atropin.

Tropometer (*trof-om'-et-er*) [*τροφή*, a twin; *μέτρον*, a measure]. An instrument for measuring the various rotations of the eyeball.

Trousseau's Phenomenon or **Sign** (*troo-*

sân) [A. Trousseau, a French physician]. Spasm of the muscles of an extremity produced by compression of its main artery or nerve; it is observed in some cases of tetany. **Trousseau's Spots.** See *Tache cérébrale*.

Troy Ounce. A unit in Troy weight, equal to 480 grains.

Troy Weight. See *Weights and Measures*.

True. Real; not false. **T. Aneurysm.** See *Aneurysm*. **T. Corpus luteum,** the corpus luteum of pregnancy. **T. Pelvis,** that part of the pelvic cavity situated below the iliopectineal line. **T. Ribs.** See *Ribs*. **T. Skin,** the corium. **T. Vocal Bands,** the inferior bands, or those concerned in the production of the voice.

Truncal (*trung'-kal*) [*truncus*, a trunk]. Pertaining to a trunk.

Trunk [*truncus*]. 1. The body except the head and limbs. 2. The main stem of a nerve or artery.

Truss (*trus*) [*Fr., trousse*]. An apparatus for maintaining a hernia in place after reduction.

Trypsin (*trip'-sin*) [*τριψιν*, a rubbing]. The proteolytic ferment of the pancreatic juice, which in an alkaline medium converts proteids into peptones.

Trypsinogen (*trip'-sin'-o-jen*) [*τριψιν*, a rubbing; *γεννᾶν*, to produce]. The zymogen from which trypsin is formed.

Tryptic (*trip'-tik*) [*τριψιν*, a rubbing]. Pertaining to or caused by trypsin.

Tryptone (*trip'-ton*) [*τριψιν*, a rubbing]. Peptone formed by the action of trypsin.

Tsuga (*tsu'-gah*). A genus of Coniferae, a species of which, *T. canadensis*, yields Canadian pitch.

Tubal (*tu' bal*) [*tuba*, a tube]. Pertaining to a tube, especially the Fallopian tube or the renal tubules.

Tube (*tub*) [*tuba*]. A hollow cylindric structure, especially the Fallopian T. or the Eustachian T. **T.-casts,** casts of the renal tubules; they indicate disease of the kidneys.

T., Drainage-, a hollow tube of glass, rubber, or other material inserted into a wound or cavity to allow of the escape of fluids.

T., Crooke's, T., Geissler's, T., Hittorf's. See *Alarvs*.

Tuber (*tu' ber*) [*tuber*, a bump or swelling].

1. A thickened portion of an underground stem. 2. Any rounded swelling. **T. cinereum,** a tract of gray matter extending from the optic chiasm to the corpora mamillaria, and forming part of the floor of the third ventricle.

Tubercle (*tu' ber-kl'*) [*tuberculum*, a tubercle]. 1. A small nodule. 2. A rounded prominence on a bone. 3. The specific lesion produced by the tubercle bacillus, consisting of a collection of round cells and epithelioid cells, with at times giant-cells. **T.,**

Adductor-, a slight protuberance at the end of the internal supracondylar line of femur, giving attachment to the tendon of adductor magnus. **T., Amygdaloid** prominence on the roof of the descending cornu of the lateral ventricle. **T., Anato-** a wart-like tuberculous growth sometimes appearing on the hands of dissectors.

Anterior, a tubercle at the anterior part of the extremity of the transverse process of a vertebra. **T.-bacillus.** See *Bacilli*.

T., Carotid, a prominence of the sixth cervical vertebra on the anterior part of its transverse process. **T., Conoid,** a bony projection of the clavicle on its posterior border at the union of its middle and outer thirds, to which the conoid ligament is attached. **T., Deltoid,** a projection on the anterior border of the clavicle, giving origin to a part of the deltoideus. **T., Fibrous** tubercle (3d def.) which has been modified by the formation of connective tissue within its structure. **T., Genial,** one of the tubercles on each side of the middle line of the inner surface of the lower maxilla.

Genital, the rudimentary penis or clitoris of the urogenital region of the embryo in front of the cloaca. **T., Gray.** See *T., Miliary*.

of Lower, a small eminence on the wall of the right auricle, between the orifices of the vena cavae. **T., Lymphoid,** a T. (3d def.) consisting chiefly of round or lymphoid cells. **T., Mamillary.** See *Corpora albican*.

T., Miliary. See *T., 3d def.* **T., Olfactory.** See *Bulb, Olfactory*. **T., Painful** nodule in the subcutaneous tissue of the region of the joints. **T., Posterior,** a tubercle at the posterior end of the lumbar and seventh of the thoracic vertebrae. **T., Pterygoid,** a T. on the inner surface of the inferior maxilla; it gives attachment to the interpterygoid muscle. **T. of Rolando,** one of the rounded masses close under the surface of the lateral columns of the medulla oblongata, formed by the enlarged dorsal horns of the gray matter. **T., Scalene-**, a T. on the first rib, giving attachment to the anterior scalene muscle.

Tubercular (*tu' bur'-ku-lar*) [*tubercular*, tubercle]. Characterized by the formation of tubercles.

Tuberculin (*tu' bur'-ku-lin*) [*tuberculum*, tubercle]. A glycerol extract of cultures of the bacillus of tuberculosis. It is a brown neutral liquid, soluble in water, and is used as a means of diagnosing tuberculosis, especially in the domestic animals; when injected into tuberculous individuals a reaction is produced which differs from that given by healthy individuals. It is also known as *Robt. Koch's T.*

din (*tu' bur'-ku-le si'-din*) [*tuberculin*, tubercle].

culum, tubercle; *cadere*, to kill]. An albumose obtained from tuberculin by precipitation with platinum chlorid. It is said to possess the beneficial effects of tuberculin without producing an injurious reaction.

Tuberculosis (*tu-bur-ku-lo'-sis*) [*tuberculum*, a tubercle]. An infectious disease due to the bacillus tuberculosis, discovered by Koch. The lesion produced by the growth of the bacillus is the tubercle (miliary or gray tubercle or nodule), a small, grayish, translucent nodule, from $\frac{1}{16}$ to 2 mm. in diameter, firmly imbedded in the surrounding tissues. By the coalescence of neighboring tubercles larger masses, the so-called tuberculous infiltrations, are produced. The tendency of tuberculous lesions is to undergo cheesy necrosis. For this degeneration two factors are responsible: the absence of blood-vessels and the action of peculiar poisons elaborated by the bacillus. The breaking down of tuberculous areas in the interior of organs gives rise to cavities, which may be seen in muscles, bones, brain, lymphatic glands, and elsewhere, but are most pronounced in the lungs. On surfaces—skin and mucous membranes—tuberculosis often leads to the formation of ulcers. The most frequent seats of tuberculosis are the lung, the intestinal tract, the lymphatic glands, the serous membranes, the bones, the skin, the testicle, the epididymis, the brain, the Fallopian tubes, the uterus, the spleen. The symptoms of tuberculosis vary with the localization of the disease. A few general phenomena are common to nearly all forms, viz.: emaciation, loss of strength, anemia, fever, and sweats. **T.**, **Acute Miliary**, an acute febrile disease, characterized by the formation of minute tubercles in great numbers in various parts of the body. It is due to the discharge into the circulatory stream of tubercle-bacilli. Three forms are usually described: (1) a general or typhoid form; (2) one with marked pulmonary symptoms; (3) one in which cerebral symptoms predominate. **T.**, **Avian**, tuberculosis affecting birds. **T.**, **Bovine**, tuberculosis occurring in cattle; pearl-disease. **T.**, **General Miliary**. See **T.**, **Acute Miliary**. **T.**, **Laryngeal**, tuberculosis of the larynx, usually secondary to tuberculosis of the lungs, but in rare cases primary. **T.**, **Miliary**, tuberculosis characterized by the formation deposit of miliary tubercles. **T.**, **Surgical**, tuberculosis of parts amenable to surgical treatment, as the bones and joints.

Tuberculous (*tu-bur'-ku-lus*) [*tuberculum*, tubercle]. Affected with or caused by tuberculosis.

Tuberculum (*tu-bur'-ku-lum*) [L.]. See **Tubercle**.

Tuberose (*tu'-ber-os*) [*tuber*]. Resembling a tuber.

Tuberosity (*tu-ber-os'-it-e*) [*tuber*, a swelling]. A protuberance on a bone. **T.**, **Greater**, a rough projection on the outer side of the head of the humerus. **T.** of the Ischium, a thick, downward projection of the ischium, on which the body rests in sitting. **T.**, **Lesser**, a small tuberosity in front of the head and on the inner side of the bicipital groove of the humerus.

Tuberous (*tu'-ber-us*) [*tuberosus*]. Like a tuber, as, *e. g.*, **T.** angioma.

Tuboabdominal (*tu-bo-ab-dom'-in-al*) [*tuba*, a tube; *abdomen*, abdomen]. Pertaining to a Fallopian tube and to the abdomen.

Tuboovarian (*tu-bo-o-va'-re-an*) [*tuba*, a tube; *ovary*]. Pertaining to the Fallopian tube and the ovary.

Tubouterine [*tuba*, tube; *uterus*, uterus]. Pertaining to the Fallopian tube and the uterus.

Tubular (*tu'-bu-lar*) [*tubulus*, a small tube]. 1. Shaped like a tube. 2. Pertaining to or affecting tubules, as, *e. g.*, **T.** nephritis. 3. Produced in a tube, as, *e. g.*, **T.** breathing. **T.** **Membrane**. See **Neurilemma**.

Tubulature (*tu'-bu-lat-tar*) [*tube*, tube]. The short tube of a retort or receiver.

Tubule (*tu'-bal*) [*tubulus*, dim. of *tubus*, a tube]. A small tube. **T.**, **Seminiferous**, any one of the tubules of the testicles. **T.**, **Uriniferous**, one of the numerous winding tubules of the kidney.

Tubulus (*tu'-bu-lus*) [L.]. A tubule.

Tufter's Treatment. A treatment for aneurysm, consisting in absolute rest, a dry diet, and the administration of potassium iodid.

Tuft, Malpighian. See **Malpighian Bodies**.

Tugging, Tracheal. See **Tracheal Tugging**.

Tumefaction (*tu-me-fak'-shun*) [*tumere*, to swell; *facere*, to make]. A swelling.

Tumor (*tu'-mor*) [*tumere*, to swell]. 1. A swelling. 2. A new growth not the result of inflammation. The classification given on next page is based, at least as regards classes A and B, on the blastodermic origin of the dominant tissue of the tumor. **T.** **albus**, white swelling; tuberculous enlargement of a joint. **T.**, **Benign**, one which does not give rise to metastasis or recur after removal. **T.**, **Gubler's**, a prominence on the back of the wrist seen in wrist-drop. **T.**, **Gummy**, a syphilitic gumma. **T.**, **Heterologous**, one composed of tissue differing from that in which it grows. **T.**, **Homologous**, one composed of tissue resembling that from which it grows. **T.**, **Malignant**, one which gives metastasis or recurs, or does both, and eventually destroys life. **T.**, **Phantom**. See **Phantom-tumor**. **T.**, **Splenic**, a term sometimes applied to an enlarged spleen.

A. MESODERMIC TUMORS.

- | | | | |
|------------------|---|-----------------------------------|---|
| 1. Sarcoma. | { | Round-cell. | { Large.
Small.
Lympho-
sarcoma. |
| | | Spindle-cell. | { Large.
Small. |
| | | Giant-cell. | |
| | | Melanotic. | |
| | | Alveolar. | |
| | | Endothelioma. | |
| | | Angiosarcoma. | |
| | | Cylindroma. | |
| | | Chloroma. | |
| | | Psammoma. | |
| 2. Fibroma. | { | Hard. | |
| | | Soft. | |
| 3. Myxoma. | | | |
| 4. Lipoma. | | | |
| 5. Chondroma. | { | Hyaline. | |
| | | Fibrous. | |
| | | O. durum or O. eburneum. | |
| 6. Osteoma. | { | O. spongiosum; O. medul-
lary. | |
| | | Telangiectatic. | |
| 7. Hemangioma. | | Cavernous. | |
| | | Simple. | |
| 8. Lymphangioma. | | Cystic. | |
| | | Liomyoma. | |
| 9. Myoma. | | Rhabdomyoma. | |

B. ECTODERMIC AND ENTODERMIC TUMORS.

- | | | | |
|-----------------|---|------------------------|----------------------------|
| 1. Glioma. | | | |
| 2. Neuroma. | { | N. myelinicum. | |
| | | N. amyelinicum. | |
| | | Adenoma. | { Tubular.
Racemose. |
| | | | { Squamous. |
| 3. Epithelioma. | | Carcinoma. | { Cylindric.
Glandular. |
| | | | { C. simplex. |
| | | Epithelial
cystoma. | { C. papilli-
ferum. |

C. TERATOID TUMORS OR TERATOMATA.

1. Dermoid cyst.
2. Cholesteatoma.

Tungsten (*tung'-sten*) [Swed., heavy stone]. A metallic element having a specific gravity of 19.1, an atomic weight of 183.5. Symbol W (from the German name *Wolfram*). It forms **Tungstic Acid**, H_2WO_4 , the latter combining with bases to form **Tungstates**, which are used as reagents.

Tunic (*tu'-nik*) [*tunica*]. A coat or membrane.

Tunica (*tu'-nik-ah*) [L.]. A tunic. **T. adventitia**, the outer coat of an artery. **T. albuginea oculi**, the sclerotic coat of the eye. **T. albuginea ovarii**, the compact connective tissue immediately under the epithelium of the cortex of the ovary. **T. albuginea testis**, the fibrous covering of the testis. **T. intima**, the inner coat of an artery. **T. media**, the middle coat of an artery. **T. ruyschiana**, the layer of capillary vessels of the choroid coat of the eye. **T. vaginalis**, the serous covering of the testis derived from the peritoneum. **T. vasculosa**, the vascular layer of the testis, called also the pia mater of the testis.

Tunnel-anemia. See *Ankylostomiasis*.

Tunnel-disease. See *Caisson-disease*.

Tupelo (*tu'-pel-o*). The *Nyssa grandidentata*, of the order Cornaceæ. Its root has been used for making tents (**T.-tent**).

Turbinal (*tur'-bin-al*) [*turbo*, top]. 1. Turbinated. 2. A turbinated bone.

Turbinated (*tur'-bin-a-ted*) [*turbo*, a top]. Top-shaped; scroll-shaped. **T. Bone**, one of the three (superior, middle, and inferior) bony projections upon the outer wall of each nasal fossa. They are covered by an erectile vascular mucous membrane.

Türk's Column. See *Column*.

Turgescence (*tur-jes'-ens*) [*turgescere*, to become swelled]. Swelling.

Turgid (*tur'-jid*) [*turgidus*, swollen]. Swollen, congested.

Turmeric (*tur'-mer-ik*) [*turmerica*, turmeric]. The *Curcuma longa*, a plant of the Zingiberaceæ. Its rhizome contains a yellow coloring matter, curcumin, $C_{14}H_{14}O_4$. Its solution in alcohol is turned reddish-brown by alkalis, and has been used as a test for the latter. Paper tinged with an alcoholic solution (**T.-paper**) may be used instead.

Turning (*turn'-ing*). See *Version*.

Turpentine (*tur'-pen-tin*) [*τερεβινθος*, terebinth]. A concrete or liquid oleoresin obtained from various species of Coniferae. The ordinary or white **T.** (Terebinthina, U. S. P., Thus americanum, B. P.), derived from *Pinus palustris* and other species of *Pinus*, contains a volatile oil, oil or spirits of **T.** (Oleum terebinthinæ, U. S. P., B. P.). This when pure consists only of carbon and hydrogen, but on exposure absorbs oxygen. Oil of **T.** is stimulant, diuretic, and anthelmintic; in large doses it acts as a cathartic; locally it is a rubefacient. In overdoses it acts as an irritant, especially to the kidneys, producing bloody urine and strangury. It is used as a stimulant in typhoid and other low fevers; in tympanites; as a hemostatic; in chronic renal diseases, dysentery, and whooping-cough; as an inhalation in bronchitis; as a cathartic in the form of enema, and as a teniafuge. Dose $\mathfrak{m}\mathfrak{v}$ -xxx (0.32-2.0). Preparations: Confectio terebinthinæ (B. P.). Dose \mathfrak{ss} -j (2.0-4.0). Enema terebinthinæ (B. P.); Linimentum terebinthinæ (U. S. P., B. P.); Linimentum terebinthinæ aceticum (B. P.); Oleum terebinthinæ rectificatum (U. S. P.). Dose $\mathfrak{m}\mathfrak{v}$ -xxx (0.32-2.0). Unguentum terebinthinæ (B. P.). **T., Canada** (Terebinthina canadensis, U. S. P., B. P.), is obtained from *Pinus balsamea*, and under the name of Canada balsam is used as a mounting medium in microscopy. **T., Chian**, collected on the island of Chios, from the *Pistacia terebinthus*, was formerly used in various diseases. **T., Common European, T.,**

Bordeux, is obtained from several species of pine, chiefly *Pinus sylvestris* and *Pinus maritima*. It yields large quantities of oil of T. T., **Strasburg**, is derived from *Abies picea*. T., **Venice**, is obtained from *Larix europæa*, and yields oil of T.

Turpeth (*tur'-peth*) [Pers., *turbad*, a purgative root]. The *Ipomœa turpethum*, a purgative plant resembling jalap, found in Asia. T.-mineral, subsulphate of mercury, used as an emetic. See *Mercury*.

Tussilago (*tus-il-a'-go*). A genus of plants of the order Compositæ. The leaves of T. farfara, coltsfoot, and also other parts of the plant are used as a demulcent in pulmonary affections associated with cough.

Tussis (*tus'-is*) [L.]. A cough. T. convulsiva, whooping-cough.

Tussive (*tus'-iv*) [*tussis*, cough]. Pertaining to or caused by cough.

Tutamina oculi (*tu-tam'-in-ah ok'-u-li*). The appendages of the eyes.

Twelfth Cranial Nerve. The hypoglossal nerve. See *Nerves*, *Table of*.

Twin. One of two individuals born at the same birth.

Tyloma (*ti-lo'-mah*) [τύλος, callus]. A callus.

Tylosis (*ti-lo'-sis*) [τύλος, callus]. A state characterized by the formation of callus.

Tympanic (*tim-pan'-ik*) [τύμπανον, a drum]. Pertaining to the tympanum. T. Bone, T. Plate, the thin plate of bone separating the tympanum from the cranial cavity. T. Membrane. See *Membrane*. T. Tegmen, the bony plate forming the roof of the tympanum.

Tympanites (*tim-pan-i'-tē*) [τύμπανον, drum]. A distention of the abdominal walls from accumulation of gas in the intestines or peritoneal cavity.

Tympanitic (*tim-pan-i'-ik*) [τύμπανον, a drum]. Caused by or of the nature of tympanites. T. Resonante, the note obtained on percussing a cavity distended with gas.

Tympanitis (*tim-pan-i'-tis*) [τύμπανον, drum; ιτις, inflammation]. Inflammation of the tympanum; otitis media.

Tympanoeustachian (*tim-pan-o-u-sta'-ke-an*) [τύμπανον, drum; Eustachius, an anatomist]. Pertaining to the tympanum and the Eustachian tube.

Tympanohyal (*tim-pan-o-hi'-al*) [τύμπανον, drum; ὑαλῆς, hyoid]. A small cartilage of the human fetus subsequently fusing with the styloid process of the temporal bone.

Tympanum (*tim'-pan-um*) [τύμπανον, drum]. The middle ear.

Typhlitis (*tif-li'-tis*) [τυφλός, cecum; ιτις, inflammation]. Inflammation of the cecum.

Typhoid (*ti'-foid*) [τύφος, stupor; εἶδος,

like]. Resembling typhus. T. Fever, enteric fever, abdominal typhus, ileotyphus. An infectious disease caused by the bacillus typhosus discovered by Eberth. It is introduced into the body with the food and drinking water, and is found in the intestines, the spleen, and the fecal discharges, but may also occur in the various complicating lesions. The principal lesions of typhoid fever are an enlargement and necrosis of Peyer's patches, and enlargement of the spleen and the mesenteric glands. The mucous membrane of the intestines is also the seat of a catarrhal inflammation. After a period of incubation of from two to three weeks the disease sets in with weakness, headache, vague pains, a tendency to diarrhoea, and nose-bleed. The temperature gradually rises, being higher each evening than the previous evening, and reaches its maximum (104°-105° F.) in from one to two weeks. It then remains at this level for from one to two weeks, and finally sinks by lysis. The pulse is soft and dicrotic, but often not as rapid as would be expected from the high temperature. The tongue is at first coated on the dorsum and red at the tip and edges, but soon becomes dry, brown, and tremulous, and, like the teeth and lips, covered with sordes. There is usually complete anorexia, the bowels are loose, the stools having a peculiar "pea-soup" color. At times constipation exists. Slight congestion of the lungs with cough is usually present. On the seventh, eighth, or ninth day the peculiar eruption appears—it consists of small, slightly-elevated, rose-colored spots, disappearing on pressure, and coming out in successive crops. Nervous symptoms are prominent in typhoid fever, and are headache, slight deafness, stupor, muttering delirium, carphologia, subsultus tendinum, and coma vigil. Complications are frequent, the most important being intestinal hemorrhage, perforation of the bowel, peritonitis, pneumonia, and nephritis. Relapses are fairly common, although second attacks are rare. Abortive typhoid is characterized by an abrupt onset of symptoms, which subside quickly, convalescence following in a few days. Walking typhoid is a variety in which the symptoms are so mild that the patient disregards them and walks about. Grave accidents may in such a case develop suddenly. T. State, the condition of stupor and hebetude, with dry, brown tongue, sordes on the teeth, rapid, feeble pulse, incontinence of feces and urine, and rapid wasting, seen in typhoid fever and other continued fevers.

Typhomalarial (*ti'-fo-mal-a'-re-al*) [τύφος, stupor; malaria]. Exhibiting symptoms of both typhoid and malarial fevers. T. Fever, a fever having symptoms both of typhoid and

of malarial fever, but probably malarial in nature.

Typhomania (*ti-fo-ma'-ne-ah*) [*τύφος*, stupor; *μανία*, madness]. The lethargic state, with delirium, sometimes observed in typhus, typhoid, and other low fevers.

Typhotoxin (*ti-fo-toks'-in*) [*τύφος*, stupor; *τοξικόν*, poison]. A ptomain produced by the typhoid bacillus. It is isomeric with the base $C_7H_{11}NO_3$, obtained from putrefying horseflesh, and induces lethargy, paralysis, and death.

Typhous (*ti'-fus*) [*τύφος*, stupor]. Pertaining to or having the nature of typhus.

Typhus (*ti'-fus*) [*τύφος*, stupor]. Jail-fever, ship-fever. An acute contagious disease chiefly characterized by a petechial rash, marked nervous symptoms, and a high fever, ending by crisis in from ten to fourteen days. The only peculiar lesion noted postmortem is a dark fluid state of the blood, and a staining of the endocardium and intima of the blood-vessels. After a period of incubation of from a few hours to two weeks, the disease sets in abruptly with pains in the head, back, and limbs, the fever rising rapidly to 104° or 105° F. The nervous symptoms resemble those of typhoid fever. The eruption appears on the fourth or fifth day as rose-col-

ored spots scattered all over the body, and quickly becoming hemorrhagic. It does not disappear on pressure. The chief complications are hyperpyrexia, pneumonia, and nephritis.

Tyreine (*ti'-re-in*) [*τυρός*, cheese]. Coagulated casein.

Tyroid (*ti'-roid*) [*τυρός*, cheese; *ειδός*, like]. Cheese-like.

Tyroma (*ti-ro'-mah*) [*τυρός*, cheese]. 1. A caseous mass. 2. A tuberculous tumor.

Tyrosine (*ti'-ro-sin*) [*τυρός*, cheese], $C_9H_{11}NO_3$. A crystalline amidoacid, a decomposition-product of proteids.

Tyrosis (*ti-ro'-sis*) [*τυρός*, cheese]. Caseation.

Tyrotaxon (*ti-ro-toks'-ik-on*) [*τυρός*, cheese; *τοξικόν*, poison]. A ptomain obtained from poisonous cheese, milk, ice-cream, etc. It induces vertigo, nausea, vomiting, chills, rigors, severe pains in the epigastric region, dilatation of the pupils, griping and purging, a sensation of numbness or pins and needles, especially in the limbs, and marked prostration, or even death. The poison is thought to be the cause of the summer-diarrhea of infants. See *Ptomaines*, Table of.

Tyson, Glands of. The sebaceous glands secreting the smegma of the prepuce and of the labia pudendi.

U

Uffelmann's Test. A test for the presence of lactic acid in the stomach, performed by adding a little of the filtered gastric contents to a mixture of ferric chlorid and dilute phenol solution. If lactic acid is present the purple color is changed to yellow.

Ulcer (*ul'-ser*) [*ulcus*]. A loss of substance occurring on the skin or mucous membrane, and due to a gradual necrosis of the tissues.

U., **Atheromatous**. A loss of substance in the wall of an artery or the endocardium, due to the breaking down of an atheromatous patch. **U.**, **Chancroidal**. See *Chaneroid*.

U., **Curling's**, an ulcer of the duodenum observed after severe burns of the body. **U.**, **Follicular**, a small ulcer on a mucous membrane having its origin in a lymph-follicle.

U., **Fungous**, one covered by fungous granulations. **U.**, **Indolent**, one with an indurated, elevated edge, and a nongranulating floor, usually occurring on the leg. **U.**, **Inflamed**, one surrounded by marked inflammation. **U.**, **Jacob's**. See *U.*, *Rodent*. **U.**, **Marjolin's**, an ulcer having for its seat an

old cicatrix. **U.**, **Perforating**, an ulcer that perforates the tissues of a part, particularly the foot or the stomach. **U.**, **Phagedenic**, one which rapidly eats away the tissues. **U.**, **Phlegmonous**. Synonym of *U.*, *Inflamed*. **U.**, **Rodent**, a form of ulcer, probably epitheliomatous, which gradually involves and eats away soft tissues and bones. **U.**, **Round**, the peptic ulcer of the stomach. **U.**, **Serpiginous**, one healing in one place while spreading in another. **U.**, **Tuberculous**, one due to the tubercle-bacillus. **U.**, **Varicose**, an ulcer due to varicose veins.

Ulcerate (*ul'-ser-at*) [*ulcus*, ulcer]. To become converted into or affected with an ulcer.

Ulceration (*ul'-ser-a'-shun*) [*ulcus*, a sore]. The formation of an ulcer; a process of liquefaction-necrosis or molecular death on a free surface.

Ulcerative (*ul'-ser-a-tiv*) [*ulcus*, ulcer]. Pertaining to ulceration; characterized by ulceration.

Ulcus (*ul'-kus*) [*U.*]. See *Ulcer*.

Ulitis (*u-lit'-tis*) [*ούλον*, gum; *ετις*, inflammation]. Inflammation of the gums.

Ulmus (*ul'-mus*) [L.]. The *Ulmus fulva*, or elm, of the order *Urticaceæ*. Its bark (*Ulmus*, U. S. P., elm, slippery elm) is used as a demulcent in diarrhea, dysentery, and diseases of the urinary tract; as a poultice in inflammations, and in the form of tents for dilating the os uteri. Preparation: *Mucilago ulmi*, U. S. P.

Ulna (*ul'-nah*) [L., a cubit]. The bone on the inner side of the forearm, articulating with the humerus and the head of the radius above and with the radius below.

Ulnar (*ul'-nar*) [*ulnaris*, ulnar]. 1. Pertaining to or in relation with the ulna, as, *e. g.*, the U. artery, U. nerve. 2. Pertaining to the U. artery or U. nerve.

Ulnaris (*ul'-na'-ris*) [L.]. 1. Ulnar. 2. The U. muscle, a muscle on the ulnar side of the forearm.

Ulnocarpal (*ul-no-kar'-pal*) [*ulna*, ulna; *καρπός*, carpus]. Pertaining to the ulna and the carpus.

Ulnoradial (*ul-no-ra'-de-al*) [*ulna*, ulna; *radius*, radius]. Pertaining to the ulna and the radius.

Uloid (*u'-loid*) [*ούλή*, scar; *είδος*, like]. Scar-like.

Ulorrhagia (*ul-or-a'-je-ah*) [*ούλον*, gum; *ῥηγνύναι*, to burst forth]. Bleeding from the gums.

Ultimate (*ul'-tim-at*) [*ultimus*, superl. of *ultra*, beyond]. Farthest; most remote; last; final. U. Principle. See *Principle*.

Ultimum (*ul'-tim-um*) [L.]. Last. U. moriens (last dying). 1. The right auricle; so called from the belief that it is the last part of the heart to cease its contractions. 2. The upper part of the trapezius muscle which usually escapes in progressive muscular atrophy.

Umbilical (*um-bil'-ik-al*) [*umbilicus*, the navel]. 1. Pertaining to the umbilicus, as, *e. g.*, the U. cord, U. vessels. 2. Pertaining to the U. cord or U. vessels. U. Region, the central of the regions into which the abdomen is divided for purposes of physical diagnosis. U. Ring, the aperture, closed in the adult, through which the umbilical vessels pass in fetal life. U. Souffle, the peculiar sound heard occasionally over the umbilical cord of the fetus. U. Vesicle, the part of the yolk-sac remaining outside of the embryo and supplying nutriment to it through the omphalomesenteric duct.

Umbilicated (*um-bil'-ik-a-ted*) [*umbilicus*, navel]. Having a depression like that of the navel.

Umbilication (*um-bil'-ik-a'-shun*) [*umbilicus*, navel]. 1. A depression like that of the navel. 2. The state of being umbilicated.

Umbilicus (*um-bil'-i'-kus*) [L.]. The navel, the round, depressed cicatrix in the median line of the abdomen, marking the site of the aperture which in fetal life gave passage to the umbilical vessels.

Umbo (*um'-bo*) [L.]. A boss, or bosselation; any central convex eminence, as, *e. g.*, the U. of the membrana tympani.

Unavoidable Hemorrhage. See *Hemorrhage*.

Uncia (*un'-se-ah*) [L.]. An ounce.

Unciform (*un'-se-form*) [*uncus*, a hook; *forma*, form]. Hook-shaped. U. Bone, a hook-shaped bone in the second row of the carpus. U. Process, a hook-shaped process on the ethmoid and other bones.

Uncinate (*un'-sin-at*) [*uncus*, hook]. Hooked. U. Convolution, U. Gyrus, the continuation of the hippocampal convolution, or gyrus fornicatus, ending in a hook-like process near the end of the temporal lobe.

Uunction (*unk'-shun*) [*unguere*, to anoint]. 1. The act of anointing. 2. An ointment.

Unctuous (*unk'-tu-us*) [*unguere*, to anoint]. Greasy, oily.

Uncus (*ung'-kus*) [L.]. 1. A hook. 2. The hook-like anterior extremity of the uncinate gyrus of the brain.

Undulation (*un-du-la'-shun*) [*undulatio*, from *unda*, a wave]. A wave-like motion.

Ungual (*ung'-gwal*) [*unguis*, a claw]. 1. Pertaining to a nail. 2. Resembling a nail in size, as the U. bone (the lacrimal bone).

Unguent (*un'-gwent*) [*unguentum*, ointment]. See *Ointment*.

Unguiculate (*ung'-gwik'-u-lat*) [*unguis*, nail]. Having nails or claws.

Unguis (*ung'-guis*) [L.]. 1. A nail. 2. The lacrimal bone.

Uni- (*u'-ne-*) [*unus*, one]. A prefix denoting one.

Uniaxial (*u-ne-aks'-e-al*) [*unus*, one; *axis*, axis]. Having but one axis.

Unicellular (*u-ne-sel'-u-lar*) [*unus*, one; *cellula*, dim. of *cella*, a cell]. Composed of but one cell.

Unicorn (*u'-ne-korn*) [*unus*, one; *cornu*, horn]. Having a single horn.

Unilateral (*u-ne-lat'-er-al*) [*unus*, one; *latus*, side]. Pertaining to or affecting but one side.

Unilocular (*u-ne-lok'-u-lar*) [*unus*, one; *loculus*, dim. of *locus*, a place]. Having but one loculus or cavity.

Uninuclear, Uninucleated (*u-ne-nu'-kle-ar, u-ne-nu'-kle-a-ted*) [*unus*, one; *nucleus*, nucleus]. Having but a single nucleus.

Unilocular (*u-ne-ok'-u-lar*) [*unus*, one; *oculus*, eye]. Pertaining to or performed with one eye.

Union (*ūn'-yun*) [*unio*, oneness]. See *Healing*.

Unipara (*u-nip'-ar-ah*) [*unus*, one; *parere*, to bear]. A woman who has borne but one child.

Uniparous (*u-nip'-ar-us*) [*unus*, one; *parere*, to produce]. Having borne but one child.

Unipolar (*u-ne-pole'-lar*) [*unus*, one; *polus*, a pole]. 1. Having but one pole or process. 2. Pertaining to one pole.

Unisexual (*u-ne-seks'-u-al*) [*unus*, one; *sexus*, sex]. Provided with the sexual organs of one sex only.

Unit (*u'-nit*) [*unus*, one]. 1. A single thing; a group considered as a whole, or as forming one of many similar groups composing a more complex body. **U.**, **Physiologic**, a term used by Herbert Spencer to express a unit between the chemic and the morphologic units in complexity, and of an aggregation of which units the body is composed, and which represents the character of the species. 2. A quantity with which others are compared. The most important units are: **The Electric Units**—*U. of Capacity*, the farad; *U. of Current*, the ampere; *U. of Electromotive Force*, the volt; *U. of Power*, the watt; *U. of Quantity*, the coulomb; *U. of Resistance*, the ohm. **U. of Force**, the dyne; **U. of Heat**, the calorie; **U. of Light**, the light of a standard candle, *i. e.*, a spermaceti-candle burning 120 grains per hour; **U. of Length**, **U. of Volume**, **U. of Weight**. See *Weights and Measures*. **U. of Work**, the erg. **U.**, **Serum-**, **U.**, **Immunizing**, according to Behring, one cubic centimeter of an antitoxic blood-serum, of which 0.1 c.c. protects a guinea-pig of 500 grams against 10 times the fatal dose of diphtheria-toxin.

Unitary (*u'-nit-a-re*) [*unus*, one]. 1. Pertaining to or having the qualities of a unit. 2. Pertaining to monsters having the organs of a single individual.

Univalent (*u-ni-val-ent*) [*unus*, one; *valere*, to be worth]. Having a valence of one; capable of replacing a single hydrogen-atom in combination.

Unorganized (*un-or'-gan-izd*) [*un*, not; *organon*, organ]. Without organs; not arranged in the form of an organ or organs.

Unpolarized (*un-pole'-lar-izd*) [*un*, not; *polus*, pole]. Not polarized.

Unstriated (*un-strit-a-ted*) [*un*, not; *stria*, stripe]. Not striated, as, *e. g.*, **U. muscle**.

Ununited (*un-u-nit-ted*) [*un*, not; *unus*, one]. Not united, as, *e. g.*, an **U. fracture**.

Unwell. 1. Ill; sick. 2. Menstruating.

Urachus (*u'-ra-kus*) [*ōipov*, urine; *ἔχω*, to hold]. The allantoic stalk connecting in the fetus the bladder with the allantois, in after-life represented by a fibrous cord passing from the apex of the bladder to the umbilicus.

Ural (*u'-ral*) [*ōipov*, urine]. Chloral-urethane, $\text{CCl}_3\text{CH} : \text{OH.NHCO}_2\text{C}_2\text{H}_5$. An hypnotic. Dose gr. x-xx (0.65-1.3).

Uramin (*u-ram'-in*). See *Guanidin*.

Uranalysis (*u-ran-al'-is-is*). See *Urinalysis*.

Uraniscoplasty (*u-ran-is'-ko-plas-te*) [*ōipov-iskos*, palate; *πλάσσειν*, to form]. A plastic operation for the repair of cleft palate.

Uraniscorrhaphy (*u-ran-is-kor'-a-fe*) [*ōipov-iskos*, palate; *ραφή*, suture]. Suture of a palatal cleft; staphylorrhaphy.

Uranium (*u-ra'-ne-um*) [*Uranus*, Uranus]. A heavy white metal. See *Elements, Table of*. Its phosphate and nitrate are used as tests for phosphoric acid. The salts are very poisonous.

Uranoplasty (*u'-ran-o-plas-te*). Same as *Uraniscoplasty*.

Urare, **Urari** (*u-rah'-re*). See *Curare*.

Urate (*u'-rat*) [*ōipov*, urine]. A salt of uric acid.

Uratic (*u-rat'-ik*) [*ōipov*, urine]. Pertaining to or characterized by urates. **U. Diathesis**, a condition in which there is a tendency to the deposition of urates in the joints and elsewhere; a tendency to gout.

Uratoma (*u-ra-to'-mah*) [*urate*; *δμα*, tumor]. A concretion composed of urates, and occurring chiefly about the joints; a tophus.

Urea (*u-re'-ah*) [*ōipov*, urine], $\text{CO}(\text{NH}_2)_2$. The chief nitrogenous constituent of urine, and principal end-product of tissue-metamorphosis; it occurs also in the blood, the lymph, and the liver.

Ureometer (*u-re-am'-e-ter*) [*urea*; *μετρον*, measure]. An apparatus for determining the amount of urea contained in a liquid.

Ureametry (*u-re-am'-et-re*) [*urea*; *μετρον*, measure]. The determination of the amount of urea in a liquid.

Uredema (*u-re-de'-mah*) [*ōipov*, urine; *οἰδημα*, a swelling]. Distention of tissues from extravasation of urine.

Uredo (*u-re'-do*) [*uredo*, a blight]. 1. A genus of fungi. 2. Urticaria.

Ureid (*u'-re-id*) [*urea*]. A compound of urea and an acid radicle.

Uremia (*u-re'-me-ah*) [*ōipov*, urine; *αἷμα*, blood]. The symptoms due to the retention in the blood of excrementitious substances normally excreted by the kidneys; it is characterized by headache, vertigo, vomiting, amaurosis, convulsions, coma, sometimes hemiplegia, and a urinous odor of the breath.

Uremic (*u-rem'-ik*, *u-re'-mik*) [*ōipov*, urine; *αἷμα*, blood]. Due to or characterized by uremia.

Ureometer (*u-re-am'-et-er*) [*urea*; *μετρον*, measure]. See *Ureameter*.

Ureometry (*u-re-am'-et-re*). See *Ureametry*.

Uresis (*u-re'-sis*) [*ōipov*]. Same as *Urinization*.

-uret (*-u'-ret*). A suffix denoting a binary compound of carbon, sulphur, etc., with another element; in modern chemistry the suffix *id* is used.

Ureter (*u-re'-ter*) [*οὐρητήρ*, ureter]. The long, narrow tube conveying the urine from the pelvis of the kidney to the bladder.

Ureteral (*u-re'-ter-al*) [*οὐρητήρ*, ureter]. Pertaining to the ureter.

Ureteralgia (*u-re-ter-al'-je-ah*) [*οὐρητήρ*, ureter; *ἄλγος*, pain]. Neuralgic pain in the ureter.

Ureteric (*u-re-ter'-ik*) [*οὐρητήρ*, ureter]. Pertaining to the ureter.

Ureteritis (*u-re-ter-i'-tis*) [*οὐρητήρ*, ureter; *τις*, inflammation]. Inflammation of a ureter.

Ureterolith (*u-re-ter-o-lith*) [*οὐρητήρ*, ureter; *λίθος*, stone]. Calculus in the ureter.

Ureterotomy (*u-re-ter-ot'-o-me*) [*οὐρητήρ*, ureter; *τομή*, a cutting]. Incision of the ureter.

Ureterouterine (*u-re-ter-o-u'-ter-in*) [*οὐρητήρ*, ureter; *uterus*, uterus]. Pertaining to the ureter and the uterus.

Ureterovaginal (*u-re-ter-o-vaj'-in-al*) [*οὐρητήρ*, ureter; *vagina*, vagina]. Pertaining to the ureter and the vagina.

Urethane (*u'-reth-an*) [*οὐρον*, urine; *ether*]. 1. $\text{NH}_4\text{CO}_2\text{C}_2\text{H}_5$, ethyl carbamate. An hypnotic. Dose gr. xx-lx (1.3-2.6). 2. In a wider sense, any ester of carbamic acid is called a urethane.

Urethra (*u-re'-thrah*) [*οὐρήθρα*, urethra]. The canal through which the urine is discharged, extending from the neck of the bladder to the meatus urinarius. It is divided in the man into the *prostatic portion*, the *membranous portion*, and the *spongy or penile portion*, and is from 8 to 9 inches long. In the woman it is about 1½ inches in length.

Urethral (*u-re'-thral*) [*οὐρήθρα*, urethra]. Pertaining to the urethra; produced in or arising from the urethra, as, *e.g.*, U. fever.

U. Arthritis, gonorrheal rheumatism.

Urethralgia (*u-re-thral'-je-ah*) [*οὐρήθρα*, urethra; *ἄλγος*, pain]. Neuralgic pain in the urethra.

Urethritis (*u-re-thri'-tis*) [*οὐρήθρα*, urethra; *τις*, inflammation]. Inflammation of the urethra. **U.**, Anterior, inflammation of the part situated anterior to the anterior layer of the triangular ligament. **U.**, Posterior, inflammation of the prostatic and membranous portions. **U.**, Simple, a nonspecific inflammation of the urethra. **U.**, Specific, that due to the gonococcus; gonorrhea.

Urethro- (*u-re'-thro-*) [*οὐρήθρα*, urethra]. A prefix denoting pertaining to the urethra.

Urethrocele (*u-re'-thro-sel*) [*οὐρήθρα*, urethra; *κήλη*, a tumor]. A protrusion of the wall of the female urethra.

Urethrometer (*u-re-throm'-et-er*) [*οὐρήθρα*, urethra; *μέτρον*, measure]. An instrument for determining the caliber of the urethra or for measuring the lumen of a stricture.

Urethroplasty (*u-re'-thro-plas-te*) [*οὐρήθρα*, urethra; *πλάσσειν*, to form]. Plastic operation upon the urethra.

Urethrorrhagia (*u-re-thror-al'-je-ah*) [*οὐρήθρα*, urethra; *ρηνύναι*, to burst forth]. Hemorrhage from the urethra.

Urethrorrhea (*u-re-thror-el'-ah*) [*οὐρήθρα*, urethra; *ροία*, a flow]. A morbid discharge from the urethra.

Urethroscope (*u-re'-thro-skōp*) [*οὐρήθρα*, urethra; *σκοπεῖν*, to view]. An instrument for inspecting the interior of the urethra.

Urethroscopy (*u-re-thros'-ko-pē*) [*οὐρήθρα*, urethra; *σκοπεῖν*, to view]. Inspection of the urethra.

Urethrospasm (*u-re'-thro-spasm*) [*οὐρήθρα*, urethra; *σπασμός*, spasm]. A spasmodic stricture of the urethra.

Urethrostenosis (*u-re-thro-sten-ō'-sis*) [*οὐρήθρα*, urethra; *στένωσις*, a constriction]. Stricture of the urethra.

Urethrotome (*u-re'-thro-tōm*) [*οὐρήθρα*, urethra; *τομή*, a cutting]. An instrument used for performing urethrotomy.

Urethrotomy (*u-re-throt'-o-me*) [*οὐρήθρα*, urethra; *τομή*, a cutting]. The operation of cutting a stricture of the urethra. **U.**, External, division of a stricture by an incision from without. **U.**, Internal, division of a urethral stricture from within the urethra.

Urethrovaginal (*u-re-thro-vaj'-in-al*) [*οὐρήθρα*, urethra; *vagina*, vagina]. Pertaining to the urethra and the vagina.

Urethrovvesical (*u-re-thro-ves'-ik-al*) [*οὐρήθρα*, urethra; *vesica*, bladder]. Pertaining to the urethra and the bladder.

Uric Acid (*u'-rik*) [*οὐρον*, urine], $\text{C}_5\text{H}_4\text{N}_4\text{O}_6$. A dibasic acid; one of the nitrogenous end-products of metabolism. It is found in the urine and in the spleen. Its accumulation in the blood (lithemia, uricacidemia) is associated with marked disturbances, and according to some writers produces the phenomena of gout. It may also form urinary calculi.

Uricacidemia (*u-rik-as-id-el-me-ah*) [*οὐρον*, urine; *acidus*, acid; *αἷμα*, blood]. The presence of an excess of uric acid in the blood; also the condition associated with an excess of uric acid in the blood, and characterized by various nervous symptoms, and perhaps also responsible for the phenomena of gout.

Uricemia (*u-re-sel-me-ah*). See *Uricacidemia*.

Uridrosis (*u-rid-ro'-sis*) [*οὐρον*, urine; *ἰδρως*, sweat]. The secretion of a sweat containing urea, uric acid, or other urinary constituents which may be deposited in

crystalline form upon the skin (U. crystallina).

Urina (*u-ri'-nah*) [L.]. Urine.

Urinal (*u'-rin-al*) [*urina*, urine]. A vessel for receiving urine.

Urinalysis (*u-rin-al'-is-is*) [*urina*, urine; *λίσσις*, a loosening]. Analysis of the urine.

Urinary (*u'-rin-a-re*) [*urina*, urine]. Pertaining to the urine. **U. Organs**, the organs concerned in the secretion and excretion of the urine, including the kidneys, ureters, bladder, and urethra. **U. Sediment**, the deposit formed after urine has been allowed to stand for some hours in a vessel.

Urinat (*u'-rin-at*) [*urin*, urine]. To discharge urine from the bladder.

Uration (*u-rin-a'-shun*) [*urina*, urine]. The act of voiding urine.

Urine (*u'-rin*) [*urina*]. The fluid excreted by the kidneys. In health urine has an amber color, a slightly acid reaction, a faint odor, a saline taste, and a specific gravity of about 1018. The average quantity excreted in 24 hours is about 1200 c.c., or 40 ounces, of which about 62.5 grams are solids. Among the solids urea is the most important, of which 35 grams are excreted in the day. Other solid constituents are uric acid (0.75 grams in the day), hippuric acid, creatinin and other extractives, sodium chlorid, phosphates and sulphates of calcium, magnesium, potassium, and sodium. The most important abnormal constituents, present in disease, are albumin, sugar, blood, pus, acetone, diacetic acid, fat, chyle, tube-casts, various cells, and bacteria.

Urinemia (*u-rin-i'-me-ah*) [*urina*, urine; *aima*, blood]. The presence of urinary constituents in the blood; uremia.

Uriniferous (*u-rin-if'-er-us*) [*urina*, urine; *ferre*, to bear]. Carrying or conveying urine, as, *e. g.*, U. tubule.

Uriniparous (*u-rin-ipl'-ar-us*) [*urina*, urine; *parere*, to produce]. Secreting urine.

Urinogenital (*u-rin-o-jen'-it-al*) [*urina*, urine; *γεννᾶν*, to produce]. Urogenital.

Urinology (*u-rin-ol'-o-je*). See *Urology*.

Urinometer (*u-rin-om'-et-er*) [*urina*, urine; *μετρεω*, measure]. An hydrometer for ascertaining the specific gravity of urine.

Urinometry (*u-rin-om'-et-er*) [*urina*, urine; *μετρεω*, measure]. The determination of the specific gravity of the urine by means of the urinometer.

Urinose, Urinous (*u'-rin-ös, u'-rin-us*) [*urina*, urine]. Having the characters of urine, as, *e. g.*, a U. odor.

Uro- (*u'-ro-*) [*οὔρον*, urine]. A prefix denoting pertaining to urine or uric acid.

Urobacillus (*u-ro-bas-il'-us*) [*οὔρον*, urine; *bacillus*]. A bacillus occurring in urine. See *Bacteria*, Table of.

Urobilin (*u-ro-bil'-in*) [*οὔρον*, urine; *bilis*,

bile], $C_{22}H_{30}N_4O_7$. A yellowish-brown, amorphous pigment derived from bilirubin. It is the principal pigment of the urine, and is increased in febrile and other conditions. U.-jaundice, a jaundice supposed to be due to the presence of U. in the blood.

Urobilinuria (*u-ro-bil-in-u'-re-ah*) [*οὔρον*, urine; *bilis*, bile; *urina*, urine]. The presence of an excess of urobilin in the urine.

Urocanin (*u-ro-kan'-in*). See *Urocaninic Acid*.

Urocaninic Acid (*u-ro-kan-in'-ik*) [*οὔρον*, urine; *canis*, dog], $C_8H_8N_2O_4 + 2H_2O$. An acid found in the urine of dogs when there is a diminution in the amount of urea. By heating it is decomposed into carbonic acid, water, and a base, Urocanin, $C_{11}H_{10}N_4O$.

Urocele (*u'-ro-sel*) [*οὔρον*, urine; *κύημα*, a tumor]. A swelling of the scrotum from extravasation of urine.

Urochrome (*u'-ro-krom*) [*οὔρον*, urine; *χρῶμα*, color]. A yellow coloring matter found in urine.

Urocyanogen (*u-ro-si-an'-o-gen*) [*οὔρον*, urine; *κύανος*, blue]. A blue pigment found in urine.

Uredema (*u-re-dé-mah*). See *Uredema*.

Uroerythrin (*u-ro-er'-ih-rin*) [*οὔρον*, urine; *ἐρυθρός*, red]. An amorphous, reddish pigment with an acid reaction, occurring in the urine in rheumatic and other diseases.

Urofuscohematin (*u-ro-fus-ko-hem'-at-in*) [*οὔρον*, urine; *fuscus*, dark; *aima*, blood]. A red pigment derived from hematin, occurring in the urine.

Urogenital (*u-ro-jen'-it-al*) [*οὔρον*, urine; *γεννᾶν*, to produce]. Pertaining to the urinary and genital organs. **U. Sinus**, the anterior part of the cloaca, into which the urogenital ducts open.

Uroglauca (*u-ro-glau'-sin*) [*οὔρον*, urine; *γλαυκός*, green]. A blue pigment, at times occurring in urine, as, *e. g.*, in scarlet fever, and supposed to result from the oxidation of a chromogen.

Urogravimeter (*u-ro-grav-im'-et-er*). See *Urinometer*.

Urohematin (*u-ro-hem'-at-in*) [*οὔρον*, urine; *hematin*]. The coloring-matter of the urine, so called because it is a modified hematin.

Urolith (*u'-ro-lith*) [*οὔρον*, urine; *λίθος*, a stone]. A calculus occurring in the urine.

Urology (*u-rol'-o-je*) [*οὔρον*, urine; *λογία*, science]. The study of the urine.

Urolutein (*u-ro-lu'-te-in*) [*οὔρον*, urine; *luteus*, yellow]. A yellow pigment sometimes found in urine.

Uromelanin (*u-ro-mel'-an-in*) [*οὔρον*, urine; *melanin*]. A black pigment sometimes found in the urine, derived from the decomposition of urochrome.

Uromelus (*u-rom'-e-lus*) [*οὔρον*, tail; *μύτος*,

limb]. A monster in which there is more or less complete fusion of the limbs, with but a single foot.

Urometer (*u-rom'-et-er*). See *Urinometer*.

Uropittin (*u-ro-pit'-in*) [*οὔρον*, urine; *πίττα*, pitch], $C_9H_6N_2O_3$. A nitrogenous derivative of urochrome.

Uroplania (*u-ro-pla'-ne-ah*) [*οὔρον*, urine; *πλάνη*, a wandering]. The presence of urine in other localities than the urinary organs; the discharge of urine from an abnormal orifice.

Uropoiesis (*u-ro-poi-e'-sis*) [*οὔρον*, urine; *ποιέειν*, to make]. The secretion of the urine by the kidneys.

Uropoietic (*u-ro-poi-et'-ik*) [*οὔρον*, urine; *ποιέειν*, to make]. Concerned in uropoiesis.

Urorhodin (*u-ro-rhod'-in*) [*οὔρον*, urine; *ρόδον*, rose]. A red pigment found in urine and derived from uroxanthin.

Urorosein (*u-ro-ro'-se-in*) [*οὔρον*, urine; *ρόσα*, rose]. A rose-colored pigment found in the urine in various diseases.

Urorubin (*u-ro-ru'-bin*) [*οὔρον*, urine; *rubet*, red]. A red pigment obtained by treating urine with hydrochloric acid, and also in the preparation of uropittin.

Urorubrohematin (*u-ro-ru-bro-hem'-at-in*) [*οὔρον*, urine; *rubet*, red; *hematin*]. A pigment obtained from urine of leprosy patients.

Urosacin (*u-ro'-sas-in*). See *Urorhodin*.

Uroscopy (*u-ro-sk'-o-pe*) [*οὔρον*, urine; *σκοπεῖν*, to view]. Examination of the urine.

Urosepsis (*u-ro-sep'-sis*) [*οὔρον*, urine; *σῆψις*, sepsis]. The condition of intoxication due to the extravasation of urine.

Uroseptic (*u-ro-sep'-tik*) [*οὔρον*, urine; *sepsis*]. Relating to or characterized by urosepsis.

Urostealith (*u-ro-ste'-al-ith*) [*οὔρον*, urine; *στίπας*, fat; *λίθος*, stone]. A fat-like substance occurring in some urinary calculi.

Urotheobromin (*u-ro-the-o-bro'-min*). See *Paraxanthin*.

Urotoxic (*u-ro-toks'-ik*) [*οὔρον*, urine; *τοξικόν*, poison]. 1. Pertaining to poisonous substances eliminated in the urine. 2. Pertaining to poisoning by urine or some of its constituents. **U. Coefficient**, the number of urotoxics formed in 24 hours by one kilogram of an individual. The normal urotoxic coefficient in man is about 0.4, i.e., a man produces for each kilogram of body-weight .4 urotoxics, or sufficient poison to kill 400 grams of animal.

Urotoxy (*u-ro-toks'-e*) [*οὔρον*, urine; *τοξικόν*, poison]. The unit of toxicity of urine—the amount necessary to kill a kilogram of living substance.

Uroxanthin (*u-ro-san'-thin*) [*οὔρον*, urine; *ξανθός*, yellow]. A yellow pigment occurring in human urine and yielding indigo-blue on oxidation.

Urrhodin (*u'-rod-in*). See *Urorhodin*.

Urain (*ur'-sin*). See *Arbutin*.

Urtica (*ur'-tik-ah*) [L.]. Nettle. 1. A genus of plants of the order Urticaceæ. *U. dioica*, the common nettle, and *U. urens*, the dwarf nettle, are used as diuretics, local irritants, and hemostatics. 2. A wheel.

Urticaria (*ur'-tik-a'-re-ah*) [*urtica*, a nettle]. Nettle-rash, hives; a disease of the skin characterized by the development of wheals, which give rise to sensations of burning and itching. They appear suddenly in large or small numbers, remain for from a few minutes to several hours, and disappear suddenly. The disease may be acute or chronic, and is due to agencies acting upon the vasomotor system, such as gastrointestinal disorders, the ingestion of certain foods, as shell-fish, strawberries, etc. *U. bullosa*, *U. characterized* by the formation of bullæ. *U. factitia*, *U. Factitious*, dermatographia, dermatographism, the form produced in individuals with an irritable skin by any slight external irritation. *U. Giant*-, *U. gigans*. See *U. edematosa*. *U. hæmorrhagica*, purpura urticans, a variety characterized by hemorrhage into the wheals from rupture of the extremely congested capillaries. *U. medicamentosa*, *U. from the use of certain drugs*. *U. edematosa*, giant-urticaria, a variety characterized by the sudden appearance of large, soft, edematous swellings of the skin and subcutaneous tissue, which may measure several inches in diameter. *U. papulosa*, lichen urticatus, a form occurring in children, in which, as a result of the inflammatory effusion, a small, solid papule remains after the subsidence of the wheal. *U. pigmentosa*, a rare type which begins within the first few months of life and consists of large, reddish, wheal-like tubercles that eventually change to a brownish-red or yellowish color. *U. tuberosa*, a form in which the wheals assume a tuberosus form and become very large, as big as a walnut, hen's egg, or even larger. *U. vesiculosa*, *U. characterized* by the presence of vesicles. **Urticarial**, **Urticarious** (*ur'-tik-a'-re-al*, *ur'-tik-a'-re-us*) [*urticaria*, nettle-rash]. Pertaining to urticaria.

Urtication (*ur'-tik-a'-shun*) [*urtica*, nettle]. 1. Flagellation with nettles, a method of treatment formerly employed in paralysis and to produce local irritation. 2. A sensation as if one had been stung by nettles.

Ustilago (*us'-til-a'-go*) [L.]. A genus of parasitic fungi, the smuts. *U. maydis*, corn-smut, a fungus parasitic upon maize or Indian corn. In properties it resembles ergot of rye.

Ustulation (*us-tu-la'-shun*) [*ustulatio*]. The act of roasting, drying, or parching.

Ustus (*us'-tus*) [*uvere*, to burn]. Calcined; burned.

Uterine (*u'-ter-in*) [*uterus*]. Pertaining to the uterus.

Uteritis (*u'-ter-i'-tis*) [*uterus*, *uterus*; *itis*, inflammation]. See *Metritis*.

Utero- (*u'-ter-o-*) [*uterus*, *uterus*]. A prefix denoting pertaining to the uterus.

Uteroabdominal (*u'-ter-o-o-ab-dom'-in-al*) [*uterus*; *abdomen*]. Pertaining to the uterus and the abdomen.

Uterofixation (*u'-ter-o-fiks-a'-shun*). See *Hysteropexy*.

Uterogestation (*u'-ter-o-jes-ta'-shun*) [*uterus*; *gestatio*, gestation]. Gestation within the cavity of the uterus.

Uteroovarian (*u'-ter-o-o-va'-re-an*) [*uterus*; *ovary*]. Pertaining to the uterus and the ovaries.

Uteropexia, Uteropexy (*u'-ter-o-peks'-e-ah*, *u'-ter-o-peks-e*). See *Hysteropexy*.

Uteroplacental (*u'-ter-o-pla-sen'-tal*) [*uterus*; *placenta*]. Pertaining to the uterus and the placenta.

Uterosacral (*u'-ter-o-sa'-kral*) [*uterus*; *sacrum*]. Pertaining to the uterus and the sacrum.

Uterotomy (*u'-ter-o'-o-me*). See *Hysterotomy*.

Uterus (*u'-ter-us*) [L.]. The womb, the organ of gestation, receiving the ovum in its cavity, retaining and supporting it during the development of the fetus, and becoming the principal agent in its expulsion during parturition. It is a pear-shaped, muscular organ, three inches long, two inches wide, and one inch thick, and is divided into three portions, the fundus, the body, and the cervix. The fundus is the upper and broad portion; the body gradually narrows to the neck, which is the contracted portion. The orifice, os uteri, communicates with the vagina. The inner surface is covered with mucous membrane continuous with that of the vagina. The outer surface of the fundus and body is covered with peritoneum. The whole organ is suspended in the pelvis by means of the broad ligaments. The Fallopian tubes enter, one on either side of the fundus, at the cornua of the organ. **U. bicornis**, a uterus divided into two horns or compartments on account of an arrest of development. **U. cordiformis**, a heart-shaped uterus, a form due to faulty development. **U. duplex**, a uterus that is double from failure of the Müllerian ducts to unite. **U., Gravid**, a pregnant uterus. **U. masculinus**, the little passage formed by the sinus pocularis in the substance of the prostate. **U. unicornis**, a

uterus having but a single lateral half with usually only one Fallopian tube; it is the result of faulty development.

Utricle (*u'-trik'-l*) [*utriculus*, dim. of *uter*, a small bag]. 1. A delicate membranous sac communicating with the semicircular canals of the ear. 2. The sinus pocularis of the prostate gland.

Utricular (*u'-trik'-u-lar*) [*utriculus*]. 1. Pertaining to the utricle. 2. Pertaining to the uterus.

Utriculus (*u'-trik'-u-lus*) [L.]. See *Utricle*.

U. hominis. See *Uterus masculinus*.

Uva (*u'-vah*) [L.]. A grape. **U. ursi**, the *Arctostaphylos uva ursi*, or bearberry, of the order Ericaceæ. Its leaves (*Uva ursi*, U. S. P., *Uvæ ursi folia*, B. P.) contain a bitter, crystalline glucosid, arbutin, $C_{12}H_{18}O_6$, splitting up into glucose and hydroquinone, $C_6H_4O_2$. *Uva ursi* is astringent and tonic, and is used in chronic nephritis, pyelitis, cystitis, incontinence of urine, gleet, leukorrhea, etc. Dose gr. xx- $\overline{3}$ (1.3-4.0). Preparations and doses: *Extractum uvæ ursi* (U. S. P.), gr. xv-xxx (1.0-2.0); *Extractum uvæ ursi fluidum* (U. S. P.), f $\overline{3}$ (4.0); *Infusum uvæ ursi* (B. P.), f $\overline{3}$ -ij (32.0-64.0).

Uvea (*u'-ve-ah*) [*uva*, grape]. The pigmented layer of the eye, comprising the iris, ciliary body, and choroid.

Uveal (*u'-ve-al*) [*uvea*, a grape]. Pertaining to the uvea.

Uveitis (*u'-ve-i'-tis*) [*uvea*; *itis*, inflammation]. Inflammation of the uvea.

Uvula (*üv'-u-lah*) [L.]. The conic appendix hanging from the free edge of the soft palate and formed by muscles (*azygos uvulæ*, *levator* and *tensor palati*), mucous membrane, and connective tissue. **U. of the Cerebellum**, a small lobule of the inferior vermis of the cerebellum, forming the posterior boundary of the fourth ventricle. **U., Vesical**, a prominence at the internal orifice of the urethra.

Uvulapptosis (*üv-u-lap-to'-sis*) [*uvula*, *uvula*; *πτῶσις*, falling]. A relaxed and pendulous condition of the uvula.

Uvular (*üv'-u-lar*) [*uvula*]. Pertaining to the uvula.

Uvulatome (*üv'-u-lat-öm*) [*uvula*, *uvula*; *τομή*, cutting]. An instrument used in performing uvulotomy.

Uvulotomy (*üv-u-lat'-o-me*) [*uvula*, *uvula*; *τομή*, a cutting]. The operation of cutting off the uvula.

Uvulitis (*üv-u-li'-tis*) [*uvula*, *uvula*; *itis*, inflammation]. Inflammation of the uvula.

Uvuloptosis (*üv-u-lap-to'-sis*). See *Uvulapptosis*.

V

V. 1. Abbreviation of vision, or acuity of vision. 2. The symbol of vanadium.

Vaccina (*vak-si'-nah*). See *Vaccinia*.

Vaccinal (*vak'-sin-al*) [*vacca*, a cow]. Pertaining to vaccination or to vaccine.

Vaccinate (*vak'-sin-at*) [*vacca*, a cow]. 1. To inoculate with the virus of vaccinia. 2. To inoculate with any virus in order to produce immunity against an infectious disease.

Vaccination (*vak-sin-a'-shun*) [*vacca*, a cow]. Inoculation with the virus of cow-pox in order to protect from small-pox.

Vaccinator (*vak'-sin-a-tor*) [*vacca*, a cow]. One who vaccinates.

Vaccine (*vak'-sen*) [*vacca*, cow]. The virus used in performing vaccination. **V.**, **Bovine**, that derived from the cow. **V.**, **Humanized**, that from vaccinal vesicles of man. **V.-lymph**. See *Vaccine*.

Vaccinella (*vak-sin-el'-ah*) [*vacca*, cow]. Spurious vaccinia.

Vaccinia (*vak-sin'-e-ah*) [*vacca*, cow]. Cow-pox, a contagious disease of cows transmissible to man by vaccination and conferring immunity against small-pox. In the human subject inoculated with cow-pox a small papule appears at the site of inoculation in from one to three days, which becomes a vesicle about the fifth day, and at the end of the first week is pustular, umbilicated, and surrounded by a red areola. Desiccation begins in the second week and a scab forms, which soon falls off, leaving a white, pitted cicatrix.

Vacciniola (*vak-sin-i'-o-lah*) [dim. of *vaccinia*]. A secondary eruption, sometimes following vaccinia and resembling the eruption of small-pox.

Vaccinophobia (*vak-sin-o-fo'-be-ah*) [*vacca*, cow; *phóbos*, fear]. Morbid dread of vaccination.

Vaccinosyphilis (*vak-sin-o-sif'-il-is*) [*vacca*, cow; *syphilis*, syphilis]. Syphilis conveyed by vaccination with contaminated virus.

Vacuolar (*vak'-u-o-lar*) [*vacuus*, empty]. Pertaining to or of the nature of a vacuole.

Vacuolation (*vak'-u-o-la'-shun*) [*vacuus*, empty]. The formation of vacuoles; the state of being vacuolated.

Vacuole (*vak'-u-ol*) [*vacuus*, empty]. A clear space filled with air or fluid in the interior of a cell.

Vacuum (*vak'-u-um*) [L.]. A space from which the air has been exhausted.

Vagabonds' Disease. Parasitic melanoderma, a pigmentation of the skin from chronic irritation by pediculi.

Vagal (*va'-gal*) [*vagare*, to wander]. Pertaining to the vagus nerve.

Vagina (*va-ji'-nah*) [L.]. 1. A sheath. 2. The musculomembranous canal extending from the vulval opening to the cervix uteri, ensheathing the latter and the penis during copulation. **V. cordis**, the pericardium. **V. femoris**, the fascia lata.

Vaginal (*vaj'-in-al*) [*vagina*, a sheath]. 1. Pertaining to or of the nature of a sheath, as, *e. g.*, the **V. tunic** (tunica vaginalis of the testicle). 2. Pertaining to the vagina.

Vaginalis (*vaj-in-a'-lis*) [*vagina*, a sheath]. Vaginal.

Vaginismus (*vaj-in-is'-mus*) [*vagina*, sheath]. Painful spasm of the vagina.

Vaginitis (*vaj-in-i'-tis*) [*vagina*, sheath; *itis*, inflammation]. Inflammation of the vagina.

Vagino- (*vaj'-in-o-*) [*vagina*, vagina]. A prefix denoting pertaining to the vagina.

Vagino-vesical (*vaj-in-o-ves'-ik-al*). See *Vesicovaginal*.

Vagitus (*va-ji'-tus*) [*vagire*, to cry]. The cry of an infant. **V. uterinus**, the cry of a child while still in the uterus.

Vagotomy (*va-got'-o-me*) [*vagus*, *vagus*; *tomē*, a cutting]. Division of the vagus nerve.

Vagus (*va'-gus*) [*vagare*, to wander]. The pneumogastric nerve. See *Nervus*, *Table of V.-pneumonia*, pneumonia following section of the vagi in the lower animals, and due to the aspiration of food into the air-passages.

Valence, Valency (*va'-lens*, *va'-len-se*) [*valere*, to be worth]. The relative combining capacity of an atom compared with that of the atom of hydrogen.

Valerian (*val'-e-re-an*). The *Valeriana officinalis* and other species of the order *Valerianæ*. Its root (*Valeriana*, U. S. P., *Valerianæ rhizoma*, B. P.) contains a volatile oil, from which valeric acid is obtained. **V.** is employed as a mild nervous stimulant in hysteria, migraine, low fevers, etc. Dose of the oil, \mathfrak{m} iv-v (0.26-0.32). Preparations and doses: *Extractum valerianæ fluidum*, U. S. P., $\mathfrak{f}\mathfrak{z}$ (4.0); *Infusum valerianæ*, B. P., $\mathfrak{f}\mathfrak{z}$ -ij (32.0-64.0); *Tinctura valerianæ*, U. S. P., B. P., and *Tinctura valerianæ ammoniata*, U. S. P., B. P., $\mathfrak{f}\mathfrak{z}$ -ij (4.0-12.0).

Valerianate (*val'-e-re-an-at*). A salt of valeric acid. The following are official: those of ammonium, iron, quinin, and zinc.

Valerianic Acid (*val-e-re-an'-ik*). See *Acid*, *Valeric*.

Valeric Acid (*va-le'-rik*). See *Acid*.

Valgus (*val'gus*) [L.]. Bowed out. Genu valgum, bow-legged; Talipes valgus, Pes valgus, or simply V., splay-foot, a condition in which the arch of the foot is depressed so that the inner side of the sole rests upon the ground.

Vallecula (*val-ek'-u-lah*) [*vallis*, valley]. A shallow groove or depression. **V. cerebelli**, the depression between the cerebellar hemispheres.

Valleix's Points. Painful points observed in neuralgia at the point of exit of the nerve. See *Points douloureux*.

Vallet's Mass. See *Ferrum*.

Valley of the Cerebellum. See *Vallecula cerebelli*.

Valli-Ritter's Law. See *Law*.

Valsalva's Experiment [*Valsalva*, an Italian anatomist]. Inflation of the tympanic cavity by expiring forcibly when the mouth and nose are tightly closed.

Valsalva's Sinus. See *Sinus*.

Value, Globular (*valeur globulaire*). A fraction of which the numerator is the percentage of hemoglobin, and the denominator the percentage of red corpuscles. It indicates the percentage of hemoglobin in a corpuscle.

Valve (*valv*) [*valva*, door]. 1. A device placed in a tube or canal so as to permit free passage one way, but not in the opposite direction. 2. A fold of membrane acting as a valve, as V. of the heart. **V., Aortic**, the valve consisting of three semilunar segments, situated at the junction of the aorta with the heart. **V., Bauheini's**, the ileocecal V. **V., Bicuspid.** See *V., Mitral*. **V., Coronary**, the valve protecting the orifice of the coronary sinus and preventing regurgitation of blood during the contraction of the right auricle. **V., Eustachian**, that between the inferior vena cava and the right auricle of the fetus. **V. of Hasner**, an imperfect V. at the inferior meatus of the nose. **V., Heister's**, a fold of mucous membrane at the neck of the gall-bladder. **V., Ileocecal**, the folds of mucous membrane at the junction of the ileum and cecum. **V. of Kerkring**, any one of the valvæ conniventes. **V., Mitral**, the V. that controls the opening from the left auricle to the left ventricle; it is constituted of two leaflets. **V., Pulmonary**, the valve composed of three semilunar leaflets, and situated at the junction of the pulmonary artery and the right ventricle. **V., Semilunar**, the three valves guarding the orifice of the pulmonary artery and aorta. **V.-test** (*Azoulay's*), auscultation of the heart while the patient is lying with the arms raised perpendicularly and the legs lifted obliquely. **V. of Thebesius**, the coronary valve or fold of the endocardium of the right auricle which protects the coronary sinus. **V., Tricuspid**, that which controls the opening from the

right auricle to the right ventricle; it consists of three segments. **V. of Varolius**, the ileocecal valve. **V. of Vieussens**, a thin leaf of medullary substance forming the roof of the anterior portion of the fourth ventricle of the brain.

Valvula (*val'-vu-lah*) [dim. of *valva*, a valve]. A small valve. **Valvulæ conniventes** (valves of Kerkring), the transverse folds of mucous membrane of the small intestine.

Valvular (*val'-vu-lar*) [*valvula*, a small valve]. Pertaining to or originating at a valve.

Valvulitis (*val-vu-li'-tis*) [*valvula*, a small valve; *itis*, inflammation]. Inflammation of a valve, especially of a cardiac valve.

Vanadium (*van-a'-de-um*) [*Vanadis*, a goddess of the Scandinavian mythology]. A rare metallic element. See *Elements, Table of*.

Vanilla (*van-il'-ah*) [L.]. A genus of plants of the order Orchideæ. The fruit of *V. planifolia* (V., U. S. P.) contains an aromatic crystalline principle, **Vanillin**, $C_8H_8 \cdot OCH_3 \cdot OH \cdot COH = C_8H_8O_3$, which is the methyl-ether of protocatechuic aldehyd. V. is used as a flavoring agent and as an ingredient of a test-solution for hydrochloric acid (phloroglucin-vanillin test).

Vanillism (*van-il'-izm*) [*vanilla*]. A form of dermatitis, characterized by marked itching, occurring among vanilla-workers.

Van Swieten's Liquor or Solution (*van swet'-tens*). A solution of mercuric chlorid 2 grains, alcohol 3 drams, distilled water sufficient to make 4 ounces.

Vapor (*va'-por*) [L.]. A gas, especially the gaseous form of a substance which at ordinary temperatures is liquid or solid.

Vapores uterini. Synonym of *Hysteria*.

Vaporarium (*va-por-a'-re-um*) [L.]. A vapor-bath; an establishment for giving vapor-baths.

Vaporization (*va-por-i-za'-shun*) [*vapor*, vapor]. The conversion of a solid or liquid into a vapor.

Vaporize (*va'-por-iz*) [*vapor*, vapor]. To convert into vapor.

Varicella (*var-is-el'-ah*) [dim. of *variola*, small-pox]. Chicken-pox; an acute, contagious disease of childhood, characterized by an eruption of transparent vesicles, which appear in successive crops on different parts of the body. The incubation-period is about two weeks. The disease is usually mild, and runs its course in two or three days. In rare cases complications, such as nephritis, develop.

Varicelloid (*var-is-el'-oid*) [*varicella*, *varicella*; *eidos*, like]. Resembling varicella.

Varices (*var'-is-ēs*) [L.]. Plural of *Varix*.

Variciform (*var-is'-if-orm*) [*varix*, a swollen vein; *forma*, a form]. Having the form of a varix.

Varicocele (*var'-ik-o-sel'*) [*varix*, varix; *κήλη*, a tumor]. Dilatation of the veins of the spermatic cord, forming a soft, elastic swelling.

Varicose (*var'-ik-ös*) [*varix*, varix]. 1. Of blood-vessels, swollen, knotted, and tortuous. 2. Due to V. veins, as V. ulcer. V. **Aneurysm**. See *Aneurysm*.

Varicosity (*var-ik-ös'-it-e*) [*varix*, varix]. The condition of being varicose; a varicose portion of a vein.

Variola (*va-ri'-o-lah*) [*varius*, variegated, spotted]. Small-pox, a contagious infectious disease ushered in with severe febrile symptoms, which, in the course of two or three days, are followed by a papular eruption spreading over all parts of the body. During the succeeding two weeks the eruption passes through the stage of vesicles and pustules, the latter going on to the formation of crusts. The falling off of the crusts leaves a pitted appearance of the skin (pock-marks). The period of incubation is about thirteen days. V. **confuens**, V. **confluent**, a severe form, in which the pustules spread and run together. V., **Discrete**, a form in which the pustules preserve their distinct individuality. V., **Hemorrhagic**, small-pox in which hemorrhage occurs into the vesicles, which gives them a blackish appearance. V., **Malignant**, black small-pox, a severe and very fatal form of the hemorrhagic type. V., **Mitigated**, V., **Modified**. See *Varioloid*.

Variolation, **Variolization** (*var-e-o-la'-shun*, *var-e-o-li-sa'-shun*) [*variola*, variola]. The inoculation of small-pox.

Varioliform (*var-i'-o-lif-orm*) [*variola*, variola; *forma*, form]. Resembling variola.

Varioloid (*var'-e-o-loid*) [*variola*, variola; *ειδος*, like]. A mild form of variola occurring in persons that have been vaccinated or inoculated with small-pox virus.

Variolous (*var-i'-o-lus*) [*variola*]. Pertaining to or having the nature of variola.

Varix (*va'-riks*) [*varus*, crooked]. A dilated and tortuous vein. V., **Aneurysmal**. See *Aneurysmal Varix*, under *Aneurysm*, *Arteriovenous*. V. **lymphaticus**, dilatation of the lymphatic vessels, especially that due to the *Filaria sanguinis hominis*.

Varus (*va'-rus*). Bowed in. Genu varum, in-knee; Talipes varus, Pes varus, or simply V., cross foot, a condition in which the foot is turned inward.

Vas (*vas*) [L.]. A vessel. V. **aberrans**, a blind tube projecting from the lower part of the epididymis. V. **deferens**, the excretory duct of the testis.

Vasa (*va'-zah*) [L.]. Plural of *vas*, a vessel.

V. **afferentia**, the branches of a lymphatic or lacteal vessel entering a lymphatic gland.

V. **brevia** (short vessels), the gastric branches of the splenic artery. V. **effluentia**. 1. The terminal ducts of the rete testis. 2. The efferent vessels of lymphatic glands. V. **intestini tenuis**, small vessels arising from the superior mesenteric artery and distributed to the jejunum and ileum. V. **recta**, the tubules of the rete testis. V. **vasorum**, the vessels supplying the arteries and veins with blood. V. **vorticosa**. See *Vena vorticosa*.

Vascular (*vas'-ku-lar*) [*vasculum*, a small vessel]. Consisting of, pertaining to, or provided with vessels.

Vascularity (*vas'-ku-lar'-it-e*) [*vasculum*, a small vessel]. The quality of being vascular.

Vascularization (*vas'-ku-lar-i-sa'-shun*) [*vasculum*, a small vessel]. The process of becoming vascular.

Vasculum (*vas'-ku-lum*) [L.]. A small vessel. V. **aberrans**. See *Vas aberrans*.

Vaselin (*vas'-el-in*) [Ger., *Wasser*, water; *έλαιον*, oil]. See *Petrolatum*.

Vasifactive (*va'-ze-fak'-tiv*) [*vas*, vessel; *facere*, make]. Giving rise to new blood-vessels.

Vaso- (*va'-so-*) [*vas*, a vessel]. A prefix denoting pertaining to a vessel.

Vasoconstrictor (*va-so-kon-strik'-tor*) [*vas*, vessel; *constringere*, to constrict]. See *Vasomotor*.

Vasodentine (*va-so-den'-tin*) [*vas*, vessel; *dens*, a tooth]. Dentine possessing blood-vessels.

Vasodilator (*va-so-di-la'-tor*) [*vas*, vessel; *dilator*, a dilator]. See *Vasomotor*.

Vasofactive (*va-so-fak'-tiv*). See *Vasoformative*.

Vasoformative (*va-so-for'-mat-iv*) [*vas*, vessel; *formare*, to form]. Forming or producing vessels.

Vasohypertonic (*va-so-hi-per-ton'-ik*). See *Vasomotor*.

Vasohypotonic (*va-so-hi-po-ton'-ik*). See *Vasomotor*.

Vasoinhibitory (*va-so-in-hib'-it-o-re*) [*vas*, vessel; *inhibere*, to inhibit]. Inhibiting vasomotor action, especially vasoconstrictor action.

Vasomotor (*va-so-mo'-tor*) [*vas*, vessel; *motor*, from *movere*, to move]. Regulating the tension of blood-vessels. V. **Centers**, centers situated in the medulla oblongata and spinal cord, and governing the caliber of the blood-vessels. V. **Nerves**, the nerves passing to the blood-vessels; they are of two kinds, the vasoconstrictor (vasohypertonic) nerves, or those stimulation of which causes contraction of the blood-vessels, and the vasodilator (vasohypotonic) nerves, stimula-

tion of which causes dilatation of the vessels.

Vasotonic (*va-zo-ton'ik*) [*vas*, vessel; *tonic*]. Pertaining to the normal tone of the blood-vessels.

Vastus (*vas'tus*). 1. Large; extensive.

2. A large muscle of the thigh. See *Muscles*, Table of.

Vater's Ampulla (*fah't-er-a*) [*Vater*, a German anatomist]. See *Ampulla*.

Vater's Corpuscles. See *Pacinian Corpuscles*.

Vectis (*vek'tis*) [*vehere*, to carry]. An instrument similar to the single blade of a forceps, used in hastening the delivery of the fetal head in labor.

Vegetable (*vej-et-ahl*) [*vegetare*, to quicken]. 1. A plant, especially one used as food. 2. See *Vegetal*. **V. Albumin**. See *Phytalbumose*. **V. Proteids**. See *Proteid*. **V. Sulphur**. See *Lycopodium*.

Vegetal (*vej-et-ahl*) [*vegetare*, to quicken]. Of or pertaining to plants; characteristic of plants.

Vegetarian (*vej-et-ah'-re-an*) [*vegetare*, to quicken]. One who lives on vegetable food alone.

Vegetarianism (*vej-et-ah'-re-an-izm*) [*vegetare*, to quicken]. 1. The doctrine that vegetable food is the only kind proper for man. 2. The practice of living only on vegetable food.

Vegetation (*vej-et-ah'-shun*) [*vegetare*, to quicken]. An outgrowth resembling a plant in outline, as the fibrinous projections on the cardiac valves in endocarditis, papillomata, polypoid growths, etc. **V., Adenoid**, growths of lymphoid tissue in the nasopharyngeal cavity.

Vegetative (*vej-et-ah'-tiv*) [*vegetare*, to quicken]. Having the power of growth, like a plant.

Vehicle (*veh'-ih-k'l*) [*vehiculum*, from *vehere*, to carry]. An excipient or substance serving as a medium of administration of medicines.

Vein (*van*) [*vena*]. A blood-vessel carrying blood from the tissues to the heart. Veins, like arteries, have three coats, but less well developed; many also possess valves. **V., Angular**, a continuation of the frontal vein downward to become the facial at the lower margin of the orbit. **V., Axillary**, a large vein formed by the junction of the brachial veins. **Veins, Azygos** (three in number), situated in front of the bodies of the thoracic vertebrae; they are a means of communication between the superior and inferior venæ cavæ. **V., Basilar**, a large vein passing back over the crus cerebri to unite with the veins of Galen. **V., Basilic**, a vein on the inner side of the arm. **Veins, Brachial**, the veins

accompanying the brachial artery. **V., Brachiocephalic**. See *V., Innominate*.

Veins of Breschet, the veins of the diploe.

V., Cephalic, a large vein of the arm, formed by the union of the median cephalic and superficial radial and opening into the axillary vein. **V., Coronary**. 1. The great cardiac vein, a vein opening into the coronary sinus of the heart. 2. See *V., Gastric*.

Veins, Emissary, small veins passing through the cranial foramina and connecting the cerebral sinuses with external veins. **V., Facial**, a continuation of the angular vein; it joins the internal jugular at the level of the hyoid bone. **V., Femoral, Common**, a short, thick trunk corresponding to the femoral artery; it becomes the external iliac at Poupart's ligament. **V., Femoral, Deep**, a vein accompanying the femoral artery; it empties into the superficial femoral. **V., Femoral, Superficial**, a name given to the femoral vein before it is joined by the deep femoral vein to form the common femoral vein.

Veins of Galen, two large veins of the brain, continuations of the internal cerebral veins, and opening into the straight sinus. **V., Gastric**, a vein accompanying the artery of the same name. **Veins, Hemizygos**, small, accessory veins of the azygos veins. **Veins, Hemorrhoidal**, a plexus of veins surrounding the rectum. **V., Iliac, Common**, a vein formed opposite the sacroiliac synchondrosis by the confluence of the external and internal iliac veins. **V., Iliac, External**, a continuation upward of the common femoral; it extends from the lower border of Poupart's ligament to the lower border of the sacroiliac synchondrosis. **V., Iliac, Internal**, a short trunk extending from the great sacroiliac synchondrosis. **Veins, Innominate**, two large valveless veins returning the blood from the head, neck, and upper extremity. **V., Jugular, Anterior**, a vein beginning at the level of the chin and ending at the clavicle in the external jugular vein. **V., Jugular, External**, a vein formed at the angle of the lower jaw by the union of the posterior auricular and temporomaxillary veins; it empties into the subclavian. **V., Jugular, Internal**, a continuation of the lateral sinus, beginning at the jugular fossa, accompanying the internal and common carotid arteries, and joining the subclavian vein to form the innominate. **V. of Marshall**. See *V., Oblique*. **V., Median Basilic**, a vein uniting with the superficial ulnar to form the basilic. **V., Median Cephalic**, a vein uniting with the superficial radial to form the cephalic. **V., Median, Deep**, a vein formed by the union of the outer vena comes of the ulnar artery and the muscular and radial recurrent veins. **V., Median, Superficial**,

one starting at the anterior plexus of the wrist and uniting with the deep median to form the median. **V.**, **Oblique** (*of Marshall*), a vein crossing the dorsal portion of the left auricle of the heart. It is the remnant of the left duct of Cuvier. **V.**, **Ophthalmic**, a short trunk carrying the blood from the eye and emptying into the cavernous sinus. **V.**, **Popliteal**, one formed by the union of the venæ comites of the anterior and posterior tibial arteries; it accompanies the popliteal artery, and becomes the femoral vein at the junction of the lower with the middle third of the thigh. **V.**, **Portal**, a short trunk entering the liver at the transverse fissure and formed by the junction of the superior mesenteric and splenic veins. **Veins**, **Pulmonary**, four veins, two from each lung, returning the aerated blood from the lungs to the heart. **V.**, **Radial**, **Superficial**, a vein accompanying the musculocutaneous nerve up the radial side of the forearm. **V.**, **Renal**, a vein accompanying the renal artery. **V.**, **Saphenous**, **Long or Internal**, a long superficial vein running up the inner aspect of the leg and thigh, terminating in the femoral vein below Poupert's ligament. **V.**, **Saphenous**, **Short or External**, a superficial vein running up the outer aspect of the foot, leg, and back of the calf, and emptying into the popliteal vein. **V.**, **Spermatic**, one returning the blood from the testicle; on the right side it terminates in the inferior vena cava and on the left in the left renal vein. **V.**, **Splenic**, one returning the blood from the spleen, and forming the portal vein by its union with the superior mesenteric vein. **V.**, **Subclavian**, a continuation of the axillary vein, uniting with the internal jugular vein to form the innominate vein at the sternoclavicular articulation. **V.**, **Temporomaxillary**, one formed by the union of the temporal and internal maxillary veins in the parotid gland; it terminates in the external jugular. **V. of Trolard**, a vein of the cerebrum passing along the posterior branch of the fissure of Sylvius and emptying into the superior petrosal sinus. **V.**, **Ulnar**, one running from the wrist up the anterior and inner surface of the forearm. **V.**, **Umbilical**, a vein conveying the blood from the placenta to the fetus.

Velamentous (*vel-am-en'-tus*) [*velamen*, a veil]. Resembling a veil.

Vella's Fistula. An intestinal fistula for obtaining gastric juice.

Vellication (*vel-ik-a'-shun*) [*vellicare*, to twitch]. Spasmodic twitching of muscular fibers.

Velosynthesis (*ve-lo-sin'-thes-is*). See *Staphylo-rhaphy*.

Velpeau's Bandage (*vel'-pōs*) [*Velpeau*,

a French surgeon]. A bandage used in fractured clavicle.

Velum (*ve'-lum*) [*velum*, veil, sail]. A veil or veil-like structure. **V.**, **Anterior Medullary**. See *Vieussens*, *Valve of*. **V. interpositum**, the membranous roof of the third ventricle. **V. palati**, the soft palate. **V.**, **Posterior Medullary**, the commissure of the flocculus of the cerebellum.

Vena (*ve'-nah*) [*L.*]. A vein. See *Vein*. **V. cava inferior**, a vein formed by the junction of the two common iliac veins and emptying into the right auricle of the heart. It receives lumbar, right spermatic, renal, suprarenal, phrenic, and hepatic veins. **V. cava superior**, a vein formed by the union of the innominate veins, and conveying the blood from the upper half of the body to the right auricle. **V. comes**, **Venæ comites**, a vein or veins accompanying an artery in its course. **Venæ Galeni**, the two trunks formed by the union of the vein of the corpus striatum and that of the choroid plexus; they empty into the straight sinus. **Venæ Thebesii**, small veins by which blood passes from the walls of the heart to the right auricle. **Venæ vorticossæ**, the venous network and trunks of the choroid coat of the eye.

Venenation (*ven-en-a'-shun*) [*venenum*, a poison]. The condition of a poisoning.

Venenatus (*ven-en-a'-tus*) [*venenum*, a poison]. Poisonous.

Venereal (*ven-e'-re-al*) [*Venus*, the goddess of love]. Pertaining to or produced by sexual intercourse. **V. Diseases**, gonorrhea, syphilis, and chancre.

Venery (*ven'-er-e*) [*Venus*, the goddess of love]. Sexual intercourse.

Venesection (*ven-e-sek'-shun*) [*vena*, vein; *secare*, to cut]. See *Blood-letting*.

Venom (*ven'-om*) [*venenum*, poison]. Poison, especially a poison secreted by certain reptiles and insects.

Venomous (*ven'-om-us*) [*venenum*, a poison]. Poisonous; secreting venom.

Venosity (*ven-os'-it-e*) [*vena*, vein]. A condition in which the arterial blood shows venous qualities.

Venous (*ve'-nus*) [*venosus*]. Pertaining to or produced in a vein. **V. Hum**, the murmur or rushing sound heard in auscultation of a vein.

Venter (*ven'-ter*) [*L.*]. 1. The belly or abdomen. 2. The belly of a muscle. 3. The cavity of the abdomen. 4. The concavity of any expanded part, as, *e. g.*, the V. of the scapula, V. of the ilium.

Ventilation (*ven-til-a'-shun*) [*ventilare*, to fan]. The act or process of supplying fresh air; the act or process of purifying the air of a place.

Ventrad (ven'-trad) [*venter*, belly; *ad*, toward]. Toward the ventral aspect.

Ventral (ven'-tral) [*venter*, belly]. Pertaining to the belly.

Ventricle (ven'-trik-l) [*ventriculus*, dim. of *venter*, a belly]. A small cavity or pouch.

V. of Arantius, a culdesac at the lower end of the fourth V. **Ventricles of the Brain**, cavities in the interior of the brain, comprising the two lateral ventricles, the third, and the fourth ventricles. **V. of Cord**, the central canal of the spinal cord. **V. of Larynx**, a depression between the true and false vocal bands. **V., Left, of Heart**, that upon the dorsal and left side of the heart, and which, through the aorta, forces the blood throughout the body. **V., Right, of Heart**, that forcing the blood through the pulmonary artery into the lungs.

Ventricular (ven'-trik'-u-lar) [*ventriculus*, a ventricle]. Pertaining to a ventricle. **V. Aqueduct**. See *Aqueductus Sylvii*.

Ventrofixation (ven'-tro-fiks'-at'-shun) [*venter*, belly; *fixation*]. The stitching of a displaced uterus to the abdominal wall.

Ventroinguinal (ven'-tro-in'-gwin-al) [*venter*, belly; *inguinal*]. Pertaining to the abdomen and the groin.

Ventrotomy (ven'-troit'-o-me) [*venter*, belly; *tomē*, a cut]. Celiotomy.

Venule, Venula (ven'-ul, ven'-u-lah) [*venula*, dim. of *vena*, a vein]. A small vein.

Veratrin (ver-at'-rin). *Veratrina* (U. S. P.), a mixture of alkaloids obtained from the seeds of *Asagrea officinalis* (Sabadilla), of the order Liliaceae. It is a local irritant, and produces tetanic convulsions followed by paralysis; it first stimulates, then paralyzes the vasomotor center. It is used externally, in the form of an ointment, in rheumatism, gout, and neuralgia. Preparations: Unguentum veratrinæ (U. S. P., B. P.); Oleatum veratrinæ (U. S. P.).

Veratroidin (ver-at'-roi'-din). See *Veratrum*.

Veratrum (ver-at'-rum). A genus of plants of the order Liliaceae. The rhizome and roots of *V. viride*, American hellebore (*V. viride*, U. S. P.; *Veratri viridis* rhizoma, B. P.), contain the alkaloids jervin and veratroidin. The former is a depressant to the vasomotor centers and the motor centers of the spinal cord. In toxic doses it produces slowness of the pulse, fall in blood-pressure, relaxation, epileptiform convulsions, paralysis, and death from failure of the respiration. Veratroidin is irritant and produces vomiting and purging; it is also depressant to the motor centers of the spinal cord and to the pulse. When veratrum viride is administered the combined action of the alkaloids is obtained, and consists chiefly in slowing of the pulse and lessening of blood-pressure, with vomiting in the case of large doses. It is employed in

sthenic inflammations, as pneumonia, peritonitis, in puerperal eclampsia, and in excessive cardiac hypertrophy. Preparations and doses: Extractum veratri viridis fluidum (U. S. P.), \mathfrak{m} j-ij (0.065-0.20); Tinctura veratri viridis (U. S. P., B. P.), \mathfrak{m} j-ij (0.065-0.20). **V. album**, white hellebore, resembles *V. viride* in general properties.

Verbascum (ver-bas'-kum) [L.]. Mullein, a genus of plants of the order Scrophulariaceae. The leaves and flowers of *V. thapsus* have been used as demulcent in catarrhal inflammation of mucous membranes and as an application to hemorrhoids.

Verdigris (ver'-dig-ris) [Fr., *verd de gris*, probably from L., *viridis*, green; *as*, brass]. Copper acetate.

Verheyen, Stars of. Venous plexuses of stellate form situated on the surface of the kidney, beneath its capsule.

Vermicide (ver'-mis-id) [*vermis*, a worm; *caedere*, to kill]. An agent that destroys intestinal worms.

Vermicular (ver-mik'-u-lar) [*vermis*, worm]. Worm-like.

Vermiculation (ver-mik-u-lar'-shun) [*vermis*, worm]. A worm-like motion; peristaltic motion.

Vermiform (ver'-mif-orm) [*vermis*, worm; *forma*, a form]. Worm-shaped. **V. Appendix**. See *Appendix*. **V. Process, Inferior, V.**, Superior, the inferior and superior surfaces of the middle lobe of the cerebellum.

Vermifuge (ver'-mif-aj) [*vermis*, worm; *fugare*, to expel]. An agent that expels intestinal worms.

Vermination (ver-min'-at'-shun) [*vermis*, worm]. Infestation with worms.

Vermineous (ver'-min-us) [*vermis*, worm]. Infested with worms.

Vermis (ver'-mis) [L.]. 1. A worm. 2. The middle lobe of the cerebellum.

Vernal (ver'-nal) [*vernalis*, of the spring]. Pertaining to the spring. **V. Conjunctivitis**, a form of conjunctivitis recurring each spring or summer, and disappearing with frost.

Vernin (ver'-nin), $C_{16}H_{20}N_6O_6$. A leuko-main-base found in young vetch, clover, ergot, etc., and yielding guanin on heating with hydrochloric acid.

Vernix caseosa [L., cheesy varnish]. A sebaceous deposit covering the surface of the fetus.

Verruca (ver'-ru'-kah) [L.]. Wart. **V. necrogenica**, anatomic tubercle, dissection-tubercle, a warty excrescence found on the fingers of those who frequently handle the tissues of tuberculous subjects.

Verrucose, Verrucous (ver'-u-kos, ver'-u-kus) [*verruca*, wart]. Warty; covered with or having warts.

Version (*ver'-shun*) [*vertere*, to turn]. Turning; an operation whereby one part of the fetus is made to replace another at the mouth of the uterus. **V.**, **Bipolar**, **V.** by acting upon both poles of the fetus. **V.**, **Cephalic**, turning of the fetus so as to bring the head to present. **V.**, **Combined**, bipolar **V.**, consisting of a combination of external and internal **V.** **V.**, **External**, that effected by external manipulation. **V.**, **Internal**, that performed by entering the hand within the uterus. **V.**, **Podalic**, that in which one or both feet are brought to the mouth of the uterus. **V.**, **Spontaneous**, the process whereby, without external influence, a transverse position is changed into a longitudinal one.

Vertebra (*ver'-teb-rah*) [*vertere*, to turn]. One of the bones forming the spinal or vertebral column. There are 33 vertebræ, divided into 7 cervical, 12 thoracic or dorsal, 5 lumbar, 5 sacral (the sacrum), 4 coccygeal (the coccyx). A typical vertebra consists of a body and an arch, the latter being formed by two pedicles and two laminae. The arch supports seven processes: 4 articular, 2 transverse, and 1 spinous. **V.**, **Basilar**, the last lumbar vertebra. **V. dentata**, the axis. **V. prominens**, the seventh cervical vertebra.

Vertebral (*ver'-teb-ral*) [*vertebra*]. 1. Pertaining to or characteristic of a vertebra; made up of or possessing vertebræ. 2. Pertaining to the vertebral artery. **V. Artery**. See *Arteries*, Table of.

Vertebralarterial (*ver-teb-rar-tel'-re-al*) [*vertebra*, vertebra; *ἀρτηρία*, artery]. Giving passage to the vertebral artery, as the **V. foramina** in the transverse processes of the cervical vertebræ.

Vertebrate, **Vertebrated** (*ver'-teb-rat*, *ver'-teb-ra-ted*) [*vertebra*, vertebra]. Having a vertebral column; resembling a vertebral column in flexibility, as, e. g., a **V. catheter**.

Vertebro- (*ver'-teb-ro-*) [*vertebra*, vertebra]. A prefix denoting pertaining to a vertebra.

Vertebrochondral (*ver-teb-ro-kon'-dral*) [*vertebra*, vertebra; *χονδρός*, cartilage]. Connecting the costal cartilages with the vertebræ.

Vertebrocostal (*ver-teb-ro-kos'-tal*) [*vertebra*, vertebra; *costa*, rib]. Pertaining to the vertebræ and the ribs.

Vertebroiliac (*ver-teb-ro-ill'-e-ak*) [*vertebra*, vertebra; *ilium*, ilium]. Pertaining to the vertebræ and the ilium.

Vertebrosacral (*ver-teb-ro-sa'-kral*) [*vertebra*, vertebra; *sacrum*, sacrum]. Pertaining to the vertebræ and the sacrum.

Vertebrosteral (*ver-teb-ro-stur'-nal*) [*vertebra*, vertebra; *sternum*]. Extending from the spinal column to the sternum. **V. Ribs**, the true ribs.

Vertex (*ver'-teks*) [*L.*]. The crown or top of the head. **V.-presentation**, a presentation of the vertex of the fetal skull.

Vertical (*ver'-tik-al*) [*vertex*]. 1. Pertaining to the vertex. 2. Perpendicular.

Verticomenal (*ver-tik-o-men'-tal*) [*vertex*, vertex; *mentum*, chin]. Pertaining to the vertex and the chin.

Vertiginous (*ver-tij'-in-us*) [*vertere*, to turn]. Resembling or affected with vertigo.

Vertigo (*ver'-tig-o*) [*L.*, from *vertere*, to turn]. Giddiness, dizziness; a sensation of lack of equilibrium. It may be due to disease of the ears (*auditory* or *aural V.*), the eyes (*ocular V.*), the brain (*cerebral V.*), the stomach (*gastric V.*), the blood, etc. **V.**, **Epileptic**, vertigo associated with or preceding an attack of epilepsy. **V.**, **Essential**, one not due to any discoverable cause. **V.**, **Labyrinthine**. See *Ménière's Disease*. **V.**, **Objective**, one in which objects seem to the patient to move. **V.**, **Paralyzing**. See *Gerlier's Disease*. **V.**, **Subjective**, one in which the patient has a sensation as if he himself were moving.

Verumontanum (*ver-u-mon-tal'-num*) [*veru*, a spit; *mons*, a mountain]. The caput gallinaginis, a longitudinal ridge on the floor of the prostatic urethra.

Vesalius, Foramen of. A small foramen in the greater wing of the sphenoid bone, transmitting a small vein.

Vesica (*ves'-ik-ah*) [*L.*]. The bladder.

Vesical (*ves'-ik-al*) [*vesica*, a bladder]. Pertaining to the bladder. **V. Calculus**, a stone in the bladder. **V. Crisis**, severe paroxysmal pain in the bladder occurring in locomotor ataxia.

Vesicant (*ves'-ik-ant*) [*vesicare*, to blister]. 1. Blistering. 2. A blistering agent.

Vesication (*ves'-ik-a'-shun*) [*vesicare*, to blister]. The formation of a blister; a blister.

Vesicatory (*ves'-ik-at-o-re*) [*vesicare*, to blister]. 1. Blistering. 2. A blistering agent.

Vesicle (*ves'-ik-l*) [*vesicula*, dim. of *vesica*, bladder]. 1. A small bladder; especially a small sac containing fluid. **V.**, **Allantoic**, the internal hollow portion of the allantois. **V.**, **Auditory**, an ectodermic sac, a part of the cerebral vesicle, from which the internal ear is formed. **V.**, **Blastodermic**. See *Blastoderm*. **V.**, **Cerebral**, an expansion of the anterior extremity of the neural tube of the embryo, subsequently forming the brain. **V.**, **Germinal**, the nucleus of the ovum. **V.**, **Graafian**. See *Follicle*, *Graafian*. **V.**, **Optic**, a hollow process of the cerebral vesicle forming the essential part of the eye. **V.**, **Otic**. See *V.*, *Auditory*. **V.**, **Seminal**, one of the two little sacs situated at the base of the bladder and serving as reservoirs for the semen. **V.**, **Umbilical**. See *Yolk-sac*.

2. A small blister on the skin, as, *e. g.*, an herpetic or small-pox vesicle.

Vesico- (*ves' ik-o*) [*vesica*, bladder]. A prefix denoting pertaining to the bladder.

Vesicoabdominal (*ves-ik-o-ab-dom' in-al*) [*vesica*, bladder; *abdomen*, abdomen]. Pertaining to the abdomen and the urinary bladder.

Vesicocele (*ves' ik-o-sel*) [*vesica*, bladder; *κηλη*, hernia]. Hernia of the bladder.

Vesicocervical (*ves-ik-o-ser' vik-al*) [*vesica*, bladder; *cervix*, cervix]. Pertaining to the cervix uteri and the urinary bladder.

Vesicoprostatic (*ves-ik-o-prov-lat' ik*) [*vesica*, bladder; *προστάτης*, prostate]. Pertaining to the prostate gland and the urinary bladder.

Vesicorectal (*ves-ik-o-rek' tal*) [*vesica*, bladder; *rectum*, rectum]. Pertaining to the bladder and the rectum.

Vesicospinal (*ves-ik-o-spi' nal*) [*vesica*, bladder; *spina*, spine]. Pertaining to the urinary bladder and the spinal cord.

Vesicumbilical (*ves-ik-o-um- bil' ik-al*) [*vesica*, bladder; *umbilicus*, umbilicus]. Pertaining to the umbilicus and the urinary bladder.

Vesicouterine (*ves-ik-o-ut' ter-in*) [*vesica*, bladder; *uterus*, uterus]. Pertaining to the uterus and the urinary bladder.

Vesicovaginal (*ves-ik-o-vaj' in-al*) [*vesica*, bladder; *vagina*, vagina]. Pertaining to the bladder and the vagina.

Vesicula (*ves-ik' u-lah*) [dim. of *vesica*, a bladder]. A vesicle. **V. graafiana**. See *Follicle*, *Graafian*. **Vesiculæ Nabothi**. See *Ovule*, second definition. **V. prostatica**, the sinus pularis. **Vesiculæ seminales**. See *Vesicle*, *Seminal*.

Vesicular (*ves-ik' u-lar*) [*vesicula*, a little bladder]. 1. Pertaining to or composed of vesicles. 2. Produced in vesicles, as, *e. g.*, *V. breathing*, *V. murmur*. **V. Column**, a column of ganglion-cells at the base of the posterior horn of the spinal cord. **V. Column, Posterior**. See *Column of Clarke*. **V. Eczema**, eczema attended with the formation of vesicles. **V. Rale**, the crepitant rale.

Vesiculated (*ves-ik' u-la-ted*) [*vesicula*, a little bladder]. Composed of vesicles.

Vesiculation (*ves-ik-u-la' shun*) [*vesicula*, a little bladder]. The formation of vesicles; the state of becoming vesiculated.

Vesiculitis (*ves-ik-u-lit' tis*) [*vesicula*, vesicle; *itis*, inflammation]. Inflammation of the seminal vesicles.

Vesiculobronchial (*ves-ik-u-lo-brong' ke-al*) [*vesicula*, vesicle; *βρόγχος*, bronchus]. Both vesicular and bronchial.

Vesiculocavernous (*ves-ik-u-lo-kav' er-nus*) [*vesicula*, vesicle; *cavernous*]. Both vesicular and cavernous.

Vesiculotympanitic (*ves-ik-u-lo-tim-pan- it' ik*) [*vesicula*, vesicle; *τύμπανον*, drum]. Both vesicular and tympanitic.

Vessel (*ves' el*) [Fr., from *vasculum*, vessel]. A receptacle for fluids, especially a tube or canal for conveying blood or lymph.

Vestibular (*ves- tib' u-lar*) [*vestibulum*, porch]. Pertaining to the vestibule.

Vestibule (*ves- tib' al*) [*vestibulum*, a porch]. An approach; an antechamber. **V. of the Ear**, the oval cavity of the internal ear, which forms the entrance to the cochlea. **V. of the Nose**, the anterior part of the nostrils. **V. of the Vagina**, **V. of the Vulva**, a triangular space below the clitoris and between the nymphæ.

Vestigial (*ves- tij' e-al*) [*vestigium*, a vestige]. Of the nature of a vestige or trace; rudimentary. **V. Fold**, a fibrous band of the pericardium representing the obliterated left innominate vein.

Vesuvius (*ves- u' vin*) [*Vesuvius*, a volcano near Naples]. Bismarck brown, triamido-benzol; it is used as a stain in microscopy.

Veterinarian (*vet' er-in- a' re-an*) [*veterinarius*, from *veterina*, a beast of burden]. One who practices veterinary medicine.

Veterinary (*vet' er- in- a- re*) [*veterinarius*, from *veterina*, a beast of burden]. Pertaining to the domestic animals. **V. Medicine**, medicine as applied to the domestic animals.

Via (*vi' ah*) [L.]. A way. **Via naturales**, the natural passages. See *Prima via*.

Viability (*vi- ab' il- it- e*) [*viabilis*, viable]. The state of being viable.

Viable (*vi' ab- l*) [*viabilis*, viable]. Capable of living; likely to live; applied to a fetus capable of living outside of the uterus.

Vial (*vi' al*) [φιάλη]. A small bottle.

Vibex (*vi' beks*) [L., *pl.*, *Vibices*]. A linear ecchymosis.

Vibrate (*vi' brat*) [*vibrare*, to shake]. To move to and fro.

Vibratile (*vi' bra- til*) [*vibrare*, to shake]. Moving to and fro; vibrating.

Vibration (*vi- bra' shun*) [*vibrare*, to shake]. The act of moving to and fro.

Vibrio (*vi' b- re-o*) [*vibrare*, to vibrate]. A genus of Schizomycetes. See *Bacteria*, *Table of*.

Vibrissa (*vi' ris- ah*) [L.]. One of the hairs near the opening of the anterior nares.

Viburnum (*vi- bur' num*). A genus of the Caprifoliaceæ. **V. opulus**, cranberry-tree, cramp-bark (*V. opulus*, U. S. P.), is used in dysmenorrhea, scurvy, asthma, etc. Dose of Extractum viburni opuli fluidum, U. S. P., f3j-ij (4.0-8.0). **V. prunifolium**, U. S. P., is used in dysmenorrhea, threatened abortion, menorrhagia, etc. Dose of Extractum viburni prunifolii fluidum, U. S. P., f3j-ij (4.0-8.0).

Vicarious (*vi- ka' re-us*) [*vices*, changes]. Taking the place of something else; of an

habitual discharge, occurring in an abnormal situation, as, *e. g.*, V. menstruation.

Vice (*vis*) [*vitium*, vice]. 1. A physical defect, as, *e. g.*, a vice of conformation. 2. A moral defect; a bad habit.

Vichy Water (*ve-she*). A mildly laxative and antacid mineral water obtained from Vichy, in France, and used in rheumatic and gouty conditions and in disorders of the liver.

Vicq d'Azyr's Bundle (*vik-dah-sers*) [*Vicq d'Azyr*, a French anatomist]. A bundle of nerve-fibers passing from the mammillary body to the anterior nucleus of the optic thalamus.

Vidian (*vid'-e-an*). Described by or named after Vidus, an Italian anatomist, as, *e. g.*, the V. canal in the sphenoid bone; the V. artery, which traverses the V. canal; the V. nerve, which also passes through the V. canal, and is formed by the union of the great petrosal nerve and carotid branches of the sympathetic, and enters Meckel's ganglion.

Vienna Paste (*ve-en'-ah*). See *Potassium*.

Viessens, Valve of [*Viessens*, a French anatomist]. A thin lamina of gray matter stretched between the two processus cerebelli ad testes, and forming the roof of the aqueduct of Sylvius and of part of the fourth ventricle.

Vigouroux's Sign (*vig-oo-rooz*). See *Signs and Symptoms, Table of*.

Villatte's Liquor. A preparation used for injecting into carious bones, consisting of zinc sulphate and copper sulphate, each 15 grains, liquor plumbi subacetatis, half a dram, and dilute acetic acid, three and a half drams.

Villous, Villous (*vil'-ds*, *vil'-us*) [*villus*, a tuft of hair]. Pertaining to a villus; covered with villi; characterized by the formation of villus-like projections.

Villosity (*vil-os'-it-e*) [*villus*, a tuft of hair]. The state of being villous.

Villus (*vil'-us*) [L., a tuft of hair]. 1. One of the minute club-shaped projections from the mucous membrane of the intestine, consisting of a lacteal vessel, an arteriole, and a vein, enclosed in a layer of epithelium. 2. One of the vascular tufts of the chorion.

Vinasse (*ve-nas'*) [Fr.]. Potash obtained from the marc or residue of the wine-press.

Vinculum (*vin'-ku-lum*) [L.]. A band.

Vincula accessoria tendinum, the slender tendinous filaments which connect the phalanges with the flexor tendons.

Vinegar (*vin'-e-gar*) [Fr., *vin*, wine; *aigre*, sour]. 1. An impure solution of acetic acid, obtained by acetous fermentation of wine, beer, cider, etc., or by the dry distillation of wood. It is used as a condiment. 2. A solution of a medicinal substance in vinegar or acetic acid. The vinegars are those of

cantharides (Acetum cantharidis, B. P.), of ipecac (Acetum ipecacuanhæ, B. P.), of opium (Acetum opii, U. S. P.), and of squill (Acetum scillæ, U. S. P., B. P.).

Vinous (*vi'-nus*) [*vinum*, wine]. Having the nature of wine; containing wine.

Vinum (*vi'-num*) [L.]. See *Wine*.

Viola (*vi'-o-lah*) [L.]. A genus of plants of the Violaceæ, including V. tricolor, heartsease, V. odorata, V. cucullata. V. odorata as well as other species are used in bronchitis.

Violet (*vi'-o-let*) [*viola*, violet]. 1. One of the colors of the spectrum, very closely resembling the purple of violets and possessing the greatest refrangibility of the spectral colors. 2. A violet dye-stuff. V., *Gentian*, a violet anilin dye used for staining in histologic and bacteriologic work. V., *Methyl-*. See *Methyl-violet*.

Violinist's Cramp, Violin-player's Cramp. An occupation-neurosis occurring in violin-players, and characterized by spasm of the fingers used in playing.

Virgin (*vir'-jin*) [*virgo*]. A woman who has never had sexual intercourse.

Virginal (*vir'-jin-al*) [*virgo*, a virgin]. Pertaining to virginity.

Virginité (*vir-jin'-it-e*) [*virgo*, a virgin]. The condition of being a virgin.

Virile (*vir'-il*) [*virilis*, from *vir*, a man]. Pertaining to or characteristic of the man. V. Member, the penis.

Virility (*vir-il'-it-e*) [*virilitas*]. The condition of being virile; procreative power.

Virtual Focus (*vir'-tu-al fo'-kus*). See *Focus*.

Virus (*vi'-rus*) [L.]. The poison of an infectious disease, especially one found in the secretion or tissues of an individual or animal suffering from an infectious disease. V., *Humanized*, vaccine-lymph taken from the vaccine-pustule of a human subject.

Vis [L.]. Force; energy; power. V. a *fronte*, a force that attracts. V. a *tergo*, a force that pushes something before it. V. *formativa*, energy manifesting itself in the formation of new tissue to replace that which has been destroyed. V. *medicatrix naturæ*, the healing power of nature apart from medicinal treatment.

Visceral (*vis'-er-al*) [*viscus*, a viscus]. Pertaining to a viscus. V. *Clefts*. See *Cleft, Visceral*.

Viscid (*vis'-id*) [*viscidus*]. Sticky; adhesive; glutinous.

Viscosity (*vis-id'-it-e*) [*viscidus*, viscid]. The state of being viscid. Same as *Viscosity*.

Viscin (*vis'-in*) [*viscum*, bird-lime]. A mucilaginous extract of mistletoe.

Viscose (*vis'-kds*) [*viscum*, bird-lime]. A gummy product of viscous fermentation.

Viscose, Viscous (*vis' - kōs, vis' - kus*). Viscid.

Viscosity (*vis-kōs' - it - ē*). The state of being viscous.

Viscum (*vis' - kum*) [L.]. Mistletoe, a genus of plants of the order Loranthaceæ, growing as parasites upon trees. *V. album*, European mistletoe, and *V. flavescens*, or *Phoradendron flavescens*, American mistletoe, contain a viscid principle, *viscin*, which is the chief constituent.

Viscus (*vis' - kus*) [L. : *pl., Viscera*]. Any one of the organs enclosed within one of the four great cavities, the cranium, thorax, abdominal cavity, or pelvis; especially one within the abdominal cavity.

Visible (*vis' - ibl*) [*videre, to see*]. Capable of being seen.

Visibility (*vis - ib - it' - ē*) [*videre, to see*]. The state of being visible.

Vision (*vi'zh' - un*) [*videre, to see*]. The act of seeing; sight. *V.*, **Binocular**. See *Binocular*. *V.*, **Central**, vision with the macula lutea. *V.*, **Direct**. See *V., Central*. *V.*, **Double**. See *Diplopia*. *V.*, **Indirect**, vision with other parts of the retina than the macula. *V.*, **Solid**, or *V.*, **Stereoscopic**, the perception of relief or depth of objects obtained by binocular vision.

Visual (*vi'zh' - u - al*) [*visualis, from videre, to see*]. Pertaining to vision. *V. Purple*, a pigmentary substance in the retina reacting to light in a peculiar manner, and thought to be intimately connected with vision. See *Rhodopsin*.

Visuoauditory (*vis - u - o - au' - dit - o - rē*) [*videre, to see; audire, to hear*]. Pertaining to hearing and seeing; of nerve-fibers, connecting the visual and auditory centers.

Vital (*vi' - tal*) [*vita, life*]. Pertaining to life. *V. Capacity*, the volume of air that can be expelled from the lungs after a full inspiration.

V. Statistics, statistics of the births, deaths, marriages, and diseases in a community.

Vitalism (*vi' - tal - izm*) [*vita, life*]. The doctrine that ascribes the phenomena exhibited by living organisms to the action of a vital force distinct from mechanic or chemic force.

Vitalist (*vi' - tal - ist*) [*vita, life*]. A believer in vitalism.

Vitals (*vi' - talz*) [*vita, life*]. A vulgar term for the organs essential to life.

Vitellary (*vit' - el - a - rē*) [*vitellus, yolk*]. Pertaining to the vitellus.

Vitellin (*vit' - el' - in*) [*vitellus, yolk*]. A globulin found in egg-yolk.

Vitelline (*vit' - el' - in, vit' - el - in*) [*vitellus, yolk*]. Pertaining to the vitellus or yolk.

V. Artery, an artery passing from the yolk-sac to the primitive aorta of the embryo.

V. Duct, the omphalomesenteric duct.

V. Membrane, the true membrane of the

ovum, lying inside of the zona pellucida. *V.*

Veins, veins returning the blood from the yolk-sac to the primitive heart of the embryo.

Vitellus (*vit' - el' - us*) [L.]. A yolk, specifically the yolk of the egg of the common fowl, *Gallus domesticus*.

Vitiligo (*vit' - il' - i' - gō*) [L.]. Piebald skin, a disease of the skin characterized by a disappearance of the natural pigment, occurring in patches and leaving whitish areas.

Vitiligoidea (*vit' - il' - ig - oi' - de - ak*) [*vitiligo; eidos, like*]. Xanthoma.

Vitreous (*vit' - re - us*) [*vitreus, from vitrum, glass*]. Glassy. *V. Chamber*, the portion of the globe of the eye posterior to the crystalline lens.

V. Degeneration, hyaline degeneration. *V. Humor*, *V. Body*, or simply the **Vitreous**, the transparent, jelly-like substance filling the posterior chamber of the eye.

V. Table, the hard, brittle, inner table of the skull.

Vitriol (*vit' - re - ol*) [*vitriolum, from vitrum, glass*]. 1. Sulphuric acid, more commonly called oil of *V.* 2. Any crystalline salt of sulphuric acid. *V.*, **Blue**, copper sulphate. *V.*, **Green**, ferrous sulphate or copperas. *V.*, **White**, zinc sulphate.

Vitrum (*vit' - rum*) [L.]. Glass.

Vivification (*viv' - if - ik - a' - shun*) [*vivus, living; facere, to make*]. The act of making alive or of converting into living tissue.

Viviparity (*viv' - ip - ar' - it - ē*) [*vivus, alive; parere, to bring forth*]. The bringing forth of living offspring; the state of being viviparous.

Viviparous (*viv' - ip' - ar - us*) [*vivus, alive; parere, to bring forth*]. Bringing forth the young alive, distinguished from *oviparous*.

Vivisection (*viv' - is - ek' - shun*) [*vivus, alive; secare, to cut*]. The dissection of a living animal; experimentation upon an animal while still alive.

Vivisector (*viv' - is - ek' - tor*) [*vivus, alive; secare, to cut*]. One who practises vivisection.

Vlemmickx's Solution. An application used for treating acne, consisting of lime 1 part, sulphur 2 parts, and water 20 parts.

Vocal (*vo' - kal*) [*vox, the voice*]. Pertaining to the voice; pertaining to the organs producing the voice. *V. Bands*, *V. Cords*. See *Larynx*. *V. Fremitus*, the thrill conveyed to the hand when applied to the chest during speaking. *V. Resonance*, the resonance produced by the voice as heard on auscultating the lung.

Voice (*vois*) [*vox*]. The sounds, especially articulate sounds, produced by the vibration of the vocal bands and modified by the resonance-organs.

Vola (*vo' - lah*) [L.]. The palm of the hand or the sole of the foot.

Volar (*vo'-lar*) [*vola*]. Pertaining to the palm or the sole.

Volatile (*vol'-at-il*) [*volatilis*, from *volare*, to fly]. Passing into vapor at ordinary temperatures; evaporating. **V. Oils**. See *Oils, Essential*.

Volatilization (*vol-at-il-i-sa'-shun*) [*volare*, to fly]. The act of volatilizing.

Volatilize (*vol'-at-il-iz*) [*volare*, to fly]. To convert into vapor by means of heat; to pass into vapor.

VolSELLA (*vol-sel'-ah*) [*vellere*, to pluck]. A forceps having one or more hooks at the end of each blade.

Volt (*volt*) [from *Volta*, an Italian scientist]. The unit of electromotive force, or the force sufficient to cause a current of one ampere to flow against a resistance of one ohm. **V.-ampere**, the amount of pressure developed by a current of one ampere having an electromotive force of one volt; also called a watt.

Voltaic (*vol-ta'-ik*). Described by or named after Volta, an Italian scientist. **V. Electricity**, galvanism.

Voltaism (*vol'-ta-izm*). See *Galvanism*.

Voltameter (*volt-am'-et-er*) [*volt*; *μτρov*, a measure]. An instrument for ascertaining the electromotive force of a current in volts.

Voltmeter (*volt'-me-ter*) [*volt*; *μτρov*, measure]. A synonym of *Voltameter*.

Voltolini's Disease (*volt-to-le'-nez*). See *Diseases, Table of*.

Volumetric (*vol-u-met'-rik*) [*volumen*, a roll; *μτρov*, measure]. Pertaining to measurement by volume. **V. Analysis**. See *Analysis*.

Volumometer (*vol-u-mom'-e-ter*) [*volumen*, volume; *μτρov*, measure]. An apparatus used for the purpose of measuring changes in volume.

Voluntary (*vol'-un-ta-re*) [*voluntarius*]. Under the control of the will; performed by an exercise of the will. **V. Muscle**, striped muscle.

Volvulus (*vol'-vu-lus*) [*volvere*, to roll]. A twisting of the bowel so as to occlude the lumen, occurring most frequently in the sigmoid flexure.

Vomer (*vo'-mer*) [*L.*, a plowshare]. The thin plate of bone situated vertically between the nasal fossæ, and forming the posterior portion of the septum of the nose.

Vomerine (*vo'-mer-in*) [*vomer*, vomer]. Pertaining to the vomer.

Vomica (*vom'-ik-ah*) [*vomere*, to vomit]. A cavity formed by the breaking down of tissue; especially a cavity in the lung.

Vomit (*vom'-it*) [*vomere*, to vomit]. 1. To expel from the stomach by vomiting. 2. Vomited matter. **V., Bilious**, vomit stained with bile. **V., Black**, the characteristic vomit of yellow fever, a dark fluid consisting of blood and the contents of the stomach. **V., Coffee-ground**, vomit consisting of broken-down blood and the contents of the stomach; it is frequently seen in carcinoma of the stomach.

Vomiting (*vom'-it-ing*) [*vomere*, to vomit]. The forcible ejection of the contents of the stomach through the mouth. **V., Pernicious**, a variety of vomiting occasionally seen in pregnancy and becoming at times so excessive as to threaten life. **V., Stercoraceous**, the ejection of fecal matter in the vomit, usually due to intestinal obstruction.

Vomito negro (*vo-me'-to na'-gro*). 1. Black vomit. 2. Yellow fever.

Vomiturition (*vom-it-u-rish'-un*) [*vomiturio*]. Ineffectual attempt at vomiting; retching.

Vomitus (*vom'-it-us*) [*vomere*, to vomit]. 1. Vomited matter. 2. The act of vomiting.

V. cruentus, bloody vomit.

Vox (*voxs*) [*L.*]. Voice.

Vulcanite (*vul'-kan-it*) [*Vulcan*, the god of fire]. Vulcanized caoutchouc.

Vulcanize (*vul'-kan-iz*) [*Vulcan*, the god of fire]. To subject india-rubber to the process of vulcanization, a process wherein it is treated with sulphur at a high temperature, and thereby is rendered either flexible or very hard (vulcanite).

Vulnerary (*vul'-ner-a-re*) [*vulnus*, a wound]. 1. Pertaining to wounds; healing wounds.

2. An agent useful in healing wounds.

Vulnus (*vul'-nus*) [*L.*]. A wound.

Vulsella (*vul-sel'-ah*). See *Volsella*.

Vulva (*vul'-vah*) [*volvere*, to roll up]. The external organs of generation in the woman.

Vulval, **Vulvar** (*vul'-val*, *vul'-var*) [*vulva*]. Pertaining to the vulva.

Vulvitis (*vul'-vi'-tis*) [*vulva*, vulva; *ιτις*, inflammation]. Inflammation of the vulva.

Vulvo- (*vul'-vo-*) [*vulva*, vulva]. A prefix denoting pertaining to the vulva.

Vulvovaginal (*vul'-vo-vaj'-in-al*) [*vulva*, vulva; *vagina*, vagina]. Pertaining to the vulva and the vagina. **V. Gland**, a small gland situated on each side of the vulva near the vagina; the gland of Bartholin.

Vulvovaginitis (*vul'-vo-vaj-in-i'-tis*) [*vulva*, vulva; *vagina*, vagina; *ιτις*, inflammation]. Inflammation of the vulva and the vagina.

W

W. The symbol of tungsten (wolfram).

Wachendorf's Membrane. The pupillary membrane.

Wafer (*waf'-fer*). A thin layer composed of moistened flour, and used to enclose powders that are taken internally.

Wagner's Corpuscles. Oval-shaped bodies at the termination of nerve-fibers, concerned in the sense of touch.

Wahoo (*wah'-hoo'*). See *Euonymus*.

Waldenburg's Apparatus (*wahl'-den-bergs*). An apparatus constructed on the principle of a gasometer, and used for compressing or rarefying air, which is inhaled, or into which the patient exhales.

Wallerian Degeneration (*wal-le'-re-an*) [from *Waller*, an English physician]. Degeneration of nerves after separation from their trophic centers.

Wall-eye. A colloquial name for leukoma of the cornea.

Walnut (*wolf'-nut*). See *Juglans*.

Wandering (*won'-der-ing*). 1. Moving about, as, *e. g.*, W. cells. 2. Abnormally movable, as, *e. g.*, W. spleen.

Warburg's Tincture. A very complex liquid first prepared by Dr. Warburg, and employed as an antiperiodic and diaphoretic. It is used especially in pernicious forms of malaria. Its composition is as follows: Aqueous extract of aloes, 28 grains; rhubarb, 448 grains; angelica-seed, 448 grains; elecampane, 224 grains; saffron, 224 grains; fennel, 224 grains; gentian, 112 grains; zedoary-root, 112 grains; cubeb, 112 grains; myrrh, 112 grains; white agaric, 112 grains; camphor, 112 grains; quinin sulphate, 1280 grains; dilute alcohol, enough to make 8 pints.

Ward's Paste. Confection of black pepper.

Wardrop's Disease. Malignant onychitis.

Wardrop's Operation. See *Operations, Table of*.

Warehousemen's Itch. Palmar eczema occurring among the workmen in warehouses.

Warm-blooded. A term applied to animals that maintain a uniform temperature, whatever the changes in the surrounding medium.

Warming Plaster. See *Cantharis* and *Pitch*.

Warren's Fat Columns. Slender columns of fatty tissue passing from the subcutaneous adipose tissue to the base of the hair-follicles. They are well developed over the dorsum of the body, particularly near the median line.

Wart [*AS., warte, wart*]. A hyperplasia of the papillae of the skin forming a small projection. See *Verruca*. **W., Anatomic,**

W., Postmortem-. See *Tubercle, Anatomic*.

Warty. Resembling a wart; covered with warts.

Wash. See *Lotion*. **W., Black.** See *Mercury*. **W., Yellow.** See *Mercury*.

Washerwoman's Itch. Eczema of the hands occurring in washerwomen.

Washing Soda. Sodium carbonate.

Wasting Palsy. See *Progressive Muscular Atrophy*.

Watchmaker's Cramp. An occupation-neurosis occurring in watchmakers, and characterized by painful cramps of the muscles of the hands.

Water (*waw'-ter*). 1. A transparent, inodorous, tasteless fluid, boiling at 212° F. (100° C.), and freezing at 32° F. (0° C.). Chemically it is hydrogen monoxid, H₂O. 2. A mineral water. 3. A solution of a medicinal substance in water, as, *e. g.*, W. of ammonia (Aqua ammoniac), chlorin-W. (Aqua chlori), etc. **W.-bag**, a rubber bag in which water of varying degrees of temperature can be introduced for topical application. **W.-bed.** See *Bed*. **W.-brash.** See *Fyrosis*. **W. of Crystallization**, the water contained in certain crystals, to which their crystalline structure is due. **W.-glass**, solution of sodium silicate. **W.-hammer Pulse.** See *Corrigan's Pulse*. **W., Hard**, water containing soluble calcium salts, and not readily forming a lather with soap. **W., Javelle**, a solution of potassium hypochlorite, KClO. **W., Mineral**, a natural water containing mineral substances in solution.

Waterborne. Of a disease, produced by contaminated drinking water.

Water on the Brain. Hydrocephalus.

Watt (*wot*) [after James Watt, a Scottish engineer]. One voltampere. See *Volt*.

Wave (*wav*) [*AS., wafian*, to waver in mind].

1. A movement in a body which is propagated with a continuous motion, each particle of the body vibrating through a fixed path, usually a closed curve. 2. One of the curves in a series of curves representing a wave-like motion. **W.-length**, the distance between corresponding points, usually the crests, of two adjacent waves. **W., Tidal**, in the sphygmogram, the wave succeeding the percussion-wave, and due to the volume of blood poured out from the heart reaching the arteries.

Wavy Respiration. Cog-wheel respiration, a type of breathing in which inspiration or expiration is not continuous, but jerky and interrupted.

Wax (*waks*). See *Cera*.

Waxy (*waks'-e*). Pertaining to or resembling wax. **W. Cast**, a tube-cast composed of amyloid or similar material. **W. Degeneration**, amyloid degeneration. **W. Kidney**, **W. Liver**, amyloid kidney or liver. **Wean** (*wen*) [*AS.*, *wenian*, to accustom]. To cease to give suck to an infant by accustoming it to take food from other sources than the mother's breast.

Webbed Fingers, Webbed Toes. Union of adjacent fingers or toes by a thin band of tissue.

Weber (*web'er*). Same as *Coulomb*.

Weber's Law (*web'-ers*) [*H. Weber*, a German physiologist]. A law that when a stimulus is increased in geometric proportion, the sensation is increased in arithmetic proportion, or, in other words, the reaction varies as the logarithm of the stimulus.

Weber's Test. A test for hearing performed by placing a vibrating tuning-fork on the vertex. In case of disease of the external or middle ear of one side, it will be heard best on the affected side; in case of disease of the internal ear of one side, it will be heard best on the unaffected side.

Weeping Eczema. Moist eczema.

Weeping Sinew. See *Ganglion*, 2d definition.

Weigert's Fibrin-stain (*wi'-gurts*) [*Wei-*

gert, a German pathologist]. A stain for fibrin, consisting in the application of a solution of gentian-violet, then one of iodine in potassium iodid, followed by one of anilin-oil and xylol. Fibrin is stained blue.

Weigert's Method. 1. A method of staining the myelin of nerve-fibers with hematoxylin.

2. A method of staining the neuroglia according to a modified Weigert's fibrin-method, after the tissue has been fixed in formalin, subjected to a mordant of copper acetate, acetic acid, and chromalum, and a reducing agent composed of potassium permanganate and a solution of sodium sulphite containing a chromogen derived from naphthalin.

Weight (*wat*). The force with which bodies tend to approach the earth's center. **W.**, **Atomic.** See *Atomic*. **W.**, **Molecular.** See *Molecular*. **Weights and Measures:** The U. S. standard unit of weight is the Troy Pound; the standard unit of liquid measure is the Winchester Wine Gallon, containing 231 cubic inches. The Imperial Gallon, adopted by Great Britain, contains 277.274 cubic inches. The standard unit of the U. S. and British linear measure is the yard. The actual standard of length of the U. S. is a brass scale 82 inches long in the U. S. Treasury Department. The yard is between the 27th and the 63d inch of this scale.

TABLE OF WEIGHTS AND MEASURES.

TROY WEIGHT.	
Grains.	
24 = 1 dwt.	
480 = 20 dwts. = 1 oz.	
5760 = 240 dwts. = 12 oz. = 1 lb. = 22.816 cu. in. of distilled water at 62° Fahr.	
AVOIRDUPOIS WEIGHT.	
Drams. Ounces.	
16 = 1 = 437.5 grains Troy.	
256 = 16 = 1 lb. = 1.2153 lb. Troy.	
6,400 = 400 = 25 lbs. = 1 quarter.	
25,600 = 1,600 = 100 " = 4 quarters = 1 cwt.	
512,000 = 32,000 = 2000 " = 80 " = 20 " = 1 ton.	
APOTHECARIES' WEIGHT.	
Grains	APOTHECARIES' MEASURE.
20 = 1 scruple.	60 minims = 1 fluidram.
60 = 3 scruples = 1 dram.	8 fluidrams = 1 fluidounce.
480 = 24 " = 8 drams = 1 oz.	16 fluidounces = 1 pint.
5760 = 288 " = 96 " = 12 ounces = 1 lb.	2 pints = 1 quart.
	4 quarts = 1 gallon.
LIQUID, OR WINE-MEASURE.	
Gills.	
1 = 7.2187 cu. in.	
4 = 1 pint = 28.875 cu. in.	
8 = 2 pints = 1 quart = 55.75 cu. in.	
32 = 8 " = 4 quarts = 1 gallon.	
2016 = 504 " = 252 " = 63 gallons = 1 hogshead.	
4032 = 1008 " = 504 " = 126 " = 2 hogsheads = 1 pipe.	
8064 = 2016 " = 1008 " = 252 " = 4 " = 2 pipes = 1 tun.	
SQUARE MEASURE.	
Sq. Inches. Sq. Feet.	
144 = 1	
1296 = 9 = 1 sq. yard.	
39,204 = 272.25 = 30.25 sq. yards = 1 perch.	
1,568,160 = 10,800 = 1210 " " = 40 perches = 1 rood.	
6,272,640 = 43,560 = 4840 " " = 160 " = 4 roods = 1 acre.	

660 WEIGHT OF WEIGHTS AND MEASURES.—Continued.

LINEAR MEASURE.

1 yard.					
3 yards	=	1 fathom.			
5.5 "	=	2.75 fathoms	=	1 perch.	
220 "	=	110 "	=	40 perches	= 1 furlong.
1760 "	=	880 "	=	320 "	= 8 furlongs = 1 mile.

DRY MEASURE.

		Pinta.	
		1 = 33.6 cubic inches.	
		2 = 1 quart = 67.2 cubic inches.	
		8 = 4 quarts = 1 gallon = 268.8 cubic inches.	
		16 = 8 " = 2 gallons = 1 peck = 537.6 cubic in.	
		64 = 32 " = 8 " = 4 pecks = 1 bushel.	

WEIGHTS.

			Avoir.		Avoir. Lb.
			Oz.		
		1	.03528	=	.0022047
		10	.3528	=	.022047
		100	3.52758	=	.2204737
		1000	35.2758	=	2.204737
		10,000		=	22.04737
		100,000		=	220.4737
		1,000,000		=	2204.737

METRIC DRY AND LIQUID MEASURE.

	Lit.	U. S. Cu. In.		U. S.
Milliliter	= .001	= .061	Liquid	.00845 gill.
			Dry	.0018 pint.
Centiliter	= .01	= .61	Liquid	.0845 gill.
			Dry	.018 pint.
Deciliter	= .1	= 6.1	Liquid	.845 gill = 0.2113 pint.
			Dry	.18 pint.
Liter	= 1	= 61.02	Liquid	2.113 pints = 1.057 quarts.
			Dry	1.8 pints = .908 quart.
Decaliter	= 10	= 610.16	Liquid	2.641 gallons.
			Dry	9.08 quarts = 1.135 pints.
Hectoliter	= 100	U. S. Cu. Ft.	Liquid	26.414 gallons.
		= 3.531	Dry	2.837 bushels.
Kiloliter	= 1000	= 35.31	Liquid	264.141 gallons.
			Dry	28.374 bushels.
Myrialiter	= 10,000	= 353.1	Liquid	2641.4 gallons.
			Dry	283.7 bushels.

METRIC LINEAR MEASURE.

	Meter.	U. S. In.	Ft.	
Millimeter*	= .001	= .03937	= .00328	
Centimeter†	= .01	= .3937	= .03280	Yd.
Decimeter	= .1	= 3.937	= .32807	= .10936
Meter	= 1	= 39.3685	= 3.2807	= 1.0936
Decameter	= 10	=	= 32.807	= 10.936 Mile.
Hectometer	= 100	=	= 328.07	= 109.36 = .661347
Kilometer	= 1000	=	= 3,280.7	= 1093.6 = .6213466
Myriameter	= 10,000	=	= 32,807	= 10,936 = 6.213466

* Nearly the $\frac{1}{25}$ of an inch. † Full $\frac{1}{2}$ of an inch.

METRIC SQUARE MEASURE.

	Sq. Meter.	U. S. Sq. In.	Sq. Ft.	Sq. Yd.	Acre.
Sq. Centimeter	= .01	= .155	=	.01196	
Sq. Decimeter	= .1	= 15.5	= .10763	=	.00025
Centiare	= 1	= 1549.88	= 10.763	= 1.196	= .0247
Are	= 10	= 154,988	= 107.63	= 11.959	= 2.47
Hectare	= 100	=	= 107,630	=	= 247
Sq. Kilometer	= 3,607	Sq. Mile.			= 247
Sq. Myriameter	= 38,607	"			= 24,708

TABLE FOR CONVERTING METRIC WEIGHTS INTO TROY WEIGHTS.

GRAMS.	EXACT EQUIVALENTS IN GRAINS.	APPROXIMATE EQUIVALENTS IN TROY WEIGHTS.				GRAMS.	EXACT EQUIVALENTS IN GRAINS.	APPROXIMATE EQUIVALENTS IN TROY WEIGHTS.				GRAMS.	EXACT EQUIVALENTS IN GRAINS.	APPROXIMATE EQUIVALENTS IN TROY WEIGHTS.			
		Ounces.	Drams.	Scruples.	Grains.			Ounces.	Drams.	Scruples.	Grains.			Ounces.	Drams.	Scruples.	Grains.
0.01	0.1543	$\frac{1}{16}$	3.0	46.297	2	$6\frac{1}{2}$	23.0	354.944	..	5	2	5
0.02	0.3086	$\frac{1}{8}$	4.0	61.729	..	1	..	$1\frac{1}{2}$	24.0	370.376	..	6	..	$10\frac{1}{2}$
0.03	0.4630	$\frac{3}{16}$	5.0	77.162	..	1	..	$1\frac{7}{8}$	25.0	385.809	..	6	1	$5\frac{1}{2}$
0.04	0.6173	$\frac{1}{4}$	6.0	92.594	..	1	1	$1\frac{1}{2}$	26.0	401.241	..	6	2	$1\frac{1}{2}$
0.05	0.7717	$\frac{5}{16}$	7.0	108.026	..	1	2	8	27.0	416.673	..	6	2	$16\frac{1}{2}$
0.06	0.9260	$\frac{3}{8}$	8.0	123.459	..	2	..	$3\frac{1}{2}$	28.0	432.106	..	7	..	$12\frac{1}{2}$
0.07	1.0803	1	9.0	138.891	..	2	..	$18\frac{1}{2}$	29.0	447.538	..	7	1	$7\frac{1}{2}$
0.08	1.2347	$1\frac{1}{8}$	10.0	154.323	..	2	1	$14\frac{1}{2}$	30.0	462.970	..	7	2	3
0.09	1.3890	$1\frac{1}{4}$	11.0	169.756	..	3	2	$9\frac{1}{2}$	31.0	478.403	..	7	2	$18\frac{1}{2}$
0.1	1.543	$1\frac{1}{2}$	12.0	185.188	..	3	..	$5\frac{1}{2}$	32.0	493.835	1	$13\frac{1}{2}$
0.2	3.086	3	13.0	200.621	..	3	1	$\frac{3}{4}$	40.0	617.294	1	2	..	$17\frac{1}{2}$
0.3	4.630	$4\frac{1}{2}$	14.0	216.053	..	3	3	16	45.0	694.456	1	3	1	$10\frac{1}{2}$
0.4	6.173	$6\frac{1}{2}$	15.0	231.485	..	3	2	$11\frac{1}{2}$	50.0	771.617	1	4	2	$11\frac{1}{2}$
0.5	7.716	$7\frac{1}{2}$	16.0	246.918	..	4	..	$6\frac{1}{2}$	60.0	925.941	1	7	1	6
0.6	9.259	$9\frac{1}{2}$	17.0	262.350	..	4	1	$2\frac{1}{2}$	70.0	1080.264	2	2	..	$\frac{1}{2}$
0.7	10.803	$10\frac{1}{2}$	18.0	277.782	..	4	1	$17\frac{1}{2}$	80.0	1234.583	2	4	1	$14\frac{1}{2}$
0.8	12.346	$12\frac{1}{2}$	19.0	293.215	..	4	2	$13\frac{1}{2}$	90.0	1388.911	2	7	..	9
0.9	13.889	14	20.0	308.647	..	5	..	8 $\frac{1}{2}$	100.0	1543.235	3	1	2	$3\frac{1}{2}$
1.0	15.432	$15\frac{1}{2}$	21.0	324.079	..	5	1	$4\frac{1}{2}$						
2.0	30.865	1	$10\frac{1}{2}$	22.0	339.512	..	5	1	$19\frac{1}{2}$						

TABLE FOR CONVERTING TROY WEIGHTS INTO METRIC WEIGHTS.

GRAINS.	GRAMS.	GRAINS.	GRAMS.	GRAINS.	GRAMS.	GRAINS.	GRAMS.	GRAINS.	GRAMS.	GRAINS.	GRAMS.	GRAINS.	GRAMS.	GRAINS.	GRAMS.
1	0.0648	16	1.037	31	2.009	46	2.980	61	3.952	76	4.924	91	5.896	300	19.440
2	0.1296	17	1.102	32	2.073	47	3.045	62	4.017	77	4.980	92	5.961	360	23.228
3	0.1944	18	1.166	33	2.138	48	3.110	63	4.082	78	5.054	93	6.026	400	25.920
4	0.2592	19	1.231	34	2.203	49	3.175	64	4.147	79	5.118	94	6.090	480	31.103
5	0.3240	20	1.296	35	2.268	50	3.234	65	4.211	80	5.183	95	6.155	500	32.396
6	0.3888	21	1.361	36	2.332	51	3.304	66	4.276	81	5.248	96	6.220	600	38.875
7	0.4536	22	1.426	37	2.397	52	3.369	67	4.341	82	5.313	97	6.285	700	45.354
8	0.5184	23	1.490	38	2.462	53	3.434	68	4.406	83	5.378	98	6.350	800	51.833
9	0.5832	24	1.555	39	2.527	54	3.499	69	4.471	84	5.442	99	6.414	900	58.313
10	0.6480	25	1.620	40	2.592	55	3.564	70	4.535	85	5.507	100	6.479	960	62.307
11	0.7128	26	1.685	41	2.656	56	3.628	71	4.600	86	5.572	120	7.776	1000	64.792
12	0.7776	27	1.749	42	2.721	57	3.693	72	4.665	87	5.637	150	9.719		
13	0.8424	28	1.814	43	2.786	58	3.758	73	4.730	88	5.702	180	11.664		
14	0.9072	29	1.869	44	2.851	59	3.823	74	4.795	89	5.766	200	12.958		
15	0.972	30	1.944	45	2.916	60	3.888	75	4.859	90	5.831	240	15.552		

Weil's Disease (*witz*). See *Diseases, Table of*.

Weir Mitchell Treatment (*weir mit-chel*) [*S. Weir Mitchell*, an American neurologist]. The rest-cure, a treatment for certain functional nervous conditions, consisting in absolute rest in bed, with massage, electricity, and the administration of abundant food, especially milk.

Wen. A sebaceous cyst.

Werthoff's Disease [*P. G. Werthof*, a German physician]. Purpura hæmorrhagica. See *Diseases, Table of*.

Wernicke's Sign. See *Reaction, Hemipic Pupillary*.

Westphal's Nucleus. See *Nucleus*.

Westphal's Paradoxical Contraction. The contraction of a muscle, as the tibialis anti-

cus, caused by the passive approximation of its extremities, observed in cases of combined sclerosis of the posterior and lateral columns, in paralysis agitans, and other diseases.

Westphal's Sign or Symptom. The loss of the knee-jerk in locomotor ataxia.

Wet-pack. See *Pack*.

Wharton's Duct [Thomas *Wharton*, an English anatomist]. The duct of the submaxillary gland.

Wharton's Jelly, Whartonian Jelly. The mucoid tissue of the umbilical cord.

Wheal (*hweel*) [AS., *hwéle*, wheal]. A whitish or pinkish elevation, developing suddenly upon the skin, and lasting usually but a short time. Wheals are produced by urticaria, the bites of insects, or the sting of a nettle. It is also called pomphus, or urtica.

Whealhouse's Operation. A variety of external urethrotomy.

Wheat (*hwét*). See *Triticum*.

Wheeze (*hwets*). To breathe hard; to breathe with difficulty and with an audible whistling sound.

Whey (*hwa*). The liquid part of milk separating from the curd. **W.-cure**, the administration of whey as a method of treating certain diseases.

Whip-worm (*hwip'-worm*). The *Trichocephalus dispar*.

Whisky, Whiskey (*hwis'-ke*) [Gael., *uisge, water*]. See *Spiritus frumenti*. **W.-nose**. See *Acne rosacea*.

Whisper (*hwis'-per*). A low, soft, sibilant sound produced by the passage of the breath through the glottis without throwing the vocal cords into vibration.

Whispering Pectoriloquy. See *Pectoriloquy*.

White (*hwit*) [AS., *hwit*, white]. 1. Having a color produced by reflection of all the rays of the spectrum; opposed to black. 2. Any white substance, as W. of egg. **W. Cell, W. Corpuscle**, the leukocyte. **W. Commissure**, the anterior commissure of the spinal cord. **W. Lead**, basic lead carbonate. **W. Leg**. See *Phlegmasia alba dolens*. **W. Leprosy**, vitiligo. **W. Line**, the linea alba. **W. Matter**, the part of the brain and spinal cord consisting of medullated nerve-fibers, and having a white color. **W. Mustard**, sinapis alba. See *Mustard*. **W., Pearl-**, bismuth subnitrate. **W. Pneumonia**, pneumonia alba. **W. Precipitate**, hydrargyrum ammoniatum. See *Mercury*. **W. Softening**, softening of nerve-substance in which the affected area presents a whitish color, due to fatty degeneration following anemia. **W. Substance of Schwann**, the myelin-sheath of medullated nerve-fibers. **W. Swelling**, tuberculous arthritis. **W. Vitriol**, zinc sulphate.

Whites (*hwite*). See *Leukorrhœa*.

Whitlow (*hwit'-lo*). See *Panaris*. **W., Painless**. See *Morvan's Disease*, in *Diseases*, *Table of*.

Whooping-cough (*hoo'-ping-kawf*). An infectious disease characterized by catarrhal inflammation of the air-passages and peculiar paroxysms of cough ending in a loud whooping inspiration. It is most frequent in children, and is probably due to a specific micro-organism. The disease begins with symptoms of coryza and bronchitis, which in from one to two weeks are followed by the so-called paroxysmal stage, characterized by a violent cough, at the close of which occurs a loud whoop and often vomiting. This stage lasts from three to four weeks and is succeeded by the stage of decline. Complications, such as catarrhal pneumonia, emphysema, atelectasis, hernia, cerebral hemorrhage, and others, may occur.

Wickersheimer's Fluid (*wik'-ers-hi-mers*). A fluid employed for the preservation of anatomic specimens, consisting of arsenious acid, sodium chlorid, potassium sulphate, carbonate, and nitrate, dissolved in a mixture of glycerol, methyl-alcohol, and water.

Wilde's Incision [*Wilde*, an Irish otologist]. See *Operations*, *Table of*.

Williams' Tracheal Tone. The peculiar resonance sometimes found in the second intercostal space in cases of very large pleural effusion. It is a dull tympanitic resonance, becoming higher on opening the mouth, and arising from the vibrations of air in a large bronchus surrounded by compressed lung.

Willis, Circle of [*T. Willis*, an English anatomist]. Fibrous bands which extend transversely across the inferior angle of the superior longitudinal sinus.

Willis, Circle of. See *Circle*.

Willow (*wil'-o*) [AS., *welig*, willow]. *Salix*, a genus of trees of the order Salicaceæ, the bark and leaves of which contain salicin. *Salix alba* and *Salix nigra* have been used in spermatorrhea, neuralgia, and malaria.

Winckel's Disease. An epidemic disease of children, characterized by cyanosis, jaundice, and hemoglobinuria. See *Diseases*, *Table of*.

Windage (*win'-dej*). The compression of air said to be produced by the passage of a ball close to the body, and to give rise to an injury called wind-contusion.

Windlass, Spanish. A form of tourniquet consisting of a handkerchief tied about a part and twisted by means of a stick.

Window (*win'-do*). 1. An aperture in a wall for the admission of light and air. 2. A small aperture in a bone. See *Fenestra*.

Wine (*win*) [*vinum*]. 1. The fermented

juice of the grape. **W.**, **Red** (*Vinum rubrum*, U. S. P.), is the fermented juice of fresh colored grapes. The most important varieties are claret or Bordeaux and port. **W.**, **White** (*Vinum album*, U. S. P.), is the fermented juice of grapes freed from seeds, stems, and skins. The most important varieties are sherry (*Vinum xericum*), Madeira, Catawba, etc. Wines consist chiefly of water and alcohol, the latter varying from 6 to 22 per cent. (from 10 to 14 per cent. in the official wines). In addition, they contain volatile oil, enanthic ether, grape-sugar, traces of glycerol, coloring matter, tannic, malic, phosphoric, carbonic, and acetic acids, potassium bitartrate, and calcium tartrate. Wine is used as a beverage in most civilized countries, and in medicine as a stimulant like alcohol. 2. A solution of a medicinal substance in wine.

Wineglass. A measure holding nearly two fluidounces.

Wing. See *Ala*.

Winogradsky, Nitrifying Bacillus of. See *Bacteria, Table of*.

Winslow, Foramen of [J. B. Winslow, a Danish anatomist]. See *Foramina, Table of*.

Winslow, Ligament of. The posterior ligament of the knee-joint.

Wintergreen. See *Gaultheria*.

Winter-itch. See *Pruritus hiemalis*.

Wintrich's Sign. The change in the pitch of the percussion-note over a pulmonary cavity when the mouth is open or closed.

Wirsung, Canal of [*wër'-soong*]. The excretory duct of the pancreas.

Wisdom-tooth. The last molar tooth, which is the last of all the teeth to appear.

Witch-hazel. See *Hamamelis*.

Wolfian (*wool'-fe-an*). Described by C. F. Wolff, a German anatomist. **W. Body**, the mesonephros, or primordial kidney, the central portion of the segmental organs of the embryo. Its excretory duct is the **W. Duct**.

Wood. The hard part of trees; the part within the bark. **W.-alcohol**, methyl-alcohol. **W.-oil.** See *Gurjun Balsam*. **W.-spirit**, methyl-alcohol. **W.-vinegar**, vinegar obtained by the dry distillation of wood.

Wooden Tongue. See *Actinomyces*.

Wool-fat. See *Lanolin*. **Wool-sorters' Disease**, anthrax.

Woorara (*woo-rar'-ah*). See *Curare*.

Word-blindness. See *Aphasia*. **W.-deafness.** See *Deafness*.

Working Distance. In a microscope, the distance between the object and the objective.

Worm (*wurm*). 1. A member of the class Vermes, of the division Invertebrata, especially one parasitic in man or animals. **W.**, **Bladder-**, the *Tænia echinococcus*. **W.**, **Guinea-**, **W.**, **Medina-**, the *Filaria medi-nensis*. See *Filaria*. **W.**, **Pin-**, **W.**, **Seat-**. See *Oxyuris*. **W.**, **Thread-**. See *Oxyuris* and *Trichocephalus*. **W.**, **Whip-**. See *Trichocephalus*.

Wormian Bone (*wurm'-me-an*) [Olaus Worm, a Danish physician, 1588-1654]. Any one of the small supernumerary bones found in the sutures of the skull.

Wormseed (*wurm'-sed*). 1. See *Chenopodium*. 2. See *Santonica*.

Wormwood (*wurm'-wood*). See *Absinthum*.

Worsted Test. See *Holmgren's Test*.

Woulfe's Bottles. An apparatus consisting of a series of three-necked bottles connected by suitable tubes and used for washing gases or saturating liquids therewith.

Wound (*woond*) [AS., *wund*, wound]. A solution of continuity of an external or internal surface of the body. **W.**, **Contused**, one produced by a blunt body. **W.**, **Incised**, one caused by a cutting instrument. **W.**, **Lacerated**, one in which the tissues are torn. **W.**, **Open**, one having a free external opening. **W.**, **Poisoned**, one in which septic materials are introduced. **W.**, **Punctured**, one made by a pointed instrument. **W.**, **Subcutaneous**, one with a very small external opening in the skin.

Wrisberg, Cartilages of [H. A. Wrisberg, a German anatomist]. See *Cartilage*.

Nerve of: 1. A small branch of the brachial plexus supplying the skin of the arm. 2. A small nerve arising from the medulla oblongata and coursing between the facial and auditory nerves. See *Nerves, Table of*.

Wrist (*rist*) [AS., *wrist*]. The part joining the fore-arm and the hand. See *Carpus*.

W.-clonus. See *Reflexes, Table of*. **W.-drop**, a paralysis of the extensor muscles of the wrist and fingers causing a dropping of the hand. **W.-joint**, the articulation between the fore-arm and the hand.

Writer's Cramp. An occupation-neurosis occurring in those who write a great deal, and characterized by painful spasm of the fingers when an effort at writing is made.

Writing Hand. A peculiar position assumed by the hand in paralysis agitans.

Wry-neck (*ri'-neck*). See *Torticollis*.

Wurali (*wu-ra'-le*). Same as *Curare*.

X

- Xanthelasma** (*zan-thel-as'-mah*) [*ξανθός*, yellow; *ελασμα*, a lamina]. See *Xanthoma*.
- Xanthic** (*zan'-thik*) [*ξανθός*, yellow]. 1. Yellow. 2. Pertaining to xanthin.
- Xanthin** (*zan'-thin*) [*ξανθός*, yellow], $C_8H_8N_4O_2$. A leukomatin found in nearly all the tissues and liquids of the animal economy, and also in many plants; it occurs in minute quantities in urine, also in guano. It is formed in the decomposition of nuclein by dilute acids. It is a colorless powder, almost insoluble in cold water, but readily soluble in dilute acids and alkalis, and acts as a muscle-stimulant, especially to the heart.
- Xanthinuria** (*zan-thin-u'-re-ah*) [*ξανθός*, yellow; *urina*, urine]. The presence of xanthin in excess in the urine.
- Xantho-** (*zan'-tho-*) [*ξανθός*, yellow]. A prefix meaning yellow.
- Xanthochromia** (*zan-tho-kro'-me-ah*) [*ξανθός*, yellow; *χρῶμα*, a color]. A yellowish discoloration of the skin.
- Xanthocreatinin** (*zan-tho-kre-at'-in-in*) [*ξανθός*, yellow; *κρέας*, flesh], $C_9H_{10}N_4O$. A leukomatin found in muscle, crystallizing in yellow crystals and resembling kreatinin. It produces depression, somnolence, fatigue, frequent defecation, and vomiting. See *Leukomains*, Table of.
- Xanthocyanopia**, **Xanthocyanopsia** (*zan-tho-si-an-o'-pe-ah*, *zan-tho-si-an-op'-se-ah*) [*ξανθός*, yellow; *κύανος*, blue; *ὄψις*, sight]. A defect of color-vision in which yellow and blue are perceived, while red is imperceptible.
- Xanthoderma**, **Xanthodermia** (*zan-tho-der'-mah*, *zan-tho-der'-me-ah*) [*ξανθός*, yellow; *δέρμα*, skin]. A yellow discoloration of the skin.
- Xanthodontous** (*zan-tho-don'-tus*) [*ξανθός*, yellow; *ὀδούς*, tooth]. Having yellow teeth.
- Xanthokreatinin**. See *Xanthocreatinin*.
- Xanthoma** (*zan-tho'-mah*) [*ξανθός*, yellow; *ῥμα*, tumor]. Xanthelasma; a new growth of the skin occurring as flat or slightly raised patches, or nodules from a pin's head to a bean in size, and of a yellowish color. The flat lesions (*X. planum*) usually occur about the eyelids; the elevated or tubercular variety (*X. tuberculatum*, *X. tuberosum*) on the neck, trunk, and extremities. Histologically the lesions consist of connective tissue undergoing a partial fatty degeneration. **X. diabetorum**, a rare disease of the skin associated with diabetes mellitus, the lesions of which are denser and firmer than those of true xanthoma, and are dull-red, discrete, and solid, with a yellowish point at the apex.
- Xanthopathy** (*zan-thop'-ath-e*) [*ξανθός*, yellow; *πάθος*, a suffering]. See *Xanthoderma*.
- Xanthophane** (*zan'-tho-fan*) [*ξανθός*, yellow; *φαίνειν*, to show]. A yellow pigment found in the retinal cones.
- Xanthoproteic** (*zan-tho-pro'-te-ik*) [*ξανθός*, yellow; *protein*]. Derived from or related to xanthoprotein. **X. Reaction**, the deep-orange color obtained by the addition of ammonia to proteids that have been heated with strong nitric acid.
- Xanthoprotein** (*zan-tho-pro'-te-in*) [*ξανθός*, yellow; *protein*]. A yellowish substance formed from proteids by the action of nitric acid.
- Xanthopsia** (*zan-thop'-se-ah*) [*ξανθός*, yellow; *ὄψις*, vision]. Yellow vision, the condition in which objects look yellow.
- Xanthoxylum** (*zan-thok'-il-um*) [*ξανθός*, yellow; *ξύλον*, wood]. Prickly ash, a genus of trees of the order Rutaceæ. The bark of *X. americanum* and *X. Clava-Herculis* (*X.*, U. S. P.) contains a crystalline principle, **xanthoxylum**, and is irritant, stimulant, and slightly diaphoretic. It is used in chronic rheumatism and as an emmenagogue. Dose gr. x-3ss (0.65-2.0). Dose of Extractum xanthoxyli fluidum, U. S. P., f3ss-j (2.0-4.0).
- Xenogenesis** (*zen-o'-jen'-es-is*) [*ξένος*, stranger; *γένεσις*, birth]. See *Heterogenesis*.
- Xero-** (*ze'-ro*) [*ξηρός*, dry]. A prefix meaning dry.
- Xeroderma**, **Xerodermia** (*ze-ro-der'-mah*, *ze-ro-der'-me-ah*) [*ξηρός*, dry; *δέρμα*, skin]. 1. An abnormal dryness of the skin. 2. A disease characterized by dryness and harshness of the skin, discoloration, and a fine scaly desquamation; by some it is considered a mild form of ichthyosis. **X. pigmentosum**, atrophoderma pigmentosum, angioma pigmentosum atrophicum, melanosis lenticularis progressiva, Kaposi's disease; a rare disease of the skin, usually beginning in childhood, and characterized by disseminated pigment-spots, telangiectasis, atrophy of muscles, and contraction of the skin, generally followed by the development of ulcers, and ending in death.
- Xeroma** (*ze-ro'-mah*). See *Xerophthalmia*.
- Xerophthalmia** (*ze-roff'-thal'-me-ah*) [*ξηρός*, dry; *ὀφθαλμός*, eye]. A dry and thickened condition of the conjunctiva, sometimes following chronic conjunctivitis or disease of the lacrimal apparatus.
- Xerosis** (*ze-ro'-sis*) [*ξηρός*, dry]. A state of dryness, especially of the skin (see *Xeroder-*

ma) or of the conjunctiva (see *Xerophthalmia*).

Xerostomia (*ze-ro-sto'-me-ah*) [*ξηρός*, dry; *στόμα*, mouth]. Dry mouth, a peculiar condition characterized by suppression of the secretion of the salivary and buccal glands.

Xerotic (*ze-ro't-ik*) [*ξηρός*, dry]. Characterized by xerosis; dry.

X-rays, or Röntgen-rays. The peculiar ether-rays or waves discovered by Prof. Röntgen of Würzburg. A vacuum-tube of glass (called a *Geissler Tube*, a *Hittorf's* or a *Crookes' Tube*) is used with two wires sealed through the glass. These wires are connected with the two poles of a battery, and Röntgen found that the rays from the cathode (*Cathode Rays*) had peculiar penetrative powers through matter opaque to other ether-rays, and by means of these rays photographs ("Shadowgrams") may be taken of bones, metallic substances, etc., situated in the tissues. Herz, and especially Lenard, prior to Röntgen, had discovered this penetrating power of the cathode rays, but failed to make the application suggested by Röntgen.

Xiphisternum (*sif-is-tur'-num*) [*ξίφος*,

sword; *στέρνον*, breast-bone]. The xiphoid cartilage.

Xiphocostal (*sif-o-kos'-tal*) [*ξίφος*, sword; *costa*, rib]. Pertaining to the xiphoid cartilages and the ribs.

Xiphodymus (*sif-od'-im-us*) [*ξίφος*, sword; *διδυμος*, double]. A double monster with united pelvic and thoracic cavities, two legs, and with an occasional rudimentary third leg.

Xiphopagus (*sif-op'-ag-us*) [*ξίφος*, sword; *πάγος*, fixed]. A monster united by the xiphoid cartilages.

Xylene (*zi'-len*) [*ξύλον*, wood]. See *Xylol*.

Xylo- (*zi'-lo-*) [*ξύλον*, wood]. A prefix meaning pertaining to or derived from wood.

Xyloidin (*si-loi'-din*) [*ξύλον*, wood; *ειδος*, like], $C_6H_5NO_7$. A white, inflammable substance obtained by the action of nitric acid upon starch or various forms of woody fiber.

Xylol (*zi'-lol*) [*ξύλον*, wood; *oleum*, oil], C_6H_{10} . Dimethyl-benzene, a volatile hydrocarbon used in microscopy as a clearing-agent and as a solvent for Canada balsam.

Xyphoid (*si'-oid*) [*ξίφος*, sword; *ειδος*, like]. Sword-shaped, ensiform.

Y

Y-ligament. The iliofemoral ligament. See *Ligaments, Table of*.

Yard. A measure used in the U. S. and in England, equal to three feet.

Yarrow (*yar'-o*). See *Achillea*.

Yawning (*yaw'-ning*) [AS., *gāntan*, to yawn]. An involuntary stretching of the muscles accompanied by a deep inspiration, occurring during the drowsy state preceding the onset of sleep.

Yaws (*yaws*). See *Frambesia*.

Yeast (*yest*) [AS., *gist*, yeast]. The name applied to various species of *Saccharomyces*. Yeast acts as a ferment producing the alcoholic fermentation. **Y.**, Beer-, **Y.**, Brewer's, the *Cerevisiæ fermentum* of the B. P., produced by the *Saccharomyces cerevisiæ*. It is used as a stimulant and locally as a poultice (Catalplasma fermenti, B. P.) and deodorant to gangrenous ulcers. **Y.-poultice**, a poultice containing yeast.

Yellow (*yell'-o*) [AS., *geolo*, yellow]. Of a color like that of gold; producing such a color. **Y. Fever**, an acute infectious disease of tropical and subtropical regions of America, and probably due to a specific organism. After a period of incubation varying from a

few hours to several days, the disease begins with a chill and pain in the head, back, and limbs. The temperature rises rapidly to from 103°-105° F., vomiting occurs, the bowels are confined, the urine scanty and albuminous. A remission follows, after which, in severe cases, the temperature rises to its original height, jaundice develops, and the vomited material becomes dark from the presence of blood (black vomit). Hemorrhages may occur from the intestinal mucous membrane. The disease is very fatal, death occurring in the typhoid state or from uremia. **Y. Precipitate**, yellow oxid of mercury. **Y.-root**. See *Hydrastis*. **Y. Spot**, the macula lutea. **Y. Wash**, a lotion of yellow oxid of mercury. **Yerba Santa** (*yar'-bah san'-tah*) [Sp., sacred herbs]. See *Eriodictyon*.

Yolk [AS., *geolera*, yolk]. 1. The nutritive part of an ovum, consisting chiefly of **Y.-cells** or **Y.-granules**. **Y.-sac**, the umbilical vesicle, the sac containing the food of the embryo. 2. The yellow portion of the egg of birds. The yolk of the hen's egg is the Vitellus of the U. S. P. (*Ovi vitellus*, B. P.), and forms a constituent of *Glyceritum vitelli* (U. S. P.).

Young's Rule. A rule of dosage in children. The dose is obtained by adding 12 to the age and dividing the result by the age, and making the quotient the denominator of a fraction the numerator of which is 1. The fraction represents the proportion of the adult dose to be given to the child.

Young-Helmholtz Theory of Color-vision. Color-vision depends upon the presence in

the retina of three different sets of fibers which respond to stimulation by a sensation of red, green, or violet respectively. All other colors are simply combinations of the three primary colors. The excitation of any one set is a matter of wave-length. The longest waves excite the red, the shortest the violet, and those of intermediate length the green fibers.

Z

Zanaloin (*sa-nal'-o-in*) [*Zanzibar*; αλόη, aloes]. The aloin derived from Zanzibar aloes.

Zea (*se'-ah*) [*zēa*, a sort of grain]. A genus of grasses. **Z. mays**, maize, Indian corn; also its stigmas (Z., U. S. P.). It has been used as a diuretic in cystitis, gonorrhea, and cardiac dropsy.

Zeiss's Glands. Sebaceous glands the ducts of which open on the free edge of the lids.

Zenker's Degeneration. A form of degeneration of muscular fibers by which they are rendered brittle, so that they readily give way if the muscle be forcibly contracted.

Zeoscope (*ze'-o-skōp*) [*zēiv*, to boil; σκοπεῖν, to view]. An apparatus for determining the alcoholic strength of a liquid by means of its boiling point.

Zinc. A bluish-white metal having a specific gravity of from 6.9 to 7.2, an atomic weight of 65.1, and a valence of 2. Symbol, Zn. In nature it occurs in two principal forms, as a sulphid, called *blende*, and as a carbonate and silicate, termed *calamine*. When melted and poured into water it becomes granular (Zincum granulatum, B. P.). Zinc is used to prepare zinc sulphate and zinc chlorid, and for generating hydrogen. The compounds of zinc are poisonous, and the slow ingestion of it produces a chronic intoxication resembling, but less severe than, that produced by lead. **Z. acetate** (Zinci acetas, U. S. P., B. P.), $\text{Zn}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 2\text{H}_2\text{O}$, is used locally in ophthalmia and gonorrhea, in solutions of from 1-2 grains to the fluidounce (0.065-0.13 to 30 c.c.) of water. **Z. bromid** (Zinci bromidum, U. S. P.), ZnBr_2 , has been used in epilepsy. Dose gr. j-ij (0.061-0.13). **Z. carbonate** (Zinci carbonas precipitatus, U. S. P.; Zinci carbonas, B. P.), $(\text{ZnCO}_3)_2 \cdot 3\text{Zn}(\text{HO})_2$, is used, generally in the form of prepared calamine, as a dusting-powder on excoriated surfaces or in the form of a cerate. **Z. chlorid** (Zinci chloridum, U. S. P., B. P.),

ZnCl_2 , is used chiefly as an escharotic in carcinoma and spreading ulcers, as an injection in gonorrhea, and as an astringent in conjunctivitis. It is also employed as a disinfectant and for preserving anatomic preparations. Liquor zinci chloridi (U. S. P., B. P.) is used as disinfectant and preservative. *Burnett's disinfecting fluid* is a solution of zinc chlorid. **Z. cyanid**, $\text{Zn}(\text{CN})_2$, has been used in chorea, epilepsy, neuralgia, gastralgia, and uterine colic. Dose gr. $\frac{1}{4}$ -j (0.016-0.065). **Z. ferrocyanid**, $\text{Zn}_2\text{Fe}(\text{CN})_6 + 3\text{H}_2\text{O}$, is used like Z. cyanid. Dose gr. j-iv (0.065-0.26). **Z. iodid** (Zinci iodidi, U. S. P.), ZnI_2 , has been used in chorea, scrofula, and hysteria, and locally as an astringent, like the chlorid. Dose gr. ss-ij (0.032-0.13). **Z. lactate**, $\text{Zn}(\text{C}_3\text{H}_5\text{O}_2)_2 \cdot 3\text{H}_2\text{O}$, has been used in epilepsy. Dose gr. ij-x (0.13-0.65). **Z. oxid** (Zinci oxidum, U. S. P., B. P.), ZnO , is an amorphous white powder, and is used internally in chorea, epilepsy, whooping-cough, and gastrointestinal catarrh; locally, as a desiccant to excoriated surfaces, in the form of powder or ointment (Unguentum zinci oxidi, U. S. P.; Unguentum zinci, B. P.). Dose gr. ij-viii (0.13-0.52). **Z. phosphid** (Zinci phosphidum, U. S. P.), Zn_3P_2 , is used in medicine as a substitute for phosphorus. Dose gr. $\frac{1}{8}$ (0.003). **Z. sulphate** (Zinci sulphas, U. S. P., B. P.), ZnSO_4 , white vitriol, is tonic, astringent, and emetic. It is used in gastric catarrh, as an emetic, and locally in ophthalmia, gonorrhea, leukorrhea, and as a caustic in ulcers, condylomata, etc. In overdoses it is a gastrointestinal irritant. Dose gr. $\frac{1}{4}$ - $\frac{1}{2}$ (0.016-0.032); as an emetic, gr. x-xxx (0.65-2.0). **Z. sulphocarbonate** (Zinci sulphocarbonas, B. P.), $\text{Zn}(\text{C}_2\text{H}_3\text{SO}_4)_2 \cdot \text{H}_2\text{O}$, is used as an antiseptic and astringent. **Z. valerianate** (Zinci valerianas, U. S. P., B. P.), $\text{Zn}(\text{C}_8\text{H}_7\text{O}_2)_2 \cdot 2\text{H}_2\text{O}$, is used in neuralgia, epilepsy, hysteria, and diabetes insipidus. Dose gr. j-ij (0.065-0.13).

Zingiber (*sin'-jib-er*) [*ζιγγίβερις*, ginger]. Ginger, a genus of plants of the Scitamineæ. The rhizome of *Z. officinale* contains a volatile oil, and is used as a stimulant and carminative in dyspepsia, flatulence, and intestinal atony; externally it is rubefacient. Dose, gr. x-xx (0.65-1.3). Preparations and doses: Extractum zingiberis fluidum (U. S. P.), \mathfrak{m} x-xx (0.65-1.3); Oleoresina zingiberis (U. S. P.), \mathfrak{m} j (0.065); Pulvis aromaticus (U. S. P., Pulvis cinnamomi compositus, B. P.), gr. x-xxx (0.65-2.0); Pulvis rhei compositus (U. S. P., B. P.), \mathfrak{z} ss- \mathfrak{z} j (2.0-4.0); Syrupus zingiberis (U. S. P., B. P.), \mathfrak{f} \mathfrak{z} j (4.0); Tinctura zingiberis (U. S. P., B. P.), \mathfrak{z} ss-j (2.0-4.0); Tinctura zingiberis fortior (B. P.), \mathfrak{m} v-xx (0.32-1.3); Trochisci zingiberis (U. S. P.), each containing \mathfrak{m} ij of the tincture.

Zinn, Ligament of. See *Ligament*.

Zinn, Zonule of. See *Zonule*.

Zn. The symbol for zinc.

Zoanthropy (*zo-an'-thro-pe*) [*ζῶον*, animal; *ἀνθρωπος*, a man]. A form of insanity in which the person imagines himself transformed into or inhabited by an animal.

Zoetrope (*zo'-e-tro-pe*) [*ζῶη*, life; *τροπέω*, to turn]. An instrument that by the revolution of a series of pictures before the eye gives the appearance of a person or animal in motion, the pictures successively representing single phases of a compound act.

Zona (*zo'-nah*) [L.]. 1. A belt or girdle. *Z. arcuata*, the inner zone of the basilar membrane, extending from the lower edge of the spiral groove of the cochlea to the external edge of the base of the outer rods of Corti. *Z. cartilaginea*, the limbus of the spiral lamina. *Z. ciliaris*, the ciliary processes collectively. *Z. denticulata*, the inner zone of the basilar membrane, together with the limbus of the spiral lamina. *Z. fasciculata*, the central portion of the cortex of the suprarenal capsule, composed of tube-like transverse bands. *Z. glomerulosa*, a part of the cortical portion of the suprarenal capsule, having a net-like appearance on section, situated near the surface of the organ. *Z. orbicularis*, a thickening of the capsular ligament around the acetabulum. *Z. pectinata*, the outer portion of the basilar membrane, extending from the rods of Corti to the spiral ligament. *Z. pellucida*, the thick, solid, elastic envelop of the ovum, corresponding to the cell-wall of a cell; it is also called the vitelline membrane. *Z. tecta*, the inner portion of the basilar membrane, bearing the organ of Corti. 2. See *Herpes zoster*. *Z. ophthalmica*. Herpes zoster of the ophthalmic nerve.

Zonal (*zo'-nal*) [*zona*, zone]. Pertaining to a girdle or to a band-like structure.

Zone (*zōn*) [*zona*, band]. See *Zona*. *Z.*, Cornu-radicular, the external part of Burdach's column abutting on the middle third of the internal border of the posterior horn, and representing approximately the posterior root-zone. *Z.*, Hysterogenous, a region, as the ovarian or submammary region, where pressure in hysteric women calls forth an hysteric attack.

Zoneesthesia (*zon-es-the'-ze-ah*) [*ζώνη*, a belt; *αἰσθησις*, sensation]. See *Girdle-sensation*.

Zonular (*zon'-u-lar*) [*zonula*, a little zone]. Pertaining to or in the shape of a zone or band, as, e. g., *Z. cataract*.

Zonule (*zon'-ul*) [*zonula*, a little zone]. A small band. *Z. of Zinn*, the suspensory ligament of the crystalline lens of the eye.

Zoo- (*zo'-o*) [*ζῶον*, an animal]. A prefix meaning animal, or pertaining to an animal.

Zoobiology (*zo-o-bi-ol'-o-gy*) [*ζῶον*, animal; *βίος*, life; *λόγος*, science]. Animal biology.

Zoochemia, Zoochemistry (*zo-o-ke'-me-ah*, *zo-o-kem'-is-tre*) [*ζῶον*, animal; *χημεία*, chemistry]. The chemistry of animal life and tissues.

Zoodynamics (*zo-o-di-nam'-iks*) [*ζῶον*, animal; *δύναμις*, power]. Animal physiology.

Zoogenesis (*zo-o-jen'-es-is*) [*ζῶον*, animal; *γεννᾶν*, to beget]. The generation of animal forms.

Zoogenous (*zo-oj'-en-us*) [*ζῶον*, animal; *γεννᾶν*, to beget]. Developed or derived from animals.

Zooglea (*zo-og'-le-ah*) [*ζῶον*, animal; *γλοιός*, a sticky substance]. A stage in the life-history of certain bacteria, in which they lie embedded in a gelatinous matrix.

Zoograft (*zo'-o-graft*) [*ζῶον*, animal; *graft*]. A graft of tissue derived from an animal.

Zoografting (*zo-o-graf'-ting*) [*ζῶον*, animal; *graft*]. Grafting with tissue taken from the lower animals.

Zooid (*zo'-oid*) [*ζῶον*, animal; *εἶδος*, like]. 1. Resembling an animal. 2. Any one of a series of more or less independent forms resulting from the development of a single ovum.

Zoology (*zo-ol'-o-jē*) [*ζῶον*, animal; *λόγος*, a treatise]. That branch of biology treating of the form, nature, and classification of animals.

Zoonomia, Zoonomy (*zo-o-no'-me-ah*, *zo-on'-o-me*) [*ζῶον*, animal; *νόμος*, law]. The principles or laws of animal life; zoobiology.

Zooparasite (*zo-o-par'-as-it*) [*ζῶον*, animal; *παράσιτος*, parasite]. An animal parasite.

Zoopathology (*zo-o-path-ol'-o-jē*) [*ζῶον*, animal; *πάθος*, suffering; *λόγος*, science]. The science of the diseases of animals.

Zoophagous (*zo-off'-og-us*) [*ζῶον*, animal; *φαγεῖν*, to eat]. Subsisting on animal food.

Zoophobia (*zo-o-fō'-be-ah*) [*ζῶον*, animal; *φόβος*, fear]. Morbid dread of certain animals.

Zoosperm (zo'-o-sperm) [ζῷον, animal; σπέρμα, seed]. See *Spermatozoön*.

Zoospore (zo'-o-spör) [ζῷον, animal; σπόρος, seed]. A motile spore.

Zootomist (zo'-ot'-o-mist) [ζῷον, animal; τομή, a cutting]. One who dissects animals; a comparative anatomist.

Zootomy (zo'-ot'-o-me) [ζῷον, animal; τομή, a cutting]. The dissection of animals.

Zoster (zos'-ter) [ζωστήρ, a girdle]. See *Herpes zoster*.

Zygapophysis (zig-ap-off'-is-is) [ζυγόν, yoke; apophysis]. The articular process of a vertebra.

Zygoma (zi-go'-mah) [ζύγωμα, the cheek-bone]. The arch formed by the union of the zygomatic process of the temporal bone and the malar bone.

Zygomatic (zi-go-mat'-ik) [ζύγωμα, zygoma]. Pertaining to the zygoma.

Zygomaticus (zi-go-mat'-ik-us) [ζύγωμα, zygoma]. One of several small subcutaneous muscles arising from or in relation with the zygoma. See *Muscles*, Table of.

Zymase (zi'-mäs). See *Microzyme*.

Zyme, Zymin (zim, zi'-min) [ζύμη, leaven]. An organized ferment. Cf. *Enzyme*.

Zymic (zi'-mik) [ζύμη, leaven]. Of or pertaining to organized ferments.

Zymo- (zi'-mo) [ζύμη, leaven]. A prefix

meaning pertaining to or produced by fermentation.

Zymogen (zi'-mo-jen) [ζύμη, leaven; γεννᾶν, to produce]. The substance existing in the glands secreting a digestive juice, and which, when set free, splits into a ferment (enzyme) and a proteid.

Zymogenic (zi-mo-jen'-ik) [ζύμη, leaven; γεννᾶν, to produce]. 1. Causing fermentation. 2. Pertaining to or producing a zymogen.

Zymology (zi-mol'-o-jē) [ζύμη, leaven; λόγος, science]. The science dealing with fermentation.

Zymometer (zi-mom'-et-er) [ζύμη, leaven; μέτρον, measure]. An instrument for measuring fermentation.

Zymophyte (zi'-mo-fit) [ζύμη, leaven; φυτόν, plant]. A microorganism producing fermentation.

Zymose (zi'-mids) [ζύμη, leaven]. See *Enzyme*.

Zymosimeter (zi-mo-sim'-et-er). See *Zymometer*.

Zymosis (zi-mo'-sis) [ζύμη, leaven]. 1. Fermentation. 2. The condition of one affected with a zymotic disease. 3. An infectious disease.

Zymotic (zi-mot'-ik) [ζύμη, leaven]. Pertaining to zymosis; produced by zymosis.

Z. Disease, an infectious disease.

APPENDIX A.

MINERAL SPRINGS OF THE UNITED STATES.

BY JUDSON DALAND, M. D.,
Of Philadelphia.

The writer has given considerable thought to the question of classification of mineral waters, and after a thorough study of those hitherto brought forward, is strongly of the opinion that the heading under which a spring is placed should indicate, as near as may be, the chief therapeutic action of that spring.

While a purely chemical classification, such as the excellent one suggested by Dr. A. C. Peale, of the U. S. Geological Survey,* is scientifically more accurate, it is feared that the practical physician would be confused when he came to make his selection, by the large number of subdivisions necessitated by this plan of division.

Though realizing fully the imperfection of the present classification, it has seemed that the needs of those consulting this work would be best subserved by its adoption.

Occasionally a spring possesses the marked characteristics of more than one class. Such springs are entered under both groups.

All analyses have been reduced to the U. S. gallon of 231 cubic inches, so as to simplify the comparative study of the value of the different waters, and for the sake of brevity, the principal ingredients are represented by their chemical formulæ as follows:—

Lithium bicarbonate, . . . $\text{Li}(\text{CO}_3)_2$,	Ferrum bicarbonate, . . . $\text{Fe}(\text{HCO}_3)_2$,
Ammonium bicarbonate, . NH_4HCO_3 ,	carbonate, FeCO_3 ,
Sodium biborate, $2\text{B}_2\text{O}_3 + 10\text{H}_2\text{O}$,	sulphate, FeSO_4 ,
carbonate, Na_2CO_3 ,	oxide, FeO ,
bicarbonate, . . . Na_2HCO_3 ,	Calcium sulphate, CaSO_4 ,
chloride, NaCl ,	bicarbonate, . . . $\text{Ca}(\text{HCO}_3)_2$,
sulphate, Na_2SO_4 ,	carbonate, CaCO_3 ,
Magnesium carbonate, . . MgCO_3 ,	chloride, CaCl_2 ,
bicarbonate, . $\text{Mg}(\text{HCO}_3)_2$,	phosphate, . . . $\text{Ca}_3(\text{PO}_4)_2$,
chloride, . . MgCl_2 ,	H_2S . Sulphuretted hydrogen gas,
sulphate, . . MgSO_4 .	CO_2 . Carbonic acid gas.

I. DIURETIC WATERS.

These springs owe their action chiefly to the increased quantity of water taken by patients, who thus flush out all the emunctories, especially stimulating the flow of urine.

* In a paper read before the American Climatological Association.

In practical medicine, waters of this class are used very extensively, and justly so. They are of undoubted value in many conditions. It is needless to dwell upon the great benefit to be derived from the increased use of ordinary, pure drinking water, as this is universally recognized, but in practice it is found that patients take these diuretic waters more willingly, in larger amounts and for a longer time than the ordinary drinking water, of which they soon become satiated. Ordinarily it is best to take a large tumbler of this water before breakfast, between meals and at bed-time, thus using two to three pints in the twenty-four hours. I may be permitted to mention a few of the ordinary conditions in which these waters produce their best results; and as first in importance are the diatheses, lithamic, gouty, rheumatic, oxalic, phosphatic and uric acid. Good results have also been obtained in renal calculus, acute and chronic congestion of the kidneys, acute nephritis and at times in the course of chronic Bright's disease and diabetes. As the use of pure water, in large amounts, produces an increase in the amount of bile and at the same time renders it more thin, as was proven several years ago by Bidder and Schmidt, these waters are used with excellent result in acute and chronic hepatic congestion, catarrhal inflammation of the mucous membrane lining the biliary passages, known as catarrhal jaundice and gall stone, both during the attack and in the interval.

I append a few analyses in full of typical springs belonging to this group, followed by a full list of springs of this class, with amount of each *active* ingredient in a U. S. gallon of 231 cubic inches:—

SWEET SPRINGS, BEDFORD, PENNSYLVANIA.

(Analyst, F. A. Genth.)

One U. S. gallon (231 cu. in.) contains—
 Carbonate of calcium,0.522 grs.
 " magnesium,0.135 "
 Silicic acid,0.654 "

1.311 "

Free carbonic acid .66 cu. in. Temp.
 52.7° F.

SUMMIT MINERAL SPRING, HARRISON, CUMBERLAND COUNTY, MAINE.

(Analyst, F. L. Bartlett.)

One U. S. gallon (231 cu. in.) contains—
 Carbonate of sodium and potassium,1.4019 grs.
 Chloride of sodium,1713 "
 Carbonate of calcium,9904 "
 Carbonate of magnesium,2476 "
 Oxide of iron and aluminium, distinct trace.
 Silicium,9800 grs.
 Silicic acid (in solution).
 Organic and volatile matter,2375 "

Temperature, 46° F. Total, . 4.0287 "

POLAND SPRING, SOUTH POLAND, MAINE.

(Analyst, C. F. Chandler.)

One U. S. gallon (231 cu. in.) contains—
 Sulphate of potassa,0.1562 grs.
 Chloride of sodium,0.2636 "
 Carbonate of sodium,0.1333 "
 " calcium,1.2287 "
 " magnesium,0.5412 "

Oxide of iron and alumina, traces.
 Silica,1.1197 "
 Organic and volatile matter,0.2332 "

Total,3.6759 "

BERKELEY SPRINGS, MORGAN COUNTY, WEST VIRGINIA.

(Analyst, A. A. Hayes.)

One U. S. gallon (231 cu. in.) contains—
 Calcium carbonate,5.000 grs.
 Sodium chloride,896 "
 Calcium chloride,208 "
 Magnesium sulphate,360 "
 Calcium crenate,3.640 "
 Ferrous crenate,080 "
 " silicate,640 "
 Loss,064 "

Temperature, 74° F. Total, . 10.888 "

CAPON SPRINGS, HAMPSHIRE COUNTY,
WEST VIRGINIA.

(Analyst, J. W. Mallet.)

One U. S. gallon (231 cu. in.) contains—	
Sulphate of calcium,477 grs.
“ potassium,136 “
Crenate of iron,080 “
Chloride of sodium,240 “
Carbonate of calcium,	6.701 “
“ sodium,475 “
“ magnesium,160 “
“ iron,033 “
Alumina,014 “
Organic matter,164 “
	<hr/>
	9.774 “
Carbonic acid,	6.898 “
Nitrogen,	2.962 “
Oxygen,	1.416 “
Temperature, 65.50° F.	

CRESSON MAGNESIA SPRING, CAMBRIA
COUNTY, PENNSYLVANIA.

(Analyst, F. A. Genth.)

One U. S. gallon (231 cu. in.) contains—	
Sulphate of calcium,	0.109 grs.
Chloride of magnesium,	0.559 “
“ calcium,	1.304 “
“ sodium,	1.229 “
Bicarbonate of iron,	0.017 “
“ manganese,	trace
“ magnesium,	0.414 “
“ calcium,	0.022 “
“ sodium,	1.425 “
“ potassium,	0.206 “
Phosphate of calcium,	0.004 “
Alumina,	0.008 “
Silicic acid,	0.914 “
Nitrous acid,	trace
Carbonic acid,	0.663 “
	<hr/>
	6.874 “

BEDFORD CHALYBEATE SPRING, BEDFORD
COUNTY, PENNSYLVANIA.

One U. S. gallon (231 cu. in.) contains—	
Sulphate of calcium,	2.741 grs.
Phosphate of calcium,033 “
Carbonate of calcium,	8.885 “
“ magnesium,	1.201 “
“ iron,442 “
“ manganese,	trace
“ sodium,395 “
“ potassium,132 “
Chloride of sodium,120 “
Silicic acid,793 “
	<hr/>
	14.942 “
Carbonic acid,	12 cu. in.

All Healing Spring, Catoosa County, Georgia. $MgCO_3$ 7, $CaSO_4$ 39.

Alum Spring, Onslow County, North Carolina.

Artesian Well, Winslow, Camden County, New Jersey.

Ashley Springs, North Carolina.

Augusta Springs, Augusta County, Virginia. See Stribling Springs.

Bath or Berkeley Springs, Morgan County, Virginia. 74° F.

Birchdale Spring, Concord, N. H.

Black Barren Spring, Lancaster County, Pennsylvania.

Bryant's Springs, Lincoln County, Kentucky. Numbers I, II and III.

- Bullfinch Springs, Mecklenburg County, Virginia. CaSO_4 and $\text{Ca}(\text{HCO}_3)_2$ 39.
 Luray Springs, 6.68 cubic inches. Temperature 60°F .
 Capon Springs, Hampshire County, West Virginia. 65.5°F .
 Chalk Creek Springs, Chalk Creek, Colorado. 130°F .
 Chalybeate Spring, Bedford Springs, Bedford County, Pennsylvania. CaCO_3 9, CaSO_4 3, FeCO_3 trace, CO_2 21 cubic inches. 52.70°F .
 Chenoweth Chalybeate Spring, Bedford County, Pennsylvania.
 Cherry Hill Phosphate Spring, Otsego County, New York. $\text{Ca}_3(\text{PO}_4)_2$ 14.
 Cherry Valley Phosphate Spring, Otsego County, New York.
 Clymerara Spring, Manor of Maryland, Maryland.
 Cola Springs, Browns Mills, New Jersey.
 Concord Spring, Concord, New Hampshire.
 Daggy Spring. See Greencastle.
 Dew Drop Spring. See Greencastle.
 Epsom Spring, in Wyandotte Cave, Indiana. 24.
 Estill Chalybeate Spring, Estill County, Georgia. O_2 41 cubic inches.
 Estill Red Sulphur Spring, Estill County, Georgia. CO_2 40 H_2S .56 cubic inches.
 Farmville Lithia Water, Farmville, Virginia. NaSO_4 3, $\text{Li}(\text{CO}_3)_2$ 3, $\text{Fe}(\text{CO}_3)_2$ 1, $\text{K}_2\text{Al}_2(\text{SO}_4)_4$ 2.
 Fayette Spring, Fayette County, Pennsylvania.
 Field Spring, Lincoln County, Kentucky.
 Florida Spring, Montgomery County, New York. $(\text{NaHCO}_3)_2$ 22.
 Glen Alpine Springs, Burke County, North Carolina.
 Greencastle Daggy Spring. CaCO_3 14.
 Greencastle Dew Drop Spring, Putnam County, Indiana. CaCO_3 12.
 Grove Spring, Lincoln County, Kentucky. CaCO_3 10.
 Hawkin's Mineral Springs, Wayne County, Indiana.
 Healing Spring (New), Bath County, Virginia. CaCO_3 18, CO_2 5 cubic inches. 88°F .
 " " (Old) " " " CaCO_3 18, CO_2 4 " " 85°F .
 Hot Springs, Garland County, Arkansas. 93° to 105°F .
 Howard Well, Lincoln County, Kentucky.
 Hubbardston Well, Iona County, Michigan. CaCO_3 17.
 Jerusalem Mineral Spring, Cannan, New Hampshire.
 Jordan Chalybeate Spring, Rockbridge County, Virginia. CO_2 6.
 Kimberling Springs, Bland County, Virginia.
 Lebanon Spring, Columbia County, New York. 73°F .
 Leslie Spring, Ingham County, Michigan. CaCO_3 18. CO_2 12 cubic inches.
 Limestone Spring, Bedford Springs, Bedford County, Pennsylvania. CaCO_3 7, CO_2 21 cubic inches. 51.8°F .
 McVittey's Saltillo Springs, Huntingdon County, Pennsylvania. $\text{Ca}(\text{HCO}_3)_2$ 10.
 Magnesia Spring (Cresson Springs), Cambria County, Pennsylvania.
 Magnesia Spring. See Versailles Spring.
 Marion Artesian Well, Grant County, Indiana.
 Marranetta Mineral Spring, Harrisonburg, Virginia. CaCO_3 15, MgCO_3 7. 50°F .
 Massanutten Spring, Rockingham County, Virginia. 59.7°F .
 Mineral Spring, ten feet from May's, Bedford County, Pennsylvania.
 Mineral Spring, at the Mound in Madison County, Indiana.

Minnequa Spring, Bradford County, Pennsylvania.
 Missioquoi Springs, Franklin County, Vermont. NaHCO_3 8. CO_2 ?
 Monitor Spring. See Versailles Spring.
 New Spring. See Healing Spring.
 Northport Mineral Spring, Northport, Maine.
 Nuckoll's Springs, Grayson County, North Carolina.
 Old Spring. See Healing Spring.
 Orkney Springs (Bear's Wallow Spring), Shenandoah County, Virginia. CO_2 5. •
 Orkney Springs (Healing Spring), Shenandoah County, Virginia. CO_2 3.
 Ourry Spring. See Versailles Spring.
 Perry Springs, Pike County, Illinois. CaCO_3 14. 48° to 50° F.
 Pine Grove Mineral Spring, Dover, Strafford County, New Hampshire. 45° F.
 Plummer Bromine Arsenic Spring, Ashe County, North Carolina.
 Poland Mineral Spring, South Poland, Maine.
 Rawley Springs, Rockingham County, Virginia.
 Red Sulphur Spring. See Estill Springs.
 Red Sulphur Springs, Monroe County, West Virginia. 54° F.
 Rock Spring, Lancaster County, Pennsylvania.
 Saltillo Spring. See McVitte's Spring.
 Santa Barbara Hot Sulphur Springs, near Santa Barbara, California. Na_2CO_3 17. 112° to 122° F.
 St. Helena White Sulphur Springs, No. 7, Napa County, California. $\text{CaCl} + \text{Na}_2\text{SO}_4$ 27. 64.5° to 97° F.
 Stone Sulphur Spring, Lincoln County, Kentucky.
 Stremmel's Gettysburg Lithia Spring, Adams County, Pennsylvania.
 Stribling's Chalybeate Spring, Augusta County, Virginia.
 Stribling's Sulphur Spring, Augusta County, Virginia.
 Sweet Springs, Bedford, Pennsylvania. Temperature 52.7° F.
 Thompson's Springs, Ashe County, North Carolina.
 Underwood Spring, Falmouth Foreside, Maine.
 Van Cleuve's Mineral Spring, Crawfordsville, Indiana. CaCO_3 10.
 Versailles Magnesia Springs, Brown County, Illinois. CO_2 24 cubic inches. 58° F.
 Versailles Monitor Springs, Brown County, Illinois.
 Versailles Ourry Springs, Brown County, Illinois.
 Virginia Arsenic, Bromine and Lithia Spring, Roanoke, Virginia. CaCO_3 6, Mg SO 43.
 Warner Spring, Albion, Michigan. $\text{Ca}(\text{HCO}_3)_2$ 17.
 White Cliff Mineral Springs, Monroe County, Tennessee. Three grains of $\text{Mg}(\text{HCO}_3)_2$.
 Yellow Springs, Green County, Ohio. 52° F.

II. IRON WATERS.

These owe their virtues to the presence of iron, which usually exists in the form of the bicarbonate of the protoxide, held in solution by an excess of carbonic acid gas. The amount of carbon dioxide is often considerable, rendering the water decidedly acidulous. In many of our mineral springs, like those at Cresson, for instance, the iron exists as ferrous sulphate, with more or less ferric sulphate from the oxidation of the former.

The ordinary indications for the use of iron are well met by the use of the waters belonging to this class.

Those springs containing sulphate of magnesium and sodium, and alkalies, in addition to iron, such as those chalybeate springs, containing alum, give excellent results in chronic gastric or intestinal catarrhal inflammation, especially when associated with diarrhoea. Thorp's Spring, Texas, or the New Almaden Vichy, of California, should be of special service in chlorosis as well as in the ordinary anæmias. The following list comprises the most important members of this group, together with several examples where the analysis is given in detail :—

IRON AND ALUM WATERS.

CHURCHILL ALUM, VIRGINIA.

One U. S. Gallon (231 cu. in.) contains—	
Sulphate of magnesium, . . .	86.064 grs.
“ “ calcium, . . .	88.832 “
“ “ potassium, . . .	2.440 “
“ “ ammonium, . . .	0.640 “
“ “ sodium, . . .	1.944 “
Chloride of sodium, . . .	4.624 “
Persulphate of the sesquioxide of iron, . . .	51.264 “
Bisulphate of the sesquioxide of iron, . . .	83.352 “
Sulphate of protoxide of iron, . . .	24.184 “
Persulphate of alumina, . . .	72.928 “
Silica, . . .	10.424 “
Phosphoric acid, . . .	trace
	<hr/> 426.696 “

BATH ALUM SPRINGS, BATH COUNTY, VA.

(Analyst, A. A. Hayes.)

One U. S. Gallon (231 cu. in.) contains—	
Chloride of sodium, . . .	
Sulphate of calcium, . . .	2.539 grs.
“ “ magnesium, . . .	1.282 “
“ “ potassium, . . .	0.258 “
Crenate of ammonium, . . .	1.776 “
Silicate of sodium, . . .	3.150 “
Protosulphate of iron, . . .	21.776 “
Alumina, . . .	12.293 “
Silica, . . .	<hr/> 43.084 “
Free carbonic acid, 8 cubic inches.	

CRESSON ALUM SPRINGS, CAMBRIA COUNTY, PENNA.

One U. S. Gallon (231 cu. in.) contains—	
Sulphate of ferric oxide, . . .	83.390 grs.
“ “ aluminium, . . .	21.205 “
“ “ ferrous oxide, . . .	16.253 “
“ “ magnesium, . . .	27.699 “
“ “ calcium, . . .	40.202 “
“ “ lithium, . . .	0.047 “
“ “ sodium, . . .	0.704 “
“ “ potassium, . . .	0.426 “
Chloride of sodium, . . .	0.023 “
Bicarbonate of iron, . . .	3.784 “
“ “ manganese, . . .	trace
“ “ calcium, . . .	
Phosphate of calcium, . . .	trace
Silicic acid, . . .	1.868 “
	<hr/> 145.565 “

SIMPLE IRON WATER.

SHARON CHALYBEATE SPRING, SCHOHARIE
COUNTY, N. Y.

One U. S. Gallon (231 cu. in.) contains—

Sulphate of magnesium, 20.56 grs.
“ “ calcium, 16.27 “
“ “ sodium, 12.35 “
“ “ iron, 24.00 “
	<u>73.18 “</u>

Sulphuretted hydrogen, 1 cubic inch.
Temperature, 48° F.

SCHUYLER CHALYBEATE SPRING, SCHUY-
LER COUNTY, ILL.

(Analyst, Dr. Blaney.)

One U. S. Gallon (231 cu. in.) contains—

	Grains.	Grains.
Sulphate of calcium,	. . . 73.936	73.986
“ “ magnesium,	. . . 2.984	2.982
“ “ iron,	. . . 69.960	69.959
Alkaline sulphate,	. . . 7.832	
“ phosphate,	. . .	7.836
Silica, 1.312	1.315
	<u>156.024</u>	<u>156.028</u>

IRON AND APERIENTS.

NEW ALMADEN VICHY, SANTA CLARA
COUNTY, CALIFORNIA.

One U. S. Gallon (231 cu. in.) contains—

Sodium bicarbonate, 201 grs.
“ chloride, 33 “
Calcium sulphate,	} 74 “
“ carbonate,	}
Magnesium sulphate, 12 “
Ferri carbonate, 5 “
	<u>325 “</u>

Free carbonic acid gas, 238 cu. in.

THORP'S SPRING, HOOD COUNTY, TEXAS.

One U. S. Gallon (231 cu. in.) contains—

Ferri carbonate, 40 grs.
Sodium sulphate, 80 “
“ carbonate, 100 “
	<u>220 “</u>

Adirondack Spring, Whitehall, New York. CaCO_3 15, MgCO_3 13, FeCO_3 4, CO_2 54.15 cubic inches. 52° F.Bath Alum Spring, Bath County, Va. No. 1. Alum 9, CaFeO 14, CO_2 11 cu. in. No. 2. FeO 21, Alum 12 CO_2 8 cu. in.Bedford Alum Springs, Bedford County, Virginia. FeSO_4 23½, MgSO_4 13, CaSO_4 19. Catter Spring. See Rock Enon Spring.Cave Spring, Bath County, Virginia. FeCO_3 2.Church Hill Alum Springs, Near Richmond, Va. $\text{Fe}_2(\text{SO}_4)_3$ 135, FeSO_4 24, MgSO_4 86, CaSO_4 89, $\text{Al}_2(\text{SO}_4)_3$ 73.Columbia Springs, Columbia County, N. Y. NaCl 84, CaSO_4 65, MgCl_2 31, CaCl_2 22, FeCl_3 3½, H_2S 4.48 cubic inches.Congress Spring, Santa Clara Co., California. NaCO_3 123, NaCl 118, FeCO_3 13.Crab Orchard Acid Spring, Genesee County, New York. H_2SO_4 83, CaSO_4 40, FeSO_4 14. Cresson Springs, Cambria County, Pennsylvania.Iron Spring. FeSO_4 and $\text{Fe}(\text{HCO}_3)_2$ 29, CaSO_4 49, MgSO_4 23.Alum Spring. FeSO_4 and $\text{Fe}(\text{HCO}_3)_2$ 53, CaSO_4 40, MgSO_4 28, and $\text{Al}_2(\text{SO}_4)_3$ 21 grains.Fauquier White Sulphur Springs, Fauquier County, West Virginia. FeSO_4 2. 55° F.Geyser Spa Spring, Sonoma County, California. NaHCO_3 23, $\text{Mg}(\text{HCO}_3)_2$ 10, FeCO_3 4.Iron Ute Manitou Spring, Colorado. CaCO_3 70, Na_2CO_3 70, FeCO_3 7, CO_2 ? 44° F.

- Jordan Rockbridge Alum, Rockbridge County, Virginia. FeSO_4 18½, CO_2 6 cu. in.
- Kittanning Mineral Spring, Armstrong County, Pennsylvania. FeSO_4 24½, MgSO_4 27, CaSO_4 65.
- Linwood Spring, Linwood, Iowa. FeCO_3 27, NaCl 93, NaHCO_3 40, MgCl_2 23.
- Manitou Spring. See Iron Ute.
- Napa Soda Spring, Napa County, California. FeCO_3 16, MgCO_3 52. 68° F.
- Napa Soda Spring, Napa County, California. MgCO_3 26, NaHCO_3 13, $\text{Fe}_2(\text{CO}_3)_3$ 8. 68° F.
- New Almaden Vichy, Santa Clara County, California. NaHCO_3 201, NaCl 33, CaSO_4 and $\text{Ca}(\text{HCO}_3)_2$ 73, MgSO_4 12, FeCO_3 5, CO_2 238 cubic inches.
- New London Alum Springs, Campbell County, Virginia. FeSO_4 23½, MgSO_4 13, CaSO_4 19.
- Oak Orchard Spring, Genesee County, N. Y. H_2SO_4 133, FeSO_4 33, CaSO_4 13.
- Ocean Spring, Jackson County, Mississippi. FeO 5, NaCl 48.
- Ojo Caliente Spring, Ojo Caliente, New Mexico. Na_2CO_3 115, NaCl 32, FeCO_3 6. 100° F.
- Owasso, Chalybeate, Owasso, Shiawassee County, Michigan. FeCO_3 12, MgCO_3 11.
- Pacific Congress Spring, ten miles S. W. Santa Clara, California. FeCO_3 14, NaCl and Na_2CO_3 242. 50° F.
- Pacific Congress Spring, near Santa Clara, California. NaCl 119, Na_2CO_3 123, FeCO_3 14. 50° F.
- Ranch Spring, Estes Park, Colorado. $\text{Fe}(\text{HCO}_3)_2$ 2. 58° F.
- Richfield Iron Spring, Otsego County, New York. CaSO_4 and $\text{Ca}(\text{HCO}_3)_2$ 17, $\text{Mg}(\text{HCO}_3)_2$ 12, FeO 5, CO_2 16. 46° F.
- River Spring, Estes Park, Colorado. $\text{Fe}(\text{HCO}_3)_2$ 4. 58° F.
- Rockbridge Alum Spring, Rockbridge County, Virginia. FeSO_4 5, Alumina 18.
- Rock Enon Spring, Frederick County, Va. FeCO_3 14, MgSO_4 13.
- Saline Chalybeate Spring, Schuyler County, Illinois. $\text{Fe}_2(\text{SO}_4)_3$ 70, CaSO_4 74.
- Schuyler County Springs, Schuyler County, Illinois. CaSO_4 74, FeSO_4 70.
- Sharon Chalybeate Spring, Schoharie County, New York. FeSO_4 24, MgSO_4 21. 48° F.
- Sparta Artesian Well, Sparta, Monroe County, Wisconsin. FeCO_3 9 grains.
- Stribling Springs, Augusta County, Virginia. $\text{K}_2\text{Al}_2(\text{SO}_4)_6$ 16, FeSO_4 9, CaSO_4 12.
- Stryker's Mineral Water, Stryker, Ohio. K_2SO_4 185, MgCl_2 119, NaCl 232, FeCO_3 10.
- Tar Spring, Crawford County, Indiana. FeCO_3 4, CaCO_3 21.
- Thorp's Spring, Hood County, Texas. FeCO_3 40, $\text{Na}_2\text{S}_2\text{O}_3$ 80, Na_2CO_3 100.
- Variety Springs, Augusta County, Virginia. FeSO_4 5 (NH_4) $_2\text{SO}_4$ 12, CaSO_4 13.

III. CARBONIC ACID WATERS.

These owe their chief virtue to the presence of carbonic acid gas, though many of them contain alkalies and alkaline earths in sufficient quantity to give them a special action.

Though springs of this class are very numerous, it is remarkable to observe how frequently information as to the exact amount of free carbonic gas is wanting, and that has been noted where the analysis, as regards the mineral constituents, has been conducted with great care and accuracy. Then, too, it would appear, in many instances, the water is sent some distance for analysis, so that considerable escapes and the water at the spring actually contains a larger quantity of the gas than is shown by the analysis. Many springs containing free carbonic acid gas also contain such large quantities of active inorganic

substances as to compel us to classify them under some one of the other groups. The waters belonging to this class are peculiarly agreeable both to the palate and to the eye. Their bright, sparkling appearance constantly tempts one to their use, and thus patients receive large quantities of water almost without effort. Fever patients find these waters very agreeable, and the addition of this gas acts as a sedative to the gastro-mucous membrane, and when taken cold and in sips relieves nausea and tends to check vomiting. This water when added to milk is admirably received by some patients with irritable stomach, and occasionally milk will be accepted in this form when it is absolutely refused in its pure state. Another very popular use to which these waters are put is in diluting wines.

Wherever a water surcharged with the gas is indicated, these springs are of especial value, particularly those containing a moderate amount of the alkalies. The following analyses are offered as examples of this group, to which belong the well-known Apollinaris, Giesshübel and Vichy springs:—

SWEET SPRINGS, WEST VIRGINIA.

(Analyst, W. B. Rogers.)

One U. S. gallon (231 cu. in.) contains—

Chloride of sodium,	0.136	grs
“ “ magnesium,	0.312	“
“ “ calcium,	0.144	“
Sulphate of sodium,	6.344	“
“ “ magnesium,	10.392	“
“ “ calcium,	13.168	“
Carbonate of sodium,		“
“ “ calcium,	30.556	“
“ “ magnesium,	0.824	“
Oxide of iron,	0.144	“
Silicic acid,	0.021	“
Iodine,	trace	
Earthy phosphates,	trace	
	<u>62.041</u>	

Free carbonic acid, 88 cubic inches.

GISSHÜBLER (NEAR CARLSBAD), BOHEMIA.

(One U. S. gallon (231 cu. in.) contains—

Sodium carbonate,	56.768	grs.
Calcium “	12.672	“
Magnesium “	2.720	“
Ferrous “	0.032	“
Potassium “	5.248	“
“ sulphate,	1.776	“
“ chloride,	3.008	“
Silica,	5.248	“
Alumina,	0.136	“
	<u>87.936</u>	

APOLLINARIS SPRING, NEUENAUH, RHENISH PRUSSIA.

(Analyst, Mohr.)

One U. S. gallon (231 cu. in.) contains—

Chloride of sodium,	20.79	grs.
Sulphate of soda,	12.2	“
Carbonate of soda,	55.07	“
“ “ lime,	14.6	“
“ “ magnesia,	23.93	“
Oxide of iron,	0.22	“
Silicic acid,	1.08	“
	<u>127.89</u>	

Carbonic acid, 376 cubic inches.

CLYSMIC NATURAL SPRING WATER, WAU-

KESHA, WISCONSIN.

(Analyst, Rathbene.)

One U. S. gallon (231 cu. in.) contains—

Chloride of sodium,	1.170	grs.
Sulphate of potassa,	0.456	“
Sulphate of soda,	0.560	“
Bicarbonate of lime,	16.044	“
Bicarbonate of magnesia,	13.563	“
Bicarbonate of iron,	0.038	“
Bicarbonate of soda,	1.261	“
Phosphate of soda,	0.032	“
Silica,	0.722	“
Organic matter,	1.616	“
Alumina,	trace	
Carbonic acid (?)	<u>35.462</u>	

SUMMIT SODA SPRINGS, ALPINE COUNTY, CALIFORNIA.

One U. S. gallon (231 cu. in.) contains—

Bicarbonate of calcium,	43.20	grs.
Carbonate of magnesium,	4.20	“
“ “ sodium,	9.50	“
Chloride “ “	26.22	“
Oxide of iron,	1.75	“
Silica,	2.06	“
Alumina,	1.75	“
Potassium,	trace	
	<u>88.68</u>	

Carbonic acid, 186.35 cubic inches.

Bladen Sulphur Spring, Choctaw County, Alabama. CO_2 86 cubic inches, and H_2S 1.30 cubic inches.

Bladen Vichy Spring, Choctaw County, Alabama. CO_2 110 cubic inches.

Blue Lick Springs, Maysville, Kentucky. NaCl 8, MgCl 76, CaSO_4 59, CO_2 46 cubic inches, H_2S 6 cubic inches.

Clarendon Gaseous Spring, Rutland County, Vermont. CO_2 46.16 cubic inches. 50°F .
Clysmic Spring, Waukesha, Wisconsin. $\text{Mg}(\text{HCO}_3)_2$ 14, $\text{Ca}(\text{HCO}_3)_2$ 16, CO_2 amount not given.

Saratoga Vichy Springs, Saratoga County, New York. NaCl 128, Na_2CO_3 82, $\text{Ca}(\text{HCO}_3)_2$ 95, MgCO_3 41, CO_2 3.83 cubic inches.

Summit Soda Springs, Alpine County, California. NaCl 26, $\text{Ca}(\text{HCO}_3)_2$ 43, CO_2 186.35 cubic inches.

Sweet Chalybeate Spring, Alleghany County, West Virginia. CaSO_4 33, CO_2 104 cubic inches. 75°F .

Sweet Springs, Monroe County, West Virginia. CaCO_3 and CaSO_4 43, CO_2 85.86 cubic inches. 74°F .

IV. PURGATIVE WATERS.

These waters depend for their action largely upon sulphate of sodium and magnesium and associated salts held in solution. An examination of the analyses of the springs thus classified shows that chloride of sodium, carbonates of the alkalis and alkaline earths with free carbonic acid are very frequently associated.

Some of these springs contain so small an amount of salts that their action is but mildly laxative, while others are actively purgative.

When a gentle saline laxative is indicated these waters often give better results than the same amount of Epsom or Glauber's salts dissolved in pure water, and are therefore of greater service.

In congestion of the liver, chronic gastric catarrh, with atony, jaundice, lithæmia, gout, and in the obese, the regular use of a water of this class, properly selected and administered, yields excellent results.

Most of the Saratoga waters are purgative and are more agreeable than many others, from the large quantity of free carbonic acid gas present. Analyses in full are given of Congress, Hathorn and Seltzer, as examples of these important and valuable springs.

Following the typical examples here given will be found analyses of Carlsbad, Hunyadi Janos, Friedrichshall and Ofner Rakozy.

CRAB ORCHARD SPRINGS, LINCOLN COUNTY, KY.

(Analyst, Prof. A. Peters.)

One U. S. gallon (231 cubic inches) contains—

Carbonate of magnesium,	7.640 grs.	Sulphate of magnesium,	205.280 grs.
“ “ iron, . . .	trace	“ “ calcium, . . .	10.792 “
“ “ calcium, . .	53.184 “	Silica,	3.264 “
Chloride of sodium, . .	17,728 “	Loss and moisture, . . .	34.584 “
Sulphate of potassium, .	9.912 “		401.454 “
“ “ sodium, . .	59.072 “		

Carbonate of calcium, magnesium, iron and silica.

PAGASSA HOT SPRINGS, 25 MILES FROM HEADWATERS OF SAN JUAN RIVER, COL.
(Analyst, Prof. O. Loew.)

One U. S. gallon (231 cubic inches) contains—

	No. 1. Grains.	No. 2. Grains.	No. 3. Grains.	No. 4. Grains.
Carbonate of sodium,	2.745	1.945		
“ “ lithium,	0.415	trace	trace	
“ “ calcium,	34.456	34.748	31.834	34.298
“ “ magnesium,	2.832	2.289	2.149	2.127
Sulphate of potassium,	4.164	4.076	3.872	4.146
“ “ sodium,	129.449	128.597	130.769	131.160
Chloride sodium,	17.082	17.146	18.227	17.409
Silicic acid,	3.329	3.043	3.230	2.230
Organic matter	trace	trace	trace	trace
	<u>194.472</u>	<u>191.844</u>	<u>190.181</u>	<u>191.370</u>

Temperature from 101 to 140° F.

LOUISVILLE, OR DUPONT'S ARTESIAN WELL, LOUISVILLE, KENTUCKY.

One U. S. gallon (231 cubic inches) contains—

Carbonate of sodium,	1.896 grs.	Sulphate of sodium,	72.296 grs.
“ “ magnesium,	1.632 “	“ “ magnesium,	77.336 “
“ “ iron,	0.256 “	“ “ aluminum,	01.800 “
“ “ calcium,	4.160 “	“ “ calcium,	29.432 “
Chloride of potassium,	4.224 “	Phosphate of sodium,	1.544 “
“ “ sodium,	621.520 “	Iodide of magnesium,352 “
“ “ magnesium,	14.776 “	Bromide of magnesium,464 “
“ “ aluminum,	1.208 “	Silica,888 “
“ “ calcium,	65.728 “	Organic matter,712 “
“ “ lithium,	0.104 “	Loss,	8.120 “
Sulphate of potassium,	3.224 “		<u>911.672 “</u>
Carbonic acid, 6.16 cubic inches.		Sulphuretted hydrogen, 2.00 cubic inches	
Nitrogen, 1.36 cubic inches.			

SARATOGA SPRINGS, SARATOGA COUNTY, N. Y.

(Analyst, C. F. Chandler.)

One U. S. gallon (231 cubic inches) contains—

	Congress. Grains.	Hathorn. Grains.	Seltzer. Grains.
Carbonate of sodium,	7.472	2.976	20.416
“ “ magnesium,	75.152	104.756	23.904
“ “ iron,	0.248	0.808	1.240
“ “ calcium,	99.992	118.520	62.432
“ “ lithium,	2.992		0.568
“ “ strontium,	trace	trace	trace
“ “ barium,	0.760	1.424	trace
Chloride of potassium,	8.048	9.592	1.336
“ “ sodium,	400.440	509.960	134.288
Sulphate of potassium,	0.888	trace	0.552
Phosphate of sodium,	0.016	0.008	trace
Iodide of “	0.138	0.200	0.032
Bromide of “	8.552	1.556	0.632
Fluoride of calcium,	trace	trace	trace
Biborate of sodium,	trace		trace
Alumina,	trace	0.128	0.376
Silicon,	0.840	1.256	2.560
Organic matter,	trace	trace	trace
	<u>602.636</u>	<u>750.992</u>	<u>248.336</u>
Carbonic acid gas,	362.00 cu. in.	376.00 cu. in.	320.00 cu. in.

- Alburgh Springs, Grand Isle County, Vermont, 15 grains of Na_2SO_4 and K_2SO_4 .
 Avon New Bath, Livingston County, New York. CaCO_3 27, Na_2SO_4 and NaCl 44. 50°F .
 Ballston Springs, Saratoga County, New York—
 Ballston Artesian Lithia Well. NaCl 750, CaCO_3 165, MgCO_3 107, Li_2CO_3 6, CO_2 426 cubic inches. 52°F .
 Franklin Artesian Well. NaCl 659, $\text{Ca}(\text{HCO}_3)_2$ 202, $\text{Mg}(\text{HCO}_3)_2$ 178, NaHCO_3 94, KCl 34, LiHCO_3 7, CO_2 460 cubic inches. 52°F .
 Sans Souci. NaCl 144, CaCO_3 43, $\text{Mg}(\text{HCO}_3)_2$ 39, FeCO_3 6.
 United States. NaCl 425, CaCO_3 29, CO_2 244 cubic inches. 50°F .
 Washington Lithia Well. NaCl 645, $\text{Mg}(\text{HCO}_3)_2$ 158, CaHCO_3 178, Li_2HCO_3 11, $\text{Fe}(\text{HCO}_3)_2$ 2, CO_2 338 cubic inches. 49°F .
 Bedford Springs, Bedford County, Pennsylvania—
 Anderson magnesia. CaSO_4 100, MgSO_4 40. 57.2°F .
 Anderson magnesia. Iron CaSO_4 90, MgSO_4 39, FeCO_3 trace. 57.2°F .
 Bedford Sulphur. CaSO_4 73, MgSO_4 33, H_2S trace, CO_2 6 cubic inches. 62.6°F .
 Beer Springs, Oregon, MgSO_4 48, MgCO_3 12, CaCO_3 15, NaCl 9.
 Benham's Carburetted Saline Spring, Crawford County, Indiana. NaCl 4850, CaCO_3 640, MgCO_3 173, MgSO_4 84, Na_2SO_4 21, FeCO_3 9.
 Bryant's Mineral Well, Lincoln County, Kentucky. MgSO_4 53, CaSO_4 56.
 Butterworth Springs, Kent County, Michigan. CaSO_4 75, MgCl_2 42. 54°F .
 Charleston Artesian Well, Charleston, South Carolina. Na_2CO_3 52, MgSO_4 12. 87°F .
 Cherry Valley Bath House Spring, Otsego County, New York. CaSO_4 85, MgCO_3 and MgSO_4 42, Na_2SO_4 11.
 Cooper's Artesian Well, Hinds County, Mississippi. MgSO_4 23, Na_2SO_4 12, CaSO_4 42. 50°F .
 Crab Orchard, Foley's Spring, Lincoln County, Kentucky,
 Crab Orchard, Sowder's Spring, Lincoln County, Kentucky. MgSO_4 173, CaSO_4 91, NaCl 57, CO_2 ?.
 Crab Orchard Springs, Lincoln County, Kentucky. MgSO_4 205, Na_2SO_4 59.
 Crab Orchard Salts. Contain in 100 grains MgSO_4 63.
 Eaton's White Sulphur Spring, Crawford County, Indiana. NaCl 121, MgSO_4 42, MgCO_3 16, CaCO_3 46. 59°F .
 Estill Spring. See Irvine Spring.
 Foley's Spring. See Crab Orchard Springs.
 Fruit Port Artesian Well, Ottawa County, Michigan. NaCl 464, CaCl_2 111, MgCl 47, Na_2SO_4 46, FeCO_3 5.44 grains. 48°F .
 Glenn Springs, Spartanburgh County, South Carolina. MgSO_4 96, Na_2SO_4 40, H_2S 24 cubic inches.
 Greenville Spring. See Harrodsburg Springs.
 Harrodsburg Springs, Mercer County Kentucky—
 Greenville Spring. MgSO_4 129, CaSO_4 88, MgCO_3 23.
 Saloon Spring. MgSO_4 223, CaSO_4 82, $\text{Fe}(\text{HCO}_3)_2$ 4.
 Hartford Sulphur Spring, Crawford County, Indiana. NaCl 59, MgSO_4 and Na_2SO_4 20, CaCO_3 20.
 Hot Springs, Canon City, Colorado. Na_2SO_4 166, Na_2CO_3 155, Ca_2CO_3 66. 102°F .
 Hot Sulphur Springs, Middle Park, ninety miles from Denver, Colorado. Na_2SO_4 25, Na_2CO_3 22. Temperature 110° to 117°F .

Idaho Warm Springs, Clear Creek County, Colorado. Na_2CO_3 31, Na_2SO_4 29. 106°F .
 Indian Spring, Butts County, Georgia. MgSO_4 572, CaSO_4 57, K_2SO_4 27. 63°F .
 Irvine Spring, Estill County, Kentucky. MgSO_4 263, CaSO_4 and CaCO_3 62.
 Le Roy Springs, Le Roy Station on the Northern Pacific Railroad, Wyoming. Na_2SO_4 117, NaCl 270, MgCO_3 51, CaCO_3 and CaSO_4 100.
 Louisville, or Du Pont's Artesian Well, Louisville, Kentucky. NaCl 622, MgSO_4 77, Na_2SO_4 72.
 Midland Spring, Midland County, Michigan. K_2SO_4 68, Na_2SO_4 18. 47°F .
 Mineral Spring, in dry wash leading from Detrital Valley (Forty-mile Desert) to the Colorado River, Arizona. NaCl 232, MgSO_4 101, CaSO_4 76, Na_2SO_4 30.
 Pagosa Hot Springs, twenty-five miles from the headwater of the San Juan River, Colorado—

No. 1. Na_2SO_4 129.

No. 3. Na_2SO_4 131.

No. 2. Na_2SO_4 129.

No. 4. Na_2SO_4 131.

Porter's Sulpho-Chalybeate Spring, Denver, Colorado. Na_2SO_4 27, CaCO_3 10.

Rhea Springs, Rhea County, Tennessee. 70 grains of CaSO_4 .

Saloon Spring. See Harrodsburg Springs.

Saratoga Springs, Saratoga County, New York—

Champion. $\text{Ca}(\text{HCO}_3)_2$ 227, NaCl 702, $\text{Mg}(\text{HCO}_3)_2$ 194, CO_2 462. 49°F .

Columbian. NaCl 267, CaCO_3 68, MgCO_3 28, FeCO_3 6, CO_2 272 cubic inches.

Congress. NaCl 400, CaCO_3 100, MgCO_3 75, NaBr 9, Li_2CO_3 3, CO_2 392 cubic inches. 52°F .

Crystal. NaCl 336, CaCO_3 71, MgCO_3 45, Li_2CO_3 3, CO_2 312 cubic inches. 50°F .

Empire. NaCl 507, CaCO_3 76, MgCO_3 25, CO_2 344 cubic inches.

Eureka. NaCl 117, CaCO_3 41, MgCO_3 29, FeCO_3 3, CO_2 232 cubic inches.

Excelsior. NaCl 371, CaCO_3 77, MgCO_3 32, FeCO_3 3, NaCO_3 15.

Geysier. NaCl 562, CaCO_3 118, MgCO_3 83, Na_2CO_3 49, Li_2CO_3 4, CO_2 456 cubic inches. 46°F .

Glacier. NaCl 702, CaCO_3 158, MgCO_3 115, Li_2CO_3 6, KCl 40, CO_2 465.44 cubic inches.

Hamilton. NaCl 299, CaCO_3 98, MgCO_3 39, FeCO_3 5, Na_2CO_3 34, CO_2 320.

Hathorn. NaCl 510, CaCO_3 118, MgCO_3 105, CO_2 376 cubic inches.

High Rock. NaCl 390, CaCO_3 92, MgCO_3 33, CO_2 408 cubic inches. 52°F .

Pavilion. NaCl 460, CaCO_3 83, MgCO_3 45, Li_2CO_3 6, CO_2 328 cubic inches.

Putnam Spring. NaCl 214, $\text{Ca}(\text{HCO}_3)_2$ 68, $\text{Mg}(\text{HCO}_3)_2$ 51, CO_2 348 cubic inches. 51°F .

Red Spring. NaCl 70, CaCO_3 59, MgCO_3 21.

Saratoga Alum. NaCl 565, CaCO_3 57, MgCO_3 20, CO_2 212 cubic inches.

Seltzer. NaCl 134, CaCO_3 62, MgCO_3 24, CO_2 320 cubic inches. 50°F .

Star. NaCl 398, CaCO_3 86, MgCO_3 37, CO_2 400 cubic inches. 52°F .

Triton Spring. NaCl 238, CaCO_3 91, MgCO_3 42, Li_2CO_3 3 grains, CO_2 360 cubic inches. 60°F .

United States. NaCl 142, CaCO_3 65, MgCO_3 43, Li_2CO_3 3, CO_2 240 cubic inches.

Union. NaCl 458, CaCO_3 67, MgCO_3 65, CO_2 384.96 cubic inches. 48°F .

Vichy. See Alkaline Waters.

Walton or Iodine. NaCl 187, MgCO_3 75, CaCO_3 26, CO_2 330 cubic inches.

Washington Spring. NaCl 183, $\text{Ca}(\text{HCO}_3)_2$ 84, $\text{Mg}(\text{HCO}_3)_2$ 66, CO_2 364 cubic inches. 45°F .

Seltzer Spring, Boulder County, Colorado. Na_2SO_4 108, $\text{Ca}(\text{HCO}_3)_2$ 43. Temperature, 40°F .
Seltzer, Springdale, Colorado. Na_2SO_4 230, Ca_3CO_3 85, FeCO_3 8.

Spring Lake Well, Ottawa County, Michigan. NaCl 406, CaCl_2 113, Na_2SO_4 47, MgCl 36. 52°F .

Sowder's Spring. See Crab Orchard Springs.

Warm Spring, at edge of Salt Lake City, Utah. MgSO_4 60, NaCl 234.

Warren Springs, Warren County, North Carolina. CaCl 11, CaSO_4 and CaCO_3 and CaCl_2 10, MgSO_4 8.

White Creek Springs, twelve miles from Nashville, Tennessee. 55 grains of CaCO_3 and CaSO_4 .

V. ALKALINE WATERS.

These owe their alkalinity to the preponderance of carbonate and bicarbonate of sodium, and also, though in small quantities, the carbonate of potassium, lithium, calcium and magnesium. These waters are frequently surcharged with carbonic acid gas, which enhances their therapeutic value very considerably, as it renders them more palatable and converts the alkaline taste, so unpleasant to many, into an agreeable acidity. These springs may be cold or hot, and may contain lime, salt or iron in varying amount.

Typical examples of this class, that is, springs containing sodium carbonate or bicarbonate with free carbonic acid gas, are, unfortunately, less numerous than we would wish, in view of their enormous importance in so many chronic diseases. Most of our alkaline springs contain other mineral substances which render them less valuable in the treatment of disease. The conditions in which these waters produce their best effects are the following: Chronic gastric catarrh, especially with hyperacidity, and catarrhal inflammation of mucous membrane of the biliary passages. Good results have also been obtained in acute catarrhal nephritis, and not a few observers have seen diabetics improve under its use. In acute cystitis and urethritis springs of this class do good by diluting the urine and diminishing its acidity, and so alleviate suffering and hasten recovery; at times brilliant results are obtained in certain diathetic conditions, such as lithæmia, gout and chronic rheumatism. From this brief summary it is evident that these waters constitute a most important addition to our means of combating disease.

Our country is so rich in mineral waters, it is in the highest degree probable that numerous springs belonging to this class are in existence and only need analysis to make known their value. It is to be hoped that before long many additions will be made to our meagre list of springs belonging to this group.

The following analyses are examples of this class, and I would particularly call attention to that of Vichy, in France, as a type of an alkaline water:—

SARATOGA VICHY SPRING, SARATOGA CO., N. Y.

Contains in one U. S. gallon of 231 cubic inches—

Chloride of sodium,	128.689
“ “ potassium,	14.113
Bromide of sodium,	0.990
Iodide of sodium,	trace.
Fluoride of calcium,	trace.
Bicarbonate of lithia,	1.760
“ “ soda,	82.873
“ “ magnesia,	41.503
	<hr/> 269.928

	269.928
Bicarbonate of lime,	95.522
“ “ strontia,	trace.
“ “ baryta,	0.593
“ “ iron,	0.052
Sulphate of potassa,	trace.
Phosphate of soda,	trace.
Alumina,	0.473
Silica,	0.758
Organic matter,	trace.
Total,	367.326

Carbonic acid gas in one gallon, 383.071 cubic inches.

Temperature 50° F.

VICHY GRANDE GRILLE AND HÔPITAL
SPRINGS.

(Analyst, Mossiér.)

One U. S. gallon (231 cu. in.) contains—

	Grande-Grille. Grains.	Hôpital. Grains.
Carbonate of soda,	259.03	254.75
“ “ lime,	12.23	18.62
“ “ magnesia,	2.28	2.05
“ “ iron,	0.60	2.73
Chloride of Sodium,	23.94	8.36
Sulphate of soda,	42.33	47.65
	340.41	334.16

Free carbonic acid 90–100 cu. in.

OJO CALIENTE SPRING, OJO CALIENTE, NEW
MEXICO.

(Analyst, O. C. Marsh.)

One U. S. gallon (231 cu. in.) contains—

Sodium carbonate,	115.019 grs.
Lithium “	0.123 “
Calcium “	2.435 “
Magnesium “	1.273 “
Iron “	5.910 “
Potassium sulphate,	3.019 “
Sodium “	7.942 “
“ chloride,	32.209 “
Silica,	0.123 “
	168.053 “

Carbonic acid undetermined.

Temperature 100° F.

Adams Springs, Lake County, California. MgCO_3 99, Na_2CO_3 57, CaCO_3 29, CO_2 300 cubic inches.

Borax Springs. See Hot Borate Springs.

California Seltzer Springs, Mendocino County, Cal. CaCO_3 67, MgCO_3 43, Na_2CO_3 35.

Carlisle, near Pueblo, Colorado. 60° F.

Clinton Spring, Cliff Street, New York City. NaCl 58, MgCO_3 35, CaCO_3 30, CO_2 68½ cubic inches.

Congress Spring, Santa Clara County, California. Na_2CO_3 123, NaCl 118, FeCO_3 13.

Hot Borate Springs, Borax Lake, Lake County, California. NH_4HCO_3 108, 103, NaCl 85, NaHCO_3 77.

Indian Spring, Martin County, Indiana. NaCl 39, $\text{CaCO}_3 + \text{CaSO}_4$ 53, $\text{MgCO}_3 + \text{MgSO}_4$ 49. 53° F.

Iron Duke, Canon City, Colorado. Na_2CO_3 156, NaCl 170.

Olympian Springs, Bath County, Kentucky. NaCl 166, MgCl_2 55.

Richfield Sulphur Spring, Otsego County, New York. CaSO_4 and $\text{Ca}(\text{HCO}_3)_2$ 92, MgSO_4 and $\text{Mg}(\text{HCO}_3)_2$ 39, NaCl 21, H_2S 4 cubic inches. 48° F.

Saratoga Vichy Springs, Saratoga County, New York. NaCl 128, Na_2HCO_3 82, CaHCO_3 95, CO_2 383 cubic inches.

Sharon Magnesia Spring, Schoharie County, New York. CaSO_4 76, $\text{MgSO}_4 + \text{Mg}(\text{HCO}_3)_2$ 53, H_2S 3.3 cubic inches. 48°F .

Snowden Mineral Spring, Valley of Yoncalla, two miles south of Drain's Station, Oregon. MgCl 145, NaCl 173, CaCl_2 115.

St. Louis Magnetic Spring, St. Louis, Gratiot County, Michigan. NaHCO_3 86, $\text{Ca}(\text{HCO}_3)_2$ 56, CaSO_4 54. 50°F .

Vichy Springs. See New Almaden Springs.

West Baden Springs, Orange County, Indiana. NaCl 78, CaCO_3 41, MgCO_3 39, MgSO_4 36, MgCl_2 11.

Wilhoit Springs, Clackamas County, Oregon. MgCO_3 85, Na_2CO_3 88, NaCl 201, FeO 6, CO_2 338.41 cubic inches.

Wilson's Saline Chalybeate, Lexington, Fayette County, Kentucky. CaCl_2 33, MgCl_2 18, MgCO_3 15, FeCO_3 $2\frac{1}{2}$, CO_2 33 cubic inches.

VI. SALINE WATERS.

These waters are solutions of common salt, containing, also, small quantities of the chlorides of the alkalis and alkaline earths, and occasionally minute traces of iodine and bromine. It is plain that the effects produced by these springs are due to chloride of sodium and the increased amount of water consumed. This salt, when taken into the stomach, dissolves albumin and starches; promotes the digestion and absorption of food; supplies the intestines with chyme rich in albumin and starches; enters the blood, which carries it to all the tissues of the body, after supplying its own needs.

In solutions of suitable strength it produces an agreeable effect on mucous membranes, favoring free and healthy secretion. It is quickly absorbed, but when taken highly concentrated acts as an irritant and produces emesis and diarrhoea. The therapeutic dose is from one to five drachms daily.

The usefulness of these waters is very restricted, and is indicated by what has been said as to its physiological action. Good results may be expected in certain dyspepsias, with defective gastric secretion and sluggishness of the bowels, as well as chronic inflammatory exudations which are being absorbed slowly or not at all. When rapid absorption is necessary the water should be given hot, and in certain cases of dyspepsia these results are best obtained by its administration early in the morning, before breakfast.

When it is desired to administer these waters, in ordinary cases, preference should be given to those springs charged with carbonic acid, as this gas greatly increases the palatability of this class of waters.

These waters, as well as the alkaline earths, when carbonated, are especially valuable in many acute affections attended with fever, particularly when nourishment is taken in small amounts, as they supply the body with mineral substances absolutely needed. The proper supply of salines in disease is too often overlooked.

The following analyses are good examples of these waters:—

HALLECK SPRING, ONEIDA COUNTY, NEW YORK.
(Analyst, Prof. Noyes.)

One U. S. gallon (231 cu. in.) contains—

Chloride of sodium,	624.00	grs.
“ calcium,	104.00	“
“ magnesium,	32.00	“
Sulphate of calcium,	40.00	“
	<u>800.00</u>	“

AKESION SPRING, SALINE COUNTY, MISSOURI.
(Analyst, C. P. Williams.)

One U. S. gallon (231 cu. in.) contains—

Chloride of sodium,	756.114	grs.
“ potassium,	28.594	“
“ lithium,	0.294	“
“ magnesium,	87.313	“
“ calcium,	74.791	“
Bromide of magnesium,	0.131	“
Nitrate of magnesium,	0.178	“
“ ammonium,	0.172	“
Sulphide of sodium,	2.609	“
Sulphate of calcium,	57.934	“
“ barium,	0.150	“
Phosphate of calcium,	0.243	“
Carbonate of calcium,	40.251	“
“ iron, (ferrous)	0.267	“
“ manganese,	0.199	“
“ alumina,	0.167	“
Organic matter,	0.513	“
	1049.920	“

SALT SPRING, NEAR ALBA, BRADFORD
COUNTY, PENNSYLVANIA.

(Analyst, Genth.)

One U. S. gallon (231 cu. in.) contains—

Chloride of sodium,	4693.608	grs.
“ iron,	trace	
“ calcium,	780.224	“
“ magnesium,	110.960	“
Bicarbonate of iron,	5.256	“
“ calcium,	1.752	“
Silicic acid,	2.950	“
Bromine not determined.		

5594.750 “

WIESBADEN, THE KOCHBRUNNEN, NASSAU,
GERMANY.

One U. S. gallon (231 cu. in.) contains—

Chloride of sodium,	419.92	grs.
“ potassium,	8.96	“
“ lithium,	0.08	“
“ calcium,	28.88	“
“ magnesium,	12.48	“
Bromide of magnesium,	0.16	“
Sulphate of lime,	5.52	“
Carbonate of lime,	25.68	“
“ protoxide of iron,	0.32	“

502.00 “

Temperature, 156° F. Carbonic acid, 52
cubic inches.

Akesion Spring, Brownville, Saline County, Missouri. NaCl 756, MgCl₂ 87, CaCl₂,
CaSO₄ and CaCO₃ 173.

Albany Artesian Well, Ferry Street, Albany, New York. NaCl 504, FeCO₃ 8, MgCO₃,
16, Na₂CO₃ 40, CO₂ 224 cubic inches. 51° to 52° F.

Ballston Condo Dentonean, Saratoga County, New York. NaCl 480, CaCO₃ 123, MgCO₃,
93, CO₂ 357 cubic inches.

Ballston Spa, Saratoga County, New York. NaCl 144, CaCO₃ 43, Mg(HCO₃)₂ 39.

Bangor's Company's Well, Bangor, Michigan. NaCl 159, MgCl 09, CaSO₄ 56, CaCl.

Black Sulphur Spring, Bath County, Kentucky. 58 grains NaCl.

- Blue Lick Springs (Lower), Nicholas County, Kentucky. NaCl 516, CaSO₄ 41, MgCl₂ 40, CaCO₃ 23, CO₂ 98, H₂S 17.
- Blue Lick Springs (Upper), Nicholas County, Kentucky. NaCl 516, CaSO₄ 144, MgCl₂ 37, CaCO₃ 25, CO₂ 48 cubic inches, H₂S 8 cubic inches.
- Borland Well, Wood County, West Virginia. NaCl 240, NaHCO₃ 112, Na₂SO₄ 38.
- Boulder Spring Water, Colorado. NaCl 514, Ca₂CO₃ 125, MgCO₃ 100, Na₂SO₄ 62, CO₂.
- Bryant's Spring, No. 4, Lincoln County, Kentucky. Na₂SO₄ 12, NaCl 54.
- Cañon City Mineral Springs, Cañon City, Fremont County, Colorado.
- Aqua Vida. NaCl 121, Na₂CO₃ 73, MgCO₃ 18.
- Big Ute. NaCl 132, CaCO₃ 43, MgCO₃ 15.
- Congress. NaCl 38, MgCO₃ 18.
- Iron Duke. NaCl 80, Na₂CO₃ 74, MgCO₃ 15.
- Little Ute. NaCl 114, Na₂CO₃ 74, MgCO₃ 14.
- Condo Dentonean Springs. See Ballston Springs.
- Des Chutes Hot Springs, Wasco County, Oregon. Na₂CO₃ and NaCl 55. 143° to 145° F.
- East Clarion Salt Spring, Elk County, Pennsylvania. NaCl 337, CaCl₂ 52, MgCl₂ 15.
- East Saginaw Company's Well, Michigan. NaCl 103, MgCl₂ 58, CaCl₂ 14, CaSO₄ 92.
- Halleck's Spring, Oneida County, New York. NaCl 624, CaCl₂ 104, MgCl₂ 32, CaSO₄ 40.
- Highgate Springs, Franklin County, Vermont. NaCl 23, Na₂CO₃ 14.
- Iola Mineral Well, Kansas. NaCl 782, Ca(HCO₃)₂ 49, Mg(HCO₃)₂ 21, KCl 14.
- Kanawha Saline Spring, Kanawha Valley, West Virginia. NaCl 240, Na₂HCO₃ 112, Na₂SO₄ 38.
- La Fayette Artesian Well, Tippecanoe County, Indiana. NaCl 325, MgCO₃ and MgCl₂ 59, CaSO₄ 56. 55° F.
- Lansing Spring, Lansing, Ingham County, Michigan. NaCl 267, Na₂CO₃ 65, CaCO₃ 62, CO₂ 198 cubic inches. 53.5° F.
- Liberty Hot Springs, or Wagonwheel Gap Springs, Morrison, 12 miles from Denver, Colorado.
- No. 1. Na₂CO₃ 41, NaCl 17. Temperature 150° F.
- No. 2. Na₂CO₃ 84, NaCl 19. Temperature 140° F.
- Livingston Artesian Well, Livingston, Sumter County, Alabama. NaCl 295 grains. 68° F.
- Lodi Artesian Well, Wabash County, Indiana. NaCl 502, CaCl₂ and CaSO₄ 104, MgCl₂ 54.
- Lower Blue Lick Springs. See Blue Lick Springs.
- Lubec Spring, Lubec Bay, Maine. NaCl 90, MgCl₂ 60, NaSO₄ 25.
- Mineral Spring, Walnut Hill, Fayette County, Kentucky. NaCl 234, MgCl₂ 19.
- Mineral Water, Paint Lick, Madison County, Kentucky. NaCl 27, Na₂CO₃ 16.
- Ott's Well, Crawford County, Indiana. NaCl 3781, CaCO₃ 323, MgCO₃ 86, FeCO₃ 12, MgSO₄ and Na₂SO₄ 58.
- Parnassus Springs, 12 miles southward of Pueblo, Colorado.
- No. 1. Na₂CO₃ 74, NaCl 60. Temperature 66° F.
- No. 2. Na₂CO₃ 69, NaCl 61. Temperature 72.5° F.
- No. 3. Na₂CO₃ 43, NaCl 31. Temperature 59° F.
- Salina Spring (Mr. Peterson), near Tarentum, Pennsylvania. NaCl 1927, CaCl₂ 501, MgCl₂ 126, MgCO₃ 80, CaCO₃ 161.
- Salina Wells, Syracuse, New York. NaCl 90, MgCl₂ 7, CaCl₂ 14, CaSO₄ 36.
- Salt Spring, near Alba, Bradford County, Pennsylvania. NaCl 4694, CaCl₂ 780, MgCl₂ 111, Fe(HCO₃)₂ 5.

Salt Sulphur, Bath County, Kentucky. NaCl 166, MgCl 55.
 Salt Water, near Conemaugh, near Saltzburg, Indiana County, Pennsylvania. NaCl 4165.
 CaCl₂ 918, MgCl₂ 232.
 Sheboygan Well, Sheboygan, Wisconsin. 307 grains of NaCl. 50° F.
 Siloam Springs, Garfield County, Colorado. NaCl 1137, Ca₂SO₄ 78. 103° F.
 South Park Springs, Colorado. NaCl 198, Na₂CO₃ 156.
 South Park Sulphur Springs, South Park, Colorado. NaCl 97, Na₂CO₃ 77.
 Steamboat Springs, Colorado. NaCl 240.
 St. Louis Artesian Well, St. Louis, Missouri. NaCl 347, MgCl₂ 34, CaCl₂ 27, CO₂ 6, H₂S 24.
 Sweet Springs, Saline County, Missouri. NaCl 90, MgCl₂ 22.
 Syracuse Salt Wells, New York. NaCl 94, MgCl₂ 85, CaCl₂ 49, CaSO₄ 32.
 Tarentum Salt Well. See Salina Springs.
 Terre Haute Spring, Clay County, Indiana. NaCl 316 grains.
 Thomas Well's Brine, Fountain County, Indiana. NaCl 4573, CaCl₂ 215, CaCO₃ 107,
 MgCl₂ 53.
 Upper Blue Lick Springs. See Blue Lick Springs.
 Verona Spring, Oneida County, New York. NaCl 720, CaSO₄ 60, CaCl₂ and MgCl₂ 68.
 Wagonwheel Gap Springs. See Liberty Hot Springs.
 Waterloo Well, Waterloo, Wisconsin. NaCl and NaNO₃ 60.
 Williams Mineral Well, Woodford County, Kentucky. NaCl 137, MgCl₂ 16.
 Yampa Spring, Glenwood, Colorado.

VII. CALCAREOUS WATERS.

The efficacy of these waters depends upon the presence of the sulphate and carbonate of calcium and other alkaline earths. They are generally produced as follows: The excess of carbonic acid in the water, resulting from the decomposition of the carbonate by means of sulphuric acid (this acid is produced by the addition of iron pyrites), acts upon the sulphates and carbonates of the alkaline earths to dissolve them.

Most of the lime in these fountains exists in the form of a sulphate, which is popularly known as gypsum. It is this salt that gives these waters the property called *hardness*, *i. e.*, that peculiar reaction to soap and the special effect upon the palate.

These springs have been used in cystitis and diabetes, but with doubtful results. Occasionally diuresis may be excited and in rachitic children favorable results may be obtained, though the constipating effect would be antagonized. It should be remembered that 10 grains of sulphate of lime to the gallon tends to retard digestion and causes constipation, and consequently they are often contra-indicated. In general, it may be said that springs of this class are of doubtful utility, except when they contain other more active mineral substances.

I append a few examples of this class of water:—

CATOOSA SPRINGS, CATOOSA COUNTY, GEORGIA. NO. I ALL-HEALING SPRING.
 One U. S. gallon (231 cu. in.) contains—

Calcium sulphate,	39.000 grs.
Magnesium sulphate,	} 34.000 "
" carbonate,	
Carbonic acid,	4.422 "
	<hr/> 77.422

TATE EPSOM SPRINGS, GRANGER COUNTY, TENNESSEE.
(Analyst, T. S. Anterelle.)

One U. S. gallon (231 cu. in.) contains—	
Sulphate of calcium,	160.66 grs.
“ magnesium,	31.97 “
“ sodium,	8.50 “
“ potassium,	1.54 “
Chloride of sodium,	40.27 “
“ iron,	2.92 “
“ manganese,	0.69 “
Iodide of sodium,	trace
Phosphate of calcium,	1.14 “
Carbonate of calcium,	21.56 “
Silica,	2.70 “
Nitric acid,	0.02 “
	<hr/>
	271.97 “

Temperature 55° F.

Alleghany Springs, Montgomery County, Virginia. CaSO_4 96, MgSO_4 42. 53° F.

Aurora Saline Springs, Aurora, Oregon. CaCl 474, NaCl 356.

Beauty Spring. See Capon Springs.

Beloit Iodo-magnesian Springs, Beloit, Wis. $\text{Ca}(\text{HCO}_3)_2$ 15, $\text{Mg}(\text{HCO}_3)_2$ 12. 48° F.

Bishop's Well, New Brunswick, New Jersey. CaSO_4 149, MgSO_4 31.

Blue Ridge Spring, Botetourt County, Virginia. CaSO_4 100, MgSO_4 48.

Bordin's Well, Eaton Rapids, Michigan. CaCO_3 and CaSO_4 85 grains.

Butterworth Spring, Grand Rapids, Mich. CaSO_4 75, MgCl 41.

Capon Springs (Beauty Spring), Hampshire Co., W. Va. CaCO_3 66, CO_2 68 cubic inches. 60° F.

Capon Springs (Main Spring), Hampshire Co., W. Va. CaCO_3 66, CO_2 68 cubic inches. 60° F.

Carlisle Spring, on Arkansas River, twenty miles above Pueblo, Colorado. CaCO_3 22, MgCO_3 11. 65° F.

Catoosa Springs, Catoosa County, Georgia.

All-Healing. CaSO_4 39, MgSO_4 and MgCO_3 34.

Alum. CaSO_4 41, MgSO_4 and MgCO_3 36, Al 1.

Black Sulphur. CaSO_4 41, MgSO_4 and MgCO_3 36.

Buffalo. CaSO_4 45, MgSO_4 and MgCO_3 42.

Chalybeate. CaSO_4 42, MgSO_4 and MgCO_3 35, FeCO_3 trace.

Congress. CaSO_4 39, MgSO_4 and MgCO_3 34.

Cosmetic. CaSO_4 42, MgSO_4 and MgCO_3 36.

Magnesia. CaSO_4 42, MgSO_4 and MgCO_3 36.

Red Sweet. CaSO_4 44, MgSO_4 and MgCO_3 38.

White Sulphur. CaSO_4 45, MgSO_4 and MgCO_3 40.

Cave Spring. See Chittenango Spring.

Chalybeate Spring, Estill County, Kentucky. CaSO_4 17, CaCO_3 9, MgSO_4 10, FeCO_3 2, CO_2 32 cubic inches.

Cherry Valley North Spring, Otsego County, New York. CaSO_4 150, MgCO_3 10.

- Chittenango Cave Spring, Madison County, New York. CaSO_4 106, MgCO_3 14, CO_2 25.6 cubic inches, H_2S 3.2 cubic inches. 49° F.
- Chittenango Magnesia Spring, Madison County, New York. CaSO_4 81, MgCO_3 13, CO_2 36 cubic inches. 49° F.
- Chittenango White Sulphur Spring, Madison Co., N. Y. CaSO_4 80, CO_2 36 cubic inches.
- Clifton Springs, Ontario County, New York. CaSO_4 and CaCO_3 79, MgSO_4 and MgCO_3 30.
- Colorado or Manitou Springs, Colorado Springs, Colorado.
- Iron Ute. CaCO_3 33, Na_2CO_3 33, MgCO_3 8. 43.3° F.
- Little Chief. CaCO_3 42, Na_2SO_4 29, NaCl 27. 43° F.
- Manitou. CaCO_3 62, Na_2CO_3 26, MgCO_3 12. 56° F.
- Navajoe. CaCO_3 72, Na_2CO_3 70, MgCO_3 19. 50.2° F.
- Shoshone. CaCO_3 61, Na_2CO_3 50. 48.5° F.
- Ute Soda. CaCO_3 22, Na_2CO_3 13.
- Eaton Rapids Well, Eaton Rapids in Eaton County, Michigan.
- Bordine Well. CaCO_3 and CaSO_4 , 85 grains.
- Frost Well. CaCO_3 and CaSO_4 , 42 grains.
- Mosher Well. CaCO_3 and CaSO_4 , 65 grains.
- Shaw Well. CaCO_3 and CaSO_4 , 69 grains.
- Stirling Well. CaSO_4 , 46 grains.
- East Clarion Spring, Elk County, Pennsylvania, NaCl 337, CaCl_2 52, MgCl_2 15, BaCl_2 1 $\frac{1}{4}$.
- Frost Well. CaCO_3 and CaSO_4 , 42 grains.
- Gettysburg Katalysine Spring, Adams County, Pennsylvania. $\text{Ca}(\text{HCO}_3)_2$ 16, MgSO_4 7. 57° F.
- Greenbrier White Sulphur. See White Sulphur Spring.
- Guilford Spring, Guilford Centre, Windham County, Vermont. 15.18 grains CaCO_3 .
- Holston Springs, Scott County, Virginia. CaSO_4 20, MgSO_4 13. 68.5° F.
- Liberty Hot or Wagonwheel Gap Springs, Morrison, twelve miles from Denver, Colorado, No. 2. CaCO_3 18.
- Magnesia Springs. See Chittenango Spring.
- Main Spring. See Capon Springs.
- Manitou Springs, Manitou, Colorado. CaCO_3 135. 56° F.
- M'Carthy's Spring, Huntingdon County, Pennsylvania. 95 grains of CaSO_4 and $\text{Ca}(\text{HCO}_3)_2$, 42 grains MgSO_4 .
- Mineral Park Bitter Spring, Arizona. CaSO_4 69, MgSO_4 38.
- Montvale Spring, Blount County, Tennessee. 75 grains of CaSO_4 . 60° F.
- Mosher Well, Eaton Rapids, Michigan. CaCO_3 and CaSO_4 , 65 grains.
- Newburg Springs, Orange County, Vermont. 17.60 grains CaCO_3 .
- Red Sulphur, Estill County, Kentucky. CaCO_3 and CaSO_4 22, CO_2 40 cubic inches, H_2S .56 cubic inches.
- Richfield Magnesia Spring, Otsego County, N. Y. CaSO_4 and $\text{Ca}(\text{HCO}_3)_2$ 55, MgSO_4 and $\text{Mg}(\text{HCO}_3)_2$ 23, CO_2 5 cubic inches. 53° F.
- Saltillo Springs. See M'Vitney's and M'Carthy's Springs.
- Shaw Well, Eaton Rapids, Michigan. CaCO_3 and CaSO_4 , 69 grains.
- Sterling Well, Eaton Rapids, Michigan. CaSO_4 , 46 grains.
- Tate Epsom Springs, Granger County, Tennessee. CaSO_4 , 161 grains. 55° F.
- Warm Spring, Bath County, Virginia. CaSO_4 15. 98° F.
- Warren Springs (cold sulphur), Warren County, North Carolina. CaSO_4 31.

Waukesha Springs, Waukesha County, Wisconsin.

Arcadian Spring. $\text{Ca}(\text{HCO}_3)_2$ 16, $\text{Mg}(\text{HCO}_3)_2$ 11.

Bethesda Spring. $\text{Ca}(\text{HCO}_3)_2$ 17. 60° F.

Fountain Spring. $\text{Ca}(\text{HCO}_3)_2$ 14.

Glenn Mineral Spring. $\text{Ca}(\text{HCO}_3)_2$ 16.

Hygeia Spring. $\text{Ca}(\text{HCO}_3)_2$ 17.

Silurian Spring,

Waukesha Mineral Rock Spring. $\text{Ca}(\text{HCO}_3)_2$ 10.

White Sulphur Spring. See Chittenango Spring.

White Sulphur Springs, Greenbrier County, West Va. CaSO_4 74, MgSO_4 19, H_2S 1.54 to 3.03 cubic inches. 62° F.

Yellow Sulphur Springs, Montgomery County, Virginia. CaSO_4 63, MgSO_4 21. 55° F.

VIII. SULPHURETTED AND HOT SPRINGS.

These springs owe their properties to the presence of sulphuretted hydrogen and heat. The sulphurets of sodium and potassium are often present. Sulphuretted hydrogen, next to carbonic acid, occurs more frequently in mineral springs than any other gas.

It frequently issues from the earth in the neighborhood of some active volcanoes, and is probably evolved through the decomposition of the sulphurets by hot water. When this gas is brought in contact with the air it is readily changed, the hydrogen combining with the oxygen and the sulphur is precipitated, giving to the water a milky appearance.

This gas is irrespirable, and quickly produces death if inhaled in considerable quantities, and when employed for its local effect upon the respiratory mucous membrane must be largely diluted with air. The first symptom of its poisonous action is a small, feeble pulse, with excessive weakness, and later pulse and respiration may both fail, with unconsciousness, followed by profuse perspiration.

Internally, in moderate quantities, peristalsis and perspiration are both augmented. When the gas is locally applied to mucous membranes, it produces stimulation of secretion. A sulphur spring of moderate strength contains not less than 12 cu. in. of sulphuretted hydrogen in the gallon, though many springs contain so small an amount that therapeutically they are inert, and the good effects observed are due to the influence of the increased use of water, change of scene and climate, cessation of work, regular meals, good hygiene and hope, all of which contribute strongly to restore health at all springs, especially the Diuretic Springs.

Many of these waters are hot, though some are cold. A few contain an excess of carbonic acid, and any member of the other classes may be charged with sulphuretted hydrogen on an increase of temperature, and consequently many of these springs are also classified under other heads.

Frequently these springs are used as baths, and with good results, especially in chronic skin affections, as eczema, rheumatism and gout.

Internally, they have been used for chronic mucous membrane inflammation, especially of bronchial tubes and upper respiratory tract, and also in hepatic congestion and catarrhal phthisis. The simple hot springs are largely used as baths, and are of unquestionable value in certain cases of lithæmia, chronic rheumatism and gout, as well as in dysmenorrhœa, syphilis and psoriasis, pityriasis and eczema. Given as a douche, good effects have been observed in neuralgias, paralyses and enlarged joints with contracted muscles or tendons. The principal hot springs are found in Virginia, Colorado and New Mexico.

The following analyses are types of sulphuretted and hot springs :—

ALPENA SPRINGS, ALPENA COUNTY, MICHIGAN.		SHARON WHITE SULPHUR SPRING, SCHO- HARIE COUNTY, NEW YORK.	
(Analyst, Prof. S. P. Duffield.)		One U. S. Gallon (231 cu. in.) contains—	
One U. S. gallon (231 cu. in.) contains—		Bicarbonate of magnesium, . . . 24.00 grs.	
Carbonate of potassium, . . .	trace	Sulphate of " . . .	34.00 "
" sodium,	10.912 grs.	" calcium,	85.40 "
" magnesium, . . .	37.288 "	Hydrosulphate of calcium and	
" iron,	1.360 "	magnesium,	3.00 "
" calcium,	38.296 "	Chloride of sodium and magne-	
Chloride of sodium,	68.256 "	sium,	2.70 "
Sulphate of calcium,	30.056 "		
Alumina and silica,	3.088 "		149.10 "
	189.256 "	Sulphuretted hydrogen, 20.5 cu. in.	
Sulphuretted hydrogen, . . .	35.36 cu. in.	Temperature, 48° F.	
Carbonic acid,	8.40 "		
Nitrogen,	0.24 "		

HAGERS SPRING, HAGERS COUNTY, TENNESSEE.

One U. S. gallon (231 cu. in.) contains—	
Sulphate of calcium and hydro-	
chlorate of sodium,	38.00 grs.
Sulphuretted hydrogen, . . .	51.52 cu. in.
Carbonic acid,	6.72 "

HOT SPRINGS.

LAS VEGAS HOT SPRINGS, LAS VEGAS, NEW MEXICO.				HOT SPRINGS, GARLAND COUNTY, ARKANSAS.	
(Analyst, Prof. F. V. Hayden, U. S. Geologist.)				One U. S. Gallon (231 cu. in.) contains—	
One U. S. Gallon (231 cu. in.) contains—				Carbonate of magnesium, 0.128 grs.	
	No. 1.	No. 2.	No. 3.	"	" calcium, 3.968 "
	Grains.	Grains.	Grains.		
Carbonate of sodium,	1.72	1.17	5.00	Chloride of sodium, 0.008 "
" calcium	} 1.08	} 10.63	} 11.43	Sulphate " potassium, 0.232 "
" magne- sium,				" sodium, 0.376 "
				" " calcium, 0.112 "
Sulphate of sodium,	14.12	15.43	16.21	Sesquioxide of iron, 0.104 "
Chloride of sodium,	27.26	24.37	27.37	Iodine and bromine, trace "
Potassium,	trace.	trace.	trace.	Silicate calcium, 0.464 "
Lithium,	trace.	trace.	trace.	Silica, 1.864 "
Silicic acid,	1.04	trace.	2.51	Alumina, 0.448 "
Iodine,	trace.	trace.	trace.	Organic matter, 0.064 "
Bromine,	trace.	trace.	trace.		
	45.22	51.60	62.49		7.768 "
Temperature, . . .	130° F.	123° F.	123° F.	Temperature, 93-150° F.	

HOT SPRINGS, BATH COUNTY, VA.

(Analyst, Wm. B. Rogers.)

One U. S. Gallon (231 cu. in.) contains—

Sulphate of magnesium, . . .	3.534 grs.
“ “ calcium, . . .	3.007 “
“ “ sodium, . . .	3.148 “
Chloride of magnesium, . . .	0.242 “
Carbonate of calcium, . . .	16.200 “
“ “ magnesia, . . .	3.058 “
“ “ iron, . . .	0.221 “
Silica, . . .	0.103 “
	<hr/> 29.513 “

Temperature, 100–106° F.

SULPHURETTED AND THERMAL WATERS.

Alpena Well, Alpena County, Michigan. NaCl 68, MgCO₃ 37, CaCO₃ 38, H₂S 35 cubic inches. 52° F.

Arkansas Hot Springs. See Hot Springs of Arkansas.

Arrowhead Springs, San Bernardino, California. H₂S .644 grain, NaSO₄ 42.Auburn Spring, 4 miles west of Auburn, New York. CaSO₄ 120, MgSO₄ 26, H₂S 12 cubic inches.Avon Lower Spring, Livingston County, New York. CaCO₃ and CaSO₄ 87, MgSO₄ 50, H₂S 10 cubic inches. 45° to 47° F.Avon Sylvan Spring, Livingston County, New York. NaCl 97, CaCO₃ and CaSO₄ 107, and MgCl₂ and MgCO₃ and MgSO₄ 91, H₂S 20.64 cubic inches.Avon Upper Spring, Livingston County, New York. CaSO₄ 84, Na₂SO₄ and NaCl 34, MgSO₄ 10, H₂S 12 cubic inches. 51° F.

Bath Spring, Steamboat Springs, Colorado. 103° F.

Blount Springs, Blount County, Alabama. H₂S 30.67 cubic inches.Calistoga Hot Sulphuretted Spring, at the terminus of the Napa branch of the Pacific Railroad. NaCl 22, H₂S 10 cubic inches. 97° F.Castilian Spring, Sumner County, Tennessee. 52 cubic inches H₂S.

Chalk Creek Hot Springs, Chalk Creek, Colorado. 130° F.

Chittenango White Sulphur Spring, Madison County, New York. CaSO₄ 115, MgCO₃ 12, H₂S 12.8 cubic inches, CO₂ 18.4 cubic inches. 49° F.Delaware Sulphur Spring, Delaware County, Ohio. H₂S 96 cubic inches.Des Chutes Hot Springs, Wasco County, Oregon. Na₂CO₃ and NaCl 55. 143–145° F.French Lick Springs, Orange County, Indiana. NaCl 141, CaSO₄ 61, Na₂SO₄ 23, MgSO₄ 18, H₂S 25 1/2 cubic inches.

Gardner Magnesia Spring. See Sharon Spring.

Gentlemen's Pleasure Bath. See Hot Springs, Bath County, Virginia. 78° F.

Hager's Spring, Hagers County, Tennessee. H₂S 52 cubic inches.Healing Springs, Bath County, Virginia. CaCO₃ 18, CO₂ 5. 88° F

Hot Springs, Bath County, Virginia—

1. Ladies' Boiler Bath. 110° F.
2. Ladies' Sulphur Bath. 102° F.
3. Gentlemen's Pleasure Bath. 78° F.

Hot Springs, Cañon City, Fremont County, Colorado. CaCO_3 32, MgCO_3 12, NaCl 18. Temp. 95–102° F.

Hot Springs, Garland County, Arkansas. 93–105° F.

Hot Sulphur Springs, Middle Park, 90 miles from Denver, Colorado. Na_2SO_4 25, Na_2CO_3 22. Temp. 110–117° F.

Idaho Warm Spring Clear Creek County, California. Na_2CO_3 31, Na_2SO_4 29. 106° F.

Jordan's White Sulphur, Frederick County, Virginia. H_2S 2 cubic inches. 57° F.

Ladies' Boiler Bath. See Hot Springs, Bath County, Virginia. 110° F.

Ladies' Sulphur Bath. See Hot Springs, Bath County, Virginia. 102° F.

Liberty Hot Springs, Morrison, 12 miles from Denver, Colorado. Temp. 140–150.

Main Spring. See Paso Robles Springs.

Massena Springs, St. Lawrence County, New York. NaCl 80, CaSO_4 61, MgCl_2 30, H_2S 5 cubic inches.

Middle Park Springs, Colorado—

1. Bath Spring, Na_2SO_4 51, Na_2CO_3 46, NaCl 26. 117° F.

2. Red Sulphur, Na_2CO_3 120, NaCl 30, CaCO_3 20. 109.5° F.

3. Saline, Na_2CO_3 103, Na_2SO_4 18. 91° F.

Mud Spring. See Paso Robles Springs.

Ojo Caliente Spring, Ojo Caliente, New Mexico, Na_2CO_3 115, NaCl 32, F. CO_2 6, CO_2 ? Temp. 100° F.

Orkney Springs, Shenandoah County, Virginia. H_2S 4.88 cubic inches. 59.7° F.

Pago Hot Springs, 20 miles from headwater of San Juan River, Colorado—

No. 1. Na_2SO_4 129.

No. 2. Na_2SO_4 129.

No. 3. Na_2SO_4 131.

No. 4. Na_2SO_4 131.

Paroquet Springs, Bullitt County, Kentucky. NaCl 310, MgCl_2 48, H_2S 30 cubic inches.

Paso Robles Hot Sulphur Springs, San Luis, Obispo County, California—

1. Main Spring, NaCl and Na_2CO_3 49, CO_2 47. 112° F.

2. Mud Spring, NaCl and Na_2CO_3 108, CO_2 79. 122° F.

Pluto's Well. See French Lick Springs.

Proserpine Spring. See French Lick Spring.

Rochester or Longmuir's Sulphur Well, Rochester, New York. Na_2SO_4 56, NaCl 52, H_2S 17.26 cubic inches. 52° F.

Salt Lake Hot Spring, Utah. Temp. ?

Salt Sulphur Springs, Monroe County, West Virginia. CaSO_4 68, CaCO_3 33, MgSO_4 and Na_2SO_4 44, H_2S 19.12 cubic inches. 65.5° F.

Sharon Gardner Magnesia Spring, Schoharie County, New York. CaSO_4 93½, MgSO_4 20, H_2S 6 cubic inches. 48° F.

Sharon Red Spring, Schoharie County, New York. CaSO_4 and CaCO_3 186, MgSO_4 19, H_2S 10.5 cubic inches. 48° F.

Sharon White Sulphur Springs, Schoharie County, New York. CaSO_4 85, MgSO_4 34, $\text{Mg}(\text{HCO}_3)_2$ 24, H_2S 20.5 cubic inches. 48° F.

Siloam Springs, Bath Springs, Colorado, NaCl 1100. 103° F.

St. Helena, White Sulphur, Napa County, California.

No. 2. NaCl 22, H_2S 6.15 cubic inches. 64½° to 97° F.

No. 6. NaCl 23, Na_2SO_4 11, H_2S 4.25 cubic inches. 64½° to 97° F.

Unnamed Springs, Glenwood, Colorado, NaCl 2200. 124.2° F.

Wagonwheel Gap Springs. See Liberty Hot Springs.

Hot and Cold Springs in Animas Valley, Colorado.
 Hot Springs, near Boise City, Idaho. 196° F.
 Hot Springs, near Idaho City, Idaho.
 Hot Springs, near Pyramid Lake, Nevada. 208° F.
 Hot Spring, near Utah Central Railroad, Utah. 175° F.
 Hot Sulphuretted Spring, on Gila River, Arizona.
 Jemez Spring, in San Diego Cañon, fifty miles west of Santa Fe, New Mexico.
 Lake Tahoe Hot Springs, on border of Lake Tahoe, California.
 Malhuer River Springs, Baker County, Oregon. 193° F.
 Monroe Hot Springs, Castle Creek, sixty miles south of Prescott, Arizona. 150° F.
 Ouray Mineral Springs, Ouray, Colorado. Temperature, 120° to 134° F.
 Ouray Mineral Spring in Uncompahgre Park, Colorado. 120° to 140° F.
 Ouray Spring, Uncompahgre Park, 9 miles northwest of Ouray, Colorado. Temperature, 140° F.
 Platte Springs, near Fort Steele, Wyoming. 115° F.
 Puncta Springs, 60 miles west of Cañon City, Colorado. 120° F.
 San Bernardino Hot Springs, San Bernardino County, California.
 South Arkansas Mineral or Poncho Hot Springs, Lake County, Colorado. 120° F.
 Thermal Springs in Arizona.
 Warm Springs, 11 miles from Ogden City, Utah. 129° F.
 Warm Sulphur Springs, Elko, Elko County, Nevada.

UNANALYZED SPRINGS.

The following list of more than 300 unanalyzed springs shows clearly how much yet remains before we shall possess a full knowledge of our wealth of mineral waters. It is highly probable that many of these springs are of great value, and only await analysis to demonstrate their usefulness. When this is accomplished the medical profession will be quick to recognize their worth and make use of them in combating diseases.

Abenquis Springs, Walpole, Cheshire County, New Hampshire.
 Addison Mineral Spring, Addison Point, Washington County, Maine.
 Alagone Spring, Western Springs, Cook County, Illinois.
 Alkaline Spring, Yazoo County, Mississippi. 62° F.

Alum Spring, Marion County, Mississippi.
 Alum Spring, Marshall County, Mississippi.
 Alum Spring, Pike County, Mississippi.
 Amherst Spring, Hillsborough County, New Hampshire.
 Anchosa Spring, Anchosa Creek, near Quitman, Mississippi.
 Aqua Caliente, or Warner's Ranch Springs, San Diego County, California.
 Azule Mineral Spring, San José, Santa Clara County, California.
 Armstrong Spring, eight miles west of Searcy, Arkansas.
 Aspinock Mineral Springs, Windham County, Connecticut.
 Auburn Mineral Spring, Auburn, Androscoggin County, Maine.
 Bailey Springs, Lauderdale County, Alabama. 72° to 80° F.
 Bartlett Springs, Lake County California.
 Bath Chalybeate Spring, Bristol, Pennsylvania.
 Bedford Spring, Trimble County, Kentucky.
 Beersheba Springs, Grundy County, Tennessee.
 Bellbrook Magnesian Springs, Bellbrook, Greene County, Ohio.
 Belle Cheney Springs, Calcasieu Parish, Louisiana.
 Belmont Lithia Springs, New White House Station, New Kent County, Virginia.
 Berkshire Soda Springs, near Great Barrington, Massachusetts.
 Big Bone Springs, Boone County, Kentucky.
 Big Lick, Gallatin County, Kentucky.
 Birchdale Medical Springs, Bradford, Merrimack County, New Hampshire.
 Black Earth Mineral Spring, Black Earth, Dane County, Wisconsin.
 Black Sulphur Springs in Alabama.
 Black Sulphur Springs, Van Buren County, Arkansas.
 Blossburg Spring, Tioga County, Pennsylvania.
 Blue Spring, near New Amsterdam, Harrison County, Indiana.
 Bogard's Springs, Bogards Valley, Indiana.
 Box Mountain Sulphur, west of Todd's Gap, Kentucky. 63° F.
 Bozrah Mineral Spring, New London County, Connecticut.
 Bradford Spring, Merrimack County, New Hampshire.
 Brandywine Spring, Claiborne County, Mississippi.
 Brunswick Springs, Brunswick, Essex County, Vermont.
 Buenventer Springs, Kentucky.
 Burdell's Well, Caldwell County, Texas.
 Burner's or Seven Springs, Shenandoah County, Virginia.
 Burning Spring, Washington County, Alabama.
 Byron Acid Spring, Genesee County, New York.
 Catalytic Springs, one-half mile from Catalytic, Georgia.
 Caledonia Spring, Franklin County, Pennsylvania.
 Campbellsville Sulphur Water.
 Campo Chalybeate Spring, California.
 Carlisle Springs, Cumberland County, Pennsylvania.
 Carroll White Sulphur Springs, Alleghany County, Maryland. 43° F
 Cascado Spring, Michigan.
 Castalian Mineral Water, Olancho, Ingo County, California.
 Castilian Springs, Holmes County, Mississippi.

Chalybeate Springs abundant in Kentucky.

Chalybeate Springs in Pulaski County, Kentucky.

Chalybeate Springs, several on Rockcastle River, Kentucky.

Chalybeate Spring, Webster County, Kentucky.

Chalybeate Springs, Whitley County, Kentucky.

Chalybeate Springs, Clark County, Mississippi.

Chalybeate Spring, near Enterprise, Mississippi. 64.4° F.

Chalybeate Springs, Itawamba County, Mississippi.

Chalybeate Spring, near Warren's Mill, on Mackay's Creek, Mississippi.

Chalybeate Spring, Winston County, Mississippi.

Chalybeate Spring, Yallabusha County, Mississippi. 75.2° F.

Chalybeate Spring in Amherst, New Hampshire.

Chalybeate Spring, four miles from Pittsburgh, Pennsylvania. Temperature, 54°

Chappaqua Spring, Rensselaer County, New York.

Chase City Mineral Water, Chase City, Mecklenburgh County, Virginia.

Cheltenham Spring, St. Louis County, Missouri.

Chick's Springs, Greenville District, South Carolina.

Choteau Spring, Cooper County, Missouri.

Chrystal Mineral Spring, Stoneham, Middlesex County, Massachusetts.

Clear Creek Sulphur Spring, Kentucky.

Cold Sulphur Springs, Rockbridge County, Virginia.

Coleman's Well, Jackson County, Mississippi.

Colfax Springs, Colfax, Jasper County, Iowa.

Columbia Springs, Marion County, Mississippi.

Commonwealth Mineral Spring, Waltham, Middlesex County, Massachusetts.

Corydon Artesian Well, Harrison County, Indiana.

Corydon Saline Sulphur Well, half a mile east of Corydon, Harrison County, Ind

Cotton Wood Hot Springs, ninety miles west of Colorado Springs, Colorado.

Crabtree Sour Wells, Sulphur, Hopkins County, Texas.

Crittenden Springs, Crittenden County, Kentucky.

Crystal Spring, Napa County, California.

Cullum Springs, Bladen Springs, Choctaw County, Alabama.

Curdwell's Springs, Caldwell County, Texas.

Dovepark Springs, Dovepark, Clark County, Arkansas.
 Drennon Spring, Henry County, Kentucky.
 Dryden Springs, Tompkins County, New York.
 Dunbar's Mineral Springs, College Springs, Page County, Iowa.
 Eggleton's Springs, Giles County, Virginia.
 Electro-magnetic Springs (Fountain Park), Woodstock, Champaign County, Ohio.
 Elk Spring, Pike County, Missouri.
 Elkmont Springs, Giles County, Tennessee. 58° F.
 Elko Warm Spring, Idaho.
 Elgin Springs, Addison County, Vermont.
 Englewood Spring, Minneapolis, Hennepin County, Minnesota.
 Ephrata Spring, Lancaster County, Pennsylvania.
 Epsom Spring, Trimble County, Kentucky.
 Esculapia Springs, Lewis County, Kentucky.
 Eureka Springs, Eureka Springs, Carroll County, Arkansas.
 Excelsior Spring, Syracuse, New York.
 Everett Crystal Spring, Everett, Middlesex County, Massachusetts.
 Fairchild's Potash Sulphur Springs, Potash Sulphur, Garland County, Arkansas.
 Fauquier White Sulphur Springs, Fauquier County, Virginia.
 Ferguson's Chalybeate Spring, on southwest quarter of section 21, township 11, range 6.
 Flint's Springs, St. Joseph County, Michigan.
 Fox Spring, Fleming County, Kentucky.
 Frankfort Springs, Beaver County, Pennsylvania.
 Franklin Spring, at head of Well's Creek, Mississippi. 64° F.
 Garnet Springs, near Toccoa Falls, Georgia.
 Garrett Spring, one and a half miles from Spartanburg, South Carolina.
 Geron's Spring, in Northern Alabama.
 Geuda Springs, Geuda Springs, Sumner County, Kansas.
 Geyser Springs, Sonoma County, California.
 Gihon Mineral Spring, Delavan, Walworth County, Wisconsin.
 Glen Flora Mineral Springs, Waukegan, Lake County, Illinois.
 Gower's Spring, Gainesville, Georgia.
 Grand Ledge Spring, Eaton County, Michigan.
 Grayson Springs, Grayson County, Kentucky.
 Grayson Sulphur Springs, Carroll County, Virginia.
 Green Spring, Green Spring, Seneca County, Ohio.
 Green's Spring, Jefferson County, Illinois.
 Greenwood Spring, Monroe County, Mississippi.
 R. B. Grigsby's White Sulphur Mineral Water, Nelson County, Kentucky.
 Harbon Springs, twenty miles from Calistoga, California.
 Hardinsville Sulphur Spring, Franklin County, Kentucky.
 Hartford Cold Spring, Hartford, Oxford County, Maine.
 Healing Springs, Washington County, Alabama.
 Highland Rock Spring, Hartford County, Connecticut.
 Highland Tonica Spring, Hartford County, Connecticut.
 Holly Spring, Woonsocket, Providence County, Rhode Island.
 Hopkinton Springs, Middlesex County, Massachusetts.

~~Johnson Springs, Clark County, Alabama~~

Johnson's Wells, near Meridianville, Alabama.

Jones' White Sulphur and Chalybeate Springs, eleven miles from Warrenton, Carolina.

Kansas Artesian Mineral Wells, Manhattan, Riley County, Kansas.

Kellum Sulphur, Grimes County, Texas.

King's Mineral Springs, Muddy Fork, Clark County, Indiana.

Kirk Spring, Lewis County, Kentucky.

Kittrell's Springs, Granville County, North Carolina.

Knightstown Spring, Henry County, Indiana.

Lampasas Springs, Lampasas, Lampasas County, Texas.

Lauderdale Spring, Mississippi.

Lane's Spring, Stanislaus County, California.

Lava Springs in Grand Cañon of Colorado, Arizona.

Leinster Poison Springs, Statesville, Iredell County, North Carolina.

Lee's Springs, twenty miles northeast from Knoxville, Tennessee.

Lena-pi Magnesic Springs, Delaware, Delaware County, Ohio.

Limestone Springs, twenty-one miles from Spartansburg, South Carolina.

Little Geyser Springs, Lononia County, California.

Litton's Seltzer Spring, Litton's Station, Sonoma County, California.

Liverpool Well, New York.

Lower Soda Springs, Linn County, Oregon.

Low's Well, Saratoga County, New York.

Lunenburg Spring, Lunenburg, Essex County, Vermont.

Magnesia Spring, Tallulah Falls, Georgia.

Mammoth Spring, Fulton County, Arkansas. 60° F.

Mammoth Well, Nelson County, Kentucky.

Manasquan Spring, Point Pleasant, Ocean County, New Jersey.

Massie's or Red Sulphur Spring, Chillicothe, Georgia. 54.5° F.

McCallister's Soda Springs, 35 miles east of Jacksonville, Jackson County, Oregon.

Mershon or White Sulphur Spring, Georgia. 55° F.

Middleton Springs, Middleton, Rutland County, Vt.

Milford or Ponemah Springs. Milford, Hillsborough County, New Hampshire.

Mineral Springs, Washington County, Illinois.
 Mineral Springs at Cowpens Furnace, near Pacolet, South Carolina.
 Mineral Springs at the base of Henry's Knob, South Carolina.
 Mineral Spring near Parson's Mountains, Abbeville, South Carolina.
 Mineral Spring on the Soluda, near Pinson's Ford, South Carolina.
 Mineral Springs near Bingham City, Utah.
 Mineral Water, two miles from Dowlingsville, Grant County, Kentucky.
 Mineral Wells, Mineral Wells, Palo Pinto County, Texas.
 Mineral Wells, Parkersburg, Wood County, West Virginia.
 Mississippi Springs, Hinds County, Mississippi.
 Missisquoi Springs, viz.: Central, Missisquoi, Vermont, and Sheldon, are in Franklin County, Vermont.
 Monagaw Sulphuretted Springs, St. Clair County, Missouri.
 Montgomery White Sulphur Springs, Montgomery County, Virginia.
 Mooresville Mineral Spring, Mooresville, Livingstone County, Missouri.
 Morrison Springs, Morrison, twelve miles from Denver, Colorado.
 Moultonborough Mineral Springs, Moultonborough, Carroll County, New Hampshire.
 Mountain Valley Springs, 10 miles north of Hot Springs, Arkansas.
 Mt. Clemens Spring, Macombe County, Michigan.
 New Point Comfort Springs, Blue Lick, Clark County, Indiana.
 Ochu Mineral Springs, Providence, Providence County, Rhode Island.
 Oliver Springs, Daviess County, Kentucky.
 Oliver Springs, Anderson County, Tennessee.
 Ottumwa Mineral Springs, Ottumwa, Wapello County, Iowa.
 Oxford Mineral Spring, New Haven County, Connecticut.
 Panacea Springs, Panacea Springs, Halifax County, North Carolina.
 Parkersburg Mineral Wells, Wood County, Virginia.
 Pearson Springs, California.
 Perry County Springs, 11 miles from Carlisle, Pennsylvania. 72° F.
 Piedmont Sulphur Springs, 10 miles north-northeast of Navasota, Texas.
 Ponce de Leon Spring, Fulton County, Georgia.
 Powhatan Lithia, and Alum Springs, Ballsville, Powhatan County, Virginia.
 Pulaski Alum Springs, Pulaski County, Virginia.
 Quitman Red Sulphur, near Quitman, Missouri.
 Randolph Mineral Springs, Randolph, Randolph County, Missouri.
 Reed's Mineral Spring, Washington County, New York.
 Reuben Jesse's Mineral Water, Woodford County, Kentucky.
 Rockbridge Baths, Rockbridge County, Virginia.
 Rockingham Springs, Rockingham, Rockingham County, Virginia.
 Rinnah Wells Spring, Andalusia, Rock Island County, Illinois.
 Roanoke Red Sulphur Springs, Roanoke County, Virginia.
 Robinson's Spring, 20 miles from Nashville, Tennessee.
 Rob's Chalybeate Spring, McCracken County, Kentucky.
 Rochester Spring, 12 miles from Harrodsville, Boyle County, Kentucky.
 Roper's Wells, Butler County, Alabama.
 Russell Chalybeate Spring, Kentucky.
 Russell Sulphur Spring, Kentucky.

ington County).

Shannondale Springs, Jefferson County, Virginia.

Sheathil Rock Spring, Waukesha County, Wisconsin.

Shenandoah Alum Springs, Shenandoah County, Virginia.

Shelby Springs, Shelby County, Alabama.

Shocco Springs, Warren County, North Carolina.

Siloam Springs, Siloam Springs, Benton County, Arkansas

Skaggs Springs, California.

Slatersville Magnesic Spring, Slatersville, Tompkins County, New York.

Social Hill Mineral Water, Kentucky.

Sour Lake Springs, Sour Lake, Hardin County, Texas.

Springdale Seltzer Springs, Boulder County, Colorado.

St. Croix Mineral Spring, East Farmingham, Polk County, Wisconsin.

Steamboat Springs, 10 miles from Carson City, Washoe County Nevada.

Stone Spring, Lincoln County, Kentucky.

Stoveall's Spring, Marion County, Mississippi.

Sudduth or Mud Spring, Kentucky.

Sulphuretted Chalybeate Spring, half mile northeast of Pittsfield, New Hampshire.

Sulphuretted Springs are found in Jasper, Lawrence and Pike Counties, Indiana.

Sulphur Spring, Tallulah Falls, Georgia.

Sulphur Spring, 1 ½ miles east of Zoar, Georgia.

Sulphur Spring, southwest center of section 15, Illinois.

Sulphur Spring, Allen County, Kansas.

Sulphur Spring, Union County, Kentucky.

Sulphur Spring, in Calcasieu Parish, Louisiana.

Sulphur Spring, near Covington, Louisiana.

Sulphur Spring, near Enterprise, Mississippi.

Sulphur Springs, St. Clair County, Alabama.

Sulphur Springs, near Philadelphia, Mississippi.

Swayne's Mammoth Springs, Henry County, Tennessee.

Sweeney's Chalybeate Spring, Kentucky.

Syracuse Salt Well, Syracuse, New York.

Talladega Spring, Talladega County, Alabama.

Tipton Well, Jackson County, Mississippi.
 Tolenas Soda Springs, near Suisun City, Solano County, California.
 Topek Mineral Wells, Topek, Shawnee County, Kansas.
 Trinity Springs, Martin County, Indiana. 57° F.
 Tuscan or Lick Springs, near Red Bluff, Tehama County, California.
 Unity Springs, Newport, Sullivan County, New Hampshire.
 Valhermoso Springs, 18 miles from Huntsville, Alabama.
 Victor Spring, Genesee County, New York.
 Volcano Springs, Lander County, Nevada.
 Wallawhatoola Alum Springs, Millboro Depot, Bath County, Virginia.
 Warren White Sulphur Springs, Warren County, North Carolina.
 Washington Bell's Sulphur Water, Marion County, Kentucky.
 Weldon Spring, St. Alban's, Franklin County, Vermont.
 West Baden Springs, West Baden, Orange County, Indiana.
 Western Saratoga, Union County, Illinois.
 White Rock Mineral Spring, Waukesha, Waukesha County, Wisconsin.
 White Sulphur Springs, Sulphur Springs, De Kalb County, Alabama.
 White Sulphur Mineral Water, Marion County, Kentucky.
 White Sulphur, 6 miles east of Gower Springs, Georgia.
 White Sulphur Spring, Catawba County, North Carolina.
 White Sulphur Spring, Neshoba County, Mississippi.
 White Sulphur Spring of San Juan Capistras, 40 miles north of San Diego, California.
 White Sulphur Springs, Breckenridge County, Kentucky.
 White Sulphur Springs, Greene County, New York.
 White Sulphur Springs, Ohio County, Kentucky.
 White Sulphur Wells, Metcalf County, Kentucky.
 Wilbur Springs, near Colusa, California.
 Williamstown Springs, Anderson County, South Carolina.
 Wilson's Springs, Spartanburg District, South Carolina.
 Winchester Springs, Franklin County, Tennessee.
 Wolf Trap Lithia Springs, Wolf Trap, Halifax County, Virginia.
 Wooley's Springs, Limestone County, Alabama.
 Wootan Wells, Wootan Wells, Robertson County, Texas.
 Wyandotte Spring, Wayne County, Michigan.
 Yates Mineral Spring, Boyle County, Kentucky.
 Yellow Spring, Chester County, Pennsylvania.
 Yelvington Spring, Daviess County, Kentucky.
 Ypsilanti Mineral Spring, Washtenaw County, Michigan.
 York Springs, Adams County, Pennsylvania.
 Zonian Springs, near Elgin, Kane County, Illinois.

In preparing this article, the following works have been freely consulted: "Report on Mineral Springs," by Committee of American Medical Association, 1880; Dr. William Pepper, Reporter; Moorman and Walton's "Mineral Springs of the United States;" Peale's "Mineral Waters," 1887, Braun's "Curative Effects of Baths and Waters," 1875; U. S. Dispensatory, 1884; "Mineral Springs of Colorado," by Charles Denison, M.D., and Thesis on Mineral Springs of the United States, 1882, by Judson Daland, M.D.

1

BASED ON RECENT MEDICAL LITERATURE.

Gould's Medical Dictionaries.

BY GEORGE M. GOULD, A.M., M.D.,

Ophthalmic Surgeon to the Philadelphia Hospital, Editor of "The Medical News."

THE STANDARD MEDICAL REFERENCE BOOKS.

The Illustrated Dictionary of Medicine and Allied Sciences.

BEING AN EXHAUSTIVE LEXICON OF MEDICINE AND THOSE SCIENCES COLLATERAL TO IT: BIOLOGY (ZOOLOGY AND BOTANY), CHEMISTRY, DENTISTRY, PHARMACOLOGY, MICROSCOPY, ETC. With many Useful Tables and numerous Fine Illustrations. Large, Square Octavo. 1600 pages.

Full Sheep, or Half Dark-Green Leather, *net*, \$10.00;
Half Russia, Thumb Index, *net*, \$12.00

The Student's Medical Dictionary.

INCLUDING ALL THE WORDS AND PHRASES GENERALLY USED IN MEDICINE, WITH THEIR PROPER PRONUNCIATION AND DEFINITIONS, BASED ON RECENT MEDICAL LITERATURE. With Tables of the Bacilli, Micrococci, Leucomains, Ptomaines, etc., of the Arteries, Muscles, Nerves, Ganglia, and Plexuses; Mineral Springs of the U. S., etc. Tenth Edition. Small Octavo. 700 pages.

Half Dark Leather, *net*, \$3.25; Half Morocco, Thumb Index, *net*, \$4.00

"One pleasing feature of the book is that the reader can almost invariably find the definition under the word he looks for, without being referred from one place to another, as is too commonly the case in medical dictionaries. The tables of the bacilli, micrococci, leucomains, and ptomaines are excellent, and contain a large amount of information in a limited space. The anatomical tables are also concise and clear. . . . We should unhesitatingly recommend this dictionary to our readers, feeling sure that it will prove of much value to them."—*The American Journal of Medical Science.*

The Pocket Pronouncing Medical Lexicon.

12,000 WORDS PRONOUNCED AND DEFINED.

Double the Number in any Other Similar Book. Containing all the Words, their Definition and Pronunciation, that the Student generally comes in contact with; also elaborate Tables of the Arteries, Muscles, Nerves, Bacilli, etc., etc.; a Dose List in both English and Metric Systems, etc., arranged in a most convenient form for reference and memorizing. Thin 64mo. Full Limp Leather, Gilt Edges, *net*, \$1.00; Thumb Index, *net*, \$1.25

These books may be ordered through any bookseller, or upon receipt of price the publishers will deliver free to the purchaser's address. *Full descriptive circulars and sample pages sent free upon application.*

P. BLAKISTON, SON & CO., Publishers, Philadelphia.

SECOND EDITION.

DISEASES OF THE SKIN.

CROCKER.

THEIR DESCRIPTION, PATHOLOGY, DIAGNOSIS, AND TREATMENT, with Special Reference to the Skin Eruptions of Children. By H. RADCLIFFE CROCKER, M. D., F. R. C. P., Physician to the Department of Skin Diseases, University College Hospital, London. 92 Illustrations. Second Edition, Revised and Enlarged. Octavo. Cloth, \$4.50 net.

"It commends itself by its very honesty. What the author believes he says, and, if he does not know a thing, he says that he does not know. We know that he is a man of large experience in the care of skin cases, as he is the Dermatologist to University College Hospital. That he is an acute and careful observer, and a systematic recorder of his observations, his book teaches us. As one of the physicians to the East London Hospital for Children he has enjoyed special advantages in the study of the skin diseases of children, and in the work now before us special attention is given to the symptomatology of such diseases—a unique feature of the book. He is, also, a skilled pathologist. Nearly all the drawings from microscopical sections are his own, and very beautiful they are as works of art. Another special and commendable feature of the book is that the general pathology of the various diseases is printed in the ordinary type of the book, while the minute anatomy is given in smaller type. This will be appreciated by many readers. Great prominence is given to constitutional conditions as causes of skin diseases, and to the importance of treating the patient quite as much as his cutaneous disorder. Dietetic matters and general regimen are insisted upon, and wisely. The definitions of the terms used in dermatology are good and useful. The general practitioner often fails to convey a proper idea of the disease to his consultant because he does not know the correct terms to use in describing a case. The chapters on etiology are excellent. It is one of the most thorough treatises on diseases of the skin that it has ever been our good fortune to review, and, as such, we heartily commend it to our readers."—*New York Medical Journal*.

"The author has endeavored to embody in the work before us a succinct statement of our present knowledge of dermatology, which should serve as a reference to the general practitioner, as well as a working manual for the student. No one familiar with Dr. Crocker's ability and clinical opportunities will question his eminent fitness and thorough equipment for his task, in the accomplishment of which he has brought to bear the matured results of his twelve years' experience in the Skin Department of the University College Hospital of London."—*Journal of Cutaneous and Genito-Urinary Diseases, New York*.

. A descriptive circular of Crocker, giving the opinions of fifteen of the leading Dermatologists and teachers, will be sent free upon application.

P. BLAKISTON, SON & CO., PUBLISHERS,
1012 WALNUT STREET, PHILADELPHIA.

POTTER'S HANDBOOK.

MATERIA MEDICA, PHARMACY AND THERAPEUTICS.

FIFTH EDITION. REVISED. 805 PAGES. OCTAVO.

With Thumb Index. Cloth, net, \$4.00; Leather, net, \$5.00.

A Handbook of Materia Medica, Pharmacy and Therapeutics
—including the Physiological Action of Drugs, Special Therapeutics of Diseases, Official and Extemporaneous Pharmacy, etc. By SAML. O. L. POTTER, M.A., M.D., Professor of the Practice of Medicine in Cooper Medical College, San Francisco; Late A. A. Surgeon, U. S. Army, Author of "Speech and its Defects," and the "Quiz-Compendis" of Anatomy and Materia Medica, etc.

DR. POTTER has become well-known by his *Compendis* of Anatomy, and of *Materia Medica*, both of which have reached four editions. In this book, more elaborate in its design, he has shown his abilities to much better advantage, and all who examine or use it will agree that he has produced a work containing more correct information in a practical, concise form than any other publication of the kind. The plan of the work is new, and its contents have been combined and arranged in such a way that it offers a compact statement of the subjects in hand.

PART I.—*MATERIA MEDICA* AND THERAPEUTICS, the drugs being arranged in alphabetical order, with the synonym of each first; then the description of the plant, its preparations, physiological action, and lastly its *Therapeutics*. This part is preceded by a section on the classification of medicines.

PART II.—PHARMACY AND PRESCRIPTION WRITING. This is written for the use of physicians who desire or of necessity must put up their own prescriptions, and includes a great deal of valuable information.

PART III.—SPECIAL THERAPEUTICS. An Alphabetical List of Diseases—giving the drugs that have been found serviceable in each disease, and the authority recommending the use of each. Over 600 prescriptions are given in this part, many being over the names of eminent men.

THE APPENDIX contains many tables, formulæ and much collateral material.

From the Therapeutic Gazette.

"No new remedy of any acknowledged value is omitted from this list. Under each the section on physiological action and therapeutics has been written with care. . . . In the enumeration of drugs suited to different disorders a very successful effort at discrimination has been made, both in the stage of disease and in the cases peculiarly suited to the remedy. It is no mere list of diseases followed by a catalogue of drugs, but is a digest of modern therapeutics, and as such will prove of immense use to its possessor."

P. BLAKISTON, SON & CO., PUBLISHERS,
1012 WALNUT STREET, PHILADELPHIA.

COLORED ILLUSTRATIONS.

"A BRIDGE BETWEEN PHYSIOLOGY AND PRACTICAL MEDICINE."

HUMAN PHYSIOLOGY.

BY LANDOIS AND STIRLING.

FOURTH AMERICAN, FROM THE SEVENTH GERMAN EDITION.

A Text-Book of Human Physiology, including Histology and Microscopical Anatomy, with special reference to the requirements of Practical Medicine. By Dr. L. LANDOIS, Professor of Physiology and Director of the Physiological Institute, University of Greifswald. Translated from the Fifth German Edition, with additions by WM. STIRLING, M.D., Sc.D., Brackenbury Professor of Physiology and Histology in Owen's College and Victoria University, Manchester; Examiner in the Honors School of Science, University of Oxford, England. Fourth Edition, revised and enlarged. 845 Illustrations.

Thoroughly Revised and Enlarged. *In Press.*

From the Prefaces to the English Edition.

The fact that Prof. Landois' book has passed through four large editions in the original, and that in barely six months' time a second edition of the English has been called for, shows that in some special way it has met a want. The characteristic which has thus commended the work will be found mainly to lie in its eminent practicality; and it is this consideration which has induced me to undertake the task of putting it into English. Landois' work, in fact, forms a *Bridge* between Physiology and the Practice of Medicine. It never loses sight of the fact that the student of to-day is the practicing physician of to-morrow. In the same way, the work offers to the busy physician in practice a ready means of refreshing his memory on the theoretical aspects of Medicine. He can pass backward from the examination of pathological phenomena to the normal processes, and in the study of these, find new indications and new lights for the appreciation and treatment of the cases under consideration. With this object in view, all the methods of investigation which may, to advantage, be used by the practitioner, are carefully and fully described. Many additions, and about one hundred illustrations, have been introduced into this edition, and the whole work carefully revised.

PRESS NOTICES.

"Most effectively aids the busy physician to trace from morbid phenomena back the course of divergence from physical operations, and to gather in this way new lights and novel indications for the COMPREHENSION AND TREATMENT of the maladies with which he is called upon to cope."—*American Journal of Medical Sciences.*

"We have no hesitation in saying that THIS IS THE WORK to which the PRACTITIONER will turn whenever he desires light thrown upon the phenomena of a COMPLICATED OR IMPORTANT CASE."—*Edinburgh Medical Journal.*

"One of the MOST PRACTICAL WORKS on Physiology ever written, forming a 'bridge' between Physiology and Practical Medicine. . . . Its chief merits are its completeness and conciseness. . . . EXCELLENTLY CLEAR, ATTRACTIVE AND SUCCINCT."—*British Medical Journal.*

"As a work of reference, LANDOIS and STIRLING's Treatise OUGHT TO TAKE THE FOREMOST PLACE among the text-books in the English language. The wood-cuts are noticeable for their number and beauty."—*Glasgow Medical Journal.*

"Landois' Physiology is, without question, the best text-book on the subject that has ever been written."—*New York Medical Record.*

"The chapter on the Brain and Spinal Cord will be a most valuable one for the general reader, the translator's notes adding not a little to its importance. The sections on Sight and Hearing are exhaustive. . . . The Chemistry of the Urine is thoroughly considered. . . . The text is smooth, accurate, and unusually free from Germanisms; in fact, it is good English."—*New York Medical Journal.*

P. BLAKISTON, SON & CO., PUBLISHERS,
1012 WALNUT STREET, PHILADELPHIA.

"AN EMINENTLY PRACTICAL BOOK."

MOULLIN'S SURGERY,

SECOND EDITION, ENLARGED.

A Complete Practical Treatise on Surgery, with Special Reference to Treatment.

By C. W. MANSELL MOULLIN, M.A., M.D. OXON., F.R.C.S.,
SURGEON AND LECTURER ON PHYSIOLOGY TO THE LONDON HOSPITAL, ETC.

THIRD AMERICAN EDITION,

Edited by JOHN B. HAMILTON, M.D., LL.D.,

Professor of the Principles of Surgery and Clinical Surgery, Rush Medical College, Chicago; Professor of Surgery, Chicago Polyclinic; Formerly Supervising Surgeon-General, U. S. Marine Hospital Service; Surgeon to Presbyterian Hospital, St. Joseph's Hospital, and Central Free Dispensary, Chicago, etc.

About Six Hundred Illustrations,

Two hundred of which are original with this work, and many of which are printed in several colors.


Royal Octavo. 1200 Pages. Handsome Cloth, \$6.00; Leather, Raised Bands, \$7.00; Half Russia, Crushed, Marble Edges, \$8.00.

THESE PRICES ARE ABSOLUTELY NET.

This new edition has been very thoroughly revised, important additions have been incorporated,—including a chapter on military surgery,—many of the old illustrations have been recut, and about one hundred new figures added.

Our claim that *Moullin's Surgery* is the best text-book for the student and general work of reference for the practitioner is based upon the reviews of a large number of journals that have pronounced it eminently practical, and upon the fact that so many teachers have seen fit to recommend it. But beyond this we may say that broad principles are stated in a clear, authoritative manner, that the relative value of the different subjects has been carefully considered, and that about the whole there is an air of responsibility that renders plain the fact that the author knows whereof he speaks, not only from his own experience, but from a large acquaintance with American and foreign literature. There is also a uniformity of style, an elegance of diction, that attracts and interests the reader, while it makes plain the subject under discussion.

"The aim to make this valuable treatise practical by giving special attention to questions of treatment has been admirably carried out. Many a reader will consult the work with a feeling of satisfaction that his wants have been understood, and that they have been intelligently met. He will not look in vain for details, without proper attention to which he well knows that the highest success is impossible."—*The American Journal of Medical Sciences, Philadelphia.*

 A complete circular, with sample pages, press notices, etc., will be sent free upon application.

P. BLAKISTON, SON & CO., PUBLISHERS,
1012 WALNUT STREET, PHILADELPHIA.

JUST READY.

Morris's Human Anatomy.

A NEW TEXT-BOOK.


791 Illustrations, 214 of which are Printed in Several Colors, and most of which are original. OCTAVO. 1286 PAGES.

Handsome Cloth, *net*, \$6.00; Full Sheep, *net*, \$7.00; Half Russia, Marbled Edges, *net*, \$8.00.

"While it must be admitted that there can be scarcely anything new in descriptive anatomy, it is refreshing to notice a departure in the arrangement and treatment of this subject which is PRACTICAL, USEFUL, and INTERESTING. In a word, the natural method is adopted, and several new features of illustration are introduced. For instance, the origin and insertion of muscles with exact areas of attachment are noted in different colored outlines, red for the former and blue for the latter. Thus the reader is enabled at a glance to discriminate in the case of a given bone the difference between the two points, and obtain thereby the direction and extent of action of each muscle. A similar principle is carried out in other parts of the work, which makes the illustrations for the most part unique and invaluable, the schematic drawings especially demanding attention and commendation in this connection. The work as a whole is filled with practical ideas, and the salient points of the subjects are properly emphasized. The surgeon will be particularly edified by the section on the topographical anatomy, which is full to repletion of excellent and useful illustrations."—*Medical Record, New York.*


"The accuracy of description and thoroughness of detail in the work before us is no doubt due, in a large measure, to the fact that the editor in preparing the book has availed himself of the valuable services of many eminent authors, who are known to have devoted special attention to the subjects allotted to them. In looking over the arrangements of authors and subjects, we find many names which afford us an insight into the character of the work. . . . A glance at these names, of ten of the foremost anatomists and surgeons of the English-speaking world, impresses one with the conviction that Professor Morris knew his men, and that their combined efforts have given us a treatise on anatomy which for a long time to come, we believe, is destined to fill an important place as a students' text-book and work of reference.

"The value of the book is greatly enhanced by the numerous illustrations which adorn its pages, over two hundred of which are printed in colors from drawings made expressly for this work by special artists, who have spared no pains to make each cut clear and representative of the normal structures and relations. . . . A feature of the book which will undoubtedly facilitate the work of students is found in the illustrations of the bones, where the origin of the muscles are indicated by red lines, the insertions by blue lines, and the attachment of ligaments by dotted black lines. So far as type, paper, and size is concerned, the book is all that could be desired, and, taken all in all, the production is one which reflects much credit on both its authors and publishers."—*Buffalo Medical and Surgical Journal, Buffalo, N. Y.*

 Descriptive circular with samples of pages and illustrations sent free upon application.

P. BLAKISTON, SON & CO., PUBLISHERS,
1012 WALNUT STREET, PHILADELPHIA.









H121 Gould, George M.
G71 Student's medical
1896 dictionary, 33028
10th ed.

NAME

DATE _____

2000

